
TURTLES OF THE WORLD

Annotated Checklist and Atlas of Taxonomy, Synonymy, Distribution, and Conservation Status (10th Ed.)

Turtle Taxonomy Working Group

Anders G.J. Rhodin, John B. Iverson,
Uwe Fritz, Natalia Gallego-García, Arthur Georges,
H. Bradley Shaffer, and Peter Paul van Dijk



Chelonian Research Monographs, Number 10

Published by
Chelonian Research Foundation and Turtle Conservancy

in association with
IUCN SSC Tortoise and Freshwater Turtle Specialist Group, Turtle Taxonomy Fund,
Turtle Conservation Fund, and IUCN Species Survival Commission

Chelonian Research Monographs
Contributions in Turtle and Tortoise Research

Editorial Board

Anders G.J. Rhodin
Chelonian Research Foundation and
Turtle Conservancy
Arlington, Vermont 05250 USA
[RhodinCRF@aol.com]

John B. Iverson
Earlham College and
Turtle Survival Alliance
Richmond, Indiana 47374 USA
[johni@earlham.edu]

Peter Paul van Dijk
Turtle Conservancy and
Chelonian Research Foundation
Herndon, Virginia 20170 USA
[peterpaul@turtleconservancy.org]

CHELONIAN RESEARCH MONOGRAPHS (CRM) (ISSN 1088-7105) is an international peer-reviewed scientific publication series for monograph-length manuscripts, collected proceedings of symposia, edited compilations, and other longer research documents focused on turtles and tortoises. The series accepts contributions dealing with any aspects of chelonian research, with a preference for conservation or biology of freshwater and terrestrial turtles and tortoises. Bibliographic and other reference materials are also of potential interest. Submit manuscripts directly to Anders Rhodin at the e-mail address above. The series is published on an occasional basis, from 1996–2016 by Chelonian Research Foundation, and from 2017 and on by Chelonian Research Foundation and Turtle Conservancy.

Published CRM Issues

CRM 1. The Galápagos Tortoises: Nomenclatural and Survival Status. 1996. By Peter C.H. Pritchard. 85 pp. Open access pdf download available at www.chelonian.org/wp-content/uploads/file/CRM%201/CRM_1_1996_Pritchard_Galapagos_Tortoises.pdf.

CRM 2. Asian Turtle Trade: Proceedings of a Workshop on Conservation and Trade of Freshwater Turtles and Tortoises in Asia. 2000. Edited by Peter Paul van Dijk, Bryan L. Stuart, and Anders G.J. Rhodin. 164 pp. Open access pdf download available at www.chelonian.org/wp-content/uploads/file/CRM%202/CRM_2_Asian_Turtle_Trade_2000.pdf.

CRM 3. Biology and Conservation of Florida Turtles. 2006. Edited by Peter A. Meylan. 376 pp. Open access pdf download available at www.chelonian.org/wp-content/uploads/file/CRM%203/CRM_3_2006_Meylan_Florida_Turtles2.pdf.

CRM 4. Defining Turtle Diversity: Proceedings of a Workshop on Genetics, Ethics, and Taxonomy of Freshwater Turtles and Tortoises. 2007. Edited by H. Bradley Shaffer, Nancy N. FitzSimmons, Arthur Georges, and Anders G.J. Rhodin. 200 pp. Open access pdf download available at www.chelonian.org/wp-content/uploads/file/CRM%204/CRM_4_Defining_Turtle_Diversity_2007.pdf.

CRM 5 (Installments 1–18, 127 accounts to date). Conservation Biology of Freshwater Turtles and Tortoises: A Compilation Project of the IUCN/SSC Tortoise and Freshwater Turtle Specialist Group. 2008–2024. Edited variously by Anders G.J. Rhodin, John B. Iverson, Peter Paul van Dijk, Kurt A. Buhlmann, Peter C.H. Pritchard, Craig B. Stanford, Eric V. Goode, Raymond A. Saumure, and Russell A. Mittermeier. 2680 pp. to date. All accounts available as open access pdf downloads at www.iucn-tftsg.org/cbftt/.

CRM 6. Turtles on the Brink in Madagascar: Proceedings of Two Workshops on the Status, Conservation, and Biology of Malagasy Tortoises and Freshwater Turtles. 2013. Edited by Christina M. Castellano, Anders G.J. Rhodin, Michael Ogle, Russell A. Mittermeier, Herilala Randriamahazo, Rick Hudson, and Richard E. Lewis. 184 pp. Open access pdf download available at www.chelonian.org/wp-content/uploads/file/CRM%206/CRM_6_Castellano_etal_2013.pdf.

CRM 5 (Installment 8). Turtles and Tortoises of the World During the Rise and Global Spread of Humanity: First Checklist and Review of Extinct Pleistocene and Holocene Chelonians. 2015. TEWG [Turtle Extinctions Working Group: Anders G.J. Rhodin, Scott Thomson, Georgios L. Georgalis, Hans-Volker Karl, Igor G. Danilov, Akio Takahashi, Marcelo S. de la Fuente, Jason R. Bourque, Massimo Delfino, Roger Bour, John B. Iverson, H. Bradley Shaffer, and Peter Paul van Dijk]. 66 pp. Open access pdf download available at www.iucn-tftsg.org/wp-content/uploads/file/Accounts/crm_5_000e_fossil_checklist_v1_2015.pdf.

CRM 7. Turtles of the World: Annotated Checklist and Atlas of Taxonomy, Synonymy, Distribution, and Conservation Status (8th Ed.). 2017. TTWG [Turtle Taxonomy Working Group: Anders G.J. Rhodin, John B. Iverson, Roger Bour, Uwe Fritz, Arthur Georges, H. Bradley Shaffer, and Peter Paul van Dijk]. 292 pp. Open access pdf download available at www.iucn-tftsg.org/wp-content/uploads/file/Accounts2/TTWG_checklist_v8_2017.pdf.

CRM 8. Turtles of the World: Annotated Checklist and Atlas of Taxonomy, Synonymy, Distribution, and Conservation Status (9th Ed.). 2021. TTWG [Turtle Taxonomy Working Group: Anders G.J. Rhodin, John B. Iverson, Roger Bour, Uwe Fritz, Arthur Georges, H. Bradley Shaffer, and Peter Paul van Dijk]. 472 pp. Open access pdf download available at www.iucn-tftsg.org/wp-content/uploads/crm.8.checklist.atlas_.v9.2021.e3.pdf.

CRM 9. Range-Wide Demographic Collapse and Extinction Dynamics of the Endemic Burmese Roofed Turtle, *Batagur trivittata*, in Myanmar. 2024. Steven G. Platt, Win Ko Ko, Kalyar Platt, Tint Lwin, Swann Htet Naing Aung, Khin Myo Myo, Me Me Soe, Myo Min Win, Kyaw Thu Zaw Wint, Htun Thu, Shine Hsu Hsu Naing, Brian D. Horne, and Thomas R. Rainwater. 26 pp. Open access pdf download available at www.chelonian.org/wp-content/uploads/file/CRM9/Platt_CRM_9_Batagur_trivittata_2024.pdf.


CRM 10. Turtles of the World: Annotated Checklist and Atlas of Taxonomy, Synonymy, Distribution, and Conservation Status (10th Ed.). 2025. TTWG [Turtle Taxonomy Working Group: Anders G.J. Rhodin, John B. Iverson, Uwe Fritz, Natalia Gallego-García, Arthur Georges, H. Bradley Shaffer, and Peter Paul van Dijk]. 575 pp. Open access pdf download available at www.iucn-tftsg.org/wp-content/uploads/crm.10.checklist.atlas.v10.2025.pdf.

CHELONIAN RESEARCH MONOGRAPHS (CRM) issues are variously available for purchase from Chelonian Research Foundation and Turtle Conservancy. Contact either Turtle Conservancy (www.turtleconservancy.org) or Chelonian Research Foundation (www.chelonian.org/crm; 564 Chittenden Dr., Arlington, VT 05250 USA; 978-807-2902; RhodinCRF@aol.com) for prices, titles, and to place orders for available print copies. Chelonian Research Foundation (founded in 1992) and Turtle Conservancy (founded in 2005 as Chelonian Conservation Center, renamed in 2010) are nonprofit tax-exempt organizations under section 501(c)(3) of the Internal Revenue Code.

Copyright © 2025 by Chelonian Research Foundation and Turtle Conservancy

TURTLES OF THE WORLD

Annotated Checklist and Atlas of Taxonomy, Synonymy, Distribution, and Conservation Status (10th Ed.)



Stephen D. Nash

COVER AND TITLE PAGES ILLUSTRATIONS

***Front Cover*:**

Left: African Spurred Tortoise, *Centrochelys sulcata* (Testudinidae), Senegal. Photo by Tomas Diagne.

Right: Williams' Side-necked Turtle, *Phrynops williamsi* (Chelidae), Brazil. Photo by Vinicius T. de Carvalho.

***Back Cover*:**

Top Left: Sri Lankan Flapshell Turtle, *Lissemys ceylonensis* (Trionychidae), Sri Lanka. Photo by Anslem de Silva.

Top Right: Painted Terrapin, *Batagur borneoensis* (Geoemydidae), Malaysia. Photo by Douglas B. Hendrie.

Bottom Left: Kemp's Ridley Sea Turtle, *Lepidochelys kempii* (Cheloniidae), Florida, USA. Photo by Blair Witherington.

Bottom Right: Vallarta Mud Turtle, *Kinosternon (Kinosternon) vogti* (Kinosternidae), Mexico. Photo by Carolina Sánchez Arias.

***Title Pages*:**

p. i: Radiated Tortoise, *Astrochelys radiata* (Testudinidae), Madagascar. Drawing by Stephen D. Nash.

p. iv: Ploughshare Tortoise, *Astrochelys yniphora* (Testudinidae), Madagascar. Drawing by Stephen D. Nash.

p. vi, *Left*: Llanos Sideneck Turtle, *Podocnemis vogli* (Podocnemididae), Venezuela. Photo by Anders G.J. Rhodin.

p. vi, *Right*: Asian Giant Softshell Turtle, *Pelochelys cantorii* (Trionychidae), Cambodia. Photo by Annette Olsson.

p. xvi, *Left*: Pantanal Swamp Turtle, *Acanthochelys macrocephala* (Chelidae), Paraguay. Photo by Thomas and Sabine Vinke.

p. xvi, *Right*: European Pond Turtle, *Emys orbicularis orbicularis* (Emydidae), Ukraine. Photo by Uwe Fritz.

RECOMMENDED CITATION

TTWG [Turtle Taxonomy Working Group: Rhodin, A.G.J., Iverson, J.B., Fritz, U., Gallego-García, N., Georges, A., Shaffer, H.B., and van Dijk, P.P.]. 2025. Turtles of the World: Annotated Checklist and Atlas of Taxonomy, Synonymy, Distribution, and Conservation Status (10th Ed.). In: Rhodin, A.G.J., Iverson, J.B., van Dijk, P.P., Stanford, C.B., Goode, E.V., Buhlmann, K.A., and Mittermeier, R.A. (Eds.). Conservation Biology of Freshwater Turtles and Tortoises: A Compilation Project of the IUCN SSC Tortoise and Freshwater Turtle Specialist Group. Chelonian Research Monographs 10:1–575. doi:10.3854/crm.10.checklist.atlas.v10.2025.

Chelonian Research Monographs. Series Edited by Anders G.J. Rhodin, John B. Iverson, and Peter Paul van Dijk.
Volume Number 10. Turtles of the World: Annotated Checklist and Atlas of Taxonomy, Synonymy, Distribution, and Conservation Status (10th Ed.). In: Conservation Biology of Freshwater Turtles and Tortoises: A Compilation Project of the IUCN SSC Tortoise and Freshwater Turtle Specialist Group.

Published online 27 July 2025

ISSN (monograph series): 1088-7105
ISBN (this volume): 978-0-9910368-6-8 (digital online); 978-0-9910368-7-5 (hard cover); 978-0-9910368-8-2 (soft cover)

Co-Published by Chelonian Research Foundation and Turtle Conservancy.
Copyright © 2025 by Chelonian Research Foundation, Arlington, Vermont, USA
and Turtle Conservancy, Ojai, California, USA.
Printed by Mercury Print Productions, Rochester, New York, USA.


DEDICATION

We dedicate this book
and our ongoing turtle work to our close friend

Russell A. Mittermeier

for his long-standing and continuing contributions to turtle taxonomy and conservation.

He continues to inspire us.




Cristina G. Mittermeier

RUSSELL A. MITTERMEIER

We thank him for his long-term support and encouragement and inspirational leadership dating back to the early 1970s when he and Anders Rhodin first became friends at Dartmouth and then worked together on turtle taxonomy at Harvard's Museum of Comparative Zoology and began describing new turtle species together.

His career since then has been transformational for global species conservation issues, helping to found the IUCN SSC Tortoise and Freshwater Turtle Specialist Group and chairing the IUCN SSC Primate Specialist Group, serving as President of Conservation International for 25 years, being honored with both the Behler Turtle Conservation Award and the Indianapolis Prize for Animal Conservation in 2018, and currently serving as Chief Conservation Officer at Re:wild.



Stephen D. Nash

TURTLES OF THE WORLD

Annotated Checklist and Atlas of Taxonomy, Synonymy, Distribution, and Conservation Status (10th Ed.)

Turtle Taxonomy Working Group

Anders G.J. Rhodin, John B. Iverson, Uwe Fritz, Natalia Gallego-García, Arthur Georges, H. Bradley Shaffer, and Peter Paul van Dijk

Supplement No. 3 of Chelonian Research Monographs No. 5:

Conservation Biology of Freshwater Turtles and Tortoises: A Compilation Project of the IUCN SSC Tortoise and Freshwater Turtle Specialist Group

Edited by

Anders G.J. Rhodin[1,2], John B. Iverson[3], Peter Paul van Dijk[1,2], Craig B. Stanford[5], Eric V. Goode[2], Kurt A. Buhlmann[6], and Russell A. Mittermeier[4]

[1]*Chelonian Research Foundation, Arlington, Vermont, USA;*
[2]*Turtle Conservancy, Ojai, California, USA;*
[3]*Earlham College, Richmond, Indiana, USA;*
[4]*Re:wild, Austin, Texas, USA;*
[5]*University of Southern California, Los Angeles, California, USA;*
[6]*University of Georgia, Aiken, South Carolina, USA*


Chelonian Research Monographs
Number 10

2025
Chelonian Research Foundation and Turtle Conservancy









Llanos Sideneck Turtle,
Podocnemis vogli (Podocnemididae),
Hato El Cedral, Mantecal, Apure, Venezuela;
photo by Anders G.J. Rhodin.

Asian Giant Softshell Turtle,
Pelochelys cantorii (Trionychidae),
Kratie, Mekong River, Cambodia;
photo by Annette Olsson.

TURTLES OF THE WORLD

Annotated Checklist and Atlas of Taxonomy, Synonymy, Distribution, and Conservation Status (10th Ed.)

Table of Contents

New or Resurrected or Taxonomically Changed Taxa Recognized in this Checklist = *
Extinct Taxa = †

ABSTRACT 1

INTRODUCTION
Methodology 2
Body Size (Carapace Length) and Sexual Size Dimorphism 5
Type Specimens 8
Etymologies 9
Taxonomic Changes 10
TTWG Guidelines for Taxonomic Changes 11
Distributions 13
GIS Maps and Historical Ranges 17
Conservation Status 23
Status Updates and Discussion 24
Conservation Status 24
Genetic Pollution 27
Request for Updates 28
Acknowledgments 29
Photos and Photographers 30

TABLES
1. Extinct modern turtles 3
2. Summary of new and proposed taxonomic changes for modern taxa recognized in this Checklist 6
3. Summary of new and proposed taxonomic changes for extinct Holocene taxa recognized in this Checklist 6
4. Summary of proposed taxonomic changes not recognized in this Checklist 7
5. Top 50 largest modern turtle taxa 8
6. Top 50 smallest modern turtle taxa 8
7. Frequency of types of name etymologies 10
8. Top turtle-rich countries for extant species 12
9. Top turtle-rich countries for extant taxa 12
10. Top turtle-rich countries for endemic taxa 12
11. Top 50 turtle taxa with smallest ranges 20
12. Top 50 turtle taxa with largest ranges 20
13. Top 50 turtle species with largest ranges 21
14. Top 50 turtle taxa with smallest EOOs 23
15. Top 50 turtle taxa with largest EOOs 24
16. Top 75 most endangered turtle taxa 25

Linnaean Classification Content **32**

CHECKLIST
Modern Turtle and Tortoise Taxa Extant Since 1500 CE **33**

Order Testudines **33**
Suborder Pleurodira **33**
Superfamily Cheloidea **33**
Family Chelidae **33**
Subfamily Chelinae **34**
Acanthochelys 34
A. macrocephala 34
A. pallidipectoris 35
A. radiolata 35
A. spixii 36
Chelus 37
C. fimbriata 37
C. orinocensis 38
Mesoclemmys 38
M. dahli 38
M. gibba 39
M. nasuta 40
M. perplexa 40
M. raniceps 41
M. tuberculata 41
M. vanderhaegei 42
M. wermuthi 43
M. zuliae 43
Phrynops 44
P. geoffroanus 44
P. hilarii 45
P. tuberosus 46
P. williamsi 46
Platemys 47

P. platycephala 47
P. p. platycephala 47
P. p. melanonota 48
Ranacephala 49
R. hogei 49
Rhinemys 49
R. rufipes 49
Subfamily Hydromedusinae **50**
Hydromedusa 50
H. maximiliani 50
H. tectifera 51
Subfamily Chelodininae **52**
Chelodina 52
(*Chelodina*) 52
C. (C.) canni 52
C. (C.) gunaleni 53
C. (C.) longicollis 53
C. (C.) mccordi 54
C. (C.) m. mccordi 54
C. (C.) m. timorensis 55
C. (C.) novaeguineae 55
C. (C.) pritchardi 56
C. (C.) reimanni 56
C. (C.) steindachneri 57
(*Chelydera*) 57
C. (C.) burrungandjii 57
C. (C.) expansa 58
C. (C.) kuchlingi 59
C. (C.) kurrichalpongo 59
C. (C.) parkeri 60
C. (C.) rugosa 60
(*Macrochelodina*) 61
C. (M.) oblonga 61
Subfamily Emydurinae **62**
Elseya 62
(*Elseya*) 62
E. (E.) branderhorsti 62
E. (E.) dentata 63
E. (E.) flaviventris 63
(*Hanwarachelys*) 64
E. (H.) novaeguineae 64
E. (H.) rhodini 65
E. (H.) schultzei 66
(*Pelocomastes*) 67
E. (P.) albagula 67
E. (P.) irwini 68
E. (P.) lavarackorum 68
Elusor 69
E. macrurus 69
Emydura 70
E. australis 70
E. gunaleni 71
E. macquarri 71
E. m. macquarri 72
E. m. emmotti 73
E. m. krefftii 73
E. m. nigra 74
E. subglobosa 74
E. s. subglobosa 75
E. s. worrelli 76
E. tanybaraga 76
Myuchelys 77
M. bellii 77
M. georgesi 78
M. latisternum 78
M. purvisi 79
Rheodytes 80
R. leukops 80
Subfamily Pseudemydurinae **80**
Pseudemydura 80
P. umbrina 80
Superfamily Pelomedusoidea **81**
Family Pelomedusidae **81**
Pelomedusa 81
P. barbata 81
P. galeata 82
P. gehafie 83
P. kobe 83
P. neumanni 84
P. olivacea 85
P. schweinfurthi 85
P. somalica 86
P. subrufa (sensu lato) 86
P. subrufa (sensu stricto) 87
P. variabilis 88
Pelusios 88
P. adansonii 89
P. bechuanicus 89
P. broadleyi 90
P. carinatus 90
P. castaneus 91
P. castanoides 92
P. c. castanoides 92
P. c. intergularis 93
P. chapini 93
P. cupulatta 94
P. gabonensis 94
P. marani 95
P. nanus 95
P. niger 96
P. rhodesianus 96
P. sinuatus 97

P. s. sinuatus ... 97
P. s. bottegi ... 98
P. subniger ... 98
P. s. subniger ... 99
P. s. parietalis ... 99
P. upembae ... 100
P. williamsi ... 100
P. w. williamsi ... 100
P. w. laurenti ... 101
P. w. lutescens ... 101
Family Podocnemididae ... **102**
Subfamily Erymnochelyinae ... **102**
Erymnochelys ... 102
E. madagascariensis ... 102
Subfamily Peltocephalinae ... **103**
Peltocephalus ... 103
P. dumerilianus ... 103
Subfamily Podocnemidinae ... **104**
Podocnemis ... 104
P. erythrocephala ... 104
P. expansa ... 105
P. lewyana ... 106
P. sextuberculata ... 106
P. unifilis ... 107
P. vogli ... 108
Suborder Cryptodira ... **109**
Infraorder Durocryptodira ... **109**
Superfamily Chelonioidea ... **109**
Family Cheloniidae ... **109**
Subfamily Carettinae ... **109**
Caretta ... 109
C. caretta ... 110
Eretmochelys ... 112
E. imbricata ... 112
Lepidochelys ... 113
L. kempii ... 114
L. olivacea ... 115
Subfamily Cheloniinae ... **116**
Chelonia ... 116
C. mydas ... 117
Natator ... 120
N. depressus ... 120
Family Dermochelyidae ... **120**
Dermochelys ... 120
D. coriacea ... 121
Superfamily Chelydroidea ... **123**
Family Chelydridae ... **123**
Chelydra ... 123
C. acutirostris ... 124
C. rossignonii ... 124
C. serpentina ... 125
Macrochelys ... 126
M. suwanniensis ... 126
M. temminckii ... 127
Family Dermatemydidae ... **127**
Dermatemys ... 128
D. mawii ... 128
Family Kinosternidae ... **129**
Subfamily Kinosterninae ... **129**
Kinosternon * ... 129
(*Kinosternon*) * ... 129
K. (K.) abaxillare ... 130
K. (K.) acutum ... 130
K. (K.) alamosae ... 131
K. (K.) albogulare * ... 131
K. (K.) chimalhuaca ... 132
K. (K.) cora ... 132
K. (K.) creaseri ... 133
K. (K.) cruentatum * ... 133
K. (K.) herrerai ... 134
K. (K.) hirtipes ... 134
K. (K.) h. hirtipes ... 135
K. (K.) h. chapalaense ... 135
K. (K.) h. magdalense ... 136
K. (K.) h. megacephalum † ... 136
K. (K.) h. murrayi ... 137
K. (K.) h. tarascense ... 137
K. (K.) integrum ... 138
K. (K.) mexicanum * ... 138
K. (K.) oaxacae ... 139
K. (K.) scorpioides * ... 139
K. (K.) sonoriense ... 141
K. (K.) s. sonoriense ... 141
K. (K.) s. longifemorale ... 142
K. (K.) vogti ... 142
(*Cryptochelys*) * ... 143
K. (C.) angustipons ... 143
K. (C.) dunni ... 143
K. (C.) leucostomum ... 144
K. (C.) l. leucostomum ... 144
K. (C.) l. postinguinale ... 145
(*Thyrosternum*) * ... 146
K. (T.) baurii ... 146
K. (T.) durangoense ... 147
K. (T.) flavescens ... 147
K. (T.) steindachneri ... 148
K. (T.) stejnegeri ... 148
K. (T.) subrubrum ... 149
K. (T.) s. subrubrum ... 149
K. (T.) s. hippocrepis ... 150
Sternotherus ... 151
S. carinatus ... 151

S. depressus 152
S. intermedius 152
S. minor 153
S. odoratus 153
S. peltifer 154
Subfamily Staurotypinae 155
Claudius 155
C. angustatus 155
Staurotypus 155
S. salvinii 156
S. triporcatus 156
Superfamily Testudinoidea 157
Family Emydidae 157
Subfamily Deirochelyinae 157
Chrysemys 157
C. dorsalis 157
C. picta 158
C. p. picta 158
C. p. bellii 159
C. p. marginata 160
Deirochelys 160
D. reticularia 160
D. r. reticularia 161
D. r. chrysea 161
D. r. miaria 162
Graptemys 162
G. barbouri 162
G. caglei 163
G. ernsti 163
G. flavimaculata 164
G. geographica 164
G. gibbonsi 165
G. nigrinoda 166
G. oculifera 166
G. ouachitensis 167
G. pearlensis 168
G. pseudogeographica 168
G. p. pseudogeographica 169
G. p. kohnii 169
G. pulchra 170
G. sabinensis 170
G. versa 171
Malaclemys 171
M. terrapin 171
M. t. terrapin 172
M. t. centrata 173
M. t. littoralis 173
M. t. macrospilota 174
M. t. pileata 174
M. t. rhizophorarum 175
M. t. tequesta 175
Pseudemys 176
P. alabamensis 176
P. concinna 176
P. c. concinna 177
P. c. suwanniensis 178
P. floridana 178
P. gorzugi 179
P. nelsoni 179
P. peninsularis 180
P. rubriventris 180
P. texana 181
Trachemys 181
T. adiutrix 182
T. callirostris 182
T. c. callirostris 183
T. c. chichiriviche 183
T. decorata 184
T. decussata 184
T. d. decussata 184
T. d. angusta 185
T. dorbigni 186
T. gaigeae 187
T. grayi 187
T. g. grayi 187
T. g. emolli 188
T. g. panamensis 189
T. hartwegi 189
T. medemi 190
T. nebulosa 190
T. n. nebulosa 191
T. n. hiltoni 191
T. ornata 192
T. scripta 192
T. s. scripta 193
T. s. elegans 194
T. s. troostii 195
T. stejnegeri 195
T. s. stejnegeri 195
T. s. malonei 196
T. s. vicina 196
T. taylori 197
T. terrapen 197
T. venusta 198
T. v. venusta 199
T. v. cataspila 199
T. v. iversoni 200
T. v. uhrigi 200
T. yaquia 201
Subfamily Emydinae 201
Actinemys 201
A. marmorata 201

A. pallida ... 202
Clemmys ... 203
C. guttata ... 203
Emydoidea ... 204
E. blandingii ... 204
Emys ... 205
E. orbicularis ... 205
E. o. orbicularis ... 205
E. o. eiselti ... 207
E. o. galloitalica ... 208
E. o. hellenica ... 208
E. o. ingauna ... 209
E. o. occidentalis ... 209
E. o. persica ... 210
E. o. ssp. indet. ... 210
E. trinacris ... 211
Glyptemys ... 211
G. insculpta ... 211
G. muhlenbergii ... 212
Terrapene ... 213
T. carolina ... 213
T. c. carolina ... 214
T. c. bauri ... 216
T. c. major ... 216
T. coahuila ... 217
T. mexicana ... 217
T. nelsoni ... 218
T. n. nelsoni ... 218
T. n. klauberi ... 219
T. ornata ... 219
T. triunguis ... 220
T. yucatana ... 221
Family Platysternidae ... **221**
Platysternon ... 221
P. megacephalum ... 221
P. m. megacephalum ... 222
P. m. peguense ... 222
P. m. shiui ... 223
Family Geoemydidae ... **223**
Subfamily Batagurinae ... **223**
Batagur ... 223
B. affinis ... 224
B. a. affinis ... 224
B. a. edwardmolli ... 225
B. baska ... 225
B. borneoensis ... 226
B. dhongoka ... 227
B. kachuga ... 228
B. trivittata ... 229
Geoclemys ... 230
G. hamiltonii ... 230
Hardella ... 231
H. thurjii ... 231
Malayemys ... 232
M. khoratensis ... 232
M. macrocephala ... 233
M. subtrijuga ... 233
Morenia ... 234
M. ocellata ... 234
M. petersi ... 235
Orlitia ... 235
O. borneensis ... 236
Pangshura ... 236
P. smithii ... 237
P. s. smithii ... 237
P. s. pallidipes ... 237
P. sylhetensis ... 238
P. tecta ... 238
P. tentoria ... 239
P. t. tentoria ... 239
P. t. circumdata ... 240
P. t. flaviventer ... 240
Siebenrockiella ... 241
S. crassicollis ... 241
S. leytensis ... 242
Subfamily Geoemydinae ... **242**
Cuora ... 242
(“*Cuora amboinensis*” Complex) * ... 242
C. amboinensis * ... 242
C. a. amboinensis * ... 243
C. a. aurantiae * ... 244
C. couro * ... 244
C. c. couro * ... 245
C. c. kamaroma * ... 245
C. lineata * ... 246
C. philippinensis * ... 246
C. praschagi * ... 247
(“*Cistoclemmys*” Complex) * ... 247
C. aurocapitata ... 247
C. a. aurocapitata ... 248
C. a. dabieshani ... 248
C. bourreti ... 249
C. cyclornata ... 249
C. c. cyclornata ... 250
C. c. annamitica ... 250
C. c. meieri ... 251
C. flavomarginata ... 251
C. f. flavomarginata ... 252
C. f. evelynae ... 252
C. galbinifrons ... 253
C. mccordi ... 254
C. mouhotii ... 254

C. m. mouhotii 254
C. m. obsti 255
C. pani 256
C. picturata 256
C. trifasciata 257
C. t. trifasciata 257
C. t. luteocephala 258
C. yunnanensis 258
C. zhoui 259
Cyclemys 259
C. atripons 259
C. dentata 260
C. enigmatica 261
C. fusca 261
C. gemeli 262
C. oldhamii 262
C. pulchristriata 263
Geoemyda 263
G. japonica 264
G. spengleri 264
Heosemys 265
H. annandalii 265
H. depressa 266
H. grandis 266
H. spinosa 267
Leucocephalon 268
L. yuwonoi 268
Mauremys 268
M. annamensis 269
M. caspica 270
M. japonica 271
M. leprosa 271
M. l. leprosa 272
M. l. saharica 273
M. mutica 274
M. m. mutica 274
M. m. kami 275
M. nigricans 275
M. reevesii 276
M. rivulata 277
M. sinensis 278
Melanochelys 279
M. tricarinata 279
M. trijuga 280
M. t. trijuga 280
M. t. coronata 281
M. t. edeniana 282
M. t. indopeninsularis 282
M. t. parkeri 283
M. t. thermalis 283
Notochelys 284
N. platynota 284
Sacalia 285
S. bealei 285
S. quadriocellata 285
Vijayachelys 286
V. silvatica 286
Subfamily Rhinoclemmyinae 287
Rhinoclemmys 287
R. annulata 287
R. areolata 288
R. diademata 288
R. funerea 289
R. melanosterna 289
R. nasuta 290
R. pulcherrima 291
R. p. pulcherrima 291
R. p. incisa 292
R. p. manni 292
R. p. rogerbarbouri 293
R. punctularia 293
R. p. punctularia 294
R. p. flammigera 295
R. rubida 296
R. r. rubida 296
R. r. perixantha 296
Family Testudinidae 297
Subfamily Manouriinae 297
Manouria 297
M. emys 297
M. e. emys 298
M. e. phayrei 298
M. impressa 299
Subfamily Xerobatinae 300
Gopherus 300
G. agassizii 300
G. berlandieri 301
G. evgoodei 302
G. flavomarginatus 302
G. morafkai 303
G. polyphemus 303
Gopherus, sp. indet. 304
Subfamily Testudininae 304
Aldabrachelys 304
A. gigantea 304
A. g. gigantea 305
A. g. arnoldi 306
A. g. daudinii † 306
A. g. hololissa 307
Astrochelys 308
A. radiata 308
A. yniphora 309

Centrochelys 309
C. sulcata 309
Chelonoidis 310
C. carbonarius 310
C. chilensis 312
C. denticulatus 312
C. niger 314
C. n. niger † 315
C. n. abingdonii † 316
C. n. becki 317
C. n. chathamensis 317
C. n. darwini 318
C. n. donfaustoi 318
C. n. duncanensis 319
C. n. guntheri 320
C. n. hoodensis 320
C. n. microphyes 321
C. n. phantasticus 321
C. n. porteri 322
C. n. vandenburghi 323
C. n. vicina 323
C. n. unnamed subsp. † 324
C. n. subsp. indet. 324
Chersina 324
C. angulata 324
Chersobius 325
C. boulengeri 325
C. signatus 326
C. solus 327
Cylindraspis † 327
C. indica † 327
C. inepta † 328
C. peltastes † 329
C. triserrata † 329
C. vosmaeri † 330
Geochelone * 331
G. elegans * 331
G. e. elegans * 332
G. e. actinodes * 333
G. platynota 333
Homopus 334
H. areolatus 334
H. femoralis 335
Indotestudo 335
I. elongata 335
I. forstenii 336
I. travancorica 337
Kinixys 337
K. belliana 338
K. erosa 338
K. homeana 339
K. lobatsiana 339
K. natalensis 340
K. nogueyi 340
K. spekii 341
K. zombensis 342
K. z. zombensis 342
K. z. domerguei 343
Malacochersus 343
M. tornieri 343
Psammobates 344
P. geometricus 344
P. oculifer 345
P. tentorius 346
P. t. tentorius 346
P. t. trimeni 347
P. t. verroxii 348
Pyxis 349
P. arachnoides 349
P. a. arachnoides 350
P. a. brygooi 350
P. a. oblonga 351
P. planicauda 351
Stigmochelys 352
S. pardalis 352
Testudo 353
(*Testudo*) 354
T. (T.) graeca 354
T. (T.) g. graeca 355
T. (T.) g. armeniaca 356
T. (T.) g. buxtoni 356
T. (T.) g. cyrenaica 357
T. (T.) g. ibera 358
T. (T.) g. marokkensis 359
T. (T.) g. nabeulensis 359
T. (T.) g. terrestris 360
T. (T.) g. tripolitania * 361
T. (T.) g. whitei 361
T. (T.) g. zarudnyi 362
T. (T.) kleinmanni 362
T. (T.) marginata 363
(*Agrionemys*) 364
T. (A.) horsfieldii 364
T. (A.) h. horsfieldii 365
T. (A.) h. bogdanovi 365
T. (A.) h. kazachstanica 366
T. (A.) h. kuznetzovi 366
T. (A.) h. rustamovi 367
(*Chersine*) 367
T. (C.) hermanni 367
T. (C.) h. hermanni 368
T. (C.) h. boettgeri 368

Infraorder Trionychia **369**
Superfamily Trionychoidea **369**
Family Carettochelyidae **369**
Carettochelys 369
C. insculpta 370
Family Trionychidae **370**
Subfamily Cyclanorbinae **370**
Cyclanorbis 370
C. elegans 371
C. senegalensis 371
Cycloderma 372
C. aubryi 372
C. frenatum 373
Lissemys 374
L. ceylonensis 374
L. punctata 374
L. p. punctata 375
L. p. andersoni 376
L. p. vittata 376
L. scutata 377
Subfamily Trionychinae **377**
Amyda 377
A. cartilaginea 378
A. c. cartilaginea 378
A. c. maculosa 379
A. ornata * 380
A. o. ornata 380
A. o. jongli * 381
A. o. phayrei 381
Apalone 382
A. ferox 382
A. mutica 383
A. m. mutica 384
A. m. calvata 384
A. spinifera 385
A. s. spinifera 385
A. s. aspera 386
A. s. atra 387
A. s. emoryi 388
A. s. guadalupensis 388
A. s. pallida 389
Chitra 389
C. chitra 389
C. c. chitra 390
C. c. javanensis 390
C. indica 391
C. vandijki 391
Dogania 392
D. subplana 392
Nilssonia 393
N. formosa 393
N. gangetica 394
N. hurum 395
N. leithii 396
N. nigricans 397
Palea 397
P. steindachneri 397
Pelochelys 398
P. bibroni 398
P. cantorii 398
P. signifera 399
Pelodiscus * 400
P. axenaria 400
P. huangshanensis 401
P. maackii 401
P. parviformis 402
P. shipian * 402
P. sinensis 403
P. variegatus 405
Rafetus 405
R. euphraticus 405
R. swinhoei 406
Trionyx 407
T. triunguis 408

Testudines, sp. indet. **410**

SUPPLEMENT
Extinct Holocene Turtle and Tortoise Taxa Extant from ca. 10,000 BCE to 1500 CE **411**

Order Testudines **411**
Suborder Meiolaniformes † **411**
Family Meiolaniidae † **411**
Meiolania † 411
M. mackayi † 411
Genus indet. † 412
Meiolaniidae sp. [New Caledonia] † 412
Suborder Cryptodira **412**
Infraorder Durocryptodira **412**
Superfamily Testudinoidea **412**
Family Testudinidae **412**
Subfamily Manouriinae **412**
Manouria 412
M. oyamai † 412
Subfamily Xerobatinae **413**
Hesperotestudo † 413
H. crassiscutata † 413
H. incisa † 414
H. turgida † 415
Subfamily Testudininae **416**
Aldabrachelys 416

A. abrupta † 416
A. grandidieri † 417
Astrochelys * 418
A. rogerbouri † * 418
Chelonoidis * 419
C. alburyorum † 419
C. a. alburyorum † 419
C. a. keegani † 420
C. a. sementis † 420
C. cubensis † 421
C. dominicensis † 421
C. gersoni † * 422
C. marcanoi † 422
C. monensis † 423
Solitudo † * 423
S. sicula † * 423
Genus indet. 424
?Meiolania damelipi † 424
Testudinidae sp. [Fiji] † 424

ANNOTATIONS
Current 2025 Checklist 425
Previous Checklist Annotations
2007 Annotations 442
2008 Annotations 446
2009 Annotations 447
2010 Annotations 450
2011 Annotations 453
2012 Annotations 455
2014 Annotations 459
2017 Annotations 464
2021 Annotations 474


DISTRIBUTION UPDATES
Distribution Updates 2011 490
Distribution Updates 2012 490
Distribution Updates 2014 491

LITERATURE CITED
1. General Bibliographic Citations 492
2. *Chelonian Research Monographs* 551
Conservation Biology of Freshwater Turtles and Tortoises (CBFTT) 551
CBFTT (TTWG, TEWG) Checklists 551
CBFTT Species Accounts 552
3. IUCN Red List and Green List Assessments ... 555

APPENDIX 1
Eponyms Used for Turtle and Tortoise Genera, Subgenera, Species, and Subspecies 564

APPENDIX 2
Institutional (Museum) Acronyms Used for Turtle and Tortoise Type Specimens 572

Turtle Taxonomy Working Group – 2025
Author Biographies 574



Pantanal Swamp Turtle,
Acanthochelys macrocephala (Chelidae), from
Filadelfia, Chaco, Paraguay;
photo by Thomas and Sabine Vinke.

European Pond Turtle,
Emys orbicularis orbicularis (Emydidae), from
Dnieper River Delta, Ukraine;
photo by Uwe Fritz.

Conservation Biology of Freshwater Turtles and Tortoises:
A Compilation Project of the IUCN SSC Tortoise and Freshwater Turtle Specialist Group
Rhodin, Iverson, van Dijk, Stanford, Goode, Buhlmann, and Mittermeier, Eds.
Chelonian Research Monographs (ISSN 1088-7105) No. 10, doi:10.3854/crm.10.checklist.atlas.v10.2025
© 2025 by Chelonian Research Foundation and Turtle Conservancy *• Published 18 July 2025*

Turtles of the World: Annotated Checklist and Atlas of Taxonomy, Synonymy, Distribution, and Conservation Status (10th Ed.)

Turtle Taxonomy Working Group*

Anders G.J. Rhodin[1,2], John B. Iverson[3,4], Uwe Fritz[5], Natalia Gallego-García[6], Arthur Georges[7], H. Bradley Shaffer[8], and Peter Paul van Dijk[1,2]

[1]Chelonian Research Foundation, 564 Chittenden Dr., Arlington, Vermont 05250 USA [rhodincrf@aol.com];
[2]Turtle Conservancy, 1794 McNell Rd., Ojai, California 93023 USA [peterpaul@turtleconservancy.org];
[3]Department of Biology, Earlham College, Richmond, Indiana 47374 USA [johni@earlham.edu];
[4]Turtle Survival Alliance, Director and Field Conservation Committee, North Charleston, South Carolina 29406 USA;
[5]Museum für Tierkunde, Senckenberg Dresden, A.B. Meyer Building, 01109 Dresden, Germany [uwe.fritz@senckenberg.de];
[6]Turtle Survival Alliance, North Charleston, South Carolina 29406 USA [ngallego@turtlesurvival.org];
[7]Institute for Applied Ecology, University of Canberra, ACT 2601, Australia [arthur.georges@canberra.edu.au];
[8]Department of Ecology and Evolutionary Biology and La Kretz Center for California Conservation Science, Institute of the Environment and Sustainability, University of California, Los Angeles, California 90095 USA [brad.shaffer@ucla.edu]

**Authorship of this article is by this working group of the IUCN SSC Tortoise and Freshwater Turtle Specialist Group, which for the purposes of this document consisted of the following contributors*:

Abstract. – This is our 10th edition of an annotated checklist and atlas of all recognized taxa of the world's modern turtle and tortoise fauna, documenting recent changes and controversies through mid-2025, and including all primary synonyms, updated from nine previous checklists. We provide an updated comprehensive listing of taxonomy and nomenclature, including type localities, type specimens, detailed distribution maps, as well as calculated Estimated Indigenous (Historical) Ranges, conservation status, and maximum known sex-based carapace lengths for all taxa. We strive to record the most recent justified taxonomic assignment of taxa in a hierarchical framework, providing detailed annotations, including alternative arrangements for a few taxa. We include current published and provisional IUCN Red List status assessments for all species, as well as current listings on CITES appendices and the USA Endangered Species Act. The diversity of turtles and tortoises in the world that has existed in modern times (since 1500 CE) and currently generally recognized as distinct and included in this checklist, now consists of 364 species. Of these, 60 are polytypic, representing 129 additional recognized subspecies (one unnamed), or 493 total taxa of modern chelonians, a 1.4% increase from 486 taxa in our previous checklist. Of these, 5 species and 5 subspecies (one unnamed), or 10 taxa (2.0%), are extinct. We also include a supplementary listing of 20 taxa of terrestrial chelonians that went extinct during the Holocene from ca. 10,000 BCE (ca. 12,000 ybp) to 1500 CE. As of the current IUCN 2025 Red List, 174 turtle species (62.8% of the 277 species formally Red-Listed) are officially regarded as globally Threatened (Critically Endangered [CR], Endangered [EN], or Vulnerable [VU]). We also record additional "*in press*" and other provisional Red List assessments by the IUCN SSC Tortoise and Freshwater Turtle Specialist Group, allowing us to evaluate the overall current threat levels for all 364 species of turtles and tortoises. Of these, 196 (53.8%) are Threatened (CR, EN, or VU), a 7.1% increase from 183 in 2021; if we provisionally adjust for predicted threat rates of 29 Data Deficient (DD) species, then ca. 58.5% of all extant turtles are Threatened, up from 55.9% in 2021. These numbers and percentages of Threatened species have all increased since our last checklist. Turtles and tortoises are among the most threatened of the major vertebrate lineages on earth.

Key Words. – Reptilia, Testudines, turtle, tortoise, terrapin, chelonian, taxonomy, nomenclature, systematics, distribution, maximum size, sexual dimorphism, conservation status, etymology, type localities, type specimens, IUCN Red List, CITES, ESA, extinction, Holocene

INTRODUCTION

The global diversity of turtles and tortoises (chelonians, order Testudines) that has existed in modern times (since 1500 CE [=AD]), and currently generally recognized as described and distinct by specialists in turtle taxonomy and systematics, consists of 364 species, of which 60 are polytypic, with 129 additional recognized subspecies, or 493 total taxa of modern (living and extinct) turtles and tortoises (including one unnamed subspecies). Of these, 7 are marine turtle species, leaving us with 357 species and 486 taxa of terrestrial tortoises and freshwater turtles. Of the 493 total taxa, 5 species plus 5 subspecies (one unnamed), or 10 total taxa of tortoises and freshwater turtles (2.0% of all modern taxa) have gone extinct since 1500 CE (see Table 1 and Fig. 1), leaving us currently with 359 species and 124 additional subspecies, or 483 total taxa of living turtles and tortoises, of which 352 species and 476 taxa are freshwater or terrestrial.

In this checklist we present a full taxonomic listing of all recognized modern turtle and tortoise taxa, including synonymized names and type localities, detailed distribution maps, and annotations concerning recently described new taxa, nomenclatural and taxonomic updates, and significant taxon-related controversies or developments. To make the checklist more useful to a broader audience of biologists and managers, we also include the current conservation status and maximum carapace lengths for all taxa.

The 493 modern turtle and tortoise taxa we recognize here are based on a synonymy of 1,520 separate named turtle and tortoise species and subspecies, including all primary description names, secondary *nomen novum* replacement names, undescribed *nomen nudum* names, and several other nomenclaturally unavailable names. These names also include those fossil taxa that have been synonymized with modern taxa.

Within the monophyletic order Testudines (turtles and tortoises) we recognize 2 suborders, 2 infraorders, 6 superfamilies, 14 families, 22 subfamilies, 97 genera, and 12 subgenera of modern turtles, for a total of 156 supraspecific groupings. These groups are based on 519 valid and synonymized names, for a total listing here of 2,038 taxonomic names applied to all modern turtle taxa and groups.

As there is always some disagreement among experts as to which taxa are distinct and valid, and at what systematic level or rank (species or subspecies, genus or subgenus), these numbers are variable depending on the authorities presenting their data or interpretations. For prior discussions and listings of all recognized modern turtle taxa, with extensive annotations regarding areas of recent taxonomic change, instability, or controversy, see the previous publications by the Turtle Taxonomy Working Group (TTWG 2007a,b, 2009, 2010, 2011, 2012, 2014, 2017, 2021), Rhodin et al. (2008), and the turtle checklist produced for CITES by Fritz and Havaš (2007). All annotations from previous TTWG checklists are included in the back of the present checklist.

For a listing of extinct Pleistocene and Holocene turtle and tortoise taxa, see our companion checklist by the Turtle Extinctions Working Group (TEWG 2015). We also include an updated Supplement for extinct Holocene taxa at the end of this checklist, where we list 18 species or 20 taxa (including two unnamed taxa) from 2 families. As such, the total number of turtle and tortoise taxa that have existed from the beginning of the Holocene through modern times is 382 species or 513 taxa.

Methodology

The Turtle Taxonomy Working Group (TTWG), created in 2005, functions under the auspices of the IUCN SSC Tortoise and Freshwater Turtle Specialist Group (TFTSG), which operates under the umbrella of the IUCN (International Union for Conservation of Nature) and its Species Survival Commission (SSC). We first compiled our checklist of modern turtle taxa in 2007 (TTWG 2007b), and have updated it frequently to reflect more recent changes, as required by subsequent publications with taxonomic novelties or proposed changes, as well as adding primary synonyms for all recognized taxa, type species and type locality designations, and distribution maps (Rhodin et al. 2008; TTWG 2009, 2010, 2011, 2012, 2014, 2017, 2021). The previous and current checklists took a longer time to update and produce because of extensive further expansion in content. This is now the 10th installment in this series. It is current through approximately May 2025.

We list all primary and synonymized description names, as well as all *nomen nudum*, *nomen dubium*, *nomen oblitum*, *nomen novum* names and various unavailable, suppressed, or rejected names of which we are aware. In general, we exclude obvious *ex errore* and typographical *lapsus* names, especially the profusion of inadvertent misspellings in modern literature (notably in the popular literature and other non-taxonomic biological sciences).

Our listing of alternative *nomen novum* names takes a broadly encompassing approach and lists both justified and unjustified subsequent emendations and replacements, including substantial name changes caused by early authors' occasional tendencies to create new or "better" names that they felt were more appropriate or more correct. Occasionally, early authors appear not to have remembered what the previously used names were, or simply came up with new spelling variations, with these new names sometimes becoming temporarily established in the literature. This was especially true for the many names and unjustified spelling emendations created and recorded by Gray between 1825 and 1874. Some of the alternative names may have simply been

Table 1. Modern freshwater turtles and tortoises that are recognized to have gone extinct since 1500 CE (one genus, five species, and five subspecies = 10 taxa), with approximate or known extinction dates. For species that apparently went extinct during the Holocene from ca. 10,000 BCE to 1500 CE, see the Supplement at the end of this publication. For species that apparently went extinct during the Pleistocene, see TEWG (2015).

Kinosternidae
- ***Kinosternon hirtipes megacephalum***
  - Viesca Mud Turtle
  - Mexico (Coahuila); ca. 1970

Testudinidae
- ***Aldabrachelys gigantea daudinii***
  - Daudin's Giant Tortoise
  - Seychelles (Mahé?); ca. 1850
- ***Chelonoidis niger abingdonii***
  - Pinta Giant Tortoise, Abingdon Island Giant Tortoise
  - Ecuador (Galápagos: Pinta [Abingdon]); 24 June 2012
- ***Chelonoidis niger niger***
  - Floreana Giant Tortoise, Charles Island Giant Tortoise
  - Ecuador (Galápagos: Floreana [Charles]); ca. 1850
- ***Chelonoidis niger* unnamed subsp.**
  - Santa Fé Giant Tortoise, Barrington Island Giant Tortoise
  - Ecuador (Galápagos: Santa Fé [Barrington]); ca. 1890 [?]
- ***Cylindraspis***
  - Mascarene Giant Tortoises
  - Indian Ocean Islands; ca. 1840
- ***Cylindraspis indica***
  - Reunion Giant Tortoise
  - Réunion; ca. 1840
- ***Cylindraspis inepta***
  - Mauritius Giant Domed Tortoise
  - Mauritius (Mauritius); ca. 1735
- ***Cylindraspis peltastes***
  - Rodrigues Domed Tortoise
  - Mauritius (Rodrigues); ca. 1800
- ***Cylindraspis triserrata***
  - Mauritius Giant Flat-shelled Tortoise
  - Mauritius (Mauritius); ca. 1735
- ***Cylindraspis vosmaeri***
  - Rodrigues Giant Saddleback Tortoise
  - Mauritius (Rodrigues); ca. 1800




Figure 1. While beyond the time-frame of extinctions for modern turtles, it is worth noting the findings of White et al. (2010), who documented the continued existence into the late Holocene of a giant terrestrial chelonian on Efate Island, Vanuatu, in the southern Pacific Ocean. The species persisted until as recently as 940–810 BCE (see TEWG 2015). White et al. (2010) tentatively assigned it to the extinct terrestrial horned turtle family Meiolaniidae, and named it ?*Meiolania damelipi*, and provided clear evidence of human butchering and consumption of the species, further corroborated and expanded by Hawkins et al. (2016). This exploitation represented the final anthropogenic extinction event for these southern Pacific giant terrestrial chelonians. For a more complete analysis of turtle extinctions caused by humans and/or climate change during the Pleistocene and Holocene, see our companion publication by the TEWG (2015) and the Supplement on extinct Holocene turtles at the end of this book. More recent work by Rabi et al. (2019) has re-assigned this species to the Testudinidae, but its presence on Vanuatu is still extraordinary. No complete skeletons of ?*M. damelipi* are known, and it has not yet been assigned to a defined genus of Testudinidae. The ecologically similar *Meiolania platyceps* from Lord Howe Island, Australia, which went extinct in the Late Pleistocene, has been beautifully reconstructed (photo above) by the American Museum of Natural History (Burke et al. 1983).

misinterpretations of handwritten script that led to typesetting errors.

However, prior to the establishment of the International Commission on Zoological Nomenclature in 1895, and the publication of the first edition of the Code of Zoological Nomenclature in 1905, these kinds of new names and changes were fairly common and we do not generally consider them to be simple errors; therefore we have instead recorded many of them as *nomen novum* names. Many early names were also unjustifiably emended in an attempt to comply with perceived rules about word constructions and the use of non-Greek vs. Greek letters (e.g., *c* vs. *k*, as in *Cinosternon* vs. *Kinosternon*, *Cinixys* vs. *Kinixys*). In addition, we do not list variations in spelling of the two alternate eponymic endings (*-ii* vs. *-i*), always using the original valid orthography.

Furthermore, in accord with Article 13 of the ICZN Code regarding availability of new names proposed after 1930, unless they were expressly created as justified *nomen novum* names, alternative names appearing post-1930 are regarded as invalid and unavailable. We list a few of these misspelled or unjustified new names that have appeared in the major turtle taxonomic literature and label them as *nomen invalidum* names.

We include listings of subsequent new combination names to reflect how taxa have been rearranged into new genera or different specific or subspecific levels. The new combination names are listed in *lighter gray text* following each associated primary name, arranged more or less chronologically from oldest to most recently created combinations, but without attributing authorship or date of first use of the new combination. We have attempted to list all known subsequent combination names, but these

listings may be incomplete. A few older *ex errore* misspelled names are included in these listings.

Original and synonymized taxon names (including higher-category names) are listed using their original spelling and genus-species combination as used by the author at the time of first publication of the name. Our synonymies for genus- and species-level taxa follow, to our best efforts, the strict and established nomenclatural rules established by the fourth edition of the International Code of Zoological Nomenclature (ICZN 1999).

However, for the higher-level suprageneric categories used in this checklist, we have also provided some synonyms and previously-used names for the same or included groupings whose usage may not necessarily correspond to nomenclatural guidelines under the ICZN. Since the ICZN does not regulate names above the superfamily rank, our listings of these names are intended to document historical use to aid understanding and resolving the difficult questions of what names are most appropriately used for these suprageneric categories and to what author they should be attributed.

For example, the names we list under the Order-level name for turtles (Testudines) are not all strict synonyms, as some were proposed at different levels of groupings, from "Family" to Order to various supraordinal categories. Many were utilized primarily for including various fossil turtle-like ancestors in an expanded concept of turtles, including some rank-free PhyloCode names (see Joyce et al. 2021). The names we list in other infraordinal suprageneric categories are not always strict synonyms either, as based on nomenclatural acceptability or availability of the utilized group name, but instead provide a partial historical record for names previously used for the same or similar grouping.

Our main checklist includes all currently recognized named taxa (species and subspecies) of modern turtles (extant after 1500 CE). By "currently recognized" we mean those taxa that have not been demonstrably refuted or justifiably synonymized in published literature, or whose description or recommended resurrection has yet to receive wide community acceptance. We have attempted to describe all recent published taxonomic recommendations in our annotations, even though we have not included all proposed changes in the checklist.

Since there are sometimes also different interpretations for some genera and polytypic species as to which names are valid and whether to lump or split the contained taxa, we occasionally list alternative name usages. Our most important criterion for accepting proposed changes is that they be accompanied by adequate data and sound arguments justifying the taxonomic revision. Consequently, some proposed taxonomic changes from lists or publications with unsupported or untested revisions have not been incorporated. We also list and annotate recent systematic papers that do not necessarily commit nomenclatural or taxonomic acts, but that present data on phylogenetic or phylogeographic relationships that either serve to further support currently understood relationships, or are indicators of distinct lineages or potential taxonomic changes to come. Occasionally we also include annotations concerning dates of publication or other bibliographic considerations.

Currently recognized modern turtle and tortoise genera and terminal taxa (species and subspecies) are listed in bold italics. Original description names and synonymized names are in non-bold italicized text. Higher suprageneric group-level names are listed in bold non-italic text and are presented in an indented phylogenetic hierarchy. For competing alternative generic names, we list them in phylogenetic order from most to least inclusive. All original names include authorship, year, and page number.

All described genera include type species designations (original and subsequent), and all described species include verbatim original and subsequently restricted or emended type localities. This has been undertaken through a comprehensive re-examination of all original literature rather than relying on secondary sources. As a result, many of these type designations and localities are somewhat different from those previously cited in Fritz and Havaš (2007) or elsewhere.

In order for readers to easily identify taxon entry fields with new or revised data since the previous checklist (TTWG 2021), we indicate such updated entries with a single asterisk (*) after the heading for the pertinent entry field(s): Distribution, Introduced, Estimated Indigenous Ranges (AOO and EOO), Size (Max SCL), CBFTT Account, IUCN Red List, TFTSG Provisional Red List, CITES, USA ESA, Type Locality, Type Specimen, Etymology, or Comment. New taxa added or resurrected in the current checklist are indicated by an asterisk (*) after the Synonymy heading.

More extensive comments on names that have undergone recent taxonomic change or phylogenetic analysis or are associated with instability or uncertainty or other changes are indicated by superscript numbers that refer to annotations at the end of this and earlier checklists. See the section on **Annotations** at the end of the checklist for all detailed explanations.

Summaries of all major taxonomic changes in this checklist as compared to our previous one are provided in Tables 2–3; all minor changes are only included in the annotations. Table 4 lists those recently described or resurrected taxa that we have not accepted as distinct.

Turtle taxa that were originally described based on Pleistocene or Holocene fossil, subfossil, or archeological material, but subsequently recognized as representing extant taxa or synonymized with modern turtle taxa, are included in the checklist and marked with a cross (†), and include stratigraphic horizon and location data. Fossil taxa synonymized with extant polytypic species are listed under the geographically most appropriate subspecies; however, such synonymizations may not

be accurate for some Pleistocene or older fossils which could conceivably represent distinct chronospecies or extinct subspecies. For further details on extinct fossil turtle and tortoise taxa from the Pleistocene and Holocene, see TEWG (2015).

Starting with the previous checklist (TTWG 2021), we also include a **Supplement** listing the turtle and tortoise species and taxa that went extinct during the Holocene epoch, from about 10,000 BCE [=BC] to 1500 CE [=AD]. At present, we recognize 20 such extinct Holocene taxa.

Those modern species and subspecies for which in-depth informational accounts have been published in this TFTSG monograph series on *Conservation Biology of Freshwater Turtles and Tortoises* (CBFTT), are indicated by a **CBFTT Account** heading, with interactive hyperlinks provided to the online published accounts. New CBFTT accounts added since TTWG (2021) are marked with an asterisk (*). The CBFTT series will gradually hopefully publish accounts for all non-marine turtle and tortoise species—accounts also include recognized subspecies within the account, but some subspecies have separate accounts and are so indicated. As of May 2025, CBFTT has published 128 accounts by 267 authors covering 173 turtle and tortoise taxa; these are all available online as downloadable open-access doi-designated pdf's on the TFTSG website at **www.iucn-tftsg.org/cbftt/**.

The checklist includes English common names for all recognized taxa, as well as for all recognized genera and suprageneric categories. We have tried to provide the most widely used names, although occasionally we have provided two or more names. We do not support the practice of designating "official" or "standard" common names, as that is the domain for scientific names. Instead, common names tend to evolve and vary from area to area and over time, as well as with language and cultural context. However, in the field of conservation, the use of reasonably widely recognized and appropriately descriptive common names is critically important for communication purposes, and so we include English common names, as well as some of the more common native language names used in countries where the species occurs. Though also important for the global conservation community, and officially included in IUCN Red List and CITES documentation, we do not at this time include non-native Spanish or French common names in this checklist. Indigenous vernacular names for certain species are often extensive and imprecise, and in general we do not list such names here, although a few commonly-used ones are listed.

Body Size (Carapace Length) and Sexual Size Dimorphism

We introduced sex-based maximum body size (carapace length) data for all turtle taxa in TTWG (2021), with separate values for males and females for most taxa. We record primarily maximum straight-line carapace length (Max SCL; see Iverson and Lewis 2018), based on cited sources, but also maximum curved carapace length (Max CCL) for some larger species in which SCL is hard to measure. Revised or new maximum sizes in the present checklist are marked with an asterisk (*).

Most authors who report SCL do not indicate whether the measurement is midline or maximum (which is greater in species with extended anterior marginals with an anterior midline notch or recess and/or serrated posterior marginals with a supracaudal notch), but when they do, we record the maximum value. Many undefined SCL measurements in the literature are probably midline and therefore likely a bit smaller than the maximum carapace length. We strongly urge all authors to standardize and accurately describe their methods of measuring body size (Iverson and Lewis 2018).

We do not record adult size means or ranges, nor calculate single-taxon sexual size dimorphism (SSD) ratios or indices (see Lovich and Gibbons 1992), as all these values are typically population-dependent, usually vary geographically, latitudinally, and altitudinally, are affected by sample sizes and potentially biased collection methodology, and are adversely impacted by long-term exploitation levels and other environmental and anthropogenic pressures and threats. Additionally, SSD indices based on single maximum values do not reflect population-level dynamics as accurately as the evaluation of body size means determined via large unbiased sample sizes.

The various ecological, geographic, environmental, phylogenetic, evolutionary, behavioral, and theoretical aspects of body size and sexual size dimorphism in turtles and tortoises have been extensively analyzed and discussed by, among others, Berry and Shine (1980), Gibbons and Lovich (1990), Bonnet et al. (2001, 2010), Lindeman (2003), Ben Kaddour et al. (2008), Gosnell et al. (2009), Stephens and Wiens (2009), Lovich et al. (2010, 2014a), Jaffe et al. (2011), Ceballos et al. (2013), Halámková et al. (2013), Ceballos and Iverson (2014), Itescu et al. (2014), Angielczyk et al. (2015), Tesche and Hodges (2015), Regis and Meik (2017), Agha et al. (2018), Rodrigues et al. (2018), Santilli and Rollinson (2018), Vlachos and Rabi (2018), Figgener et al. (2022), Joos et al. (2022), and Farina et al. (2023). The nearly complete sex-based maximum size data recorded here for most turtles and tortoises should help provide improved data for further analyses.

We note that one of the attributes of these data is that for exploited species, the historical maximum sizes, which we record here, may seldom be reached in affected populations, and that this is usually an indication of exploitation pressures on the largest specimens (e.g., Close and Seigel 1997; Boundy and Kennedy 2006; Wolak et al. 2010; Eisemberg et al. 2011; among others). We also note that there are occasional individuals

Table 2. Summary of new or resurrected taxa (*) recognized in this checklist with major proposed taxonomic changes from TTWG 2021. See the annotations for a full discussion of all these changes; minor changes associated only with overlooked or previously synonymized names or dates of authorship or other primarily nomenclatural changes are not listed here, but only in the annotations. This table does not include added synonymized fossil taxa, *nomina nuda*, or names not considered valid in the 2021 checklist (i.e., newly added synonyms).

TTWG 2021 (357 species, 486 taxa)	**TTWG 2025** (364 species, 493 taxa)	Change
Chelodina (Chelydera) kuchlingi + *Chelodina (Chelydera) walloyarrina*	*Chelodina (Chelydera) kuchlingi*	two species synonymized into one
Emydura victoriae	*Emydura australis* *	older species name resurrected
Kinosternon	(*Kinosternon*) + (*Cryptochelys*) + (*Thyrosternum*)	genus split into 3 subgenera
Kinosternon scorpioides scorpioides	*Kinosternon scorpioides*	subspecies elevated to species
Kinosternon scorpioides albogulare	*Kinosternon albogulare*	subspecies elevated to species
Kinosternon scorpioides cruentatum	*Kinosternon cruentatum* + *Kinosternon mexicanum* *	taxon split, 1 taxon elevated to species, 1 species resurrected
Cuora amboinensis couro	*Cuora couro couro*	subspecies allocated to different species
Cuora amboinensis kamaroma	*Cuora couro kamaroma*	subspecies allocated to different species
Cuora amboinensis lineata	*Cuora lineata*	subspecies elevated to species
Cuora amboinensis	*Cuora amboinensis aurantiae* *	taxon split, 1 new subspecies described
Cuora amboinensis	*Cuora philippinensis* *	taxon split, 1 new species described
Cuora amboinensis	*Cuora praschagi* *	taxon split, 1 new species described
Geochelone elegans	*Geochelone elegans elegans* + *G. elegans actinodes* *	taxon split, 1 subspecies resurrected
Testudo (Testudo) graeca nabeulensis	*T. (T.) graeca nabeulensis* + *T. (T.) graeca tripolitania* *	taxon split, 1 new subspecies described
Amyda ornata	*Amyda ornata jongli* *	1 new subspecies described
Pelodiscus axenaria	*Pelodiscus axenaria* + *Pelodiscus shipian* *	taxon split, 1 new species described

that exceed typical maximal sizes in the usual range; however, we choose to include data from such record-size individuals, as they are always of interest.

We provide sources for all our size data, with most values accessed from primary references cited either in our CBFTT review accounts or in compilations by others, including Pritchard (2001), Ernst and Lovich (2009), Hansen et al. (2010), Legler and Vogt (2013), Ceballos et al. (2013), Itescu et al. (2014), and Cann and Sadlier (2017). Many are from various other primary cited sources, including some from IUCN Red List assessments; and in general, we strive to provide primary sources for data as much as possible.

Many accounts in the literature provide only generalized or approximate maximum carapace lengths; we attempt to record only those that are based on actual measurements of documented specimens. Generalized records of large animals may also be suspect unless specific measurement methodology is noted, as they often represent rounded-off estimates of SCL, or measurements of curved carapace length (CCL), or even total body length. Many of the maximum sizes provided in Itescu et al. (2014) were from Ernst et al. (2006b) and Bonin et el. (2006), but many of those were generalized size approximations and were not specifically referenced, especially those in Bonin et al. (2006), many of which appear to be overestimated. They are therefore only included here if more precise size data are not readily available. Many of the sizes we record do not agree with numbers previously cited in the various reviews noted above, but we have verified ours by checking primary sources. Some sizes provided in some reviews appear to be rounded up or erroneously interpreted or inaccurately copied from the primary sources.

For some species, notably giant tortoises, large sea turtles, large softshells, and a few others, we also provide maximum CCL measurements. For large species such as these, SCL is difficult to measure in the field, so CCL measurements are often recommended and reported (see Bolten 1999 for sea turtles). For species with only CCL measurements available, we provide a few estimated SCL values. For some polytypic species, we only have data available for the whole species, not for the included subspecies. Whenever possible, we have incorporated previously unpublished data on maximum sizes from our own measurements of museum and live specimens, as well as unpublished measurements received from colleagues, all of which are recorded and cited where appropriate.

Table 3. Summary of new or resurrected extinct Holocene taxa (*) included in the Supplement to this checklist, with taxonomic changes from TTWG 2021.

TTWG 2021 (17 taxa)	**TTWG 2025** (20 taxa)	Change
———	*Astrochelys rogerbouri* *	1 new species described
Chelonoidis marcanoi	*Chelonoidis marcanoi* + *Chelonoidis gersoni* *	taxon split, 1 new species described
———	*Solitudo* *	1 new genus described
———	*Solitudo sicula* *	1 new species described

Table 4. Summary of new or resurrected taxa proposed since TTWG 2021 that have **not** been recognized as distinct in this checklist, but instead synonymized with currently accepted taxa. See Annotations for discussions of these decisions.

Proposed Taxon	Synonymized Under
Mesoclemmys jurutiensis	*Mesoclemmys raniceps*
Mesoclemmys sabiniparaensis	*Mesoclemmys vanderhaegei*
Chelodina (Chelodina) ipudinapi	*Chelodina (Chelodina) novaeguineae*
Elseya (Elseya) kalumburu	*Elseya (Elseya) dentata*
Elseya (Elseya) sahul	*Elseya (Elseya) dentata*
Elseya (Hanwarachelys) caelatus berau	*Elseya (Hanwarachelys) novaeguineae*
Elseya (Hanwarachelys) freweri	*Elseya (Hanwarachelys) rhodini*
Elseya (Hanwarachelys) nabire	*Elseya (Hanwarachelys) schultzei*
Elseya (Hanwarachelys) papua	*Elseya (Hanwarachelys) rhodini*
Elseya (Hanwarachelys) timika timika	*Elseya (Hanwarachelys) rhodini*
Elseya (Hanwarachelys) timika aru	*Elseya (Hanwarachelys) rhodini*
Elseya (Hanwarachelys) timika wamena	*Elseya (Hanwarachelys) rhodini*
Elseya (Hanwarachelys) timika paniai	*Elseya (Hanwarachelys) rhodini*
Elseya (Pelocomastes) gondwana	*Elseya (Pelocomastes) lavarackorum*
Elseya (Pelocomastes) oneiros	*Elseya (Pelocomastes) lavarackorum*
Elseya (Pelocomastes) irwini stirlingi	*Elseya (Pelocomastes) irwini*
Elseya (Solomonemys)	*Elseya (Hanwarachelys)*
Elseya (Solomonemys) eidolon	*Elseya (Hanwarachelys) schultzei*
Elseya (Solomonemys) auramemoria	*Elseya (Hanwarachelys) schultzei*
Pelodiscus huike	*Pelodiscus axenaria*
Pelodiscus jiangxiensis	*Pelodiscus shipian*

We provide maximum shell size for all 364 recognized species and 485 (98.4%) of the 493 recognized taxa of the world's modern turtles. The average turtle species has a maximum SCL of 35.6 cm and the average turtle taxon has a maximum SCL of 35.5 cm (slightly smaller since many polytypic species are relatively smaller than monotypic species, with the giant tortoises of the Galápagos and Indian Ocean being notable exceptions).

We rank the 50 largest modern turtle and tortoise taxa by straight-line carapace length (SCL) in Table 5. The Leatherback, *Dermochelys coriacea*, is the world's largest turtle and largest marine species at ca. 203.2 cm SCL (213.4 cm CCL). The Siamese Narrow-headed Softshell Turtle, *Chitra chitra chitra*, is the largest freshwater species at 140.0 cm, and the Seychelles Giant Tortoise, *Aldabrachelys gigantea hololissa*, the largest terrestrial species at 138.0 cm. Several of the largest taxa (8 of 50) have gone extinct, a typical pattern for reptiles and vertebrates in general (see Ripple et al. 2017).

We note, interestingly, that other than all 7 marine turtles (Dermochelyidae and Cheloniidae) among these 50 largest chelonians, 38 of the remaining 43 taxa are either tortoises (Testudinidae, 25 taxa) or softshell turtles (Trionychidae, 13 taxa), with only 2 Chelydridae, 2 Geoemydidae, and a single Podocnemididae. Among the other chelonian families, the largest (and only) Dermatemydidae is *Dermatemys mawii* at 60.0 cm, the largest Pelomedusidae is *Pelusios sinuatus bottegi* at 55.0 cm, the largest Chelidae is *Chelus orinocensis* at 52.6 cm, the largest (and only) Carettochelyidae is *Carettochelys insculpta* at 52.5 cm, the largest Emydidae is *Trachemys grayi emolli* at 54.8 cm, the largest Kinosternidae is *Staurotypus triporcatus* at 40.2 cm, and the largest Platysternidae is *Platysternon megacephalum peguense* at 25.5 cm SCL.

The largest known turtle that ever lived was *Stupendemys geographicus* Wood 1976 (Podocnemididae), a freshwater or estuarine turtle from the Miocene of northern South America closely related to extant *Peltocephalus dumerilianus*. It has been recorded as reaching an impressive SCL of 240 cm (Cadena et al. 2020), but not much larger than the largest Leatherback.

We rank the 50 smallest modern taxa by SCL in Table 6. The recently described Vallarta Mud Turtle, *Kinosternon vogti*, appears to be the world's smallest turtle and smallest freshwater species at 10.2 cm SCL, but the known sample size is still very small, and the other recently described species, *Pelodiscus shipian*, is just about as small at 10.3 cm. The South African tortoises, *Chersobius signatus* and *C. boulengeri*, are the smallest terrestrial species at 11.0 cm each, and the Kemp's Ridley, *Lepidochelys kempii*, is the smallest marine species at 76.0 cm. Only one of the 50 smallest taxa has gone extinct.

We also note that among the 50 smallest chelonians, 32 of the taxa are either mud turtles (Kinosternidae, 20 taxa) or tortoises (Testudinidae, 12 taxa), with 7 Emydidae, 4 Geoemydidae, 3 Pelomedusidae, 2 Trionychidae, and one each of Platysternidae and Chelidae. Among the other polytypic chelonian families, the smallest Podocnemididae is *Podocnemis erythrocephala* at 32.2 cm, the smallest Chelydridae is *Chelydra acutirostris* at 48.0 cm, and the smallest Cheloniidae is *Lepidochelys kempii* at 76.0 cm SCL.

Although we do not record sexual size dimorphism (SSD) index values for individual species in the checklist, for the reasons provided above, we calculated them for the 327 species (89.8% of 364) and 440 taxa (89.2% of 493) for which we have sex-apportioned data, using the method recommended by Lovich and Gibbons (1992). We found that, overall, based on maximum recorded SCL, turtle species have an average female-biased SSD index of 0.173, and turtle taxa have a slightly lower average female-biased SSD index of 0.172.

Based on the maximum SCL data we provide, the Angulate Tortoise, *Chersina angulata*, may be the most male-biased species, at an SSD of -0.625, with the Suwannee Alligator Snapping Turtle, *Macrochelys suwanniensis*, second-most at -0.520, and the African Spurred Tortoise, *Centrochelys sulcata*, the third-most at -0.488. However, incomplete data suggest that the extinct Reunion Giant Tortoise, *Cylindraspis indica*, may have been the most male-biased species at -1.340.

The Crowned River Turtle, *Hardella thurjii*, is apparently the most female-biased species at 2.185, with the Pink-ringed Tent Turtle, *Pangshura tentoria*

Table 5. Top 50 largest modern turtle taxa by maximum straight-line carapace length (SCL, cm) of the larger sex; † = extinct.

Taxon	Family	SCL
Dermochelys coriacea	Dermochelyidae	203.2
Chelonia mydas	Cheloniidae	141.0
Chitra chitra chitra	Trionychidae	140.0
Aldabrachelys gigantea hololissa	Testudinidae	138.0
Chelonoidis niger porteri	Testudinidae	135.8
Chelonoidis niger vandenburghi	Testudinidae	129.5
Chitra chitra javanensis	Trionychidae	129.0
Caretta caretta	Cheloniidae	125.0
Aldabrachelys gigantea gigantea	Testudinidae	123.0
Cylindraspis indica †	Testudinidae	117.0
Chelonoidis niger donfaustoi	Testudinidae	112.0
Rafetus swinhoei	Trionychidae	109.5
Cylindraspis vosmaeri †	Testudinidae	109.0
Podocnemis expansa	Podocnemididae	109.0
Trionyx triunguis	Trionychidae	108.4
Chelonoidis niger darwini	Testudinidae	107.0
Chelonoidis niger becki	Testudinidae	105.0
Cylindraspis inepta †	Testudinidae	104.0
Pelochelys bibroni	Trionychidae	102.0
Aldabrachelys gigantea daudinii †	Testudinidae	101.0
Chitra vandijki	Trionychidae	100.0
Pelochelys cantorii	Trionychidae	100.0
Pelochelys signifera	Trionychidae	100.0
Chelonoidis niger abingdonii †	Testudinidae	99.1
Chitra indica	Trionychidae	99.0
Chelonoidis niger hoodensis	Testudinidae	97.0
Aldabrachelys gigantea arnoldi	Testudinidae	96.0
Chelonoidis niger niger †	Testudinidae	96.0
Natator depressus	Cheloniidae	95.0
Chelonoidis niger chathamensis	Testudinidae	95.0
Eretmochelys imbricata	Cheloniidae	94.0
Chelonoidis niger guntheri	Testudinidae	94.0
Nilssonia gangetica	Trionychidae	94.0
Chelonoidis niger phantasticus	Testudinidae	88.0
Centrochelys sulcata	Testudinidae	86.0
Chelonoidis niger duncanensis	Testudinidae	85.0
Chelonoidis niger vicina	Testudinidae	84.2
Amyda ornata phayrei	Trionychidae	83.0
Chelonoidis denticulatus	Testudinidae	82.0
Nilssonia nigricans	Trionychidae	81.0
Macrochelys suwanniensis	Chelydridae	80.1
Orlitia borneensis	Geoemydidae	80.0
Lepidochelys olivacea	Cheloniidae	79.3
Macrochelys temminckii	Chelydridae	77.5
Lepidochelys kempii	Cheloniidae	76.0
Batagur borneoensis	Geoemydidae	76.0
Cylindraspis triserrata †	Testudinidae	75.0
Stigmochelys pardalis	Testudinidae	75.0
Amyda cartilaginea maculosa	Trionychidae	75.0
Chelonoidis niger microphyes	Testudinidae	73.8

Table 6. Top 50 smallest modern turtle taxa by maximum straight-line carapace length (SCL, cm) of the larger sex; † = extinct.

Taxon	Family	SCL
Kinosternon vogti	Kinosternidae	10.2
Pelodiscus shipian	Trionychidae	10.3
Chersobius signatus	Testudinidae	11.0
Chersobius boulengeri	Testudinidae	11.0
Kinosternon subrubrum hippocrepis	Kinosternidae	11.1
Kinosternon steindachneri	Kinosternidae	11.4
Chersobius solus	Testudinidae	11.4
Glyptemys muhlenbergii	Emydidae	11.5
Sternotherus intermedius	Kinosternidae	11.6
Pelodiscus huangshanensis	Trionychidae	11.6
Kinosternon hirtipes megacephalum †	Kinosternidae	11.7
Kinosternon acutum	Kinosternidae	12.0
Kinosternon angustipons	Kinosternidae	12.0
Homopus areolatus	Testudinidae	12.0
Pelusios nanus	Pelomedusidae	12.2
Kinosternon creaseri	Kinosternidae	12.5
Kinosternon subrubrum subrubrum	Kinosternidae	12.5
Sternotherus depressus	Kinosternidae	12.5
Sternotherus peltifer	Kinosternidae	12.6
Kinosternon cora	Kinosternidae	12.8
Geoemyda spengleri	Geoemydidae	12.8
Pyxis arachnoides brygooi	Testudinidae	12.8
Pelodiscus axenaria	Trionychidae	12.9
Emys orbicularis eiselti	Emydidae	13.1
Psammobates tentorius tentorius	Testudinidae	13.1
Kinosternon alamosae	Kinosternidae	13.5
Clemmys guttata	Emydidae	13.6
Kinosternon hirtipes magdalense	Kinosternidae	13.6
Kinosternon hirtipes tarascense	Kinosternidae	13.6
Kinosternon baurii	Kinosternidae	13.8
Kinosternon hirtipes hirtipes	Kinosternidae	14.0
Psammobates tentorius verroxii	Testudinidae	14.0
Testudo (Testudo) kleinmanni	Testudinidae	14.4
Sternotherus minor	Kinosternidae	14.5
Psammobates tentorius trimeni	Testudinidae	14.5
Terrapene nelsoni nelsoni	Emydidae	14.7
Psammobates oculifer	Testudinidae	14.7
Sternotherus odoratus	Kinosternidae	15.0
Emys orbicularis ingauna	Emydidae	15.0
Platysternon megacephalum shiui	Platysternidae	15.1
Kinosternon hirtipes chapalaense	Kinosternidae	15.2
Sacalia quadriocellata	Geoemydidae	15.2
Pyxis arachnoides oblonga	Testudinidae	15.4
Pseudemydura umbrina	Chelidae	15.5
Graptemys sabinensis	Emydidae	15.5
Chrysemys dorsalis	Emydidae	15.6
Sacalia bealei	Geoemydidae	15.6
Rhinoclemmys rubida perixantha	Geoemydidae	15.6
Pelomedusa schweinfurthi	Pelomedusidae	15.7
Pyxis arachnoides arachnoides	Testudinidae	15.7

circumdata, the second-most at 2.080, and Barbour's Map Turtle, *Graptemys barbouri*, the third-most at 1.515.

As SCL data for many taxa are generalized or missing or only provide values without sexual differentiation (and often unclear as to whether midline or maximum SCL), we invite and encourage our readers to submit documented corrections, updates, and missing values and reference sources as necessary. Any input will be gratefully accepted and appropriately acknowledged.

Type Specimens

We introduced information on type specimens in the previous checklist, recording the types of all currently recognized species, subspecies, and synonymized taxa.

We list the primary name-bearing holotype, lectotype, neotype, or syntype specimen(s) as designated, with notes as warranted, but not secondary allotype, paratype, or paralectotype specimens, except as occasionally necessary. A more complete listing of all type material, including paratypes, can be found in Iverson (2022a). We document the status, designation, and museum location (if known) of name-bearing type specimens for all turtle and tortoise taxa with type localities (whether designated or not). Of these, many have no known or located type specimens, or the types are apparently lost, or there are only illustrated iconotypes (figured type specimens). Where possible, we document figured type specimens for taxa with no located type specimens. The acronyms we use for the museum collections holding turtle type specimens are defined at the end of the checklist, and are also documented in Sabaj (2019) and Uetz et al. (2019). We note that several type specimens listed by Uetz et al. (2019) unfortunately are not accurate or lack up-to-date lectotype or neotype designations.

Since DNA-based character identification of historical and ancient name-bearing type specimens is emerging as an increasingly important component of current taxonomic differential diagnoses and for resolution of nomenclatural issues and controversies (Renner 2016; Kehlmaier et al. 2019a), those historical turtle type specimens that are known to have been genetically sampled and analyzed to date are indicated and referenced in the checklist.

Several older turtle type specimens, collected and described in the 18th through the 20th centuries, have already been genetically analyzed (e.g., Austin and Arnold 2001; Austin et al. 2002, 2003; Blanck et al. 2006a; Praschag et al. 2008; Stuart and Fritz 2008; Murphy et al. 2011; Stuckas and Fritz 2011; Parham et al. 2012; Poulakakis et al. 2012, 2015; Stuckas et al. 2013; Fritz et al. 2014; Petzold et al. 2014; Tiedemann et al. 2014; Kindler et al. 2016; Kehlmaier et al. 2019a, 2019b). We urge that many more ancient and historical turtle type specimens undergo genotyping with DNA-based sampling and characterization.

A few holotype specimens of recently described taxa have also been genotyped at the time of their description (e.g., Blanck et al. 2006a; Petzold et al. 2014; Poulakakis et al. 2015; Ihlow et al. 2016). However, this is still uncommon and even though most recent descriptions of new taxa are at least partly based on DNA characterization, holotype specimens are not yet being routinely genetically analyzed. We recommend that authors that describe new taxa that are based at least partly on either mitochondrial or nuclear DNA characterization, should also consider genetically analyzing the selected holotype specimen. All genetically analyzed type specimens, whether ancient and historical or newly described, should have their genetic samples accessioned into GenBank (https://www.ncbi.nlm.nih.gov/genbank/) or the European Nucleotide Archive (ENA; https://www.ebi.ac.uk/ena), or other similar public repository, and their accession numbers published. We solicit input from our readers to update and correct any errors of omission or commission.

Etymologies

We are introducing etymologies for all named genera, subgenera, species, and subspecies in this edition of the checklist. These have been systematically sourced and collated by Iverson and Rhodin over the years, and recently augmented by an extensive listing submitted to us by Holger Vetter. In recording etymologies, we distinguish and record nine basic types of sources of names that have been used for turtle taxa.

Latin. — A name based on Latin (e.g., *Podocnemis unifilis* from the Latin words *unus* and *filum*). The generic Latin names for turtles or tortoises are *testudo* and *testudines*.

Greek. — A name based on Greek (e.g., *Acanthochelys macrocephala* from the Greek words *makros* and *kefale*). The generic Greek names for turtles or tortoises are *emys*, *chelone*, *chelys*, and *klemmys*.

The Latin and Greek meanings and origins of names have been partially sourced from Jaeger (1955) and Brown (1956).

Vernacular. — A name typically based on local indigenous tribal languages (e.g., Australian Aboriginal or Native Amazonian). The name is sometimes Latinized (e.g., *Peltocephalus tracaxus*, *Cuora couro*), but often not (e.g., *Chelodina burrungandjii*, *Testudo arrau*, *Amyda ornata jongli*). Occasional vernacular names are based on colloquial words in more widespread languages (e.g., Old French—*Testudo coui*, or English from Old French—*Gopherus*).

Eponym. — A name that honors a person or persons. Eponyms are typically Latinized and end either in *-i* (e.g., *Chelodina parkeri*) or *-ii* (e.g., *Lepidochelys kempii*) or *-ia* (e.g., *Bellia*, *Nilssonia*) when masculine, *-ae* when feminine (e.g., *Cuora flavomarginata evelynae*), and *-arum* or *-orum* when plural (e.g., *Elseya lavarackorum*). However, they may also end in *-anus* (e.g., *Phrynops geoffroanus*), *-ana* (e.g., *Kinixys belliana*), or *-anum* (e.g., *Kinosternon berendtianum*) (all meaning "belonging to"). Occasionally, they are simply rendered as the actual name of the honored person (e.g., *Clemmys sigriz*).

Eponyms are elsewhere sometimes also informally referred to as patronyms, but patronyms (or patronymics) are technically family names based on the given name of the father (e.g., Thomas Robertson, son of Robert Johnson) and not correct as a designation for the eponymous names used in scientific nomenclature.

Biographical information that we include on persons honored with eponyms has been gathered from multiple sources, including the original descriptions, Wikipedia, other online data repositories, personal contacts, and

biographical publications by Adler (1989, 2007, 2012), Beolens et al. (2011), and Li Vigni (2013).

Toponym. — A name that refers to a geographic place or region where the taxon occurs. Toponyms are typically Latinized and include the name of the place or region and end in *-ensis* or *-ense* (e.g., *Erymnochelys madagascariensis, Kinosternon durangoense*) (meaning "belonging to," implying "coming from"). They may also end in *-anus* (e.g., *Pelusios rhodesianus*), *-ana* (e.g., *Kinixys lobatsiana*), or *-anum* (e.g., *Kinosternon cobanum*) (also meaning "belonging to"). Other toponyms are rendered in the genitive and end in *-ae* or *-is*, meaning "belonging to, pertaining to" (e.g. *Kinosternon alamosae, Emydura australis*).

Ethnonym. — A name that honors a defined ethnic group or indigenous tribe, typically one that at some point in time inhabited the region in which the taxon occurs (e.g., *Malaclemys terrapin tequesta, Emys orbicularis ingauna, Kinosternon hirtipes tarascense*).

Mythonym. — A name that honors a mythological or fictional character (e.g., *Testudo hercules* or *Elseya oneiros*).

Bionym. — A name that refers to a biological entity (e.g., *Malaclemys rhizophorarum*, a scientific name referring to mangroves).

Anonym. — An anonymous or undefined name for which there is no clear or obvious etymology. Some may be based on an apparent Eponym, possibly honoring an unidentified or fictitious person (e.g., *Ocadia, Manouria, Sacalia*), or possibly an unrecorded indigenous Vernacular name (e.g., *Batagur*), or some obscure Greek or Latin or other language word, or possibly a fabricated name or word-game (e.g., *Anota, Tanoa*, an anagram) with no apparent meaning. Most of the Anonyms that we record were coined by J.E. Gray in the 1800s.

Etymology Analysis. — We provide etymologies for 1,972 names of turtle and tortoise genera, subgenera, species, and subspecies in this checklist (see Table 7), and we include an alphabetized list of the 350 people honored with Eponyms in Appendix 1 at the back of the checklist.

Table 7. Frequency of use of the nine types of etymologies that have been used for turtle genus-level and species-level names.

Category	No. of Uses	Percentage of Use
Latin	637	32.3%
Eponym	476	24.1%
Greek	411	20.8%
Toponym	311	15.8%
Vernacular	95	4.9%
Anonym	21	1.0%
Ethnonym	11	0.6%
Mythonym	8	0.4%
Bionym	2	0.1%
Totals	**1,972**	**100.0%**

Taxonomic Changes

A primary purpose of this checklist is to record taxonomic changes published in the literature, to evaluate the strength of the data supporting those proposed changes, and to recommend whether the community should adopt or reject the proposed changes. It is important to note that the recommendations we make here as the TTWG regarding the validity or non-validity of any included or excluded taxonomic names or systematic relationships are not prescriptive, nor are they official recommendations by the TFTSG or the IUCN, as such matters are generally best left to specialists working in these areas. However, we have tried to be consistent in our listing of what appear to be valid taxa and relationships, based on criteria of published scientific descriptions and proposed taxonomic changes accompanied by data and sound argumentation (see TTWG 2007a and inset on TTWG Guidelines on Taxonomic Changes on p. 11).

Our hope is that through this process, the TTWG and the TFTSG may help stabilize and guide the fluid state of chelonian systematics and nomenclature, and provide a standard reference source for updated taxonomy, systematic relationships, distributions, and conservation status of all turtles and tortoises. The list should also provide an impetus for ongoing and future work aimed at clarifying and resolving areas of taxonomic disagreement and/or uncertainty, as well as documented distribution patterns.

The very first checklist (TTWG 2007b) was compiled on the 'last published revision' principle, though reflecting some alternative arrangements through our use of the '*Xxxx* **or** *Yyyy*' arrangement. As the checklist has developed over the years and is increasingly adopted as the taxonomic standard by other groups and entities (IUCN Red List, Reptile Database, and others), and informs nomenclatural deliberations in CITES, ITIS, and other institutions, the TTWG author team has increasingly felt a need to evaluate both the scientific merit and the wider implications of adopting proposed taxonomic novelties. Evaluations have always been on a case-by-case basis, bringing the diverse perspectives of the authorship team to bear on the merits of each proposed change. We have considered drafting criteria for adoption or rejection, but concluded that every case is unique, making it unrealistic and undesirable to rely on a single set of "rules". Instead, we have formulated guidelines and considerations of what increases (or decreases) the scientific credibility of a proposed taxonomic novelty, and therefore the likelihood of its adoption into (or rejection from) the TTWG turtle checklist (see inset on p. 11).

We have previously (TTWG 2007a) presented proactive guidelines for researchers proposing taxonomic novelties; these remain valuable guidance also when we evaluate new published names or arrangements. But updating the checklist has required additional considerations, which we describe here. None of these are all-or-nothing decisions; instead, almost every proposed taxonomic novelty, and the underlying supporting data as presented in the publication, falls somewhere on a

TTWG Guidelines for Taxonomic Changes

Taxonomy is both a summary of scientific knowledge and a language for biological communication. As such, it is critical that taxonomic changes be carefully considered and based on strong, comprehensive underlying data to ensure that changes are stable and long-lasting. We fully recognize that taxonomy and the systematics research on which it is based, is a dynamic field and that change is a sign of healthy science. However, we also recognize that taxonomic and nomenclatural stability are of immense value to the wider community of biologists, conservationists, legislative authorities, and the public at large. Pauly et al. (2009) argued that taxonomy should aim for stability and monophyly; in cases where these two objectives are in conflict, well-supported monophyly usually prevails over stability. Given the dynamic nature of turtle taxonomy, we believe that a series of best practices can and should be followed that should lead to changes that are stable, informative, and long-lasting. We summarize these best practices both to identify many of the key points in our group discussions on newly-proposed name changes, and as a set of considerations for authors who are considering new name changes. We hope the community finds them useful. For additional related discussions, see TTWG (2007a), Pauly et al. (2009), Kaiser et al. (2013), Rhodin et al. (2015), and Thomson et al. (2018).

1. A proposed taxonomic change must meet the ICZN criteria for nomenclatural validity. Published names gain much greater credibility by being published in a peer-reviewed scientific journal or equivalent publication standard. These standards include the 2012 emendations of the Code (ICZN 2012) regarding accepted methods of electronic publication of new names.

2. Taxonomic changes above the species level should preferably be suggested and adopted only when a currently recognized higher taxon is demonstrably non-monophyletic. We share the view of the global systematics community that phylogeny should be reflected in higher taxonomic categories, and that changes should be proposed to "fix" a non-monophyletic grouping. As discussed in no. 5 below, non-monophyly should be based on multiple lines of statistically well-supported evidence. As pointed out by Pauly et al. (2009), the use of novel levels within a taxonomic hierarchy (subgenera, supergenera, etc.) allows for the recognition of new and/or previously known clades while still maintaining taxonomic stability within a group.

3. Taxonomic changes should incur the fewest possible name changes while resulting in a final set of monophyletic taxa. We share the view that taxonomic stability, and therefore the fewest possible nomenclatural changes, is always a desirable outcome.

4. Avoid naming monotypic higher groups when possible. As has been repeatedly stated in the literature, monotypic genera, families, etc. provide only very limited information on group membership, and therefore are less informative than alternative schemes where higher groups have multiple species within them. This may imply merging or lumping, rather than splitting, to resolve issues of non-monophyly. On the other hand, monotypic higher taxa emphasize the unique position of its contained (surviving) taxon. Monotypic higher taxa have been recognized among turtles for over two centuries, and many (though not all) contain additional extinct taxa as well as a single surviving species. We do not advocate eliminating traditionally recognized monotypic higher taxa, since that would lead to taxonomic destabilization, but caution against proliferation of monotypic higher taxa.

5. Taxonomic arrangements that are supported by several independent character sets, provide strong support for each, and report reasonable concordance between different datasets are more compelling than results from a single character set. Independence in evolutionary studies is a complex concept. In systematics, independence means that characters are not constrained to covary. For example, when multiple genetically independent nuclear genes, or nuclear genes and morphological characters, imply the same phylogenetic relationships or species boundaries, they presumably do so because both reflect the evolutionary history of the contained lineages. However, two mitochondrial genes are far less independent, since they are physically linked in the same non-recombining piece of mtDNA, and natural selection, drift, introgression, or any other process acts simultaneously on that linked set of nucleotides. Single characters (e.g., only mitochondrial or nuclear DNA, or only geographic distribution patterns) may reflect the history of the species, or they may reflect the history of that one character. We strongly recommend that individual characters (each nuclear gene, composite set of mtDNA data, morphological, behavioral, and other characters) be analyzed separately to test for concordance among multiple independent data sets.

6. Independent datasets may or may not provide convincing evidence for monophyly and taxonomic changes. When one dataset conflicts strongly with several other independent ones, there may still be strong support for the hypothesis supported by multiple independent data sets. However, character conflict may often suggest that additional analyses or data are needed before taxonomic changes should be endorsed and accepted.

7. Sampling should be comprehensive at the appropriate level. Broad taxon sampling for species trees, with multiple specimens from across the geographic range of each taxon, can help avoid spuriously high statistical support values for apparent clades (see Spinks et al. 2013 for a recent chelonian example).

8. Species delimitation studies should include broad geographic sampling of all relevant taxa. Comprehensive geographic sampling for each character from individuals across the ranges of all species being considered is often critically important to correctly diagnose new species. We recognize that comprehensive geographic sampling may be difficult for rare species, but every effort should be made to be as comprehensive as possible.

9. Studies that only evaluate a taxonomic or geographic subset of the relevant group, or only make changes to some taxa without evaluating the relevance of these changes to related taxa, are less likely to be convincing and stand the test of time, and therefore are less likely to be widely adopted. For example, a study that elevates a particular subspecies to species rank, without examining variation among the remainder of the species is unlikely to be adopted until further supporting and clarifying information is published.

10. The TTWG primarily reacts to taxonomic changes proposed in the published literature, although we also will take under consideration publications that are under review but not yet published if they add additional information to a proposed change. Any information that the TTWG members have access to can be used to argue for or against adoption of a new taxonomic arrangement proposed in a validly published publication, although in almost all cases we rely on information that is either published or under review in a peer-reviewed journal. The TTWG will not use information from an as-yet unpublished study or manuscript to initiate a taxonomic change. In very rare cases, the TTWG may decide to make a new nomenclatural act, such as creating new nomenclatural combinations.

Table 8. Top turtle-rich countries for all **Extant Species** of terrestrial tortoises, freshwater turtles, and nesting sea turtles.

Tortoises & Freshwater Turtle Species	Tortoises & Freshwater Turtles & Nesting Sea Turtles
1. USA, 59	1. USA, 64
2. Mexico, 49	2. Mexico, 56
3. Brazil, 32	3. Brazil, 37
Indonesia, 32	Indonesia, 37
5. China, 31	5. India, 35
India, 31	6. China, 34
7. Colombia, 29	Colombia, 34
8. Myanmar, 28	8. Australia, 32
9. Vietnam, 27	9. Myanmar, 31
10. Australia, 26	Vietnam, 31
Thailand, 26	11. Bangladesh, 30
12. Bangladesh, 25	Thailand, 30
13. Laos, 19	13. Malaysia, 23
Malaysia, 19	14. Venezuela, 22
15. South Africa, 18	15. Ecuador, 20
Venezuela, 18	South Africa, 20
17. Congo (DRC), 17	17. Laos, 19
Nepal, 17	18. Congo (DRC), 17
19. Bolivia, 16	Guatemala, 17
Ecuador, 16	Nepal, 17
21. Cambodia, 14	Panama, 17
Guatemala, 14	22. Bolivia, 16
23. Cameroon, 13	Cameroon, 16
Peru, 13	Papua New Guinea, 16
25. Angola, 12	25. Mozambique, 15
Central African Republic, 12	Peru, 15
Kenya, 12	
Nigeria, 12	
Papua New Guinea, 12	
Tanzania, 12	
Uganda, 12	

Table 9. Top turtle-rich countries for all **Extant Taxa** (species and subspecies) of terrestrial tortoises, freshwater turtles, and nesting sea turtles.

Tortoises & Freshwater Turtles Taxa	Tortoises & Freshwater Turtles & Nesting Sea Turtles
1. USA, 83	1. USA, 88
2. Mexico, 61	2. Mexico, 66
3. India, 39	3. India, 43
4. China, 35	4. Indonesia, 39
Indonesia, 35	5. China, 38
6. Brazil, 32	6. Brazil, 37
7. Vietnam, 31	7. Australia, 36
8. Australia, 30	8. Vietnam, 35
9. Colombia, 29	9. Colombia, 34
Myanmar, 29	10. Myanmar, 32
11. Thailand, 28	Thailand, 32
12. Ecuador, 27	12. Bangladesh, 31
13. Bangladesh, 26	Ecuador, 31
14. Malaysia, 22	14. Malaysia, 26
15. Laos, 20	15. Venezuela, 24
South Africa, 20	16. South Africa, 22
Venezuela, 20	17. Laos, 20
18. Congo (DRC), 17	18. Nepal, 18
Nepal, 17	19. Congo (DRC), 17
20. Bolivia, 16	Guatemala, 17
21. Cambodia, 14	21. Bolivia, 16
Guatemala, 14	Cameroon, 16
Peru, 14	Mozambique, 16
24. Cameroon, 13	Panama, 16
Tanzania, 13	Papua New Guinea, 16
Uganda, 13	Peru, 16

Table 10. Top 10 turtle-rich countries for **Extant Endemic Taxa** (species and subspecies) of terrestrial tortoises and freshwater turtles.

1. USA, 59	6. Indonesia, 9
2. Mexico, 32	7. India, 8
3. Australia, 27	Madagascar, 8
4. China, 14	9. Brazil, 6
5. Ecuador, 12	Myanmar, 6

continuum between 'adopt unreservedly' and 'reject outright'.

The collective weight of evidence supporting any proposed change (availability of the name; strength and nature of the supporting evidence; phylogenetic context; agreement with other studies; effect on taxonomic stability) is deliberated by the TTWG team (often very extensively and often with different philosophical views of the value of the underlying evidence). In order to provide a more comprehensive and international approach to TTWG deliberations and decisions, especially as regards issues of phylogenetic analysis, in 2017 we expanded our previous authorship team (Rhodin, Iverson, van Dijk, Shaffer, and Bour) to also include Georges and Fritz, and in 2022, after Bour's passing, we added Gallego-García. We have also frequently sought the advice of experts outside the TTWG. Our deliberations lead to conclusions on whether to:

1) adopt fully, reject, or recommend modification of a proposed taxonomic change,

2) include it as an '*Xxxx* (or *Yyyy*)' arrangement, or

3) suspend adoption until additional, independent supporting or alternative data are published.

It is important to note that decisions and recommendations within the TTWG are not always unanimous, and our checklist is not necessarily reflective of the individual taxonomic views or conclusions of all team members. In the accompanying text box (see inset on p. 11), we summarize our guidelines and recommendations for making taxonomic changes.

Nomenclatural and taxonomic changes often have disruptive effects for legislation and other 'users' of checklists. A degree of disruption is inevitable as phylogenetic knowledge accumulates; but we are more likely to adopt proposed changes that have significant 'disruptive' effects on widely-used names if such changes are strongly supported by robust data. In contrast, we are inclined to suspend adoption of novel names and arrangements if they are based on weaker data sets or do not greatly improve our overall phylogenetic understanding. As an example, we would be reluctant to adopt a proposal to transfer a single species out of an




Figure 2. Updated Global Species Richness Map for all 352 extant freshwater and terrestrial turtle and tortoise species (not including extinct species or marine turtles) as presented in this 2025 checklist and atlas using fine-scaled HUC level 12 hydrobasin polygons. See comparative fine-scaled map in TTWG (2021) and the earlier map by Buhlmann et al. (2009) that used broad-scaled HUC level 06 hydrobasin polygons. For enlarged and more detailed regional and focused species richness maps, see Figs. 3–8.

established genus to form a new, monotypic genus, a move that would involve new names and combinations without significant improvement of our understanding of the overall relationships of the group of species. We repeat our recommendation (TTWG 2007a) that taxonomy should not be driven by politics or opportunism, but that the wider implications of taxonomic and nomenclatural decisions be understood and carefully considered.

We have noted for many years that the ICZN (2012) has emended its Code regarding accepted methods of electronic publication of new names. The revision permits electronic publication after 2011 only after the work (not the new name) is first registered in ZooBank (http://zoobank.org/; The Official Registry of Zoological Nomenclature). ZooBank must register the precise electronic archive where the work is to be published, as well as the ISSN or ISBN of the work. In addition, amendments to the Code also clarify that preliminary electronic versions of works due for publication on paper are unavailable, and that abstracts of meetings, presentation texts and posters are unavailable for nomenclatural purposes, and preliminary electronic versions do not bring forward the date of publication, unless the electronic version meets the requirements for availability. Authors intending to publish taxonomic novelties in electronic archives are cautioned to read the text of International Commission on Zoological Nomenclature (2012) carefully, and to follow the guidelines precisely, or be at risk of having their work judged as inadmissible.

Distributions

We summarize distributions for all taxa in the checklist, listing all nations and territories in which they occur as native populations (see Tables 8–10 for the top turtle-rich nations). For several larger nations we also list political or geographic subunits (e.g., states, provinces, regions, or larger islands). We attempt to also indicate nations or territories where species have been extirpated or where they occur as non-native introduced or invasive species, or where there are uncertainties as to occurrence. Where nations or subunits have been added or deleted due to new data acquired since our last checklist, we have marked those distribution entries with an asterisk (*).

For introductions, we attempt to distinguish between two forms: (1) modern introductions (since ca. 1500 CE) for non-native species that appear to have relatively well-established or potentially reproducing populations in extra-limital areas primarily as a result of relatively recent trade for food or pets or planned conservation introductions (labeled “introduced” or “modern”), and (2) earlier historical or prehistorical introductions for those species that appear to have native populations, but where population genetics studies find evidence of founder effects suggestive of possible introduction by humans, or other dispersal events, during the last ca. 2000–3000 years (labeled “prehistorical introduction?”). However, many non-native records in the literature also appear to represent one or a few translocated individual turtles, possibly representing released trade or pet specimens, and not necessarily reproductive populations; many of these are also included on our maps as orange dots.

For freshwater and terrestrial turtles and tortoises, we compiled native and introduced distributions and locality records from a combination of multiple published and database sources. For native distributions we used Iverson (1992) and Fritz and Havaš (2007) as

Figure 3. Updated Species Richness Map for North and South America as presented in this checklist and atlas using fine-scaled HUC level 12 hydrobasin polygons. For composite global species richness map, see Fig. 2.




Figure 4. Updated Species Richness Map for Europe and Africa as presented in this checklist and atlas using fine-scaled HUC level 12 hydrobasin polygons. For composite global species richness map, see Fig. 2.




Figure 5. Updated Species Richness Map for Asia and Australia as presented in this checklist and atlas using fine-scaled HUC level 12 hydrobasin polygons. For composite global species richness map, see Fig. 2.





Figure 6. Updated Focused Species Richness Map for only Europe as presented in this checklist and atlas using fine-scaled HUC level 12 hydrobasin polygons.



starting points, and then added data from other recent literature by numerous authors, data from our extensive database compiled from Iverson's work by Buhlmann et al. (2009), further data from our published CBFTT species accounts, our TFTSG-organized IUCN Red Listing workshops, and data from Iverson's continuing intensive compilation of distribution records from the literature. Recently we have also added numerous distribution records from the Global Biodiversity Information Facility (GBIF).

For introduced species, we used Kraus (2009) as a starting point and have added data from other publications and online sources and databases. We have attempted to primarily list introductions that are based on recorded populations rather than just single animals (but not necessarily with evidence of reproduction), but have not been rigorous in this, as it can be difficult to determine what the actual situation may be in each recorded case. Many locality records no doubt represent individual translocated non-native animals. See Reshetnikov et al. (2023) for an excellent analysis and discussion of the invasive capability of *Trachemys scripta elegans*, and Mu and Li (2023) for a discussion of the increased risk of further invasiveness by *Chelydra serpentina* in China as predicted by climate change models.


Finally, we also solicited and received input from many members of the TFTSG for corrections and additions to all the native distributions and introductions data. Despite this effort, it is likely that we have committed errors of omission or commission, and we request that any corrections or updates be brought to our attention so that they can be included in future editions of this checklist.

For sea turtles, we compiled distributions from a combination of IUCN Red List data, CMS (Convention on Migratory Species) listings, and the extensive GIS shapefiles and spreadsheets of nesting sites and foraging ranges that the IUCN SSC Marine Turtle Specialist Group (MTSG) includes in its SWOT mapping application (The State of the World's Sea Turtles) (http://seaturtlestatus.org/learn/maps/all), updated in 2024 (Kot et al. 2023), generously shared with us by the MTSG. Based on these data, we list sea turtle distributions in three distributional categories: 1) nesting: native regularly nesting populations, 2) foraging: native permanently foraging or regularly migrating populations (but no evidence yet of regular nesting), and 3) vagrant: temporarily foraging or migrating animals not necessarily considered native. Our marine turtle maps also show the oceanic Regional Management Units developed by the MTSG for each species, with recent revisions and updates (Wallace et al. 2023).

GIS Maps and Historical Ranges

This edition of the checklist includes 631 separate distribution maps for species, subspecies, families,

Figure 7. Updated Focused Species Richness Map for only Africa as presented in this checklist and atlas using fine-scaled HUC level 12 hydrobasin polygons.




suborders, and species richness. We provide enhanced and updated maps that include specific locality points upon which species and subspecies ranges are based, as well as detailed topographic features.

Native distribution locality points are indicated by **yellow dots**; probable or possible non-native localities (trade specimens, translocations or introductions, and/or questionable identifications) are indicated by **orange dots**; extirpated populations or extinct taxa are indicated by **red dots**. Type localities of recognized or synonymized taxa are indicated by **larger color-coded stars**, usually yellow for the accepted type locality of the recognized taxon and orange for erroneous or incorrect type localities (e.g., out of range) for the recognized taxon, and other labeled colors for any synonymized taxa.

Estimated Indigenous (Historical) Ranges for all species and taxa are continuously revised and recalculated based on new data and improved geographic analysis, in general restricting ranges more tightly around recorded occurrence localities and accounting for geographic watershed limits and habitat preferences of the species involved (e.g., whether primarily riverine or more lacustrine, semi-terrestrial, or fully terrestrial. Many such ranges previously calculated in TTWG (2021) have been further refined and recalculated in this edition; these updates are marked with an asterisk (*).

We have followed the definition of the **indigenous range** of a species as established by the IUCN (2013) as "the known or inferred distribution generated from **historical** (written or verbal) records, or physical evidence of the species' occurrence. Where direct evidence is inadequate to confirm previous occupancy, the existence of suitable habitat within ecologically appropriate proximity to proven range may be taken as adequate evidence of previous occupation." Following the recommendations of Akçakaya et al. (2018) and Stephenson et

Figure 8. Updated Focused Species Richness Map for only Australia and New Guinea as presented in this checklist and atlas using fine-scaled HUC level 12 hydrobasin polygons.



al. (2019), we have selected a benchmark cutoff date of approximately 1500 CE for the determination of the Estimated Indigenous (Historical) Range of all turtle species, that being the year from which the IUCN Red List considers modern extinctions and the approximate start of European expansion and global development patterns beginning to impact native turtle ranges.

Edited or new maps in the current checklist with new locality points and revised presumed historical ranges differing from the previous checklist are indicated by a double asterisk (**); those with only new locality dots are indicated by a bullet (•).

The total number of locality records providing the basis for the maps in the current checklist is 120,753 points, or an average of 332 records per species, ranging from only single records for each of the five extinct *Cylindraspis* species and *Pelodiscus huangshanensis* to 3,349 for *Trachemys scripta*, 3,411 for *Emys orbicularis*, 6,651 for *Chrysemys dorsalis*, 9,300 for *Sternotherus odoratus*, 9,492 for *Chrysemys picta*, and 9,752 for *Emydura macquarii* as the six species with the most records entered. For those species with very large numbers of records, very many represent specimens from the same locality, or close-by, and show up only as single dots on our maps.

Map production began with point locality datasets from Iverson (1992), based on the many museum-held voucher specimens and published records amassed by him over the years and updated on the EmySystem website (http://emys.geo.orst.edu/). These datasets were then supplemented by newer data and converted into shapefiles and edited and corrected and updated for content by Iverson, Ross Kiester, Tom Akre, Kurt Buhlmann, Peter Paul van Dijk, Arthur Georges, Anders Rhodin, Russ Mittermeier, and Whit Gibbons, and analyzed by Buhlmann et al. (2009).

The original maps created this way were based on constructing projected historical ranges. This was

Table 11. Top 50 modern turtle **taxa** with the smallest Estimated Indigenous (Historical) Range (AOO) sizes (sq. km); † = extinct.

Taxon	Family	Size
Chelonoidis niger duncanensis	Testudinidae	10
Chelonoidis niger abingdonii †	Testudinidae	15
Chelonoidis niger ssp. Santa Fé †	Testudinidae	18
Chelonoidis niger hoodensis	Testudinidae	21
Chelonoidis niger niger †	Testudinidae	78
Chelonoidis niger donfaustoi	Testudinidae	89
Chelonoidis niger phantasticus	Testudinidae	109
Chelonoidis niger microphyes	Testudinidae	111
Cylindraspis peltastes †	Testudinidae	113
Cylindraspis vosmaeri †	Testudinidae	113
Aldabrachelys gigantea daudinii †	Testudinidae	159
Aldabrachelys gigantea arnoldi	Testudinidae	162
Chelonoidis niger darwini	Testudinidae	173
Aldabrachelys gigantea gigantea	Testudinidae	190
Chelodina mccordi timorensis	Chelidae	224
Aldabrachelys gigantea hololissa	Testudinidae	229
Chelonoidis niger becki	Testudinidae	234
Pelusios castanoides intergularis	Pelomedusidae	240
Pelusios subniger parietalis	Pelomedusidae	240
Chelodina mccordi mccordi	Chelidae	242
Kinosternon vogti	Kinosternidae	263
Chelonoidis niger chathamensis	Testudinidae	282
Chelonoidis niger porteri	Testudinidae	306
Chelonoidis niger vandenburghi	Testudinidae	317
Chelonoidis niger vicina	Testudinidae	357
Chelonoidis niger guntheri	Testudinidae	499
Apalone spinifera atra	Trionychidae	535
Pelusios williamsi laurenti	Pelomedusidae	551
Cuora flavomarginata evelynae	Geoemydidae	668
Kinosternon hirtipes magdalense	Kinosternidae	694
Mauremys mutica kami	Geoemydidae	705
Trachemys stejnegeri malonei	Emydidae	726
Kinixys zombensis domerguei	Testudinidae	835
Terrapene coahuila	Emydidae	840
Astrochelys yniphora	Testudinidae	884
Trachemys taylori	Emydidae	887
Myuchelys georgesi	Chelidae	1,002
Geoemyda japonica	Geoemydidae	1,613
Kinosternon hirtipes megacephalum †	Kinosternidae	1,659
Emys orbicularis ingauna	Emydidae	1,701
Cylindraspis inepta †	Testudinidae	1,882
Cylindraspis triserrata †	Testudinidae	1,882
Pseudemydura umbrina	Chelidae	1,884
Kinosternon sonoriense longifemorale	Kinosternidae	2,089
Emydura macquarii nigra	Chelidae	2,150
Cylindraspis indica †	Testudinidae	2,536
Chelodina kuchlingi	Chelidae	2,778
Malaclemys terrapin rhizophorarum	Emydidae	3,372
Cuora picturata	Geoemydidae	3,541
Emydura gunaleni	Chelidae	3,789

Table 12. Top 50 modern turtle **taxa** with the largest Estimated Indigenous (Historical) Range (AOO) sizes (sq. km).

Taxon	Family	Size
Kinosternon scorpioides	Kinosternidae	6,451,732
Chelonoidis denticulatus	Testudinidae	5,543,761
Chelydra serpentina	Chelydridae	5,364,328
Chelonoidis carbonarius	Testudinidae	5,323,930
Podocnemis unifilis	Podocnemididae	5,213,904
Platemys platycephala platycephala	Chelidae	5,097,698
Stigmochelys pardalis	Testudinidae	4,450,736
Mesoclemmys gibba	Chelidae	4,445,462
Emys orbicularis orbicularis	Emydidae	3,618,563
Centrochelys sulcata	Testudinidae	3,416,077
Mesoclemmys raniceps	Chelidae	3,410,211
Phrynops geoffroanus	Chelidae	3,407,292
Chelus fimbriata	Chelidae	3,173,941
Kinixys erosa	Testudinidae	2,920,544
Pelomedusa subrufa	Pelomedusidae	2,913,756
Chrysemys picta bellii	Emydidae	2,732,375
Kinixys spekii	Testudinidae	2,522,956
Sternotherus odoratus	Kinosternidae	2,352,387
Pelusios gabonensis	Pelomedusidae	2,289,713
Pelusios sinuatus bottegi	Pelomedusidae	2,288,663
Kinixys belliana	Testudinidae	2,268,595
Pelusios rhodesianus	Pelomedusidae	2,250,980
Trionyx triunguis	Trionychidae	2,198,985
Rhinoclemmys punctularia punctularia	Geoemydidae	2,188,265
Kinixys nogueyi	Testudinidae	2,154,510
Trachemys scripta elegans	Emydidae	2,010,329
Pelomedusa olivacea	Pelomedusidae	2,009,750
Cyclanorbis senegalensis	Trionychidae	1,998,159
Pelusios castaneus	Pelomedusidae	1,966,366
Mesoclemmys wermuthi	Chelidae	1,953,977
Terrapene ornata	Emydidae	1,930,983
Podocnemis expansa	Podocnemididae	1,706,101
Pelusios subniger subniger	Pelomedusidae	1,664,250
Chrysemys picta marginata	Emydidae	1,603,643
Pelochelys cantorii	Trionychidae	1,514,742
Pelusios adansonii	Pelomedusidae	1,475,150
Terrapene carolina carolina	Emydidae	1,424,405
Lissemys punctata andersoni	Trionychidae	1,343,471
Kinosternon flavescens	Kinosternidae	1,337,830
Cyclemys dentata	Geoemydidae	1,333,998
Indotestudo elongata	Testudinidae	1,308,716
Cuora couro couro	Geoemydidae	1,286,966
Testudo horsfieldii kazachstanica	Testudinidae	1,213,129
Chelonoidis chilensis	Testudinidae	1,118,096
Apalone spinifera spinifera	Trionychidae	1,086,381
Dogania subplana	Trionychidae	1,078,975
Psammobates oculifer	Testudinidae	1,078,671
Chelus orinocensis	Chelidae	1,032,210
Terrapene triunguis	Emydidae	1,017,532
Pelomedusa variabilis	Pelomedusidae	996,640

done by selecting GIS-defined hydrologic unit compartments (HUCs; www.hydrosheds.org), originally at coarse level 06 hydroshed basins with verified locality points, and then adding HUCs that connected known point localities in the same watershed or physiographic region and that had similar habitats and elevations as the verified HUCs. As such, these first maps represented predicted ranges that were generally much larger than reality, and required further verification and adjustment.

These distribution shapefiles were then further revised and formatted sequentially over the years by Rhodin, first using ArcGIS Desktop 10.1, then ArcMap 10.8.2, and finally ArcGIS Pro 3.3 (www.esri.com), initially as part of the IUCN-associated BioFresh initiative (http://atlas.freshwaterbiodiversity.eu/), first using finer

Table 13. Top 50 modern turtle **species** with the largest Estimated Indigenous (Historical) Range (AOO) sizes (sq. km).

Taxon	Family	Size
Kinosternon scorpioides	Kinosternidae	6,446,998
Chelonoidis denticulatus	Testudinidae	5,543,761
Chelydra serpentina	Chelydridae	5,364,328
Chelonoidis carbonarius	Testudinidae	5,323,930
Platemys platycephala	Chelidae	5,283,733
Podocnemis unifilis	Podocnemididae	5,213,904
Chrysemys picta	Emydidae	4,569,908
Emys orbicularis	Emydidae	4,567,751
Stigmochelys pardalis	Testudinidae	4,450,736
Mesoclemmys gibba	Chelidae	4,445,462
Centrochelys sulcata	Testudinidae	3,416,077
Mesoclemmys raniceps	Chelidae	3,410,211
Phrynops geoffroanus	Chelidae	3,407,292
Chelus fimbriata	Chelidae	3,173,941
Kinixys erosa	Testudinidae	2,920,544
Pelomedusa subrufa	Pelomedusidae	2,913,756
Pelusios sinuatus	Pelomedusidae	2,570,192
Kinixys spekii	Testudinidae	2,522,956
Trachemys scripta	Emydidae	2,447,717
Kinixys nogueyi	Testudinidae	2,445,542
Sternotherus odoratus	Kinosternidae	2,354,096
Pelusios gabonensis	Pelomedusidae	2,289,713
Kinixys belliana	Testudinidae	2,268,595
Pelusios rhodesianus	Pelomedusidae	2,250,980
Lissemys punctata	Trionychidae	2,239,325
Rhinoclemmys punctularia	Geoemydidae	2,213,334
Trionyx triunguis	Trionychidae	2,198,985
Testudo horsfieldii	Testudinidae	2,122,553
Testudo graeca	Testudinidae	2,013,047
Pelomedusa olivacea	Pelomedusidae	2,009,750
Cyclanorbis senegalensis	Trionychidae	1,998,159
Pelusios castaneus	Pelomedusidae	1,966,366
Mesoclemmys wermuthi	Chelidae	1,953,977
Terrapene ornata	Emydidae	1,930,983
Cuora couro	Geoemydidae	1,804,560
Apalone spinifera	Trionychidae	1,770,022
Podocnemis expansa	Podocnemididae	1,706,101
Pelusios subniger	Pelomedusidae	1,664,490
Terrapene carolina	Emydidae	1,632,524
Pelochelys cantorii	Trionychidae	1,514,742
Melanochelys trijuga	Geoemydidae	1,511,818
Pelusios adansonii	Pelomedusidae	1,475,150
Kinosternon subrubrum	Kinosternidae	1,444,909
Cyclemys dentata	Geoemydidae	1,345,504
Indotestudo elongata	Testudinidae	1,308,716
Kinosternon flavescens	Kinosternidae	1,284,360
Chelonoidis chilensis	Testudinidae	1,118,096
Dogania subplana	Trionychidae	1,078,975
Psammobates oculifer	Testudinidae	1,078,671
Chelus orinocensis	Chelidae	1,032,210

geographic scales (IUCN hydroshed basins at HUC levels 10 and 12) and the WGS 1984 coordinate system (available at https://www.iucnredlist.org/resources/spatialtoolsanddata), and then eventually only fine-scale HUC level 12 polygons based on WGS 1984 locality points.

This allowed elimination of many higher-altitude regions from the projected ranges (notably in areas such as the Himalayan and Andean foothills and other mountainous regions), while keeping lower altitude HUC distributions in the same overall drainage basins, and in general significantly tightening up and reducing many of the projected ranges around known locality points. Similarly, many primarily riverine species had their ranges limited to tighter distributions around the rivers from which they had been recorded.

As a result, the average difference in presumed ranges of the species recorded in Buhlmann et al. (2009) to their corresponding ranges recorded in the current checklist is a 46.6% reduction, with the average species range in Buhlmann et al. recorded as 1,108,310 sq. km, and the average range per species now calculated as 591,486 sq. km. The maps have also been further revised through input of data provided by authors of published CBFTT accounts and participants in TFTSG-organized IUCN Red Listing workshops, but still represent projected and assumed historical ranges. For some taxa with extremely limited distributions (e.g., *Chelonoidis niger* subspecies in the Galápagos and a few other small-island taxa), custom-drawn polygons smaller than HUC level 12 were used as appropriate.

Sea turtle maps were generated from GIS data generously supplied to us by the IUCN SSC Marine Turtle Specialist Group (MTSG) and SWOT, and show documented nesting sites as yellow dots and generalized foraging distributions for each species as shaded oceanic distributional ranges delimited as either Regional Management Units (RMUs) or Distinctive Population Segments (DPSs) (see Seminoff et al. 2015).

For some relatively cryptic, poorly known, or possibly questionable species, the ranges depicted in this checklist are at best general approximations of their potential distributions. Some species that fall into this category include *Pelodiscus axenaria*, *Pelodiscus huangshanensis*, *Cuora mccordi*, *Cuora yunnanensis*, *Cuora zhoui*, *Cyclemys enigmatica*, *Pelomedusa* spp., *Rafetus swinhoei*, and *Chelodina gunaleni*.

Widespread species with significant documented phylogeographic differentiation in the form of recognized subspecies or genetically-defined lineages and Evolutionarily Significant Units (ESUs) or Management Units (MUs) may eventually warrant recognition as multiple taxa at the species level. Some species that fall into this category include *Kinosternon hirtipes*, *Kinosternon integrum*, *Kinosternon scorpioides*, *Melanochelys trijuga*, *Chelonoidis carbonarius*, and *Phrynops geoffroanus.*

Native populations and recorded specimens are marked with yellow locality dots set in partially transparent distribution range polygons (using HUCs), using red polygons at 50% transparency for species and nominate subspecies, and other colors at 50% transparency for other subspecies. Populations that represent possible prehistorical introductions (whether genetically verified or hypothesized as such) or possible prehistorical or more recent natural range extensions are also indicated with yellow dots.

A composite global species richness map for all extant tortoise and freshwater turtle species is depicted in Fig. 2, with enlarged regional detailed maps in Figs. 3–5. We also provide focused continental richness maps for Europe, Africa, and Australia / New Guinea in Figs. 6–8, as each of those continents has less species richness than North and South America and Asia. We note that the three most prominent global areas of high species richness are the southeastern USA in North America, the Amazon basin in South America, and the Brahmaputra-Ganges basins in Asia, as previously noted by Buhlmann et al. (2009) and Mittermeier et al. (2015). See Ennen et al. (2020, 2021) for further in-depth analysis of global biogeographic distribution and conservation patterns.

Apparently introduced or translocated populations near the native range that are most likely of modern and recent historical origin are generally included, and are indicated by orange dots. Remote introduced populations are not included. Questionable locality records near the native range (possibly misidentified and/or regional trade specimens or translocated or released animals) are also indicated by orange dots. Extinct taxa (species or subspecies) are designated by red dots.

It is critically important to note that the maps published here depict **Estimated Indigenous (Historical) Ranges**, defined as the **historical** Area of Occupancy, or AOO, as per IUCN (2013) guidelines, as they are based on a combination of older historical museum and literature data and more recent locality data, and **do not** indicate the actual **current** Area of Occupancy of these species. Most turtle and tortoise species have had their historical ranges decrease considerably as a result of extensive habitat loss and degradation and/or overexploitation. For example, the ranges depicted here for *Batagur trivittata* and *Geochelone platynota* from Myanmar, and *Psammobates geometricus* from South Africa, show their **indigenous** ranges, rather than their **current** ranges, which have all been reduced by >90%. In general, all species assessed as Critically Endangered or Endangered on the IUCN Red List or the TFTSG Provisional Red List have had their current AOO ranges greatly decreased from historical extents.

We have calculated the estimated indigenous (historical) range, or original Area of Occupancy (AOO, in sq. km) for all taxa of tortoises and freshwater turtles—but not marine turtles—and include that value under each taxon. We have calculated these ranges precisely in either ArcMap 10.8.2 using Goode Homolosine (Land) equal area projections or in ArcGIS Pro 3.3 using Cylindrical Equal Area (sphere) projections of our GIS range polygons generated using fine-scaled HUC level 12. As such, these distribution ranges reflect apparently precise and reasonably accurate **Estimated Indigenous (Historical) Range** sizes, but **not** the actual current Area of Occupancy (AOO). Furthermore, each individual HUC polygon, especially along the edges of the depicted range, may be larger than the actual distribution of the species within that polygon, thereby somewhat overestimating the total presumed indigenous range.

We have also calculated the indigenous IUCN-defined Extent of Occurrence (EOO, in sq. km) for all tortoises and freshwater turtles and also include that value under each taxon. EOO is defined by IUCN as "the area contained within the shortest continuous imaginary boundary which can be drawn to encompass all the known, inferred or projected sites of...occurrence of a taxon" and generally includes areas of habitat discontinuity (IUCN 2022). We calculated estimated indigenous EOO precisely in ArcGIS based on our GIS fine-scaled HUC level 12 range polygons (not based on our plotted locality dots), and utilized the IUCN EOO Calculator (v1.5; https://www.iucnredlist.org/resources/spatialtoolsanddata), which calculates the area of the resultant minimum convex polygon hull using the Cylindrical Equal Area (world) projection. The IUCN EOO Calculator measures overall geographical extent of a taxon and does not exclude areas of habitat discontinuity such as marine or uninhabitable high-altitude environments within the EOO of a freshwater or terrestrial low-altitude species. We note that although these calculations reflect apparently precise and reasonably accurate estimated indigenous Extent of Occurrence (EOO), they do not necessarily reflect the actual current EOO.

Importantly, the precision and accuracy of these calculated Estimated Indigenous Ranges (AOO and EOO) are also clearly affected by our choice of how tightly or broadly to limit the selected range around our point localities, and how best to reflect apparent discontinuities in the range, and should not be interpreted as explicitly indicative of the apparent precision. We can clearly expect significant potential variance in the numbers provided, but have chosen not to round them off.

As an example of some of these potential discrepancies and caveats, the Estimated Indigenous Range for the Coahuilan Box Turtle, *Terrapene coahuila*, based on our map using fine-scale HUC level 12 polygons, is 1,170 sq. km, whereas the potential turtle habitat of the closed Cuatro Ciénegas Basin, where the species exists, is only about 840 sq. km (Howeth and Brown 2011 CBFTT), and the species' apparent Area of Occupancy (AOO) has been estimated to have shrunk from about 600 sq. km in the 1960s to only about 360 sq. km in the early 2000s (van Dijk et al. 2007 IUCN), and now considered to be less than 6 sq. km (Castañeda Gaytán 2020).

The average indigenous range size (historical AOO) for all 486 recognized tortoise and freshwater turtle taxa (species plus subspecies, excluding marine turtles) was 426,296 sq. km, ranging from the smallest range of 10 sq. km for the Pinzón Giant Tortoise, *Chelonoidis niger duncanensis*, to the largest range of 6,451,732 sq. km for the Scorpion Mud Turtle, *Kinosternon scorpioides*. Tables 11–13 list the top 50 ranked taxa with the smallest and largest Estimated Indigenous Ranges.

The present numbers differ somewhat from the calculations provided in our previous checklist (TTWG

2021), since we continuously update and revise our distribution maps with new locality data and reanalysis of estimated distribution patterns. For the present checklist, we have revised the Estimated Indigenous (Historical) Ranges of 348 taxa (71.6% of all 386 recognized non-marine taxa), all indicated by an asterisk (*). Of these, the estimated range of 189 taxa increased, and the estimated range of 159 taxa decreased. Overall, estimated ranges of all taxa decreased by an average of 5.63%, reflecting generally slightly smaller estimated ranges after our revisions, although some ranges increased substantially.

The taxa with the smallest indigenous ranges are the most likely to be endangered and threatened, with many already extinct (see Ripple et al. 2017; Newsome et al. 2020). Widespread species appear to be under less threat, but are more difficult to assess in terms of the degree to which they may be undergoing similar relative percentages of population reductions as the restricted-range species. Additionally, widespread species likely harbor potentially unrecognized regional or cryptic diversity that should be investigated for included evolutionarily significant lineages that might warrant taxonomic recognition and increased conservation concern, as was recently demonstrated for the previously considered widespread species, *Pelomedusa subrufa* sensu lato, now a constellation of several recognized separate species (Petzold et al. 2014).

The average Indigenous (Historical) Extent of Occurrence (historical EOO; minimum convex polygon hull) for all 486 recognized tortoise and freshwater turtle taxa (species plus subspecies, excluding marine turtles) was 1,151,984 sq. km, ranging from the smallest range of 11 sq. km for the Pinzón Giant Tortoise, *Chelonoidis niger duncanensis*, to the largest range of 24,355,804 sq. km for the Nile Softshell Turtle, *Trionyx triunguis*. Tables 14–15 list the top 50 ranked taxa with the smallest and largest estimated indigenous EOO.

Table 14. Top 50 modern turtle taxa with the smallest Estimated Indigenous (Historical) Extent of Occurrence (EOO) (sq. km); † = extinct.

Taxon	Family	Size
Chelonoidis niger duncanensis	Testudinidae	11
Chelonoidis niger abingdonii †	Testudinidae	19
Chelonoidis niger ssp. Santa Fé †	Testudinidae	19
Chelonoidis niger hoodensis	Testudinidae	22
Chelonoidis niger niger †	Testudinidae	86
Chelonoidis niger donfaustoi	Testudinidae	90
Chelonoidis niger microphyes	Testudinidae	121
Cylindraspis peltastes †	Testudinidae	135
Cylindraspis vosmaeri †	Testudinidae	135
Chelonoidis niger phantasticus	Testudinidae	142
Chelonoidis niger darwini	Testudinidae	183
Chelodina mccordi timorensis	Chelidae	245
Chelonoidis niger becki	Testudinidae	271
Aldabrachelys gigantea daudinii †	Testudinidae	286
Kinosternon vogti	Kinosternidae	306
Chelonoidis niger vandenburghi	Testudinidae	318
Chelonoidis niger porteri	Testudinidae	319
Chelonoidis niger chathamensis	Testudinidae	368
Chelonoidis niger vicina	Testudinidae	402
Aldabrachelys gigantea gigantea	Testudinidae	406
Aldabrachelys gigantea arnoldi	Testudinidae	492
Chelodina mccordi mccordi	Chelidae	623
Chelonoidis niger guntheri	Testudinidae	672
Pelusios williamsi laurenti	Pelomedusidae	732
Kinosternon hirtipes magdalense	Kinosternidae	925
Trachemys stejnegeri malonei	Emydidae	1,055
Terrapene coahuila	Emydidae	1,063
Trachemys taylori	Emydidae	1,138
Myuchelys georgesi	Chelidae	1,277
Kinixys zombensis domerguei	Testudinidae	1,520
Cylindraspis inepta †	Testudinidae	2,090
Cylindraspis triserrata †	Testudinidae	2,090
Apalone spinifera atra	Trionychidae	2,116
Emys orbicularis ingauna	Emydidae	2,139
Kinosternon hirtipes megacephalum †	Kinosternidae	2,142
Pseudemydura umbrina	Chelidae	2,388
Cuora flavomarginata evelynae	Geoemydidae	2,529
Cylindraspis indica †	Testudinidae	2,700
Pelusios subniger parietalis	Pelomedusidae	2,789
Aldabrachelys gigantea hololissa	Testudinidae	2,803
Pelusios castanoides intergularis	Pelomedusidae	2,803
Astrochelys yniphora	Testudinidae	2,886
Kinosternon sonoriense longifemorale	Kinosternidae	3,177
Chelodina kuchlingi	Chelidae	3,231
Emydura macquarii nigra	Chelidae	4,380
Mauremys mutica kami	Geoemydidae	4,910
Emydura gunaleni	Chelidae	5,189
Cuora picturata	Geoemydidae	5,774
Kinosternon hirtipes hirtipes	Kinosternidae	6,507
Pyxis planicauda	Testudinidae	6,605

Conservation Status

We include current IUCN Red List conservation status for all species. The status categorizations listed here are current as of the IUCN 2025 Red List of Threatened Species™, version 2025.1 (www.iucnredlist.org). The TFTSG is the official global IUCN Red List Authority responsible for continuously updating IUCN Red List assessments of all tortoises and freshwater turtles, and this process is handled through multiple consensus-building workshops and consultations.

As many species on the Red List need updating, either because their previous evaluations are more than ten years old, or because of recent conservation status or taxonomic changes, we have also included the results of TFTSG Provisional Red List assessments (through May 2025) to indicate their current provisional status, which should be released on the official IUCN Red List site in the near future. In addition, many species that were determined by the TFTSG to be Least Concern in 1996 were never formally listed (as per IUCN Red List protocol at the time), but the original determinations as prepared at that time are still available and are indicated here.

In addition, we include regulatory status listings on CITES (Convention on International Trade in Endangered Species of Wild Fauna and Flora) Appendices I,

Table 15. Top 50 modern turtle taxa with the largest Estimated Indigenous (Historical) Extent of Occurrence (EOO) (sq. km).

Taxon	Family	Size
Trionyx triunguis	Trionychidae	24,355,804
Pelochelys cantorii	Trionychidae	14,176,406
Kinosternon scorpioides	Kinosternidae	13,294,360
Stigmochelys pardalis	Testudinidae	11,956,566
Phrynops geoffroanus	Chelidae	11,495,510
Chelonoidis carbonarius	Testudinidae	11,429,957
Chelonoidis denticulatus	Testudinidae	10,304,369
Chelydra serpentina	Chelydridae	9,022,386
Emys orbicularis orbicularis	Emydidae	8,873,890
Pelomedusa subrufa	Pelomedusidae	8,493,444
Podocnemis unifilis	Podocnemididae	8,111,508
Cyclanorbis senegalensis	Trionychidae	7,374,057
Platemys platycephala platycephala	Chelidae	7,223,458
Podocnemis expansa	Podocnemididae	7,126,911
Mesoclemmys gibba	Chelidae	7,072,096
Kinixys belliana	Testudinidae	7,015,110
Chelus fimbriata	Chelidae	6,709,315
Centrochelys sulcata	Testudinidae	6,524,249
Pelusios subniger subniger	Pelomedusidae	6,503,796
Cuora couro couro	Geoemydidae	6,391,521
Dogania subplana	Trionychidae	6,040,981
Pelusios castaneus	Pelomedusidae	6,025,930
Pelusios rhodesianus	Pelomedusidae	5,777,726
Rhinoclemmys punctularia punctularia	Geoemydidae	5,735,675
Kinixys erosa	Testudinidae	5,563,008
Kinixys spekii	Testudinidae	5,561,920
Chrysemys picta bellii	Emydidae	5,445,182
Pelusios adansonii	Pelomedusidae	5,350,089
Mauremys leprosa saharica	Geoemydidae	5,343,917
Pelusios gabonensis	Pelomedusidae	5,236,162
Mesoclemmys raniceps	Chelidae	4,860,277
Indotestudo elongata	Testudinidae	4,802,341
Pelusios sinuatus bottegi	Pelomedusidae	4,799,189
Sternotherus odoratus	Kinosternidae	4,565,464
Chitra indica	Trionychidae	4,503,800
Cyclemys dentata	Geoemydidae	4,433,372
Apalone spinifera spinifera	Trionychidae	4,377,847
Cyclanorbis elegans	Trionychidae	4,284,609
Mauremys reevesii	Geoemydidae	4,277,816
Kinixys nogueyi	Testudinidae	4,105,375
Pelusios castanoides castanoides	Pelomedusidae	3,895,232
Peltocephalus dumerilianus	Podocnemididae	3,692,687
Notochelys platynota	Geoemydidae	3,522,719
Nilssonia gangetica	Trionychidae	3,349,942
Pelodiscus sinensis	Trionychidae	3,303,687
Heosemys spinosa	Geoemydidae	3,298,125
Lissemys punctata andersoni	Trionychidae	3,218,279
Terrapene ornata	Emydidae	3,109,956
Mesoclemmys vanderhaegei	Chelidae	3,090,178
Siebenrockiella crassicollis	Geoemydidae	3,036,820

II, or III, current as of the 25 Nov 2024 listing (http://cites.org/eng/app/appendices.php). Finally, we also include regulatory status listings for those species and subspecies that are included on the USA Endangered Species Act (ESA) (https://ecos.fws.gov/ecp/). As such, this document brings together most important aspects of taxonomy, names, distribution, and conservation status of all turtles and tortoises of the world.

STATUS UPDATES AND DISCUSSION

Conservation Status

To assess and summarize the current conservation status of turtles and tortoises in the broadest strokes, we first provide an update and analysis of the most current IUCN Red List (www.iucnredlist.org). Additionally, since some species have not yet been formally assessed for the IUCN, or assessed and submitted, but not yet formally published, for these taxa we provide additional analyses based on their "*in press*" or "*in prep*." or provisional assessments as determined by the IUCN SSC Tortoise and Freshwater Turtle Specialist Group (TFTSG). The official assessments of conservation status of turtles are provided to the IUCN Red List by the TFTSG, which is continuously producing provisional and finalized assessments for unevaluated taxa as well as previously assessed taxa needing updates (should be undertaken every 10 years). Quantifying the overall conservation status and percentage of threatened species of turtles is important in understanding how seriously they are endangered, and how they compare with other imperiled clades. Mittermeier et al. (2015) provided an analysis of global Turtle Hotspots as part of such an evaluation, and Rhodin et al. (2018b) provided a global overview of the conservation status of all turtles and tortoises as assessed at that time on both the IUCN Red List and the TFTSG Provisional Red List. Subsequent work by Ennen et al. (2020, 2021) has analyzed the conservation status of turtles based on hydrobasin distributions.

The current IUCN 2025 Red List (version 2025.1) formally lists 277 turtle and tortoise species (only a 1.1% increase in number of species listed from 274 in 2021), in addition to 9 separate subspecies and 25 regional subpopulations, using a slightly different taxonomy from the one presented in this checklist. Of the species listed, 8 are listed as Extinct (EX) [including *Pelusios seychellensis*, no longer considered valid, instead synonymized under *Pelusios castaneus*, and two species of Galapagos tortoises, considered by us to be subspecies], 68 Critically Endangered (CR) (a 1.5% increase from 67 in 2021), 48 Endangered (EN) (a 4.3% increase from 46 in 2021), 58 Vulnerable (VU) (no change from 2021), 38 Near Threatened (NT) (a 5.6% increase from 36 in 2021), 1 Lower Risk/conservation dependent (LR/cd; an old category being phased out) (no change), 43 Least Concern (LC) (a

Table 16. The world's Top 75 most endangered non-marine turtle taxa in need of conservation action, based on our review and analysis of prior ranked listings by the Turtle Conservation Fund (TCF 2003), the IUCN Tortoise and Freshwater Turtle Specialist Group (TFTSG 2007), the Turtle Conservation Coalition (TCC 2011, 2018), and the latest unranked listing by TCC (2025, in press), plus our own review of IUCN Red List assessments by the TFTSG. The current ranking placement of each taxon in this TTWG 2025 list is subjectively estimated and approximate.

	Species or Subspecies	Family	TCF 2003	TFTSG 2007	TCC 2011	TCC 2018	TCC 2025
1	*Chelonoidis niger phantasticus*	Testudinidae	–	–	–	51-65	1-67
2	*Rafetus swinhoei*	Trionychidae	12	3	2	1	1-67
3	*Cuora zhoui*	Geoemydidae	–	26-100	6	6	1-67
4	*Cuora yunnanensis*	Geoemydidae	–	2	3	3	1-67
5	*Kinosternon hirtipes hirtipes*	Kinosternidae	–	–	–	–	1-67
6	*Cuora mccordi*	Geoemydidae	–	26-100	7	7	1-67
7	*Astrochelys yniphora*	Testudinidae	14	9	10	2	1-67
8	*Cuora aurocapitata*	Geoemydidae	–	18	8	9	1-67
9	*Cuora trifasciata*	Geoemydidae	4	7	9	12	1-67
10	*Kinosternon vogti*	Kinosternidae	–	–	–	–	1-67
11	*Chelodina mccordi mccordi*	Chelidae	3	11	12	14	1-67
12	*Cyclanorbis elegans*	Trionychidae	–	–	–	11	1-67
13	*Chitra chitra*	Trionychidae	2	8	13	15	1-67
14	*Batagur baska*	Geoemydidae	1	12	4	4	1-67
15	*Psammobates geometricus*	Testudinidae	16	22	22	8	1-67
16	*Kinosternon sonoriense longifemorale*	Kinosternidae	–	–	–	–	1-67
17	*Batagur trivittata*	Geoemydidae	9	4	5	5	1-67
18	*Dermatemys mawii*	Dermatemydidae	23	24	15	18	1-67
19	*Mesoclemmys dahli*	Chelidae	21	26-100	–	10	1-67
20	*Erymnochelys madagascariensis*	Podocnemididae	13	16	16	19	1-67
21	*Geochelone platynota*	Testudinidae	6	5	11	13	1-67
22	*Pseudemydura umbrina*	Chelidae	19	17	20	23	1-67
23	*Terrapene coahuila*	Emydidae	–	19	34	37	1-67
24	*Kinosternon hirtipes magdalense*	Kinosternidae	–	–	–	–	1-67
25	*Cuora pani*	Geoemydidae	–	26-100	26	29	1-67
26	*Sternotherus depressus*	Kinosternidae	–	26-100	48	28	1-67
27	*Mauremys annamensis*	Geoemydidae	10	10	14	17	1-67
28	*Pyxis planicauda*	Testudinidae	15	14	30	33	1-67
29	*Ranacephala hogei*	Chelidae	–	26-100	21	24	1-67
30	*Podocnemis lewyana*	Podocnemididae	–	26-100	24	26	1-67
31	*Myuchelys georgesi*	Chelidae	–	–	–	16	1-67
32	*Kinosternon hirtipes chapalaense*	Kinosternidae	–	–	–	–	1-67
33	*Mesoclemmys zuliae*	Chelidae	–	26-100	–	–	1-67
34	*Leucocephalon yuwonoi*	Geoemydidae	7	15	19	22	1-67
35	*Batagur affinis*	Geoemydidae	–	–	17	20	1-67
36	*Batagur kachuga*	Geoemydidae	–	26-100	18	21	1-67
37	*Siebenrockiella leytensis*	Geoemydidae	17	25	23	25	1-67
38	*Batagur borneoensis*	Geoemydidae	8	20	25	27	1-67
39	*Testudo kleinmanni*	Testudinidae	18	13	27	30	1-67
40	*Heosemys depressa*	Geoemydidae	5	6	28	31	1-67
41	*Chersobius solus*	Testudinidae	–	–	43	44	1-67
42	*Chitra vandijki*	Trionychidae	–	26-100	31	34	1-67
43	*Mauremys nigricans*	Geoemydidae	–	26-100	32	35	1-67
44	*Gopherus flavomarginatus*	Testudinidae	–	26-100	39	42	1-67
45	*Malacochersus tornieri*	Testudinidae	–	26-100	–	48	1-67
46	*Kinixys homeana*	Testudinidae	–	–	–	51-65	1-67
47	*Orlitia borneensis*	Geoemydidae	–	26-100	–	51-65	1-67
48	*Astrochelys radiata*	Testudinidae	–	26-100	35	38	1-67
49	*Glyptemys muhlenbergii*	Emydidae	24	26-100	40	43	1-67
50	*Emys orbicularis eiselti*	Emydidae	–	–	–	–	1-67
51	*Chelodina mccordi timorensis*	Chelidae	–	–	–	–	1-67
52	*Chersobius boulengeri*	Testudinidae	–	–	–	–	1-67
53	*Pelodiscus huangshanensis*	Trionychidae	–	–	–	–	1-67
54	*Trachemys taylori*	Emydidae	–	26-100	–	–	1-67
55	*Vijayachelys silvatica*	Geoemydidae	–	26-100	–	–	1-67
56	*Pyxis arachnoides*	Testudinidae	–	26-100	38	41	1-67
57	*Nilssonia nigricans*	Trionychidae	–	26-100	47	46	1-67
58	*Cuora cyclornata*	Geoemydidae	–	–	–	50	1-67
59	*Pelochelys cantorii*	Trionychidae	–	26-100	49	47	1-67
60	*Elusor macrurus*	Chelidae	20	21	41	–	1-67
61	*Manouria emys*	Testudinidae	–	26-100	42	51-65	1-67
62	*Sacalia bealei*	Geoemydidae	–	26-100	–	51-65	1-67
63	*Nilssonia leithii*	Trionychidae	–	26-100	–	51-65	1-67
64	*Platysternon megacephalum*	Platysternidae	–	26-100	–	51-65	1-67
65	*Chersobius signatus*	Testudinidae	11	26-100	–	–	1-67
66	*Indotestudo forstenii*	Testudinidae	–	26-100	–	–	1-67
67	*Staurotypus salvinii*	Kinosternidae	–	–	–	–	1-67
68	*Chitra indica*	Trionychidae	–	23	33	36	–
69	*Cuora bourreti*	Geoemydidae	–	26-100	36	39	–
70	*Nilssonia formosa*	Trionychidae	–	26-100	46	45	–
71	*Palea steindachneri*	Trionychidae	–	26-100	–	–	–
72	*Kinosternon cora*	Kinosternidae	–	–	–	–	–
73	*Mauremys mutica*	Geoemydidae	–	26-100		51-65	–
74	*Apalone spinifera atra*	Trionychidae	–	26-100	–	–	–
75	*Terrapene nelsoni*	Emydidae	–	26-100	–	–	–

10.4% decrease from 48 in 2021), and 13 Data Deficient (DD) (a 30.0% increase from 10 in 2021).

An additional 77 assessments have been finalized by the TFTSG and have been submitted to the Red List and are currently "*in press"* and are expected to be published over the coming year. Another 22 formal Red List assessments are still "*in prep*." with anticipated submission over the next few months. All these assessments are indicated as such in the checklist.

According to IUCN Red List protocol, Threatened species are defined as those in the three categories of Critically Endangered, Endangered, and Vulnerable, meaning that 174 species are officially regarded as Threatened (62.8% of the 277 species listed, a 0.6% increase from the 62.4% of the 274 species listed in 2021), with 116 species (41.9% of all those listed) considered Critically Endangered or Endangered. This represents a 1.7% increase in the species that are currently considered Critically Endangered or Endangered as compared to the 113 species (41.2% of all those listed) assessed as such in 2021.

Of the 364 species recognized as distinct in our checklist, 63 are not yet officially listed on the IUCN Red List. Many of these apparently "unassessed" species have already been evaluated by the TFTSG, first in 1996, when Least Concern (LC) species were not formally listed (as some are now), and then more recently through a series of draft assessments. Of these species, the TFTSG evaluated 53 as Least Concern in 1996 (Behler 1996), and these are marked as such in this checklist, with some of them still needing updating with new Red List assessments.

Further status assessments have been accomplished through an ongoing series of regional IUCN Red Listing workshops held by the TFTSG. These workshops have assessed both previously unevaluated species and updated older previously evaluated species. Since 1999 the TFTSG has held Red Listing workshops in or for Asia, Mexico, the Mediterranean, India, Madagascar, Australia, New Guinea, USA, northern South America, southern South America, the Galápagos Islands, Asia a second time, Sub-Saharan Africa, India a second time, Mexico and Central America again, South America again, and the USA three times more. Although not all assessments are yet official IUCN Red List evaluations, we can use all of these "*in press*" and "*in prep*." and provisional evaluations to determine overall threat rates to all turtles and tortoises. The current assessments that are based on the findings and results of these workshops, as well as on recently published literature, but have not yet been finalized and published on the IUCN Red List, are included in this checklist as TFTSG Provisional Red List status assessments.

Combining the formally published IUCN Red List assessments noted above with all "*in press*" and "*in prep*." and provisional TFTSG status evaluations for previously unlisted species and updated assessments for currently listed but outdated assessments, yields the following total current status numbers for all 364 species of turtles and tortoises (not counting subspecies): 5 Extinct (EX), no change from 2021, 74 Critically Endangered (CR), a 1.4% increase from 73 in 2021, 60 Endangered (EN), a 27.7% increase from 47 in 2021, 62 Vulnerable (VU), a 5.1% increase from 59 in 2021, 45 Near Threatened (NT), a 7.1% increase from 42 in 2021, 89 Least Concern (LC), an 8.2% decrease since 97 in 2021, and 29 Data Deficient (DD), a 6.5% decrease from 31 in 2021. This yields 134 species (36.8%) that are Critically Endangered or Endangered, an 8.9% increase from 123 (34.7%) in 2021, and 196 (53.8%) that are Threatened (Critically Endangered, Endangered, or Vulnerable), a 7.1% increase from 183 (51.3%) in 2021.

We can provisionally adjust these numbers to account for Data Deficient species which may also be Threatened. We follow the calculation method of determining percentage of Threatened species utilized by Hoffmann et al. (2010): the number of Threatened species (196) is divided by the number of data-sufficient species (335), i.e., the total number of species (364) minus those that are Data Deficient (29). This assumes that Data Deficient species will have the same percentage of Threatened species as data-sufficient species. Using this calculation methodology, 58.5% of all assessed data-sufficient turtles and tortoises are Threatened. For comparison, using the same methods, Hoffmann et al. (2010) determined that 41% of amphibians, 33% of cartilaginous fishes, 25% of mammals, and 13% of birds were Threatened. An updated analysis of all Red-Listed species in these vertebrate groups by Rhodin et al. (2018b) demonstrated that turtles were more threatened than birds, lizards, mammals, crocodilians, or amphibians, but less than primates. A more recent review by Cox et al. (2021) continues to demonstrate that turtles are more threatened than birds, mammals, or amphibians, as well as more than lizards, crocodiles, or snakes.

No matter how we analyze these various percentages of Red-Listed Threatened species, turtles and tortoises, with anywhere from ca. 54–63% of their 364 species Threatened, are among the most threatened of the major vertebrate lineages on earth, and surpassed among the larger vertebrate lineages only by the Primates, currently with 66.2% of their 527 Red-Listed species assessed as Threatened, although Caudata (salamanders) have a similar ca. 56–59% of their 758 species Threatened; Crocodylia have ca. 48% of their 23 species Threatened, Anura (frogs and toads) have ca. 35% of their 2,045 species Threatened, and Squamata (snakes and lizards) have only ca. 17% of their 10,015 species Threatened (www.iucnredlist.org/search, accessed May 2025).

Using the methods developed by Stuart and Thorbjarnarson (2003), Rhodin (2006), and Rhodin et al. (2018b), we also calculated the Average Threat Level (ATL) for various groupings of turtles and tortoises with IUCN or TFTSG Red List assessments (including all species and several subspecies). We assigned a numerical threat level per taxon based on current assessments,

with LC = 1, NT = 2, DD = 2.5, VU = 3, EN = 4, CR = 5, CR(PEW) = 5.5, EW = 6, CR(PE) = 7, and EX = 8. Taxa assessed as DD were assigned an estimated value of 2.5 based on the potential predicted average threat level after evaluation (see Hoffmann et al. 2010). We then summed all values and calculated the ATL for each grouping (where ATL equals the total summed threat value per grouping divided by the total number of assessed taxa in that grouping).

The ATL for all Testudines at this time is 3.053, a 2.4% increase since the ATL of 2.982 in 2021, which was a 0.74% increase from the ATL of 2.960 in 2018 (Rhodin et al. 2018b; TTWG 2021). For the most endangered turtle family, the Testudinidae, the ATL at this time is 4.254, a 1.4% increase since the ATL of 4.197 in 2021, which was a 3.37% increase from the ATL of 4.060 in 2018. For the second most threatened grouping, the two combined Asian subfamilies Geoemydinae and Batagurinae, their ATL at this time remains at 3.903, with no change from the ATL of 3.903 in 2021, which was a 0.33% increase from the ATL of 3.890 in 2018. For the other two most threatened families in 2018 and 2021, the Podocnemididae and Cheloniidae, the Podocnemididae have now had their ATL decrease from 3.875 to 3.750 (one species, *Podocnemis expansa*, previously provisionally assessed by the TFTSG as CR has instead now been formally assessed as EN, but is still "*in press*" with the IUCN), and the Cheloniidae remain unchanged with an ATL of 3.750. The next most endangered family at this time is the Trionychidae, which increased 2.18% from an ATL of 3.440 in 2018 to 3.515 in 2021, but has now decreased 4.1% to an ATL of 3.371. The Emydidae currently have an ATL of 2.483, up by 3.0% from the ATL of 2.410 in 2021. The Chelydridae have an ATL of 2.400, up by 4.3% from an ATL of 2.300 in 2021, Rhinoclemmydinae have an ATL of 2.222, an increase of 21.2% from an ATL of 1.833 in 2021. The least threatened family in 2018 and 2021, the Kinosternidae, increased 3.6% from an ATL of 1.650 in 2018 to 1.710 in 2021, and its ATL has now increased dramatically by 29.9% to 2.221 at present, and it is no longer the least threatened turtle family. The Chelidae have an ATL of 2.175, an increase of 1.4% from an ATL of 2.145 in 2021, and the Pelomedusidae, now the least threatened turtle family, has an ATL of 1.813, a decrease of 0.1% from an ATL of 1.815 in 2021.

As part of the process of determining the relative threatened status of the world's tortoises and freshwater turtles, the Turtle Conservation Fund (TCF 2003), the IUCN Tortoise and Freshwater Turtle Specialist Group (TFTSG 2007), and the Turtle Conservation Coalition (TCC 2011, 2018, 2025) have to date published five consensus listings of "*Turtles in Trouble*", which listed the top ca. 25–65 most threatened species in each of those years. These documents have become widely cited, especially as a basis for justifying and supporting conservation grant proposals and action plans. Based on these works and our own assessment, we provide a ranked listing (Table 16) of what we currently consider to be the 75 most endangered tortoise and freshwater turtle taxa in need of conservation action.

Continuing to evaluate further changes to these "*Turtles in Trouble*" compilations, both in terms of improved status for those species benefitting from conservation efforts, and documenting potentially deteriorating survival prospects for other species, will be critically important for future conservation efforts for these highly threatened species. This continuing process is undertaken by the TFTSG in collaboration with other turtle conservation organizations, notably Turtle Conservancy, Turtle Survival Alliance, Chelonian Research Foundation, and Re:wild.

Increasing support for conservation efforts targeting endangered freshwater turtles and tortoises has been generated over the years through separate and joint efforts of all of the above organizations and others. A welcome groundswell of turtle conservation work has risen to meet the challenge of increasing threat levels to native populations of freshwater turtles and tortoises. Helping to provide leadership in these efforts has been: 1) the Turtle Conservation Fund (TCF 2002, 2019) that from 2002 to date has provided over US$ 1.62 million in grants to 379 deserving projects in 66 countries focused on urgently needed conservation efforts for endangered tortoises and freshwater turtles, and 2) the Mohamed bin Zayed Species Conservation Fund (Rhodin et al. 2018a) that from 2009 to date has provided over US$ 1.88 million to 179 deserving projects in 48 countries.

In addition, there has been fairly robust support for conservation work focused on marine turtles over the years, especially from the Marine Turtle Conservation Fund administered by the U.S. Fish and Wildlife Service (USFWS). In 2021, the USFWS initiated a major expansion of its program to also include support for international conservation work on freshwater turtles and tortoises—a most welcome and critically needed expansion of support. In addition, USFWS has supported several turtle Red Listing workshops in North and South America over the last 2–3 years. Unfortunately, further USFWS support is currently on hold under the present USA political administration, but we hope and anticipate that it will be reinstated in the future.

Genetic Pollution

Aside from overt and highly impactful conservation threats such as overexploitation and habitat destruction, the global turtle fauna is also increasingly facing another significant threat: genetic pollution caused by human-facilitated hybridization and introgression from introduced and invasive species (Rhymer and Simberloff 1996; Simison et al. 2013; Spencer et al. 2014; García-Díaz et al. 2015; Capinha et al. 2017; Nori et al. 2017; Georges et al. 2018b; Reshetnikov et al. 2023).

This is not entirely new, but the current extent is unprecedented. Some taxa have historically already been affected.

This is most evident for the American Red-eared Slider, *Trachemys scripta elegans*, considered the most invasive of all turtles (Capinha et al. 2017; Reshetnikov et al. 2023), having been introduced to or recorded from at least 102 countries outside its native range in the United States and northern Mexico. It is considered to be among the "Top 100" most invasive species worldwide (https://www.iucngisd.org/gisd/100_worst.php) and hybridizes readily with other *Trachemys* species in the many North and South American countries that harbor native populations of *Trachemys* spp.

This is also true for the Asian softshell turtles of the genus *Pelodiscus*. These turtles have been farmed and traded for centuries, with the corollary of translocating different species and local genetic lineages, leading to the admixture of different taxa and lineages in farms and in the wild (Fritz et al. 2010b; Suzuki and Hikida 2014; Gong et al. 2018). Similarly, the historical introduction of *Mauremys reevesii* to Japan resulted in massive hybridization with the native *M. japonica* (Suzuki et al. 2014). Another historical case of human-mediated admixture of genetic lineages is known from European pond turtles (*Emys orbicularis*). Here, the non-native populations on the Balearic Islands, most probably introduced in Roman times (Valenzuela et al. 2016), are of admixed origin (Lenk et al. 1999). Another population with genetic signatures of an old or ancient introduction of *E. o. hellenica* was discovered near Rome (Lenk et al. 1999; Vamberger et al. 2015) within the range of another subspecies (*E. o. galloitalica*).

However, unlike in historical times, when only a few turtle species were affected, genetic pollution has become a major issue for nature conservation in recent years, facilitated by the massive pet and food trade and increased human mobility. Today, genetic pollution is also caused by well-meaning augmentation of endangered local turtle populations with genetically mismatched individuals (typically, but not exclusively, from non-coordinated actions by turtle enthusiasts), the release of surplus or abandoned genetically divergent pet turtles, and also by large-scale releases of confiscated turtle shipments, especially in Southeast Asia.

Examples of restocking with mismatched genetic individuals include endangered populations of *E. orbicularis* at the northern edge of its range (Fritz et al. 2004; Velo-Antón et al. 2011: genetic evidence for restocking with several different subspecies), and in southern France (Vamberger et al. 2015; Raemy et al. 2017: restocking with non-native *E. o. hellenica* instead of native *E. o. galloitalica*), also northern edge populations of *Mauremys leprosa* in southern France (Palacios et al. 2015: restocking with *M. l. saharica* and northern African *M. l. leprosa* instead of European *M. l. leprosa*), of *M. rivulata* in Croatia (Vamberger et al. 2014: restocking with Cretan individuals), and of *Testudo graeca* in Doñana National Park in Spain (Graciá et al. 2017b: restocking with non-native *T. g. marokkensis* from Morocco instead of native *T. g. whitei* from Spain).

Examples of genetic pollution caused by abandoned pet turtles include *Chrysemys picta bellii* from British Columbia introgressed by non-native subspecies (Jensen et al. 2014b) and Antillean *Trachemys* introgressed by Red-eared Sliders (*Trachemys scripta elegans*; Parham et al. 2013). Also some of the above-mentioned cases for European Pond Turtles refer at least partially to genetic pollution by abandoned pet turtles. Hybridization in the wild from released trade animals has been recorded in Taiwan for *Mauremys reevesii* and *M. sinensis* (Fong and Chen 2010).

This issue of potentially increasing genetic pollution needs to be kept in mind as conservationists devise management plans designed to reinforce or restore dwindling or extirpated populations of turtles impacted by overexploitation and habitat loss. The need to maintain well-defined and relatively pure non-hybrid genetic lineages, subspecies, and species is important and needs to be kept in focus. However, also maintaining increased heterozygosity of non-hybrid transplanted populations is also important for improved survival (Scott et al. 2020).

Request for Updates

Please help the TTWG and the TFTSG keep this Turtles of the World Checklist and Atlas up-to-date by e-mailing us (addresses noted above) and including pdf's of any relevant articles about new taxonomic or distributional information and/or revisions that should be included and annotated here in upcoming checklists, whether you are an author on a paper providing updated information, or have become aware of data that you believe should be included. Also please inform us of any errors or discrepancies in any of our data, especially for geographic distributions in countries or states, and for cited references and names, so that we may update or correct them as necessary. For sea turtle distribution data, please submit additions and corrections via the SWOT website. We want this checklist and atlas to be as accurate, up-to-date, and comprehensive as possible, and ask for your assistance to help us accomplish this goal.

The maps published here all represent work in progress, and will continue to be updated and revised in future checklists as we acquire new and improved locality data. We strongly encourage and welcome our readers and professional colleagues, especially field-based turtle specialists and other enthusiasts, to inform us about proposed corrections and changes to these maps, and to submit specific locality data on the presence or absence of species in various locations for consideration of incorporation into the next checklist.

We are especially interested in receiving new point locality data from presumed range areas not well represented by locality points on our maps, especially from gap areas in the presumed range, as well as verified range extensions. Please help us improve and update these maps by submitting any new locality information along with geographic coordinates and the associated citations to Rhodin at rhodincrf@aol.com.

Acknowledgments

We extend special thanks to James Parham, a major contributor and co-author on earlier versions of the checklist, and a source of incisive initial input and analytic debate. Special thanks also for extensive detailed input from Holger Vetter, Peter Uetz, and Scott Thomson, as well as Peter Pritchard and Roger Bour, close colleagues and collaborators whom we honored with a posthumous dedication in our previous edition.

We also thank various members of the IUCN SSC Tortoise and Freshwater Turtle Specialist Group (TFTSG) as well as many other individuals who have provided pertinent data or references or localities for maps, or helped in various ways with informational input or review of this and earlier checklists, or help with etymologies and eponyms: Rafael Acuña-Mesén, Kraig Adler, Erik Åhlander, Leandro Alcalde, Natalia Ananjeva, Matthew Aresco, Omar Attum, Mark Auliya, Roy Averill-Murray, Ernst Baard, Sherif Baha El Din, Albert Bertolero, John Bickham, Torsten Blanck, Bill Branch, Ronald Brooks, Ortwin Bourquin, James Buskirk, Joseph Butler, Gisella Caccone, Patrick Campbell, María Julia Cassano, Linda Cayot, Eng Heng Chan, Ylenia Chiari, Laurent Chirio, Werner Conradie, Jago Cooper, Henrique Caldeira Costa, Marcelo de la Fuente, Jeffrey Dawson, Tomas Diagne, Andrew DiMateo, David Emmett, Taylor Edwards, Alejandro Fallabrino, Balazs Farkas, Camila Ferrara, Bruno Ferronato, Oscar Flores-Villela, Natàlia Rizzo Friol, Eugene Gaffney, Justin Gerlach, Hanyeh Ghaffari, James Gibbs, Scott Gillingwater, Shiping Gong, Gracia Gonzalez, Eric Goode, Steve Gotte, Michael Granatosky, Ute Grimm, Cris Hagen, Bonnie Malone Hall, Brian Henen, Jerry Herman, Scott Hillard, Craig Hilton-Taylor, Judith Hirt, Margaretha Hofmeyr, Marinus Hoogmoed, Brian Horne, Flora Ihlow, Sixto Inchaustegui, Djoko Iskandar, Dale Jackson, Fredric Janzen, Robert Jones, Mehdi Joseph-Ouni, Alice Karl, Ross Kiester, Kenneth Krysko, Gerald Kuchling, Minh Le, Thomas Leuteritz, Colin Limpus, Peter Lindeman, Jeffrey Lovich, Luca Luiselli, Roderic Mast, Erica Mejlon, Fabrice Méral, Peter Mikulíček, David Mifsud, Russell Mittermeier, Patrick Moldowan, Robert Murphy, Simon Nemtzov, Dario Ottonello, Vivian Páez, Olivier Pauwels, Ayudha Bahana Perdamaian, Fabio Petrozzi, Hans-Dieter Philippen, Nicolas Pilcher, Håkan Pohlstrand, Peter Praschag, Peter Pritchard, Awal Riyanto, Willem Roosenburg, José Rosado, Jose Vicente Rueda-Almonacid, Raymond Saumure, Chuck Schaffer, Jason Schaffer, Hans-Hermann Schleich, Sabine Schoppe, Norman Scott, Peter Scott, Will Selman, Haitao Shi, Franco Souza, Phil Spinks, Bryan Stuart, Travis Thomas, Robert Thomson, Oguz Türkozan, Melita Vamberger, Richard Vogt, Bruce Weissgold, Chris Wiese, and Reza Yadollahvand.

Distributional data on sea turtles from the IUCN SSC Marine Turtle Specialist Group (MTSG), SWOT (The State of the World's Sea Turtles), and Seminoff et al. (2015) were gratefully received from Bryan Wallace, Jeffrey Seminoff, and Kyle Van Houtan, with updates from the SWOT website in 2024.

The Panaphil Foundation provided partial funding support for PPvD. TFTSG-organized IUCN Red Listing workshops and other redlisting efforts have been generously supported by the Frankel Family Foundation, Conservation International, Turtle Conservancy, Moore Family Foundation, George Meyer and Maria Semple, Matt Frankel, Turtle Conservation Fund, Chelonian Research Foundation, Galapagos Conservancy, U.S. Fish and Wildlife Service, Andrew Sabin Family Foundation, and Mohamed bin Zayed Species Conservation Fund.

Initial map production and GIS-analysis by the TFTSG and Chelonian Research Foundation was supported by the BioFresh Project of the IUCN Freshwater Biodiversity Unit, IUCN Species Programme, and Forschungsverbund Berlin e.V., Leibniz-Institut für Gewässerökologie und Binnenfischerei. We thank Kevin Smith, Savrina Carrizo, Will Darwall, Ackbar Joolia, Marcelo Tognelli, Adrian Hughes, Diego Juffe, and Eresha Fernando of IUCN, as well as Melita Vamberger, for welcome technical assistance and helpful suggestions in our production of maps and use of the Esri ArcMap and ArcPro software applications. We thank Sarah C. Sweat of the Tennessee Aquarium Conservation Institute and Nyssa R. Hunt and Tegan Childers of the University of Tennessee Chattanooga for invaluable help in the production of the color-ramped species richness maps in the previous checklist, and Nyssa Hunt for producing the updated species richness maps in the current checklist. We thank Bryan Wallace, Rod Mast, and Ei Fujioka of SWOT and Duke University for help with sea turtle maps.

Ongoing support for map production and checklist preparation and publication has come from Chelonian Research Foundation (CRF) and Turtle Conservancy (TC). Publication costs of previous checklists have been shared by CRF, TC, TFTSG, Turtle Conservation Fund, Re:wild (formerly Global Wildlife Conservation), Surprise Spring Foundation, Senckenberg Gesellschaft für Naturforschung, Turtle Taxonomy Fund, and John B. Iverson. Printing of the current checklist has been supported by Chelonian Research Foundation, Turtle Conservancy, IUCN SSC Tortoise and Freshwater Tutle Specialist Group, Turtle Conservation Fund, Turtle Taxonomy Fund, and Re:wild.

New and updated locality and unpublished distributional data and suggested edits of our maps have been

submitted by many people over the years, including the authors themselves and: Jeff Briggler, Henrik Bringsøe, James Buskirk, John Carr, Vinícius T. de Carvalho, María Julia Cassano, Henrique Caldeira Costa, Fábio A.G. Cunha, Enrique Yure Domic Rivadeneira, Mike Dreslik, Wendy Estes-Zumpf, Vincenzo Ferri, Carl Franklin, Kathy Gipe, Jim Godwin, Pedro Gómez Murillo, Shiping Gong, Luke Groff, Carol Hall, Marinus Hoogmoed, Mark Howery, Brian Hubbs, Daniel Jablonski, Hossein Javanbakht, Ethan Kessler, Yurii V. Kornilev, Benjamin Leprince, Peter Lindeman, Jeffrey Lovich, Luca Luiselli, Jacob Marlin, Bryce Maxell, Håkan Pohlstrand, Georgi Popgeorgiev, Eduardo Reyes-Grajales, Viviana Ricardez, Stephen Richards, Phil Rosen, Rocio Sanchez, Pavel Široký, Travis W. Taggart, Travis Thomas, Stanley Trauth, Richard Vogt, Kai Wang, and Katherine Young-Valencia. We gratefully acknowledge their valuable contributions and apologize for any possible oversights.

Previously unpublished measurement data on maximum sizes of various turtle and tortoise taxa have been provided to us by the following people and organizations, and included and cited in the checklist as appropriate: Rafael Acuña-Mesén, Arizona Game and Fish Dept., Roy Averill-Murray, James Berry, Torsten Blanck, Brian Bock, Michael Bogan, Andrew Brinker, Pelf Nyok Chen, Ylenia Chiari, Andrew Coleman, David Floyd, Peter Floyd, Scott Floyd, Carl Franklin, Galapagos Conservancy, Galapagos National Park Directorate, Justin Gerlach, Hanyeh Ghaffari, Giant Tortoise Restoration Initiative, James Gibbs, Anna N. Gnetneva, Miguel Angel Grageda García, Cris Hagen, Philip Hall, Scott Hillard, Rick Hudson, Gerald Johnston, Michael Jones, Alice Karl, Ross Kiester, Michael Knoerr, Gerald Kuchling, John Legler, Colin Limpus, Luca Luiselli, Tom Mann, Pearson McGovern, Brian Mealey, Russell Mittermeier, Don Moll, Eric Munscher, Hidetoshi Ota, Dario Ottonello, Vivian Páez, Fabio Petrozzi, Thong Van Pham, Ny Aina Rakotoarisoa, José Rosado, Elizabeth Schwartz, Peter Scott, Gabriel Segniagbeto, Richard Seigel, Will Selman, Washington Tapia, Ertan Taskavak, Turner Endangered Species Fund, Ryan Walker, Chris Wiese, Lisabeth Willey, and Marco Zuffi. We gratefully acknowledge their valuable contributions.

Photos and Photographers

We also most gratefully acknowledge and very much appreciate the use of 1437 turtle and tortoise photos by the following 338 photographers that enhance this checklist. We include both previously unpublished and republished photos from previous CRF publications, with each acknowledged and (if republished) identified as to its original source: *Conservation Biology of Freshwater Turtles and Tortoises* species accounts (CBFTT), *Chelonian Conservation and Biology* journal articles (CCB), *Chelonian Research Monographs* books (CRM 1–6), *Turtle and Tortoise Newsletter* (TTN), *TurtleLog* (TL), *Turtle Conservation Fund* 2002 prospectus (TCF), and the *Turtle Conservation Coalition* 2011 and 2018 Top 25+ documents (TCC). We strive to use primarily photos from natural native settings and to include locality data when available, although some photos are from captivity or the trade.

We include photos of 492 (99.8%) of all 493 recognized modern taxa. We lack photos of only one modern taxon: *Testudo horsfieldii kuznetzovi* (Testudinidae). We urge those of you who may have photos of any of this missing taxon to submit them for consideration of inclusion in the next checklist. We are also always looking for better quality photos with locality data for those taxa we have already illustrated; if you have photos that may offer such improvement, please submit them for consideration.

Contributing Photographers. — We acknowledge all photographers for the use of their photos, notably the following who contributed 10 or more photos each: John B. Iverson (126), Anders G.J. Rhodin (80), Torsten Blanck (35), Peter Paul van Dijk (33), Russell A. Mittermeier (31), Gerald Kuchling (28), Peter Praschag (26), Arthur Georges (25), Flora Ihlow (22), Victor J.T. Loehr (22), Roger Bour (21), Uwe Fritz (21), Indraneil Das (20), Eduardo Reyes-Grajales (20), John L. Carr (17), Tomas Diagne (17), James H. Harding (17), Jérôme Maran (16), Andreas Nöllert (15), Peter C.H. Pritchard (15), Eric V. Goode (13), Sabine Schoppe (13), Richard D. Bartlett (12), James C. Godwin (12), Tomáš Mazuch (12), Edward O. Moll (12), Justin Gerlach (11), Peter V. Lindeman (11), Craig B. Stanford (11), Richard C. Vogt (11), Kurt A. Buhlmann (10), and Luca Luiselli (10).

We also thank all the other photographers whose contributions are equally appreciated: Collette Adams, Guta Agostini, Thomas S.B. Akre (2), Nancy Albury (6), Leandro Alcalde (3), Phil Allman, Bhaba Amatya, Ben Anders (2), Archive of the Museum of Zoology Senckenberg Dresden, Marine Arakelyan, Matthew Aresco (2), Harald Artner (7), Benjamin K. Atkinson, Mark Auliya (4), Roy Averill-Murray, Cesar Ayres (2), Salman Baloch, Chittaranjan Baruah (5), Aaron S. Baxter, Virginia C.D. Bernardes, Rafael Bernhard, James F. Berry, Kristin H. Berry (2), Albert Bertolero (5), Subramanian Bhupathy, Michael S. Bogle, Ryan M. Bolton, Douglas B. Booher, Nik Borrow, Jason R. Bourque (6), Deborah Bower (2), Richard C. Boycott (7), William R. Branch (6), Raissa Bressan, Elizângela S. Brito, Grover J. Brown, Rafe M. Brown, Grégory Bulté (2), Marius Burger, Russell L. Burke (2), R. Bruce Bury (2), James R. Buskirk (6), Alejandra Cadavid (2), Matt Cage, Young Cage, Eduardo Campos, John Cann (9), Vinícius T. de Carvalho (7), Gamaliel Castañeda Gaytán (2), Claudia Ceballos, Luis Ceriaco, Eng Heng Chan, Joyee Chan (2), Tien-Hsi Chen (5), Laurent Chirio (3), Viacheslav Chkhikvadze (2), B.C. Choudhury, Andrew T. Coleman, Marilyn Connell (4), Paul Crow, Stefania D'Angelo, Neil D'Cruze (2), Fábio A.G. Cunha

(4), Shekar Dattatri, Jeffrey E. Dawson (4), Tui De Roy/Roving Tortoise Photos (2), Anslem de Silva (6), Veerappan Deepak (6), Carlos del Valle, David Dennis (2), Mathew Denton, Frank Deschandol (3), Bernard Devaux (2), Christo Deysel, Larry Ditto (2), James L. Dobie, C. Kenneth Dodd, Jr. (7), Clifford Dorse, Charlotte Ducotterd (2), Marc Dupuis-Desormeaux (2), L.M. Ehrhart, Carla C. Eisemberg, Johannes Els, David Emmett (2), Kevin Enge, Carl H. Ernst, Mark Feldman, Pablo Feliz (2), Camila Ferrara, Vincenzo Ferri (4), Darren Fielder (3), German Forero-Medina (4), Manuel Merchán Fornelino, Raúl Fournier (2), Alastair Freeman (5), Jennifer Frey, Galápagos Conservancy (2), Carlos A. Galvis-Rizo, Georg Gassner (2), Maren Gaulke (4), Saurav Gawan, Sebastian Gehring, David J. Germano, Hanyeh Ghaffari (2), Rupali Ghosh, Paul M. Gibbons, Scott D. Gillingwater (4), Alan Giraldo, Shiping Gong (8), Yan-An Gong (3), Gracia Gonzalez-Porter (2), Robert H. Goodman, Jr., Terry E. Graham, Martin Grimm, Hedelvy J. Guada, Danny Gunalen, Joko Guntoro (2), Václav Gvoždík (8), Wulf Haacke (2), Cris Hagen (6), Adrian Hailey, Norbert Halasz, Philip M. Hall, William M. Hammond, Kristen M. Hart, James Harvey (5), George L. Heinrich (4), Douglas B. Hendrie (3), Kelly Herbinson, Richard Herren, Terry Hibbitts, Pierson Hill (2), Kate Hodges (2), Margaretha D. Hofmeyr (3), Dan C. Holland, Marinus S. Hoogmoed (3), Brian D. Horne (4), Janet Hostetter, Mian Hou, Rick Hudson (3), Bonggi R. Ibarrondo (2), Alexander A. Inozemtsev, Iriomote Wildlife Conservation Center, Dale R. Jackson, S. Jayakumar (3), John Jensen, Carlos Alberto Jimenez, Chris Johnson, Michael T. Jones (7), Robert L. Jones (5), Nobuhiro Kawazoe (4), Matthew G. Keevil, Rod Kennett (3), Rebekah Kim, Michael Lau, Dayoung Lee, Edgar Lehr (6), Chris Leone (2), Francesco Luigi Leonetti, Thomas E.J. Leuteritz, Romain Levasseur, Colin J. Limpus, Duncan Limpus (4), Marco A. López-Luna (3), Jeffrey E. Lovich, Gary Luciano, Rodrigo Macip-Ríos (3), William E. Magnusson, Kevin Main, Thomas M. Mann, Barry Mansell (4), Georgia Mantziou, Maximilian Maurer (3), Peter May, William P. McCord (4), Timothy E.M. McCormack (3), Suzanne E. McGaugh, Pearson McGovern (8), Brian K. Mealey, Khaled Merabet, Melvin Mérida, Sébastien Métrailler (2), Peter A. Meylan, Joseph C. Mitchell, Asghar Mobaraki (6), Mónica A. Morales-Betancourt (3), Jiří Moravec, Enrico Morawa, Robert H. Mount, Eric Munscher, Rahul Naik, Volker Naths, Nguyen Thanh Luan, Matthew L. Niemiller, Sergio Nolasco-Perez, Michael Ogle (2), Annette Olsson, Ronald Orenstein (2), Dario Ottonello, Vivian P. Páez (3), Charles W. Painter, James F. Parham (5), Fred Parker (6), Olivier S.G. Pauwels, Miguel Pedrono, John Pemberton, Jarmo Perälä, Thong Van Pham (4), Hans-Dieter Philippen (3), Pino Piccardo, Carmine Pilcher (2), Nicolas J. Pilcher, Kalyar Platt (3), Håkan Pohlstrand, John R. Polisar, Hynek Prokop (2), Basem Rabia (3), Michael Redmer, Renae Reed (2), Viviana Ricardez (7), Maurice Rodrigues (4), John Roe, Fernando J.M. Rojas-Runjaic, Philip C. Rosen (7), Simon Rouot (3), José Vicente Rueda-Almonacid (2), Amtyaz Safi (4), Chris Sagebiel (3), Carolina Sánchez Arias (2), Daniel O. Santana (3), Dian Sartika, Joel Sartore (2), Chuck Schaffer (3), Jason Schaffer (5), Brendan Schembri, Hermann Schleich (2), Alfred Schleicher, Jeffrey R. Schmid, Fabian Schmidt (3), Norbert Schneeweiss, Christoph Schneider (3), Willi Schneider (6), Alice Schumacher (2), Elizabeth R. Schwartz, Peter A. Scott, Gabriel H. Segniagbeto, Michael E. Seidel, Will Selman (4), H. Bradley Shaffer (2), Haitao Shi (9), Shailendra Singh (6), Pavel Široký (3), Shashwat Sirsi, Paco Soberón, Sitha Som, Franco L. Souza (3), Phil Spark (2), Kai Squires, Nathanael Stanek (4), Brett Stearns (2), Bev Steveson, Richard Struijk, James N. Stuart (3), Yik-Hei Sung (2), Emerson Sy, Chris Tabaka, Hiromi Tada, Akio Takahashi (3), Washington Tapia (3), Travis Thomas (2), Robert C. Thomson, Stanley E. Trauth (3), Claire Treilibs, Apostolis Trichas (2), Ashutosh Tripathi, Trustees of the Natural History Museum London (3), Anton Tucker (2), Oguz Türkozan, Dennis W. Uhrig (2), United States Geological Survey (2), Melita Vamberger (6), Robert P. Van Dam, Carla Van Ness, Juliette Velosoa, Miguel Vences (3), Sabine Vinke (4), Thomas Vinke (4), Raju Vyas, Timothy Walsh (2), Joseph P. Ward (2), Annett Werner, Nikhil Whitaker, Blair Witherington (7), Dawn Wilson, Win Ko Ko, Yuichirou Yasukawa (3), Katherine Young-Valencia (2), Vyacheslav Yusupov, Ting Zhou, Cassiano Zaparoli (2), Robert T. Zappalorti (2), Brian Zarate (2), Thomas Ziegler (3), Anders Zimny (2), Stephen M. Zozaya (2), and George R. Zug.

Hierarchical Linnaean Classification of Suprageneric Categories of Modern Turtles (Order, Suborder, Infraorder, Superfamily, Family, Subfamily) Used in this Checklist, with Numbers of Included Families, Genera, Species, and Taxa

	Families	Genera	Species	Taxa
Order Testudines	**14**	**97**	**364**	**493**
••Suborder Pleurodira	3	20	95	106
__Superfamily Cheloidea	1	15	60	66
____Family Chelidae	1	15	60	66
_____Subfamily Chelinae		7	22	23
_____Subfamily Hydromedusinae		1	2	2
_____Subfamily Chelodininae		1	15	16
_____Subfamily Emydurinae		5	20	24
_____Subfamily Pseudemydurinae		1	1	1
__Superfamily Pelomedusoidea	2	5	35	40
____Family Pelomedusidae	1	2	27	32
____Family Podocnemididae	1	3	8	8
_____Subfamily Erymnochelyinae		1	1	1
_____Subfamily Peltocephalinae		1	1	1
_____Subfamily Podocnemidinae		1	6	6
••Suborder Cryptodira	11	77	269	387
•Infraorder Durocryptodira	9	63	233	339
__Superfamily Chelonioidea	2	6	7	7
____Family Cheloniidae	1	5	6	6
_____Subfamily Carettinae		3	4	4
_____Subfamily Cheloniinae		2	2	2
____Family Dermochelyidae	1	1	1	1
__Superfamily Chelydroidea	3	7	40	48
____Family Chelydridae	1	2	5	5
____Family Dermatemydidae	1	1	1	1
____Family Kinosternidae	1	4	34	42
_____Subfamily Kinosterninae		2	31	39
_____Subfamily Staurotypinae		2	3	3
__Superfamily Testudinoidea	4	50	186	284
____Family Emydidae	1	12	57	91
_____Subfamily Deirochelyinae		6	42	67
_____Subfamily Emydinae		6	15	24
____Family Platysternidae	1	1	1	3
____Family Geoemydidae	1	19	75	99
_____Subfamily Batagurinae		8	20	24
_____Subfamily Geoemydinae		10	46	61
_____Subfamily Rhinoclemmydinae		1	9	14
____Family Testudinidae	1	18	53	91
_____Subfamily Manouriinae		1	2	3
_____Subfamily Xerobatinae		1	6	6
_____Subfamily Testudininae		16	45	82
•Infraorder Trionychia	2	14	36	48
__Superfamily Trionychoidea	2	14	36	48
____Family Carettochelyidae	1	1	1	1
____Family Trionychidae	1	13	35	47
_____Subfamily Cyclanorbinae		3	7	9
_____Subfamily Trionychinae		10	28	38

CHECKLIST

Modern Turtle and Tortoise Taxa Extant Since 1500 CE [1,2,3]*

Order Testudines Batsch 1788 (07:1, 10:4, 12:6, 17:1, 21:3, 21:4) (4)*

Turtles, Tortoises, and Terrapins; Chelonians

(includes 14 families, 97 genera, 364 species, 493 taxa)

Testudinata Klein 1751:96 (invalid pre-1758 name)

Testudines Linnaeus 1758:194 (invalid vernacular name)

Testudinata Klein *in* Behn 1760:tab.gen. (*nomen oblitum*) (21:4) (4)*

Testudines Batsch 1788:437

Testudinea Batsch 1796:179

Cheloniens Brongniart 1800a:196 (invalid vernacular name)

Chelonii Latreille 1800:xi

Chelonia Ross and Macartney 1802:tab.iii

Cataphractae Link 1807:51

Testudinata Oppel 1811:3

Perostia Rafinesque 1814:66

Cataphracta Hemprich 1820:101

Chelonea Fleming 1822:268

Fornicata Haworth 1825:373

Chelynae Wagler 1828:861

Sterrichrotes Ritgen 1828:269

Chelonites Burmeister 1837:730

Chelonides Swainson 1839:112

Tylopoda Mayer 1849:197

Testudina Fry 1850:21

Chersemydes Strauch 1862:16

Rhynchochelones Dollo 1886:79

Cheloniae Hoffmann 1890:372


Testudoformes Chang 1957:50

Chelonomorpha Kuhn 1960:30

Casichelydia Gaffney 1975:4

Testudinomorpha Laurin and Reisz 1995:197

Pantestudines Joyce, Parham, and Gauthier 2004:996



Testudines Species Richness

Suborder Pleurodira Cope 1864 (08:20, 21:8, 21:9)

Side-necked Turtles


(includes 3 families, 20 genera, 95 species, 106 taxa)

Pleurodères Duméril and Bibron 1834:354 (invalid vernacular name)

Pleurodera Lichtenstein and von Martens 1856:2 (08:20, 21:8)

Pleurodera Cope 1864:181

Pleurodira Cope 1865:186



Pleurodira Species Richness

Superfamily Cheloidea Gray 1825 (21:3) (5)*

Cheloid Side-necked Turtles

(includes 1 family, 15 genera, 60 species, 66 taxa)

Chelidina Gray 1825:211

Chelydoidea Fitzinger 1826:7

Chelyoidea Baur 1893a:212

Cheloides Gaffney, Tong, and Meylan 2006:33

Cheloidea Rhodin, Iverson, Bour, Fritz, Georges, Shaffer, and van Dijk [Turtle Taxonomy Working Group] 2021:28

Chelidoidea Gable, Byars, Literman, and Tollis 2022:1333 (5)*

Family Chelidae Gray 1825 (12:39, 17:86, 21:3, 21:9)

Chelid Side-necked Turtles

(includes 5 subfamilies, 15 genera, 60 species, 66 taxa)

Chelides Cuvier 1816:14 (invalid vernacular name)

Chelydes Schmid 1819:17 (*nomen oblitum*)

Chelidina Gray 1825:211

Chelydoidea Fitzinger 1826:7

Chelydae Gray 1831d:7

Chelydidae Gray 1831d:37

Chelina Bonaparte 1831:63

Hydraspidina Bonaparte 1836:3 (*partim*)

Hydraspididae Agassiz 1857a:249

Chelydidi Portis 1890:17

Chelyidae Baur 1893a:211

Chelodinidae Baur 1893a:211

Hydromedusidae Baur 1893a:211

Rhinemydidae Baur 1893a:212

Chelidae Lindholm 1929:289

Cheluidae Storr 1978:303 (17:86)



Chelidae Species Richness

Subfamily Chelinae Gray 1825 (12:39)

South American Side-necked Turtles

(includes 7 genera, 22 species, 23 taxa)

Chelides Cuvier 1816:14 (invalid vernacular name)

Chelydes Schmid 1819:17 (*nomen oblitum*)

Chelidina Gray 1825:211

Chelina Bonaparte 1831:63

Chelidinae Georges, Birrell, Saint, McCord, and Donnellan 1998:235

Chelinae van Dijk, Iverson, Shaffer, Bour, and Rhodin [Turtle Taxonomy Working Group] 2012:289

Acanthochelys Gray 1873c (14:39)

Side-necked Swamp Turtles

(includes 4 species, 4 taxa)

Acanthochelys Gray 1873c:305

Type species: *Acanthochelys spixii* [= *Platemys spixii* Duméril and Bibron 1835], by original monotypy.

Etymology:* Greek, *akantha*, thorn or prickle, and *chelys*, turtle; referring to the spiny, tuberculate neck skin.

Acanthochelys macrocephala (Rhodin, Mittermeier, and McMorris 1984) (07:84)

Pantanal Swamp Turtle, Big-headed Pantanal Swamp Turtle




Thomas and Sabine Vinke / CBFTT / nr. Filadelfia, Chaco, Paraguay





left: Sébastien Métrailler / CBFTT / Chaco, Paraguay
right: Anders G.J. Rhodin / CBFTT / Bolivia / megacephalic female



** (orange dot = probable trade; stars = type localities)

Distribution: Bolivia (*Santa Cruz*), Brazil (*Mato Grosso*, *Mato Grosso do Sul*), Paraguay

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 304,133 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 434,972 sq. km

Size (Max SCL):* male 28.3 cm, female 29.5 cm (Rhodin et al. 2009 CBFTT; Brito et al. 2020)

CBFTT Account: Rhodin, Métrailler, Vinke, Vinke, Artner, and Mittermeier (2009)

IUCN Red List: Near Threatened (NT A4c) (Rhodin, Vinke, Métrailler, and Mittermeier 2018). Previously: Near Threatened (NT) (TFTSG 1996); the synonymized taxon *Acanthochelys chacoensis* listed as Vulnerable (VU) (TFTSG *in* Baillie and Groombridge 1996)

Synonymy:

Phrynops schoepffii Fitzinger *in* Siebenrock 1904b:27 (*partim*, *nomen nudum et novum*)

Etymology:* Eponym honoring Johann David Schoepff (or Schöpf) (1752–1800), Germany; physician, botanist, herpetologist, and turtle taxonomist.

Comment:* Name synonymized *in partim* under *Acanthochelys macrocephala* and *Mesoclemmys vanderhaegei*.

Platemys macrocephala Rhodin, Mittermeier, and McMorris 1984a:38

Acanthochelys macrocephala

Type locality:* "Caiçara, Rio Paraguai, Mato Grosso, Brazil (16°03' S 57°43' W)." [= Fazenda de Caiçara, Cáceres, Mato Grosso] [yellow star].

Type specimen: NMW 1293, holotype, see Tiedemann et al. (1994), Fritz and Pauler (1999), and Gemel et al. (2019).

Etymology:* Greek, *makros*, long or large, and *kefale*, head; referring to the large wide head.

Comment:* Collected by Johann Natterer in 1826 in Fazenda de Caiçara, near the origin of Rio Sangradorzinho, near Cáceres, Mato Grosso.

Phrynops chacoensis Fritz and Pauler 1992:299 (07:84)

Acanthochelys chacoensis, *Mesoclemmys chacoensis*

Type locality: "Paraguayanischer Chaco, 22°30'3" S, 59°44'30" W" [Paraguay] [green star].

Type specimen: SMNS 3984–85 (shell and soft parts of one specimen), holotype, see Schlüter and Hallermann (1997).

Etymology:* Toponym for the Chaco region of South America.


Acanthochelys pallidipectoris (Freiberg 1945)
Chaco Side-necked Turtle



Thomas and Sabine Vinke / CBFTT / Boquerón, Paraguay



left: Sébastien Métrailler / CBFTT / Salta, Argentina
right: Thomas and Sabine Vinke / CBFTT / Boquerón, Paraguay



** (orange dot = probable trade or translocated; star = type locality)

Distribution:* Argentina (*Chaco*, *Formosa*, *Salta*, *Santa Fe*), Bolivia (*Tarija*), Paraguay

Introduced: Argentina (*Mendoza*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 307,421 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 459,256 sq. km

Size (Max SCL): male 18.0 cm, female 17.5 cm (Vinke et al. 2011 CBFTT)

CBFTT Account: Vinke, Vinke, Richard, Cabrera, Paszko, Marano, and Métrailler (2011)

IUCN Red List: Endangered (EN A2cde+3cde+4cde; C1+2a(i)) (Vinke and Vinke 2016). Previously: Vulnerable (VU) (TFTSG 1996)

Synonymy:

Platemys pallidipectoris Freiberg 1945:19

Acanthochelys pallidipectoris

Type locality: "Pcia. [Presidencia] Roque Sáenz Peña, Chaco" [Argentina] [yellow star].


Type specimen: MACN 1731, holotype, see also Freiberg (1947).

Etymology:* Latin, *pallidus*, pale or yellowish, and *pectoris*, breast or chest; referring to the pale yellow plastron.


Acanthochelys radiolata (Mikan 1820) (14:39, 17:87) (6)*
Brazilian Radiolated Swamp Turtle, *Cágado-Amarelo*



Russell A. Mittermeier / Alagoas, Brazil



Russell A. Mittermeier / San Miguel dos Campos, Alagoas, Brazil



** (orange dot = misidentified or uncertain locality; stars = type localities)

Distribution: Brazil (*Alagoas*, *Bahia*, *Espírito Santo*, *Minas Gerais*, *Rio de Janeiro*, *Sergipe*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 154,757 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 791,793 sq. km

Size (Max SCL): male 18.4 cm, female 19.6 cm (Garbin et al. 2016; A.G.J. Rhodin and R.A. Mittermeier, unpubl. data)

IUCN Red List:* Data Deficient (DD) (Schiavetti, Dias, Miranda, Montag, and Fraga, in press). Previously: TFTSG Provisional Red List: Data Deficient (DD) (2011, 2018); Near Threatened (NT) (TFTSG 1996)

Synonymy:

Emys radiolata Mikan 1820:[unpaginated]

Chelodina radiolata, *Rhinemys radiolata*, *Chelys (Hydraspis) radiolata*, *Chelys radiolata*, *Hydraspis radiolata*, *Platemys radiolata*, *Platemys radiolata radiolata*, *Acanthochelys radiolata*

Type locality: "Sebastianopoli...Brasilia" [= Rio de Janeiro, Brazil]. Erroneously emended to "São Paulo: Sebastianópolis (= São Sebastião) (23º45'S, 45º25'W)" [orange star] by Rhodin et al. (1984:784); corrected to "Sebastianopolis...used to be the name... of São Sebastião do Rio de Janeiro" [= Rio de Janeiro] by Vanzolini (1994:8) [yellow star].

Type specimens: NMW 184, 1295, 23390, syntypes (3), see Gemel et al. (2019); 23390 erroneously listed as holotype by Tiedemann and Häupl (1980) and Tiedemann et al. (1994).

Etymology:* Latin, *radiolus*, small ray; referring to the carapacial striations.

Platemys gaudichaudii Duméril and Bibron 1835:427
Hydraspis gaudichaudii
Type locality: "Brésil" [Brazil] [no star].
Type specimens: MNHN 1946, 2101, syntypes (2).
Etymology:* Eponym honoring Charles Gaudichaud-Beaupré (1789–1854), France; botanist who collected the type.

Hydraspis affinis Gray 1844:41 (*nomen dubium et oblitum*)
Type locality: "Brazils" [Brazil] [no star].
Type specimen: ZMB s/n (possibly ZMB 165), holotype, not located (G. Peters, in litt.); see also Fritz et al. (1994:161).
Etymology:* Latin, *affinis*, related or adjacent.
Comment: Description based on a lost manuscript ("Gray, D. C. 22, n. 10" = Gray, J.E. No date. Descriptive Catalogue of Reptiles), but later annotated in pencil by Gray himself as being *H. radiolata* (= *Acanthochelys radiolata*) (A.F. Stimson, in litt.).

Platemys werneri Schnee 1900:463
Type locality: "Umgebung von São Paulo" [Brazil] [brown star, locality uncertain, possible trade specimen].
Type specimens: NMW 23389, one of two syntypes, see Tiedemann and Häupl (1980), Tiedemann et al. (1994), and Gemel et al. (2019); other syntype apparently lost.
Etymology:* Eponym honoring Franz Werner (1867–1939), Austria; zoologist and herpetologist who worked as a professor at the University of Vienna.


Platemys radiolata quadrisquamosa Luederwaldt 1926:437
Platemys quadrisquamosa
Type locality: "Rio Doce (Est. do Espirito Santo)...[&]...Belmonte (Bahia)" [Brazil; green stars].
Type specimens: MZUSP 62, 64, and 337, syntypes (3).
Etymology:* Latin, *quattuor*, four, and *squamosa*, having scales; referring to the four enlarged scales on the hind legs.

Acanthochelys spixii (Duméril and Bibron 1835)


Black Spiny-necked Turtle, Spix's Sideneck Turtle, *Tortuga de Canaleta, Cágado do Cerrado*



Russell A. Mittermeier / São Paulo, Brazil



Russell A. Mittermeier / São Paulo, Brazil



** (orange dots = introduced or questionable; stars = type localities)

Distribution: Argentina (*Corrientes*), Brazil (*Bahia, Goiás, Minas Gerais, Paraná, Rio Grande do Sul, Santa Catarina, São Paulo*), Uruguay

Introduced:* Argentina (*Buenos Aires [?], Chaco [?], Mendoza*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 593,661 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 1,861,836 sq. km

Size (Max SCL): male 17.8 cm, female 18.0 cm (Bager et al. 2016)

IUCN Red List:* Vulnerable (VU A2c+4c) (Marques, Estrades, Fallabrino, Horta, Portelinha, Yves, Valadão, Cabrera, Florencia David, and Miorando, in press). Previously: TFTSG Provisional Red List: Near Threatened (NT) (2011); Near Threatened (NT) (TFTSG 1996)

Synonymy:

Emys depressa Spix 1824:4 (junior homonym, not = *Emys depressa* Wied-Neuwied *in* Merrem 1820 [= *Phrynops geoffroanus*])
Type locality: "provinciarum Rio de Janeiro et fluminis Sti Francisci" [Brazil]. Restricted to "Rio de Janeiro" [Brazil] by Ernst (1983:1) [orange star]. Restricted to "Rio São Francisco, near Rio dos Pandeiros, Minas Gerais, Brazil" by Rhodin et al. (1984b:783) by restriction of *nomen novum* replacement name *Platemys spixii* Duméril and Bibron 1835 [yellow star].
Type specimen: ZSM 3003/0, lectotype, designated by Hoogmoed and Gruber (1983:345), see Franzen and Glaw (2007).
Etymology:* Latin, *de-*, down, and *pressus*, pressed; referring to the depressed flattened shell.

Emys aspera Cuvier *in* Gray 1830e:16 (*nomen oblitum*) (10:7)
Type locality: "America." Restricted to "Brésil" [Brazil] by Bour and Pauler (1987:14).
Type specimen: MNHN 8751, holotype, see Bour and Pauler (1987).
Etymology:* Latin, *asper*, rough or thorny; referring to the neck spines.

Platemys spixii Duméril and Bibron 1835:409 (*nomen novum*)
Hydraspis spixii, Acanthochelys spixii, Platemys radiolata spixii
Type locality: "Brésil" [Brazil]. Restricted to "Rio São Francisco, near Rio dos Pandeiros, Minas Gerais, Brazil" by Rhodin et al. (1984b:783) [yellow star].
Type specimen: ZSM 3003/0, lectotype by default, designated by Hoogmoed and Gruber (1983:345); MNHN 8751, recorded as "holotype" by Ernst (1983), is not a type, since *spixii* is a *nomen novum* for *depressa*, discussed by Rhodin et al. (1984b) and Iverson (1992).
Etymology:* Eponym honoring Johann Baptist von Spix (1781–1826), Germany; zoologist, herpetologist, turtle taxonomist, and explorer of the Brazilian Amazon.
Comment:* Justified replacement name for *depressa*.

Chelus Duméril 1805 (10:11)

Matamata Turtles

(includes 2 species, 2 taxa)

Chelus Duméril 1805:76 (10:11)

Type species: *Chelus fimbriata* [= *Testudo fimbriata* Schneider 1783], by original monotypy.

Etymology:* Greek, *chelys*, turtle.

Chelys Oppel 1811:6 (*nomen novum*)

Etymology:* Greek, *chelys*, turtle.

Chelyda Rafinesque 1815:75 (*nomen novum*)

Etymology:* Greek, *chelys*, turtle.

Matamata Merrem 1820:21 (*nomen novum*)


Etymology:* Vernacular (Amazonian Native, Tupi Guarani), *mata-matá*, local indigenous name for the species.


Chelus fimbriata (Schneider 1783) (17:88, 21:10, 21:11)

Amazon Matamata, *Mata-Matá Amazonica*



Rafael Bernhard / Cachoeira do Caldeirão, Rio Madeira, Amazonas, Brazil





left: Peter C.H. Pritchard / CBFTT / Leticia, Amazonas, Colombia
right: Anders G.J. Rhodin / CBFTT / Leticia, Amazonas, Colombia



** (orange dots = possibly translocated or trade; stars = type localities)

Distribution: Bolivia (*Beni, Pando, Santa Cruz*), Brazil (*Amapá, Amazonas, Goiás, Maranhão, Mato Grosso, Pará, Rondônia, Tocantins*), Colombia (*Amazonas, Caquetá, Putumayo, Vaupés*), Ecuador, French Guiana, Peru (*Amazonas, Cuzco (?), Huanuco, Junin, Loreto, Madre de Dios, Pasco (?), Puno (?), San Martin, Ucayali*), Suriname

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 3,173,941 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 6,709,315 sq. km

Size (Max SCL): male 42.8 cm, female 43.7 cm (Dixon and Soini 1977; A.G.J. Rhodin, unpubl. data)

CBFTT Account: Pritchard (2008)

IUCN Red List:* Least Concern (LC) (Portelinha, Marques, Cunha, Brito, Valadão, Portilla, Páez, and Lamar, in press). Previously: TFTSG Provisional Red List: Least Concern (LC) (2011); Least Concern (LC) [Not Listed] (Behler 1996)

CITES:* Appendix II (2023)

Synonymy:

Testudo terrestris Fermin 1765:51 (junior homonym and *nomen suppressum*, not = *Testudo terrestris* Garsault 1764 (*nomen oblitum*) [= *Emys orbicularis orbicularis*], not = *Testudo terrestris* Forskål 1775 (*nomen protectum*) [= *Testudo (Testudo) graeca terrestris*])

Type locality: "Surinam" [yellow star].

Type specimen: Not located or figured.

Etymology:* Latin, *terrestris*, of the earth, terrestrial.

Comment: Name suppressed by ICZN (1963), see Wermuth (1956) and Mertens and Wermuth (1961).

Testudo fimbriata Schneider 1783:349 (*nomen protectum*)

Chelus fimbriata, Chelys fimbriata, Matamata fimbriata, Chelus fimbriatus

Type locality:* "Surinam..[&]..Aprouague..[&]..Remire" [Suriname and Approuague River and Rémiré-Montjoly, French Guiana] [yellow stars]

Type specimen: Not known or located.

Etymology:* Latin, *fimbria*, fringe, *fimbriata*, fringed; referring to the multiple fringelike skin flaps on the head and neck.

Comment: Name conserved by ICZN (1963), see Wermuth (1956) and Mertens and Wermuth (1961).

Testudo fimbria Gmelin 1789:1043 (*nomen novum*)

Chelys fimbria

Etymology:* Latin, *fimbria*, fringe, *fimbriata*, fringed; referring to the multiple fringelike skin flaps on the head and neck.

Comment: Unjustified replacement name for *fimbriata*.

Testudo matamata Bruguière 1792:257

Emydes matamata, Chelus matamata, Chelys matamata

Type locality: "Cayenne" [French Guiana] [green star].

Type specimen: Not located, holotype, type specimen figured (pl.13).

Etymology:* Vernacular (Amazonian Native, Tupi Guarani), *mata-matá*, local indigenous name for the species.

Testudo bispinosa Ruiz de Xelva *in* Daudin 1801:94

Testudo bi-spinosa, Chelys bispinosa, Matamata bispinosa

Type locality: "Brésil" [Brazil].

Type specimen: Possibly MNHN, not located.

Etymology:* Latin, *bi-*, two, and *spinosa*, spined; referring to the two supracaudal marginal scutes that point out like spines.

Testudo rapara Gray 1831d:44 (*nomen nudum*)

Etymology:* Vernacular (Suriname Creole, Sranan Tongo), *raparapa*, local indigenous name for the species.

Testudo raparara Gray 1844:44 (*nomen nudum et novum*)

Etymology:* Vernacular (Suriname Creole, Sranan Tongo), *raparapa*, local indigenous name for the species.

Chelys boulengerii Baur 1890b:968

Type locality: Not known; recorded as "South America" in NHMUK catalogue.

Type specimen: NHMUK 1881.9.27.3, shell and skull, figured by Boulenger (1889:f.52-53), lectotype, designated by Vargas-Ramírez et al. (2020:13).

Etymology:* Eponym honoring George Albert Boulenger (1858–1937), Belgium / United Kingdom; herpetologist and curator at the British Museum of Natural History, London.

Testudo corticolis Ferreira *in* Carvalho 1972:210 (*nomen nudum*) (21:10)

Testudo corticollis

Etymology:* Latin, *corti*, error for *torti*, twisted, and *collum*, neck; referring to the horizontal bending of the neck.

Testudo torticollis Ferreira *in* Ceríaco and Bauer 2017:42 (*nomen nudum et novum*) (21:10)


Etymology:* Latin, *tortus*, twisted, and *collum*, neck; referring to the horizontal bending of the neck.

Chelus orinocensis Vargas-Ramírez, Caballero, Morales-Betancourt, Lasso, Amaya, Martínez, Viana, Vogt, Farias, Hrbek, Campbell, and Fritz 2020 (21:11)

Orinoco Matamata, *Matamata del Orinoco, Caripatúa*



Mónica A. Morales-Betancourt / La Macarena, Guayabero R., Meta, Colombia



Mónica A. Morales-Betancourt / Pto. Carreño, Bita R., Vichada, Colombia



** (orange dots = probably introduced or trade; star = type locality)

Distribution: Brazil (*Amazonas, Roraima*), Colombia (*Arauca, Casanare, Guainía, Meta, Vaupés, Vichada*), Guyana, Trinidad, Venezuela (*Amazonas, Anzoátegui, Apure, Barinas, Bolívar, Cojedes, Delta Amacuro, Guárico, Monagas, Sucre*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 1,032,210 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 1,963,235 sq. km

Size (Max SCL): male 48.5 cm, female 52.6 cm (Kabisch 1997; Meier and Schaefer 2003; Barrio-Amorós and Manrique 2006; Pritchard 2008 CBFTT; Vargas-Ramírez et al. 2020)

CBFTT Account: Pritchard (2008) [as part of *Chelus fimbriata* sensu lato]

IUCN Red List:* Least Concern (LC) (Morales-Betancourt, Lasso, and Rojas-Runjaic, in press). Previously: TFTSG Provisional Red List: Least Concern (LC) (2011), as part of *Chelus fimbriata*; Least Concern (LC) [Not Listed] (Behler 1996), as part of *Chelus fimbriata*

CITES:* Appendix II (2023)

Synonymy:

Chelus orinocensis Vargas-Ramírez, Caballero, Morales-Betancourt, Lasso, Amaya, Martínez, Viana, Vogt, Farias, Hrbek, Campbell, and Fritz 2020:13
Type locality: "Colombia, Vichada, Pto. Carreño, Bita River (5.7610 N / 68.5858 W)" [yellow star].
Type specimen: IAVH R-8755, holotype.
Etymology:* Toponym for the Orinoco River region of Colombia and Venezuela.

Mesoclemmys Gray 1873c (07:100, 21:12)

Toad-headed Turtles

(includes 9 species, 9 taxa)

Mesoclemmys Gray 1873c:305
Type species: *Mesoclemmys gibba* [= *Emys gibba* Schweigger 1812], by original monotypy.
Etymology:* Greek, *mesos*, middle, and *klemmys*, turtle; referring to the ability to retract and hide the head and neck and that the skull is intermediate in shape between *Hydraspis* (= *Phrynops*) and *Platemys*.

Batrachemys Stejneger 1909:126
Type species: *Batrachemys nasuta* [= *Emys nasuta* Schweigger 1812], by original monotypy.
Etymology:* Greek, *batrachos*, frog, and *emys*, turtle; referring to the frog-like broad face.

Bufocephala McCord, Joseph-Ouni, and Lamar 2001:732
Type species: *Bufocephala vanderhaegei* [= *Phrynops tuberculatus vanderhaegei* Bour 1973], by original designation.
Etymology:* Latin, *bufo*, toad, and Greek, *kefale*, head; referring to the toad-like broad face.

Mesoclemmys dahli (Zangerl and Medem 1958) (12:40, 21:13)


Dahl's Toad-headed Turtle, *Carranchina*



German Forero-Medina / CBFTT / Colombia



left: Uwe Fritz / Colombia; *right*: Russell A. Mittermeier / Colombia



** (orange dots = possible translocations, star = type locality)

Distribution: Colombia (*Atlántico, Bolívar, Cesar, Córdoba, Magdalena, Sucre*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 28,459 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 58,711 sq. km

Size (Max SCL): male 22.9 cm, female 29.7 cm (Forero-Medina et al. 2013 CBFTT)

CBFTT Account: Forero-Medina, Castaño-Mora, Cárdenas-Arevalo, and Medina-Rangel (2013)

IUCN Red List:* Critically Endangered (CR A2c+4c)

(Gallego-García and Forero-Medina, in press). Previously: TFTSG Provisional Red List: Critically Endangered (CR) (2017); Critically Endangered (CR B1+2c [ver. 2.3]) (TFTSG 1996)

Synonymy:

Phrynops (Batrachemys) dahli Zangerl and Medem 1958:376

Phrynops dahli, Batrachemys dahli, Phrynops nasutus dahli, Mesoclemmys dahli

Type locality: "Vicinity of Sincelejo, Bolivar, Colombia" [yellow star].

Type specimen: FMNH 75980, holotype, see Marx (1976).


Etymology:* Eponym honoring George Dahl (1905–1979), Sweden / Colombia; biologist and ichthyologist who worked and lived in Colombia and collected the types.

Mesoclemmys gibba (Schweigger 1812) (21:14)


Gibba Turtle, *Hedionda, Cágado-de-Poça*



Frank Deschandol / French Guiana



Anders G.J. Rhodin / Maipuco, Rio Maranon, Peru



** (orange dot = questionable, possibly translocated or misidentified; stars = type localities)

Distribution: Bolivia, Brazil (*Acre, Amapá, Amazonas, Maranhão, Mato Grosso, Pará, Rondônia, Roraima, Tocantins*), Colombia (*Amazonas, Arauca, Caquetá, Casanare, Guainía, Guaviare, Meta, Putumayo, Vaupés, Vichada*), Ecuador, French Guiana, Guyana, Peru (*Amazonas, Loreto, Madre de Dios, Puno, Ucayali*), Suriname, Trinidad, Venezuela (*Amazonas, Bolívar, Delta Amacuro, Monagas*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 4,445,462 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 7,072,096 sq. km

Size (Max SCL): male 20.0 cm, female 23.3 cm (Mittermeier et al. 1978; Pritchard and Trebbau 1984; Ernst et al. 2006b; Itescu et al. 2014)

IUCN Red List:* Least Concern (LC) (Ettmar, in press). Previously: TFTSG Provisional Red List: Least Concern (LC) (2011); Least Concern (LC) [Not Listed] (Behler 1996)

Synonymy:

Emys gibba Schweigger 1812:299

Rhinemys gibba, Hydraspis cayennensis gibba, Platemys gibba, Hydraspis (Podocnemis) gibba, Hydraspis gibba, Phrynops gibbus, Mesoclemmys gibba, Phrynops (Mesoclemmys) gibba, Mesoclemmys gibbus

Type locality: Not known. Restricted to "Amérique méridional" by Duméril and Duméril (1851:20); and to "environs de Cayenne, Guyane française" [French Guiana] by Bour and Pauler (1987:7) [yellow star].

Type specimen: MNHN 8756, holotype, see Ernst (1981b), Iverson (1992), Bour (2005d), and Ettmar (2019); MNHN 8356 (a typo) erroneously listed as holotype by Bour (2005d) and Ceríaco and Bour (2012).

Etymology:* Latin, *gibbus*, hunched, humped, bent; referring to the deep humped shell with posterior vertebral keel.

Emys stenops Spix 1824:12 (21:14)

Hydraspis stenops

Type locality:* "fluminis Solimoëns" [Rio Solimões, Amazonas, Brazil]. Restricted to "l'Amazone entre Tefé et Tabatinga au Brésil (Amazonas)" [Brazil] by Bour and Pauler (1987:7) [= possibly near São Paulo de Olivença] [green star].

Type specimen: ZSM 2454/0, holotype, see Hoogmoed and Gruber (1983), Bour (2005d), and Franzen and Glaw (2007).

Etymology:* Greek, *stenos*, narrow, and *ops*, eye; referrring to the narrow interorbital distance.

Platemys miliusii Duméril and Bibron 1835:431

Phrynops miliusii, Hydraspis miliusii

Type locality: "Cayenne" [French Guiana] [yellow star].

Type specimen: MNHN 8755, holotype, see Bour (2005d), who recorded it erroneously as MNHN 8355.

Etymology:* Eponym honoring Baron Pierre Bernard Milius (1773–1829), France; nobleman, naval officer, and Governor of French Guiana, who obtained the type specimen.

Hydraspis gordoni Gray 1868:563

Type locality: "Trinidad, near the mountain of Tamana" [pink star].

Type specimen: NHMUK 1947.3.4.18 (formerly 1881.69.2.1), holotype, see Bour (2005d).

Etymology:* Eponym honoring Arthur Hamilton-Gordon (1829–1912), United Kingdom; nobleman and Governor of Trinidad, who obtained the type.

Hydraspis bicolor Gray 1873c:304

Type locality: "Demerara Falls" [Guyana] [blue star].

Type specimen: NHMUK 1946.1.22.86 (formerly 1872.10.16.80), holotype, see Bour (2005d), who recorded it erroneously as NHMUK 1946.2.22.86.

Etymology:* Latin, *bi-*, two, and *color*, tint or hue.

Mesoclemmys nasuta (Schweigger 1812)
Guyanan Toad-headed Turtle




Jérôme Maran / French Guiana





Left: Fábio A.G. Cunha / Pará, Brazil
Right: Eduardo Campos / Macapá, Amapá, Brazil



** (star = type locality)

Distribution:* Brazil (*Amapá, Maranhão, Pará*), French Guiana, Suriname

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 356,146 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 594,289 sq. km

Size (Max SCL):* male 31.7 cm, female 27.8 cm (Pritchard and Trebbau 1984; Métrailler and Le Gratiet 1996; Cunha et al. 2025)

IUCN Red List:* Data Deficient (DD) (Cunha, Brito, and Valadão, in press). Previously: TFTSG Provisional Red List: Data Deficient (DD) (2011, 2018); Least Concern (LC) [Not Listed] (Behler 1996)

Synonymy:

Emys nasuta Schweigger 1812:298
Rhinemys nasuta, Hydraspis (Rhinemys) nasuta, Hydraspis nasuta, Platemys nasuta, Batrachemys nasuta, Phrynops (Batrachemys) nasuta, Phrynops nasuta, Phrynops nasutus, Phrynops nasutus nasutus, Phrynops nasuta nasuta, Batrachemys nasutus, Mesoclemmys nasuta
Type locality: Not known. Restricted to "Amérique méridionale" by Duméril and Bibron (1835:437); to "Guyanes et au nord-est de l'Amazonie" by Lescure and Fretey (1976:1318); and to "rivières Ouaqui et Inini, bassin du Maroni en amont de Maripasoula, Guyane française" [French Guiana] by Bour and Pauler (1987:6) [yellow star].
Type specimen: MNHN 4140, holotype, see Bour (2007d) and Ceríaco and Bour (2012).
Etymology:* Latin, *nasuta*, having a big nose; referring to the prominent snout.

Emys barbatula Gravenhorst 1829:15
Hydraspis barbatula
Type locality: Not known.
Type specimen: MNHW, holotype, specimen figured (pl.5,f.3-4), apparently destroyed during World War II, see Ettmar (2019).
Etymology:* Latin, *barbatus*, bearded; referring to the chin barbels.

Platemys schweiggerii Duméril and Bibron 1835:435 (*nomen novum*)
Type specimen: MNHN 4140, holotype by default, see Bour (2007d) and Ceríaco and Bour (2012).
Etymology:* Eponym honoring Augustus F. Schweigger (1783–1821), Germany; naturalist, herpetologist, and turtle taxonomist.
Comment:* Unjustified replacement name for *nasuta*.

Phrynops walbaumi Fitzinger *in* Siebenrock 1904b:20 (*nomen nudum*)
Etymology:* Eponym honoring Johann J. Walbaum (1724–1799), Germany; physician, naturalist, and taxonomist.

Mesoclemmys perplexa Bour and Zaher 2005
Cerrado Side-necked Turtle, Piaui Side-necked Turtle




Vinicius T. de Carvalho / Piauí, Brazil





Vinicius T. de Carvalho / Piauí, Brazil



** (orange dots = questionable or misidentified; star = type locality)

Distribution: Brazil (*Ceará, Goiás [?], Maranhão, Piauí, Tocantins [?]*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 195,476 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 291,836 sq. km

Size (Max SCL): male 16.9 cm, female 21.4 cm (Maran 2006b)

IUCN Red List:* Data Deficient (DD) (Bock, Páez, Carvalho, and Rhodin, in press). Previously: TFTSG Provisional Red List: Data Deficient (DD) (2011, 2018)


Synonymy:


Mesoclemmys perplexa Bour and Zaher 2005:298
Type locality: "the regions called "Olho d'Água da Santa" and "Baixão do Fausto", southern part of the Parque Nacional da Serra das Confusões, State of Piauí, Brasil" [Brazil] [yellow star].

Type specimen: MZUSP 4111, holotype, see Ettmar (2019).
Etymology:* Latin, *perplexa*, confused, obscure, ambiguous; referring to the Serra das Confusões type locality and the ambiguous generic allocation of this species.

Mesoclemmys raniceps (Gray 1856b) (12:41, 17:89, 21:15) (Z)*
Amazon Toad-headed Turtle


Jiří Moravec / nr. Anguilla, Rio Nanay, Loreto, Peru


Fábio A.G. Cunha / Barreirinha, Amazonas, Brazil


** (stars = type localities)


Distribution: Bolivia, Brazil (*Acre, Amazonas, Mato Grosso, Pará, Rondônia, Roraima*), Colombia (*Amazonas, Caquetá, Guainía, Putumayo, Vaupés*), Ecuador, Peru (*Loreto, Madre de Dios, Pasco, Ucayali*), Venezuela (*Amazonas*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 3,434,777 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 4,860,277 sq. km
Size (Max SCL): male 33.4 cm, female 33.5 cm (Brito et al. 2019a)
IUCN Red List:* Least Concern (LC) (Cunha, Vasconcelos, Viana, Brito, and Valadão, in press). Previously: TFTSG Provisional Red List: Data Deficient (DD) (2011, 2018); Least Concern (LC) [Not Listed] (Behler 1996)
Synonymy:
Hydraspis raniceps Gray 1856b:55
Platemys raniceps, Phrynops raniceps, Batrachemys raniceps, Batrachemys raniceps raniceps, Mesoclemmys raniceps
Type locality:* "Brazils; Para" [Pará, Brazil; probably = Belém, Pará, Brazil] [yellow star].
Type specimen: NHMUK 1947.3.5.92 (formerly 1851.8.12.1), lectotype, designated by Bour and Pauler (1987:8).
Etymology:* Latin, *rana*, frog, and *-ceps*, head; referring to the large, broad head.


Batrachemys heliostemma McCord, Joseph-Ouni, and Lamar 2001:734 (21:15)
Mesoclemmys heliostemma
Type locality: "base of Pico da Neblina (situated on the Venezuela/Brazil border) on the left bank of Río Baria (= Río Mawarinuma) [4º95'N, 66º10'W], a tributary of the Rio Negro, Amazonas, Venezuela." Original GPS coordinates incorrect, emended to 0º50'N, 66º10'W by TTWG (2017:181) [green star].
Type specimen: USNM 541895, holotype, see Reynolds et al. (2007).
Etymology:* Greek, *helios*, the sun, and *stemma*,wreath, crown; referring yellow-orange head markings.

Mesoclemmys jurutiensis Cunha, Sampaio, Carneiro, and Vogt 2021:155 (Z)*
Type locality:* "the region of the Capiranga Community, Juruti-Velho Village, in Juruti municipality, State of Pará, Brazil (WGS84, 02º36'16.20"S, 56º24'40.80"W; 90 m elev. asl)" [blue star].
Type specimen:* INPA H41283, holotype.
Etymology:* Toponym for Juruti, Pará, Brazil.

Mesoclemmys tuberculata (Luederwaldt 1926)
Tuberculate Toad-headed Turtle


Daniel O. Santana / CBFTT / Areia Branca, Sergipe, Brazil


left: Daniel O. Santana / CBFTT / Areia Branca, Sergipe, Brazil
right: Daniel O. Santana / CBFTT / Tobias Barreto, Sergipe, Brazil


** (star = type locality)

Distribution: Brazil (*Alagoas, Bahia, Ceará, Maranhão, Minas*

Gerais, Paraíba, Pernambuco, Piauí, Rio Grande do Norte, Sergipe)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 608,127 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 1,325,430 sq. km

Size (Max SCL):* male 20.1 cm, female 25.8 cm (Santana et al. 2016 CBFTT; Santana et al. 2023)

CBFTT Account: Santana, Marques, Vieira, Moura, Faria, and Mesquita (2016)

IUCN Red List: Least Concern (LC) (Santana and Cunha, in press). Previously: TFTSG Provisional Red List: Data Deficient (DD) (2011, 2018); Least Concern (LC) [Not Listed] (Behler 1996)


Synonymy:

Rhinemys tuberculata Luederwaldt 1926:428
Batrachemys tuberculata, Phrynops (Batrachemys) tuberculata, Phrynops tuberculata, Phrynops (Batrachemys) tuberculatus, Phrynops tuberculatus, Phrynops tuberculatus tuberculatus, Mesoclemmys tuberculata
Type locality: "Brasil: Estado da Bahia e Pará" [Brazil]. Restricted to "Villa Nova, Bahia" by Mertens and Wermuth (1955:400); and to "Fortaleza, Ceará" [Brazil] by Bour and Pauler (1987:9) [yellow star].
Type specimen: MZUSP 43, lectotype, designated by Bour and Pauler (1987:9).
Etymology:* Latin, *tuberculum*, small hump, nodule; referring to the humplike skin nodules on the head and neck.

Mesoclemmys vanderhaegei (Bour 1973) (R)*
Vanderhaege's Toad-headed Turtle, *Cágado-Cabeça-de-Sapo*


Elizângela S. Brito / CBFTT / Cáceres, Mato Grosso, Brazil


left: Anders G.J. Rhodin / *right*: Russell A. Mittermeier
Chapada dos Guimarães, Mato Grosso, Brazil / female


**
(orange dots = questionable; stars = type localities)

Distribution:* Argentina (*Corrientes, Misiones*), Bolivia, Brazil (*Goiás, Maranhão, Mato Grosso, Mato Grosso do Sul, Minas Gerais, Pará, São Paulo, Tocantins*), Paraguay

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 979,971 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 3,090,178 sq. km

Size (Max SCL): male 28.5 cm, female 28.0 cm (Marques et al. 2014 CBFTT)

CBFTT Account: Marques, Böhm, Brito, Cabrera, and Verdade (2014)

IUCN Red List:* Vulnerable (VU A2c+4c) (Brito, Valadão, Cunha, Vinke, Vinke, Cabrera, Marques, Carvajal-Bacarreza, and Strüssmann, in press). Previously: TFTSG Provisional Red List: Near Threatened (NT) (2018); Near Threatened (NT) (TFTSG 1996)

Synonymy:

Phrynops schöpfii Fitzinger *in* Diesing 1839:237 (*nomen nudum*)
Type locality: "Cuiabá, Mato Grosso, Brésil" [Brazil] [orange star]
Type specimen: NMW 15762, see Bour and Pauler (1987).
Etymology:* Eponym honoring Johann David Schoepff (or Schöpf) (1752–1800), Germany; physician, botanist, herpetologist, and productive turtle taxonomist.

Phrynops schoepffii Fitzinger *in* Siebenrock 1904b:22 (*partim, nomen nudum et novum*)
Etymology:* Eponym honoring Johann David Schoepff (or Schöpf) (1752–1800), Germany; physician, botanist, herpetologist, and productive turtle taxonomist.
Comment: Justified emendation of *P. schöpfii*. Name synonymized *in partim* under *Acanthochelys macrocephala* and *Mesoclemmys vanderhaegei*.

Phrynops paraguayensis Vanzolini *in* Donoso-Barros 1965:13 (*nomen nudum*)
Type locality: "rio Paraguay," see Bour and Pauler (1987) [no star].
Type specimen: Possibly MZUSP, not located.
Etymology:* Toponym for Paraguay.


Phrynops tuberculatus vanderhaegei Bour 1973:175
Phrynops (Batrachemys) vanderhaegei, Phrynops vanderhaegei, Batrachemys vanderhaegei, Bufocephala vanderhaegei, Mesoclemmys vanderhaegei
Type locality: "environs d'Asunción au Paraguay." Restricted to "Tobati (25º15' S, 57º04' W), La Cordillera, Paraguay" by Bour and Pauler (1987:10) [yellow star].
Type specimen: MNHN 1977.50, holotype, see Bour and Pauler (1987).
Etymology:* Eponym honoring Maurice Vanderhaege (1938–), France; amateur herpetologist.

Mesoclemmys sabiniparaensis Cunha, Sampaio, Carneiro, Vogt, Mittermeier, Rhodin, and Andrade 2022:164 (R)*
Type locality:* "near Serra das Andorinhas State Park, municipality of São Geraldo do Araguaia, state of Pará, Brazil (WGS84, 06º12'57"S, 48º27'29"W, 212 m above sea level)" [green star].
Type specimen:* MPEG-H1251, holotype.
Etymology:* Eponym honoring Andrew Sabin (1945–), USA; conservationist and philanthropist; and Toponym for Pará, Brazil.

Mesoclemmys wermuthi (Mertens 1969) [21:15]
Wermuth's Toad-headed Turtle



Richard C. Vogt / Iquitos, Peru / captivity



left: Roger Bour / Iquitos, Peru / captivity
right: Fábio A.G. Cunha / nr. Juruti, Pará, Brazil



(star = type locality)

Distribution: Bolivia, Brazil (*Acre, Amazonas, Mato Grosso, Pará, Rondônia*), Colombia (*Amazonas*), Peru (*Loreto, Madre de Dios, Pasco, Ucayali*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 1,953,977 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 2,725,690 sq. km

Size (Max SCL): male 23.2 cm, female 33.8 cm (Molina et al. 2012; A.G.J. Rhodin and R.A. Mittermeier, unpubl. data)

IUCN Red List:* Data Deficient (DD) (Cunha, Brito, Valadão, Oliveira, and Vasconcelos, in press). Previously: TFTSG Provisional Red List: Data Deficient (DD) (2020)

Synonymy:


Hydraspis maculata Gray 1873c:305 (*nomen oblitum*) [17:89, 21:15]
Mesoclemmys maculata
Type locality: "Tropical America." Restricted to "S. America" by Gray (1873j:65); and to "Venezuela" [no star] by Boulenger (1889:219) and Pritchard and Trebbau (1984:127), in error, see Rivas et al. (2015).
Type specimen: NHMUK 1946.1.22.14 (formerly 1866.8.14.233), holotype, see Bour and Pauler (1987), Rivas et al. (2015), Ettmar (2019), and Cunha et al. (2019).
Etymology:* Latin, *macula*, spot, spotted; referring to the white spots on the pleural scutes and a dark brown spot on the plastron.

Phrynops wermuthi Mertens 1969b:132 [21:15]
Phrynops tuberculatus wermuthi, Phrynops (Batrachemys) nasutus wermuthi, Phrynops nasutus wermuthi, Phrynops nasuta wermuthi, Batrachemys raniceps wermuthi, Mesoclemmys wermuthi
Type locality: "Peru...zweifellos der amazonische Teil des Landes." Restricted to "Iquitos (3°50' S 73°15' W), Loreto, Peru" by Bour and Pauler (1987:8) [yellow star].
Type specimen: SMF 66246, holotype, see Mertens (1970) and Bour and Pauler (1987).
Etymology:* Eponym honoring Heinz Wermuth (1918–2002), Germany; herpetologist and curator at the Staatliches Museum für Naturkunde in Stuttgart.


Mesoclemmys zuliae (Pritchard and Trebbau 1984) [12:40]
Zulia Toad-headed Turtle, *Cabezón de Zulia*



Peter C.H. Pritchard / nr. Maracaibo, Zulia, Venezuela



Peter C.H. Pritchard / nr. Maracaibo, Zulia, Venezuela



(star = type locality)

Distribution:* Colombia (?) (*Norte de Santander [?]*), Venezuela (*Mérida, Tachira [?], Zulia*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 21,344 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 30,972 sq. km

Size (Max SCL): male 20.8 cm, female 27.9 cm (Pritchard and Trebbau 1984; Ceballos et al. 2013)

IUCN Red List:* Critically Endangered (CR A2ce+4ce) (Rivas, Lasso-Alcalá, Barros, Franklin, Sibira, Viloria, Escalona, and Páez, in press). Previously: TFTSG Provisional Red List: Vulnerable (VU) (2011); Vulnerable (VU B1+2c [ver. 2.3]) (TFTSG 1996)

Synonymy:

Phrynops (Batrachemys) zuliae Pritchard and Trebbau 1984:135
Phrynops zuliae, Batrachemys zuliae, Mesoclemmys zuliae
Type locality: "Caño Madre Vieja near El Guayabo, Distrito Colón, Edo. Zulia, Venezuela (8°53' N, 72°30' W)" [yellow star].
Type specimen: UF 53439, holotype.
Etymology:* Toponym for Zulia State, Venezuela.

Phrynops Wagler 1830b (07:100)
Stripe-faced Side-necked Turtles
(includes 4 species, 4 taxa)
Phrynops Wagler 1830b:135
Type species: *Phrynops geoffroanus* [= *Emys geoffroana* Schweigger 1812], by original monotypy.
Etymology:* Greek, *phrynos*, toad or frog, and *ops*, face or eye; referring to toad- or frog-like appearance of the face.
Spatulemys Gray 1872b:463
Type species: *Spatulemys lasalae* [= subjective synonym of *Platemys hilarii* Duméril and Bibron 1835], by original monotypy.
Etymology:* Greek, *spathe*, spatula, and *emys*, turtle; referring to the spatula-like elongated broad plastron with a rounded front.
Parahydraspis Wieland 1923:2
Type species: *Parahydraspis paranaensis* † Wieland 1923, by original monotypy.
Etymology:* Greek, *para-*, near relationship, *hydor, hydro-*, water, and *aspis*, a shield.

Phrynops geoffroanus (Schweigger 1812) (10:44, 14:40, 17:90, 21:16) (9)*
Geoffroy's Side-necked Turtle, *Cágado de Barbicha*


Vinícius T. de Carvalho / Piracicaba, São Paulo, Brazil



Russell A. Mittermeier / São Paulo, Brazil / captivity


**
(orange dots = possible translocations; stars = type localities)

Distribution: Argentina (*Corrientes, Misiones*), Bolivia (*Beni*), Brazil (*Acre, Alagoas, Amapá, Amazonas, Bahia, Ceará, Espírito Santo, Goiás, Maranhão, Mato Grosso, Mato Grosso do Sul, Minas Gerais, Pará, Paraíba, Paraná, Pernambuco, Piauí, Rio de Janeiro, Rio Grande do Norte, Rio Grande do Sul, Rondônia, Santa Catarina, São Paulo, Sergipe, Tocantins*), Colombia (*Amazonas, Caquetá, Casanare, Guainía, Meta, Putumayo, Vaupés, Vichada*), Ecuador, Paraguay, Peru (*Cusco, Huanuco, Junin, Loreto, Madre de Dios, Pasco*), Venezuela (*Amazonas*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 3,407,292 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 11,495,510 sq. km
Size (Max SCL): male 39.5 cm, female 46.3 cm (Souza and Abe 2001; A.G.J. Rhodin and R.A. Mittermeier, unpubl. data)
IUCN Red List:* Least Concern (LC) (Páez, Bock, Rhodin, Mittermeier, Cisneros-Heredia, and Rocha e Silva, in press).
Previously: TFTSG Provisional Red List: Least Concern (LC) (2011); Least Concern (LC) [Not Listed] (Behler 1996)
Synonymy:
Emys geoffroana Schweigger 1812:302,350
Chelodina geoffroana, Phrynops geoffroanus, Platemys geoffroana, Hydraspis (Phrynops) geoffroana, Hydraspis geoffroana, Phrynops geoffroana, Rhinemys geoffroana, Phrynops geoffroana geoffroana, Phrynops geoffroanus geoffroanus, Phrynops (Phrynops) geoffroanus
Type locality: "Brasilia" [Brazil] [yellow star, location estimated, restricted here to "Cuiabá, Mato Grosso, Brazil" (V.T. de Carvalho, A.G.J. Rhodin, and R.A. Mittermeier, unpubl. data)]. (9)*
Type specimen: MNHN 9417, holotype, see Bour (2008c) and Ceríaco and Bour (2012).
Etymology:* Eponym honoring Étienne Geoffroy Saint-Hilaire (1772–1844), France; zoologist who obtained the type.
Emys depressa Wied-Neuwied *in* Merrem 1820:22 (senior homonym, not = *Emys depressa* Spix 1824 [= *Acanthochelys spixii*]) (21:16)
Testudo depressa, Chelys (Hydraspis) depressa, Chelys depressa, Hydraspis depressa, Platemys depressa
Type locality:* "Brasilia" [Brazil]. Restricted to "Flusse Mucurí" [Rio Mucurí, Bahia, Brazil] by Wied-Neuwied (1825a:29) [blue star].
Type specimen:* Not located, specimen figured by Wied-Neuwied (1825b:pl.4-5), lectotype, designated by Bour (2008c:39).
Etymology:* Latin, *de-*, down, and *pressus*, pressed; referring to the depressed flattened shell.
Emys viridis Spix 1824:3
Chelys (Hydraspis) viridis, Chelys viridis, Hydraspis viridis
Type locality: "fluminis Carinhanhae, confluentis Sti Francisci" [Rio Carinhanha, at junction with Rio São Francisco, Bahia, Brazil]. Emended to "Brazil, back-swamp lakes of Rio Carinhanha at confluent with São Francisco, probably at or near 14°18'S, 43°47'W" by Vanzolini (1981:xxiv) [green star].
Type specimen: ZSM 3008/0, holotype, see Hoogmoed and Gruber (1983) and Franzen and Glaw (2007).
Etymology:* Latin, *viridis*, green; referring to the olive green carapace and a yellowish green plastron.
Emys tritentaculata Saint-Hilaire *in* Cuvier 1829:11 (*nomen nudum et dubium*) (14:40)
Etymology:* Latin, *tria*, three, and *tentaculum*, thread; referring to the assertion that this taxon has three chin barbels.
Emys geoffroyana Gray 1830e:16 (*nomen novum*)
Phrynops geoffroyana, Hydraspis geoffroyana, Platemys geoffroyana
Etymology:* Eponym honoring Étienne Geoffroy Saint-Hilaire (1772–1844), France; zoologist who obtained the type specimen.
Comment: Invalid replacement name or error for *geoffroana*.
Platemys geoffreana Duméril and Bibron 1835:418 (*nomen novum*)
Etymology:* Eponym honoring Étienne Geoffroy Saint-Hilaire (1772–1844), France; zoologist who obtained the type specimen.
Comment: Invalid replacement name or error for *geoffroana*.
Platemys waglerii Duméril and Bibron 1835:422
Hydraspis waglerii, Phrynops waglerii

Type locality: "Brésil" [Brazil] [pink star, location estimated, restricted here to "Piracicaba, São Paulo, Brazil" (V.T. de Carvalho, A.G.J. Rhodin, and R.A. Mittermeier, unpubl. data)]. (2)*

Type specimen: MNHN 8758, holotype, see Mertens (1967b) and Bour (2008c).

Etymology:* Eponym honoring Johann G. Wagler (1800–1832), Germany; ornithologist and herpetologist at the Ludwig Maximilian University of München.

Platemys neuwiedii Duméril and Bibron 1835:425 (*nomen novum*)

Type locality: "Brésil" [Brazil].

Type specimen: Not located, specimen figured by Wied-Neuwied (1825b:pl.4-5), lectotype, by default, designated by Bour (2008c:39).

Etymology:* Eponym honoring Prince Alexander Philipp Maximilian zu Wied-Neuwied (1782–1867), Germany; nobleman, naturalist, and explorer of Brazil and the United States.

Comment:* Invalid replacement name for *depressa*.

Emys lyrae Reuss *in* Gray 1844:39 (*nomen nudum et dubium*)

Etymology:* Greek, *lyra*, lyre; possibly referring to the body shape of this species resembling a lyre.

Hydraspis boulengeri Bohls 1895:53

Type locality: "nördlichen Theile Paraguays...im Aquidaban, Tagatiya und anderen linken Nebenflüssen des Paraguaystromes" [Paraguay]. Restricted to "Departmento Concepción: Río Saladillo, 23ºS" [Paraguay] by Cacciali et al. (2016:45) [dark gray star].

Type specimen: NHMUK 1947.3.5.94 (formerly 1896.5.11.1), holotype.

Etymology:* Eponym honoring George Albert Boulenger (1858–1937), Belgium / United Kingdom; herpetologist and curator at the British Museum of Natural History, London.

Hydraspis lutzi Ihering *in* Luederwaldt 1926:441

Phrynops lutzi

Type locality: "Mogy-guassù (Est. de S. Paulo)" [= Mogi Guaçu, São Paulo, Brazil] [light gray star, partially hidden].

Type specimen: MZUSP 31, holotype.

Etymology:* Eponym honoring Bertha Lutz (1894–1976), Brazil; female herpetologist, feminist, and politician, who collected the type.

Phrynops hilarii (Duméril and Bibron 1835)


Hilaire's Side-necked Turtle, Saint-Hilaire's Side-necked Turtle



Russell A. Mittermeier / Rio Pelotas, Rio Grande do Sul, Brazil



left: Anders G.J. Rhodin / Uruguay / captivity / juvenile
right: Russell A. Mittermeier / Rio Pelotas, Rio Grande do Sul, Brazil



** (orange dots = possibly introduced or translocated; star = type locality)

Distribution: Argentina (*Buenos Aires, Chaco, Córdoba, Corrientes, Entre Rios, Formosa, Misiones, Santa Fe, Santiago del Estero*), Brazil (*Rio Grande do Sul, Santa Catarina*), Paraguay, Uruguay

Introduced: Argentina (*Formosa, La Rioja, Mendoza, Salta, San Juan, Santiago del Estero, Tucumán*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 637,615 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 1,594,666 sq. km

Size (Max SCL): male 35.6 cm, female 40.8 cm (Ernst et al. 2006b; A.G.J. Rhodin and R.A. Mittermeier, unpubl. data; A. Bager, unpubl. data)

IUCN Red List:* Least Concern (LC) (Alcalde, Carreira, and Fagundes, in press). Previously: TFTSG Provisional Red List: Least Concern (LC) (2011); Least Concern (LC) [Not Listed] (Behler 1996)

Synonymy:

Platemys hilarii Duméril and Bibron 1835:428

Hydraspis hilarii, Hydraspis geoffroyana hilarii, Phrynops hilarii, Phrynops (Phrynops) hilarii, Phrynops geoffroana hilarii, Phrynops geoffroanus hilarii

Type locality: "Brésil" [Brazil] [no star, location unknown]

Type specimen: MNHN 8757, holotype, see Bour (2008c).

Etymology:* Eponym honoring Augustin F.C. Prouvençal de Saint-Hilaire (1779–1853), France; botanist who collected in Argentina and Brazil.

Hydraspis hilairii Gray 1844:40 (*nomen novum*)

Platemys hilairii, Phrynops (Phrynops) geoffroanus hilairii, Phrynops geoffroanus hilairii

Etymology:* Eponym honoring Augustin F.C. Prouvençal de Saint-Hilaire (1779–1853), France; botanist who collected in Argentina and Brazil.

Comment: Unjustified replacement name for *hilarii*.

Spatulemys lasalae Gray 1872b:463

Spatulemys lasala

Type locality: "Rio Parana, Corrientes" [Argentina] [green star, location estimated].

Type specimen: NHMUK 1947.3.5.93 (formerly 1872.11.6.1), holotype.

Etymology:* Eponym honoring Colonel P. Perez de Lasala (1800s), Argentina; naval officer who presented the type.

Phrynops tuberosus (Peters 1870) (10:44, 17:90)
Guianan Shield Side-necked Turtle; Cotingo River Side-necked Turtle



Peter C.H. Pritchard / Canaima National Park, Rio Caroni, Bolivar, Venezuela



Peter C.H. Pritchard / Canaima National Park, Rio Caroni, Bolivar, Venezuela



** (star = type locality)

Distribution: Brazil (*Roraima*), Guyana, Venezuela (*Bolívar*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 83,664 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 162,669 sq. km

Size (Max SCL): male 28.6 cm, female 32.6 cm (Pritchard and Trebbau 1984; Ceballos et al. 2013)

IUCN Red List: Not Evaluated (NE)

TFTSG Provisional Red List: Least Concern (LC) (2011)

Synonymy:

Platemys tuberosa Peters 1870:311

Hydraspis tuberosa, Phrynops tuberosa, Phrynops geoffroana tuberosa, Phrynops geoffroanus tuberosus, Phrynops (Phrynops) tuberosus, Phrynops tuberosus

Type locality: "Cotingaflusse am Roraimagebirge in British-Guyana" [Guyana]. Emended to "Amazon drainage, the Río Cotingo... Brazil...state of Roraima" by Bour (2008c:38) [yellow star].

Type specimen: ZMB 166, holotype, see Fritz et al. (1994) and Bour (2008c).

Etymology:* Latin, *tuber*, hump or bump; referring to the sculpted bumpy carapace of the holotype of this species.

Phrynops williamsi Rhodin and Mittermeier 1983
Williams' Side-necked Turtle



Vinícius T. de Carvalho / Picada Café, rio Cadeia, Rio Grande do Sul, Brazil



Vinícius T. de Carvalho / Picada Café, rio Cadeia, Rio Grande do Sul, Brazil



** (orange dot = questionable, possibly translocated; star = type locality)

Distribution: Argentina (*Corrientes, Entre Rios [?], Misiones*), Brazil (*Paraná, Rio Grande do Sul, Santa Catarina*), Paraguay, Uruguay

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 218,807 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 621,144 sq. km

Size (Max SCL): male 27.8 cm, female 35.4 cm (Rhodin et al. 1988; Ceballos et al. 2013; Bressan and Verrastro 2020)

IUCN Red List: Vulnerable (VU A4cd) (Rhodin, Bressan, Buskirk, Cabrera, Carreira, Estrades, Mittermeier, Vinke, and Vinke 2018). Previously: Least Concern (LC) [Not Listed] (Behler 1996)

Synonymy:

Phrynops williamsi Rhodin and Mittermeier 1983:58

Phrynops (Phrynops) williamsi

Type locality:* "Rio Cadea, Rio Grande do Sul, Brazil" [= Rio Cadeia, Rio Grande do Sul, Brazil] [yellow star].

Type specimen: MCZ 64135, holotype.

Etymology:* Eponym honoring Ernest E. Williams (1914–1998), USA; herpetologist and curator at the Museum of Comparative Zoology, Harvard University, and mentor to the authors.

Platemys Wagler 1830b (14:39)

Twist-necked Turtles

(includes 1 species, 2 taxa)

Platemys Wagler 1830b:135

Type species: *Platemys planiceps* [= *Testudo planiceps* Schneider 1792] [= objective synonym of *Testudo platycephala* Schneider 1792], by original monotypy.

Etymology:* Greek, *platys*, flat, and *emys*, turtle; referring to the flattened head and shell.

Platyemys Agassiz 1847:294,295 (*nomen novum*)

Etymology:* Greek, *platys*, flat, and *emys*, turtle; referring to the flattened head and shell.

Comment:* Unjustified emendation of *Platemys*.

Platemys platycephala (Schneider 1792) (14:41, 17:91)

Twist-necked Turtle

(includes 2 subspecies)



(subspecies: *platycephala* = red, *melanonota* = blue)

Distribution: Bolivia (*Beni*, *La Paz*, *Pando*, *Santa Cruz*), Brazil (*Acre*, *Amapá*, *Amazonas*, *Maranhão*, *Mato Grosso*, *Pará*, *Rondônia*, *Roraima*), Colombia (*Amazonas*, *Caquetá*, *Guainía*, *Guaviare*, *Meta*, *Putumayo*, *Vaupés*, *Vichada*), Ecuador, French Guiana, Guyana, Peru (*Amazonas*, *Huánuco*, *Loreto*, *Madre de Dios*, *Puno*, *Ucayali*), Suriname, Venezuela (*Amazonas*, *Bolívar*, *Delta Amacuro*, *Monagas*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 5,283,733 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 7,377,448 sq. km

Size (Max SCL): male 18.0 cm, female 16.7 cm (see subspp.)


IUCN Red List:* Least Concern (LC) (Cunha, Brito, Valadão, and Rhodin, in press). Previously: TFTSG Provisional Red List: Least Concern (LC) (2011); Least Concern (LC) [Not Listed] (Behler 1996)

Platemys platycephala platycephala (Schneider 1792)


Eastern Twist-necked Turtle, Common Twist-necked Turtle



Russell A. Mittermeier / nr. Leticia, Colombia



Anders G.J. Rhodin / Suriname



(subspecies: *platycephala* = red, *melanonota* = blue; stars = type localities)

Distribution: Bolivia (*Beni*, *La Paz*, *Pando*, *Santa Cruz*), Brazil (*Acre*, *Amapá*, *Amazonas*, *Mato Grosso*, *Pará*, *Roraima*), Colombia (*Amazonas*, *Caquetá*, *Putumayo*, *Vaupés*), French Guiana, Guyana, Peru (*Huánuco*, *Loreto*, *Madre de Dios*, *Puno*, *Ucayali*), Suriname, Venezuela (*Amazonas*, *Bolívar*, *Delta Amacuro*, *Monagas*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 5,097,698 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 7,223,458 sq. km

Size (Max SCL): male 18.0 cm, female 16.5 cm (Ceballos et al. 2013)

Synonymy:

Testudo platycephala Schneider 1792:261

Platemys platycephala, *Platemys platycephala platycephala*

Type locality: "Ost-indien." Restricted to "Cayenne, French Guiana" by Ernst (1984:350) [yellow star].

Type specimen: Not located, type specimen figured (pl.7), iconotype.

Etymology:* Greek, *platos*, flat, and *kephale*, head; referring to the flattened head.

Testudo planiceps Schneider 1792:pl.7 (senior homonym and *nomen novum*, not = *Testudo planiceps* Gray 1854b [= *Chelonoidis porteri*])

Emys planiceps, *Hydraspis planiceps*, *Clemmys planiceps*, *Platemys planiceps*, *Chelys (Hydraspis) planiceps*, *Chelys planiceps*

Type locality: "Ost-indien." Restricted to "Cayenne, French Guiana" by Ernst (1984:350) [yellow star].
Type specimen: Not located, type specimen figured (pl.7), iconotype.
Etymology:* Latin, *planis*, flat, plane, and *ceps*, head; referring to the flattened head.
Comment: Erroneous alternate name for *Testudo platycephala*.

Testudo martinella Daudin 1802:377
Platemys martinella
Type locality: "Cayenne et dans l'intérieur de la Guiane" [French Guiana] [yellow star].
Type specimen: MNHN 8760, holotype, see Ernst (1987).
Etymology:* Eponym honoring Joseph Martin (1788–1826), France / French Guiana; botanist and director of the botanical garden in Cayenne, French Guiana, who collected the type.

Emys discolor Thunberg *in* Schweigger 1812:302,348
Hydraspis (Phrynops) discolor, Hydraspis discolor
Type locality: Not known. Restricted to "Cayenne, French Guiana" by Ernst (1984:350) [yellow star].
Type specimen: UPSZTY 723 (formerly UUZM 723), holotype, listed by Thunberg (1828), but not listed by Lönnberg (1896), Andersson (1900), Holm (1957), or Wallin (2001); discovered and identified by A.G.J. Rhodin (unpubl. data).
Etymology:* Latin, *dis-*, twice, and *color*, hue or tint; referring to the two colors of the carapace and plastron.

Emys canaliculata Spix 1824:10
Hydraspis canaliculata, Platemys canaliculata
Type locality:* "fluminis Solimoens" [Rio Solimões, Amazonas, Brazil] [= between Tefé and Tabatinga, possibly near São Paulo de Olivença] [green star].
Type specimen: ZSM 3007/0, lectotype, designated by Hoogmoed and Gruber (1983:349), see also Franzen and Glaw (2007).
Etymology:* Latin, *canalicus*, small canal; referring to the noticeable midline vertebral depression.

Emys carunculata Saint-Hilaire *in* Cuvier 1829:11 (*nomen nudum et dubium*) (14:41)
Etymology:* Latin, *caruncula*, fleshy papillae; referring to the small blunt tubercles on the dorsal neck.


Emys constricta Cuvier *in* Gray 1830e:17 (*nomen nudum*)
Etymology:* Latin, *constricta*, narrow, contracted, compact; referring to the relatively narrow nuchal scute.

Hydraspis pachyura Boie *in* Gray 1830e:17 (*nomen nudum*)
Emys pachyura
Etymology:* Greek, *pachys*, thick, coarse, and *ura*, tail; referring to the relatively broad tail.

Hydraspis constricta Gray 1831d:43
Type locality: Not known. Restricted to "Belem, Brazil" by Ernst (1984:350) [blue star].
Type specimens: MNHN, holotype or syntypes, apparently lost.
Etymology:* Latin, *constricta*, narrow, contracted, compact; referring to the relatively narrow nuchal scute.

Platemys platycephala melanonota Ernst 1984 (17:91)
Black-backed Twist-necked Turtle


James H. Harding / No data / captivity


**
(subspecies: *platycephala* = red, *melanonota* = blue; star = type locality)

Distribution: Colombia (*Putumayo*), Ecuador, Peru (*Amazonas, Loreto*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 186,035 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 219,890 sq. km

Size (Max SCL): male 16.8 cm, female 16.7 cm (Ernst 1984)

Synonymy:

Platemys platycephala melanonota Ernst 1984:352
Type locality: "vicinity of Galilea, on the Río Santiago, Amazonas, Perú (4°1' S, 77°47' W)" [yellow star].
Type specimen: USNM 224136, holotype, see Reynolds et al. (2007).
Etymology:* Greek, *melanos*, black, dark, and *notos*, the back; referring to the dark carapace.

Ranacephala McCord, Joseph-Ouni, and Lamar 2001 (07:100, 21:3)

Frog-headed Turtles

(includes 1 species, 1 taxon)

Ranacephala McCord, Joseph-Ouni, and Lamar 2001:732

Type species: *Ranacephala hogei* [= *Phrynops hogei* Mertens 1967a], by original monotypy and designation.


Etymology:* Latin, *rana*, frog, and Greek, *kefale*, head; referring to the froglike head.

Ranacephala hogei (Mertens 1967a)


Hoge's Side-necked Turtle



Russell A. Mittermeier / TCF / TCC / Espírito Santo, Brazil



Russell A. Mittermeier / TCF / TCC / Espírito Santo, Brazil



** (orange dots and star = uncertain or in error; star = type locality)

Distribution: Brazil (*Espírito Santo, Minas Gerais, Rio de Janeiro, São Paulo [?]*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 19,005 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 45,212 sq. km

Size (Max SCL): male 38.0 cm, female 34.0 cm (Moreira 2002; A.G.J. Rhodin and R.A. Mittermeier, unpubl. data)

IUCN Red List: Critically Endangered (CR A2bcd+4bcd) (Drummond, Coutinho, and Vogt 2016), as *Mesoclemmys hogei*. Previously: Endangered (EN) (TFTSG 1996)

USA ESA:* Endangered, as *Phrynops hogei* (USFWS 1991b)

Synonymy:

Phrynops hogei Mertens 1967a:73

Ranacephala hogei, Mesoclemmys hogei

Type locality: "Rio Pequena, südwestlich von São Paulo, Brasilien" [Brazil] [locality in error]. Emended to "Rio Pequeno, southeast of São Paulo" [Brazil] [locality probably in error] by Rhodin et al. (1982:179) [orange star].

Type specimen: SMF 62530, holotype, see Mertens (1967b).

Etymology:* Eponym honoring Alphonse R. Hoge (1912–1982), Brazil; herpetologist at the Instituto Butantan, São Paulo, where the type specimen was first located.

Rhinemys Wagler 1830b (07:100)

Red Side-necked Turtles

(includes 1 species, 1 taxon)

Rhinemys Wagler 1830b:134

Type species: *Rhinemys rufipes* [= *Emys rufipes* Spix 1824], by subsequent designation by Fitzinger (1843:29).


Etymology:* Greek, *rhinos*, nose, and *emys*, turtle; referring to the somewhat prominent nose.

Rhinemys rufipes (Spix 1824)


Red Side-necked Turtle, Red-footed Sideneck Turtle, *Cágado-Vermelho*



William E. Magnusson / CBFTT / Presidente Figuero, Amazonas, Brazil



left: Richard C. Vogt / CBFTT / Reserva Ducke, Amazonas, Brazil
right: Russell A. Mittermeier / Amazonas, Brazil



** (orange dot = probable trade; star = type locality)

Distribution: Brazil (*Amazonas, Pará*), Colombia (*Amazonas, Guainía, Vaupés*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 458,107 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 940,213 sq. km

Size (Max SCL): male 23.0 cm, female 25.6 cm (Magnusson and Vogt 2014 CBFTT)

CBFTT Account: Magnusson and Vogt (2014)

IUCN Red List:* Least Concern (LC) (Ferrara, Brito, Forero-Medina, Cunha, and Vogt, in press). Previously: TFTSG Provisional Red List: Least Concern (LC) (2011); Near Threatened (NT) (TFTSG 1996)

Synonymy:

Emys rufipes Spix 1824:7

Hydraspis rufipes, Rhinemys rufipes, Chelys (Hydraspis) rufipes, Chelys rufipes, Platemys rufipes, Phrynops rufipes

Type locality:* "fluminis Solimoëns" [Rio Solimões, Amazonas,

Brazil] [= between Tefé and Tabatinga, possibly near Tefé] [yellow star].
Type specimen: ZSM 3006/0, holotype, see Hoogmoed and Gruber (1983), Franzen and Glaw (2007), and Ettmar (2019).
Etymology:* Latin, *rufa*, red, red-brown, and *pes*, foot; referring to the red extremities.

Subfamily Hydromedusinae Baur 1893a (12:39)

South American Snake-necked Turtles
(includes 1 genus, 2 species, 2 taxa)
Hydromedusidae Baur 1893a:211
Hydromedusinae Georges, Birrell, Saint, McCord, and Donnellan 1998:235

Hydromedusa Wagler 1830b

South American Snake-necked Turtles
(includes 2 species, 2 taxa)
Hydromedusa Wagler 1830b:135
Type species: *Hydromedusa maximiliani* [= *Emys maximiliani* Mikan 1825], by original monotypy.
Etymology:* Greek, *hydor*, water, and *medusa*, a mythical beast (a Gorgon) with a head of snakes instead of hair; referring to the long snake-like neck of this turtle genus.
Chelomedusa Gray 1873c:303
Type species: *Hydromedusa (Chelomedusa) depressa* [= *Hydromedusa depressa* Gray 1856b] [= subjective synonym of *Emys maximiliani* Mikan 1825], by subsequent designation by Lindholm (1929:289). Genus established as *Hydromedusa (Chelomedusa)* without a type species.
Etymology:* Greek, *chelys*, a turtle, and *medusa*, a mythical beast (a Gorgon) with a head of snakes instead of hair; referring to the long snake-like neck of this turtle genus.

Hydromedusa maximiliani (Mikan 1825)

Brazilian Snake-necked Turtle, Maximilian's Snake-necked Turtle, *Cágado-da-Serra*



Franco L. Souza / CBFTT / Parque Estadual Carlos Botelho, São Paulo, Brazil



Franco L. Souza / CBFTT / Parque Estadual Carlos Botelho, São Paulo, Brazil



** (star = type locality)


Distribution:* Brazil (*Bahia, Espírito Santo, Minas Gerais, Paraná, Rio de Janeiro, São Paulo*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 288,138 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 560,752 sq. km
Size (Max SCL): male 17.0 cm, female 20.0 cm (Souza and Abe 1997; Souza and Martins 2009 CBFTT)
CBFTT Account: Souza and Martins (2009)
IUCN Red List:* Endangered (EN A2ce+4ce) (Famelli, Souza, Valadão, and Brito, in press). Previously: TFTSG Provisional Red List: Near Threatened (NT) (2011); Vulnerable (VU B1+2cd [ver. 2.3]) (TFTSG 1996)
Synonymy:
Emys maximiliani Mikan 1825:[unpaginated]
Chelodina maximiliani, Hydromedusa maximiliani, Hydraspis maximiliani
Type locality: "Brasiliam...Capitania St. Paulo" [São Paulo, Brazil]. Restricted to "auf dem Weg nach Portofeliz" [nr. Pôrto Feliz, São Paulo, Brazil] by J. Natterer (1820, unpubl. data) *in* Gemel et al. (2019:80) [yellow star].
Type specimen: NMW 23391, holotype, see Tiedemann and Häupl (1980), Tiedemann et al. (1994), and Gemel et al. (2019).
Etymology:* Eponym honoring Prince Alexander Philipp Maximilian zu Wied-Neuwied (1782–1867), Germany; nobleman, naturalist and explorer of Brazil and the United States.
Emys maximiliana Gray 1830e:17 (*nomen novum*)
Hydromedusa maximiliana
Etymology:* Eponym honoring Prince Alexander Philipp Maximilian zu Wied-Neuwied (1782–1867), Germany; nobleman, naturalist and explorer of Brazil and the United States.
Comment: Unjustified emendation or error for *maximiliani*.
Chelodina flavilabris Duméril and Bibron 1835:446
Hydromedusa flavilabris, Chelomedusa flavilabris
Type locality: "Brésil" [Brazil].
Type specimen: MNHN 9411, holotype.
Etymology:* Latin, *flavus*, yellow, and *labrum*, lip; referring to the cream-colored to yellow jaws.
Hydromedusa subdepressa Gray 1854a:134
Type locality: "Brazils" [Brazil].
Type specimen: NHMUK 1946.1.22.57 (formerly 1849.11.7.29), holotype.
Etymology:* Latin, *sub-*, below, almost, or near, *de-*, down, and *pressus*, pressed; referring to the nearly depressed flattened shell.
Hydromedusa depressa Gray 1856b:60 (*nomen novum*)
Chelomedusa depressa
Type locality: "Brazils" [Brazil].
Type specimen: NHMUK 1946.1.22.57 (formerly 1849.11.7.29), holotype, by default.
Etymology:* Latin, *de-*, down, and *pressus*, pressed; referring to the depressed flattened shell.
Comment: Unjustified emendation or error for *subdepressa*.
Hydromedusa bankae Giebel 1866b:19

Type locality: “Insel Banka” [Bangka Island, Sumatra, Indonesia] [in error].
Type specimen: Possibly MLUH, holotype, not located, type specimen figured (pl.4).
Etymology:* Toponym for Banka Island (= Bangka Island, Sumatra, Indonesia).


Hydromedusa tectifera Cope 1870a
South American Snake-necked Turtle, Argentine Snake-necked Turtle, *Tortuga Cuello de Vibora*



Peter C.H. Pritchard / captivity / Brazil



left: Raissa Bressan / Rio Grande do Sul, Brazil / male
right: Andreas Nöllert / No data / captivity



** (stars = type localities)

Distribution: Argentina (*Buenos Aires, Corrientes, Entre Rios, Misiones, Salta, San Luis, Santa Fe, Santiago del Estero*), Brazil (*Minas Gerais, Paraná, Rio de Janeiro, Rio Grande do Sul, Santa Catarina, São Paulo*), Paraguay, Uruguay
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 869,546 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 2,746,802 sq. km
Size (Max SCL): male 28.4 cm, female 30.6 cm (Bager et al. 2003; Chinen et al. 2004; Regis and Meik 2017; Alcalde et al. 2021 CBFTT)
CBFTT Account: Alcalde, Sánchez, and Pritchard (2021)
IUCN Red List:* Least Concern (LC) (Alcalde, Fagundes, and Carreira, in press). Previously: TFTSG Provisional Red List: Least Concern (LC) (2011); Least Concern (LC) [Not Listed] (Behler 1996)
Synonymy:
Hydromedusa tectifera Cope 1870a:147
Type locality: “tributaries of the Parana or Uraguay rivers, either in the Argentine Confederation or the Banda Oriental” [Argentina or Uruguay] [yellow star, location estimated].
Type specimen: Not located, holotype in either USNM or ANSP, apparently lost, not recorded by Cochran (1961), Malnate (1971), or Reynolds et al. (2007). NHMUK 1947.3.5.85, erroneously listed as “type” by Uetz et al. (2019), is the holotype of *Hydromedusa platanensis* Gray 1873c.
Etymology:* Latin, *tectum*, roof, and *ferre*, to bear; referrring to the roof-like extended front edge of the carapace.
Hydromedusa platanensis Gray 1873c:302
Hydromedusa platensis
Type locality: “Rio de la Plata” [Argentina or Uruguay] [yellow star].
Type specimen: NHMUK 1947.3.5.85 (formerly 1854.8.16.11), holotype.
Etymology:* Toponym for Rio de la Plata, Argentina.
Platemys antiqua † Ameghino 1882:41 (*nomen nudum*)
Type locality: “Provincia Buenos Aires” [Argentina].
Type specimen: Not designated, fossil.
Geological age: Pleistocene or Holocene.
Etymology:* Latin, *antiqua*, old, ancient; referring to this taxon being classified as extinct.
Platemys fossilis † Ameghino 1882:41 (*nomen nudum*)
Type locality: “Provincia Buenos Aires” [Argentina].
Type specimen: Not designated, fossil.
Geological age: Pleistocene or Holocene.
Etymology:* Latin, *fossilis*, excavated, petrified; referring to this taxon being a fossil.
Platemys laevis † Ameghino 1882:41 (*nomen nudum*)
Type locality: “Provincia Buenos Aires” [Argentina].
Type specimen: Not designated, fossil.
Geological age: Pleistocene or Holocene.
Etymology:* Latin, *laevis*, smooth, delicate; referring possibly to this taxon having had a smooth or smaller shell.
Platemys robusta † Ameghino 1882:41 (*nomen nudum*)
Type locality: “Provincia Buenos Aires” [Argentina].
Type specimen: Not designated, fossil.
Geological age: Pleistocene or Holocene.
Etymology:* Latin, *robusta*, smooth, delicate; referring possibly to this taxon having had a robust or larger shell.
Hydromedusa wagleri Günther 1884:423
Type locality: “Buenos Ayres” [Buenos Aires, Argentina] [green star].
Type specimen: NHMUK 1947.3.5.84 (formerly 1848.2.3.3), holotype.
Etymology:* Eponym honoring Johann G. Wagler (1800–1832), Germany; ornithologist and herpetologist at the University of München.

Subfamily Chelodininae Baur 1893a (12:39, 21:17)

Australasian Snake-necked Turtles

(includes 1 genus, 15 species, 16 taxa)

Chelodinidae Baur 1893a:211

Chelodininae Georges, Birrell, Saint, McCord, and Donnellan 1998:235

Chelodina Fitzinger 1826 (07:85, 08:2, 10:34, 21:5, 21:17) (10)*

Australasian Snake-necked Turtles

(includes 3 subgenera, 15 species, 16 taxa)

Chelodina Fitzinger 1826:6

Type species: *Chelodina longicollis* [= *Testudo longicollis* Shaw 1794], by original designation.

Etymology:* Greek, *chelys*, turtle, and *deinos*, terrible, referring to the long snake-like neck.

Hydraspis Bell 1828b:512

Type species: *Hydraspis longicollis* [= *Testudo longicollis* Shaw 1794], by original designation.

Etymology:* Greek, *hydor*, water, and *aspis*, viper; referring to the long snake-like neck.

Chelyodina Agassiz 1847:79 (*nomen novum*)

Etymology:* Greek, *chelys*, turtle, and *deinos*, terrible, referring to the long snake-like neck.

Comment:* Invalid replacement name for *Chelodina*. Frétey (2024) noted that this publication appeared in 1847, not 1846.

Hesperochelodina Wells and Wellington 1985:9 (*nomen nudum*)

Etymology:* Greek, *hesperos*, west, western, and *chelodina*, turtle genus; referring to its distribution in Western Australia.

Macrochelodina Wells and Wellington 1985:9

Type species: *Macrochelodina oblonga* [= *Chelodina oblonga* Gray 1841], by original designation.

Etymology:* Greek, *makros*, large, and *chelodina*, turtle genus; referring to the larger, thicker neck in this genus.

Macrodiremys McCord and Joseph-Ouni 2007b:57

Type species: *Macrodiremys oblonga* [= *Chelodina oblonga* Gray 1841], by original designation.

Etymology:* Greek *makros*, large, and *deire*, neck, and *emys*, turtle; referring to the larger, thicker neck in this genus.

Chelydera Thomson and Georges *in* Shea, Thomson, and Georges 2020:430

Type species: *Chelodina (Chelydera) parkeri* [= *Chelodina parkeri* Rhodin and Mittermeier 1976], by original designation.

Etymology:* Greek, *chelys*, turtle, and *deire*, neck, referring to the long neck.

Chelodina (Chelodina) Fitzinger 1826 (10:34, 21:17) (10)*

Australasian Long-necked Turtles

(includes 8 species, 9 taxa)

Chelodina Fitzinger 1826:6

Type species: *Chelodina longicollis* [= *Testudo longicollis* Shaw 1794], by original designation.

Etymology:* Greek, *chelys*, turtle, and *deinos*, terrible, referring to the long snake-like neck.

Hydraspis Bell 1828b:512

Type species: *Hydraspis longicollis* [= *Testudo longicollis* Shaw 1794], by original designation.

Etymology:* Greek, *hydor*, water, and *aspis*, viper; referring to the long snake-like neck.

Chelyodina Agassiz 1847:79 (*nomen novum*)

Etymology:* Greek, *chelys*, turtle, and *deinos*, terrible, referring to the long snake-like neck.

Comment:* Invalid replacement name for *Chelodina*. Frétey (2024) noted that this publication appeared in 1847, not 1846.

Hesperochelodina Wells and Wellington 1985:9 (*nomen nudum*)

Etymology:* Greek, *hesperos*, west, western, and *chelodina*, turtle genus; referring to its distribution in Western Australia.

Chelodina (Chelodina) canni McCord and Thomson 2002 (07:86, 21:17)

Cann's Long-necked Turtle, Cann's Snake-necked Turtle




John Cann / Kalala Station, Roper R. drainage, Northern Territory, Australia





left: Jason Schaffer / Rinyirru National Park, Queensland, Australia
right: Arthur Georges / Sunday Creek, Northern Territory, Australia



** (stars = type localities)

Distribution: Australia (*Northern Territory, Queensland*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 261,592 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 1,107,085 sq. km

Size (Max SCL): male 16.9 cm, female 27.9 cm (Ceballos et al. 2013; Cann and Sadlier 2017; A.G.J. Rhodin, unpubl. data)

IUCN Red List: Not Evaluated (NE)

TFTSG Provisional Red List: Near Threatened (NT) (2009)

Synonymy:

Chelodina rankini Wells and Wellington 1985:8 (*nomen nudum*)

Type locality: "Lower Burdekin River, north east Queensland" [Australia] [orange star].

Etymology:* Eponym honoring Peter R. Rankin (1955–1979), Australia; amateur herpetologist.

Chelodina canni McCord and Thomson 2002:256

Chelodina (Chelodina) canni, *Chelodina novaeguineae canni*, *Chelodina (Chelodina) canni canni*

Type locality: "Malogie Waterhole, near Scarlet Hill on Kalala Station (16°08' S, 133°36' E), Northern Territory, Australia" [yellow star].

Type specimen: NTM 24515, holotype; see Cann and Sadlier (2017).

Etymology:* Eponym honoring John R. Cann (1938–), Australia; herpetologist who studied turtles extensively in Australia and New Guinea.

Chelodina rankini Wells 2007a:2 (unavailable name) (07:86, 10:43)


Chelodina canni rankini, *Chelodina (Chelodina) canni rankini*

Type locality: "north-eastern Australia...Burdekin River, Queensland"

[orange star].
Type specimen: NHMUK 1908.2.25.1, holotype.
Etymology:* Eponym honoring Peter R. Rankin (1955–1979), Australia; amateur herpetologist.

Chelodina (Chelodina) gunaleni McCord and Joseph-Ouni 2007a (10:35, 21:18)
Gunalen's Snake-necked Turtle


William P. McCord / Uta R., Papua, Indonesia





(star = type locality)

Distribution: Indonesia (*Papua*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 23,956 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 34,263 sq. km
Size (Max SCL): male 16.7 cm, female 23.9 cm (McCord and Joseph-Ouni 2007a)
IUCN Red List: Not Evaluated (NE)
TFTSG Provisional Red List: Data Deficient (DD) (2009)
Synonymy:
Chelodina gunaleni McCord and Joseph-Ouni 2007a:48
Chelodina (Chelodina) gunaleni
Type locality: "Uta River basin, Mimika District, Central Papua Province (Irian Jaya), Indonesia" [yellow star].
Type specimen: AMNH 160133, holotype; genotyped by Kehlmaier et al. (2019a).
Etymology:* Eponym honoring Danny Gunalen (1975–), Indonesia; reptile exporter and herpetologist who obtained the type.

Chelodina (Chelodina) longicollis (Shaw 1794) (17:92)
Eastern Long-necked Turtle, Eastern Snake-necked Turtle


John Roe / CBFTT / Booderee National Park, Jervis Bay, New South Wales, Australia


Arthur Georges / CBFTT / Australia



(orange dots = probably introduced; star = type locality)

Distribution: Australia (*New South Wales, Queensland, South Australia, Victoria*)
Introduced: Australia (*Tasmania*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 895,986 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 2,376,875 sq. km
Size (Max SCL): male 24.9 cm, female 28.2 cm (Kennett et al. 2009 CBFTT)
CBFTT Account: Kennett, Roe, Hodges, and Georges (2009)
IUCN Red List: Least Concern (LC) [Not Listed] (Behler 1996)
TFTSG Provisional Red List: Least Concern (LC) (2009)
Synonymy:
Testudo longicollis Shaw 1794:19
Emys longicollis, Chelodina longicollis, Hydraspis longicollis, Chelys (Chelodina) longicollis, Chelys longicollis, Chelodina longicollis longicollis, Chelodina (Chelodina) longicollis
Type locality: "The vast island or rather continent of Australia, Australasia, or New Holland." Type locality presumed to be near Botany Bay or Sydney, New South Wales (Cann and Sadlier 2017:17), but not formally restricted [yellow star].


Type specimen: NHMUK 1947.3.5.86, holotype, designated by Cogger et al. (1983:61), see Cann and Sadlier (2017); genotyped by Kehlmaier et al. (2019a).
Etymology:* Latin, *longus*, long, and *collum*, neck; referring to the long neck.

Chelodina novaehollandiae Duméril and Bibron 1835:443 (*nomen novum*)
Chelodina novae-hollandiae
Type locality: "Nouvelle-Hollande" [Australia].
Type specimen: NHMUK 1947.3.5.86, holotype by default; MNHN 6969, recorded as "type" by Roux-Esteve (1979), and "holotype" by Cogger et al. (1983), Georges and Thomson (2010), and Cann and Sadlier (2017), is not a type, since *C. novaehollandiae* is a *nomen novum* replacement name for *C. longicollis*.
Etymology:* Toponym for New Holland (= Australia).

Chelodina sulcata Gray 1856a:201
Chelodina longicollis sulcata
Type locality: "Australia."
Type specimen: NHMUK 1947.3.5.87 (formerly 1851.4.24.27), holotype.
Etymology:* Latin, *sulcus*, groove or furrow, and *-ata*, having; referring to the concentric furrows on the carapace.

Chelodina sulcifera Gray 1856b:59 (*nomen novum*)
Chelodina longicollis sulcifera
Type locality: "Australia."
Type specimen: NHMUK 1947.3.5.87 (formerly 1851.4.24.27), holotype by default, see Goode (1967), Cogger et al. (1983), and Cann and Sadlier (2017).
Etymology:* Latin, *sulcus*, groove or furrow, and *-fera*, bearing; referring to the concentric furrows on the carapace.

Chelodina (Chelodina) mccordi Rhodin 1994b (10:36, 21:19) (11)*
Roti Island Snake-necked Turtle; Rote Snake-necked Turtle
(includes 2 subspecies)




(subspecies: *mccordi* = red; *timorensis* = purple)

Distribution: Indonesia (*Lesser Sundas [Roti = Rote]*), Timor-Leste
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 466 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 7,218 sq. km
Size (Max SCL): male 20.2 cm, female 24.1 cm (see subspp.)
CBFTT Account: Rhodin, Ibarrondo, and Kuchling (2008)
IUCN Red List: Critically Endangered (CR A2acde) (As-singkily, Eisemberg, Horne, Kuchling, and Rhodin 2019).
Previously: Critically Endangered (CR) (ATTWG 2000); Vulnerable (VU) (TFTSG *in* Baillie and Groombridge 1996)
CITES: Appendix II (2005); zero export quota for specimens from the wild (2013)


Chelodina (Chelodina) mccordi mccordi Rhodin 1994b (10:36, 21:19) (11)*
Roti Island Snake-necked Turtle, Rote Snake-necked Turtle



Anders G.J. Rhodin / TCF / CBFTT / nr. Busalangga, Roti, Indonesia



Anders G.J. Rhodin / CBFTT / nr. Danau Peto, Roti, Indonesia



(stars = type localities)

Distribution: Indonesia (*Lesser Sundas [Roti = Rote]*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 242 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 623 sq. km
Size (Max SCL): male 20.2 cm, female 24.1 cm (Rhodin et al. 2008 CBFTT)
IUCN Red List: Critically Endangered (Possibly Extinct in the Wild) (CR(PEW) A2acde+B2ab(ii,iv)) (As-singkily, Eisemberg, Horne, Kuchling, and Rhodin 2019)
IUCN Green List:* Critically Depleted (CD) (Horne and Bennett 2025)
CITES: Appendix II (2005); zero export quota for specimens from the wild (2013)
Synonymy:

Chelodina rottiensis Brongersma *in* Rhodin 1994b:3 (*nomen nudum*)
Etymology:* Toponym for Roti, Indonesia.

Chelodina mccordi Rhodin 1994b:4
Chelodina mccordi mccordi, *Chelodina (Chelodina) mccordi*, *Chelodina (Chelodina) mccordi mccordi*
Type locality: "Danau Naloek, near Busalangga, ca. 11 km northeast of Tudameda and ca. 8 km southwest of Ba'a, elevation ca. 115 m, southwestern Roti Island (10°48' S, 123°00' E), East Nusa Tenggara Province, Indonesia" [yellow star].
Type specimen: MCZ 176730, holotype; MCZ 176731, paratype, genotyped by Kehlmaier et al. (2019a).
Etymology:* Eponym honoring William P. McCord (1950–), USA; veterinarian and herpetologist, who obtained additional specimens

of this species from the trade.

Chelodina mccordi roteensis McCord, Joseph-Ouni, and Hagen 2007b:59 (07:87, 10:36, 21:19) (11)*

Chelodina (Chelodina) mccordi roteensis, Chelodina (Chelodina) mccordi rotiensis

Type locality: "Lake Enduy, eastern Rote Island, East Nusa Tenggara Province, Indonesia" [green star].

Type specimen: AMNH R160132, holotype; genotyped by Kehlmaier et al. (2019a).

Etymology:* Toponym for Rote (= Roti), Indonesia.

Chelodina (Chelodina) mccordi timorensis McCord, Joseph-Ouni, and Hagen 2007a (07:89, 10:36, 21:19) (11)*


Timor Snake-necked Turtle, Timor-Leste Snake-necked Turtle



Bonggi R. Ibarrondo / CCB / Lospalos, Lautém Dist., Timor-Leste



left: Carla C. Eisemberg / CCB / Lake Iralalaro, Lautém Dist., Timor-Leste
right: Bonggi R. Ibarrondo / CCB / Lospalos, Lautém Dist., Timor-Leste



(star = type locality)

Distribution: Timor-Leste

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO): 224 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 245 sq. km

Size (Max SCL): male 20.0 cm, female 23.3 cm (Kuchling et al. 2007)

IUCN Red List: Critically Endangered (CR B1ab(ii,iv)) (Assingkily, Eisemberg, Horne, Kuchling, and Rhodin 2019)

CITES: Appendix II (2005); zero export quota for specimens from the wild (2013)

Synonymy:

Chelodina timorensis McCord, Joseph-Ouni, and Hagen 2007a:54

Chelodina mccordi timorensis, Chelodina (Chelodina) mccordi timorensis, Chelodina (Chelodina) timorensis

Type locality: "Lake Ira Lalaro, Lautem District (regency), Tutuala Subdistrict, eastern East Timor (= Timor-Leste; = Timor Lorosa'E)" [yellow star].

Type specimen: AMNH 160135, holotype; genotyped by Kehlmaier et al. (2019a).

Etymology:* Toponym for Timor, Indonesia.

Chelodina mccordi timorlestensis Kuchling, Rhodin, Ibarrondo, and Trainor 2007:213

Type locality: "plain of Lake Iralalaro (= Lagoa Ira Lalaro) (ca. 08º28' S; 127º07' E, elev. ca. 334 m), east of Lospalos, Lautém District, Timor-Leste" [yellow star].

Type specimen: WAM 165888, holotype, see Ellis and Georges (2015); genotyped by Kehlmaier et al. (2019a).


Etymology:* Toponym for Timor-Leste.

Chelodina (Chelodina) novaeguineae Boulenger 1888b (21:20) (12)*


New Guinea Snake-necked Turtle



Fred Parker / Abam, Western Prov., Papua New Guinea



Anders G.J. Rhodin / Binaturi R., Western Prov., Papua New Guinea



** (stars = type localities)

Distribution: Indonesia (*Papua*), Papua New Guinea (*Southern*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 75,339 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 105,602 sq. km

Size (Max SCL): male 15.1 cm, female 21.8 cm (A.G.J. Rhodin and P.M. Hall, unpubl. data)

IUCN Red List: Least Concern (LC) (ATTWG 2000). Previously: Least Concern (LC) [Not Listed] (Behler 1996)

TFTSG Provisional Red List: Least Concern (LC) (2009)

Synonymy:

Chelodina novaeguineae Boulenger 1888b:450

Chelodina novaeguineae novaeguineae, Chelodina (Chelodina) novaeguineae

Type locality: "Katow, S. E. New Guinea" [Papua New Guinea].

Emended to "Mawatta, Binaturi River (as Katow), Papua New Guinea" by Wells and Wellington (1985:8), and "Mawatta, Binaturi River, Western Province, Papua New Guinea" by Rhodin (1994a:9) [yellow star].

Type specimen: NHMUK 1946.1.22.36 (formerly 1888.5.21.7), lectotype, designated by Wells and Wellington (1985:8), see also Rhodin (1994a); genotyped by Kehlmaier et al. (2019a).

Etymology:* Toponym for New Guinea.

Chelodina (Chelodina) ipudinapi Joseph-Ouni and McCord 2022a:9 (12)*

Chelodina ipudinapi

Type locality: "vicinity of Balimo, Aramia River, Western Province, Papua New Guinea" [green star].

Type specimen: YPM HERR.022303, holotype.

Etymology:* Vernacular (New Guinean, Gogodala), *ipudinapi*, little stinker, local indigenous name for the species; referring to the particularly strong and foul odor emitted when captured.


Chelodina (Chelodina) pritchardi Rhodin 1994a
Pritchard's Snake-necked Turtle



Anders G.J. Rhodin / Bore, Kemp Welch R., Central Prov., Papua New Guinea / holotype



Anders G.J. Rhodin / Bore, Kemp Welch R., Central Prov., Papua New Guinea / holotype



(star = type locality)

Distribution: Papua New Guinea (*Southern*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 6,947 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 11,045 sq. km

Size (Max SCL): male 18.6 cm, female 22.8 cm (Rhodin 1994a)

IUCN Red List: Vulnerable (VU B1ab(ii,iii,v)+2ab(ii,iii,v)) (Rhodin and Georges 2020). Previously: Endangered (EN) (ATTWG 2000); Vulnerable (VU) (TFTSG *in* Baillie and Groombridge 1996)

Synonymy:

Chelodina pritchardi Rhodin 1994a:4
Chelodina (Chelodina) pritchardi

Type locality: "Bore, Kemp Welch River, 13 km southeast of Kwikila, Central Province, Papua New Guinea (9°53' S, 147°46' E)" [yellow star].

Type specimen: MCZ 173543, holotype, specimen figured (f.1) and in Rhodin and Genorupa (2000:f.7) and here (TTWG:above), subsequently badly damaged in fire (Rhodin, pers. comm.); MCZ 175813 and AMNH 139735, paratypes.

Etymology:* Eponym honoring Peter C.H. Pritchard (1943–2020), United Kingdom (Northern Ireland) / USA; herpetologist and a Co-Chair of the IUCN Tortoise and Freshwater Turtle Specialist Group who founded Chelonian Research Institute, Florida, and first collected this species.


Chelodina (Chelodina) reimanni Philippen and Grossmann 1990 (21:20)
Reimann's Snake-necked Turtle



Anders G.J. Rhodin / nr. Merauke, Papua, Indonesia / captivity



Anders G.J. Rhodin / nr. Merauke, Papua, Indonesia / captivity



(star = type locality)

Distribution: Indonesia (*Papua*), Papua New Guinea (*Southern*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 49,192 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 73,219 sq. km

Size (Max SCL): male 20.9 cm, female 22.0 cm (Lyons et al. 2013)

IUCN Red List: Near Threatened (NT) (ATTWG 2000). Previously: Data Deficient (DD) (TFTSG *in* Baillie and Groombridge 1996)

TFTSG Provisional Red List: Data Deficient (DD) (2009)

Synonymy:

Chelodina reimanni Philippen and Grossmann 1990:95
Chelodina novaeguineae reimanni, Chelodina (Chelodina) reimanni


Type locality: "Merauke-River, West-Irian, Neuguinea" [Papua, Indonesia] [yellow star].

Type specimen: MTD 29178, holotype; genotyped by Kehlmaier et al. (2019a); MTD 29241, listed as "holotype" by Uetz et al. (2019), is a paratype.
Etymology:* Eponym honoring Michael J. Reimann (1952–), Germany; herpetologist who first recognized the species as distinct.

Chelodina (Chelodina) steindachneri Siebenrock 1914 (07:85)
Western Long-necked Turtle, Steindachner's Long-necked Turtle


Stephen M. Zozaya / nr. Wodgina, Western Australia, Australia


left: Gerald Kuchling / Borodale R., Western Australia, Australia
right: Brendan Schembri / Indee Station, Western Australia, Australia


(stars = type localities)

Distribution: Australia (*Western Australia*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 626,147 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 815,896 sq. km
Size (Max SCL): male 17.8 cm, female 21.2 cm (Glauert 1923; Cann and Sadlier 2017)
IUCN Red List: Least Concern (LC) [Not Listed] (Behler 1996)
TFTSG Provisional Red List: Least Concern (LC) (2009, 2018)
Synonymy:
Chelodina steindachneri Siebenrock 1914:386
Hesperochelodina steindachneri, Chelodina (Chelodina) steindachneri
Type locality: "Marloo Station am Grey River in Westaustralien" [Western Australia, Australia]. Types possibly from "Marloo Station (28°19'S, 116°10'E) in the Murchison District ca. 50 km west of...Yagloo", but locality not formally restricted (Cann and Sadlier 2017:42) [yellow star].
Type specimens: NMW 1818, 19796–98, syntypes (4), see Gemel et al. (2019), only three of these (19796–98) listed by Tiedemann and Häupl (1980), Tiedemann et al. (1994), and Cann and Sadlier (2017); 19798 listed as "holotype" by Cogger et al. (1983) and Uetz et al. (2019), but has no status as holotype (Gemel et al. 2019).
Etymology:* Eponym honoring Franz Steindachner (1834–1919), Austria; ichthyologist and director of the Naturhistorisches Museum, Vienna.
Chelodina millymillyensis Glauert 1923:54
Chelodina milly-millyensis
Type locality: "Milly Milly, Murchison River, W.A." [Australia]. Emended to "Milly Milly Creek, Milly Milly Station, Murchison River, W.A." [Australia] by Cogger et al. (1983:62) [green star].
Type specimen: WAM R1000, lectotype, designated by Cogger et al. (1983:62), see Ellis and Georges (2015).
Etymology:* Toponym for Milly Milly Station, Western Australia.


Chelodina (Chelydera) Thomson and Georges *in* Shea, Thomson, and Georges 2020 (21:5, 21:17, 21:21) (10)*

Australasian Thick-necked Turtles
(includes 6 species, 6 taxa)
Chelydera Thomson and Georges *in* Shea, Thomson, and Georges 2020:430
Type species: *Chelodina (Chelydera) parkeri* [= *Chelodina parkeri* Rhodin and Mittermeier 1976], by original designation.
Etymology:* Greek, *chelys*, turtle, and *deire*, neck, referring to the long neck.

Chelodina (Chelydera) burrungandjii Thomson, Kennett, and Georges 2000
Arnhem Snake-necked Turtle, Sandstone Snake-necked Turtle


Rod Kennett / Koolpin Gorge, Kakadu National Park, Northern Territory, Australia


Rod Kennett / Koolpin Gorge, Kakadu National Park, Northern Territory, Australia



(star = type locality)


Distribution: Australia (*Northern Territory*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 47,849 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 55,352 sq. km
Size (Max SCL): male 22.0 cm, female 27.1 cm (Thomson et al. 2011 CBFTT)

CBFTT Account: Thomson, Kennett, Tucker, FitzSimmons, Featherston, Alacs, and Georges (2011) [includes *Chelodina (Chelydera) walloyarrina* (= *kuchlingi*)]
IUCN Red List: Not Evaluated (NE)
TFTSG Provisional Red List: Least Concern (LC) (2009)
Synonymy:
Chelodina burrungandjii Thomson, Kennett, and Georges 2000:676
Macrochelodina burrungandjii, Chelodina (Macrochelodina) burrungandjii, Chelodina (Macrochelodina) oblonga burrungandjii, Chelodina (Chelydera) burrungandjii, Chelydera burrungandjii
Type locality: "Koolpin Gorge, South Alligator River (13º28' S, 132º38' E)... Arnhem Land Plateau, Northern Territory, Australia" [yellow star].
Type specimen: NTM 16010, holotype, see Cann and Sadlier (2017).
Etymology:* Vernacular (Australian Aboriginal, Gagadju), *burrungandji*, local indigenous name for the species.

Chelodina (Chelydera) expansa Gray 1857 (17:4, 17:93, 21:5)
Broad-shelled Snake-necked Turtle


Claire Treilibs / CBFTT / Paringa, South Australia, Australia


Deborah Bower / CBFTT / Australia


**
(orange dots = probably introduced or translocated; star = type locality)

Distribution: Australia (*New South Wales, Queensland, South Australia, Victoria*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 316,891 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 1,446,711 sq. km
Size (Max SCL): male 33.7 cm, female 50.0 cm (Cann and Legler 1994; Bower and Hodges 2014 CBFTT; A.G.J. Rhodin, unpubl. data)
CBFTT Account: Bower and Hodges (2014)
IUCN Red List: Least Concern (LC) [Not Listed] (Behler 1996)]
TFTSG Provisional Red List: Near Threatened (NT) (2009)
Synonymy:
Chelodina expansa Gray 1857:370
Chelodina oblonga expansa, Macrochelodina expansa, Chelodina (Macrochelodina) expansa, Chelodina (Chelydera) expansa, Chelydera expansa
Type locality: "Australia." Restricted to "N. Australia" by Gray (1873j:63). Types possibly from the Condamine River in southeastern Queensland, west of the Great Dividing Range, but locality not formally restricted (Cann and Sadlier 2017:98) [yellow star, location estimated].
Type specimen: NHMUK 1947.3.4.21 (formerly 1856.12.3.6), lectotype, designated by Iverson et al. (2001:366); Uetz et al. (2019) erroneously listed NHMUK 1947.3.4.21 and 1947.3.5.88 as syntypes.
Etymology:* Latin, *expansa*, expanded; referring to the broad, wide shell.


Chelodina (Chelydera) kuchlingi Cann 1997d (07:90, 10:37, 14:42, 17:94, 21:23) (13)*
Kuchling's Snake-necked Turtle, Kimberley Snake-necked Turtle



John Cann / TCF / Kimberleys, Western Australia, Australia



Anton Tucker / CBFTT (as *Chelodina burrungandjii*) / Kimberleys, Western Australia, Australia




** (orange dot = questionable; stars = type localities)

Distribution:* Australia (*Northern Territory*, *Western Australia*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 199,504 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 270,910 sq. km
Size (Max SCL):* male 26.3 cm, female 36.0 cm (Thomson et al. 2011 CBFTT; Maier and Sonneman 2018)
CBFTT Account: Thomson, Kennett, Tucker, FitzSimmons, Featherston, Alacs, and Georges (2011) [as part of *Chelodina (Macrochelodina) burrungandjii* account]
IUCN Red List: Not Evaluated (NE)
TFTSG Provisional Red List: Data Deficient (DD) (2025).
Previously: Critically Endangered (CR) (2020)
Synonymy:
Chelodina kuchlingi Cann 1997d:41
Macrochelodina kuchlingi, Chelodina (Macrochelodina) kuchlingi, Chelodina (Chelydera) kuchlingi, Chelydera kuchlingi
Type locality: "Kalumburu, N.W. Australia, (14°18' S x 126°28' E)"; [in error, orange star], restricted to "Parry Creek, lower Ord River floodplain, Kimberley, Western Australia" [Australia] [yellow star] by Kuchling (2020:53).
Type specimen: WAM R29411, holotype; genotyped by Kehlmaier et al. (2019a, 2025).
Etymology:* Eponym honoring Gerald Kuchling (1952–), Austria / Australia; herpetologist, conservationist, and turtle researcher.
Macrochelodina walloyarrina McCord and Joseph-Ouni 2007b:59 (08:4, 10:38, 17:94, 21:17) (13)*
Chelodina (Macrochelodina) walloyarrina, Chelodina (Macrochelodina) oblonga walloyarrina, Chelodina burrungandjii walloyarrina, Chelodina (Chelydera) walloyarrina, Chelydera walloyarrina
Type locality: "The Fitzroy River at Fitzroy River Crossing, Western Australia, Australia, 18°10.834' S and 125°35.849' E" [green star].
Type specimen: WAM R164345, holotype, see Ellis and Georges (2015) and Cann and Sadlier (2017).
Etymology:* Vernacular (Australian Aboriginal), *wallo*, chin, and *yarrin*, bearded; referring to its long chin barbels.


Chelodina (Chelydera) kurrichalpongo (Joseph-Ouni, McCord, Cann, and Smales 2019a) (21:24)
(or *Chelodina (Chelydera) kuchlingi*)
Darwin Snake-necked Turtle



Arthur Georges / CBFTT (as *Chelodina oblonga*) / Blyth-Cadell R., Northern Territory, Australia



left: Arthur Georges / CBFTT (as *Chelodina oblonga*) / Blyth-Cadell R., Northern Territory, Australia
right: John Cann / CBFTT (as *Chelodina oblonga*) / nr. Darwin, Northern Territory, Australia



** (stars = type localities)

Distribution: Australia (*Northern Territory*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 215,759 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 484,864 sq. km
Size (Max SCL): 31.5 cm (Joseph-Ouni et al. 2019a)
IUCN Red List: Not Evaluated (NE)
TFTSG Provisional Red List: Least Concern (LC) (2020)
Synonymy:
Macrochelodina billabong Wells and Wellington 1985:9


(*nomen nudum*) (14)*
Chelodina billabong
Type locality: "Bullo River Crossing, along the Katherine to Kununurra Rd., Northern Territory" [Australia] [orange star].
Type specimen: AMS R72933, holotype, see Cann and Sadlier (2017).
Etymology:* Vernacular (Australian Aboriginal), *billabong*, a lake associated with a temporary stream; referring to the species' habitat.

Macrochelodina kurrichalpongo Joseph-Ouni, McCord, Cann, and Smales 2019a:9
Chelodina (Macrochelodina) kurrichalpongo, Chelodina (Chelydera) kurrichalpongo, Chelydera kurrichalpongo
Type locality: "McMinns Lagoon, Outer Darwin, Litchfield, Northern Territory, Australia ... latitude -12.517, longitude 131.083)" [yellow star].
Type specimen: NTM R17209, holotype.
Etymology:* Vernacular (Australian Aboriginal), *kurrichalpongo*, Dreamtime mythology's black rock-snake that gave rise to rainbow snakes; referring to the species' long black neck and colorful hatchlings.

Chelodina (Chelydera) parkeri Rhodin and Mittermeier 1976 (21:21)
Parker's Snake-necked Turtle


Fred Parker / Balimo, Aramia River, Western Prov., Papua New Guinea / female


Fred Parker / Balimo, Aramia River, Western Prov., Papua New Guinea / female



(orange dots = trade; star = type locality)


Distribution: Indonesia (*Papua*), Papua New Guinea (*Southern*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 31,664 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 47,992 sq. km
Size (Max SCL): male 27.4 cm, female 35.3 cm (Georges et al. 2006; Ceballos et al. 2013)
IUCN Red List: Near Threatened (NT A4d) (Rhodin and Georges 2020). Previously: Vulnerable (VU) (ATTWG 2000); Vulnerable (VU) (TFTSG *in* Baillie and Groombridge 1996)

Synonymy:
Chelodina parkeri Rhodin and Mittermeier 1976:465
Macrochelodina parkeri, Chelodina (Macrochelodina) parkeri, Chelodina (Chelydera) parkeri, Chelydera parkeri
Type locality: "Mawa, Lake Murray, Western District, Papua New Guinea" [yellow star].
Type specimen: AMS 21425, holotype, see Cogger (1979) and Shea and Sadlier (1999).
Etymology:* Eponym honoring Fred Parker (1941–), India / Australia / Papua New Guinea; government patrol officer (*kiap*) and herpetologist in Papua New Guinea, who first collected the species.

Chelodina (Chelydera) rugosa Ogilby 1890 (07:91, 10:37, 14:43, 17:94, 21:22, 21:25) (13, 15)*
Northern Snake-necked Turtle


Arthur Georges / CBFTT / Morehead R., Western Prov., Papua New Guinea / female


left: Arthur Georges / CBFTT / Morehead R., Western Prov., Papua New Guinea
right: Anders G.J. Rhodin / CBFTT / nr. Daru, Western Prov., Papua New Guinea / male


(stars = type localities)

Distribution: Australia (*Queensland*), Indonesia (*Papua*), Papua New Guinea (*Southern*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 242,678 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 641,077 sq. km
Size (Max SCL): male 26.8 cm, female 36.0 cm (Georges et al. 2006; Lyons et al. 2013; Kennett et al. 2014 CBFTT)
CBFTT Account: Kennett, Fordham, Alacs, Corey, and Georges (2014), as *Chelodina oblonga*
IUCN Red List: Least Concern (LC) [Not Listed] (Behler 1996);

the currently synonymized taxon *Chelodina siebenrocki* listed by IUCN as Near Threatened (NT) (ATTWG 2000), as *Chelodina rugosa siebenrocki*

TFTSG Provisional Red List: Least Concern (LC) (2018)

Synonymy:

Chelodina rugosa Ogilby 1890:56 (14:43, 21:25)

Chelodina oblonga rugosa, Macrochelodina rugosa, Macrochelodina rugosa rugosa, Chelodina (Macrochelodina) rugosa, Chelodina (Macrochelodina) oblonga rugosa, Chelodina (Chelydera) rugosa, Chelydera rugosa

Type locality: "Cape York, Q." [Queensland, Australia]. Restricted to "Cape York Peninsula north of the Jardine River" by Cann and Sadlier (2017:79); emended to "Somerset (northern Cape York Peninsula), Queensland" by Kehlmaier et al. (2024b:542) [yellow star].

Type specimen: AMS 6256, holotype, see Goode (1967), Cogger et al. (1983), and Cann and Sadlier (2017); NHMUK 1947.3.5.89, listed as "holotype" of *Chelodina rugosa* by Uetz et al. (2019), is the holotype of *Chelodina oblonga*.

Etymology:* Latin, *rugosa*, rich in folds, wrinkled; referring to the extensive carapacial rugosity.

Comment: A proposal to the ICZN by Thomson (2006, 2007) to give *Chelodina rugosa* Ogilby 1890 precedence over *Chelodina oblonga* Gray 1841 whenever the two are considered synonyms, was not approved by ICZN (2013a).

Chelodina siebenrocki Werner 1901a:602 (07:91, 21:24, 21:25)

Chelodina oblonga siebenrocki, Macrochelodina siebenrocki, Macrochelodina rugosa siebenrocki

Type locality: "Deutsch Neu-Guinea" [northern Papua New Guinea] [in error]. Emended to "New Guinea's south coast" [southern Papua New Guinea] by Rhodin and Mittermeier (1976:474) [green star, location estimated].

Type specimen: ZMB 36221 (formerly 16491), holotype, previously recorded as lost (Peters, pers. comm. *in* Rhodin and Mittermeier 1976), rediscovered by Fritz et al. (1994); genotyped by Kehlmaier et al. (2019a).

Etymology:* Eponym honoring Friedrich Siebenrock (1853–1925), Austria; zoologist, herpetologist, and curator at the Naturhistorisches Museum, Vienna.

Chelodina intergularis Fry 1915:88

Type locality: "Australia." Restricted to "Somerset (northern Cape York Peninsula), Queensland" by Kehlmaier et al. (2024b:542) [yellow star].

Type specimen: AMS 6255, holotype, see Goode (1967) and Cogger et al. (1983); genotyped by Kehlmaier et al. (2024b).

Etymology:* Latin, *inter-*, between, and *gula*, throat; referring to the abnormal intergular scute.

Chelodina (Macrochelodina) Wells and Wellington 1985

(07:85, 08:2, 10:34, 21:21)

Southwestern Thick-necked Turtles

(includes 1 species, 1 taxon)

Macrochelodina Wells and Wellington 1985:9

Type species: *Macrochelodina oblonga* [= *Chelodina oblonga* Gray 1841], by original designation.

Etymology:* Greek, *makros*, large, and *chelodina*, turtle genus; referring to the longer, thicker neck in this genus.

Macrodiremys McCord and Joseph-Ouni 2007b:57 (08:3, 10:39)

Type species: *Macrodiremys oblonga* sensu Burbidge et al. 1974 [= *Chelodina oblonga* Gray 1841 sensu Burbidge et al. 1974 = subjective synonym of *Chelodina colliei* Gray 1856a], by original designation.

Etymology:* Greek *makros*, large, and *deire*, neck, throat, and *emys*, turtle; referring to the longer, thicker neck in this genus.

Chelodina (Macrochelodina) oblonga Gray 1841 (07:88, 08:3, 10:39, 14:43, 14:44, 21:22)

Southwestern Snake-necked Turtle



Gerald Kuchling / Moore River Nature Reserve, Western Australia, Australia



left: Gerald Kuchling / Moore River Nature Reserve, Western Australia, Australia
right: Russell A. Mittermeier / Western Australia, Australia



(stars = type localities)

Distribution: Australia (*Western Australia*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 81,641 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 218,730 sq. km

Size (Max SCL): male 23.3 cm, female 31.0 cm (Bonnet et al. 2010; Cann and Sadlier 2017; A.G.J. Rhodin, unpubl. data)

IUCN Red List: Near Threatened (NT) (TFTSG *in* Baillie and Groombridge 1996), as *Chelodina oblonga*

TFTSG Provisional Red List: Near Threatened (NT) (2009)

Synonymy:

Chelodina oblonga Gray 1841:446 (14:43)

Chelodina (Macrochelodina) oblonga, Chelodina (Macrochelodina) oblonga oblonga, Macrochelodina oblonga, Macrodiremys oblonga, Chelodina (Macrodiremys) oblonga

Type locality: "Western Australia." Restricted to "Northern Territory... possibly...Port Essington" [Australia] [orange star, in error] by Thomson (2000a:747); further restricted to "southwestern Australia" by Shea et al. (2020:427) [yellow star, location estimated].

Type specimen: NHMUK 1947.3.5.89 (formerly 1840.12.9.81), lectotype, originally considered the holotype as discussed by Thomson (2000a, 2006, 2007), see also Goode (1967) and Cann and Sadlier (2017); genotyped by Kehlmaier et al. (2019a); further discussed

and designated lectotype by Shea et al. (2020:427).
Etymology:* Latin, *oblongus*, elongated; referring to the somewhat elongated carapace.

Chelodina colliei Gray 1856a:200
Macrodiremys colliei, Chelodina (Macrodiremys) colliei
Type locality: "Swan River" [Western Australia, Australia] [yellow star, location estimated].
Type specimen: NHMUK 1947.3.5.91 (formerly 1852.11.12.2), lectotype, designated by Thomson (2000a:747), see Cann and Sadlier (2017); NHMUK 1947.3.5.91 also designated as neotype of *Chelodina oblonga* Gray 1841 by McCord and Joseph-Ouni (2007:56), but that designation was invalid (Kuchling 2010; Shea et al. 2020).
Etymology:* Eponym honoring Alexander Collie (1793–1835), United Kingdom (Scotland) / Australia; surgeon and botanist who worked and lived in Western Australia and collected the species.

Subfamily Emydurinae Gaffney 1977 (21:26)

Australasian Side-necked Turtles
(includes 5 genera, 20 species, 24 taxa)
Emydurodd Gaffney 1977:24
Emydurinei Bour and Dubois 1984b:80
Emydurinae Joyce, Anquetin, Cadena, Claude, Danilov, Evers, Ferreira, Gentry, Georgalis, Lyson, Pérez-García, Rabi, Sterli, Vitek, and Parham 2021:18

Elseya Gray 1867 (07:92, 14:45, 17:95)

Australasian Snapping Turtles
(includes 3 subgenera, 9 species, 9 taxa)
Elseya Gray 1867:44
Type species: *Elseya dentata* [= *Chelymys dentata* Gray 1863a], by subsequent designation by Lindholm (1929:291).
Etymology:* Eponym honoring Joseph R. Elsey (1834–1857), United Kingdom; surgeon and naturalist who explored northern Australia.

Pelocomastes De Vis 1897:6
Type species: *Pelocomastes ampla* † De Vis 1897 [= subjective synonym of *Chelymys uberrima* † De Vis 1897 according to Thomson 2000b], by original monotypy.
Etymology:* Greek, *pelos*, mud or clay, and *komastes*, connoisseur; referring to a presumed preference for muddy habitats.

Hanwarachelys Thomson, Amepou, Anamiato, and Georges 2015:65
Type species: *Elseya (Hanwarachelys) novaeguineae* [= *Platemys novaeguineae* Meyer 1874], by original designation.
Etymology:* Vernacular (New Guinean, Tok Pisin), *han*, hand, *wara*, water, smaller river, and Greek, *chelys*, turtle; referring to the preferred habitat of smaller streams and rivers.

Solomonemys Joseph-Ouni, McCord, and Dwyer 2022a:59 (21)*
Type species: *Elseya (Solomonemys) eidolon*, by original designation.
Etymology:* Toponym for the Solomon Islands, and Greek, *emys*, turtle.

Elseya (Elseya) Gray 1867 (17:95)


Northern Snapping Turtles
(includes 3 species, 3 taxa)
Elseya Gray 1867:44
Type species: *Elseya dentata* [= *Chelymys dentata* Gray 1863a], by subsequent designation by Lindholm (1929:291).
Etymology:* Eponym honoring Joseph R. Elsey (1834–1857), United Kingdom; surgeon and naturalist who explored northern Australia.

Elseya (Elseya) branderhorsti (Ouwens 1914) (07:93)

White-bellied Snapping Turtle, Branderhorst's Snapping Turtle


Arthur Georges / Bensbach R., Western Prov., Papua New Guinea


Anders G.J. Rhodin / nr. Merauke, Papua, Indonesia / captivity


**

(stars = type localities)


Distribution: Indonesia (*Papua*), Papua New Guinea (*Southern*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 105,359 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 155,347 sq. km
Size (Max SCL): male 40.2 cm, female 48.1 cm (Georges et al. 2006; Ceballos et al. 2013)
IUCN Red List: Vulnerable (VU B1+2e [ver. 2.3]) (ATTWG 2000)
TFTSG Provisional Red List: Vulnerable (VU) (2019)
Synonymy:
Emydura branderhorsti Ouwens 1914:31
Elseya branderhorsti, Elseya (Elseya) branderhorsti
Type locality: "Sth. Nw. Guinea" [Southern New Guinea = Papua, Indonesia]. Restricted to "southeastern Papua, Indonesia, between the Lorentz River and Merauke" [orange star, location estimated] by Thomson et al. (2015:65); and to "Bensbach River of the Trans-Fly region of Papua New Guinea (8°50'58.6896" S., 141°14'52.944" E.)" [yellow star] by Thomson et al. (2015:63) by neotype designation.
Type specimen: MZB, holotype, originally live, now lost (Thomson et al. 2015); PNGM 25201, neotype, designated by Thomson et al.

(2015:63).
Etymology:* Eponym honoring Bastiaan Branderhorst (1880–1971), Netherlands / Indonesia; physician and botanist who worked in Dutch New Guinea (now Papua, Indonesia) and Java, Netherlands Indies (now Indonesia), and collected the type.

Elseya (Elseya) dentata (Gray 1863a) [16]*
Northern Snapping Turtle; North Australian Snapping Turtle


Gerald Kuchling / Carson R., Western Australia, Australia


Gerald Kuchling / Western Australia, Australia



**


(stars = type localities)

Distribution: Australia (*Northern Territory, Western Australia*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 154,942 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 411,102 sq. km
Size (Max SCL):* male 28.4 cm, female 33.0 cm (Cann and Sadlier 2017)
IUCN Red List: Least Concern (LC) [Not Listed] (Behler 1996)
TFTSG Provisional Red List: Least Concern (LC) (2011)
Synonymy:
Chelymys dentata Gray 1863a:98
Podocnemis dentata, Elseya dentata, Emydura dentata, Elseya dentata dentata, Elseya (Elseya) dentata
Type locality: "N. Australia; Upper Victoria, in Beagle's Valley." [Northern Territory, Australia]. Emended to "Beagles Valley, Victoria River, Northern Territory...(15°34' S, 130°54' E)" [yellow star] by Thomson et al. (2015:62).
Type specimen: NHMUK 1947.3.6.3 (formerly 1857.10.24.73), lectotype, designated by Wells and Wellington (1985:8), see Goode (1967) and Cann and Sadlier (2017); NHMUK 1947.3.6.2 erroneously designated lectotype by Thomson et al. (2015:62).
Etymology:* Latin, *dens*, tooth, *dentata*, toothed, serrated; referring to the serrated rear marginal scutes in juveniles.
Chelymys elseyi Gray 1864d:132 (*nomen nudum*)
Etymology:* Eponym honoring Joseph R. Elsey (1834–1857), United Kingdom; surgeon and naturalist who explored northern Australia.
Chelymys elseya Gray 1870c:76 (*nomen nudum*)
Etymology:* Eponym honoring Joseph R. Elsey (1834–1857), United Kingdom; surgeon and naturalist who explored Australia.
Elseya intermedia Gray 1872c:23
Type locality: "North Australia, upper part of Victoria" [Victoria River, Northern Territory, Australia] [yellow star, location estimated].
Type specimen: NHMUK 1947.3.4.14 (formerly 1857.10.24.72), holotype, see Cogger et al. (1983) and Cann and Sadlier (2017; genotyped by Kehlmaier et al. (2019a).
Etymology:* Latin, *inter-*, between, and *medius*, middle; referring to this taxon being similar to and intermediate between *Elseya dentata* and *Elseya spinosa* [= *Myuchelys latisternum*].
Elseya (Elseya) kalumburu Joseph-Ouni, McCord, and Cann 2022c:38 [16]*
Elseya kalumburu
Type locality:* "Kalumburu, King Edward River, Western Australia ... 126°65'E; 14°3'S." GPS coordinates incorrect, emended here to 126°39'E; 14°16'S [green star].
Type specimen:* QM J77957, holotype.
Etymology:* Toponym for Kalumburu, Western Australia.
Elseya (Elseya) sahul Joseph-Ouni, Cann, and McCord 2022d:59 [16]*
Elseya sahul
Type locality:* "Turtle Lagoon, Wongalara Sanctuary, Roper River region, Northern Territory, Australia ... 126°65'E; 14°3'S." GPS coordinates incorrect, emended here to 134°28'E; 14°6'S [blue star].
Type specimen:* NTM R36773, holotype.
Etymology:* Toponym for the Sahul Shelf, the marine continental shelf connecting northern Australia to New Guinea and the Aru Islands

Elseya (Elseya) flaviventralis Thomson and Georges 2016 (17:96)
Yellow-bellied Snapping Turtle


Arthur Georges / South Alligator R., Northern Territory, Australia


Arthur Georges / South Alligator R., Northern Territory, Australia


(stars = type localities)

Distribution: Australia (*Northern Territory*)


Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 107,600 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 161,440 sq. km
Size (Max SCL): male 26.5 cm, female 34.0 cm (Thomson and Georges 2016; Joseph-Ouni et al. 2020)
IUCN Red List: Not Evaluated (NE)
TFTSG Provisional Red List: Least Concern (LC) (2018)
Synonymy:
Elseya flaviventralis Georges, Doody, Young, and Cann 2000:7 (*nomen nudum*)
Etymology:* Latin, *flavus*, yellow, *venter*, belly, abdomen; referring to the uniformly yellowish or cream-colored plastron.
Elseya jukesi Wells 2002b:7 (*nomen nudum*)
Etymology:* Eponym honoring Brian Jukes (?–), Australia; amateur herpetologist.
Elseya jukesi Wells 2007b:5 (unavailable name) (07:94, 10:43)
Type locality: "Pul Pul Billabong, South Alligator River (13 34'S, 132 35'E)" [Northern Territory, Australia] [green star].
Type specimen: NTM 13985, holotype.
Etymology:* Eponym honoring Brian Jukes (?–), Australia; amateur herpetologist who collected the species.
Elseya (Elseya) flaviventralis Thomson and Georges 2016:20
Type locality: "Pine Creek Crossing, South Alligator River Drainage, Kakadu National Park, Northern Territory, Australia, 13°30' S 132°28' E" [yellow star].
Type specimen: NTM 13512, holotype.
Etymology:* Latin, *flavus*, yellow, *venter*, belly, abdomen; referring to the uniformly yellowish or cream-colored plastron.

Elseya (Hanwarachelys) Thomson, Amepou, Anamiato, and Georges 2015 (17:95)

New Guinea Stream Turtles
(includes 3 species, 3 taxa)
Hanwarachelys Thomson, Amepou, Anamiato, and Georges 2015:65
Type species: *Elseya (Hanwarachelys) novaeguineae* [= *Platemys novaeguineae* Meyer 1874], by original designation.
Etymology:* Vernacular (New Guinean, Tok Pisin), *han*, hand, *wara*, water, smaller river, and Greek, *chelys*, turtle; referring to the preferred habitat of smaller streams and rivers.
Solomonemys Joseph-Ouni, McCord, and Dwyer 2022a:59 (21)*
Type species:* *Elseya (Solomonemys) eidolon*, by original designation.
Etymology:* Toponym for the Solomon Islands.

Elseya (Hanwarachelys) novaeguineae (Meyer 1874) (10:40, 14:45, 17:97, 21:27) (17, 18)*


Western New Guinea Stream Turtle, New Guinea Snapping Turtle



MTD 8222, holotype / Passim, Barbussi R., West Papua, Indonesia



left: Anders G.J. Rhodin / Bintuni Bay region, West Papua, Indonesia / captivity / "*Elseya caelata*"
right: William P. McCord / Waren R., West Papua, Indonesia



(stars = type localities)

Distribution: Indonesia (*West Papua*)
Introduced: Palau (*Babeldaob*), see Aoki (1977)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 66,950 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 153,447 sq. km
Size (Max SCL): male 19.1 cm, female 22.9 cm (A.G.J. Rhodin, unpubl. data)
IUCN Red List: Least Concern (LC) (ATTWG 2000). Previously: Least Concern (LC) [Not Listed] (Behler 1996)
TFTSG Provisional Red List: Least Concern (LC) (2015)
Synonymy:
Platemys novaeguineae Meyer 1874:128
Platemys novae guineae, Emydura novaeguineae, Elseya novaeguineae, Elseya latisternum novaeguineae, Elseya dentata novaeguineae, Elseya novaeguineae novaeguineae, Myuchelys novaeguineae, Elseya (Hanwarachelys) novaeguineae, Hanwarachelys novaeguineae
Type locality: "Neu-Guinea" [West Papua, Indonesia]. Restricted to "Passim (Nordwest Neu Guinea)" [West Papua, Indonesia] [yellow star] by Meyer (1887:4); emended to "southwestern shore of Cenderawasih Bay on the southeastern Vogelkop" [West Papua, Indonesia] [yellow star] by Rhodin and Genorupa (2000:132); and further emended to "Passim, Barbussi River, ca. 3 km N. Sieb, 1 km S. Tandjung Sjeri (= Syeri), west shore Geelvink Bay (= Cenderawasih Bay), Papua, Indonesia (1°41' S, 134°05' E)" [yellow star] by

Thomson et al. (2015:67).
Type specimen: MTD 8222, holotype, see Obst (1976) and Thomson et al. (2015); MSNG CE8605, erroneously listed as a syntype by Goode (1967) is not a type; MTD 8222 genotyped by Kehlmaier et al. (2019a).
Etymology:* Toponym for New Guinea.

Elseya caelatus Joseph-Ouni and McCord 2019a:25 (21:27) (17, 18)*
Elseya caelatus caelatus, Elseya caelata, Elseya caelata caelata, Elseya (Hanwarachelys) caelata, Elseya (Hanwarachelys) caelata caelata
Type locality: "Salawati Island, West Papua Province, (New Guinean) Indonesia" [green star, location estimated].
Type specimen: AMNH R178715, holotype.
Etymology:* Latin, *caelatus*, carved; referring to the sculptured carapace.


Elseya caelatus ayamaru Joseph-Ouni and McCord 2019a:28 (21:27)
Elseya caelata ayamaru, Elseya (Hanwarachelys) caelata ayamaru
Type locality: "Ayamaru lakes, West Papua Province, (New Guinean) Indonesia" [blue star, location estimated].
Type specimen: AMNH R178717, holotype.
Etymology:* Toponym for the Ayamaru lakes, Vogelkop Peninsula, Papua, Indonesia.

Elseya (Hanwarachelys) caelatus berau Joseph-Ouni and McCord 2023a:9 (17)*
Elseya caelatus berau
Type locality:* "vicinity of Samberai, upper Wasian River, southeastern Vogelkop Peninsula, West Papua Province, Indonesia (New Guinea)" [pink star, location estimated].
Type specimen:* YPM HERR.022597, photograph in Joseph-Ouni and McCord 2023a:10(Plate1), holotype; see also Joseph-Ouni and McCord (2025).
Etymology:* Toponym for the Berau Gulf, Vogelkop Peninsula, Papua, Indonesia.

Elseya (Hanwarachelys) rhodini Thomson, Amepou, Anamiato, and Georges 2015 (17:97) (19)*
Southern New Guinea Stream Turtle; Rhodin's Stream Turtle


John Cann / Kapara R., nr. Uta, Mimika Regency, Papua, Indonesia


left: John Cann / Josaker, Eilanden R., Papua, Indonesia
right: Fred Parker / Nomad R., Western Prov., Papua New Guinea


**
(orange dots = probably introduced or trade; stars = type localities)

Distribution: Indonesia (*Aru Islands*, *Papua*); Papua New Guinea (*Southern*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 278,847 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 551,990 sq. km
Size (Max SCL): male 20.5 cm, female 27.6 cm (Thomson et al. 2015)
IUCN Red List: Least Concern (LC) (Amepou, Georges, Thomson, Eisemberg, and Rhodin 2020)
Synonymy:

Elseya (Hanwarachelys) rhodini Thomson, Amepou, Anamiato, and Georges 2015:69
Elseya rhodini, Hanwarachelys rhodini
Type locality: "Rue Creek (tributary of Wau Creek), Gulf Province, Papua New Guinea (07°11'67.3" S, 144°37'13.8" E)" [yellow star]; GPS coordinates emended to (07°11.673' S, 144°37.138' E) [yellow star] by TTWG (2017:195).
Type specimen: PNGM 25204, holotype.
Etymology:* Eponym honoring Anders G.J. Rhodin (1949–), Sweden / USA; orthopedic surgeon and herpetologist, Chair Emeritus of the IUCN Tortoise and Freshwater Turtle Specialist Group and founder of Chelonian Research Foundation, who studied in Papua New Guinea and collected and first recognized the species as distinct.

Elseya (Hanwarachelys) papua Joseph-Ouni and McCord 2022c:42 (19)*
Elseya papua, Hanwarachelys papua
Type locality:* "vicinity of Muting, northern Merauke region, Papua Province, Indonesia (New Guinea)" [green star].
Type specimen:* YPM HERR.022306, holotype.
Etymology:* Toponym for Papua Province, Indonesia.

Elseya (Hanwarachelys) timika timika Joseph-Ouni and McCord 2023b:95 (19)*

Elseya timika, Elseya timika timika, Hanwarachelys timika, Hanwarachelys timika timika

Type locality:* "lower reaches of the Timika (Mimika) River, Papua Province, Indonesia (New Guinea)." [blue star, location estimated].
Type specimen:* YPM HERR 022326, holotype.
Etymology:* Toponym for the Timika River, Papua, Indonesia.

Elseya (Hanwarachelys) timika aru Joseph-Ouni and McCord 2023b:100 [19]*

Elseya timika aru, Hanwarachelys timika aru

Type locality:* "middle reaches of the Tunguwatu River, Wokam Island, Aru Islands, Papua Province, Indonesia (New Guinea)" [pink star].
Type specimen:* YPM HERR 022328, holotype.
Etymology:* Toponym for the Aru Islands, Indonesia.

Elseya (Hanwarachelys) timika wamena Joseph-Ouni and McCord 2023b:105 [19]*

Elseya timika wamena, Hanwarachelys timika wamena

Type locality:* "North Baliem River, Baliem Valley, Papua Province, Indonesia (New Guinea)" [brown star].
Type specimen:* YPM HERR 022324, holotype.
Etymology:* Toponym for the Wamena region, Papua, Indonesia.

Elseya (Hanwarachelys) timika paniai Joseph-Ouni and McCord 2023b:109 [19]*

Elseya timika paniai, Hanwarachelys timika paniai

Type locality:* "Ned. Nieuw Guinea, Tigi meer = Lake Tigi, Paniai Lakes, Indonesia (New Guinea)" [= Danau Tigi] [purple star].
Type specimen:* RMNH RENA.27896, holotype.
Etymology:* Toponym for the Paniai Lakes, Papua, Indonesia.

Elseya (Hanwarachelys) freweri Joseph-Ouni and McCord 2023b:111 [19]*

Elseya freweri, Hanwarachelys freweri

Type locality:* "vicinity of Waurake, upper Kamoera River, Papua Province, Indonesia (New Guinea)" [gray star, location estimated].
Type specimen:* YPM HERR 022323, holotype.
Etymology:* Eponym honoring Jason Frewer (1900s–), United Kingdom / USA; commercial turtle breeder and dealer in Florida.

Elseya (Hanwarachelys) schultzei (Vogt 1911) (07:95, 10:40, 14:45, 21:28) [18, 20, 21]*


Northern New Guinea Stream Turtle, Schultze's Snapping Turtle



William P. McCord / Cyclops Mts., Papua, Indonesia / "*Elseya orestiad*"



Anders G.J. Rhodin / nr. Aiome, Ramu R., Madang Prov., Papua New Guinea



• (orange dots and star and brown star = probable trade; stars = type localities)

Distribution: Indonesia (*Papua*); Papua New Guinea (*Northern*)
Introduced:* Solomon Islands (*Guadalcanal, Malaita*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 154,896 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 495,713 sq. km
Size (Max SCL): female 22.5 cm (A.G.J. Rhodin, unpubl. data)
IUCN Red List: Not Evaluated (NE)
TFTSG Provisional Red List: Least Concern (LC) (2015)
Synonymy:

Emydura schultzei Vogt 1911:410

Elseya schultzei, Elseya novaeguineae schultzei, Elseya (Hanwarachelys) schultzei, Hanwarachelys schultzei

Type locality: "Fluss westlich der Tamimündung...Holländisch-Neu-Guinea" [Papua, Indonesia] [yellow star]. Restricted to "near Sae village, Seko coast, near Skosai, ca 5 km W. mouth of Tami River, Papua, Indonesia (2°37' S, 140°54' E)" [yellow star] by Thomson et al. (2015:68).
Type specimen: ZMB 22182, holotype, see Goode (1967), Fritz et al. (1994), and Thomson et al. (2015).
Etymology:* Eponym honoring Leonhard Schultze-Jena (1872–1955), Germany; explorer, zoologist, and anthropologist who worked in New Guinea and collected the types.

Elseya orestiad Joseph-Ouni and McCord 2019b:45 (21:28) [18]*

Type locality: "Cyclops Mountains, Papua Province, (New Guinean) Indonesia" [green star, location estimated].
Type specimen: AMNH R178719, holotype.
Etymology:* Mythonym for the *Orestiades*, Greek mythological nymphs who came out to hunt and fish; referring to the predatory nature of these turtles.

Elseya (Hanwarachelys) nabire Joseph-Ouni and McCord 2022b:29 [20]*

Elseya nabire, Hanwarachelys nabire
Type locality:* "Wanggar River, Papua Province, Indonesia (New Guinea)" [blue star].
Type specimen:* YPM HERR.022308, holotype.
Etymology:* Toponym for Nabire, Papua, Indonesia.

Elseya (Solomonemys) eidolon Joseph-Ouni, McCord, and Dwyer 2022a:61 (21)*
Elseya eidolon, Solomonemys eidolon
Type locality:* "wetland southeast of Malu, northern Malaita Island, Malaita District, Solomon Islands" [orange star] [likely trade specimen].
Type specimen:* YPM HERR.022305, holotype.
Etymology:* Greek, *eidolon*, ghost or image; referring to the uncertainty regarding this turtle's existence and to a native belief that freshwater turtles represent the incarnation of deceased, spiritual wanderers.
Comment: The single known specimen (holotype) most likely represents a translocated trade specimen originally from New Guinea.

Elseya (Solomonemys) auramemoria Joseph-Ouni, McCord, and Dwyer 2022a:64 (21)*
Elseya auramemoria, Solomonemys auramemoria
Type locality:* "middle-lower reaches of the eastern bank of the Mbonehe River, northern Guadalcanal, Guadalcanal District, Solomon Islands" [brown star] [likely trade specimen].
Type specimen:* YPM HERR.022304, holotype.
Etymology:* Greek, *aura*, air or atmosphere, and Latin, *memoria*, remembrance; referring to a native belief that freshwater turtles represent the incarnation of deceased, spiritual wanderers, and in memorial to the thousands of human lives lost during World War II battles in the Solomons.
Comment: The single known specimen (holotype) most likely represents a translocated trade specimen originally from New Guinea.

Elseya (Pelocomastes) De Vis 1897 (17:95)


Eastern Snapping Turtles
(includes 3 species, 3 taxa)
Pelocomastes De Vis 1897:6
Type species: *Pelocomastes ampla* † De Vis 1897 [= subjective synonym of *Chelymys uberrima* † De Vis 1897 according to Thomson 2000b], by original monotypy.
Etymology:* Greek, *pelos*, mud or clay, and *komastes*, connoisseur; referring to a presumed preference for muddy habitats.

Elseya (Pelocomastes) albagula Thomson, Georges, and Limpus 2006 (21:29)


White-throated Snapping Turtle, Southern Snapping Turtle



John Cann / CCB / Burnett R., Queensland, Australia



left: Marilyn Connell / Mary R., Queensland, Australia / female
right: Arthur Georges / Burnett R., Queensland, Australia



**

(star = type locality)

Distribution: Australia (*Queensland*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 64,166 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 183,961 sq. km
Size (Max SCL): male 28.3 cm, female 41.8 cm (Thomson et al. 2006; Hamann et al. 2008)
IUCN Red List: Not Evaluated (NE)
TFTSG Provisional Red List: Endangered (EN) (2018)
Synonymy:
Elseya albagula Thomson, Georges, and Limpus 2006:75
Elseya dentata albagula, Elseya (Pelocomastes) albagula, Pelocomastes albagula, Elseya albagula albagula
Type locality: "Ned Churchwood Weir, Burnett River, Queensland, Australia (25º03' S, 152º05' E)" [yellow star].
Type specimen: QM J81785, holotype.


Etymology:* Latin, *alba*, white; and *gula*, throat; referring to the white throat commonly seen in adult females.

Elseya (Pelocomastes) irwini Cann 1997c (21:29) (22)*
Irwin's Snapping Turtle, Burdekin River Snapping Turtle, Johnstone River Snapping Turtle


Anders Zimny / Broken R., Queensland, Australia



left: Duncan Limpus / Burdekin R., Queensland, Australia
right: Anders Zimny / Broken R., Queensland, Australia



**

(orange dot = probably introduced; stars = type localities)


Distribution: Australia (*Queensland*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 14,425 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 54,374 sq. km
Size (Max SCL): male 23.9 cm, female 35.7 cm (Itescu et al. 2014; Cann and Sadlier 2017; Freeman et al. 2018)
IUCN Red List: Not Evaluated (NE)
TFTSG Provisional Red List: Near Threatened (NT) (2018)
Synonymy:
Elseya stirlingi Wells and Wellington 1985:9 (*nomen nudum*)
Type locality:* "Cairns district, Queensland" [Australia]; restricted to "Barron River, near Kuranda, north-east Queensland" [orange star] by Wells (2007b:2); emended to "Cairns district, Qld (16 55'S, 145 46'E)" by Wells (2007b:4) [geographic coordinates in error, brown star]; emended here to "16 52'S, 145 41'E" [orange star].
Type specimen: AMS R93048 (formerly R68848), discussed by Iverson et al. (2001b), see Shea and Sadlier (1999), Cann and Sadlier (2017), and Joseph-Ouni et al. (2023a), who identify it as a misidentifed specimen of *Elseya latisternum*.
Etymology:* Eponym honoring Stanley Stirling (1918–2012), Australia; Australia; amateur naturalist and private zoological park owner, who collected the type.
Elseya irwini Cann 1997c:36
Elseya dentata irwini, Elseya (Pelocomastes) irwini, Pelocomastes irwini, Elseya irwini irwini, Elseya (Pelocomastes) irwini irwini
Type locality: "Burdekin River, Queensland, 19°42' S, 147°18' E, approximately 18 km upstream from Ayr" [Australia] [yellow star].
Type specimen: QM J59431, holotype, see Cann and Sadlier (2017).
Etymology:* Eponym honoring Steve Irwin (1962–2006), Australia; herpetologist, wildlife promoter, and television personality.
Elseya sterlingi Iverson, Thomson, and Georges 2001b:363 (*nomen invalidum*)
Comment: Typographic error for *stirlingi*.
Elseya stirlingi Wells 2007b:4 (unavailable name) (07:96, 10:43)
Type locality:* "South Johnstone River, Qld (17 38'S, 145 05'E)" [Queensland, Australia] [geographic coordinates with typographic error, green star]; emended here to "17 38'S, 145 50'E" [blue star].
Type specimen: QM J48059, holotype.
Etymology:* Eponym honoring Stanley Stirling (1918–2012), Australia; amateur naturalist and private zoological park owner.
Elseya (Pelocomastes) irwini stirlingi Wells *in* Joseph-Ouni, Cann, Turner, Freeman, and McCord 2023a:43 (22)*
Type locality:* "South Johnstone River, Queensland, Australia (17 38'S, 145 05'E)" [geographic coordinates with typographic error, green star]; emended here to "17 38'S, 145 50'E" [blue star].
Type specimen:* QM J48059, holotype.
Etymology:* Eponym honoring Stanley Stirling (1918–2012), Australia; amateur naturalist and private zoological park owner.

Elseya (Pelocomastes) lavarackorum (White and Archer 1994) (21:30) (23)*
Gulf Snapping Turtle, Riversleigh Snapping Turtle


John Cann / CBFTT / Riversleigh Station, Gregory R., Queensland, Australia



Alastair Freeman / CBFTT / Boodjamulla National Park, Queensland, Australia / male, female, juvenile


**

(stars = type localities)

Distribution: Australia (*Northern Territory, Queensland*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 48,232 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 166,038 sq. km

Size (Max SCL):* male 24.2 cm, female 35.2 cm (Freeman et al. 2014 CBFTT; Joseph-Ouni et al. 2020); Max SCL of fossil specimen: ca. 41 cm (White and Archer 1994)

CBFTT Account: Freeman, Thomson, and Cann (2014)

IUCN Red List: Not Evaluated (NE)

TFTSG Provisional Red List: Data Deficient (DD) (2018)

Synonymy:

Emydura lavarackorum † White and Archer 1994:159

Elseya lavarackorum, Elseya dentata lavarackorum, Elseya (Pelocomastes) lavarackorum, Pelocomastes lavarackorum, Elseya (Elseya) lavarackorum

Type locality: "Gregory River, Riversleigh Station, northwestern Queensland, approximately 200 km northwest of Mount Isa" [Australia] [yellow star].

Type specimen:* QM F24121, holotype, fossil, almost complete plastron, partial carapace and pelvic fragments, figured (f.1-4), see Thomson et al. (1997); specimen currently registered as UNSW AR14547, see Thomson et al. (2023).

Geological age: Late Pleistocene, Terrace Site, gravel sediments, ^{14}C age ca. 28,000–21,200 ybp, see Davis and Archer (1997).

Etymology:* Eponym honoring Sue and Jim Lavarack (1926– and 1922–2014), Australia; amateur paleontologists who worked on and helped direct the Riversleigh Expedition and collected the type.

Comment: Although originally described from a fossil specimen, Thomson et al. (1997, 2023) considered the extant taxon from the same drainage basin as conspecific with *E. lavarackorum*. However, Joseph-Ouni et al. (2020, 2022f) considered *E. oneiros* distinct from *E. lavarackorum*.

Elseya (Pelocomastes) oneiros Joseph-Ouni, McCord, Cann, Smales, Freeman, Sadlier, Couper, White, and Amey 2020:32 (21:30) (23)*

Elseya oneiros

Type locality: "Elizabeth Gorge, Bowthorn Station, Queensland, Australia [...] latitude -18.2167, longitude 138.3333" [green star].

Type specimen: QM J47908, holotype.

Etymology:* Mythonym for *Oneiros*, the Greek mythological dream messenger; referring to the ethnic Australian Aboriginal "dream time" in allusion.

Elseya (Pelocomastes) gondwana Joseph-Ouni, Cann, and McCord 2022d:72 (16, 23)*

Elseya gondwana

Type locality:* "Red Lily Lagoon, Roper River, Northern Territory, Australia ... 134°083'E; 14°7'S." GPS coordinates incorrect, emended here to 133°25'E; 14°53'S [blue star].

Type specimen:* NTM R16328, holotype.

Etymology:* Toponym for Gondwana, the ancient southern supercontinent upon which chelid turtles evolved.

Elusor Cann and Legler 1994

Mary River Turtles

(includes 1 species, 1 taxon)

Elusor Cann and Legler 1994:83

Type species: *Elusor macrurus* Cann and Legler 1994, by original monotypy.

Etymology:* Latin, *eludo*, remain unnoticed, avoid, elude; referring to how long the native habitat of this species escaped discovery after the species was discovered in the pet trade.

Elusor macrurus Cann and Legler 1994


Mary River Turtle



Marilyn Connell / Mary R., Queensland, Australia / male



Marilyn Connell / Mary R., Queensland, Australia / male



** (star = type locality)

Distribution: Australia (*Queensland*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 6,116 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 8,165 sq. km

Size (Max SCL): male 43.6 cm, female 34.8 cm (Cann and Legler 1994; Limpus 2008; Cann and Sadlier 2017)

IUCN Red List: Endangered (EN B1+2c [ver. 2.3]) (TFTSG 1996)

TFTSG Provisional Red List: Critically Endangered (CR) (2018)

Synonymy:

Elusor macrurus Cann and Legler 1994:83

Type locality: "Mary River. 45.5 km S and 21.0 km W Maryborough, Queensland, Australia, elevation approximately 30 m (25°58' S, 152°30' E)" [yellow star].

Type specimen: QM J51275, holotype, see Shea and Sadlier (1999) and Cann and Sadlier (2017).

Etymology:* Greek, *makros*, big, long, and *oura*, tail, rump; referring to the long and large tail.

Emydura Bonaparte 1836

Australasian Short-necked Turtles

(includes 5 species, 9 taxa)

Emydura Bonaparte 1836:7

Type species: *Emydura macquaria* [= *Emys macquaria* Cuvier 1829] [= *Chelys (Hydraspis) macquarii* Gray 1830e], by original monotypy.

Etymology:* Greek, *emys*, turtle, and *oura*, tail, rump; referring to the long and large tail in males.

Chelymys Gray 1844:42

Type species: *Chelymys macquaria* [= *Emys macquaria* Cuvier 1829] [= *Chelys (Hydraspis) macquarii* Gray 1830e], by original monotypy.

Etymology:* Greek, *chelys*, turtle, and *emys*, turtle.

Euchelymys Gray 1871a:118

Type species: *Euchelymys sulcifera* [= subjective synonym of *Chelys (Hydraspis) macquarii* Gray 1830e], by subsequent designation by Lindholm (1929:290).

Etymology:* Greek, *eu-*, true, and *chelys*, turtle, and *emys*, turtle.

Euchelemys Günther 1873b:80 (*nomen novum*)

Etymology:* Greek, *eu-*, true, and *chelys*, turtle, and *emys*, turtle.

Comment:* Unjustified emendation of *Euchelymys*.

Tropicochelymys Wells and Wellington 1985:9

Type species: *Tropicochelymys victoriae* [= *Hydraspis victoriae* Gray 1842], by original designation.

Etymology:* Latin, *tropica*, tropical, and Greek, *chelys*, turtle, and *emys*, turtle.

Emydura australis (Gray 1841) (21:17) (24)*


Northern Red-faced Turtle



Gerald Kuchling / Barnett R., Western Australia, Australia / male



Gerald Kuchling / Western Australia, Australia / male



** (stars = type localities)

Distribution: Australia (*Northern Territory, Western Australia*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 103,930 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 427,464 sq. km

Size (Max SCL): 30.0 cm (Ernst et al. 2006b; Itescu et al. 2014)

IUCN Red List: Least Concern (LC) [Not Listed] (Behler 1996)

TFTSG Provisional Red List: Least Concern (LC) (2009)

Synonymy:

Hydraspis australis Gray 1841:445 (10:41, 21:17) (24)*

Hydraspis australia, Chelymys australis, Emydura australis, Emydura australis australis

Type locality: "Western Australia?" Restricted to "Australia, Macquarie River" [in error; no star] by Gray (1872d:506). Suggested to be in the "vicinity of Prince Regent River" [Western Australia, Australia] by Cann and Sadlier (2017:343); restricted in our annotation to "vicinity of the Glenelg and Prince Regent rivers, Western Australia, Australia" [yellow star, location estimated].

Type specimen: NHMUK 1947.3.4.36 (formerly 1840.12.9.52), holotype, see Goode (1967), Cogger et al. (1983), and Cann and Sadlier (2017); genotyped by Kehlmaier et al. (2019a); see also Kehlmaier et al. (2024a).

Etymology:* Toponym for Australia.

Hydraspis victoriae Gray 1842:55 (24)*

Chelymys victoriae, Emydura victoriae, Tropicochelymys victoriae

Type locality: "Victoria River, North-west coast of New Holland" [Northern Territory, Australia] [green star, location estimated].


Type specimen: NHMUK 1947.3.5.96 (formerly 1841.10.12.2), holotype, see Cann and Sadlier (2017) and Kehlmaier et al. (2019a); NHMUK 1947.3.5.95 (formerly 1846.5.23.3), erroneously designated "lectotype" by Wells and Wellington (1985:9), but this specimen was received after the description of the species (Uetz et al. 2019), and has been identified as an *Emydura macquarii* (Iverson et al. 2001; Kehlmaier et al. 2019a); both specimens genotyped by Kehlmaier et al. (2019a); see also Kehlmaier et al (2024).

Etymology:* Eponym honoring Queen Victoria (1819–1901), United Kingdom; British monarch for whom the Victoria River in Australia was also named.

Emydura gunaleni Smales, McCord, Cann, and Joseph-Ouni 2019a (21:31)
Gunalen's Short-necked Turtle



William P. McCord and Danny Gunalen / Kais R., West Papua, Indonesia



(orange dot = possible trade; star = type locality)

Distribution: Indonesia (*West Papua*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 3,789 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 5,189 sq. km

Size (Max SCL): male 14.7 cm, female 18.5 cm (Smales et al. 2019a)


IUCN Red List: Not Evaluated (NE)

TFTSG Provisional Red List:* Data Deficient (DD) (2022)

Synonymy:

Emydura gunaleni Smales, McCord, Cann, and Joseph-Ouni 2019a:24
Type locality: "Kais River (Sungai Kais) at approximately 1°43.808'S 132°7.826'E, in Kais District of Sarong Selatan Regency, West Papua Province (Papua Barat Province), Indonesia" [yellow star].
Type specimen: AMNH R178722, holotype.
Etymology:* Eponym honoring Danny Gunalen (1975–), Indonesia; reptile exporter and herpetologist who obtained the type.

Emydura macquarii (Gray 1830e) (10:7, 17:98)
Eastern Short-necked Turtle, Southern River Turtle
(includes 4 subspecies)



(subspecies: *macquarii* = red, *emmotti* = blue, *krefftii* = green, *nigra* = pink; overlap = intergrades)

Distribution: Australia (*New South Wales, Queensland, South Australia, Victoria*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 772,938 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 2,805,474 sq. km

Size (Max SCL): male 30.0 cm, female 36.8 cm (see subspp.)

IUCN Red List: Least Concern (LC) [Not Listed] (Behler 1996)

TFTSG Provisional Red List: Least Concern (LC) (2011)


Emydura macquarii macquarii (Gray 1830e) (07:98, 10:7, 10:41, 10:42)
Macquarie River Turtle, Murray Short-necked River Turtle



Arthur Georges / Brisbane R., Queensland, Australia



Arthur Georges / Lower Murray R., South Australia, Australia



** (subspecies: *macquarii* = red, *emmotti* = blue, *krefftii* = green, *nigra* = pink; overlap = intergrades; stars = type localities)

Distribution: Australia (*New South Wales, Queensland, South Australia, Victoria*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 502,406 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 1,370,502 sq. km

Size (Max SCL): male 30.0 cm, female 34.0 cm (McCord et al. 2003; Cann and Sadlier 2017)

IUCN Red List: The currently synonymized taxon *Emydura signata* is listed as Least Concern (LC) (TFTSG 1996), as *Emydura macquarii signata*

Synonymy:

Emys macquaria Cuvier 1829:11 (*nomen nudum*)
Etymology:* Eponym honoring Lachlan Macquarie (1762–1824), United Kingdom (Scotland) / Australia; governor of New South Wales, Australia, in whose honor the Macquarie River was named.

Chelys (Hydraspis) macquarii Gray 1830e:15 (10:7)
Chelys macquarii, Emys macquarii, Emydura macquarii, Chelymys macquarii, Emydura macquarii macquarii
Type locality: "New Holland" [Australia]. Restricted to "Novâ Hollandiâ, Macquarrie River" [New South Wales, Australia] by Gray (1831d:40), and to "Upper Macquarie River, in the vicinity of Bathurst [NSW]" [Australia] [yellow star] by Cann (1998:101).
Type specimen: MNHN 9409, holotype, see Cogger et al. (1983) and Cann and Sadlier (2017).
Etymology:* Eponym honoring Lachlan Macquarie (1762–1824), United Kingdom (Scotland) / Australia; governor of New South Wales, Australia, in whose honor the Macquarie River was named.

Hydraspis macquarrii Gray 1831d:40 (*nomen novum*)
Emydura macquarrii, Emydura macquarrii macquarrii
Etymology:* Eponym honoring Lachlan Macquarie (1762–1824), United Kingdom (Scotland) / Australia; governor of New South Wales, Australia, in whose honor the Macquarie River was named.
Comment:* Unjustified emendation or error for *macquarii*.

Platemys macquaria Duméril and Bibron 1835:438 (*nomen novum*)
Hydraspis macquaria, Chelymys macquaria
Etymology:* Eponym honoring Lachlan Macquarie (1762–1824), United Kingdom (Scotland) / Australia; governor of New South Wales, Australia, in whose honor the Macquarie River was named.
Comment:* Unjustified emendation or error for *macquarii*.

Euchelymys sulcifera Gray 1871a:118
Type locality: "North Australia."
Type specimen: NHMUK 1947.3.5.97 (formerly 1858.12.8.6), lectotype, designated by Cogger et al. (1983:63).
Etymology:* Latin, *sulcus*, furrow, and *ferre*, to bear or carry; referring to the numerous longitudinal furrows on the carapace.

Emydura macquariae Boulenger 1889:ix (*nomen novum*)
Etymology:* Eponym honoring Lachlan Macquarie (1762–1824), United Kingdom (Scotland) / Australia; governor of New South Wales, Australia, in whose honor the Macquarie River was named.
Comment:* Unjustified emendation or error for *macquarii*.

Emydura signata Ahl 1932:127 (10:42)
Emydura macquarrii signata, Chelymys signata, Emydura macquarii signata
Type locality: "Umgebung von Brisbane, Australien" [Queensland, Australia] [green star].
Type specimen: ZMB 34102, holotype, see Goode (1967), Cogger et al. (1983), Fritz et al. (1994), and Cann and Sadlier (2017).
Etymology:* Latin, *signum*, the sign or feature; referring to the features that appeared to differentiate the examined specimens from other species.

Emydura canni Worrell 1970:pl.6 (*nomen nudum*)
Etymology:* Eponym honoring John R. Cann (1938–), Australia; herpetologist who studied turtles extensively in Australia and New Guinea.

Chelymys cooki Wells and Wellington 1985:8 (*nomen nudum*)
Emydura cooki
Type locality: "Macleay River, New South Wales (30 46'S x 152 18'E)" [Australia] [orange star, partially hidden].
Type specimen: AMS R44816, holotype, see Cann and Sadlier (2017).
Etymology:* Eponym honoring Robert Cook (0000–), Australia; amateur herpetologist.

Chelymys johncanni Wells and Wellington 1985:8 (*nomen nudum*)
Type locality: "Rouchel Brook, New South Wales" [Australia] [brown star, partially hidden].
Type specimen: AZM R101, holotype.
Etymology:* Eponym honoring John R. Cann (1938–), Australia; herpetologist who studied turtles extensively in Australia and New Guinea.

Emydura macquarii binjing Cann 1998:116 (10:42)
Type locality: "Clarence River and its tributaries in eastern New South Wales...29°45'S, 152°15'E" [Australia] [blue star].
Type specimen: AMS R59558, holotype, see Shea and Sadlier (1999) and Cann and Sadlier (2017).
Etymology:* Vernacular (Australian Aboriginal), *binjing*, local indigenous name for the species.

Emydura macquarii dharra Cann 1998:120 (10:42)
Type locality: "Macleay River and its tributaries in eastern New South Wales...30°54'S, 152°10'E" [Australia] [pink star].
Type specimen: AMS R59553, holotype, see Shea and Sadlier (1999) and Cann and Sadlier (2017).
Etymology:* Vernacular (Australian Aboriginal), *dharra*, local indigenous name for the species.

Emydura macquarii gunabarra Cann 1998:123 (10:42)


Type locality: "Hunter River and its tributaries in eastern New South Wales...32°09'S, 150°58'E" [Australia] [light gray star].
Type specimen: AMS 143596, holotype, see Shea and Sadlier (1999) and Cann and Sadlier (2017).
Etymology:* Vernacular (Australian Aboriginal), *gunabarra*, local indigenous name for the species.

Emydura macquarii dharuk Cann 1998:126 (10:42)
Type locality: "Norton's Basin, Nepean River, 0.5 km upstream from the junction of the Warragamba and Nepean Rivers at 33°52'S, 150°37'E...Sydney Basin in eastern New South Wales" [Australia] [dark gray star].
Type specimen: AMS 143598, holotype, see Shea and Sadlier (1999) and Cann and Sadlier (2017).
Etymology:* Ethnonym honoring the Australian Aboriginal people (known as Dharuk, Dahrug, or Darug) who lived in the area.

Emydura macquarii emmotti Cann, McCord, and Joseph-Ouni *in* McCord, Cann, and Joseph-Ouni 2003 (17:98)
Cooper Creek Turtle


Kate Hodges / Stonehenge, Thomson R., Queensland, Australia / megacephalic female


left: Arthur Georges / Eulbertie Waterhole, Cooper Creek, Tanbar, Queensland, Australia
right: Kate Hodges / Stonehenge, Thomson R., Queensland, Australia


**
(subspecies: *macquarii* = red, *emmotti* = blue, *krefftii* = green, *nigra* = pink; overlap = intergrades; stars = type localities)


Distribution: Australia (*Queensland*, *South Australia*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 87,155 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 332,145 sq. km
Size (Max SCL):* male 32.4 cm, female 36.8 cm (McCord et al. 2003; Cann and Sadlier 2017; McKnight et al. 2023)
Synonymy:
Chelymys windorah Wells and Wellington 1985:8 (*nomen nudum*)
Emydura windorah
Type locality: "Windorah district of south-west Queensland" [Australia] [orange star].
Type specimen: AZM R104, holotype.
Etymology:* Toponym for the Windorah District, Queensland, Australia.

Emydura macquarii emmotti Cann, McCord, and Joseph-Ouni *in* McCord, Cann, and Joseph-Ouni 2003:60
Emydura emmotti
Type locality: "Waterloo Station, shearing-shed waterhole, southwestern Queensland, Australia (24°13' S, 143°17' E)" [yellow star].
Type specimen: QM J51255, holotype, see Cann and Sadlier (2017).
Etymology:* Eponym honoring Angus Emmott (1962–), Australia; grazier, conservationist, and naturalist.

Emydura macquarii krefftii (Gray 1871c) (17:98)
Krefft's River Turtle


Arthur Georges / Grays Waterhole, Burnett R., Queensland, Australia


Duncan Limpus / Burnett R., Queensland, Australia / male


**
(subspecies: *macquarii* = red, *emmotti* = blue, *krefftii* = green, *nigra* = pink; overlap = intergrades; star = type locality)

Distribution: Australia (*Queensland*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 176,208 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 516,677 sq. km
Size (Max SCL): male 27.7 cm, female 34.2 cm (Georges 1985; Cann and Sadlier 2017)

Synonymy:

Chelymys krefftii Gray 1871c:366

Emydura krefftii, Emydura australis krefftii, Emydura macquarrii krefftii, Tropicochelymys krefftii, Emydura macquarii krefftii

Type locality: "Burnett's River" [Queensland, Australia] [yellow star].

Type specimen: NHMUK 1947.3.6.1 (formerly 1871.9.25.1), holotype, see Goode (1967), Cogger et al. (1983), and Cann and Sadlier (2017).

Etymology:* Eponym honoring Gerard Krefft (1830–1881), Germany / Australia; naturalist, herpetologist, and director at the Australian Museum, Sydney.

Chelymys victoriae marmorata Gray 1872d:506

Type locality: "east coast of Queensland, Burnett River" [Australia] [yellow star].

Type specimen: NHMUK 1871.9.25.5, holotype, see Cogger et al. (1983).

Etymology:* Latin, *marmor*, marble, and *-atus*, provided with; referring to the marbled appearance of the carapace.

Chelymys victoriae sulcata Gray 1872d:506

Type locality: "east coast of Queensland, Burnett River" [Australia] [yellow star].

Type specimens: NHMUK 1871.9.25.3–4, syntypes (2), see Cogger et al. (1983) and Georges and Thomson (2010).

Etymology:* Latin, *sulcus*, groove or furrow, and *-ata*, having; referring to the furrows on the carapace.

Emydura macquarii nigra McCord, Cann, and Joseph-Ouni 2003 (17:98)


Fraser Island Short-necked Turtle



Arthur Georges / Fraser Island, Queensland, Australia



left: Duncan Limpus, *right*: Arthur Georges / Fraser Island, Queensland, Australia



** (subspecies: *macquarii* = red, *krefftii* = green, *nigra* = pink; overlap = intergrades; star = type locality)

Distribution: Australia (*Queensland*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO): 2,150 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 4,380 sq. km

Size (Max SCL): male 19.7 cm, female 24.6 cm (Georges 1985; Itescu et al. 2014)

Synonymy:

Tropicochelymys insularis Wells and Wellington 1985:9 (*nomen nudum*)

Emydura insularis

Type locality: "Fraser Island, Queensland" [Australia] [yellow star, location estimated].

Type specimen: AZM R102, holotype.

Etymology:* Latin, *insula*, island, and *-aris*, pertaining to; referring to the insular distribution of the species.

Emydura macquarii nigra McCord, Cann, and Joseph-Ouni 2003:59

Emydura nigra

Type locality: "Lake McKenzie, Fraser Island, Queensland, Australia (25°27' S, 153°04' E)" [yellow star].


Type specimen: QM J48008, holotype, see Cann and Sadlier (2017).

Etymology:* Latin, *nigra*, black; referring to the shell color.

Emydura subglobosa (Krefft 1876) (21:32)

Red-bellied Short-necked Turtle

(includes 2 subspecies)



** (subspecies: *subglobosa* = red, *worrelli* = blue; orange dots = probable trade)

Distribution: Australia (*Northern Territory, Queensland*), Indonesia (*Papua, West Papua*), Papua New Guinea (*Southern*)

Introduced: Papua New Guinea (*New Britain, Northern*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 412,844 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 2,069,296 sq. km

Size (Max SCL): male 20.0 cm, female 26.1 cm (see subspp.)

IUCN Red List: Least Concern (LC) (ATTWG 2000). Previously: Least Concern (LC) [Not Listed] (Behler 1996)

TFTSG Provisional Red List: Least Concern (LC) (2009, 2018)

Emydura subglobosa subglobosa (Krefft 1876) (21:32)


New Guinea Red-bellied Short-necked Turtle, New Guinea Painted Turtle



Arthur Georges / Suki-Aramba Swamps, Fly R., Western Prov., Papua New Guinea



Anders G.J. Rhodin / Bore, Kemp Welch R., Central Prov., Papua New Guinea



** (subspecies: *subglobosa* = red; orange dots = probable trade; stars = type localities)

Distribution: Australia (*Queensland*), Indonesia (*Papua, West Papua*), Papua New Guinea (*Southern*)

Introduced: Papua New Guinea (*New Britain, Northern*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 252,202 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 623,926 sq. km

Size (Max SCL): male 20.0 cm, female 25.5 cm (Georges et al. 2006; Ceballos et al. 2013; Lyons et al. 2013)

Synonymy:

Euchelymys subglobosa Krefft 1876:390

Emydura subglobosa, Emydura australis subglobosa, Emydura macquarrii subglobosa, Chelymys subglobosa, Tropicochelymys subglobosa, Emydura subglobosa subglobosa

Type locality:* "Amama River S. E. New Guinea" [Papua New Guinea]. Restricted to "Naiabui S. E. New Guinea" [Papua New Guinea] by Boulenger (1888b:450); emended here to "Inawabui, between Bioto and Angabunga rivers, Central Province, Papua New Guinea (8º42'S, 146º38'E)" [yellow star].

Type specimen: MSNG CE2320, holotype, see Ogilby (1905), Capocaccia (1961), Goode (1967), Cogger et al. (1983), Cann and Sadlier (2017), and Smales et al. (2019a); genotyped by Kehlmaier et al. (2019a).

Etymology:* Latin, *sub-*, almost, little, and *globosa*, spherical; referring to the shape of the shell.

Emydura albertisii Boulenger 1888b:449

Emydura australis albertisii

Type locality: "Katow, S. E. New Guinea". Emended to "Mawatta, Binaturi River (9°08' S, 142°55' E), Papua New Guinea" [green star] by Cogger et al. (1983) and TTWG (2017:199).

Type specimens:* MSNG CE8430, syntypes (2), see Ogilby (1905), Capocaccia (1961), Goode (1967), Cogger et al. (1983), and Smales et al. (2019a); apparently lost.

Etymology:* Eponym honoring Luigi M. d'Albertis (1841–1901), Italy; explorer and naturalist who worked in New Guinea and collected the types.

Comment:* Capocaccia (1961) listed CE8430 as being two syntypes, but Goode (1967:74,77) listed CE8430 and CE8431 as separate syntypes and provided photographs of the larger female, labeled CE8430 in the plate, but CE8431 in the measurement table; Smales et al. (2019a:10,pl.2) noted the discrepancy and relabeled their reproduction of the plate as showing CE8431. However, G. Doria at MSNG (Smales, pers. comm.) has confirmed that CE8430 consisted of two syntype specimens that were lost in flooding of the museum in 1970, and CE8431 is a specimen of *Cuora amboinensis*.

Tropicochelymys goodei Wells and Wellington 1985:9 (*nomen nudum*)

Emydura goodei

Type locality: "Jardine River, Cape York Peninsula, Queensland" [Australia] [blue star].

Type specimen: AMS s/n, holotype.

Etymology:* Eponym honoring John Goode (1927–2008), United Kingdom / Australia; journalist and amateur herpetologist.

Emydura subglobosa angkibaanya Joseph-Ouni, McCord, Cann, Smales, and Freeman 2019b:55 (21:32)

Type locality: "Old River Crossing, Jardine River, Cape York Peninsula, Queensland, Australia ... 11°08'S, 142°21'E" [blue star].

Type specimen: AMS R37669, holotype.

Etymology:* Vernacular (Australian Aboriginal, Gudang), *angkibaanya*, local indigenous word for rainbow; referring to the taxon's bold facial and ventral coloration.

Emydura subglobosa worrelli (Wells and Wellington 1985) [07:99]
Worrell's Short-necked Turtle, Diamond-head Turtle




Jason Schaffer / Dugald R., Flinders basin, Queensland, Australia





Arthur Georges / Australia



** (subspecies: *subglobosa* = red; *worrelli* = blue; stars = type localities)

Distribution: Australia (*Northern Territory, Queensland*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 111,855 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 392,918 sq. km

Size (Max SCL): male 20.0 cm, female 26.1 cm (Cann and Sadlier 2017)

Synonymy:

Tropicochelymys leichhardti Wells and Wellington 1985:9 (*nomen nudum*)

Type locality: "Leichhardt River, Queensland" [Australia] [orange star, location estimated].

Type specimen: AZM 103, holotype.

Etymology:* Eponym honoring Friedrich Wilhelm Ludwig Leichhardt (1813–1848), Germany / Australia; explorer and naturalist in Australia in whose honor the Leichhardt River was named.

Tropicochelymys worrelli Wells and Wellington 1985:9

Emydura worrelli, Emydura subglobosa worrelli

Type locality: "Caranbirini Waterhole, ca. 21 km north of MacArthur River, Northern Territory (16°16' S x 136°05' E)" [Australia] [yellow star].


Type specimen: AMS 53689, holotype, see Cann and Sadlier (2017).

Etymology:* Eponym honoring Eric F.A. Worrell (1924–1987), Australia; naturalist and herpetologist.

Emydura tanybaraga Cann 1997b
Northern Yellow-faced Turtle



Jason Schaffer / nr. Coen, Archer River, Queensland, Australia



** (star = type locality)

Distribution: Australia (*Northern Territory, Queensland*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 96,233 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 943,540 sq. km

Size (Max SCL): female 28.5 cm (Cann and Sadlier 2017)

IUCN Red List: Not Evaluated (NE)

TFTSG Provisional Red List: Least Concern (LC) (2009, 2018)

Synonymy:

Emydura tanybaraga Cann 1997b:24

Emydura subglobosa tanybaraga

Type locality: "near Policeman Crossing, Daly River, Northern Territory (13°46' S x 130°43' E)" [Australia] [yellow star].

Type specimen: AMS R125498, holotype, see Shea and Sadlier (1999) and Cann and Sadlier (2017).

Etymology:* Vernacular (Australian Aboriginal), *tanybaraga*, local indigenous name for the species.

Myuchelys Thomson and Georges 2009 (09:45, 10:43, 17:99)

Australian Helmeted Turtles

(includes 4 species, 4 taxa)

Wollumbinia Wells 2007c:1 (unavailable name) (07:97, 10:43)

Type species: *Wollumbinia latisternum* [= *Elseya latisternum* Gray 1867], by original designation.

Etymology:* Vernacular (Australian Aboriginal, Bundjalung), *wollumbin*, rainmaker; referring to the local indigenous name for Mt. Warning in New South Wales, within the range of the genus.

Myuchelys Thomson and Georges 2009:33 (09:45)

Type species: *Myuchelys latisternum* [= *Elseya latisternum* Gray 1867], by original designation.

Etymology:* Vernacular (Australian Aboriginal), *myuna*, clear water, and Greek, *chelys*, turtle; referring to the preferred habitat.

Flaviemys Le, Reid, McCord, Naro-Maciel, Raxworthy, Amato, and Georges 2013:257 (17:99)

Type species: *Flaviemys purvisi* [= *Elseya purvisi* Wells and Wellington 1985], by original designation.

Etymology:* Latin, *flava*, yellow, and Greek, *emys*, turtle; referring to the yellow plastron.

Myuchelys bellii (Gray 1844) (07:97)


Bell's Sawshelled Turtle, Western Sawshelled Turtle



Darren Fielder / CBFTT / Bald Rock Creek, Queensland, Australia



Darren Fielder / CBFTT / Bald Rock Creek, Queensland, Australia



** (stars = type localities)

Distribution: Australia (*New South Wales, Queensland*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 16,358 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 23,915 sq. km

Size (Max SCL): male 22.7 cm, female 30.0 cm (Fielder et al. 2015 CBFTT)

CBFTT Account: Fielder, Chessman, and Georges (2015)

IUCN Red List: Endangered (EN B1+2c [ver. 2.3]) (TFTSG 1996). Previously: Endangered (EN) as *Elseya* sp. 1, Namoi River Snapping Turtle (TFTSG *in* Baillie and Groombridge 1996)

TFTSG Provisional Red List: Vulnerable (VU) (2018)

Synonymy:

Phrynops bellii Gray 1844:41

Hydraspis bellii, Elseya bellii, Wollumbinia bellii, Wollumbinia bellii bellii, Elseya latisternum bellii, Myuchelys bellii

Type locality: Not known. Restricted to "the upper reaches of both the Namoi-MacDonald and Gwydir Rivers, above the New England escarpment, in New South Wales" [Australia] [yellow star, location estimated] by Cann (1998:211).

Type specimen: OUM 8460, holotype, discussed by Cann (1998), Nowak-Kemp and Fritz (2010), Fielder (2013), and Cann and Sadlier (2017); genotyped by Kehlmaier et al. (2019a).

Etymology:* Eponym honoring Thomas Bell (1792–1880), United Kingdom; herpetologist, dental surgeon, and Professor of Zoology at King's College, London.

Elseya dorriani Wells 2002b:16 (*nomen nudum*) (07:97)

Wollumbinia bellii dorriani

Type locality:* "Bald Rock Creek, south-eastern Queensland" [Australia] [orange star].


Type specimen:* Not designated.

Etymology:* Eponym honoring Chris Dorrian (1942–2013), Australia; amateur herpetologist who discovered the species.

Myuchelys georgesi (Cann 1997a)
Bellinger River Sawshelled Turtle



John Cann / CBFTT / Bellinger River, New South Wales, Australia



Arthur Georges / CBFTT / Bellinger River, New South Wales, Australia




(star = type locality)

Distribution: Australia (*New South Wales*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 1,002 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 1,277 sq. km
Size (Max SCL): male 21.2 cm, female 24.0 cm (Blamires et al. 2005; Cann et al. 2015 CBFTT)
CBFTT Account: Cann, Spencer, Welsh, and Georges (2015)
IUCN Red List: Data Deficient (DD) (TFTSG 1996), as *Elseya sp. 3*, Bellinger River
TFTSG Provisional Red List: Critically Endangered (CR) (2015)
Synonymy:
Elseya georgesi Cann 1997a:18
Wollumbinia georgesi, Elseya latisternum georgesi, Myuchelys georgesi
Type locality: "Bellinger River 30°25' S, 152°46' E" [New South Wales, Australia] [yellow star].
Type specimen: AMS R31721, holotype, see Shea and Sadlier (1999) and Cann and Sadlier (2017).
Etymology:* Eponym honoring Arthur Georges (1953–), Australia; geneticist and herpetologist at the University of Canberra and turtle researcher.


Myuchelys latisternum (Gray 1867) [(09:47)]
Sawshelled Turtle, Common Sawshelled Turtle



Alastair Freeman / CBFTT / Chillagoe Creek, Queensland, Australia



Alastair Freeman / Porcupine Gorge, Queensland, Australia



(orange dots = introduced, translocated; stars = type localities)

Distribution: Australia (*New South Wales, Northern Territory, Queensland*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 276,563 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 1,870,207 sq. km
Size (Max SCL): male 22.4 cm, female 28.8 cm (Freeman and Cann 2014 CBFTT)

CBFTT Account: Freeman and Cann (2014)
IUCN Red List: Least Concern (LC) [Not Listed] (Behler 1996)
TFTSG Provisional Red List: Least Concern (LC) (2011)
Synonymy:

Elseya latisternum Gray 1867:44
Emydura latisternum, Elseya latisternum latisternum, Wollumbinia latisternum, Myuchelys latisternum
Type locality: "North Australia." Restricted to "Cape York" [Queensland, Australia] by Cann and Sadlier (2017:193), and to "Cape York Peninsula north of latitude 11°40'S" [Queensland, Australia] [yellow star, location estimated] by Smales et al. (2019b:705).
Type specimens: NHMUK 1867.5.13.19, lectotype, designated by Smales et al. (2019b:705); NHMUK 1867.5.13.19–20, syntypes (2), previously discovered and designated by Cann and Sadlier (2017:193); NHMUK 1947.3.4.13, previously recorded as "holotype" by Goode (1967), Cogger et al. (1983), and Cann (1998), is not a type, see Cann and Sadlier (2017) and Smales et al. (2019b).
Etymology:* Latin, *latum*, broad, wide, extended, and Greek, *sternon*, chest, sternum, thorax; referring to the relatively broad plastron.


Euchelymys spinosa Gray 1871a:118 (09:47)
Elseya spinosa
Type locality: "North Australia."
Type specimen: NHMUK 1946.1.22.77 (formerly 1866.6.19.1), referred to as holotype by Cann (1998) and Cann and Sadlier (2017); designated lectotype by Smales et al. (2019b:705).
Etymology:* Latin, *spina*, thorn, spike; referring to the numerous long, pointed tubercles on the neck.


Elseya latisternon Gray 1871b:292 (*nomen novum*)
Etymology:* Latin, *latum*, broad, wide, extended, and Greek, *sternon*, chest, sternum, thorax; referring to the relatively broad plastron.
Comment:* Unjustified emendation or error for *E. latisternum*.

Wollumbinia dorsii Wells 2009:2 (unavailable name) (09:46, 10:43)
Type locality: "Richmond River, near Wiangaree, New South Wales" [Australia] [orange star].
Type specimen: AMS R172224, holotype, see Cann and Sadlier (2017).
Etymology:* Eponym honoring Marc Dorse (1958–), Australia; educator and herpetologist.

Myuchelys purvisi (Wells and Wellington 1985) (14:46, 17:99, 21:33)
Manning River Sawshelled Turtle


Phil Spark / Barrington R., New South Wales, Australia


left: Phil Spark / Barrington R., New South Wales, Australia
right: Arthur Georges / Barnard R., New South Wales, Australia


**
(star = type locality)

Distribution: Australia (*New South Wales*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 4,598 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 6,934 sq. km
Size (Max SCL): male 17.5 cm, female 22.9 cm (Cann and Sadlier 2017)
IUCN Red List: Data Deficient (DD) (TFTSG 1996), as *Elseya sp. 2*, Manning River
TFTSG Provisional Red List: Near Threatened (NT) (2011)
Synonymy:

Elseya purvisi Wells and Wellington 1985:8
Wollumbinia purvisi, Elseya latisternum purvisi, Myuchelys purvisi, Flaviemys purvisi
Type locality: "a river 15 km S., 32.3 km E. of Nowendoc, New South Wales (31º39' S x 152º04' E. elevation 183 m)" [Australia] [yellow star].
Type specimen: AMS 44654, holotype; on loan to UU since 1976, see Cann and Sadlier (2017:208), now presumably there.
Etymology:* Eponym honoring Malcolm Purvis (?–), Australia; teacher and amateur herpetologist.

Rheodytes Legler and Cann 1980

Fitzroy River Turtles

(includes 1 species, 1 taxon)

Rheodytes Legler and Cann 1980:2

Type species: *Rheodytes leukops* Legler and Cann 1980, by original designation.


Etymology:* Greek, *reos*, current, flowing, stream, and *dytes*, diver; referring to the habitat of the species in fast flowing waters.

Rheodytes leukops Legler and Cann 1980


Fitzroy River Turtle



Stephen M. Zozaya / Connors R., upper Fitzroy R., Queensland, Australia



Jason Schaffer / Connors R., upper Fitzroy R., Queensland, Australia



(star = type locality)

Distribution: Australia (*Queensland*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 21,385 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 63,781 sq. km

Size (Max SCL): male 26.2 cm, female 26.2 cm (Cann and Legler 1994)

IUCN Red List: Vulnerable (VU A1c+2c, D2 [ver. 2.3]) (ARASG 1996)

TFTSG Provisional Red List: Endangered (EN) (2018)

Synonymy:

Rheodytes leukops Legler and Cann 1980:2

Elseya leukops

Type locality: "Fitzroy River, 63 km N and 25 km E of Duaringa, Queensland, Australia, elevation 40 m (23°09' S 149°55' E)" [yellow star].

Type specimen: QM J31701, holotype, see Cogger et al. (1983) and Cann and Sadlier (2017).

Etymology:* Greek, *lefkos*, white, bright, shining, and *ops*, eye; referring to the distinct white eye ring around the iris.

SUBFAMILY PSEUDEMYDURINAE Gaffney 1977 (17:100)

Western Swamp Turtles

(includes 1 genus, 1 species, 1 taxon)

Pseudemydurinae Gaffney 1977:24

Pseudemydura Siebenrock 1901 (17:100)

Western Swamp Turtles

(includes 1 species, 1 taxon)

Pseudemydura Siebenrock 1901:248

Type species: *Pseudemydura umbrina* Siebenrock 1901, by original monotypy.

Etymology:* Latin, *pseudo*, false, and Greek, *emys*, turtle, and *oura*, tail, rump; referring to the long and large tail in males.

Pseudemydura umbrina Siebenrock 1901 (17:100)


Western Swamp Turtle, Western Swamp Tortoise



Gerald Kuchling / TCC / Ellen Brook Nature Reserve, nr. Perth, Western Australia, Australia



Gerald Kuchling / Ellen Brook Nature Reserve, nr. Perth, Western Australia, Australia



(star = type locality)

Distribution: Australia (*Western Australia*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO): 1,884 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 2,388 sq. km

Size (Max SCL): male 15.5 cm, female 14.2 cm (Burbidge 1967; Cann and Legler 1994; Durrell and Keeley 2019)

IUCN Red List: Critically Endangered (CR A1c, B1+2c, C1+2b, D [ver. 2.3]) (TFTSG 1996)

TFTSG Provisional Red List: Critically Endangered (CR) (2011)

CITES: Appendix I (1975)

USA ESA:* Endangered (USFWS 1970)

Synonymy:

Pseudemydura umbrina Siebenrock 1901:249

Type locality: "Australien" [Australia] [no star].

Type specimen: NMW 1296 (formerly 1839 89(8450)), holotype, see Goode (1967), Tiedemann and Häupl (1980), Cogger et al. (1983); Tiedemann et al. (1994), Cann and Sadlier (2017), and Gemel et al. (2019).

Etymology:* Latin, *umbra*, darkness, shadow; referring to the dark color of the carapace and dorsal head.

Emydura inspectata Glauert 1954:125

Type locality: "Warbrook, about 24 miles north of Perth...Swan River District" [Western Australia, Australia] [green star].

Type specimen: WAM R11092, holotype, see Cogger et al. (1983) and Ellis and Georges (2015).

Etymology:* Latin, *inspectatus*, observed, inspected; possibly referring to the holotype of this taxon having attracted considerable attention at an animal exhibition in 1953, since it was interpreted as being a specimen of a supposedly unknown species.

Superfamily Pelomedusoidea Baur 1893 (21:3)

Pelomedusoid Side-necked Turtles

(includes 2 families, 5 genera, 35 species, 40 taxa)

Pelomedusoidea Baur 1893a:212

Pelomedusoides Broin 1988:105,130

Family Pelomedusidae Cope 1868 (11:19)


Pelomedusid Side-necked Turtles

(includes 2 genera, 27 species, 32 taxa)

Hydraspidina Bonaparte 1836:3 (*partim*)

Pelomedusidae Cope 1868:119

Pelomedusinae Zangerl 1948:48



Pelomedusidae Species Richness

Pelomedusa Wagler 1830b (11:19, 17:101) (25)*

African Helmeted Turtles

(includes 10 species, 10 taxa)

Pelomedusa Wagler 1830b:136 (*nomen protectum*)

Type species: *Pelomedusa galeata* [= *Testudo galeata* Schoepff 1792], by original monotypy.

Etymology:* Greek, *pelos*, mud, and *medusa*, a mythical being with a head of snakes; referring to the snake-like neck.

Comment: Name conserved by ICZN (1989), see Smith et al. (1980b).

Pentonyx Duméril and Bibron 1835:389

Type species: *Pentonyx galeata* [= *Testudo galeata* Schoepff 1792], by original designation.


Etymology:* Greek, *pente*, five, and *onyx*, claw, nail; refersring to the five claws on all four feet.

Pelomedusa barbata Petzold, Vargas-Ramírez, Kehlmaier, Vamberger, Branch, Du Preez, Hofmeyr, Meyer, Schleicher, Široký, and Fritz 2014 (17:101)


Arabian Helmeted Turtle, Arabian Marsh Turtle



Johannes Els / Taif, Saudi Arabia



Anders G.J. Rhodin / Taif, Saudi Arabia / captivity / Sharjah Zoo, United Arab Emirates



** (orange dots = possible trade or inaccurate localities; star = type locality)

Distribution:* Saudi Arabia (*Asir*, *Al-Behah*, *Makkah*, *Jizan*), Yemen

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 89,533 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 257,579 sq. km

Size (Max SCL):* male 21.6 cm, female 20.1 cm (Gasperetti et al. 1993; Petzold et al. 2014); ca. 25.0 cm (Els et al. 2022 CBFTT)

CBFTT Accounts:* Els, Alqahtani, Fritz, Shobrak, Busais, and Burriel-Carranza (2022); Boycott and Bourquin (2008) [as part of *Pelomedusa subrufa* sensu lato]

IUCN Red List: Not Evaluated (NE)

TFTSG Provisional Red List: Endangered (EN) (2016)

Synonymy:

Pelomedusa barbata Petzold, Vargas-Ramírez, Kehlmaier, Vamberger, Branch, Du Preez, Hofmeyr, Meyer, Schleicher, Široký, and Fritz 2014:530

Type locality: "Zinjibar, Abyan, Yemen, N13°7.75 E45°22.81" [yellow star].

Type specimen: MTD 24637, holotype; ZFMK 87122, paratype, genotyped by Petzold et al. (2014).

Etymology:* Latin, *barba*, beard; referring to the two to three very long barbels under the chin.

Pelomedusa galeata (Schoepff 1792) (17:101, 21:34)
South African Helmeted Turtle, Cape Terrapin




William R. Branch / Port Elizabeth, South Africa





Tomas Diagne / South Africa



(yellow dots = genotyped, gray dots = ungenotyped, presumed; stars = type localities)

Distribution: Botswana (?), Eswatini (Swaziland), Lesotho, Mozambique, South Africa

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 812,031 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 1,200,744 sq. km

Size (Max SCL): male 32.5 cm, female 29.0 cm (Boycott and Bourquin 2008 CBFTT; Petzold et al. 2014)

CBFTT Account: Boycott and Bourquin (2008) [as part of *Pelomedusa subrufa* sensu lato]

IUCN Red List: Least Concern (LC) (Hofmeyr and Fritz 2018)

Synonymy:

Testudo scabra Retzius *in* Schoepff 1792:12 (junior homonym and *nomen nudum*, not = *Testudo scabra* Linnaeus 1758 [= *Rhinoclemmys punctularia punctularia*])
Etymology:* Latin, *scabra*, rough, shabby, mangy, unclean; referring to the rough carapacial surface.

Testudo galeata Schoepff 1792:12 (*nomen protectum*)
Emys galeata, Pelomedusa galeata, Pentonyx galeata, Hydraspis (Pelomedusa) galeata, Hydraspis galeata, Pelomedusa galeata galeata
Type locality: "India orientali." Restricted to "near Cape Town" [South Africa] [yellow star] by Hewitt (1935:326).
Type specimen: LUZM Lxxx/6481 (formerly ZMUL 6481), lectotype, designated by Fritz et al. (2014b:508); NRM 7043, paralectotype, specimen figured in Schoepff (1792:pl.3,f.1), designated "holotype" by Rhodin and Carr (2009:4).
Etymology:* Latin, *galea*, helmet, hood; referring to the head scalation that looks like a helmet.
Comment: Name conserved by ICZN (1989), see Smith et al. (1980b).

Pentonyx capensis Duméril and Bibron 1835:390
Type locality: "au cap de Bonne-Espérance, dans l'Ile de Madagascar." Restricted to "Kap der Guten Hoffnung" [Cape of Good Hope, South Africa] by Mertens (1937:139), but invalid designation; and to "Cape of Good Hope" [South Africa] [green star] by Fritz et al. (2014b:510).
Type specimen: MNHN 9506, lectotype, designated by Fritz et al. (2014b:510), see also Bour (2009a) who labeled this name a *nomen novum*; genotyped by Fritz et al. (2014b) and Petzold et al. (2014).
Etymology:* Toponym for the Cape of Good Hope, South Africa.

Pelomedusa nigra Gray 1863b:99
Pelomedusa galeata nigra, Pelomedusa subrufa nigra
Type locality: "Natal" [South Africa] [blue star, location estimated].
Type specimen: NHMUK 1849.1.30.27, lectotype, designated by Fritz et al. (2014b:514); genotyped by Fritz et al. (2014b) and Petzold et al. (2014).
Etymology:* Latin, *nigra*, black; referring to the shell color.

Pelomedusa galeata orangensis Hewitt 1935:332
Pelomedusa subrufa orangensis
Type locality: "presumably from the Kimberley neighbourhood" [South Africa] [pink star].
Type specimen: KM s/n, holotype, specimen figured (pl.32), apparently lost; PEM 9404, 9408, paratypes, genotyped by Fritz et al. (2014b).
Etymology:* Toponym for the Orange River, northern South Africa.

Pelomedusa galeata devilliersi Hewitt 1935:337
Type locality: "Besondermeid, Steinkopf, Namaqualand, C.P." [South Africa] [gray star].
Type specimen: PEM 14962, holotype; genotyped by Fritz et al. (2014b) and Petzold et al. (2014).
Etymology:* Eponym honoring Cornelius G.S. de Villiers (1894–1978), South Africa; physiologist and zoologist at Stellenbosch University.


Pelomedusa gehafie (Rüppell 1835) (17:101)
Eritrean Helmeted Turtle



Tomáš Mazuch / Ghinda, Eritrea



left: Tomáš Mazuch / Ghinda, Eritrea
right: Clifford Dorse / Kurmuk, Ethiopia



** (yellow dots and star = genotyped, gray dots = ungenotyped, presumed; black dots and gray shading = taxonomically unspecified populations of *P. subrufa* sensu lato, orange/yellow dot = genotyped *P. gehafie* specimen from questionable trade locality; orange dot = probable trade or translocated; stars = type localities)

Distribution: Eritrea, Ethiopia, Sudan

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 268,028 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 598,554 sq. km

Size (Max SCL): 17.8 cm (Petzold et al. 2014)

CBFTT Account: Boycott and Bourquin (2008) [as part of *Pelomedusa subrufa* sensu lato]

IUCN Red List: Not Evaluated (NE)

TFTSG Provisional Red List: Data Deficient (DD) (2018)

Synonymy:

Pentonyx gehafie Rüppell 1835:2

Pelomedusa galeata gehafie, Pelomedusa subrufa gehafie, Pelomedusa gehafie

Type locality: "östlichen Abhange der abyssinischen Küstengebirge" [eastern slope of Abyssinian coastal mountains] [Eritrea] [yellow star, location estimated]; restricted to "Massaua" [Masawa, Eritrea] [orange star] by Mertens (1937:140), but invalid designation (Fritz et al. 2014b:511).

Type specimen: SMF 7947, lectotype, designated by Mertens (1937:140); genotyped by Fritz et al. (2014b) and Petzold et al. (2014).

Etymology:* Vernacular (Eritrean, Tigrinya), *gehafie,* local indigenous name for the species in the Massaua region.

Pelomedusa gehafiae Gray 1844:38 (*nomen novum*)

Comment: Unjustified emendation of *gehafie*.

Etymology:* Vernacular (Eritrean, Tigrinya), *gehafie,* local indigenous name for the species in the Massaua region.

Pelomedusa galeata disjuncta Vaillant and Grandidier 1910:56

Type locality: Not designated. Restricted to "shore of Lake Abaya, Sidamo, Ethiopia" [green star] by Loveridge (1941:480), invalid designation; and to "Abyssinia...the eastern slope of the coastal mountains in present-day Eritrea" [yellow star, location estimated] by Fritz et al. (2014b:515).

Type specimen: MNHN 7870, lectotype, designated by Fritz et al. (2014b:515).

Etymology:* Latin, *dis-*, not, and *junctus*, joined; referring to the disjunct separated pectoral scutes.


Pelomedusa kobe Petzold, Vargas-Ramírez, Kehlmaier, Vamberger, Branch, Du Preez, Hofmeyr, Meyer, Schleicher, Široký, and Fritz 2014 (17:101)
Tanzanian Helmeted Turtle



Melita Vamberger / Akagera National Park, Rwanda / male



Melita Vamberger / Akagera National Park, Rwanda / male



** (yellow dots and star = genotyped, gray dots = ungenotyped, presumed; orange dot = questionable, possible trade, star = type locality)

Distribution:* Rwanda, Tanzania, Uganda (?)

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 277,070 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 529,758 sq. km

Size (Max SCL):* male 21.9 cm, female 18.9 cm (Umulisa et al. 2025)

CBFTT Account: Boycott and Bourquin (2008) [as part of *Pelomedusa subrufa* sensu lato]

IUCN Red List: Not Evaluated (NE)

TFTSG Provisional Red List: Data Deficient (DD) (2018)

Synonymy:

Pelomedusa kobe Petzold, Vargas-Ramírez, Kehlmaier, Vamberger, Branch, Du Preez, Hofmeyr, Meyer, Schleicher, Široký, and Fritz 2014:535

Type locality: "Naberera, Manyara, Tanzania, S4°11.66 E36°55.74" [yellow star].

Type specimen: ZSM 334/1978, holotype; genotyped by Petzold et al. (2014).


Etymology:* Vernacular (Indigenous, Swahili), *kobe*, pond turtle; local indigenous name for the species.

Pelomedusa neumanni Petzold, Vargas-Ramírez, Kehlmaier, Vamberger, Branch, Du Preez, Hofmeyr, Meyer, Schleicher, Široký, and Fritz 2014 (17:101)


East African Helmeted Turtle, Neumann's Helmeted Turtle



Tomáš Mazuch / nr. Turmi, Omo Region, Ethiopia



left: Tomáš Mazuch / nr. Turmi, Omo Region, Ethiopia
right: Russell A. Mittermeier / Samburu, Kenya



** (yellow dots and star = genotyped, gray dots = ungenotyped, presumed, black dots and gray shading = taxonomically unspecified populations of *P. subrufa* sensu lato; orange dots = questionable, possible trade; star = type locality)

Distribution: Burundi, Congo (DRC), Ethiopia, Kenya, Rwanda, Tanzania, Uganda, Zambia

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 597,534 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 1,390,455 sq. km

Size (Max SCL): 19.4 cm (Petzold et al. 2014)

CBFTT Account: Boycott and Bourquin (2008) [as part of *Pelomedusa subrufa* sensu lato]

IUCN Red List: Not Evaluated (NE)

TFTSG Provisional Red List: Data Deficient (DD) (2018)

Synonymy:

Pelomedusa neumanni Petzold, Vargas-Ramírez, Kehlmaier, Vamberger, Branch, Du Preez, Hofmeyr, Meyer, Schleicher, Široký, and Fritz 2014:537

Type locality: "Kakamega, Kenya, N0°17.04 E34°44.52" [yellow star].

Type specimen: NMP6V 74974, holotype.

Etymology:* Eponym honoring Oscar R. Neumann (1867–1946), Germany; ornithologist and mammalogist who did field work in East Africa and worked at the Museum für Naturkunde, Berlin.


Pelomedusa olivacea (Schweigger 1812) [17:101]
Sahelian Helmeted Turtle



Laurent Chirio / Maradi, Niger



left: Luca Luiselli / Pama North, Burkina Faso
right: Pearson McGovern / Senegal



(yellow dots and star = genotyped, gray dots = ungenotyped, presumed; stars = type localities)

Distribution: Benin, Burkina Faso, Gambia, Ghana, Guinea, Guinea-Bissau, Ivory Coast (Côte d'Ivoire), Mali, Mauritania, Niger, Nigeria, Senegal, Sierra Leone, Togo

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 2,009,750 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 2,846,805 sq. km

Size (Max SCL): 16.8 cm (Petzold et al. 2014)

CBFTT Account: Boycott and Bourquin (2008) [as part of *Pelomedusa subrufa* sensu lato]

IUCN Red List: Not Evaluated (NE)

TFTSG Provisional Red List: Least Concern (LC) (2018)

Synonymy:


Emys olivacea Schweigger 1812:307 (senior homonym, not = *Emys olivacea* Gray 1856b [= *Trachemys stejnegeri stejnegeri*])
Hydraspis (Pelomedusa) olivacea, Hydraspis olivacea, Pelomedusa subrufa olivacea, Pelomedusa olivacea
Type locality: "sabulosis Nigritiae" [Senegal] [yellow star, location estimated].
Type specimen: MNHN 7971, holotype, see Bour (2009a), Ceríaco and Bour (2012), and Fritz et al. (2014); Uetz et al. (2019) erroneously listed ZFMK 17076 as neotype, but that specimen is the neotype of *Pelomedusa gasconi*.
Etymology:* Latin, *oliva*, olive, and *-aceus*, belonging to, having the nature of; referring to the olive-colored carapace.

Pelomedusa gasconi Rochebrune 1884:25
Type locality:* "Dagana, Saidé, lac de N'Guer, marigot des Maringouins" [= Merinaghene, Lac de Guiers, Senegal] [green star]. Restricted to "Dagana, Senegal" by Loveridge (1941:480), invalid designation; and to "Dakar" [Senegal] [yellow star] by Fritz et al. (2014b:514) by neotype designation.
Type specimens: None originally preserved; ZFMK 17076, neotype, designated and genotyped by Fritz et al. (2014b:514).
Etymology:* Eponym honoring Alfred Gasconi (1842–1929), France / Senegal; politician who obtained the original types.

Pelomedusa schweinfurthi Petzold, Vargas-Ramírez, Kehlmaier, Vamberger, Branch, Du Preez, Hofmeyr, Meyer, Schleicher, Široký, and Fritz 2014 [17:101]
Central African Helmeted Turtle, Schweinfurth's Helmeted Turtle



Enrico Morawa / holotype, SMF 56161, female / Liria, Central Equatoria, South Sudan
from Petzold et al. (2014:Fig. 4)



(yellow dots and star = genotyped, gray dots = ungenotyped, presumed, black dots and gray shading = taxonomically unspecified populations of *P. subrufa* sensu lato; star = type locality)

Distribution: Central African Republic, Congo (DRC), Ethiopia (?), South Sudan, Sudan (?), Uganda

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 912,375 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 1,060,593 sq. km

Size (Max SCL): 15.7 cm (Petzold et al. 2014)

CBFTT Account: Boycott and Bourquin (2008) [as part of *Pelomedusa subrufa* sensu lato]

IUCN Red List: Not Evaluated (NE)

TFTSG Provisional Red List: Least Concern (LC) (2018)

Synonymy:

Pelomedusa schweinfurthi Petzold, Vargas-Ramírez, Kehlmaier, Vamberger, Branch, Du Preez, Hofmeyr, Meyer, Schleicher, Široký, and Fritz 2014:539
Type locality: "Liria, Central Equatoria, South Sudan, N4°38.66 E32°4.83" [yellow star].
Type specimen: SMF 56161, holotype; genotyped by Petzold et al. (2014).
Etymology:* Eponym honoring Georg A. Schweinfurth (1836–1925), Germany; botanist, paleontologist, and African explorer who collected one of the types.

Pelomedusa somalica Petzold, Vargas-Ramírez, Kehlmaier, Vamberger, Branch, Du Preez, Hofmeyr, Meyer, Schleicher, Široký, and Fritz 2014 [(17:101)]
Somalian Helmeted Turtle




Tomáš Mazuch / Daarbuduq, Somalia





Tomáš Mazuch / Las Geel, Somalia



** (yellow dot and star = genotyped, gray dots = ungenotyped, presumed, black dots and gray shading = taxonomically unspecified populations of *P. subrufa* sensu lato; orange dot = probable translocated; star = type locality)

Distribution: Djibouti (?) (*extirpated?*), Ethiopia, Somalia

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 268,559 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 475,647 sq. km

Size (Max SCL): 15.7 cm (Petzold et al. 2014)

CBFTT Account: Boycott and Bourquin (2008) [as part of *Pelomedusa subrufa* sensu lato]

IUCN Red List: Not Evaluated (NE)

TFTSG Provisional Red List: Data Deficient (DD) (2018)

Synonymy:


Pelomedusa somalica Petzold, Vargas-Ramírez, Kehlmaier, Vamberger, Branch, Du Preez, Hofmeyr, Meyer, Schleicher, Široký, and Fritz 2014:540

Type locality: "Borama district, Awdal, Somaliland/Somalia, N9°55 E43°10, 4500 ft." [yellow star].

Type locality: NHMUK 1970.1481, holotype; NMW 24449, paratype, genotyped by Petzold et al. (2014).

Etymology:* Toponym for Somalia.

Pelomedusa subrufa (Bonnaterre 1789) **(sensu lato)** [(10:45, 11:19, 17:101)]
Helmeted Turtle, African Helmeted Terrapin
(includes *P. subrufa* sensu stricto and five taxonomically unspecified populations of *P. subrufa* sensu lato)




** (yellow dots = genotyped, gray dots = ungenotyped, presumed, black dots and gray shading = taxonomically unspecified populations of *P. subrufa* sensu lato, orange dots = probable trade)

Distribution: Angola, Botswana, Cameroon, Central African Republic, Chad, Congo (DRC), Congo (ROC), Gabon, Kenya, Madagascar (*prehistorical introduction?*), Malawi, Mozambique, Namibia, Niger, Nigeria, Somalia, South Africa, Sudan, Tanzania, Zambia, Zimbabwe


Pelomedusa subrufa (Bonnaterre 1789) **(sensu stricto)** (10:45, 11:19, 17:101)
Helmeted Turtle, African Helmeted Terrapin



Anders G.J. Rhodin / CBFTT / Tsihombe, Cap Sainte Marie, Madagascar



Anders G.J. Rhodin / CBFTT / Tsihombe, Cap Sainte Marie, Madagascar



** (yellow dots = genotyped, gray dots = ungenotyped, presumed, black dots and gray shading = taxonomically unspecified populations of *P. subrufa* sensu lato, orange dots = probable trade; stars = type localities)

Distribution: Angola, Botswana, Congo (DRC), Kenya, Madagascar (*prehistorical introduction?*), Malawi, Mozambique, Namibia, South Africa, Tanzania, Zambia, Zimbabwe

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 2,945,326 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 8,493,444 sq. km

Size (Max SCL): 19.7 cm (Petzold et al. 2014)

CBFTT Account: Boycott and Bourquin (2008) [*Pelomedusa subrufa* sensu lato]

IUCN Red List: Least Concern (LC) [Not Listed] (Behler 1996)

TFTSG Provisional Red List: Data Deficient (DD) (2018)

Synonymy:

Testudo planitia Meuschen 1778:11 (senior homonym and *nomen dubium*, not = *Testudo planitia* Gmelin 1789 [= *Macrochelys temminckii*])
Hydraspis (Pelomedusa) planitia, Hydraspis planitia
Type locality: Not designated.
Type specimen: Not known or located.
Etymology:* Latin, *planus*, plane, flat, and *itius*, characterized by; referring to the relatively flat shell.

Testudo subrufa Lacepède 1788:173, synopsis[table] (07:102, 09:6) (*nomen suppressum*)
Type locality: "l'Inde" [India] [in error]. Restricted to "Kap der Guten Hoffnung" [Cape of Good Hope, South Africa] [in error] [orange star] by Mertens (1937:139); and to "Taolañaro (Fort-Dauphin), République Malagasy (Madagascar)" [yellow star] by Bour (1982c:535).
Type specimen: MNHN 7970, holotype, type specimen figured (pl. opp.173), see Bour (2009a); genotyped by Fritz et al. (2014b).
Etymology:* Latin, *sub-*, under, below, and *rufus*, red; referring to the brownish-red color of the plastron.
Comment: Name suppressed by ICZN (2005a) as published in a rejected and invalid non-binomial work, see Savage (2003).

Testudo subrufa Bonnaterre 1789:28
Emys subrufa, Pelomedusa subrufa, Chelys (Hydraspis) subrufa, Chelys subrufa, Hydraspis subrufa, Pelomedusa galeata subrufa, Pelomedusa subrufa subrufa
Type locality: "l'Inde" [India] [in error]. Restricted to "Taolañaro, Madagascar" [yellow star] by Fritz et al. (2014b:507), following Bour (1982c:535).
Type specimen: MNHN 7970, holotype, type specimen figured in Lacepède (1788:pl.opp.137), see Bour (2009a); genotyped by Fritz et al. (2014b).
Etymology:* Latin, *sub-*, under, below, and *rufus*, red; referring to the brownish-red color of the plastron.

Testudo rubra Meyer 1790:83 (*nomen novum, dubium et obli-tum*) (09:8)
Etymology:* Latin, *rubra*, red; referring to the brownish-red color of the plastron.
Comment: Unjustified replacement name for *subrufa*.

Testudo badia Donndorff 1798:34 (*nomen novum*)
Etymology:* Latin, *badius*, brown or chestnut-colored; referring to the brownish color of the plastron.

Testudo rubicunda Suckow 1798:49 (*nomen novum*)
Etymology:* Latin, *rubra*, red-colored; referring to the brownish color of the plastron.

Pentonix americana Cornalia 1849:13 (*nomen dubium*) (17:105)
Pentonyx americana
Type locality: "Flum. prope Novaeboracum" [= Novum Eboracum = New York, USA] [in error].
Type specimen: MSNM s/n, holotype, destroyed during World War II, discussed by Fritz et al. (2014).
Etymology:* Toponym for America.

Pelomedusa mossambicensis Peters *in* Lichtenstein and von Martens 1856:2 (*nomen nudum*)
Etymology:* Toponym for Mozambique.

Pelomedusa mozambica Peters *in* Gray 1856b:53 (*nomen nudum*)
Etymology:* Toponym for Mozambique.

Pelomedusa galeata damarensis Hewitt 1935:338
Pelomedusa subrufa damarensis
Type locality: "Quickborn, near Okahandja, South West Africa" [Namibia] [green star].
Type specimen: PEM 14953, lectotype, designated by Fritz et al. (2014b:515); genotyped by Fritz et al. (2014b) and Petzold et al. (2014).
Etymology:* Toponym for the Damara region of central Namibia.

Pelomedusa subrufa wettsteini Mertens 1937:141
Type locality: "Majunga, West-Madagaskar" [Mahajanga, Madagascar] [blue star].
Type specimen: SMF 7958, holotype, see Mertens (1967b).
Etymology:* Eponym honoring Otto von Wettstein-Westersheim (1892–1967), Austria; herpetologist and curator at the Naturhistorisches Museum, Vienna.

Pelomedusa variabilis Petzold, Vargas-Ramírez, Kehlmaier, Vamberger, Branch, Du Preez, Hofmeyr, Meyer, Schleicher, Široký, and Fritz 2014 (17:101)


West African Helmeted Turtle



Tomas Diagne / Afram Arm, Ghana



Luca Luiselli / Adjohoun, Benin



(yellow dots and star = genotyped, gray dots = ungenotyped, presumed, black dots and gray shading = taxonomically unspecified populations of *P. subrufa* sensu lato; star = type locality)

Distribution: Benin, Burkina Faso, Cameroon, Equatorial Guinea, Ghana, Guinea, Ivory Coast (Côte d'Ivoire), Liberia, Nigeria, Togo

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 996,640 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 1,727,930 sq. km

Size (Max SCL): male 24.8 cm (Petzold et al. 2014)

CBFTT Account: Boycott and Bourquin (2008) [as part of *Pelomedusa subrufa* sensu lato]

IUCN Red List: Not Evaluated (NE)

TFTSG Provisional Red List: Data Deficient (DD) (2018)

Synonymy:

Pelomedusa variabilis Petzold, Vargas-Ramírez, Kehlmaier, Vamberger, Branch, Du Preez, Hofmeyr, Meyer, Schleicher, Široký, and Fritz 2014:543
Type locality: "Gold Coast, Ghana" [yellow star, location estimated].
Type specimen: SMF 58075, holotype.
Etymology:* Latin, *variabilis*, changeable; referring to the very variable shell morphology.

Pelusios Wagler 1830b (11:19) (17:102)

African Hinged Terrapins

(includes 17 species, 22 taxa)

Cheliphus Rafinesque 1815:75 (*nomen nudum*)
Etymology:* Greek, *chelys*, turtle, and *physa* or *physes*, breath, vesicle; reference unclear (Frétey 2024).
Comment: Discussed by Frétey (2024).

Sternothaerus Bell 1825a:305 (*partim*, *nomen suppressum*)
Type species: *Sternothaerus leachianus* Bell 1825a [= subjective synonym of *Emys castanea* Schweigger 1812 = *Pelusios castaneus*], by subsequent designation by Bell (1828c:515); not *Sternothaerus odoratus* Bell [= *Testudo odorata* Latreille *in* Sonnini and Latreille 1801], by incorrect designation by Fitzinger (1843:290).
Etymology:* Greek, *sternon*, chest, sternum, and *thairos*, hinge; referring to the plastral hinge.
Comment: Name suppressed by ICZN (1989), see Smith et al. (1980b) and Bour and Dubois (1984c).

Pelusios Wagler 1830b:137 (*nomen protectum*)
Type species: *Pelusios subniger* Wagler [= *Testudo subnigra* Lacepède 1788 (*nomen suppressum*) = *Testudo subnigra* Bonnaterre 1789], by subsequent designation by Fitzinger (1843:29).
Etymology:* Greek, *pelos*, mud, swamp; referring to the typical habitat of muddy, swampy waters.
Comment:* Name conserved by ICZN (1989), see Smith et al. (1980b) and Bour and Dubois (1984c).

Cheliphus Rafinesque 1832:64
Type species:* *Testudo subnigra* Lacepède 1788 [= *Testudo subnigra* Bonnaterre 1789 = *Pelusios subniger*], by subsequent designation by Bour and Dubois (1984c:198).
Etymology:* Greek, *chelys*, turtle, and *physa* or *physes*, breath, vesicle; reference unclear (Frétey 2024).

Pelusius Agassiz 1847:278 (*nomen novum*)
Etymology:* Greek, *pelos*, mud, swamp; referring to the typical habitat of muddy, swampy waters.
Comment:* Unjustified emendation of *Pelusios*.

Tanoa Gray 1863f:193
Type species: *Sternothaerus (Tanoa) sinuatus* (Smith 1838), by subsequent designation by Lindholm (1929:288). Genus established as *Sternothaerus (Tanoa)* without a type species.
Etymology:* Anonym, *tanoa*, probably a fabricated name, an anagram of *anota* and near-anagram of *notoa*.

Notoa Gray 1863f:195
Type species: *Sternothaerus (Notoa) subniger* (Lacepède 1788) (*nomen suppressum*) = *Sternothaerus (Notoa) subniger* (Bonnaterre 1789)], by original monotypy.
Etymology:* Anonym, *notoa*, probably a fabricated name, a near-anagram of *tanoa* and *anota*, two anagrams.

Anota Gray 1863f:196 (junior homonym, not = *Anota* Hallowell 1852 [= Sauria])
Type species: *Sternothaerus (Anota) niger* (Duméril and Bibron 1835), by original monotypy.
Etymology:* Anonym, *anota*, probably a fabricated name, an anagram of *tanoa* and near-anagram of *notoa*.

Pelusios adansonii (Schweigger 1812)
Adanson's Mud Turtle




Roger Bour / CBFTT / Lac de Guiers, Senegal





left: Tomas Diagne / Lac de Guiers, Senegal / male
right: Roger Bour / CBFTT / Lac de Guiers, Senegal



(star = type locality in error)

Distribution: Benin (?), Cameroon, Central African Republic, Chad, Ethiopia, Mali, Mauritania, Niger, Nigeria, Senegal, South Sudan, Sudan, Uganda (?)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 1,475,150 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 5,350,089 sq. km

Size (Max SCL): male 20.0 cm, female 23.8 cm (Bour 2008 CBFTT)

CBFTT Account: Bour (2008)

IUCN Red List: Least Concern (LC) [Not Listed] (Behler 1996)

TFTSG Provisional Red List: Least Concern (LC) (2013)

Synonymy:

Emys adansonii Schweigger 1812:308
Hydraspis adansonii, Pelomedusa adansonii, Sternotherus adansonii, Sternothaerus adansonii, Pentonyx andansonii, Pentonyx adansonii, Pelusios adansonii, Pelusios adansonii adansonii
Type locality: "Nigritia" [Senegal]. Restricted to "cap Vert" [Senegal] [in error] [orange star] by Duméril and Bibron (1835:395).
Type specimen: MNHN 7972, holotype, see Bour (1986) and Ceríaco and Bour (2012).
Etymology:* Eponym honoring Michel Adanson (1727–1806), France / Senegal; botanist who worked in Senegal and collected the type.

Chelys (Hydraspis) adamsonii Gray 1830e:15 (*nomen novum*)
Etymology:* Eponym honoring Michel Adanson (1727–1806), France / Senegal; botanist who worked in Senegal and collected the type.
Comment: Unjustified emendation or error for *adansonii*.

Pelusios bechuanicus FitzSimons 1932
Okavango Mud Turtle




Marius Burger / Liuwa Plain National Park, Zambia





left: Uwe Fritz / Kavango R., Namibia
right: William R. Branch / Moremi National Park, Botswana



** (star = type locality)

Distribution: Angola, Botswana, Namibia, Zambia, Zimbabwe

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 520,519 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 820,172 sq. km

Size (Max SCL): male 30.0 cm, female 33.0 cm (Broadley 1981a; Ceballos et al. 2013)

IUCN Red List: Least Concern (LC) [Not Listed] (Behler 1996)

TFTSG Provisional Red List: Least Concern (LC) (2013)


Synonymy:

Pelusios bechuanicus FitzSimons 1932:37
Pelusios castaneus bechuanicus, Pelusios bechuanicus bechuanicus
Type locality: "Thamalakane River at Maun, Ngamiland" [Botswana] [yellow star].
Type specimen: TMP 14688, holotype.
Etymology:* Toponym for Bechuanaland (now = Botswana).


Pelusios broadleyi Bour 1986
Turkana Mud Turtle



Tomas Diagne / Lake Turkana, Kenya



left and center: Roger Bour / Loiyangalani, Lake Turkana, Kenya
right: Tomáš Mazuch / Loiyangalani, Lake Turkana, Kenya



(star = type locality)

Distribution: Ethiopia, Kenya
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 12,951 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 27,638 sq. km
Size (Max SCL): male 17.9 cm, female 14.9 cm (Prokop 2010)
IUCN Red List: Vulnerable (VU D2 [ver. 2.3]) (TFTSG 1996)
TFTSG Provisional Red List: Endangered (EN) (2013)
Synonymy:
Pelusios broadleyi Bour 1986:31
Type locality: "Loiengalani [= Loyengalanij] (2°43' N, 36°43' E), Marsabit district, Kenya" [yellow star].
Type specimen: CAS 123062, holotype.
Etymology:* Eponym honoring Donald G. Broadley (1932–2016), United Kingdom / Zimbabwe; herpetologist and turtle researcher who lived and worked in Zimbabwe.


Pelusios carinatus Laurent 1956 (17:102)
African Keeled Mud Turtle



Tomas Diagne / Congo (DRC) / captivity



Václav Gvoždík / Mai-Ndombe Prov., Congo (DRC)



(orange dot = uncertain identification; star = type locality)

Distribution: Congo (DRC), Congo (ROC), Gabon
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 191,523 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 263,151 sq. km
Size (Max SCL): 30.0 cm (Branch 2008; Itescu et al. 2014)
IUCN Red List: Least Concern (LC) [Not Listed] (Behler 1996)
TFTSG Provisional Red List: Least Concern (LC) (2013)
Synonymy:
Pelusios carinatus Laurent 1956:39
Type locality: "Eala, Equateur" [Democratic Republic of Congo (DRC)] [yellow star].
Type specimen: MRAC 2821, holotype.
Etymology:* Latin, *carina*, keel, *carinatus*, keeled; referring to the distinct carapacial keel.

Pelusios castaneus (Schweigger 1812) (14:47, 17:102, 21:35) (26)*
West African Mud Turtle




Gabriel H. Segniagbeto / CBFTT / Togoville, Togo





left: Luis Ceriaco / CBFTT / Cacuaco, nr. Luanda, Angola
right: Luca Luiselli / CBFTT / Uyo, Akwa Ibom State, Nigeria



(orange dots = introduced or trade; stars = type localities)

Distribution: Angola, Benin, Burkina Faso, Cameroon, Central African Republic, Chad, Congo (DRC), Congo (ROC), Gabon, Gambia, Ghana, Guinea, Guinea-Bissau, Ivory Coast (Côte d'Ivoire), Liberia, Niger, Nigeria, Senegal, Sierra Leone, Togo

Introduced: Guadeloupe, São Tomé and Príncipe (*probable historical introduction*), Seychelles (?)

Estimated Indigenous (Historical) Ranges (AOO and EOO): Est. Indigenous Area of Occupancy (AOO): 1,966,366 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 6,025,930 sq. km

Size (Max SCL): male 22.2 cm, female 28.5 cm (Segniagbeto et al. 2015; Bour et al. 2016 CBFTT)

CBFTT Accounts: Bour, Luiselli, Petrozzi, Segniagbeto, and Chirio (2016) on *Pelusios castaneus*; Bour and Gerlach (2008) on *Pelusios seychellensis*

IUCN Red List: Least Concern (LC) [Not Listed] (Behler 1996); the currently synonymized taxon *P. seychellensis* assessed as Extinct on the IUCN Red List (Gerlach 2003).

TFTSG Provisional Red List: Least Concern (LC) (2013)

Synonymy:


Emys castanea Schweigger 1812:314
Pelusios castaneus, Chelys (Sternotherus) castaneus, Chelys castaneus, Sternotherus castaneus, Clemmys (Pelusios) castanea, Clemmys castanea, Sternothaerus castaneus, Sternothaerus nigricans castaneus, Sternothaerus nigricans castanea, Pelusios nigricans castaneus, Pelusios subniger castaneus, Pelusios castaneus, Pelusios castaneus castaneus
Type locality: Not known. Restricted to "Afrique occidentale" by Bour (1979:148), and to "vicinity of Koutchatcha (7°20' N, 1°18' E)...close to the Amou River (ca. 30 km East of Gléï), Ogou Prefecture, Plateaux Region, Togo" [yellow star] by Bour (2008e:37) by neotype designation.
Type specimen: MNHN 8366, holotype, lost, see Bour (2008e); MNHN 2008.0303, neotype, designated by Bour (2008e:37).
Etymology:* Greek, *kastano*, brown; referring to the brownish chestnut-colored carapace.

Sternothaerus leachianus Bell 1825a:306
Sternotherus leachianus
Type locality: Not known.
Type specimen: OUM 8618, holotype, see Bour (2008e) and Nowak-Kemp and Fritz (2010).
Etymology:* Eponym honoring William Elford Leach (1791–1836), United Kingdom; zoologist and marine biologist at the British Museum of Natural History, London.

Sternotherus derbianus Gray 1844:37
Sternothaerus derbianus, Sternothaerus (Tanoa) derbianus, Pelusios derbianus, Pelusios castaneus derbianus
Type locality: "W. Africa, Sierra Leone ? Gambia." Restricted to "Gambia" [green star, location estimated] by Loveridge (1941:491).
Type specimen: NHMUK 1947.3.5.77 (formerly 1841.12.25.1), holotype, see Bour (2008e).
Etymology:* Eponym honoring Edward Smith-Stanley, 13th Earl of Derby (1775–1851), United Kingdom; nobleman and naturalist who obtained the type specimen.

Sternothaerus nigricans seychellensis Siebenrock 1906c:38 (07:103, 14:47, 17:103, 21:35)
Pelusios subniger seychellensis, Sternothaerus castaneus seychellensis, Pelusios castaneus seychellensis, Pelusios seychellensis
IUCN Red List: Extinct (EX) (Gerlach 2003), as *Pelusios seychellensis*. Previously: Vulnerable (VU) (TFTSG *in* Baillie and Groombridge 1996)
TFTSG Provisional Red List: Not Listed (NL) (2021)
Type locality:* "Seychellen" [Seychelles] [in error]. Restricted to "Seychellen, Insel Gloriosa?" [Seychelles] by Siebenrock (1909a:559), and to "Insel Mahé" [Seychelles] by Siebenrock (1909b:362).
Type specimen: NMW 13247, lectotype, designated as "type specimen" by Broadley (1981a:655) and as lectotype by Bour (1983:353), see also Tiedemann and Häupl (1980), Tiedemann et al. (1994), and Gemel et al. (2019); genotyped by Stuckas et al. (2013) and Kindler et al. (2016).
Etymology:* Toponym for the Seychelles Islands.
Comment:* Only 3 known specimens, unlikely to have been native to the Seychelles, most likely originating from coastal western Central Africa (see Stuckas et al. 2013; Kindler et al. 2016; and TTWG 2021, annotation 21:35).

Pelusios castanoides Hewitt 1931 (11:19, 12:42)
Yellow-bellied Mud Turtle
(includes 2 subspecies)



(subspecies: *castanoides* = red, *intergularis* = Seychelles [clustered orange dots])

Distribution: Kenya, Madagascar (*prehistorical introduction?*), Malawi, Mozambique, Seychelles (*prehistorical introduction?*), South Africa, Tanzania

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 937,734 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 5,050,209 sq. km

Size (Max SCL): male 19.0 cm, female 23.0 cm (see subspp.)

IUCN Red List: Least Concern (LC) (TFTSG 1996)

TFTSG Provisional Red List: Least Concern (LC) (2013)

Pelusios castanoides castanoides Hewitt 1931 (11:19, 12:42)
East African Yellow-bellied Mud Turtle




Miguel Vences / Lac Alaotra, Madagascar





Miguel Vences / Makira, Madagascar



(subspecies: *castanoides* = red; stars = type localities)

Distribution: Kenya, Madagascar (*prehistorical introduction?*), Malawi, Mozambique, South Africa, Tanzania

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 937,494 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 3,895,232 sq. km

Size (Max SCL): female 23.0 cm (Branch 2008; Itescu et al. 2014)

Synonymy:


Pelusios nigricans castanoides Hewitt 1931:463
Pelusios subniger castanoides, Pelusios castaneus castanoides, Pelusios castanoides, Pelusios castanoides castanoides
Type locality: "Richards Bay, Zululand" [South Africa] [orange star]. Restricted to "Lake St. Lucia estuary, KwaZulu" [South Africa] [yellow star] by Broadley (1981a:673).
Type specimen: TMP 13433, holotype.
Etymology:* Greek, *kastano*, brown, and *oides*, similarity; referring to the brownish color and similarity to *Pelusios castaneus*.

Pelusios castaneus kapika Bour 1979:149
Pelusios castanoides kapika
Type locality: "Delta du Sambirano (Province de Diégo-Suarez), nord de Madagascar" [green star].
Type specimen: MNHN 1929.238, holotype, see Broadley (1981a).
Etymology:* Vernacular (Madagascan, Sakalava), *kapika*; local indigenous name for the species.


Pelusios castanoides intergularis Bour 1983 (11:19, 12:42)
Seychelles Yellow-bellied Mud Turtle



Justin Gerlach / CBFTT / Seychelles



Justin Gerlach / CBFTT / Seychelles



(subspecies: *intergularis* = purple; star = type locality)

Distribution: Seychelles (*Cerf, Fregate, La Digue, Mahé, Praslin, Silhouette*) (*prehistorical introduction?*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 240 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 2,803 sq. km

Size (Max SCL): male 19.0 cm, female 23.0 cm [both estimated]; **(Max CCL):** male 19.4 cm, female 23.5 cm (Gerlach 2008a CBFTT)

CBFTT Account: Gerlach (2008a)

IUCN Red List: Critically Endangered (CR A2c, B2ab) (Gerlach 2003)

Synonymy:

Pelusios castanoides intergularis Bour 1983:355
Type locality: "La Digue Island, Seychelles" [yellow star].
Type specimen: NHMUK 1874.8.7.1, holotype.
Etymology:* Latin, *inter-*, between, and *gula*, throat; referring to the long intergular scute.


Pelusios chapini Laurent 1965 (11:19, 17:102)
Central African Mud Turtle



Jérôme Maran / Mandji, Gabon



Václav Gvoždík / Oyo–Mossaka region, Congo (ROC) / trade



(star = type locality)

Distribution: Cameroon (?), Central African Republic, Congo (DRC), Congo (ROC), Gabon, South Sudan, Uganda

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 974,361 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 2,039,549 sq. km

Size (Max SCL): 38.0 cm (Schmidt 1919; Laurent 1965; Maran 2006a; Itescu et al. 2014)

IUCN Red List: Least Concern (LC) [Not Listed] (Behler 1996)

TFTSG Provisional Red List: Least Concern (LC) (2013)


Synonymy:

Pelusios castaneus chapini Laurent 1965:21
Pelusios chapini
Type locality: "Kasenyi, Lake Albert, Bunia Terr., Ituri, Congo" [Democratic Republic of Congo (DRC)] [yellow star].
Type specimen: MRAC 20937, holotype.
Etymology:* Eponym honoring James P. Chapin (1889–1964), USA; ornithologist and curator at the American Museum of Natural History, New York, who collected in the Congo.

Pelusios cupulatta Bour and Maran 2003
Ivory Coast Mud Turtle




Jérôme Maran / San Pédro, Ivory Coast (Côte d'Ivoire)





left: Roger Bour / holotype / San Pédro, Ivory Coast (Côte d'Ivoire)
right: Tomas Diagne / Ivory Coast (Côte d'Ivoire)



(orange dots = uncertain or trade; star = type locality)

Distribution: Ghana, Guinea, Ivory Coast (Côte d'Ivoire), Liberia, Sierra Leone

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 274,784 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 390,381 sq. km

Size (Max SCL): male 31.3 cm, female 27.1 cm (L. Luiselli and F. Petrozzi, unpubl. data)

IUCN Red List: Not Evaluated (NE)

TFTSG Provisional Red List: Least Concern (LC) (2013)

Synonymy:

Pelusios cupulatta Bour and Maran 2003:28
Type locality: "environs de San Pédro, Côte d'Ivoire, précisément entre San Pédro (10 km W) et Grand-Bérébi (20 km E) (4°50' N, 6°47' W)." [Ivory Coast] [yellow star].
Type specimen: MNHN 2000.2605, holotype.
Etymology:* Vernacular (Corsican), *cupulatta*, turtle; also referring to *A Cupulatta*, the turtle-focused captive breeding organization on Corsica, France.

Pelusios gabonensis (Duméril 1856) [(17:102)]
African Forest Turtle




Jérôme Maran / Gabon





Václav Gvoždík / Mbomo, Congo (ROC)



**

(orange dots = possible trade or misidentified; star = type localities)

Distribution:* Angola, Cameroon, Central African Republic, Congo (DRC), Congo (ROC), Equatorial Guinea, Gabon, Ivory Coast (Côte d'Ivoire), Liberia, Zambia

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 2,289,713 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 3,573,376 sq. km

Size (Max SCL): male 32.0 cm, female 27.2 cm (Maran 2006c); maximum estimated: 33.0 cm (Branch 2008; Itescu et al. 2014)

IUCN Red List: Least Concern (LC) [Not Listed] (Behler 1996)

TFTSG Provisional Red List: Least Concern (LC) (2013)

Synonymy:

Pentonyx gabonensis Duméril 1856:373
Pelomedusa gabonensis, Sternothaerus gabonensis, Pelusios gabonensis
Type locality: "Gabon." Emended to "mouth of the Gabon river at present-day Libreville" by Bour 2008d:41 [yellow star].
Type specimen: MNHN 4237, holotype, discussed by Bour (2008d); MNHN 4137 listed by Iverson (1992), Bour and Maran (2003), and Uetz et al. (2019).
Etymology:* Toponym for Gabon.

Pentonyx gaboonensis Gray 1863f:194 (*nomen novum*)
Etymology:* Toponym for Gabon.
Comment: Unjustified emendation or error for *gabonensis*.


Pelomedusa gabonica Peters 1864:644 (*nomen novum*)
Etymology:* Toponym for Gabon.
Comment: Unjustified replacement name for *gabonensis*.

Sternothaerus steindachneri Siebenrock 1902a:6
Type locality: "Madagascar" [in error]. Restricted to "Gabon?" [yellow star, location estimated] by Tiedemann and Häupl (1980:8).
Type specimen: NMW 23392, holotype, see Tiedemann and Häupl (1980), Tiedemann et al. (1994), and Gemel et al. (2019).
Etymology:* Eponym honoring Franz Steindachner (1834–1919), Austria; ichthyologist and director at the Naturhistorisches Museum, Vienna.

Pelusios marani Bour 2000
Gabon Mud Turtle



Jérôme Maran / Fouganou, Gabon



** (star = type locality)

Distribution: Congo (ROC), Gabon

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 63,601 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 103,067 sq. km

Size (Max SCL): male 27.2 cm, female 22.9 cm (Bour 2000; Maran 2006a)

IUCN Red List: Not Evaluated (NE)

TFTSG Provisional Red List: Data Deficient (DD) (2018)

Synonymy:

Pelusios marani Bour 2000:3
Type locality: "Yombi (01º26' S, 10º37' E), province de N'Gounié, Gabon; environ 30 km SSE de Fougamou, entre Lambaréné et Mouïla" [yellow star].
Type specimen: MNHN 2000.2601, holotype.
Etymology:* Eponym honoring Jérôme Maran (1973–), France; herpetologist at *A Cupulatta* and collector of the type.


Pelusios nanus Laurent 1956 (17:102)
African Dwarf Mud Turtle



William R. Branch / northwest Zambia



left: Christo Deysel / Kalumbila, Zambia
right: Pearson McGovern / Congo (DRC) / captivity



** (star = type locality)

Distribution: Angola, Congo (DRC), Zambia

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 886,105 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 1,421,800 sq. km

Size (Max SCL): male 11.1 cm, female 12.2 cm (Laurent 1956; R. Bour, unpubl. data)

IUCN Red List: Least Concern (LC) [Not Listed] (Behler 1996)

TFTSG Provisional Red List: Data Deficient (DD) (2018)

Synonymy:

Pelusios nanus Laurent 1956:31
Pelusios adansonii nanus
Type locality: "Dilolo, Haut Lualaba" [Democratic Republic of Congo (DRC)] [yellow star].
Type specimen: MRAC 7833, holotype.
Etymology:* Greek, *nanos*, dwarf, small; referring to the small size of this species.

Pelusios niger (Duméril and Bibron 1835)
West African Black Mud Turtle




Gerald Kuchling / Kribi, Cameroon





left: Luca Luiselli / CBFTT / Port Harcourt, Nigeria
right: Jérôme Maran / CBFTT / Gamba, Gabon



(orange dots = uncertain native or trade or introduced; star = type locality)

Distribution: Benin, Cameroon, Equatorial Guinea, Gabon, Nigeria, Togo (?)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 209,728 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 599,699 sq. km

Size (Max SCL): male 35.5 cm, female 26.5 cm (Maran 2006a; Luiselli et al. 2018 CBFTT)

CBFTT Account: Luiselli, Bour, Petrozzi, Akani, and Segniagbeto (2018)

IUCN Red List: Near Threatened (NT A4cde) (Luiselli, Petrozzi, Akani, Segniagbeto, and Bour 2019). Previously: Least Concern (LC) [Not Listed] (Behler 1996)

Synonymy:

Sternotherus niger Duméril and Bibron 1835:397
Sternothaerus niger, Sternothaerus (Anota) niger, Pelusios niger
Type locality: "probablement...originaire de l'île de Madagascar" [in error]; restricted to "Région du Littoral, République du Cameroun" [Cameroon] [yellow star, location estimated] by Bour (2009d:37).
Type specimen: MNHN 8954, holotype, discussed by Bour (2009d).
Etymology:* Latin, *niger*, black, dark colored; referring to the blackish color of the carapace.

Sternothaerus oxyrhinus Boulenger 1897b:919
Type locality: "unknown...but probably...from some part of Tropical Africa."
Type specimen: NHMUK s/n, holotype, type specimen figured (pl.53), apparently lost.
Etymology:* Greek, *oxys*, pointed, sharp, and *rhinos*, nose; referring to the pointed snout.

Sternothaerus heinrothi Kanberg 1924:195
Type locality: "Kamerun" [Cameroon] [yellow star, location estimated].
Type specimen: ZMB 8232, holotype, see Fritz et al. (1994).
Etymology:* Eponym honoring Oskar Heinroth (1871–1945), Germany; biologist, ethologist, and director of the Berlin Aquarium who presented living specimens of the species.

Pelusios rhodesianus Hewitt 1927 (11:19, 17:102)
Variable Mud Turtle, Mashona Hinged Terrapin




Richard C. Boycott / CBFTT / nr. Lake St. Lucia, KwaZulu-Natal, South Africa





Richard C. Boycott / CBFTT / nr. Lake St. Lucia, KwaZulu-Natal, South Africa



(star = type locality)

Distribution: Angola, Botswana, Burundi, Congo (DRC), Congo (ROC), Malawi, Mozambique, Namibia, Rwanda, South Africa, Tanzania, Uganda, Zambia, Zimbabwe

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 2,250,980 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 5,777,726 sq. km

Size (Max SCL): 25.5 cm (Broadley and Boycott 2008 CBFTT)


CBFTT Account: Broadley and Boycott (2008)

IUCN Red List: Least Concern (LC) (TFTSG 1996)

Synonymy:

Pelusios nigricans rhodesianus Hewitt 1927:375
Pelusios subniger rhodesianus, Pelusios rhodesianus, Pelusios castaneus rhodesianus, Pelusios rhodesianus rhodesianus
Type locality: "Mpika district, N.E. Rhodesia" [Zambia] [yellow star].
Type specimen: PEM R14956 (formerly AMG 5432), holotype, see Broadley (1981a); PEM R14957, R14959, paratypes, genotyped by Kindler et al. (2016).
Etymology:* Toponym for Rhodesia (now = Zambia).

Pelusios sinuatus (Smith 1838) (11:19, 21:36)
Serrated Hinged Terrapin
(includes 2 subspecies)



(subspecies: *sinuatus* = red, *bottegi* = blue)

Distribution: Botswana, Burundi, Congo (DRC), Eswatini (Swaziland), Ethiopia, Kenya, Malawi, Mozambique, Rwanda, Somalia, South Africa, Tanzania, Zambia, Zimbabwe

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 2,570,192 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 5,283,476 sq. km

Size (Max SCL): male: 35.0 cm, female: 55.0 cm (see subspp.)

CBFTT Account: Broadley and Boycott (2009)

IUCN Red List: Least Concern (LC) [Not Listed] (Behler 1996)

TFTSG Provisional Red List: Least Concern (LC) (2013)


Pelusios sinuatus sinuatus (Smith 1838) (11:19, 21:36)
Southern Serrated Hinged Terrapin



Richard C. Boycott / CBFTT / Ndumu Game Reserve, KwaZulu-Natal, South Africa



left: Richard C. Boycott / CBFTT / Mpumalanga, KwaZulu-Natal, South Africa
right: Richard C. Boycott / CBFTT / Ndumu Game Reserve, KwaZulu-Natal, South Africa



(subspecies: *sinuatus* = red, *bottegi* = blue; stars = type localities)

Distribution: Botswana, Eswatini (Swaziland), Mozambique, South Africa, Zimbabwe (?)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 281,529 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 400,245 sq. km

Size (Max SCL): female: 34.7 cm (Broadley and Boycott 2009 CBFTT; Vamberger et al. 2019)

Synonymy:

Sternotherus sinuatus Smith 1838:Reptilia,pl.1
Sternothaerus (Tanoa) sinuatus, Sternothaerus sinuatus, Pelusios sinuatus, Pelusios sinuatus sinuatus
Type locality: "rivers to the north of 25° south latitude" [South Africa]. Restricted to "the Crocodile-Marico confluence, northern Transvaal" [South Africa] [yellow star] by Broadley (1981a:654).
Type specimen: NMSZ 1859.13.1684 (formerly RSM), lectotype, designated as the type by FitzSimons (1937:261,pl.X) and Broadley (1981a:654), listed as "syntype" by Herman et al. (1990); NMSZ 1859.13.1686, paralectotype, listed as "?syntype" by Herman et al. (1990).
Etymology:* Latin, *sinus*, curvature, bend, *sinuatus*, indented; referring to the serrated rear edge of the carapace.

Sternotherus dentatus Peters 1848:494 (*nomen nudum*)
Sternothaerus dentatus
Etymology:* Latin, *dens*, tooth, *dentatus*, toothed, serrated; referring to the serrated rear edge of the carapace.

Pelusios sinuatus zuluensis Hewitt 1927:371
Type locality: "near the Umsinene River, Zululand" [Mzinene River, KwaZulu-Natal, South Africa] [green star, location estimated].
Type specimen: NMB 609, holotype, see Broadley (1981a).
Etymology:* Toponym for Zululand, South Africa.


Pelusios sinuatus bottegi (Boulenger 1895a) (11:19, 21:36)
Northern Serrated Hinged Terrapin



Ronald Orenstein / Nairobi National Park, Kenya



H. Bradley Shaffer / nr. Dar es Salaam, Tanzania



(subspecies: *sinuatus* = red, *bottegi* = blue; stars = type localities)

Distribution: Botswana, Burundi, Congo (DRC), Ethiopia, Kenya, Malawi, Mozambique, Rwanda, Somalia, Tanzania, Zambia, Zimbabwe

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 2,288,663 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 4,799,189 sq. km

Size (Max SCL): male: 35.0 cm, female: 55.0 cm (Spawls et al. 2002; Broadley and Boycott 2009 CBFTT; Ceballos et al. 2013)


Synonymy:

Sternothaerus bottegi Boulenger 1895a:9
Pelusios sinuatus bottegi
Type locality: "Bardera...Giuba e dei suoi affluenti" [Bardera, Jubba River, Italian Somaliland] [Somalia] [yellow star].
Type specimen: MSNG CE2319, holotype, see Capocaccia (1961).
Etymology:* Eponym honoring Vittorio Bottego (1860–1897), Italy; military officer and explorer who collected the type.

Pelusios sinuatus leptus Hewitt 1933a:45
Type locality: "Isoka, North-East Rhodesia" [Zambia] [green star].
Type specimen: PEM R 14960 (formerly AMG s/n), holotype, see Broadley (1981a).
Etymology:* Greek, *leptos*, slender, thin; referring to the narrow vertebral scutes.

Sternothaerus rudolphi † Arambourg 1947:461
Pelusios rudolphi
Type locality: "basin du Lac Rodolphe et de la basse vallée de l'Omo" [Ethiopia] [black star]. Emended to "Shungura Formation...Omo River Basin...Ethiopia" by Lapparent de Broin (2000b:59).
Type specimen: MNHN P1933-9-449, holotype, fossil, incomplete plastron; P1933-9-448, paratype, incomplete carapace; figured (f.79-80, pl.34,f.4, pl.35,f.3) and in Broin (1969:pl.25,f.4).
Geological age: Late Pliocene–Early Pleistocene, Villafranchian, Shungura Formation, ca. 3.79–1.6 million ybp, see Broin (1969) and Lapparent de Broin (2000b).
Etymology:* Eponym honoring Crown Prince Rudolf (1858–1889), Austria; royalty for whom Lake Rudolf (now Lake Turkana) was also named.

Pelusios subniger (Bonnaterre 1789) (12:43)
East African Black Mud Turtle
(includes 2 subspecies)



(subspecies: *subniger* = red, *parietalis* = Seychelles [clustered yellow dots]; orange dots = introduced)

Distribution: Botswana, Burundi, Congo (DRC), Madagascar (*prehistorical introduction?*), Malawi, Mozambique, Seychelles (*prehistorical introduction?*), South Africa, Tanzania, Zambia, Zimbabwe

Introduced: British Indian Ocean Territory (*Diego Garcia, Chagos Archipelago*), French Southern Territories (*Glorieuse*), Mauritius

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 1,664,490 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 6,523,416 sq. km

Size (Max SCL): male 15.0 cm, female 20.0 cm (see subspp.)


IUCN Red List: Least Concern (LC) (TFTSG 1996)

TFTSG Provisional Red List: Least Concern (LC) (2013)


Pelusios subniger subniger (Bonnaterre 1789) [12:43]
East African Black Mud Turtle



James Harvey / northeast South Africa



left: James Harvey / northeast South Africa
right: Sebastian Gehring / Manompana, Madagascar



(subspecies: *subniger* = red; orange dot = introduced; star = type locality)

Distribution: Botswana, Burundi, Congo (DRC), Madagascar (*prehistorical introduction?*), Malawi, Mozambique, South Africa, Tanzania, Zambia, Zimbabwe

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 1,664,250 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 6,503,796 sq. km

Size (Max SCL): 20.0 cm (Branch 2008; Itescu et al. 2014)

Synonymy:

Testudo subnigra Lacepède 1788:175, synopsis[table] (*nomen suppressum*) [09:6]
Type locality: Not known. Restricted to "Tamatave, est de Madagascar" [= Toamasina] [yellow star] by Bour (1979:152).
Type specimen: MNHN 8366, holotype, type specimen figured (pl. opp.175), discussed by Bour (2004b).
Etymology:* Latin, *sub*, almost, below, and *nigra*, black, dark colored; referring to the blackish carapace coloring.
Comment: Name suppressed by ICZN (2005a) as published in a rejected and invalid non-binomial work, see Savage (2003).

Testudo subnigra Bonnaterre 1789:30 (*nomen protectum*)
Emys subnigra, Pelusios subniger, Sternotherus subniger, Clemmys (Pelusios) subnigra, Clemmys subnigra, Sternothaerus subniger, Sternothaerus (Notoa) subniger, Pelusios subniger, Pelusios subniger subniger
Type locality: Not known. Restricted to "Tamatave, est de Madagascar" [= Toamasina] [yellow star] by Bour (1979:152).
Type specimen: MNHN 8366, holotype, type specimen figured in Lacepède (1788:pl.opp.175), discussed by Bour (2004b).
Etymology:* Latin, *sub*, almost, below, and *nigra*, black, dark colored; referring to the blackish carapace coloring.
Comment: Name conserved by ICZN (1989), see Smith et al. (1980b).

Testudo nigricans Donndorff 1798:34 (*nomen novum*)
Terrapene nigricans, Kinosternon nigricans, Sternotherus nigricans, Sternothaerus nigricans, Sternothaerus nigricans nigricans, Pelusios nigricans, Pelusios nigricans nigricans
Comment: Unjustified replacement name for *subnigra*.
Etymology:* Latin, *nigricans*, to become black, blackish; referring to the blackish carapace coloring.


Pelusios subniger parietalis Bour 1983 [12:43]
Seychelles Black Mud Turtle



Justin Gerlach / CBFTT / Seychelles



Justin Gerlach / CBFTT / Seychelles



(subspecies: *parietalis* = purple; star = type locality)

Distribution: Seychelles (*Cerf, Cousin [extirpated], Fregate, La Digue, Mahé, Praslin, Silhouette, St. Anne [extirpated]*) (*prehistorical introduction?*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 240 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 2,789 sq. km

Size (Max SCL): male 15.0, female 19.0 cm [both estimated]; **(Max CCL):** male 15.8 cm, female 20.0 cm (Gerlach 2008b CBFTT)

CBFTT Account: Gerlach (2008b)

IUCN Red List: Critically Endangered (CR A2c, B2ab(ii,iii)) (Gerlach 2003)

Synonymy:

Pelusios subniger parietalis Bour 1983:359


Type locality: "La Digue Island, Seychelles" [yellow star].
Type specimen: USNM 19802, holotype, see Reynolds et al. (2007).
Etymology:* Latin, *parietalis*, walls, sides; referring to the parietal scales with numerous furrows on the head.

Pelusios upembae Broadley 1981a
Upemba Mud Turtle


Tomas Diagne / Congo (DRC) / captivity




Tomas Diagne / Congo (DRC) / captivity


(star = type locality)

Distribution: Congo (DRC)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 18,931 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 27,931 sq. km
Size (Max SCL): 23.0 cm (Branch 2008; Itescu et al. 2014)
IUCN Red List: Data Deficient (DD) (TFTSG 1996)
TFTSG Provisional Red List: Data Deficient (DD) (2018)
Synonymy:
Pelusios bechuanicus upembae Broadley 1981a:667
Pelusios upembae
Type locality: "Kanonga River, tributary of the right bank of the Fungwe River (695 m), Upemba National Park, Shaba Province, Zaire" [Democratic Republic of Congo (DRC)] [yellow star].
Type specimen: TMP 38178, holotype.
Etymology:* Toponym for the Upemba Depression, Democratic Republic of Congo (DRC).

Pelusios williamsi Laurent 1965 [(21:37)]
Williams' Mud Turtle
(includes 3 subspecies)


(subspecies: *williamsi* = red, *laurenti* = blue, *lutescens* = green; orange dots = probable trade or questionable)


Distribution: Congo (DRC), Kenya, Rwanda, Tanzania, Uganda
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 79,575 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 315,368 sq. km
Size (Max SCL): male 19.7 cm, female 23.8 cm (see subspp.)
IUCN Red List: Least Concern (LC) [Not Listed] (Behler 1996)
TFTSG Provisional Red List: Least Concern (LC) (2013)

Pelusios williamsi williamsi Laurent 1965
Lake Victoria Mud Turtle


Jérôme Maran / Lake Victoria, Uganda / captivity



left: Harald Artner / Lake Victoria, Uganda / captivity
right: Jérôme Maran / Lake Victoria, Uganda / captivity


(subspecies: *williamsi* = red, *laurenti* = blue, *lutescens* = green; star = type locality)


Distribution: Kenya, Tanzania, Uganda
Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO): 54,068 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 159,275 sq. km
Size (Max SCL): male 19.7 cm, female 23.8 cm (Bour 1984a)
Synonymy:
Pelusios williamsi Laurent 1965:12
Pelusios williamsi williamsi, Pelusios castaneus williamsi, Pelusios williamsii
Type locality: "Kakamega, Kaimosi, Kenya" [yellow star].
Type specimen: MCZ 40021, holotype.
Etymology:* Eponym honoring Ernest E. Williams (1914–1998), USA; herpetologist and curator at the Museum of Comparative Zoology, Harvard University.

Pelusios williamsi laurenti Bour 1984a
Ukerewe Island Mud Turtle



Roger Bour / Ukerewe Island, Lake Victoria, Tanzania
Left: paratype, MCZ 30017, female; *Right*: holotype, MCZ 30016, male from Bour (1984a:Fig. 1)




(subspecies: *williamsi* = red, *laurenti* = blue; star = type locality)
Distribution: Tanzania
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 551 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 732 sq. km
Size (Max SCL): male 18.7 cm, female 22.6 cm (Bour 1984a)
Synonymy:
Pelusios williamsi laurenti Bour 1984a:29
Type locality: "Ukerewe Island (Lake Victoria), Tanzania, altitude 1150 m" [yellow star].
Type specimen: MCZ 30016, holotype.
Etymology:* Eponym honoring Raymond F. Laurent (1917–2005), Belgium; herpetologist who researched African turtles of the genus *Pelusios*.

Pelusios williamsi lutescens Laurent 1965
Albert Nile Mud Turtle



Nik Borrow / iNaturalist.org / Kanungu, Uganda



(subspecies: *williamsi* = red, *laurenti* = blue, *lutescens* = green; orange dots = probable trade or questionable; star = type locality)
Distribution: Congo (DRC), Rwanda, Uganda
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 24,956 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 69,011 sq. km
Size (Max SCL): male 17.8 cm, female 22.4 cm (Bour 1984a)
Synonymy:
Pelusios williamsi lutescens Laurent 1965:16
Pelusios castaneus lutescens
Type locality: "Semliki River, 1 km below the Lake Edward" [Democratic Republic of Congo (DRC)] [yellow star].
Type specimen: IRSNB 6822, holotype.
Etymology:* Latin, *lutescens*, yellowish, golden yellowish; referring to the predominantly yellowish colored plastron.

Family Podocnemididae Cope 1869 (07:104, 09:48, 11:20, 21:3, 21:6, 21:9, 21:38)

Podocnemid Side-necked Turtles
(includes 3 subfamilies, 3 genera, 8 species, 8 taxa)
Hydraspidina Bonaparte 1836:3 (*partim*)
Podocnemididae Cope 1869:282
Peltocephalidae Gray 1870c:70,82
Podocneminae Zangerl 1947:39
Podocnemidae Meylan 1996:31



Podocnemididae Species Richness

Subfamily Erymnochelyinae Broin 1988 (21:3, 21:9)

Madagascan Big-headed Turtles
(includes 1 genus, 1 species, 1 taxon)
Erymnochelyinae Broin 1988:105,137

Erymnochelys Baur 1888a

Madagascan Big-headed Turtles
(includes 1 species, 1 taxon)
Dumerilia Grandidier 1867:232 (junior homonym, not = *Dumerilia* Leach 1824 [= Coleoptera] or *Dumerilia* Robineau-Desvoidy 1835 [= Diptera] or *Dumerilia* Bocage 1866 [= Sauria])
Type species: *Dumerilia madagascariensis* Grandidier 1867, by original monotypy.
Etymology:* Eponym honoring André Marie Constant Duméril (1774–1860), France; naturalist, herpetologist, and turtle researcher at the Muséum National d'Histoire Naturelle, Paris.
Erymnochelys Baur 1888a:421 (*nomen novum*)
Type species: *Erymnochelys madagascariensis* [= *Dumerilia madagascariensis* Grandidier 1867], by application of ICZN Article 67.8.
Etymology:* Greek, *erymnos*, defensive, reinforced, strong, and *chelys*, turtle; referring to assumed defensive behavior based on the size of the jaws.
Comment:* Justified replacement name for *Dumerilia*.

Erymnochelys madagascariensis (Grandidier 1867)


Madagascan Big-headed Turtle, *Rere*



Anders G.J. Rhodin / TCC / CRM 6 / Madagascar / captivity / Ampijoroa



left and center: Anders G.J. Rhodin / Baly Bay N.P., Madagascar / juvenile
right: Juliette Velosoa / Madagascar / captivity / Ampijoroa



(stars = type localities)

Distribution: Madagascar
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 151,139 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 321,408 sq. km
Size (Max SCL): male 45.8 cm, female 43.0 cm (Pedrono and Smith 2013)
IUCN Red List: Critically Endangered (CR A4d) (Leuteritz, Kuchling, Garcia, and Velosoa 2008). Previously: Endangered (EN) (TFTSG *in* Baillie and Groombridge 1996)
CITES: Appendix II (1975)
Synonymy:
Dumerilia madagascariensis Grandidier 1867:232
Podocnemis madagascariensis, Erymnochelys madagascariensis, Dumerilia madagascarensis, Erymnochelys madagascarensis, Podocnemis madagascarensis
Type locality: "Mouroundava Tsidsibonque flumina in occidentali insulae Madagascar littore" [Morondava and Tsiribihina rivers on the western coast of Madagascar Island]. Restricted to "Morondava River...Madagascar" [yellow star] by Bour (2006a:37).
Type specimen: MNHN 9544, holotype, see Bour (2006a).
Etymology:* Toponym for Madagascar.
Podocnemis madagascariensis bifilaris Boettger 1893:14
Type locality: "Mojanga, West-Madagaskar" [Mahajanga, Madagascar] [green star].
Type specimen: SMF 7982, holotype, see Mertens (1967b).

Etymology:* Latin, *bi-*, two, and *filaris*, threadlike; referring to the two chin barbels.

Subfamily Peltocephalinae Gray 1870f (21:3, 21:9)

South American Big-headed Turtles

(includes 1 genus, 1 species, 1 taxon)

Peltocephalina Gray 1870f:719

Peltocephalinae Joyce, Anquetin, Cadena, Claude, Danilov, Evers, Ferreira, Gentry, Georgalis, Lyson, Pérez-García, Rabi, Sterli, Vitek, and Parham 2021:21

Peltocephalus Duméril and Bibron 1835

South American Big-headed Turtles

(includes 1 species, 1 taxon)

Peltocephalus Duméril and Bibron 1835:377

Type species: *Peltocephalus tracaxa* [= *Emys tracaxa* Spix 1824] [= subjective synonym of *Emys dumeriliana* Schweigger 1812], by original monotypy.

Etymology:* Greek, *pelta*, small, crescent-shaped leather shield, and *kefale*, head; referring to the head scaling reminiscent of a helmet.

Chelonemys Gray 1864d:134 (junior homonym, not = *Chelonemys* Jourdan 1862 [= fossil sea turtles, Eurysternidae])

Type species: *Chelonemys dumeriliana* sensu Gray 1864d (not *Emys dumeriliana* Schweigger 1812) [= subjective synonym of *Podocnemis unifilis* Troschel 1848], by original monotypy.

Etymology:* Greek, *chelone*, turtle, and *emys*, turtle.

Comment:* Frétey (2024:17) designated the generic name *Chelonemys* (not its designated type species) to be a synonym of *Peltocephalus* instead of *Podocnemis*, under which generic name it had traditionally been synonymized.

Peltocephalus dumerilianus (Schweigger 1812)

South American Big-headed Turtle, *Cabeçudo*, *Cabezón*



Richard C. Vogt / Rio Negro, Amazonas, Brazil



Russell A. Mittermeier / Rio Negro, Amazonas, Brazil / adult, juvenile



** (orange dots = probable trade; stars = type localities)

Distribution: Brazil (*Amapá*, *Amazonas*, *Pará*, *Roraima*), Colombia (*Amazonas*, *Caquetá*, *Guainía*, *Guaviare*, *Meta*, *Putumayo*, *Vaupés*, *Vichada*), Ecuador, French Guiana, Peru (*Loreto*), Venezuela (*Amazonas*, *Apure*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 978,450 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 3,692,687 sq. km

Size (Max SCL): male 50.0 cm, female 47.0 cm (Iverson and Vogt 2002; Ceballos et al. 2013)

IUCN Red List:* Vulnerable (VU A2cd+4cd) (Gentil, de Medeiros, Morales-Betancourt, Maffei, and Da Silveira, in press). Previously: TFTSG Provisional Red List: Vulnerable (VU) (2011); Vulnerable (VU A1acd [ver. 2.3]) (TFTSG 1996)

CITES: Appendix II (1975)

Synonymy:

Emys dumeriliana Schweigger 1812:300

Podocnemis dumeriliana, Hydraspis dumeriliana, Peltocephalus dumerilianus, Chelonemys dumeriliana, Peltocephalus dumeriliana

Type locality: "America meridionali." Restricted to "French Guiana" by Bour (2006a:29) [yellow star, location estimated].

Type specimen: Original holotype apparently lost, possibly MNHN 7893, its identification as either *Podocnemis unifilis* or *P. expansa* discussed by Williams (1954), Hoogmoed and Gruber (1983), Pritchard and Trebbau (1984), and Bour (2006a); MNHN 8364, neotype, designated by Bour (2006a:29).

Etymology:* Eponym honoring André Marie Constant Duméril (1774–1860), France; naturalist, herpetologist, and turtle researcher at the Muséum National d'Histoire Naturelle, Paris.

Emys macrocephala Spix 1824:5 (senior homonym, not = *Emys macrocephalus* Gray 1844 [= *Malaclemys terrapin terrapin*])

Peltocephalus macrocephala

Type locality: "Airon ad ripam fluminis Yau, confluentis Rio Negro" [Airão, Rio Jáu, Amazonas, Brazil] [green star].

Type specimen: RMNH 6164, lectotype, designated by Hoogmoed and Gruber (1983:347), see Pritchard and Trebbau (1984) and Hoogmoed et al. (2010).

Etymology:* Greek, *makros*, large, long, and *kefale*, head; referring to the large, broad head.

Emys tracaxa Spix 1824:6

Hydraspis tracaxa, Podocnemis tracaxa, Peltocephalus tracaxa, Peltocephalus tracaxus

Type locality: "fluminis Solimoëns" [Rio Solimões, Amazonas, Brazil] [= between Tefé and Tabatinga, possibly near São Paulo de Olivença] [blue star].

Type specimen: ZSM 16/0, lectotype, designated by Hoogmoed and Gruber (1983:347), see Pritchard and Trebbau (1984) and Franzen and Glaw (2007).

Etymology:* Vernacular (Native Amazonian), *tracaja*, local indigenous name for the species.

Chelys (Hydraspis) dumerilliana Gray 1830e:17 (*nomen novum*)

Etymology:* Eponym honoring André Marie Constant Duméril

(1774–1860), France; naturalist, herpetologist, and turtle researcher at the Muséum National d'Histoire Naturelle, Paris.
Comment: Unjustified emendation or error for *dumeriliana*.

Emys icterocephala Spix *in* Gray 1830e:17 (*nomen nudum*)
Comment: Unjustified replacement name or error for *macrocephala*.

Peltocephalus tracaya Troschel 1848:646 (*nomen novum*)
Etymology:* Vernacular (Native Amazonian), *tracaja*, local indigenous name for the species.
Comment: Unjustified replacement name for *tracaxa*.

Subfamily Podocnemidinae Cope 1869 (21:3, 21:9) (27)*

South American River Turtles
(includes 1 genus, 6 species, 6 taxa)
Podocnemididae Cope 1869:282
Podocnemina Gray 1870f:719
Podocnemidina Gray 1873j:73
Podocneminae Zangerl 1947:39
Podocnemidinae Bour and Dubois 1984:80

Podocnemis Wagler 1830b (21:38)

South American River Turtles
(includes 6 species, 6 taxa)
Podocnemis Wagler 1830b:135
Type species: *Podocnemis expansa* [= *Emys expansa* Schweigger 1812], by subsequent designation by Fitzinger (1843:29).
Etymology:* Greek, *podos*, leg, and *knemis*, shin, lower leg; referring to the widened scales on the lower legs.
Bartlettia Gray 1870f:720 (junior homonym, not = *Bartlettia* Adams 1867 [= Mollusca])
Type species: *Bartlettia pitipii* Gray 1870 [= subjective synonym of *Podocnemis sextuberculata* Cornalia 1849], by original monotypy.
Etymology:* Eponym honoring Edward Bartlett (1836–1908), United Kingdom / Malaysia; ornithologist and herpetologist at the British Museum of Natural History, London, and at the Sarawak Museum in Malaysia, who collected the type species.

Podocnemis erythrocephala (Spix 1824) (10:46, 10:48, 14:50, 17:104, 21:39, 21:41)


Red-headed Amazon River Turtle, *Irapuca*, *Chipiro*



Richard C. Vogt / CBFTT / Rio Jufari, Amazonas, Brazil / male



left: Virginia C.D. Bernardes / Amazonas, Brazil
right: Russell A. Mittermeier / CBFTT / Amazonas, Brazil / female



** (orange dots = questionable or probable trade, stars = type localities)

Distribution: Brazil (*Amazonas*, *Pará*, *Roraima*), Colombia (*Guainía*, *Vichada*), Venezuela (*Amazonas*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 432,578 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 1,246,551 sq. km

Size (Max SCL): male 24.4 cm, female 32.2 cm (Mittermeier et al. 2015 CBFTT)

CBFTT Account: Mittermeier, Vogt, Bernhard, and Ferrara (2015)

IUCN Red List:* Vulnerable (VU A2cd+4cd) (Andrade, Rojas-Runjaic, Ferrara, Bernhard, and Vogt, in press). Previously: TFTSG Provisional Red List: Vulnerable (VU) (2011); Vulnerable (VU) (TFTSG 1996)

CITES: Appendix II, as *Podocnemis* spp. (1975)

Synonymy:

Emys cayennensis Schweigger 1812:298 (*partim*, misidentified type [= *Podocnemis unifilis*]) (10:48, 14:50, 21:41)
Chelys (Hydraspis) cayennensis, *Chelys cayennensis*, *Hydraspis cayennensis*, *Podocnemis cayennensis*
Type locality: "Cayenna" [Cayenne, French Guiana] [orange star].
Type specimen: MNHN 8359, lectotype, designated by Bour (2006a:33), discussed by Pritchard and Trebbau (1984), Ceríaco and Bour (2012), and Vogt et al. (2013); MNHN 4152, erroneously designated lectotype by Hoogmoed and Gruber (1983:343), is a *P. expansa* (Bour 2006:32).
Etymology:* Toponym for Cayenne, French Guiana.
Comment: A proposal to the ICZN by Vogt et al. (2013) to give *Podocnemis unifilis* Troschel 1848 precedence over *Emys cayennensis* Schweigger 1812 whenever the two are considered synonyms, was approved by ICZN (2019). Name synonymized *in partim* under *Podocnemis erythrocephala* and *Podocnemis unifilis*.

Emys erythrocephala Spix 1824:9
Podocnemis erythrocephala, *Hydraspis expansa erythrocephala*, *Chelys (Hydraspis) erythrocephala*, *Chelys erythrocephala*
Type locality: "aquis ripariis fluminis Solimoens" [Rio Solimões, Amazonas, Brazil] [= between Tefé and Tabatinga] [yellow star, location estimated].
Type specimen: ZSM 2517/0, holotype, see Hoogmoed and Gruber (1983) and Franzen and Glaw (2007).
Etymology:* Greek, *erythros*, reddish, and *kephale*, head; referring to the red color of the head.

Emys bitentaculata Cuvier *in* Gray 1830e:17 (*nomen nudum et dubium*) (14:48)
Etymology:* Latin, *bi*, twice, and *tentaculum*, thread, small tentacle; referring to the two chin barbels.

Hydraspis bitentaculata Gray 1831d:42 (*nomen oblitum et dubium*) (14:48)
Type locality: "Brasilia" [Brazil].
Type specimen: MNHN s/n, holotype, apparently lost.
Etymology:* Latin, *bi*, twice, and *tentaculum*, thread, small tentacle; referring to the two chin barbels.

Podocnemis agassizii Coutinho *in* Göldi 1886:277 (10:46)
Type locality: "Rio Negro" [Amazonas, Brazil] [green star, location

estimated].
Type specimen: Not located, type specimen figured (pl.5).
Etymology:* Eponym honoring Louis Agassiz (1807–1873), Switzerland / USA; herpetologist at Harvard University who founded the Museum of Comparative Zoology.

Podocnemis coutinhii Göldi 1886:279 (*nomen novum*)
Type locality: "Rio Negro" [Amazonas, Brazil] [green star, location estimated].
Etymology:* Eponym honoring João Martins da Silva Coutinho (1830–1889), Brazil; military engineer and participant on the Thayer Amazon Expedition with Louis Agassiz.
Comment: Unjustified replacement name for *agassizii*.

Podocnemis expansa (Schweigger 1812)
Giant South American River Turtle, Giant Amazon River Turtle, *Tartaruga-da-Amazônia, Charapa, Arrau*


Richard C. Vogt / Rio Trombetas, Pará, Brazil / female


left: Camila Ferrara / Rio Purus, Amazonas, Brazil
right: Russell A. Mittermeier / Amazonas, Brazil / juvenile


(orange dots = trade or introduced and waifs; stars = type localities)

Distribution:* Bolivia (*Beni, La Paz, Pando, Santa Cruz*), Brazil (*Acre, Amapá, Amazonas, Goiás, Maranhão, Mato Grosso, Pará, Rondônia, Roraima, Tocantins*), Colombia (*Amazonas, Arauca, Caquetá, Casanare, Guainía, Meta, Putumayo, Vaupés, Vichada*), Ecuador, French Guiana, Guyana, Peru (*Loreto, Ucayali*), Venezuela (*Amazonas, Anzoátegui, Apure, Bolívar, Delta Amacuro, Guárico, Monagas*)
Introduced:* Trinidad (*waifs*), Venezuela (*Aragua, Carabobo*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 1,706,101 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 7,126,911 sq. km
Size (Max SCL): male 55.0 cm, female 109.0 cm (Ceballos et al. 2013; Ferrara et al. 2017)
IUCN Red List:* Endangered (EN A2cd+4cd) (Ferrara, Fagundes, Forero-Medina, Brito, Miorando, Lacava, Gomes Lustosa, Hernández, Cueva, Millar, Dewynter, Páez, Carvajal, and Domínguez, in press). Previously: TFTSG Provisional Red List: Critically Endangered (CR) (2011); Lower Risk/conservation dependent (LR/cd) (TFTSG 1996)
CITES: Appendix II, as *Podocnemis* spp. (1975)
USA ESA:* Endangered (USFWS 1970)
Synonymy:
Emys expansa Schweigger 1812:299
Podocnemis expansa, Chelys (Hydraspis) expansa, Chelys expansa, Hydraspis expansa
Type locality: "America meridionali." Restricted to "French Guiana" by Bour (2006a:35) by lectotype designation [yellow star, likely the Oyapock River].
Type specimen: MNHN 7997, lectotype, designated by Bour (2006a:35), see also Ceríaco and Bour (2012).
Etymology:* Latin, *expansa*, expanded; referring to the strongly expanded posterior carapacial margin.
Testudo arrau Humboldt 1819a:243 (14:49)
Emys arrau
Type locality: "entre le confluent de l'Orénoque avec l'Apure et les grandes cataractes ou *Raudales*...entre Cabruta et la mission d'Aturès...les trois pêches de l'Encaramada ou Boca del Cabullare, de Cucuruparu ou Boca de la Tortuga, et de Pararuma, un peu au-dessous de Carichana" [Venezuela]. Restricted to "zwischen dem Zusammenfluss des Apure mit dem Orinoko und den grossen Wasserfällen" [Venezuela] [green star, location estimated] by Wermuth and Mertens (1961:296).
Type specimen: Not known or located.
Etymology:* Vernacular (Amazonian Native, Arawakan or Tupi Guarani), *araú*, local indigenous name for the species.
Emys amazonica Spix 1824:1
Type locality:* "fluvio Solimoens et confluentibus Javary, Rio Branco" [Rio Solimões at the confluence of Rio Javary and Rio Branco, Amazonas, Brazil] [blue star].
Type specimen: ZSM 2446/0/1, lectotype, designated by Hoogmoed and Gruber (1983:343), discussed by Franzen and Glaw (2007) and Hoogmoed et al. (2010).
Etymology:* Toponym for the Amazon River basin, Brazil.

Podocnemis lewyana Duméril 1852 (12:44)
Magdalena River Turtle, *Tortuga del río Magdalena*




Alejandra Cadavid / CBFTT / TCC / Río Chicagua, Bolivar, Colombia





left: Alejandra Cadavid / CBFTT / TCC / Río Chicagua, Bolivar, Colombia / male
right: Vivian P. Páez / CBFTT / Colombia / female



** (orange dots = probable trade, translocated, or introduced; stars = type localities)

Distribution: Colombia (*Antioquia, Atlántico, Bolívar, Boyacá, Caldas, Cesar, Córdoba, Cundinamarca, La Guajira, Magdalena, Santander, Sucre, Tolima*)

Introduced: Venezuela (*Zulia*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 66,926 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 188,057 sq. km

Size (Max SCL): male 42.4 cm, female 50.0 cm (Gallego-García and Castaño-Mora 2008; Ceballos et al. 2013; Vergara-Ríos et al. 2015)

CBFTT Account: Páez, Restrepo, Vargas-Ramírez, and Bock (2009)

IUCN Red List: Critically Endangered (CR A2acd+4acd) (Páez, Gallego-Garcia, and Restrepo 2016). Previously: Endangered (EN) (TFTSG 1996)

CITES: Appendix II, as *Podocnemis* spp. (1975)

Synonymy:

Podocnemis lewyana Duméril 1852:242
Type locality: "Santa Fé de Bogota...et...la République de Venezuella" [Colombia and Venezuela]. Restricted to "Bogotá, Colombia" [erroneous locality] [orange star] by Williams (1954:281); further restricted here to "Río Magdalena, ca. 75 km west of Bogotá, Colombia" [yellow star] (see Medem 1969).
Type specimen: MNHN 8985, lectotype, designated "holotype" by Williams (1954:281), discussed by Bour (2006a).
Etymology:* Eponym honoring Bernard Lewy (1817–1863), Denmark / France / Colombia; chemist and naturalist who worked in Bogotá, Colombia, and obtained the type.

Podocnemis sextuberculata Cornalia 1849 (10:47, 17:105, 21:40)
Six-tubercled Amazon River Turtle, *Iaçá, Pitiú, Cupiso*



Richard C. Vogt / Paraná Floresta, Roraima, Brazil / female





Russell A. Mittermeier / Rio Negro, Amazonas, Brazil / hatchling, female



** (orange dots = probable trade; stars = type localities)

Distribution: Bolivia (*Pando*), Brazil (*Acre, Amazonas, Pará, Rondônia, Roraima*), Colombia (*Amazonas, Caquetá, Putomayo, Vaupés*), Ecuador (?), Peru (*Loreto, Ucayali*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 614,881 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 2,874,370 sq. km

Size (Max SCL): male 25.0 cm, female 34.0 cm (Fachín-Terán et al. 2003; Ferrara et al. 2017; Iverson et al. 2017)

IUCN Red List:* Endangered (EN A2ad+4cd) (Miorando, Lacava, Ferrara, Andrade, and Valadao, in press). Previously: TFTSG Provisional Red List: Vulnerable (VU) (2011); Vulnerable (VU) (TFTSG 1996)

CITES: Appendix II, as *Podocnemis* spp. (1975)

Synonymy:

Podocnemis sextuberculata Cornalia 1849:13 (17:105)

Bartlettia sextuberculata
Type locality: "Fl. Amazonum" [Amazonas, Brazil] [yellow star, location estimated, possibly nr. Manaus].
Type specimen: MSNM, holotype, type specimen figured (pl.12,f.3), apparently lost (Bour *in* Iverson 1992).
Etymology:* Latin, *sex*, six, and *tuberculata*, having small bumps; referring to the six bumps on the juvenile plastron.


Podocnemis pitiu Coutinho 1868:150 (10:47)
Type locality: "l'Amazone" [Amazonas, Brazil] [yellow star, location estimated, possibly nr. Manaus].
Type specimens: Not known or located.
Etymology:* Vernacular (Amazonian Native, Tupi Guarani), *jurará-pitiú*, local indigenous name for the species.

Bartlettia pitipii Gray 1870f:720
Bartlettia pitipiti
Type locality: "Lakes of the Upper Amazons" [Amazonas, Brazil] [green star, location estimated, possibly Lake Coari].
Type specimens: NHMUK 1946.1.22.90, 1947.3.5.82–83 (formerly 1869.6.7.8–9), syntypes (3), see Iverson et al. (2017).
Etymology:* Vernacular (Amazonian Native, Tupi Guarani), *jurará-pitiú*, local indigenous name for the species.

Podocnemis unifilis Troschel 1848 (07:105, 08:16, 09:49, 10:48, 10:49, 14:50, 21:41)
Yellow-spotted River Turtle, Yellow-spotted Sideneck Turtle, *Terecay*, *Tracajá*, *Taricaya*


Richard C. Vogt / Amazonas, Brazil / female


left: Andreas Nöllert / Rio Tiguino, Ecuador / female
right: Russell A. Mittermeier / Amazonas, Brazil / male


** (orange dots = trade or introduced; stars = type localities)

Distribution: Bolivia (*Beni*, *Cochabamba*, *La Paz*, *Pando*, *Santa Cruz*), Brazil (*Acre*, *Amapá*, *Amazonas*, *Goiás*, *Mato Grosso*, *Maranhão*, *Pará*, *Rondônia*, *Roraima*, *Tocantins*), Colombia (*Amazonas*, *Arauca*, *Caquetá*, *Casanare*, *Guainía*, *Meta*, *Putumayo*, *Vaupés*, *Vichada*), Ecuador, French Guiana, Guyana, Peru (*Huanuco*, *Loreto*, *Madre de Dios*, *Pasco*, *Ucayali*), Suriname, Venezuela (*Amazonas*, *Anzoátegui*, *Apure*, *Barinas*, *Bolívar*, *Cojedes*, *Delta Amacuro*, *Guárico*, *Monagas*)
Introduced: Brazil (*Mato Grosso do Sul*, *Pernambuco*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 5,213,904 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 8,111,508 sq. km
Size (Max SCL):* male 33.5 cm, female 50.0 cm (Pritchard and Trebbau 1984; Barrio Amorós 2001; Ceballos et al. 2013)
IUCN Red List:* Vulnerable (VU A2cd+4bcd) (Escalona, Norris, de Thoisy, Carvajal-Bacarreza, Cueva, Conway-Gómez, Andrade, Ouboter, and Acebey-Quiroga, in press). Previously: TFTSG Provisional Red List: Endangered (EN) (2011); Vulnerable (VU A1acd [ver. 2.3]) (TFTSG 1996)
CITES: Appendix II, as *Podocnemis* spp. (1975)
USA ESA:* Endangered (USFWS 1970)
Synonymy:

Emys cayennensis Schweigger 1812:298 (*partim*, misidentified type [= *Podocnemis unifilis*]) (10:48, 14:50, 21:41)
Chelys (Hydraspis) cayennensis, *Chelys cayennensis*, *Hydraspis cayennensis*, *Podocnemis cayennensis*
Type locality: "Cayenna" [Cayenne, French Guiana] [orange star].
Type specimen: MNHN 8359, lectotype, designated by Bour (2006a:33), discussed by Pritchard and Trebbau (1984), Ceríaco and Bour (2012), and Vogt et al. (2013); not MNHN 4152, designated lectotype by Hoogmoed and Gruber (1983:343).
Etymology:* Toponym for Cayenne, French Guiana.
Comment: A proposal to the ICZN by Vogt et al. (2013) to give *Podocnemis unifilis* Troschel 1848 precedence over *Emys cayennensis* Schweigger 1812 whenever the two are considered synonyms, was approved by ICZN (2019). Name synonymized *in partim* under *Podocnemis erythrocephala* and *Podocnemis unifilis*.

Testudo terekay Humboldt 1819a:243 (*nomen oblitum*) (14:49)
Emys terekay
Type locality: "Haut-Orénoque...l'Apure, l'Uritucu, la Guarico et...les Llanos de Caracas" [= Upper Orinoco, Apure, Uritucu, Guarico, and the Llanos of Caracas, Venezuela]; restricted here to "junction of Rio Apure and Rio Orinoco, Venezuela" [green star].
Type specimen: Not known or located, see Vogt et al. (2013).
Etymology:* Vernacular (Native Amazonian), *terekay*, local indigenous name for the species.

Chelys (Hydraspis) lata Bell *in* Gray 1830e:17 (*nomen oblitum*) (08:16, 10:7)

Chelys lata, Hydraspis lata
Type locality: "Demerara" [Guyana] [blue star].
Type specimen: Not located, possibly OUM, apparently lost.
Etymology:* Latin, *lata*, broad, wide, extensive, flat; referring to the somewhat flatter carapace than some other species.

Podocnemis unifilis Troschel 1848:647 (14:50, 21:41)
Type locality: "Britisch-Guiana...Rupununi und Takutu" [Guyana]; restricted here to "Rupunini River, Guyana" [yellow star, location estimated].
Type specimen: ZMB 142, lectotype, designated by Vogt et al. (2013:35).
Etymology:* Latin, *unum*, one, and *filum*, thread; referring to the single chin barbel.
Comment: A proposal to the ICZN by Vogt et al. (2013) to give *Podocnemis unifilis* Troschel 1848 precedence over *Emys cayennensis* Schweigger 1812 whenever the two are considered synonyms, was approved by ICZN (2019).


Podocnemis tracaya Coutinho 1868:149 (10:49)
Type locality: "l'Amazone" [Amazonas, Brazil] [no star].
Type specimens: Not known or located.
Etymology:* Vernacular (Native Amazonian), *tracaja*, local indigenous name for the species.

Emys quadripunctata Lidth de Jeude *in* Gray 1873j:74 (*nomen nudum*)
Type locality: "India" [in error], restricted to "Tropical America" by Gray (1873j:74).
Type specimen: NHMUK 1866.8.14.223, see Gray (1873j).
Etymology:* Latin, *quadri-*, four, and *punctata*, spotted or punctured; referring to the four prominent head spots.

Podocnemis vogli Müller 1935 (28)*
Llanos Sideneck Turtle, Savanna Sideneck Turtle, *Galápago Sabanero*


Anders G.J. Rhodin / Hato El Cedral, Mantecal, Apure, Venezuela


Anders G.J. Rhodin / Hato El Cedral, Mantecal, Apure, Venezuela


**
(orange dots = probable trade, introduced, or translocated; star = type locality)

Distribution: Colombia (*Arauca, Casanare, Guaviare, Meta, Vichada*), Venezuela (*Amazonas, Anzoátegui, Apure, Barinas, Bolívar, Cojedes, Delta Amacuro, Guárico, Monagas, Portuguesa*)
Introduced:* Venezuela (*Lara, Zulia*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 364,561 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 906,029 sq. km
Size (Max SCL):* male 27.7 cm, female 36.9 cm (Pritchard and Trebbau 1984; Thorbjarnarson et al. 1997; Viloria and Forti 2015)
IUCN Red List:* Vulnerable (VU A2cd+4cd) (Páez, Rivas, Rojas-Runjaic, Bock, Hernández, Lasso, Lasso-Alcalá, Morales-Betancourt, and Escalona, in press). Previously: TFTSG Provisional Red List: Vulnerable (VU) (2011); Least Concern (LC) [Not Listed] (Behler 1996)
CITES: Appendix II, as *Podocnemis* spp. (1975)
Synonymy:
Podocnemis vogli Müller 1935:104
Type locality: "Barinas (Staat Zamora), Venezuela" [= Barinas, Barinas Province] [yellow star].
Type specimen: ZSM 128/1928, holotype, see Franzen and Glaw (2007).
Etymology:* Eponym honoring Father Cornelius Vogl (1884–1959), Germany / Venezuela; priest and naturalist who worked in Venezuela and collected the type.

Suborder Cryptodira Cope 1869 (08:20, 21:3, 21:6, 21:8)


Hidden-necked Turtles

(includes 11 families, 77 genera, 269 species, 387 taxa)

Cryptodères Duméril and Bibron 1834:354 (invalid vernacular name)

Cryptodera Lichtenstein and von Martens 1856:1

Cryptodira Cope 1869:282



Cryptodira Species Richness

Infraorder Durocryptodira Danilov and Parham 2006 (21:3)

Hardshell Hidden-necked Turtles

(includes 9 families, 63 genera, 233 species, 339 taxa)

Durocryptodira Danilov and Parham 2006:577

Superfamily Chelonioidea Oppel 1811 (29)*

Sea Turtles

(includes 2 families, 6 genera, 7 species, 7 taxa)

Chelonii Oppel 1811:8

Chlonopteria Rafinesque 1814:66

Cheloniae Schmid 1819:14

Edigitata Haworth 1825:373

Oiacopodae Wagler 1828:861

Chelonidae Bonaparte 1831:64

Oeacopodes Burmeister 1837:731

Pterodactyli Mayer 1849:199

Chelonioidea Baur 1893b:673

Family Cheloniidae Oppel 1811 (09:5, 12:7, 12:8, 12:9) (30)*

Sea Turtles

(includes 2 subfamilies, 5 genera, 6 species, 6 taxa)

Chelonii Oppel 1811:8 (partim)

Cheloniadae Gray 1825:212


Carettidae Gray 1825:212

Mydae Ritgen 1828:269

Chelonina Bonaparte 1831:64

Cheloniidae Cope 1869:282 (21:6)

Cheloniadi Portis 1890:23



Cheloniidae Species Richness

Subfamily Carettinae Gray 1825 (12:7)

Carettine Sea Turtles

(includes 3 genera, 4 species, 4 taxa)

Carettidae Gray 1825:212

Carettinae Deraniyagala 1952:57

Caretta Rafinesque 1814 (12:8)

Loggerhead Sea Turtles

(includes 1 species, 1 taxon)

Caretta Rafinesque 1814:66

Type species: *Caretta nasuta* Rafinesque 1814 [= subjective synonym of *Testudo caretta* Linnaeus 1758], by original monotypy.

Etymology:* Vernacular (Old French), *caret*, sea turtle.

Thalassochelys Fitzinger 1835:121

Type species: *Thalassochelys caouana* Fitzinger [= *Testudo caouana* Lacepède 1788 (*nomen suppressum*) = *Testudo caouana* Bonnaterre 1789 = subjective synonym of *Testudo caretta* Linnaeus 1758], by subsequent designation by Fitzinger (1843:30).

Etymology:* Greek, *thalassa*, the sea, and *chelys*, turtle; referring to the marine habitat.

Caouana Cocteau *in* Cocteau and Bibron 1838:31

Type species: *Chelonia* (*Caouana) cephalo* [= *Testudo cephalo* Schneider 1783 = subjective synonym of *Testudo caretta* Linnaeus 1758], by original monotypy.

Etymology:* Vernacular (Old French), *caouane*, large sea turtle; probable origin in the West Indies.

Halichelys Fitzinger 1843:30

Type species: *Thalassochelys* (*Halichelys) atra* Fitzinger [= *Caretta atra* Merrem 1820 = subjective synonym of *Testudo caretta* Linnaeus 1758], by original designation.

Etymology:* Greek, *hals*, sea, salt, salt water, *chelys*, turtle; referring to the marine habitat.

Eremonia Gray 1873i:408

Type species: *Eremonia elongata* Gray [= *Caouana elongata* Gray 1844 = subjective synonym of *Testudo caretta* Linnaeus 1758], by original designation.

Etymology:* Greek, *eremos*, lonely, desolate, hermit; probably in reference to Gray noting that he never succeeded in finding a second specimen of what he considered the type species, *Caouana elongata*.

Caretta caretta (Linnaeus 1758) (10:5, 14:2, 17:5, 21:42)
Loggerhead, Loggerhead Sea Turtle




Blair Witherington / CRM 3 / Atlantic Ocean off the Florida Keys, Florida






left: Blair Witherington / CRM 3 / Atlantic Ocean off St. Lucie Co., Florida
right: Blair Witherington / CRM 3 / Gulf of Mexico off Monroe Co., Florida



**





(yellow dots = nesting sites; blue lines delimit purple Regional Management Units; stars = type localities)

Nesting: Aruba, Australia (Queensland, Western Australia), Bahamas, Bangladesh, Belize, Bermuda, Bonaire, Brazil (Bahia, Espirito Santo, Rio de Janeiro, Sergipe), Cape Verde, Cayman Islands, China, Colombia, Costa Rica, Cuba, Curacao, Cyprus, Dominican Republic, Egypt, France, Greece, Haiti, Honduras, Israel, Italy, Japan, Lebanon, Libya, Madagascar, Mauritania, Mexico, Montserrat, Mozambique, Myanmar, New Caledonia, Oman, Panama, Papua New Guinea (Trobriand Islands), Saint Lucia, Saint Vincent and the Grenadines, Sierra Leone, South Africa, Spain, Sri Lanka, Syria, Tunisia, Turkey, Turks and Caicos, USA (Alabama, Florida, Georgia, Louisiana, Mississippi, North Carolina, South Carolina, Texas), US Virgin Islands, Vanuatu, Venezuela, Yemen

Foraging: Albania, Algeria, Anguilla, Antigua and Barbuda, Argentina, Bahrain, Barbados, British Virgin Islands, Canada, Chile, Comoros, Croatia, Djibouti, Dominica, Eritrea, Fiji, French Guiana, Gambia, Grenada, Guadeloupe, Guinea-Bissau, Guatemala, Guyana, India, Indonesia, Iran, Jamaica, Kenya, Malta, Martinique, Mauritius, Monaco, Montenegro, Morocco, Namibia, Netherlands Antilles (Bonaire, Saba, St. Eustatius), Nicaragua, North Korea, Pakistan, Peru, Philippines, Portugal, Puerto Rico, Qatar, Réunion, Saint Kitts and Nevis, Samoa, Saudi Arabia, Senegal, Seychelles, Sint Maarten, Slovenia, Solomon Islands, Somalia, South Korea, Sudan, Suriname, Taiwan, Tanzania, Tonga, Trinidad and Tobago, United Arab Emirates, Uruguay, USA (California, Hawaii, Oregon), Vietnam, Western Sahara

Vagrant: Angola, Benin, Brunei, Cambodia, Cameroon, Congo (DRC), Congo (ROC), Ecuador, El Salvador, Equatorial Guinea, Gabon, Ghana, Great Britain, Guinea, Iraq, Ireland, Ivory Coast (Côte d'Ivoire), Kuwait, Liberia, Malaysia, Maldives, New Zealand, Nigeria, Thailand, Togo, Tuvalu

Estimated Indigenous (Historical) Ranges (AOO and EOO) (nesting localities intersecting HUC level 12 polygons only):
Est. Indigenous Area of Occupancy (AOO):* 59,636 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 172,370,880 sq. km

Size (Max SCL):* male 99.5 cm, female 115.0 cm (Ehrhart 1978; Hirth 1982; Pritchard and Trebbau 1984; Dodd 1988; Márquez 1990; Figgener et al. 2022); **(Max CCL):*** male 104.0 cm, female 124.0 cm (Ehrhart and Yoder 1978; Limpus 1985; Dodd 1988; Figgener et al. 2022)

Comment: Alleged maximum of 213 cm, cited by Conant (1958) and many others, is erroneous (Pritchard and Trebbau 1984); maximum of 146.7 cm cited by Dodd (1988) is total body length (Brongersma 1972)

IUCN Red List: Vulnerable (VU A2b) (Casale and Tucker 2015, 2017); Subpopulations: Mediterranean: Least Concern (LC) (2015); North East Atlantic: Endangered (EN B2ab(iii)) (2015); North East Indian Ocean: Critically Endangered (CR D) (2015); North Pacific: Least Concern (LC) (2015); North West Atlantic: Least Concern (LC) (2015); North West Indian Ocean: Critically Endangered (CR A4b) (2015); South East Indian Ocean: Near Threatened (NT) (2015); South Pacific: Critically Endangered (CR A2b) (2015); South West Atlantic: Least Concern (LC) (2015); South West Indian Ocean: Near Threatened (NT) (2015). Previously: Endangered (EN) (1996)

CITES: Appendix I, as Cheloniidae spp. (1981)

USA ESA:* Endangered or Threatened, varying for different Distinct Population Segments (USFWS 1978)

Synonymy:

Testudo caretta Linnaeus 1758:197 (21:42)

Chelone caretta, Chelonia caretta, Thalassochelys caretta, Talassochelys caretta, Caouana caretta, Caretta caretta, Caretta caretta caretta

Type locality: "insulas Americanas." Restricted to "Mari Mediterraneo, Atlantico" [no stars] by Schoepff (1793a:70); to "Bermuda Islands" [yellow star] by Smith and Taylor (1950a:315, 1950b:16); and to "Bimini, British Bahamas" [orange star] by Schmidt (1953:107).

Type specimens: Not known or located, description based on cited literature, see Brongersma (1961) and Wallin (1985); Brongersma (1961) recommended that a neotype be designated.

Etymology:* Vernacular (Old French), *caret*, sea turtle.

Comment: Description cited as sourced from Gronovius (1756:85,n.69), Browne (1756:465), Catesby (1743:39,pl.39), and Ray (1693:258).

Testudo marina Garsault 1764:pl.675 (10:5)

Type locality: Not designated.

Type specimen: Not located, type specimen figured (pl.675).

Etymology:* Latin, *marina*, living in the sea; referring to its habitat in the ocean.

Testudo cephalo Schneider 1783:303

Caretta cephalo, Chelonia cephalo, Thalassochelys cephalo

Type locality: Not designated. Restricted to "Charleston, South Carolina" [USA] [green star] by Smith and Taylor (1950a:360).

Type specimen: Not known or located.

Etymology:* Greek, *kefale*, head; referring to the large head.

Testudo caouana Lacepède 1788:95, synopsis[table] (*nomen suppressum*) (09:6)

Type locality: "les contrées chaudes du nouveau Continent."

Type specimen: Possibly MNHN, not located.

Etymology:* Vernacular (Old French), *caouane*, large sea turtle; probable origin in the West Indies.

Comment: Name suppressed by ICZN (2005a) as published in a rejected and invalid non-binomial work, see Savage (2003).

Testudo nasicornis Lacepède 1788:103, synopsis[table] (*nomen suppressum*) (09:7)

Testudo caretta nasicornis, Caretta nasicornis

Type locality: "mers du nouveau Continent, voisines de l'équateur." Restricted to "Ascension Island" [blue star] by Smith and Smith (1980:302).

Type specimen: Possibly MNHN, not located.

Etymology:* Latin, *nasus*, nose, and *cornus*, horn, referring to the horn-like nose.

Comment: Name suppressed by ICZN (2005a) as published in a rejected and invalid non-binomial work, see Savage (2003).

Testudo caouana Bonnaterre 1789:20

Chelonia caouana, Caretta caouana, Thalassochelys caouana

Type locality: "Les contrées chaudes de l'ancien & du nouveau Continent, les côtes de la Jamaïque, très-fréquemment dans la Méditerranée, auprès de Cagliari en Sardaigne & de Castel-Sardo, vers le quarante-uniéme degré de latitude" [no stars].

Type specimen: Possibly MNHN, not located.

Etymology:* Vernacular (Old French), *caouane*, large sea turtle; probable origin in the West Indies.

Testudo lauanna Meyer 1790:82 (*nomen novum et oblitum*) (09:8)

Etymology:* Vernacular (Old French), *caouane (*misspelled as *lauanne)*, large sea turtle; probable origin in the West Indies.

Comment: Unjustified replacement name for *caouana*.

Testudo gigas Walbaum *in* Donndorff 1798:35

Type locality: Not designated. Restricted to "Westindischen Meere" by Bechstein (1800:273).

Type specimen: Not known or located.

Etymology:* Greek, *gigas*, giant; referring to the large body size.

Caretta nasuta Rafinesque 1814:66

Type locality: "Sicil." [Sicily, Italy] [pink star, location estimated].

Type specimen: Not known or located.

Etymology:* Latin, *nasuta*, having a big nose; referring to the prominent snout.

Caretta atra Merrem 1820:17

Chelonia (Thalassochelys) atra, Thalassochelys atra, Thalassochelys (Halichelys) atra, Halichelys atra

Type locality: "mari ad Insulam Adscensionis" [Ascension] [blue star].

Type specimen: Not known or located.

Etymology:* Latin, *atra*, black; referring to the color of the shell.

Comment: Description cited as sourced from Linnaeus (1749b:284,n.7, *T. mydas γ*; 1754:50, *T. atra*; and 1766:351, *T. mydas γ*), a mix of *Caretta caretta* and *Chelonia mydas*.

Chelonia radiata Cuvier 1829:14

Type locality: Not designated.

Type specimen: MNHN s/n, holotype, not located; Cuvier (1829) and Cogger et al. (1983) cite Schoepff (1793b:pl.16B) [plate number corrected from Schoepff (1793a:pl.17B)] as the figured type specimen, which is a *Caretta caretta*.

Etymology:* Latin, *radius*, ray, rod, spoke; referring to the radiating pattern on the carapacial scutes.

Testudo corianna Gray 1831d:53 (*nomen novum*)

Etymology:* Vernacular (Old French), *caouane*, large sea turtle; probable origin in the West Indies.

Comment: Unjustified emendation or error for *caouana*.

Chelonia pelasgorum Valenciennes *in* Bory de Saint-Vincent 1833:planches, pl.6 (14:2)

Type locality: "sur la plage sablonneuse entre Arcadia et l'embouchure de la Neda" [Greece] [brown star].

Type specimen: MNHN 7907, holotype, figured (pl.6).

Etymology:* Toponym for Pelasgia, an historical region of Greece.

Chelonia pelasgica Bibron and Bory de Saint-Vincent 1833:64 (*nomen novum*) (14:2)

Etymology:* Toponym for Pelasgia, an historical region of Greece.

Comment: Unjustified emendation or error for *pelasgorum*.

Caouana elongata Gray 1844:53

Thalassochelys elongata, Eremonia elongata

Type locality: Not designated. Restricted to "Ascension Island" [blue star] by Smith and Smith (1980:303).

Type specimen: NHMUK 1947.3.5.41, holotype, see Cogger et al. (1983).

Etymology:* Latin, *elongata*, stretched; referring to the elongated carapace.

Thalassochelys corticata Girard 1858:431

Type locality: "Madeira" [Canary Islands] [light gray star]. Emended to "Funchal, Madeira (Portugal)" by Cochran (1961:235).

Type specimen: USNM 7778, holotype, see Cochran (1961) and Reynolds et al. (2007).

Etymology:* Latin, *corticata* or *cortex*, bark; possibly an error for *cordicata*, heart-shaped; possibly referring to the carapace, described as subcordiform.

Caretta gigas Deraniyagala 1933:66

Caretta caretta gigas

Type locality: "Ceylon" [Sri Lanka]. Emended to "Gulf of Mannar" [Sri Lanka] [dark gray star] by Deraniyagala (1939:164).

Type specimen: NHMUK 1946.1.22.64, 1947.3.5.76 (separate parts of one specimen) (formerly 1934.5.1.1), holotype, see Das (2009).

Etymology:* Greek, *gigas*, giant; referring to the large body size.

Eretmochelys Fitzinger 1843 (09:5, 12:7, 12:8, 14:3)

Hawksbill Sea Turtles

(includes 1 species, 1 taxon)

Eretmochelys Fitzinger 1843:30

Type species: *Chelonia (Eretmochelys) imbricata* Cuvier [= *Testudo imbricata* Linnaeus 1766], by original designation.

Etymology:* Greek, *eretman*, having oars, and *chelys*, turtle; referring to the oar-shaped fins.

Herpysmostes Gistel 1868:145

Type species: *Herpysmostes imbricatus* [= *Testudo imbricata* Linnaeus 1766], by original monotypy.

Etymology:* Greek, *herpestes*, creeper, sneak; referring to the creeping mode of locomotion when nesting.

Onychochelys Gray 1873i:397

Type species: *Onychochelys kraussi* Gray 1873i [= subjective synonym of *Testudo imbricata* Linnaeus 1766], by original monotypy.

Etymology:* Greek, *onyx*, claw, nail, and *chelys*, turtle; referring to the conspicuously large claw on the first front toe.

Eretmochelys imbricata (Linnaeus 1766) (07:5, 09:9, 12:9, 14:4, 17:6)


Hawksbill Turtle, Hawksbill Sea Turtle



Robert P. Van Dam / CCB / Mona Island, Puerto Rico



left: Richard Herren / CRM 3 / Hutchinson Island, St. Lucie Co., Florida
right: Blair Witherington / CRM 3 / Key West National Wildlife Refuge, Monroe Co., Florida



(yellow dots = nesting sites; blue lines delimit purple Regional Management Units; stars = type localities)

Nesting: Anguilla, Antigua and Barbuda, Aruba, Australia, Bahamas, Bangladesh, Barbados, Belize, Brazil, British Indian Ocean Territory, British Virgin Islands, Cameroon, Cayman Islands, China, Colombia, Congo (ROC), Costa Rica, Cuba, Curacao, Dominica, Dominican Republic, Ecuador, Egypt, El Salvador, Equatorial Guinea, Eritrea, Fiji, French Guiana, French Southern Territories, Gabon, Grenada, Guadeloupe, Guatemala, Guinea, Guinea-Bissau, Guyana, Haiti, Honduras, India, Indonesia, Iran, Ivory Coast (Côte d'Ivoire), Jamaica, Japan, Kenya, Liberia, Madagascar, Malaysia, Maldives, Martinique, Mexico (Campeche, Yucatán),

Micronesia, Montserrat, Mozambique, Netherlands Antilles (Bonaire, Sint Eustatius), Nicaragua, Oman, Palau, Panama, Papua New Guinea, Philippines, Puerto Rico, Qatar, Saint Kitts and Nevis, Saint Lucia, Saint Vincent and the Grenadines, Samoa, São Tomé and Príncipe, Saudi Arabia, Senegal, Seychelles, Sint Maarten, Solomon Islands, Sri Lanka, Suriname, Taiwan, Tanzania, Thailand, Trinidad and Tobago, Turks and Caicos, USA (Florida, Hawaii, Texas), US Virgin Islands, Vanuatu, Venezuela, Vietnam

Foraging: American Samoa, Argentina, Ascension, Bahrain, Benin, Bermuda, Brunei, Cambodia, Cook Islands, Djibouti, French Polynesia, Gambia, Ghana, Guam, Iraq, Israel, Kuwait, Mauritania, Mauritius, Mayotte, Myanmar, Nigeria, Northern Mariana Islands, Pakistan, Peru, Réunion, Sierra Leone, Singapore, Society Islands, Somalia, South Africa, Sudan, Togo, Tokelau, Tonga, Tuamotu, Tuvalu, United Arab Emirates, Wallis and Futuna, Yemen

Vagrant: Algeria, Angola, Cape Verde, Chile, Comoros, Congo (DRC), Kiribati, Marshall Islands, Morocco, Namibia, Nauru, New Caledonia, North Korea, Pitcairn Island, Portugal, South Korea, Spain, Uruguay

Estimated Indigenous (Historical) Ranges (AOO and EOO) (nesting localities intersecting HUC level 12 polygons only):
Est. Indigenous Area of Occupancy (AOO):* 100,167 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 194,744,608 sq. km

Size (Max SCL):* male 85.1 cm, female 114.0 cm (Witzell 1983; Márquez 1990)

IUCN Red List: Critically Endangered (CR A2bd) (Mortimer and Donnelly 2008). Previously: Critically Endangered (CR) (1996)

CITES: Appendix I, as Cheloniidae spp. (1981); subspecies *E. i. imbricata* listed as Appendix I (1975, 1977), and as Cheloniidae spp. (1981); subspecies *E. i. bissa* listed as Appendix II (1975), and Appendix I (1977), and as Cheloniidae spp. (1981)

USA ESA:* Endangered (USFWS 1970)

Synonymy:

Testudo imbricata Linnaeus 1766:350
Chelone imbricata, Chelonia imbricata, Caretta imbricata, Eretmochelys imbricata, Herpysmostes imbricatus, Herpysmostes imbricata, Chelonius imbricatus, Eretmochelys imbricata imbricata
Type locality: "Mari Americano, Asiatico." Restricted to "Bermuda Islands" [orange star] by Smith and Taylor (1950a:315, 1950b:17), and to "Belize, British Honduras" [yellow star] by Schmidt (1953:106).
Type specimens: UPSZTY 130 (formerly UUZM 130), possible syntype, listed by Thunberg (1828), identified as a *Testudo imbricata* by Lönnberg (1896), but as a *Testudo mydas* (= *Chelonia mydas*) by Wallin (1985, 2001).
Etymology:* Latin, *imbricatus*, overlapping like roof shingles; referring to the overlapping carapacial scutes.
Comment: Description cited as sourced from Gronovius (1763:16, n.72), Bontius (1658:82), Du Tertre (1667:229), and Ray (1693:258).

Testudo nasicornis Bonnaterre 1789:21 (09:7)
Type locality: "La Zone torride, les mers du nouveau Continent."
Type specimen: Not located, type specimen figured (pl.3,f.3), lectotype, designated "holotype" by TTWG (2009:62).
Etymology:* Latin, *nasus*, nose, and *cornus*, horn; referring to the horny beak.

Chelonia pseudomydas Lesson 1831b:299
Chelonia pseudo-mydas
Type locality: "l'Océan atlantique." Restricted to "Bermuda Islands" [orange star] by Smith and Taylor (1950a:315).
Type specimen: Possibly MNHN, not located.
Etymology:* Greek, *pseudo-*, false or deceptive, and *mydos*, wetness; referring to the similarity between this species and *Chelonia mydas*.

Chelonia pseudocaretta Lesson 1831b:302
Chelonia pseudo-caretta
Type locality: "l'Océan atlantique." Restricted to "Bermuda Islands" [orange star] by Smith and Taylor (1950a:315).
Type specimen: Possibly MNHN, not located.
Etymology:* Greek, *pseudo-*, false, similar, and Vernacular (Old French), *caret*, sea turtle; referring to the similarity between this species and *Caretta caretta*.

Caretta bissa Rüppell 1835:4 (07:5, 09:9)
Eretmochelys imbricata bissa
Type locality: "im rothen Meere...Abyssinien" [Red Sea...Ethiopia] [green star].
Type specimen: SMF 7886, lectotype, designated by Mertens (1967b:52).
Etymology:* Vernacular (Ethiopian), *bissa*, local indigenous name for the male of this species.

Eretmochelys squamata Agassiz 1857a:382
Caretta squamata, Eretmochelys imbricata squamata
Type locality: "Singapore and Bengal, India." Restricted to "Singapore, Straits Settlements" [blue star] by Smith and Taylor (1950a:315, 1950b:17).
Type specimen: MCZ 1416, lectotype, designated by Smith and Taylor (1950b:17).
Etymology:* Latin, *squama*, scale; referring to the carapacial scales used for tortoiseshell.

Caretta squamosa Girard 1858:442
Eretmochelys squamosa, Eretmochelys imbricata squamosa
Type locality: "Sooloo Seas and Indian Ocean." [Sulu Sea, Philippines] [pink star, location estimated]
Type specimen: USNM, not listed by Cochran (1961) or Reynolds et al. (2007).
Etymology:* Latin, *squama*, scale; referring to the carapacial scales used for tortoiseshell.
Comment:* Based on the pre-Linnaean description of *Testudo squamosa* (Ray 1693:260); not a *nomen novum* for *Caretta squamata* (Agassiz 1857a).

Caretta rostrata Girard 1858:446
Type locality: "Feejee Islands" [Fiji] [light gray star].
Type specimens: USNM 257185–86 (formerly USNM Osteo 12387–88), syntypes (2), see Cochran (1961) and Reynolds et al. (2007).
Etymology:* Latin, *rostrum*, snout, beak; referring to the prominent horny beak.

Onychochelys kraussi Gray 1873i:398
Type locality: "Ocean, French Guiana" [dark gray star].
Type specimens: NHMUK 1947.3.4.37 and 1947.3.5.43 (formerly 1873.3.13.3–4), syntypes (2), see Cogger et al. (1983).
Etymology:* Eponym honoring Christian Ferdinand Krauss (1812–1890), Germany; zoologist and director of the Staatliches Museum für Naturkunde, Stuttgart, who presented the types.

Lepidochelys Fitzinger 1843 (12:8) (31)*

Ridley Sea Turtles
(includes 2 species, 2 taxa)

Lepidochelys Fitzinger 1843:30
Type species: *Thalassochelys (Lepidochelys) olivacea* Eschscholtz 1829a [= *Chelonia olivacea* Eschscholtz 1829a], by original designation.
Etymology:* Greek, *lepidotos*, covered with scales, and *chelys*, turtle; referring to the increased number of carapace scutes compared to other sea turtles.

Cephalochelys Gray 1873i:408
Type species: *Cephalochelys oceanica* Gray 1873i [= subjective synonym of *Chelonia olivacea* Eschscholtz 1829a], by original monotypy.
Etymology:* Greek, *kefale*, head, and *chelys*, turtle; referring to the relatively large head size.


Colpochelys Garman 1880:124
Type species: *Thalassochelys (Colpochelys) kempii* Garman 1880, by

original monotypy.
Etymology:* Greek, *kolpos*, gulf, bay, and *chelys*, turtle; referring to the range of the genus in the Gulf of Mexico.


Lepidochelys kempii (Garman 1880) (31)*
Kemp's Ridley, Kemp's Ridley Sea Turtle, Atlantic Ridley



Blair Witherington / CRM 3 / Florida Bay, Monroe Co., Florida



left: Jeffrey R. Schmid / CRM 3 / Levy Co., Florida
right: Peter C.H. Pritchard / CCB / Tamaulipas, Mexico



(yellow dots = nesting sites; stars = type localities;
blue line delimits purple Regional Management Unit)

Nesting:* Mexico (Tabasco, Tamaulipas, Veracruz), USA (Florida, Georgia, Texas)
Foraging:* Mexico (Tabasco, Veracruz), USA (Alabama, Connecticut, Delaware, Florida, Georgia, Louisiana, Maryland, Massachusetts, Mississippi, New Jersey, New York, North Carolina, Rhode Island, South Carolina, Virginia)
Vagrant: Algeria, Anguilla, Bahamas, Bermuda, British Virgin Islands, Canada, Cayman Islands, Cuba, France, Ireland, Italy, Morocco, Portugal, Spain
Estimated Indigenous (Historical) Ranges (AOO and EOO) (nesting localities intersecting HUC level 12 polygons only):
Est. Indigenous Area of Occupancy (AOO):* 5,638 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 1,803,050 sq. km
Size (Max SCL):* male 72.0 cm, female 76.0 cm (Márquez 1990; Ernst and Lovich 2009)

IUCN Red List: Critically Endangered (CR A2bd) (Wibbels and Bevan 2019). Previously: Critically Endangered (CR) (1996)
CITES: Appendix I, as Cheloniidae spp. (1981)
USA ESA:* Endangered (USFWS 1970)
Synonymy:

Testudo viridisquamosa Lacepède 1788:92, synopsis[table] (*partim, nomen dubium et suppressum*) (09:6)
Testudo viridi-squamosa
Type locality: "la mer du Sud, auprès du Cap Blanco, de la nouvelle Espagne...aussi dans le golfe du Mexique...&...les rivages chauds du nouveau monde." Restricted to "Bocas del Toro, Panama, Golf von Mexico" [orange star, in error] by Wermuth (1956:405); and to "the island of Blanquilla...West Indies" [Venezuela] [brown star, in error] by Brongersma (1961:27).
Type specimen: Possibly MNHN, not located.
Etymology:* Latin, *viridis*, green, emerald, and *squamosa*, scaly; referring to the allegedly somewhat green color of the carapacial scutes.
Comment: Name suppressed by ICZN (1963), see Wermuth (1956) and Mertens and Wermuth (1961), and by ICZN (2005a) as published in a rejected and invalid non-binomial work, see Savage (2003). Name synonymized *in partim* under *Lepidochelys kempii* and *Chelonia mydas*.

Testudo viridisquamosa Bonnaterre 1789:20 (*partim, nomen dubium*)
Testudo viridi-squamosa
Type locality: "La mer du Sud, le golfe du Mexique, les rivages du Nouveau-Monde."
Type specimen: Possibly MNHN, not located.
Etymology:* Latin, *viridis*, green, emerald, and *squamosa*, scaly; referring to the allegedly somewhat green color of the carapacial scutes.
Comment:* Name synonymized *in partim* under *Lepidochelys kempii* and *Chelonia mydas*.

Testudo bomarii Meyer 1790:82 (*partim, nomen dubium et oblitum*) (09:8)
Etymology:* Eponym honoring Jacques-Christophe Valmont de Bomare (1731–1807), France; botanist and naturalist who wrote an influential encyclopedia of natural history in the 1760s.
Comment: Unjustified replacement name for *viridisquamosa*. Name synonymized *in partim* under *Lepidochelys kempii* and *Chelonia mydas*.


Testudo mydas minor Suckow 1798:30 (*partim, nomen dubium et suppressum*)
Type locality: "Amazonen-Flusse..[&]..Südsee..[&]..Cap Blanco in Mexico." Restricted to "the island of Blanquilla...West Indies" [Venezuela] [brown star, in error] by Brongersma (1961:27).
Type specimen: Not known or located.
Etymology:* Latin, *minor*, small; referring to its smaller size.
Comment: Name suppressed by ICZN (1963), see Wermuth (1956) and Mertens and Wermuth (1961). Name synonymized *in partim* under *Lepidochelys kempii* and *Lepidochelys olivacea*.

Thalassochelys (Colpochelys) kempii Garman 1880:123 (*nomen protectum*)
Lepidochelys kempii, Colpochelys kempii, Caretta kempii, Lepidochelys olivacea kempii
Type locality: "Gulf of Mexico." Restricted to "Key West, Monroe Co., Florida" [USA] [yellow star] by Smith and Taylor (1950a:358, 1950b:15).
Type specimens: MCZ 46538–39, syntypes (2), see Barbour and Loveridge (1929).
Etymology:* Eponym honoring Richard M. Kemp (1825–1908), Bahamas / USA; fisherman who obtained the types.
Comment: Name conserved by ICZN (1963), see Wermuth (1956) and Mertens and Wermuth (1961).


Lepidochelys olivacea (Eschscholtz 1829a) [17:7] [31]*
Olive Ridley, Olive Ridley Sea Turtle, Pacific Ridley




Nicolas J. Pilcher / Red Sea, nr. Assab, Eritrea




Anders G.J. Rhodin / Puerto Escondido, Oaxaca, Mexico / male, female



**





(yellow dots = nesting sites; blue lines delimit purple Regional Management Units; stars = type localities)

Nesting: Angola, Australia (Northern Territory), Bangladesh, Benin, Brazil (Bahia, Espirito Santo, Sergipe), Brunei, Cameroon, Colombia, Congo (ROC), Costa Rica, Ecuador, El Salvador, Equatorial Guinea, Eritrea, French Guiana, Gabon, Ghana, Guatemala, Guinea-Bissau, Guyana, Honduras, India, Indonesia (Java, Papua), Iran, Ivory Coast (Côte d'Ivoire), Kenya, Liberia, Malaysia, Mexico (Baja California Sur, Chiapas, Guerrero, Jalisco, Michoacán, Nayarit, Oaxaca, Sinaloa), Mozambique, Myanmar, Nicaragua, Oman, Pakistan, Panama, Peru, São Tomé and Príncipe, Sierra Leone, Sri Lanka, Suriname, Thailand, Togo, Trinidad and Tobago, Vanuatu, Vietnam

Foraging: Bahrain, Cambodia, Cape Verde, China, Comoros, Congo (DRC), Djibouti, Egypt, Gambia, Guinea, Iran, Iraq, Israel (Southern), Kuwait, Liberia, Madagascar, Maldives, Mauritius, New Caledonia, Nigeria, Papua New Guinea, Philippines, Qatar, Saudi Arabia, Senegal, Seychelles, Singapore, Solomon Islands, Somalia, Sudan, Tanzania, Timor-Leste, United Arab Emirates, USA (Hawaii), Venezuela, Yemen

Vagrant: Antigua and Barbuda, Barbados, Canada, Chile, Cuba, Dominica, Dominican Republic, Grenada, Guadeloupe, Haiti, Jamaica, Japan, Marshall Islands, Martinique, Mauritania, Micronesia, Morocco, Namibia, New Zealand, North Korea, Puerto Rico, Saint Kitts and Nevis, Saint Lucia, Saint Vincent and the Grenadines, South Africa, South Korea, Spain, Taiwan, Uruguay, US Virgin Islands

Estimated Indigenous (Historical) Ranges (AOO and EOO) (nesting localities intersecting HUC level 12 polygons only):
Est. Indigenous Area of Occupancy (AOO):* 40,710 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 122,533,080 sq. km

Size (Max SCL): male 72.0 cm, female 79.3 cm (Ceballos et al. 2013)

IUCN Red List: Vulnerable (VU A2bd) (Abreu-Grobois and Plotkin 2008). Previously: Endangered (EN) (1996)

CITES: Appendix I, as Cheloniidae spp. (1981). Previously: Appendix II (1975)

USA ESA:* Endangered, for breeding colony populations on Pacific coast of Mexico, or Threatened, wherever found, except when listed as Endangered (USFWS 1978)

Synonymy:

Testudo mydas minor Suckow 1798:30 (*partim, nomen dubium et suppressum*)

Type locality: "Amazonen-Flusse..[&]..Südsee..[&]..Cap Blanco in Mexico." Restricted to "the island of Blanquilla...West Indies" [Venezuela] [orange star] by Brongersma (1961:27).
Type specimen: Not known or located.
Etymology:* Latin, *minor*, small; referring to its smaller size.
Comment: Name suppressed by ICZN (1963), see Wermuth (1956) and Mertens and Wermuth (1961). Name synonymized *in partim* under *Lepidochelys kempii* and *Lepidochelys olivacea*.

Chelonia multiscutata Kuhl 1820:78 (*nomen suppressum*)
Type locality: Not designated.
Type specimen: Not known or located.
Etymology:* Latin, *multum*, numerous, and *scutum*, scute; referring to the multiple carapacial costal scutes often seen on this species.
Comment: Name suppressed by ICZN (1963), see Wermuth (1956) and Mertens and Wermuth (1961).

Chelonia olivacea Eschscholtz 1829a:15 (17:7)
Chelonia caretta olivacea, Caretta olivacea, Thalassochelys (Lepidochelys) olivacea, Caouana olivacea, Lepidochelys olivacea, Caretta caretta olivacea, Lepidochelys olivacea olivacea, Caretta olivacea olivacea
Type locality: "Bai von Manilla" [Philippines] [yellow star].
Type specimens: Originally in MZT, syntypes (2), see Smith and Smith (1980:327) and Flores-Villela et al. (2016:161); transferred to ZMMU, now apparently lost, see Flores-Villela et al. (2016); type specimen figured in Eschscholtz (1829b:pl.3).
Etymology:* Latin, *oliva*, olive, and *-aceus*, belonging to, having the nature of; referring to the olive-colored carapace.

Chelonia dussumierii Duméril and Bibron 1835:557
Lepidochelys dussumierii
Type locality: "mers de la Chine et sur la côte de Malabar" [China and India] [green stars, locations estimated].
Type specimens: MNHN 7908 and 8009, syntypes (2).
Etymology:* Eponym honoring Jean-Jacques Dussumier (1792–1883), France; shipowner in the merchant marine and naturalist collector.

Caouana ruppellii Gray 1844:53 (*nomen nudum*)
Type locality: "India?"
Type specimen: NHMUK s/n, holotype, apparently lost (Zug et al. 1998).
Etymology:* Eponym honoring Eduard Rüppell (1794–1884), Germany; mammalogist, naturalist, and explorer.

Chelonia subcarinata Rüppell *in* Gray 1844:53 (*nomen nudum*)
Etymology:* Latin, *sub-*, below, lesser, and *carina*, keel, *carinata*, keeled; referring to the low carapacial vertebral keel.

Chelonia polyaspis Bleeker 1857b:239 (*nomen nudum*)
Type locality: "Batavia...Java" [Indonesia]. Restricted to "Borneo" by Gray (1873j:90).
Type specimen: NHMUK 1863.12.4.119, holotype, see Gray (1873j).
Etymology:* Greek, *polys*, much, often, and *aspis*, shield; referring to the multiple carapacial costal scutes often seen on this species.

Chelonia dubia Bleeker *in* Gray 1864a:13 (*nomen nudum*)
Type locality: Not designated. Restricted to "Borneo" by Gray (1873j:90).
Type specimen: NHMUK 1863.12.4.122, holotype, see Gray (1873j).
Etymology:* Latin, *dubia*, doubtful, uncertain; probably referring to the uncertain identity of this taxon.

Cephalochelys oceanica Gray 1873i:408
Type locality: "West Coast of America – Mexico?" [blue star, location estimated].
Type specimen: NHMUK 1947.3.5.40 (formerly 1871.2.7.48), holotype, see Cogger et al. (1983).
Etymology:* Latin, *oceanica*, belonging to the ocean; referring to the marine habitat.

Thalassochelys tarapacana Philippi 1887:85
Thalassiochelys tarapacona, Thalassochelys tarapacona
Type locality: "Iquique...Chile" [pink star]. Emended to "Iquique (20º12'S; 70º10'O)" [Chile] by Ortiz and Nuñez (1986:6).
Type specimen: MNHNC 1511, holotype, see Ortiz and Nuñez (1986).
Etymology:* Toponym for the Tarapacá region of Chile.

Thalassochelys controversa Philippi 1899:732
Type locality: "Quinteros...Chile" [light gray star]. Emended to "Quinteros (32º46'S; 71º32'O)" [Chile] by Ortiz and Nuñez (1986:6).
Type specimen: MNHNC 1512, lectotype, designated by Ortiz and Nuñez (1986:6).
Etymology:* Latin, *controversa*, controversial; referring to opinions by others that this taxon was perhaps a variant of *Caretta caretta*.

Caretta remivaga Hay 1908a:194
Lepidochelys olivacea remivaga
Type locality: "Ventosa Bay, Gulf of Tehuantepec, Oaxaca, Mexico" [dark gray star].
Type specimen: USNM 243393 (formerly USNM Osteo 9973), holotype, see Cochran (1961) and Reynolds et al. (2007).
Etymology:* Latin, *remus*, oar, and *vaga*, to wander; referring to the broad range of the species and its oar-like flippers.

Subfamily Cheloniinae Oppel 1811 (12:7)

Cheloniine Sea Turtles
(includes 2 genera, 2 species, 2 taxa)
Chelonii Oppel 1811:8 (*partim*)
Cheloniadae Gray 1825:212
Mydae Ritgen 1828:269
Cheloniidae Cope 1869:282 (21:6)

Chelonia Brongniart 1800 (09:5, 12:8)

Green Sea Turtles
(includes 1 species, 1 taxon)

Chelonia Brongniart 1800b:89
Type species: *Chelonia mydas* [= *Testudo mydas* Linnaeus 1758], by subsequent designation by Bell (1828c:516).
Etymology:* Greek, *chelone*, turtle.

Chelone Brongniart 1805:610 (*nomen novum*)
Etymology:* Greek, *chelone*, turtle.

Chelona Fleming 1828:149 (*nomen novum*)
Etymology:* Greek, *chelone*, turtle.

Mydas Cocteau *in* Cocteau and Bibron 1838:22
Type species: *Chelonia (Mydas) viridis* [= *Testudo viridis* Schneider 1783 = subjective synonym of *Testudo mydas* Linnaeus 1758], by tautonymy.
Etymology:* Greek, *mydos*, dampness, wetness; referring to its aquatic habitat.

Mydasea Gervais 1843:457
Type species: *Chelonia (Mydasea) mydas* [= *Testudo mydas* Linnaeus 1758], by tautonymy.
Etymology:* Greek, *mydos*, dampness, wetness; referring to its aquatic habitat.

Euchelonia Tschudi 1846:22
Type species: *Chelonia (Euchelonia) midas* Schweigger [= *Testudo mydas* Linnaeus 1758], by original monotypy.
Etymology:* Greek, *eu-*, true, and *chelone*, turtle.

Megemys Gistel 1848:8 (*nomen novum*)
Etymology:* Greek, *mega*, large, and *emys*, turtle.

Euchelys Girard 1858:447
Type species: *Euchelys macropus* Girard [= *Testudo macropus* Walbaum 1782 = subjective synonym of *Testudo mydas* Linnaeus 1758], by original monotypy.
Etymology:* Greek, *eu-*, true, and *chelys*, turtle.

Midas Herrera 1901:68 (*nomen novum et suppressum*)
Comment: Unjustified replacement name for *Mydas*. Name suppressed by ICZN (1922).
Etymology:* Greek, *mydos*, dampness, wetness; referring to its aquatic habitat.

Chelonia mydas (Linnaeus 1758) (07:4, 09:5, 12:9, 12:10, 17:8, 21:43) (32)*
Green Turtle, Green Sea Turtle




L.M. Ehrhart / CRM 3 / Melbourne Beach, Brevard Co., Florida




Blair Witherington / CRM 3 / Atlantic Ocean off Florida Keys, Florida



**





(yellow dots = nesting sites; blue lines delimit purple Regional Management Units; stars and triangles = type localities)

Nesting: American Samoa, Angola, Anguilla, Antigua and Barbuda, Aruba, Ascension, Australia (Northern Territory, Queensland, Western Australia), Bahamas, Bangladesh, Barbados, Belize, Brazil, British Virgin Islands, Cayman Islands, China, Cocos (Keeling) Islands, Colombia, Comoros, Congo (ROC), Costa Rica, Cuba, Curacao, Cyprus, Dominica, Dominican Republic, Ecuador, Egypt, El Salvador, Equatorial Guinea, Eritrea, French Guiana, French Polynesia, French Southern Territories, Gambia, Grenada, Guadeloupe, Guam, Guinea-Bissau, Guyana, Haiti, Honduras, India, Indonesia, Iran, Jamaica, Japan, Kenya, Lebanon, Madagascar, Malaysia, Martinique, Mauritania, Mayotte, Mexico (Baja California, Campeche, Michoacán, Quintana Roo, Sinaloa, Sonora, Tabasco, Tamaulipas, Veracruz, Yucatán), Micronesia, Montserrat, Mozambique, Myanmar, Netherlands Antilles (Bonaire, Sint Eustatius), Nicaragua, Oman, Pakistan, Panama, Papua New Guinea, Peru, Philippines, Puerto Rico, Réunion, Saint Kitts and Nevis, Saint Lucia, Saint Vincent and the Grenadines, São Tomé and Príncipe, Saudi Arabia, Senegal, Seychelles, Sint Maarten, Sri Lanka, Suriname, Syria, Taiwan, Tanzania, Thailand, Trinidad and Tobago, Turkey, Turks and Caicos, USA (Florida, Hawaii), US Virgin Islands, Vanuatu, Venezuela, Vietnam, Yemen

Foraging: Argentina, Bahrain, Benin, Bermuda, British Indian Ocean Territory, Brunei, Cambodia, Cameroon, Cape Verde, Chile, Christmas Island, Congo (DRC), Cook Islands, Djibouti, Fiji, Gabon, Ghana, Greece, Guatemala, Guinea, Iraq, Israel, Ivory Coast (Côte d'Ivoire), Kuwait, Liberia, Libya, Maldives, Marshall Islands, Mauritius, Namibia, New Caledonia, Nigeria, Niue, Palau, Qatar, Samoa, Sierra Leone, Singapore, Somalia, South Africa, Sudan, Timor-Leste, Togo, Tonga, United Arab Emirates, Uruguay, Wallis and Futuna, Western Sahara

Vagrant: Algeria, Canada, Italy, Kiribati, Malta, Northern Mariana Islands, Morocco, Nauru, New Zealand, Portugal, Saint Helena, Slovenia, Solomon Islands, Spain, Tokelau, Tuamotu, Tunisia, Tuvalu

Estimated Indigenous (Historical) Ranges (AOO and EOO) (nesting localities intersecting HUC level 12 polygons only):
Est. Indigenous Area of Occupancy (AOO):* 93,934 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 238,159,328 sq. km

Size (Max SCL): male 108.0 cm, female 141.0 cm (Carr and

Hirth 1962; Hirth 1997); **(Max CCL):** male 118.0, female 153.0 (Hirth 1980; Sanches and Bellini 2002)

IUCN Red List:* Endangered (EN A2bd) (Seminoff 2004). Previously: Endangered (EN) (1996). Subpopulation (Hawaiian): Least Concern (LC) (2019) (Pilcher, Chaloupka, and Woods, 2019). Previously: Least Concern (LC) (2012). Subpopulation (Mediterranean): Near Threatened (NT B2b(ii,iii)) (Broderick, Türkozan, Demetropoulos, Mastrogiacomo, Demetropoulos, Levy, and Leader 2023, 2024)

CITES: Appendix I, as Cheloniidae spp. (1981)

USA ESA:* Endangered or Threatened, varying for different Distinct Population Segments (USFWS 1978)

Synonymy:

Testudo atra Linnaeus 1754:50 (invalid pre-1758 name)

Type locality: "Mari ad insulam Adscensionis" [Ascension] [yellow star].

Type specimen: NRM s/n (formerly MAF s/n), holotype, possibly lost, included in composite syntype series of *Testudo mydas* Linnaeus 1758, see Andersson (1900) and Wallin (1985), who identified it as *Chelonia mydas*.

Etymology:* Latin, *atra*, black; referring to the color of the shell.

Comment: Description cited as sourced from Linnaeus (1749b:284,n.7, *T. mydas γ*), a specimen in MAF, as well as Seba (1734:pl.79,f.4–6) and Grew (1681:pl.3,f.4). Unavailable pre-1758 binomial name.

Testudo mydas Linnaeus 1758:197

Chelonia mydas, Chelone mydas, Caretta mydas, Mydas mydas, Mydasea mydas, Chelonia (Euchelonia) midas, Euchelonia mydas, Megemys mydas, Chelonia mydas mydas

Type locality: "insulas Pelagi: insulam Adscensionis." [Ascension] [yellow star]. Emended to "Insel Ascension" by Mertens and Müller (1928:23).

Type specimens: UPSZTY 19, 26, 231 (formerly UUZM), syntypes (3), all three listed as *T. mydas* by Linné and Thunberg (1780) and Thunberg (1828), no. 19 listed by Lönnberg (1896) as the type, but all later identified by Wallin (1985) as = *Testudo caretta* Linnaeus (= *Caretta caretta*), see also Wallin (2001); NRM 5000, 6880–82, possible syntypes (4), see Andersson (1900) and Wallin (1985), listed on NRM website as = *Chelonia mydas*; UUZM 20, listed as "type" by Stuart (1963), Smith and Smith (1980), and Cogger et al. (1983), is a syntype of *Testudo geometrica* Linnaeus (= *Psammobates geometricus*), but the specimen is actually a *Geochelone elegans*, see Wallin (1977, 2001).

Etymology:* Greek, *mydos*, dampness, wetness; referring to its aquatic habitat.

Comment: Description by Linnaeus (1758) of three different varieties of *T. mydas*, cited as sourced from the following: *T. mydas α*: Linnaeus (1749a:138,n.22, *T. mydas α*; 1754:50, *T. atra*), Osbeck (1757:293), Gesner (1558:1131["78"], *T. marina*), Aldrovandi (1637:712,pl.714), Grew (1681:38,pl.3,f.4), Olearius (1674:25,pl.XVII,f.1), Bradley (1721:pl.IV,f.IV); *T. mydas β*: Linnaeus (1749a:137,n.21, *T. mydas β*), Seba (1734:pl.80,f.9), Marggraf (1648:241), Ray (1693:256); *T. mydas γ*: Linnaeus (1749b:284["287"],n.7, *T. mydas γ*; 1754:50, *T. atra*), Seba (1734:pl.79,f.4–6).

Testudo macropus Walbaum 1782:112 (unavailable name)

Euchelys macropus, Chelone macropus

Type locality: Not designated.

Type specimen: Not located, type specimen figured in Edwards (1751:pl.206,top) and Seligmann (1764:pl.101, top).

Etymology:* Greek, *makros*, big, large, and *pus*, foot, leg; referring to the large, flipper-shaped legs.

Comment: Description cited as sourced from Seligmann (1764:pl.101), the same figure as Edwards (1751:pl.206). Unavailable name in a non-binomial work, see Wermuth (1956).

Testudo viridis Schneider 1783:299

Chelonia viridis, Chelone viridis, Chelonia (Mydas) viridis, Mydas viridis, Chelonia mydas viridis

Type locality: Not designated. Restricted to "Charleston, South Carolina" [USA] [green star] by Smith and Taylor (1950a:360).

Etymology:* Latin, *viridis*, green, emerald; referring to the greenish color tinge of the internal body fat.

Type specimen: Not located, type specimen figured (pl.opp.364).

Testudo japonica Thunberg 1787:178

Testudo japanica, Chelonia japonica, Chelonia mydas japonica

Type locality: "Japan" [blue star, location estimated, possibly nr. Dajima, Nagasaki, Kyushu].

Type specimen: UUZM, holotype, listed by Thunberg (1828), possibly lost, not listed by Wallin (2001); type specimen figured (pl.7,f.1).

Etymology:* Toponym for Japan.

Testudo marina vulgaris Lacepède 1788:54, synopsis[table] (*nomen suppressum*) (09:6)

Type locality: "contrées équatoriales."

Type specimen: Possibly MNHN, not located, type specimen figured (pl.opp.54).

Etymology:* Latin, *vulgaris*, common or usual; referring to the apparent commonness of the species.

Comment: Name suppressed by ICZN (2005a) as published in a rejected and invalid non-binomial work, see Savage (2003).

Testudo viridisquamosa Lacepède 1788:92, synopsis[table] (*partim, nomen dubium et suppressum*) (09:6)

Testudo viridi-squamosa

Type locality: "la mer du Sud, auprès du Cap Blanco, de la nouvelle Espagne...aussi dans le golfe du Mexique...&...les rivages chauds du nouveau monde." Restricted to "Bocas del Toro, Panama, Golf von Mexico" [orange star] by Wermuth (1956:405); and to "the island of Blanquilla in the Caribbean" [Venezuela] [brown star] by Brongersma (1961:23).

Type specimen: Possibly MNHN, not located.

Etymology:* Latin, *viridis*, green, emerald, and *squamosa*, scaly; referring to the allegedly somewhat green color of the carapacial scutes.

Comment: Name suppressed by ICZN (1963), see Wermuth (1956) and Mertens and Wermuth (1961), and by ICZN (2005a) as published in a rejected and invalid non-binomial work, see Savage (2003). Name synonymized *in partim* under *Lepidochelys kempii* and *Chelonia mydas*.

Testudo viridisquamosa Bonnaterre 1789:20 (*partim, nomen dubium*)

Testudo viridi-squamosa

Type locality: "La mer du Sud, le golfe du Mexique, les rivages du Nouveau-Monde."

Type specimen: Possibly MNHN, not located.

Etymology:* Latin, *viridis*, green, emerald, and *squamosa*, scaly; referring to the allegedly somewhat green color of the carapacial scutes.

Comment:* Name synonymized *in partim* under *Lepidochelys kempii* and *Chelonia mydas*.

Testudo macropus Gmelin 1789:1038

Euchelys macropus, Chelone macropus

Type locality: Not designated.

Type specimen: Not located, type specimen figured in Edwards (1751:pl.206,top) and Seligmann (1764:pl.101,top).

Etymology:* Greek, *makros*, big, large, and *pus*, foot, leg; referring to the large, flipper-shaped legs.

Comment: Description cited as sourced from Walbaum (1782:112), which was based on Seligmann (1764:pl.101), the same figure as Edwards (1751:pl.206).

Testudo bomarii Meyer 1790:82 (*partim, nomen dubium et oblitum*) (09:8)

Etymology:* Eponym honoring Jacques-Christophe Valmont de Bomare (1731–1807), France; botanist and naturalist who wrote an influential encyclopedia of natural history in the 1760s.

Comment: Unjustified replacement name for *viridisquamosa*. Name synonymized *in partim* under *Lepidochelys kempii* and *Chelonia mydas*.

Testudo chloronotos Bechstein 1800:107 (33)*

Testudo chloronotus

Type locality:* Not designated. Restricted to "the island of Blanquilla in

the Caribbean" [Venezuela] [brown star] by Brongersma (1961:23).
Type specimen: Not designated.
Etymology:* Greek *chloros* = green, pale green, and *notos*, the back; referring to allegedly somewhat greenish scutes on the shell.

Testudo rugosa Van-Ernest *in* Daudin 1801:37 (senior homonym, not = *Testudo rugosa* Shaw 1802 [= *Trachemys decussata decussata* or *Trachemys terrapen*])
Type locality: "la mer des Indes...près de la ligne à environ trois dégrés des îles Maldives" [light gray star, location estimated].
Type specimen: Possibly MNHN, not located.
Etymology:* Latin, *ruga*, wrinkle, ridge, fold; referring to the ridged shell as described for this species.

Testudo cepediana Daudin 1801:50
Type locality: Not known.
Type specimen: Possibly MNHN, not located, type specimen figured (pl.17,f.1).
Etymology:* Eponym honoring Baron B.G.E. de Lacepède (or La Cépède) (1756–1825); France; naturalist, herpetologist, politician, and curator at the Muséum National d'Histoire Naturelle, Paris.

Chelonia virgata Schweigger 1812:291
Caretta virgata, Chelonia (Mydas) virgata, Mydas virgata, Chelone virgata
Type locality: "mari sub zona torrida." Restricted to "Bermuda Islands" [dark gray star] by Smith and Taylor (1950a:315).
Type specimens: MNHN 7900, 7914–15, 8014, 9376, 9539, syntypes (6), see Ceríaco and Bour (2012).
Etymology:* Latin, *virgata*, striped; referring to the striped pattern on the shell scutes.

Caretta cepedii Merrem 1820:18 (*nomen novum*)
Etymology:* Eponym honoring Baron B.G.E. de Lacepède (or La Cépède) (1756–1825); France; naturalist, herpetologist, politician, and curator at the Muséum National d'Histoire Naturelle, Paris.
Comment: Unjustified emendation or error for *cepediana*.

Caretta esculenta Merrem 1820:18
Chelonia esculenta
Type locality: "Oceano Atlantico."
Type specimen: Not known or located.
Etymology:* Latin, *esculenta*, edible; referring to the fact that this species is edible and commonly consumed by humans.

Caretta thunbergii Merrem 1820:19 (*nomen novum*)
Comment: Unjustified replacement name for *japonica*.
Etymology:* Eponym honoring Carl Peter Thunberg (1743–1828), Sweden; naturalist and curator of the Linnaean collection in Uppsala.

Chelonia castanea Eschscholtz 1829a:11 (*nomen oblitum*) (14:5, 17:7)
Type locality: "karaibischen Meere; küste von Surinam" [tourmaline star, location estimated].
Type specimens: Originally in MZT, syntypes (3), transferred to ZMMU, now apparently lost, see Flores-Villela et al. (2016).
Etymology:* Latin, *castaneus*, chestnut color; referring to the brownish color of the carapace.

Chelonia grisea Eschscholtz 1829a:13 (14:4, 17:7)
Chelonia griseam
Type locality: "kaspische Meer" [Caspian Sea] [orange triangle, in error].
Type specimen: Originally in MZT, holotype, transferred to ZMMU, now apparently lost, see Flores-Villela et al. (2016).
Etymology:* Latin, *grisea*, ashen, gray; referring to the yellowish-gray shell color.

Chelonia maculosa Cuvier 1829:13
Chelone maculosa
Type locality: Not designated. Restricted to "Ascension Island" [yellow star] by Smith and Taylor (1950a:315).
Type specimen: Possibly MNHN, not located.
Etymology:* Latin, *macula*, spot, *maculosa*, rich in spots; referring to the carapacial markings.

Chelonia lachrymata Cuvier 1829:13
Type locality: Not designated.
Type specimen: Possibly MNHN, not located.
Etymology:* Latin, *lacrima*, tear, tears; probably referring to the tear-like spots on the carapace scutes.

Chelonia midas Wagler 1830b:133 (*nomen novum*)
Comment: Unjustified emendation or error for *mydas*.
Etymology:* Greek, *mydos*, dampness, wetness; referring to its aquatic habitat.

Chelonia bicarinata Lesson 1831b:301
Type locality: "l'Océan atlantique."
Type specimen: Possibly MNHN, not located.
Etymology:* Latin, *bi-*, two, and *carinata*, keeled, ridged; referring to the two carapacial ridges.

Chelonia marmorata Duméril and Bibron 1835:546
Chelone marmorata
Type locality: "l'île de l'Ascension" [yellow star].
Type specimens: MNHN 7878 and 9535, syntypes (2).
Etymology:* Latin, *marmor*, marble, and *-atus*, provided with; referring to the marbled appearance of the carapace.

Chelonia formosa Girard 1858:456
Type locality: "Feejee Islands" [Fiji] [green triangle].
Type specimen: USNM 257183 (formerly USNM Osteo 12386), see Cochran (1961) and Reynolds et al. (2007).
Etymology:* Latin, *formosa*, beautiful, splendid; referring to the attractive carapace markings.

Chelonia tenuis Girard 1858:459
Type locality: "Honden Island, Paumotu Group; Tahiti and Eimo; Rosa Island." Rosa Island identified by Hirth (1980:1) as Rose Atoll, American Samoa; restricted to "Rosa Island [Rose Island, Rose Atoll, American Somoa]" [American Samoa] [blue triangle] by Reynolds et al. (2007:10).
Type specimen: USNM 257184 (formerly USNM Osteo 12390), holotype, see Cochran (1961) and Reynolds et al. (2007).
Etymology:* Latin, *tenuis*, thin, fine; referring to the allegedly thinner shell bones and scutes of this turtle.

Chelonia albiventer Nardo 1864:1420
Type locality: "Adriatico...prossimata del porto di Malamocco" [Italy] [pink triangle].
Type specimen: MSNVE s/n, holotype, see Hirth (1980).
Etymology:* Latin, *albus*, white, and *ventris*, ventral surface; referring to the white plastron.

Chelonia agassizii Bocourt 1868:122 (07:4)
Chelonia mydas agassizii
Type locality: "embouchure du Nagualate...Pacifique (Guatémala)" [mouth of Rio Nagualate, Pacific coast of Guatemala] [light gray triangle].
Type specimen: MNHN 9537, holotype, see Iverson (1992) and Bour (2006b); MNHN 1871–72 erroneously listed as syntypes by Stuart (1963), Cogger et al. (1983), and King and Burke (1989).
Etymology:* Eponym honoring Louis Agassiz (1807–1873), Switzerland / USA; herpetologist at Harvard University who founded the Museum of Comparative Zoology.

Chelonia maculata Lidth de Jeude *in* Gray 1873j:94 (*nomen nudum*)
Type locality: "Ocean, Guiana?" [Guyana] [dark gray triangle].
Type specimen: NHMUK 1867.4.2.186, see Gray (1873j).
Etymology:* Latin, *macula*, spot; referring to the spots on the carapace scutes.

Chelonia picta Lidth de Jeude *in* Gray 1873j:95 (*nomen nudum*)
Type locality: "Guiana?" [Guyana] [dark gray triangle, location estimated].
Type specimen: NHMUK 1866.8.14.236, see Gray (1873j).
Etymology:* Latin, *picta*, painted; referring to the bright colors of this turtle.

Chelonia lata Philippi 1887:84
Type locality: "Valparaiso..[&]..Insel Chiloe" [Chile] [white triangle].
Type specimen: MNHNC s/n, syntypes (3), apparently lost, discussed by Ortiz and Nuñez (1986); MNHNC 100201, erroneously listed as "holotype" by Hirth (1980).
Etymology:* Latin, *lata*, broad, wide; referring to the wider carapace of this turtle than that of *Chelonia mydas*.

Chelonia mydas carrinegra Caldwell 1962:4
Type locality: "waters adjacent to Isla Angel de la Guarda, Bahia de Los Angeles, central Gulf of California, Mexico" [purple triangle].

Type specimen: LACM 1696, holotype, see Cogger et al. (1983).
Etymology:* Eponym honoring Archie F. Carr, Jr. (1909–1987), USA; herpetologist, conservationist, and turtle researcher at the University of Florida, and Vernacular (Spanish), *negra*, black; referring to the typically black shell color.

Testudo nigrita Tamayo 1962:358 (junior homonym and *nomen nudum*, not = *Testudo nigrita* Duméril and Bibron 1835 [= *Chelonoidis niger porteri*])
Etymology:* Latin, *nigrita*, blackish; referring to relatively dark overall color of the shell.

Natator McCulloch 1908 (12:8)

Flatback Sea Turtles
(includes 1 species, 1 taxon)
Natator McCulloch 1908:127
Type species: *Natator tessellatus* McCulloch 1908 [= subjective synonym of *Chelonia depressa* Garman 1880], by original monotypy.
Etymlogy:* Latin, *natator*, the swimmer; referring to the aquatic habitat.

Natator depressus (Garman 1880)

Flatback, Flatback Sea Turtle




Carmen Pilcher / Shoalwater Bay, Queensland, Australia





left: Colin J. Limpus / Mon Repos, nr. Bundaberg, Queensland, Australia / juvenile
right: Carmen Pilcher / Shoalwater Bay, Queensland, Australia




(yellow dots = nesting sites; star = type locality;
blue lines delimit purple Regional Management Units)

Nesting: Australia (Northern Territory, Queensland, Western Australia)
Foraging: Indonesia (Papua), Papua New Guinea (Southern)
Vagrant: Indonesia (Java, Lesser Sundas), Timor-Leste
Estimated Indigenous (Historical) Ranges (AOO and EOO) (nesting localities intersecting HUC level 12 polygons only):
Est. Indigenous Area of Occupancy (AOO):* 19,999 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 3,840,045 sq. km
Size (Max SCL): male 88.4 cm; female 93.0 cm [both estimated]; **(Max CCL):** male 93.1 cm, female 100.0 cm (Limpus 2007; C. Limpus, unpubl. data; Whittock et al. 2014)
IUCN Red List: Data Deficient (DD) (RLSPS 1996). Previously: Vulnerable (VU) (1996)
CITES: Appendix I, as Cheloniidae spp. (1981). Previously: Appendix II (1975)
Synonymy:
Chelonia depressa Garman 1880:124
Chelonia depressus, Natator depressus, Natator depressa
Type locality: "East Indies and North Australia." Restricted to "North Australia" by Loveridge (1934:261) [yellow star, location estimated].
Type specimen: MCZ 4473, lectotype, designated by Loveridge (1934:261).
Etymology:* Latin, *de-*, down, and *pressus*, pressed; referring to the depressed flattened shell.

Natator tessellatus McCulloch 1908:127
Type locality: "near Port Darwin, North Australia" [yellow star].
Type specimen: AMS R4158, holotype, apparently lost, see Cogger et al. (1983) and Shea and Sadlier (1999).
Etymology:* Latin, *tessela*, cube, and *-atus*, provided with; referring to the large symmetrical scales on top of the head.

Family Dermochelyidae Baur 1888a (12:9) (30, 34)*

Leatherback Sea Turtles
(includes 1 genus, 1 species, 1 taxon)
Sphargidae Gray 1825:212
Sphargidina Bonaparte 1831:64
Dermatochelydae Fitzinger 1843:30
Sphragidoidae Agassiz 1847:348
Athecae Cope 1871:235
Dermochelydidae Baur 1888a:422
Athecata Lydekker 1889:223
Dermochelyidae Lydekker 1889:223



Dermochelyidae Species Richness

Dermochelys Blainville 1816

Leatherback Sea Turtles
(includes 1 species, 1 taxon)
Chelonias Rafinesque 1814:66 (*nomen oblitum*)
Etymology:* Greek, *chelone*, turtle.
Comment:* Formally affirmed and validated as a *nomen oblitum* by Frétey (2022).

Chelyra Rafinesque 1815:74 (*nomen nudum*)
Etymology:* Greek, *chelys*, turtle, and *lyra*, lyre or lute; referring to the body shape reminiscent of a lyre.

Dermochelys Blainville 1816:119 ["111"] (*nomen protectum*)
Type species: *Dermochelys coriacea* [= *Testudo coriacea* Vandellius 1761], by subsequent monotypy by Boulenger (1889:10), in accordance with ICZN Article 67.2.2.
Etymology:* Greek, *derma*, skin, and *chelys*, turtle; referring to the leathery skin covering the carapace.
Comment:* Name conserved by ICZN (1926, 1956b), see Frétey (2022).

Sphargis Merrem 1820:19 (*nomen suppressum*)
Type species: *Sphargis mercurialis* Merrem 1820 [= subjective synonym of *Testudo coriacea* Vandellius 1761], by original monotypy.
Etymology:* Greek, *spargosis*, to swell; perhaps referring to the large swollen body, or possibly *spharagos*, bursting with noise, perhaps referring to the body shape of a lyre or lute producing sound.
Comment:* Name suppressed by ICZN (1956b), see Frétey (2022).

Coriudo Fleming 1822:271 (*nomen suppressum*)
Type species: *Coriudo coriacea* [= *Testudo coriacea* Vandellius 1761], by original monotypy.
Etymology:* Latin, *corium*, leather or skin, and *testudo*, turtle; referring to the leathery skin covering the carapace.
Comment:* Name suppressed by ICZN (1956b), see Frétey (2022).

Scytina Wagler 1828:861 (*nomen novum*)
Etymology:* Greek, *skytinos*, leathery; referring to the leathery skin covering the carapace.

Dermochelis Cuvier 1829:14 (*nomen novum et suppressum*)
Etymology:* Greek, *derma*, skin, and *chelys*, turtle; referring to the leathery skin covering the carapace.
Comment:* Name suppressed by ICZN (1956), see Frétey (2022).

Dermatochelys Wagler 1830b:133 (*nomen novum et suppressum*) (17:9)
Etymology:* Greek, *derma*, skin, and *chelys*, turtle; referring to the leathery skin covering the carapace.
Comment:* Name suppressed by ICZN (1956b), see Frétey (2022).

Chelyra Rafinesque 1832:64
Type species: *Chelyra coriacea* [= *Testudo coriacea* Vandellius 1761], by original designation.
Etymology:* Greek, *chelys*, turtle, and *lyra*, lyre or lute; referring to the body shape reminiscent of a lyre.

Sphragis Agassiz 1847:342 (*nomen novum*)
Etymology:* Greek, *spargosis*, to swell; perhaps referring to the large swollen body, or possibly *spharagos*, bursting with noise, perhaps referring to the body shape of a lyre or lute producing sound.
Comment:* Unjustified emendation of *Sphargis*.


Dermochelys coriacea (Vandellius 1761) (12:9, 14:6, 14:7, 17:10) (35)*
Leatherback, Leatherback Sea Turtle



Chris Johnson / CRM 3 / Juno Beach, Palm Beach Co., Florida



Anders G.J. Rhodin / *left*: Guancaste, Costa Rica / *right*: Culebra, Puerto Rico



(yellow dots = nesting sites; blue lines delimit purple Regional Management Units; stars = type localities)

Nesting: Angola, Anguilla, Antigua and Barbuda, Aruba, Australia (Northern Territory), Bahamas, Bangladesh, Barbados,

Benin, Brazil, British Virgin Islands, Cameroon, Colombia, Congo (ROC), Costa Rica, Cuba, Curacao, Dominica, Dominican Republic, Ecuador, El Salvador, Equatorial Guinea, French Guiana, Gabon, Ghana, Grenada, Guadeloupe, Guatemala, Guyana, Haiti, Honduras, India, Indonesia (Java, Papua, West Papua), Ivory Coast (Côte d'Ivoire), Jamaica, Malaysia, Martinique, Mexico (Baja California Sur, Guerrero, Jalisco, Michoacán, Oaxaca), Mozambique, Netherlands Antilles (Bonaire, Sint Eustatius), Nicaragua, Panama, Papua New Guinea (Northern), Puerto Rico, Saint Kitts and Nevis, Saint Lucia, Saint Vincent and the Grenadines, São Tomé and Príncipe, Sierra Leone, Sint Maarten, Solomon Islands, South Africa, Sri Lanka, Suriname, Thailand, Togo, Trinidad and Tobago, USA (Florida, Texas), US Virgin Islands, Vanuatu, Venezuela, Vietnam

Foraging: Albania, Algeria, Argentina, Belize, Brunei, Cambodia, Canada (British Columbia, New Brunswick, Newfoundland, Nova Scotia), Chile, China, Comoros, Congo (DRC), Croatia, Cyprus, Egypt, Fiji, France, Gambia, Great Britain, Greece, Guinea, Guinea-Bissau, Ireland, Israel, Italy, Japan, Kenya, Kiribati, Lebanon, Liberia, Libya, Madagascar, Malta, Marshall Islands, Mauritania, Mauritius, Micronesia, Montenegro, Monaco, Morocco, Myanmar, Namibia, Nauru, New Zealand, Nigeria, North Korea, Palau, Peru, Philippines, Portugal, Russia, Samoa, Senegal, Seychelles, Slovenia, South Korea, Spain, Syria, Taiwan, Tanzania, Tonga, Tunisia, Turkey, Turks and Caicos, Tuvalu, Uruguay, USA (Alaska, California, Connecticut, Delaware, Georgia, Hawaii, Maine, Maryland, Massachusetts, New Jersey, New York, North Carolina, Oregon, Rhode Island, South Carolina, Washington)

Vagrant: Bahrain, Denmark, Djibouti, Eritrea, Iceland, Iran, Iraq, Kuwait, Maldives, Netherlands, Norway, Oman, Pakistan, Qatar, Saudi Arabia, Somalia, Sudan, Sweden, United Arab Emirates, Yemen

Estimated Indigenous (Historical) Ranges (AOO and EOO) (nesting localities intersecting HUC level 12 polygons only):
Est. Indigenous Area of Occupancy (AOO):* 59,468 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 131,516,216 sq. km

Size (Max SCL):* male 203.2 cm [estimated], female 199.9 cm [estimated] (Firdous 1989; Chu Cheong 1990), conversion equation used for estimates (in cm): SCL = (CCL/1.04) - 2.04 (Tucker and Frazer 1991); **(Max CCL):** male 213.4 cm, female 210.0 cm (Firdous 1989; Chu Cheong 1990; Eckert et al. 2012). Incorrect previously recorded alleged maximum measurements: SCL: 226.0 cm [estimated], CCL: 256.5 cm (Eckert and Luginbuhl 1988; see Davenport et al. 1990 and McClain et al. 2015 for corrections), female 252.0 cm (Girondot and Fretey 1996; Eckert et al. 2012; see Girondot et al. 2021 and M. Girondot, unpubl. data, for corrections)

IUCN Red List: Vulnerable (VU A2bd) (Wallace, Tiwari, and Girondot 2013); Subpopulations: East Pacific Ocean: Critically Endangered (CR A2bd+4bd) (2013); Northeast Indian Ocean: Data Deficient (DD) (2013); Northwest Atlantic Ocean: Endangered (EN) (Northwest Atlantic Leatherback Working Group 2019); Southeast Atlantic Ocean: Data Deficient (DD) (2013); Southwest Atlantic Ocean: Critically Endangered (CR D) (2013); Southwest Indian Ocean: Critically Endangered (CR C2a(ii)) (2013); West Pacific Ocean: Critically Endangered (CR A2bd+4bd) (2013). Previously: Critically Endangered (CR) (2000); Endangered (EN) (1996); Subpopulations: Northwest Atlantic Ocean: Least Concern (LC) (2013)

CITES: Appendix I (1977). Previously: Appendix II (1975)

USA ESA:* Endangered (USFWS 1970)

Synonymy:

Testudo coriacea Vandellius 1761:1
Chelone coriacea, Chelonia coriacea, Dermochelys coriacea, Coriudo coriacea, Scytina coriacea, Sphargis coriacea, Dermatochelys coriacea, Dermochelys coriacea coriacea, Chelyra coriacea
Type locality: "maris Tyrrheni oram in agro Laurentiano" [Italy] [yellow star]. Restricted to "Palermo, Sicily" [Italy] [orange star, in error] by Smith and Taylor (1950a:315, 1950b:13); to "la côte romaine (Italie), Mer Tyrrhénienne, Méditerranée occidentale" [Italy] by Fretey and Bour (1980:198); and to "Laurentum, between Lido di Ostia and Tor Paterno, shore of the Tyrrhenian Sea, Italy" [yellow star] by Bour and Dubois (1984a:359).
Type specimen: MZP (previously ZMUP) s/n, holotype, identified by Fretey and Bour (1980), discussed by Rhodin and Smith (1982), Bour and Dubois (1984a), and Smith and Rhodin (1986).
Etymology:* Latin, *corium*, leather or skin; referring to the leathery skin covering the carapace.
Comment: The author's name as given on his 1761 paper is both Vandellius and Vandelli; we and others have previously used the Vandelli spelling, but Frétey (2022) points out that Vandellius is more correct, and we have now therefore switched usage.

Testudo arcuata Catesby 1771:40
Type locality: Not designated. Restricted to "Küsten-Gebiete von Carolina und Florida" [USA] [green star, location estimated] by Mertens and Wermuth (1955:386).
Type specimen: Not designated.
Etymology:* Latin, *arcus*, an arch; referring to the arched carapace of this taxon.

Testudo lyra Lacepède 1788:111, synopsis[table] (*nomen suppressum*) (09:6)
Type locality: "Méditerranée" [no star].
Type specimen: Possibly MNHN, not located, type specimen figured opp.111).
Etymology:* Greek, *lyra*, lyre or lute; referring to the body shape reminiscent of a lyre.
Comment: Name suppressed by ICZN (2005a) as published in a rejected and invalid non-binomial work, see Savage (2003).

Testudo lyra Bonnaterre 1789:22
Chelonia lyra
Type locality: "La Méditerranée, l'Océan, les côtes du Pérou, du Mexique, & la plupart de celles d'Afrique, qui sont sous la Zone torride" [no stars]
Type specimen: Possibly MNHN, not located, type specimen figured in Lacepède (1788:opp.111).
Etymology:* Greek, *lyra*, lyre or lute; referring to the body shape reminiscent of a lyre.

Testudo tuberculata Pennant *in* Schoepff 1801:123
Sphargis tuberculata, Dermochelydis tuberculata
Type locality: Not designated. Restricted to "Palermo, Sicily" [Italy] [orange star] by Smith and Smith (1980:244).
Type specimens: Not located, type specimens figured in Pennant (1771:pl.10,f.4) and Schoepff (1801:pl.28).
Etymology:* Latin, *tuberculum*, small hump, nodule; referring to the humplike nodules on the hatchling carapace.

Chelonias lutaria Rafinesque 1814:66
Type locality: "Sicil." [Sicily, Italy] [orange star, location estimated].
Type specimen: Not designated.
Etymology:* Latin, *lutum*, mud; possibly referring to the erroneous assumption that the species resides predominantly on the sea floor.

Sphargis mercurialis Merrem 1820:19 (*nomen suppressum*)
Type locality: "Mari mediterraneo et Oceano atlantico" [no stars]
Type specimen: Not known or located.
Etymology:* Mythonym for *Mercury*, the Greek god considered to be the inventor of the lyre musical instrument; referring to the shell shaped somewhat like a lyre.
Comment: Name suppressed by ICZN (1956b).

Dermochelis atlantica LeSueur *in* Cuvier 1829:14 (*nomen nudum*)
Dermochelys atlantica, Dermatochelys atlantica

Etymology:* Toponym for the Atlantic Ocean.

Dermatochelys porcata Wagler 1830c:expl.tab.f.1 (14:6, 17:9)
Type locality: Not designated.
Type specimen: Not known or located.
Etymology:* Latin, *porcata*, ridged; referring to the several clearly thickened longitudinal ridges on the carapace.

Testudo coriacea marina Ranzani 1832:3 (14:7)
Type locality: Not designated.
Type specimen: Not located, type specimen figured (pl.4).
Etymology:* Latin, *marina*, living in the sea, marine; referring to the habitat of this species.

Sphargis coriacea schlegelii Garman 1884:303
Dermochelys schlegelii, Dermochelys coriacea schlegelii, Sphargis schlegelii
Type locality: "Tropical Pacific and Indian Oceans." Restricted to "Guaymas, Sonora, Mexico" [blue star] by Smith and Taylor (1950a:344, 1950b:13).
Type specimens: Not designated, description included specimens figured in "Temminck and Schlegel (1838:pl.1-3)" [actually Temminck and Schlegel (1834:pl.1-3)]; Pritchard (1980a) erroneously designated type specimens figured in plates 1-2 in Siebold (1826), a publication which contains no references to turtles and has no figures; type specimen figured in Temminck and Schlegel (1834:pl.1) designated lectotype by TTWG (2021:102).
Etymology:* Eponym honoring Hermann Schlegel (1804–1884), Germany / Netherlands; ornithologist and herpetologist who worked in Austria and the Netherlands and was director of the Rijksmuseum van Natuurlijke Historie, Leiden.

Sphargis angusta Philippi 1899:730
Dermatochelys angusta
Type locality: "cerca de Tocopilla" [Chile] [pink star]. Emended to "Tocopilla (22º04'S; 70º12'O)" [Chile] by Ortiz and Nuñez (1986:7).
Type specimen: MNHNC 1515, holotype, see Ortiz and Nuñez (1986).
Etymology:* Latin, *angusta*, narrow, short; referring to the allegedly narrower head and carapace as compared to *Dermochelys coriacea*.

Superfamily Chelydroidea Gray 1831d (21:3, 21:44) (29)*

Chelydroid Turtles
(includes 3 families, 7 genera, 40 species, 48 taxa)
Chelydrae Gray 1831d:4
Chelydroidae Agassiz 1857a:341,409
Chelydroidea Baur 1893b:673

Family Chelydridae Gray 1831d (09:3)

Snapping Turtles
(includes 2 genera, 5 species, 5 taxa)
Chelydrae Gray 1831d:4
Chelidridae Swainson 1839:116
Chelydroidae Agassiz 1857a:341,409
Chelydradae Gray 1869a:178
Chelydraina Gray 1870c:63
Chelydrinae Cope 1872:22
Chelydridae Cope 1875b:16



Chelydridae Species Richness

Chelydra Schweigger 1812 (07:2)

American Snapping Turtles
(includes 3 species, 3 taxa)

Chelydra Schweigger 1812:292
Type species: *Chelydra serpentina* Schweigger [= *Testudo serpentina* Linnaeus 1758], by subsequent designation by Fitzinger (1843:29).
Etymology:* Greek, *chelydros*, water serpent; referring to the long, agile, snake-like neck and tail.

Cheliurus Rafinesque 1815:75 (*nomen nudum*)
Etymology:* Greek, *chelys*, turtle, and *oura*, tail, rump; referring to the long tail.

Chelonura Fleming 1822:270 (senior homonym, not = *Chelonura* Rafinesque 1832 = *Cylindraspis*)
Type species: *Chelonura serpentina* [= *Testudo serpentina* Linnaeus 1758], by original monotypy.
Etymology:* Greek, *chelone*, turtle, and *oura*, tail, rump; referring to the long tail.

Ophichelone Jarocki 1822:21
Type species: *Ophichelone serpentina* [= *Testudo serpentina* Linnaeus 1758], by original monotypy.
Etymology:* Greek, *ofis*, snake, serpent, and *chelone*, turtle; referring to the long, agile, snake-like neck and tail.

Rapara Gray 1825:210
Type species: *Rapara serpentina* Gray [= *Testudo serpentina* Linnaeus 1758], by original monotypy.
Etymology:* Anonym, *rapara*, possibly a fabricated name, nearly a palindrome; or possibly an incorrect use of *raparapa*, a Vernacular name later used as an alternate species name for *Chelus fimbriata*.

Saurochelys Latreille 1825:92
Type species: *Saurochelys* "Tortue à longue queue" [= *Testudo serpentina* Linnaeus 1758], by original monotypy.
Etymology:* Greek, *sauros*, lizard, and *chelys*, turtle; referring to the lizard-like long neck and tail.

Cheliurus Rafinesque 1832:64
Type species: *Cheliurus serpentina* [= *Testudo serpentina* Linnaeus 1758], by original monotypy.
Etymology:* Greek, *chelys*, turtle, and *oura*, tail, rump; referring to the long tail.

Emysaurus Duméril and Bibron 1835:348
Type species: *Emysaura serpentina* [= *Testudo serpentina* Linnaeus 1758], by original monotypy.
Etymology:* Greek, *emys*, turtle, and *sauros*, lizard; referring to the lizard-like long neck and tail.

Chelidra Swainson 1839:116
Type species: *Chelidra serpentina* [= *Testudo serpentina* Linnaeus 1758], by original monotypy.
Etymology:* Greek, *chelydros*, water serpent; referring to the long, agile, snake-like neck and tail.
Comment: Unjustified emendation or error for *Chelydra*.

Emydosaurus Agassiz 1847:137
Etymology:* Greek, *emydos*, turtle, and *sauros*, lizard; referring to the lizard-like long neck and tail.
Comment:* Unjustified emendation of *Emysaurus*.

Devisia Ogilby 1905:11
Type species: *Devisia mythodes* Ogilby 1905 [= subjective synonym of *Testudo serpentina* Linnaeus 1758], by original monotypy.
Etymology:* Eponym honoring Charles Walter De Vis (1829–1915), United Kingdom / Australia; director of the Queensland Museum in Brisbane, who described several fossil turtle species.


Chelydra acutirostris Peters 1862
South American Snapping Turtle, *Tortuga Pímpano*



Katherine Young-Valencia / Quindío, Colombia



left: Katherine Young-Valencia / Quindío, Colombia / male
right: Vivian P. Páez / Colombia / captivity / Cali Zoo



** (orange dots = probable trade or translocated; star = type locality)

Distribution: Colombia (*Antioquia, Caldas, Cauca, Chocó, Córdoba, Nariño, Quindío, Valle del Cauca*), Costa Rica, Ecuador, Honduras, Nicaragua, Peru (?), Panama

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 363,578 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 1,433,152 sq. km

Size (Max SCL):* male 42.7 cm, female 39.1 cm (Medem 1977; Young-Valencia et al. 2014); **(Max CCL):** male 48.0 cm (Grünewald 2008)

IUCN Red List:* Near Threatened (NT, A2bcd+4bcd) (Young-Valencia, Carr, Giraldo, Forero Medina, Botero Botero, and Ortega Guío, in press). Previously: TFTSG Provisional Red List: Near Threatened (NT) (2011, 2018)

Synonymy:

Chelydra serpentina acutirostris Peters 1862:627
Chelydra acutirostris
Type locality: "Guayaquil" [Ecuador] [yellow star].
Type specimen: ZMB 4500, holotype, see Medem (1977) and Fritz et al. (1994).
Etymology:* Latin, *acutus*, sharp; *rostrum*, beak or snout; referring to the sharp projecting snout.


Chelydra rossignonii (Bocourt 1868)
Central American Snapping Turtle, *Servengue, Chiquiguao*



Eduardo Reyes-Grajales / Juárez, Chiapas, Mexico



left: Eduardo Reyes-Grajales / Chiapas, Mexico
right: Craig B. Stanford / Chiapas, Mexico



** (orange dots = probably introduced; star = type locality)

Distribution: Belize, Guatemala, Honduras, Mexico (*Campeche, Chiapas, Oaxaca, Tabasco, Veracruz*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 148,554 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 326,897 sq. km

Size (Max SCL): male 47.0 cm, female 33.0 cm (Legler and Vogt 2013)

IUCN Red List:* Near Threatened (NT, A2cd+4cd) (Reyes-Grajales, Walde, and Munscher, in press). Previously: Vulnerable (VU A2d) (van Dijk, Lee, Calderón Mandujano, Flores-Villela, López-Luna, and Vogt 2007)


Synonymy:

Emysaurus rossignonii Bocourt 1868:121
Chelydra rossignonii, Chelydra serpentina rossignonii
Type locality: "marais de Pansos, près le Rio Polochic (Guatémala)" [yellow star].
Type specimens: MNHN 1501, 1501A, syntypes (2); MNHN 1230, erroneously listed as "type" by Stuart (1963:47), is not a type, see Duméril and Bocourt (1870) and Bour (2005b).
Etymology:* Eponym honoring Jules Rossignon (c.1810–1883), France / Guatemala; scientist, agriculturist (coffee grower), and naturalist who lived and worked in Guatemala.

Chelydra serpentina mexicanae Cope *in* Gray 1870c:64 (*nomen nudum*)
Etymology:* Toponym for Mexico.

Chelydra serpentina (Linnaeus 1758) (08:5)
North American Snapping Turtle, Common Snapping Turtle


Anders G.J. Rhodin / Manchester, Bennington Co., Vermont / female


Anders G.J. Rhodin / Manchester, Bennington Co., Vermont / female


**

(orange dots = introduced or translocated; stars = type localities)

Distribution: Canada (*Manitoba, New Brunswick, Nova Scotia, Ontario, Québec, Saskatchewan*), USA (*Alabama, Arkansas, Colorado, Connecticut, Delaware, Florida, Georgia, Illinois, Indiana, Iowa, Kansas, Kentucky, Louisiana, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Vermont, Virginia, West Virginia, Wisconsin, Wyoming*)

Introduced:* Canada (*British Columbia*), Belgium, China, France, Germany, Hong Kong, Italy, Japan, Mexico, Micronesia, South Korea, Taiwan, USA (*Arizona, California, Nevada, Oregon, Utah, Washington*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 5,364,328 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 9,022,386 sq. km

Size (Max SCL):* male 50.3 cm, female 41.8 cm (Gerholdt and Oldfield 1987; Pritchard 1989; Iverson et al. 1997; Johnston et al. 2012, 2024b)

IUCN Red List:* Least Concern (LC) (Moldowan and van Dijk 2025). Previously: Least Concern (LC) (van Dijk 2012); Least Concern (LC) [Not Listed] (Behler 1996)

CITES:* Appendix II (2023). Previously: Appendix III (USA) (2016)

Synonymy:

Testudo serpentina Linnaeus 1758:199
Chelydra serpentina, Emys serpentina, Chelonura serpentina, Ophichelone serpentina, Rapara serpentina, Saurochelys serpentina, Chelidra serpentina, Cheliurus serpentina, Emysaurus serpentina, Emysaura serpentina, Hydraspis (Chelydra) serpentina, Emysaurus serpentinus, Chelydra serpentina serpentina
Type locality: "Calidis regionibus." Restricted to "Algiriae, Chinae aquis dulcibus" [Algeria and China] by Linnaeus (1766:354); to "Louisiana, Orleans Parish, New Orleans" [USA] [yellow star] by Smith and Taylor (1950a:358, 1950b:21); and invalidly to "vicinity of New York City" [New York, USA] [orange star] by Schmidt (1953:86).
Type specimen: UPSZTY 280 (formerly UUZM 280, MGA 49, and MAF s/n), holotype, see Linnaeus (1764), Thunberg (1828), Holm (1957), and Wallin (2001); not lost as per Andersson (1900), Gibbons et al. (1988), and Uetz et al. (2019).
Etymology:* Latin, *serpent*, snake, and *-ina*, similarity of form; referring to the long snake-like neck and tail.
Comment: Description later cited by Linnaeus (1766:354) as sourced from Linnaeus (1764:36), a work that was actually written in 1754.

Testudo serpentaria Wiedemann 1802:191 (*nomen novum*)
Etymology:* Latin, *serpent*, snake, and *-aria*, similarity of shape; referring to the long snake-like neck and tail.
Comment: Unjustified emendation or error for *serpentina*.

Chelydra lacertina Schweigger 1812:293 (senior homonym, not = *Gypochelys lacertina* Agassiz 1857a = *Macrochelys temminckii*)
Chelydra serpentina lacertina
Type locality: Not known. Restricted to "vicinity of New York City" [New York, USA] [orange star] by Schmidt (1953:86).
Type specimen: MNHN 9580, holotype, see Gibbons et al. (1988).
Etymology:* Latin, *lacerta*, lizard, and *-ina*, similarity of form; referring to the long lizard-like neck and tail.

Testudo serrata Pennant *in* Gray 1830e:14 (junior homonym and *nomen nudum*, not = *Testudo serrata* Daudin 1801 [= *Trachemys scripta scripta*], nor = *Testudo serrata* Shaw 1802 [= *Geoemyda spengleri*])
Etymology:* Latin, *serratus*, saw-shaped, serrated; referring to the posterior carapacial margin and serrated tail.

Testudo longicauda Shaw *in* Gray 1831d:36 (*nomen nudum*)
Etymology:* Latin, *longum*, long, and *cauda*, tail; referring to the long tail.

Chelydra emarginata Agassiz 1857a:417
Type locality: "Mobile and New Orleans." Restricted to "Mobile" [Alabama, USA] [green star] by Schmidt (1953:86).
Type specimens: MCZ 1472, syntype (one of several), others apparently lost, see Gibbons et al. (1988).
Etymology:* Latin, *emarginata*, having a notch; referring to the allegedly wider notch between the marginal scutes above the tail than in *Chelydra serpentina*.

Devisia mythodes Ogilby 1905:11
Type locality: "Fly River, British New Guinea" [Papua New Guinea] [in error].
Type specimen: QM J20207, holotype, see Covacevich (1971).
Etymology:* Greek, *mythos*, fable, legend, and *-odes*, likeness; referring to the highly unusual nature of the presumed geographic origin.

Chelydra laticarinata † Hay 1916a:72 (*nomen suppressum*)
Type locality: "Vero, St. Lucie County, Florida" [USA] [gray star].
Type specimen: USNM (formerly FGS 7094), holotype, fossil, peripheral bone, figured (pl.6,f.6–7).
Geological age: Pleistocene.
Etymology:* Latin, *lati-*, wide or broad, and *carina*, the keel, *carinata*, keeled; referring to the wide carapacial vertebral keel.
Comment: Name suppressed by ICZN (1986), see Smith et al. (1983).

Chelydra sculpta † Hay 1916a:73 (*nomen suppressum*)
Type locality: "Vero, St. Lucie County, Florida" [USA] [gray star].
Type specimen: USNM (formerly FGS 5510), holotype, fossil,

peripheral bone, figured (pl.4,f.7,pl.6,f.8–9).
Geological age: Pleistocene.
Etymology:* Latin, *sculpta*, carved, chiseled; referring to the peripheral bones covered with depressions and grooves in contrast to *Chelydra laticarinata*, described at the same time.
Comment: Name suppressed by ICZN (1986), see Smith et al. (1983).

Chelydra osceola Stejneger 1918:89 (*nomen protectum*) (08:5)
Chelydra serpentina osceola
Type locality: "Clearwater, Pinellas County, Florida" [USA] [blue star].
Type specimen: USNM 10369, holotype, see Cochran (1961) and Reynolds et al. (2007).
Etymology:* Toponym for Osceola, Florida, USA.
Comment: Name conserved by ICZN (1986), see Smith et al. (1983).

Macrochelys Gray 1856a (07:3, 17:2)

Alligator Snapping Turtles
(includes 2 species, 2 taxa)

Macrochelys Gray 1856a:200
Type species: *Macrochelys temminckii* [= *Chelonura temminckii* Troost *in* Harlan 1835], by original monotypy.
Etymology:* Greek, *makros*, large, and *chelys*, turtle; referring to the large size.

Macroclemys Gray 1856b:48 (*nomen novum*)
Etymology:* Greek, *makros*, large, and *klemmys*, turtle; referring to the large size.
Comment: Unjustified replacement name or error for *Macrochelys*.

Gypochelys Agassiz 1857a:248, 413
Type species: *Gypochelys lacertina* Agassiz 1857a [= subjective synonym of *Chelonura temminckii* Troost *in* Harlan 1835], by original monotypy.
Etymology:* Greek, *gypos*, vulture, and *chelys*, turtle; referring to the vulture-like shape of the horny upper jaw.


Macroclemmys Strauch 1862:35 (*nomen novum*)
Etymology:* Greek, *makros*, large, and *klemmys*, turtle; referring to the large size.
Comment: Unjustified replacement name or error for *Macroclemys*.

Macrochelys suwanniensis Thomas, Granatosky, Bourque, Krysko, Moler, Gamble, Suarez, Leone, Enge, and Roman 2014 (14:1, 17:2, 17:3, 17:4, 21:5)

Suwannee Alligator Snapping Turtle


Kevin Enge / White Springs, Suwannee R., Florida


Travis Thomas / Suwannee R., Florida


**
(orange dots = probably introduced or translocated; star = type locality)

Distribution: USA (*Florida*, *Georgia*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 15,800 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 30,617 sq. km
Size (Max SCL):* male 80.1 cm (midline SCL = 71.3 cm), female 52.7 cm (midline SCL = 49.2 cm) (Pritchard 1980; Johnston et al. 2015, 2023; G.R. Johnston, unpubl. data)
IUCN Red List:* Vulnerable (VU C1) (Thomas, Enge, and Johnston 2025). Previously: TFTSG Provisional Red List: Vulnerable (VU) (2018)
CITES:* Appendix II (2023), as part of *M. temminckii*. Previously: Appendix III (USA) (2006)
USA ESA:* Threatened (USFWS 2024a); Species Status Assessment: USFWS (2022b)
Synonymy:
Macrochelys suwanniensis Thomas, Granatosky, Bourque, Krysko, Moler, Gamble, Suarez, Leone, Enge, and Roman 2014:150
Type locality: "Santa Fe River and State Road 235, Alachua County, Florida (29.87872ºN, 82.33619ºW...elev. 23 m)" [USA] [yellow star].
Type specimen: UF 166146, holotype.
Etymology:* Toponym for the Suwannee River, Florida, USA.


Macrochelys temminckii (Troost *in* Harlan 1835) [09:4, 14:1, 17:2, 21:5] [36]*
Western Alligator Snapping Turtle, Alligator Snapping Turtle



James C. Godwin / CCB / lower Tallapoosa R., Elmore Co., Alabama / juvenile



left: James C. Godwin / Tombigbee R., Washington Co., Alabama
right: Eric Munscher / Houston, Texas



(light red = likely extirpated; orange dots = uncertain;
red dots = likely extirpated; stars = type localities)

Distribution:* USA (*Alabama, Arkansas, Florida, Georgia, Illinois, Indiana [extirpated], Iowa [extirpated], Kansas [extirpated], Kentucky, Louisiana, Mississippi, Missouri, Oklahoma, Tennessee, Texas*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 505,589 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 1,406,452 sq. km

Size (Max SCL):* male 77.5 cm (midline SCL = 73.8 cm), female 55.0 cm (midline, not max) (Pritchard 1989; Boundy and Kennedy 2006; Rosenbaum et al. 2023); maximum midline SCL: male 74.0 cm (Rosenbaum et al. 2023)

IUCN Red List:* Endangered (EN A2bcd+4bcd) (Carr, Riedle, Munscher, Pearson, Kessler, and Dreslik 2025). Previously: TFTSG Provisional Red List: Vulnerable (VU) (2011); Vulnerable (VU) (TFTSG 1996)

CITES:* Appendix II (2023). Previously: Appendix III (USA) (2006)

USA ESA:* Proposed Threatened but Not Listed; Species Status Assessment: USFWS (2021c)

Synonymy:

Testudo planitia Gmelin 1789:1045 (junior homonym, not = *Testudo planitia* Meuschen 1778 [= *Pelomedusa subrufa*]; *nomen suppressum*)
Chersine planitia
Type locality: "Surinami" [Suriname] [in error].
Type specimen: Not located, no known figured type specimens.
Etymology:* Latin, *planus*, plane, flat, and *itius*, characterized by; referring to the relatively flat shell.
Comment: Description cited as sourced from Gronovius (1756:86,n.70 and 1763:17,n.76). Name suppressed by ICZN (1963), see Wermuth (1956) and Mertens and Wermuth (1961).

Chelonura temminckii Troost *in* Harlan 1835:158 (*nomen protectum*)
Emysaurus temminckii, Macrochelys temminckii, Macroclemys temminckii, Chelydra temminckii, Gypochelys temminckii, Macrochelys temminckii temminckii
Type locality: "a tributary stream of the Mississippi, which enters that river above Memphis, in west Tennessee" [USA]. Restricted to "Wolf River, Shelby County, Tennessee, USA" [yellow star] by Bour (1987b:343).
Type specimen: RMNH 6166, holotype, designated and discussed by Hoogmoed et al. (2010:4); MNHN A.4540, designated "holotype" by Bour (1987b:343), is apparently not a type (Hoogmoed et al. (2010).
Etymology:* Eponym honoring Coenraad Jacob Temminck (1778–1858), Netherlands; zoologist and director of the Rijksmuseum van Natuurlijke Historie, Leiden.
Comment: Name conserved by ICZN (1963), see Wermuth (1956) and Mertens and Wermuth (1961).

Gypochelys lacertina Agassiz 1857a:414 (junior homonym, not = *Chelydra lacertina* Schweigger 1812 [= *Chelydra serpentina*])
Macrochelys lacertina, Macroclemys lacertina
Type locality: "New Orleans...Mississippi...and...south-western Georgia" [USA] [no stars].
Type specimens: Probably MCZ, syntypes, not located.
Etymology:* Latin, *lacerta*, lizard, and *-ina*, similarity of form; referring to the long lizard-like neck and tail.

Macrochelys apalachicolae Thomas, Granatosky, Bourque, Krysko, Moler, Gamble, Suarez, Leone, Enge, and Roman 2014:151 [14:1, 17:2, 17:3, 17:4, 21:5]
Type locality: "Apalachicola River, Gadsden County, Florida" [USA] [green star].
Type specimen: UF 3998, holotype.
Etymology:* Toponym for the Apalachicola River, Florida.

Family Dermatemydidae Gray 1870e

Dermatemydid River Turtles
(includes 1 genus, 1 species, 1 taxon)
Dermatemydae Gray 1870e:714
Dermatemydidae Baur 1888b:595



Dermatemydidae Species Richness

Dermatemys Gray 1847

Central American River Turtles

(includes 1 species, 1 taxon)

Dermatemys Gray 1847:55

Type species: *Dermatemys mawii* Gray 1847, by original monotypy.

Etymology:* Greek, *derma*, skin, and *emys*, turtle; referring to the skin-like appearance of the shell.

Chloremys Gray 1870c:50

Type species: *Chloremys abnormis* [= *Dermatemys abnormis* Cope 1868 = subjective synonym of *Dermatemys mawii* Gray 1847], by original monotypy.

Etymology:* Greek, *chloros*, green, and *emys*, turtle; referring to the carapace and extremities sometimes appearing greenish in color.

Limnochelone Werner 1901b:297

Type species: *Limnochelone micrura* Werner 1901b [= subjective synonym of *Dermatemys mawii* Gray 1847], by original monotypy.


Etymology:* Greek, *limne*, pond, lake, and *chelone*, turtle; referring to the aquatic habitat and behavior of this species.

Dermatemys mawii Gray 1847 (14:8)


Central American River Turtle, Hicatee, *Tortuga Blanca*



Melvin Mérida / CBFTT / TCC / Petén, Guatemala / female



left: Russell A. Mittermeier / CBFTT / Veracruz, Mexico / juvenile
right: Gracia Gonzalez-Porter / CBFTT / Belize R., Belize / male



** (orange dots = probably introduced or trade; stars = type localities)

Distribution: Belize, Guatemala, Mexico (*Campeche, Chiapas, Oaxaca, Quintana Roo, Tabasco, Veracruz*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 131,154 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 326,595 sq. km

Size (Max SCL):* male 47.9 cm, female 52.0 cm (Legler and Vogt 2013; E. Reyes-Grajales, unpubl. data); 60.0 cm (possibly CCL) (Vogt et al. 2011 CBFTT)

Comment:* Smith and Smith (1980) and Legler and Vogt (2013) cited 65.0 cm as the maximum known carapace length but provided no details; this may have been CCL, and the data are suspect. The largest female measured by Vogt *in* Legler and Vogt (2013) was 51.2 cm SCL.

CBFTT Account: Vogt, Polisar, Moll, and Gonzalez-Porter (2011)

IUCN Red List:* Critically Endangered (CR A2bcd+4bcd) (Lesher-Gordillo, Gallardo-Alvarez, Hernández-Marín, Reyes-Grajales, McKnight, Ligon, Marlin, Barrett, and Zenteno-Rúiz, in press). Previously: Critically Endangered (CR A2abd+4d) (Vogt, Gonzalez-Porter, and van Dijk 2006); Endangered (EN) (TFTSG *in* Baillie and Groombridge 1996)

CITES: Appendix II (1981)

USA ESA:* Endangered (USFWS 1983)

Synonymy:

Dermatemys mawii Gray 1847:55

Emys mawii

Type locality: "South America." Restricted to "Alvarado, Veracruz, Mexico" [yellow star] by Smith and Taylor (1950a:346).

Type specimen: NHMUK 1947.3.4.12, holotype, see Stuart (1963).

Etymology:* Eponym honoring Lieutenant Solomon Maw (1800s), United Kingdom; British naval officer who collected the type.

Emys berardii Duméril and Bibron *in* Duméril and Duméril 1851:11

Ptychemys berardii, Clemmys berardii, Dermatemys berardii, Pseudemys berardi

Type locality: "l'Amérique mérid., ...environs de Vera-Cruz" [Veracruz, Mexico] [green star].

Type specimens: MNHN 7835 and 9518, syntypes (2).

Etymology:* Eponym honoring Auguste Bérard (1796–1852), France; naval officer who presented the types.

Dermatemys mavei Cope 1865:187 (*nomen novum*)

Dermatemys marvei

Etymology:* Eponym honoring Lieutenant Solomon Maw (1800s), United Kingdom; British naval officer who collected the type.

Comment: Unjustified emendation of *mawii*.

Dermatemys abnormis Cope 1868:120

Chloremys abnormis

Type locality: "Belize River, Yucatan" [Belize] [blue star, location estimated]. Restricted to "Belize [city], British Honduras" by Smith and Taylor (1950a:316, 1950b:19); restriction reversed by Dunn and Stuart (1951:59).

Type specimen: ANSP 61 (formerly USNM 6545), holotype, see Malnate (1971) and Reynolds et al. (2007).

Etymology:* Latin, *abnormis*, irregular or abnormal; referring to the broad vertebral scutes in the type specimen.

Dermatemys salvinii Gray 1870c:50

Type locality: "Guatemala" [pink star, location estimated].

Type specimen: NHMUK 1946.1.22.96 (formerly 1864.2.19.6), holotype, see Stuart (1963).

Etymology:* Eponym honoring Osbert Salvin (1835–1898), United Kingdom; ornithologist, herpetologist, and explorer of Guatemala, who collected the type.

Limnochelone micrura Werner 1901b:298

Type locality: "Mexico." Restricted to "Alvarado, Veracruz, Mexico" [yellow star] by Smith and Taylor (1950a:316, 1950b:19).

Type specimen: KZM s/n, holotype, apparently destroyed during World War II.

Etymology:* Greek, *mikros*, small, and *oura*, tail, rump; referring to the very short tail.

Dermatemys mawei Neill and Allen 1959:28 (*nomen novum*)

Etymology:* Eponym honoring Lieutenant Solomon Maw (1800s), United Kingdom; British naval officer who collected the type.

Comment: Unjustified emendation of *mawii*.

Family Kinosternidae Agassiz 1857a (14:9)


Mud Turtles

(includes 2 subfamilies, 4 genera, 34 species, 42 taxa)

Cinosternoidae Agassiz 1857a:249

Kinosterna Gray 1869a:180

Kinosternidae Hay 1892:560



Kinosternidae Species Richness

Subfamily Kinosterninae Agassiz 1857a (14:9)

Kinosternine Mud Turtles

(includes 2 genera, 31 species, 39 taxa)

Cinosternoidae Agassiz 1857a:249

Kinosternina Gray 1869a:180

Kinosterninae Lindholm 1929:277

Kinosternon Spix 1824 (14:9, 17:11) (37)*

American Mud Turtles

(includes 3 subgenera, 25 species, 33 taxa)

Monoclida Rafinesque 1815:75 (*nomen nudum*)

Etymology:* Greek, *monos*, one, only, and Latin, *claudere*, to close; referring to the single plastral hinge.

Uronyx Rafinesque 1815:75 (*nomen nudum*)

Etymology:* Greek, *oura*, tail, and *onyx*, claw, nail; referring to the horn nail on the tip of the tail.

Kinosternon Spix 1824:17 (*nomen protectum*)

Type species: *Kinosternon longicaudatum* Spix 1824 [= subjective synonym of *Testudo scorpioides* Linnaeus 1766], by subsequent designation by Bell (1828c:515).

Etymology:* Greek, *kino*, to move, and *sternon*, chest, sternum; referring to the hinged plastron.

Comment: Name conserved by ICZN (1989), see Smith et al. (1980b).

Kinosternum Bonaparte 1830:166 (*nomen novum*)

Etymology:* Greek, *kino*, to move, and *sternon*, chest, sternum; referring to the hinged plastron.

Cinosternon Wagler 1830b:137 (*nomen novum*)

Etymology:* Greek, *kino*, to move, and *sternon*, chest, sternum; referring to the hinged plastron.

Monoclida Rafinesque 1832:64

Type species: *Monoclida retziana* Rafinesque 1832 [= subjective synonym of *Testudo scorpioides* Linnaeus 1766], by original monotypy.

Etymology:* Greek, *monos*, one, only, and Latin, *claudere*, to close; referring to the single plastral hinge.

Uronyx Rafinesque 1832:64

Type species: *Uronyx scorpioides* [= *Testudo scorpioides* Linnaeus 1766], by original monotypy.

Etymology:* Greek, *oura*, tail, and *onyx*, claw, nail; referring to the horn nail on the tip of the tail.

Cinosternum Burmeister 1837:731 (*nomen novum*)

Etymology:* Greek, *kino*, to move, and *sternon*, chest, sternum; referring to the hinged plastron.

Swanka Gray 1844:32

Type species: *Swanka scorpiodes* Gray [= *Testudo scorpioides* Linnaeus 1766], by original monotypy.

Etymology:* Vernacular (Native Central American, Miskito), *swanka*, indigenous name used for turtles of this genus.

Thyrosternum Agassiz 1857a:418,427

Type species: *Thyrosternum pensilvanica* [= *Testudo pensilvanica* Gmelin 1789 = subjective synonym of *Testudo subrubra* Bonnaterre 1789], by subsequent designation by Lindholm (1929:277).

Etymology:* Greek, *thyra*, gate, door, and *sternon*, chest, sternum; referring to the hinged plastron.

Platythyra Agassiz 1857a:420,429

Type species: *Platythyra flavescens* Agassiz 1857a, by original monotypy.

Etymology:* Greek, *platys*, flat, even, wide, and *thyra*, gate, door; referring to the flat, hinged shell.

Cinosternos Herrera 1901:35 (*nomen novum et suppressum*)

Etymology:* Greek, *kino*, to move, and *sternon*, chest, sternum; referring to the hinged plastron.

Comment: Unjustified replacement name for *Cinosternon*. Name suppressed by ICZN (1922).

Cryptochelys Iverson, Le, and Ingram 2013:933 (14:9, 17:11)

Type species: *Cryptochelys leucostomum* [= *Cinosternon leucostomum* Duméril and Bibron *in* Duméril and Duméril 1851], by original designation.

Etymology:* Greek, *krypto*, hidden, and *chelys*, turtle; referring to the hidden, cryptic nature of this clade of turtles not previously recognized as distinct.

Kinosternon (Kinosternon) Spix 1824 (14:9, 17:11) (37)*

Neotropical Mud Turtles

(includes 16 species, 22 taxa)

Monoclida Rafinesque 1815:75 (*nomen nudum*)

Etymology:* Greek, *monos*, one, only, and Latin, *claudere*, to close; referring to the single plastral hinge.

Uronyx Rafinesque 1815:75 (*nomen nudum*)

Etymology:* Greek, *oura*, tail, and *onyx*, claw, nail; referring to the horn nail on the tip of the tail.

Kinosternon Spix 1824:17 (*nomen protectum*)

Type species: *Kinosternon longicaudatum* Spix 1824 [= subjective synonym of *Testudo scorpioides* Linnaeus 1766], by subsequent designation by Bell (1828c:515).

Etymology:* Greek, *kino*, to move, and *sternon*, chest, sternum; referring to the hinged plastron.

Comment: Name conserved by ICZN (1989), see Smith et al. (1980b).

Kinosternum Bonaparte 1830:166 (*nomen novum*)

Etymology:* Greek, *kino*, to move, and *sternon*, chest, sternum; referring to the hinged plastron.

Cinosternon Wagler 1830b:137 (*nomen novum*)

Etymology:* Greek, *kino*, to move, and *sternon*, chest, sternum; referring to the hinged plastron.

Monoclida Rafinesque 1832:64

Type species: *Monoclida retziana* Rafinesque 1832 [= subjective synonym of *Testudo scorpioides* Linnaeus 1766], by original monotypy.

Etymology:* Greek, *monos*, one, only, and Latin, *claudere*, to close; referring to the single plastral hinge.

Uronyx Rafinesque 1832:64

Type species: *Uronyx scorpioides* [= *Testudo scorpioides* Linnaeus 1766], by original monotypy.

Etymology:* Greek, *oura*, tail, and *onyx*, claw, nail; referring to the horn nail on the tip of the tail.

Cinosternum Burmeister 1837:731 (*nomen novum*)

Etymology:* Greek, *kino*, to move, and *sternon*, chest, sternum; referring to the hinged plastron.

Swanka Gray 1844:32

Type species: *Swanka scorpiodes* Gray [= *Testudo scorpioides* Linnaeus 1766], by original monotypy.

Etymology:* Vernacular (Native Central American, Miskito), *swanka*, indigenous name used for turtles of this genus.

Cinosternos Herrera 1901:35 (*nomen novum et suppressum*)

Etymology:* Greek, *kino*, to move, and *sternon*, chest, sternum;

referring to the hinged plastron.
Comment: Unjustified replacement name for *Cinosternon*. Name suppressed by ICZN (1922).

Kinosternon (Kinosternon) abaxillare Baur *in* Stejneger 1925 (14:10, 21:45) (37)*


Central Chiapas Mud Turtle, *Casquito Pardo*



Eduardo Reyes-Grajales / Villa Hidalgo, Chiapas, Mexico



Eduardo Reyes-Grajales / *left*: Villaflores, *right*: Villa Hidalgo, Chiapas, Mexico



(orange dots = probable trade or translocated; star = type locality)


Distribution: Guatemala, Mexico (*Chiapas*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 22,225 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 28,970 sq. km
Size (Max SCL):* male 17.4 cm, female 15.7 cm (Iverson 2008, 2010; Legler and Vogt 2013; Reyes-Grajales et al. 2021; E. Reyes-Grajales, unpubl. data)
CBFTT Account:* Reyes-Grajales, Iverson, Walde, and Akre (2025)
IUCN Red List: Vulnerable (VU A2cd+4cd) (Reyes-Grajales and Guichard-Romero 2021)
CITES:* Appendix II, as *Kinosternon* spp. (2023)
Synonymy:
Kinosternon abaxillare Baur *in* Stejneger 1925:462
Kinosternon scorpioides abaxillare, Kinosternon cruentatum abaxillare, Kinosternon (Kinosternon) abaxillare
Type locality: "Tuxtla, Chiapas, Mexico." Emended to "Tuxtla Gutiérrez, Chiapas" [Mexico] by Legler and Vogt (2013:142) [yellow star].
Type specimens: USNM 7518, holotype, apparently lost, see Reynolds et al. (2007); USNM 7519–21, 7523–29, paratypes (10), 7520 also apparently lost.
Etymology:* Latin, *ab-*, without; *axilla*, armpit; referring to the lack of an axillary scute on the anterior bridge.

Kinosternon (Kinosternon) acutum Gray 1831d (14:9, 21:45) (37)*


Tabasco Mud Turtle, *Montera*, *Pochitoque Jahuactero*



John B. Iverson / CBFTT / Belize



John B. Iverson / CBFTT / Belize




(orange dot = questionable; stars = type localities)


Distribution:* Belize, Guatemala, Mexico (*Campeche, Chiapas, Oaxaca, Quintana Roo, Tabasco, Veracruz*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 162,775 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 301,292 sq. km
Size (Max SCL): male 10.5 cm, female 12.0 cm (Iverson and Vogt 2011 CBFTT; Ceballos et al. 2013)
CBFTT Account: Iverson and Vogt (2011)
IUCN Red List:* Near Threatened (NT A2c+4c) (Reyes-Grajales, Munscher, Walde, and Iverson, in press). Previously: Near Threatened (NT) (TFTSG *in* Baillie and Groombridge 1996)
CITES:* Appendix II, as *Kinosternon* spp. (2023)
Synonymy:
Kinosternon scorpioides acuta Gray 1831d:34
Kinosternon scorpoides acuta, Kinosternon acutum, Cryptochelys acuta, Kinosternon (Kinosternon) acutum
Type locality: Not designated. Restricted to "C. America" [Central America] by Gray (1844:33); to "British Honduras" [Belize] by Schmidt (1941:476); and to "Cosamaloapan, Veracruz, Mexico" [yellow star] by Smith and Taylor (1950a:347, 1950b:23).
Type specimen: NHMUK 1947.3.4.58, holotype, see Stuart (1963).
Etymology:* Latin, *acutus*, acute or pointed; referring to the pointed unnotched posterior margin of the plastron.
Cinosternum berendtianum Cope 1865:189
Cinosternon berendtianum, Kinosternon berendtianum
Type locality: "Tabasco" [Mexico] [green star, location estimated].
Type specimens: USNM 6517, 106293–94, syntypes (3); 6517 now lost, see Reynolds et al. (2007).
Etymology:* Eponym honoring Carl H. Berendt (1817–1878), Germany / USA; physician and naturalist who collected the types.

Cinosternon effeldtii Peters 1873:603
Cinosternum effeldtii
Type locality: "angeblich aus Mexico (Veracruz)." Restricted to "Cosamaloapan, Veracruz, Mexico" [yellow star] by Smith and Taylor (1950a:347).
Type specimen: ZMB 7868, lectotype, designated by Fritz et al. (1994:168); erroneously listed as "holotype" by Iverson (1980).
Etymology:* Eponym honoring Rudolf Effeldt (1821–1876), Germany; herpetologist and chiropterologist who collected the type specimen.

Kinosternon (Kinosternon) alamosae Berry and Legler 1980 [37]*
Alamos Mud Turtle, *Casquito de Alamos*


John B. Iverson / Sonora, Mexico



John B. Iverson / Sonora, Mexico



**
(orange dots = uncertain identifications; star = type locality)

Distribution: Mexico (*Sinaloa, Sonora*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 45,160 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 81,862 sq. km
Size (Max SCL): male 13.5 cm, female 12.6 cm (Berry and Legler 1980; Ceballos et al. 2013)
IUCN Red List:* Near Threatened (NT A3c) (Lara Resendiz, in press). Previously: TFTSG Provisional Red List: Least Concern (LC) (2018); Data Deficient (DD) (Frost, Hammerson, and Gadsden 2007); Least Concern (LC) [Not Listed] (Behler 1996)
CITES:* Appendix II, as *Kinosternon* spp. (2023)
Synonymy:
Kinosternon alamose Pritchard 1979:556 (*nomen suppressum*)
Type locality: "vicinity of Alamos, Sonora, and the lower Rio Yaqui" [Mexico] [yellow star].
Type specimen: Not designated.
Etymology:* Toponym for Alamos, Sonora, Mexico.
Comment: Name suppressed by ICZN (1985c), see Pritchard and Pronek (1982).
Kinosternon alamosae Berry and Legler 1980:1 (*nomen protectum*)
Kinosternon (Kinosternon) alamosae
Type locality: "Rancho Carrizal, 7.2 km north and 11.5 km west of Alamos, Sonora, Mexico (27º 05' N, 109º 03' W)" [yellow star].
Type specimen: LACM 127639, holotype.
Etymology:* Toponym for Alamos, Sonora, Mexico.
Comment: Name conserved by ICZN (1985c), see Pritchard and Pronek (1982).

Kinosternon (Kinosternon) albogulare (Duméril and Bocourt 1870) [37, 40]*
White-throated Mud Turtle


John B. Iverson / CBFTT / Rio Corobizi, Guanacaste, Costa Rica


John B. Iverson / CBFTT / Rio Corobizi, Guanacaste, Costa Rica


**
(orange dots = questionable or trade or translocated; star = type locality)

Distribution: Colombia (*Archipiélago de San Andrés, Providencia y Santa Catalina [prehistorical or modern introduction?]*), Costa Rica, El Salvador, Honduras, Nicaragua, Panama
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 126,189 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 528,842 sq. km
Size (Max SCL): male 20.5 cm, female 17.1 cm (Iverson 2010; Berry and Iverson 2011 CBFTT; J.M. Legler and J.F. Berry, unpubl. data)
CBFTT Account: Forero-Medina and Castaño-Mora (2011)
IUCN Red List:* Least Concern (LC) (Forero Medina, Iverson,

Hurtado-Gómez, López-Luna, van Dijk, and Rhodin, in press), as *Kinosternon scorpioides albogulare*. Previously: TFTSG Provisional Red List: Least Concern (LC) (2011), as *Kinosternon scorpioides albogulare*

CITES:* Appendix II, as *Kinosternon* spp. (2023)

Synonymy:

Cinosternon albogulare Duméril and Bocourt 1870:24

Cinosternum albogulare, Kinosternon cruentatum albogulare, Kinosternon scorpioides albogulare, Kinosternon albogulare, Kinosternon (Kinosternon) albogulare

Type locality: "S. Jose (Costa Rica)" [yellow star].

Type specimens: MNHN 1760 and 4349, syntypes (2), 1760 erroneously listed as "holotype" by Berry and Iverson (2001b:5); Uetz et al. (2019) listed MNHN 1760 and 4349 as syntypes.


Etymology:* Latin, *albus*, white, and *gula*, throat; referring to the unmarked chin.

Kinosternon (Kinosternon) chimalhuaca Berry, Seidel, and Iverson 1997 (07:7, 14:11, 21:46) (37)*


Jalisco Mud Turtle, *Casquito de Chamela*



John B. Iverson / Rio Purificación, Jalisco, Mexico



left: John B. Iverson, *right*: James F. Berry / Rio Purificación, Jalisco, Mexico



** (orange dot = probably translocated; star = type locality)

Distribution: Mexico (*Colima, Jalisco*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 9,828 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 14,412 sq. km

Size (Max SCL):* male 17.2 cm, female 14.0 cm (Legler and Vogt 2013; López-Luna et al. 2020; Garrido et al. 2021)

IUCN Red List:* Least Concern (LC) (Butterfield and Matías-Palafox 2025). Previously: Least Concern (LC) (van Dijk, Ponce Campos, and Garcia Aguayo 2007)

CITES:* Appendix II, as *Kinosternon* spp. (2023)

Synonymy:

Kinosternon chimalhuaca Berry, Seidel, and Iverson *in* Rogner 1996:23 (*nomen suppressum* pending)

Comment: Name pending suppression by ICZN, see Rogner et al. (2013, 2016).

Etymology:* Ethnonym honoring the Chimalhuaca, a prehistoric indigenous tribe described in the historical novel *Aztec*, believed to have lived in the area where the species occurs.

Kinosternon chimalhuaca Berry, Seidel, and Iverson 1997:331

Kinosternon (Kinosternon) chimalhuaca

Type locality: "30 m southeast of Mexico Highway 80, 1.9 km northeast of Barra de Navidad, Jalisco, Mexico (19º 15' N, 104º 43' S)" [yellow star].

Type specimen: CM 140201, holotype.


Etymology:* Ethnonym honoring the Chimalhuaca, a prehistoric indigenous tribe described in the historical novel *Aztec*, believed to have lived in the area where the species occurs.

Kinosternon (Kinosternon) cora Loc-Barragán, Reyes-Velasco, Woolrich-Piña, Grünwald, Venegas de Anaya, Rangel-Mendoza, and López-Luna 2020 (21:47) (37)*

Cora Mud Turtle, *Casquito Cora*, *Chacuanita Cora*



Marco A. López-Luna / Escuinapa, Sinaloa, Mexico



Marco A. López-Luna / Escuinapa, Sinaloa, Mexico / male



** (light red = possibly extirpated; star = type locality)

Distribution: Mexico (*Nayarit, Sinaloa*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 8,895 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 12,770 sq. km

Size (Max SCL):* male 8.8 cm, female 12.8 cm (Loc-Barragán et al. 2020; Loc-Barragán and Iverson 2025)

IUCN Red List:* Endangered (EN A2cd+4cd; B1ab(i,ii,iii)) (Loc-Barragán, Woolrich-Piña, López-Luna, and Iverson, in press). Previously: TFTSG Provisional Red List: Vulnerable (VU) (2020)

CITES:* Appendix I (2023)

Synonymy:

Kinosternon cora Loc-Barragán, Reyes-Velasco, Woolrich-Piña, Grünwald, Venegas de Anaya, Rangel-Mendoza, and

López-Luna 2020:512

Kinosternon (Kinosternon) cora

Type locality: "Mexico: Sinaloa: Ejido La Concepción (La Concha), Municipio de Escuinapa (22.531758°, -105.450767°; datum WGS 84; elev. 10 masl)" [yellow star].

Type specimen: MZFC-HE 35627, holotype.

Etymology:* Ethnonym honoring the Cora, the native Mexican ethnic group that is most widespread in the Nayarit region, where the species occurs.

Kinosternon (Kinosternon) creaseri Hartweg 1934 (14:9, 21:45) (37)*


Creaser's Mud Turtle, Yucatan Mud Turtle, *Casquito de Creaser*



John B. Iverson / nr. Felipe Carillo Puerto, Quintana Roo, Mexico



John B. Iverson / nr. Felipe Carillo Puerto, Quintana Roo, Mexico



** (star = type locality)

Distribution: Belize, Guatemala (?), Mexico (*Campeche, Quintana Roo, Yucatán*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 121,674 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 138,956 sq. km

Size (Max SCL): male 12.5 cm, female 12.1 cm (Iverson 1991; Ceballos et al. 2013; Legler and Vogt 2013)

IUCN Red List:* Near Threatened (NT A2c+4c) (Díaz-Gamboa, Butterfield, Jones, and Willey 2025). Previously: Least Concern (LC) (van Dijk, Lee, and Calderón Mandujano 2007); Near Threatened (NT) (TFTSG *in* Baillie and Groombridge 1996)

CITES:* Appendix II, as *Kinosternon* spp. (2023)

Synonymy:

Kinosternon creaseri Hartweg 1934:1

Cryptochelys creaseri, Kinosternon (Kinosternon) creaseri

Type locality: "one mile south of the Hacienda, Chichen Itza, Yucatan" [Mexico] [yellow star].

Type specimen: UMMZ 73090, holotype, see Peters (1952) and Kluge (1984).

Etymology:* Eponym honoring Edwin P. Creaser (1907–1981), USA; biologist at the University of Michigan who collected the types.

Kinosternon (Kinosternon) cruentatum (Duméril and Bibron *in* Duméril and Duméril 1851) (37, 38)*


Atlantic Red-cheeked Mud Turtle, *Chachagua*



John B. Iverson / CBFTT / Manuel, Tamaulipas, Mexico



John B. Iverson / CBFTT / *left*: Tamaulipas, Mexico / *right*: Dzibalchen, Campeche, Mexico



** (orange dots = probable trade or translocated; stars = type localities)

Distribution: Belize, Guatemala, Mexico (*Campeche, Chiapas, Oaxaca, Quintana Roo, Tamaulipas, Veracruz, Yucatán*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 256,583 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 727,821 sq. km

Size (Max SCL): male 16.6 cm, female 14.8 cm (Iverson 2010; Berry and Iverson 2011 CBFTT)

IUCN Red List:* Least Concern (LC) (Forero Medina, Iverson, Hurtado-Gómez, López-Luna, van Dijk, and Rhodin, in press), as *Kinosternon scorpioides cruentatum*

CITES:* Appendix II, as *Kinosternon* spp. (2023)

Synonymy:

Cinosternon cruentatum Duméril and Bibron *in* Duméril and Duméril 1851:16

Kinosternum cruentatum, Kinosternon cruentatum, Cinosternum cruentatum, Swanka cruentata, Thyrosternum cruentatum, Kinosternon cruentatum cruentatum, Kinosternon scorpioides cruentatum, Kinosternon (Kinosternon) cruentatum

Type locality: "Amér. septentr." Restricted to "San Mateo del Mar, Oaxaca, Mexico" [in error, orange star] by Smith and Taylor (1950a:339, 1950b:23). Restricted to "Pueblo Nuevo X-Can, Quintana Roo, Mexico (20.87°N, 87.60°W)" [yellow star] by Iverson and Berry (2024:18).

Type specimen: MNHN 1759, holotype, discussed by Bour (2004a).

Etymology:* Latin, *cruentatus*, bloody, spattered with blood, dripping blood; referring to the bright red or orange markings on the head and neck of some male specimens of some populations.


Kinosternon cruentatum consors Stejneger 1941:458

Type locality: "Cozumel Island, Yucatan, Mexico" [green star].
Type specimen: USNM 13912, holotype, see Cochran (1961) and Reynolds et al. (2007).
Etymology:* Latin, *consortia*, brotherly, consort; referring to its described close relationship to *Kinosternon cruentatum cruentatum*.


Kinosternon (Kinosternon) herrerai Stejneger 1925 (14:9, 21:45) (37)*
Herrera's Mud Turtle, *Casquito de Herrera*



John B. Iverson / nr. Gutiérrez Zamora, Veracruz, Mexico




John B. Iverson / nr. Gutiérrez Zamora, Veracruz, Mexico



** (orange dots = probable trade; stars = type localities)

Distribution:* Mexico (*Hidalgo, Puebla, Querétaro [?], San Luis Potosi, Tamaulipas, Veracruz*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 67,130 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 115,115 sq. km
Size (Max SCL): male 17.2 cm, female 15.7 cm (Carr and Mast 1988; Legler and Vogt 2013)
IUCN Red List:* Near Threatened (NT A2c+4cd) (López-Luna and Cázares-Hernández, in press). Previously: Near Threatened (NT) (van Dijk, Hammerson, Lavin, and Mendoza Quijano 2007); Least Concern (LC) [Not Listed] (Behler 1996)
CITES:* Appendix II, as *Kinosternon* spp. (2023)
Synonymy:
Kinosternon herrerai Stejneger 1925:462
Cryptochelys herrerai, Kinosternon (Kinosternon) herrerai
Type locality: "Xochimilco, Valley of Mexico" [orange star] [in error; market specimen]. Restricted to "La Laja, Veracruz, Mexico" [green star] by Smith and Taylor (1950a:349, 1950b:24); and to "vicinity of Tampico" [Tamaulipas, Mexico] [yellow star] by Smith and Brandon (1968:54).
Type specimen: USNM 61249, holotype, see Cochran (1961) and Reynolds et al. (2007).
Etymology:* Eponym honoring Alfonso Luis Herrera López (1868–1942), Mexico; zoologist and director of the National Museum of Mexico.

Kinosternon (Kinosternon) hirtipes (Wagler 1830c) (09:11, 17:9) (37, 39)*
Rough-footed Mud Turtle, Rough-legged Mud Turtle, *Casquito de Pata Rugosa*
(includes 6 subspecies)



** (subspecies: *hirtipes* = red [light red = extirpated], *chapalaense* = purple, *magdalense* = blue, *megacephalum* = pink, *murrayi* = green, *tarascense* = brown; red dots = extinct or extirpated, orange dots = questionable or misidentified)

Distribution:* Mexico (*Aguascalientes, Chihuahua, Coahuila, Distrito Federal, Durango, Guanajuato, Jalisco, México, Michoacán, Querétaro, Zacatecas*), USA (*Texas*)
Introduced: Mexico (*San Luis Potosi*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 145,976 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 645,957 sq. km
Size (Max SCL):* male 18.6 cm, female 16.3 cm (see subspp.)
IUCN Red List:* Vulnerable (VU A2c+4c) (Macip-Ríos and Iverson, in press). Previously: Least Concern (LC) (van Dijk, Hammerson, Vazquez Diaz, Quintero Diaz, Santos, and Flores-Villela 2007); Least Concern (LC) [Not Listed] (Behler 1996)
CITES:* Appendix II, as *Kinosternon* spp. (2023)

Kinosternon (Kinosternon) hirtipes hirtipes (Wagler 1830c) [09:11, 17:9] [37, 39]*


Valley of Mexico Mud Turtle



Sergio Nolasco-Perez / Mexico City, Distrito Federal, Mexico



left: Gracia Gonzalez-Porter / Mexico City, Distrito Federal, Mexico
right: Paco Soberón / Mexico City, Distrito Federal, Mexico



** (subspecies: *hirtipes* = red [light red = extirpated], *tarascense* = brown; red dots = extirpated; star = type locality)

Distribution: Mexico (*Distrito Federal*, *México*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 5,620 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 6,507 sq. km

Size (Max SCL): male 14.0 cm, female 14.0 cm (Iverson 1981; Legler and Vogt 2013)

IUCN Red List:* Critically Endangered (CR B2ab(ii,iii,iv)) (Macip-Ríos and Iverson, in press). Previously: TFTSG Provisional Red List: Critically Endangered (CR) (2019)

CITES:* Appendix II, as *Kinosternon* spp. (2023)

Synonymy:

Cinosternon hirtipes Wagler 1830b:137 (*nomen nudum*) [17:9]

Cinosternon hirtipes Wagler 1830c:expl.tab:pl.V, f.29-30 [09:11, 17:9]

Clemmys (Cinosternon) hirtipes, Kinosternum hirtipes, Kinosternon hirtipes, Cinosternum hirtipes, Thyrosternum hirtipes, Ozotheca hirtipes, Kinosternon hirtipes hirtipes, Kinosternon (Kinosternon) hirtipes, Kinosternon (Kinosternon) hirtipes hirtipes

Type locality: Not designated. Restricted to "Mexico" by Wagler (1833:unpaginated); to "Mazatlán, Sinaloa" [Mexico] [in error] by Smith and Taylor (1950a:343, 1950b:25); and to "lakes near Mexico City" [Mexico] [yellow star] by Schmidt (1953:89).

Type specimen: ZSM 2374/0, holotype, see Franzen and Glaw (2007) and Uetz (2019); Iverson (1981, 1992) erroneously listed the holotype as ZSM 1374/0.

Etymology:* Latin, *hirtus*, rough, and *pes*, foot; referring to the rough scales on the feet.

Kinosternon (Kinosternon) hirtipes chapalaense Iverson 1981 [37, 39]*


Lake Chapala Mud Turtle



John B. Iverson / Lago Zapotlan, Jalisco, Mexico



left: John B. Iverson / nr. Ocotlan, Jalisco, Mexico [AMNH, New York]
right: John B. Iverson / Lago Zapotlan, Jalisco, Mexico



** (subspecies: *chapalaense* = purple, *magdalense* = blue, *tarascense* = brown; star = type locality)

Distribution: Mexico (*Jalisco*, *Michoacán*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 5,532 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 9,485 sq. km

Size (Max SCL): male 15.2 cm, female 14.9 cm (Iverson 1981; Legler and Vogt 2013)

IUCN Red List:* Critically Endangered (CR B2ab(ii,iii,iv)) (Macip-Ríos and Iverson, in press). Previously: TFTSG Provisional Red List: Critically Endangered (CR) (2018)

CITES:* Appendix II, as *Kinosternon* spp. (2023)

Synonymy:

Kinosternon hirtipes chapalaense Iverson *in* Pritchard 1979:557 (*nomen nudum*)

Kinosternon hirtipes chapalaense Iverson 1981:51

Kinosternon hirtipes chapalense, Kinosternon (Kinosternon) hirtipes chapalaense

Type locality: "Lake Chapala, 0.25 mile off Chapala, Jalisco, Mexico [20º 18' N, 103º 12' W]" [yellow star].

Type specimen: UMMZ 97128, holotype, see Kluge (1984).

Etymology:* Toponym for Lake Chapala, Jalisco, Mexico.


Kinosternon (Kinosternon) hirtipes magdalense Iverson 1981 [37, 39]*
San Juanico Mud Turtle



John B. Iverson / Presa San Juanico, Michoacán, Mexico



John B. Iverson / Presa San Juanico, Michoacán, Mexico



(subspecies: *chapalaense* = purple, *magdalense* = blue, *tarascense* = brown; star = type locality)

Distribution: Mexico (*Michoacán*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 694 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 925 sq. km

Size (Max SCL): male 13.6 cm, female 13.2 cm (Legler and Vogt 2013)

IUCN Red List:* Critically Endangered (CR B2ab(ii,iii,iv)) (Macip-Ríos and Iverson, in press). Previously: TFTSG Provisional Red List: Critically Endangered (CR) (2018)

CITES:* Appendix II, as *Kinosternon* spp. (2023)

Synonymy:


Kinosternon hirtipes magdalense Iverson 1981:53
Kinosternon (Kinosternon) hirtipes magdalense

Type locality: "along the face of the dam at *Presa* San Juanico, Michoacán [ca. 19º 50' N, 102º 40' W]" [Mexico] [yellow star].
Type specimen: UF 45035, holotype.
Etymology:* Toponym for the Magdalena Valley, Michoacán, Mexico.


Kinosternon (Kinosternon) hirtipes megacephalum Iverson 1981 [37, 39]*
Viesca Mud Turtle
(Extinct, ca. 1970)



John B. Iverson / 3.2 km SE Viesca, Coahuila, Mexico



John B. Iverson / 3.2 km SE Viesca, Coahuila, Mexico



(subspecies: *megacephalum* = pink, *murrayi* = green; orange dot = questionable or introduced, red star = extinct = type locality)

Distribution: Mexico (*Coahuila [extinct]*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 1,659 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 2,142 sq. km

Size (Max SCL): male 9.9 cm, female 11.7 cm (Iverson 1981; Legler and Vogt 2013)

IUCN Red List:* Extinct (EX) (Macip-Ríos and Iverson, in press). Previously: TFTSG Provisional Red List: Extinct (EX) (2018)

CITES:* Appendix II, as *Kinosternon* spp. (2023)

Synonymy:

Kinosternon hirtipes megacephalum Iverson 1981:52
Kinosternon hirtipes megalocephala, Kinosternon megacephalum, Kinosternon (Kinosternon) hirtipes megacephalum

Type locality: "3.2 km SE Viesca [25º 21' N, 102º 48' W], Coahuila" [Mexico] [red star].
Type specimen: SM(BCB) 11466, holotype, see Auth et al. (2000).
Etymology:* Greek, *mega*, large, and *kephale*, head; referring to the large head.

Kinosternon (Kinosternon) hirtipes murrayi Glass and Hartweg 1951 [37, 39]*


Mexican Plateau Mud Turtle



John B. Iverson / Nazas, Durango, Mexico



John B. Iverson / *left*: Galeana, Chihuahua, Mexico / *right*: north of Durango, Durango, Mexico



** (subspecies: *megacephalum* = pink, *murrayi* = green; orange dots = questionable or misidentified, red dot = extinct; star = type locality)

Distribution:* Mexico (*Chihuahua, Coahuila, Durango, Zacatecas*), USA (*Texas*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 95,917 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 316,857 sq. km

Size (Max SCL):* male 18.6 cm, female 16.3 cm (Legler and Vogt 2013; Platt et al. 2024)

IUCN Red List:* Near Threatened (NT A2c+4c) (Macip-Ríos and Iverson, in press). Previously: TFTSG Provisional Red List: Least Concern (LC) (2018)

CITES:* Appendix II, as *Kinosternon* spp. (2023)

Synonymy:

Kinosternon murrayi Glass and Hartweg 1951:50

Kinosternon hirtipes murrayi, Kinosternon (Kinosternon) hirtipes murrayi

Type locality: "Harper Ranch, 37 miles south of Marfa, Presidio County, Texas" [USA] [yellow star].

Type specimen: TCWC 650, holotype, see Lieb (2023).

Etymology:* Eponym honoring Leo T. Murray (1902–1958), USA; naturalist at Texas A & M University and director of the Strecker Museum.

Kinosternon (Kinosternon) hirtipes tarascense Iverson 1981 [37, 39]*


Pátzcuaro Mud Turtle



Rodrigo Macip-Ríos / Lago de Pátzcuaro, Michoacán, Mexico



Rodrigo Macip-Ríos / Lago de Pátzcuaro, Michoacán, Mexico



** (subspecies: *hirtipes* = red [light red = extirpated], *chapalaense* = purple, *magdalense* = blue, *murrayi* = green, *tarascense* = brown; red dots = extirpated; orange dot = questionable or misidentified; star = type locality)

Distribution:* Mexico (*Aguascalientes, Guanajuato, Jalisco, México, Michoacán, Querétaro*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 36,554 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 68,075 sq. km

Size (Max SCL):* male 15.1 cm, female 13.5 cm (Iverson 1981; Legler and Vogt 2013; J.B. Iverson, unpubl. data)

IUCN Red List:* Critically Endangered (CR B2ab(ii,iii,iv)) (Macip-Ríos and Iverson, in press). Previously: TFTSG Provisional Red List: Critically Endangered (CR) (2018)

CITES:* Appendix II, as *Kinosternon* spp. (2023)

Synonymy:

Kinosternon hirtipes tarascense Iverson 1981:52

Kinosternon (Kinosternon) hirtipes tarascense

Type locality: "Lago de Pátzcuaro, adjacent to city of Pátzcuaro [19º 32' N, 101º 36' W]" [Michoacán, Mexico] [yellow star].

Type specimen: UF 43506, holotype.

Etymology:* Ethnonym honoring the Tarascas people, inhabitants of the Lake Patzcuaro region of Michoacán, Mexico.


Kinosternon (Kinosternon) integrum (Le Conte 1854) [37]*
Mexican Mud Turtle, *Casquito de Burro*



John B. Iverson / Alamos, Sonora, Mexico



left: Anders G.J. Rhodin / Alamos, Sonora, Mexico
right: John B. Iverson / Nayarit, Mexico



** (orange dots = probable trade or introduced or misidentified; stars = type localities)

Distribution:* Mexico (*Aguascalientes, Chihuahua, Colima, Durango, Guanajuato, Guerrero, Hidalgo, Jalisco, Michoacán, Morelos, Nayarit, Nuevo León, Oaxaca, Puebla, Querétaro, San Luis Potosi, Sinaloa, Sonora, Tamaulipas, Zacatecas*)

Introduced: Mexico (*Baja California Sur*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 380,977 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 924,909 sq. km

Size (Max SCL): male 22.3 cm, female 19.6 cm (Macip-Ríos et al. 2009; Ceballos et al. 2013; Legler and Vogt 2103)

IUCN Red List:* Least Concern (LC) (Macip-Ríos and Iverson, in press). Previously: Least Concern (LC) (van Dijk, Hammerson, Vazquez Diaz, Quintero Diaz, Santos, and Flores-Villela 2007); Least Concern (LC) [Not Listed] (Behler 1996)

CITES:* Appendix II, as *Kinosternon* spp. (2023)

Synonymy:

Kinosternum integrum Le Conte 1854:183
Cinosternum integrum, Thyrosternum integrum, Thyrosternon integrum, Cinosternon integrum, Kinosternon integrum, Swanka integra, Cinosternum scorpioides integrum, Kinosternon scorpioides integrum, Kinosternon (Kinosternon) integrum
Type locality: "Mexico." Restricted to "Acapulco, Guerrero, Mexico" [yellow star] by Smith and Taylor (1950a:331, 1950b:25).
Type specimen: Possibly ANSP, apparently lost, not listed by Malnate (1971).
Etymology:* Latin, *integrum*, complete, intact; referring to the ability to completely close the shell using the plastral hinges.

Cinosternon rostellum Bocourt 1876a:391
Cinosternum rostellum
Type locality: "Guanajuato" [Mexico] [green star].
Type specimen: MNHN 2112, holotype.
Etymology:* Latin, *rostrum*, snout, beak; referring to the somewhat smaller snout or beak on this species.

Cinosternon guanajuatense Dugès 1888:107 (*nomen nudum*)
Etymology:* Toponym for the state of Guanajuato, Mexico.

Cinosternum scorpioides integrum mexicana Siebenrock 1907:579 (unavailable name)
Type locality: "Mexiko." Restricted to "Mazatlan" [Mexico] [blue star] by Smith and Smith (1980:115).
Type specimen: Possibly NMW, not located, not listed by Tiedemann et al. (1994) or Tiedemann and Grillitsch (1999).
Etymology:* Toponym for Mexico.


Kinosternon (Kinosternon) mexicanum (Le Conte 1854) [37, 38]*
Pacific Red-cheeked Mud Turtle, *Casquito Rojo, Pochitoque Rojo*



Eduardo Reyes-Grajales / Suchiate, Chiapas, Mexico / female



Eduardo Reyes-Grajales / Suchiate, Chiapas, Mexico / *left*: female, *right*: male



** (orange dot = probable trade or questionable; star = type localities)

Distribution:* El Salvador, Guatemala, Honduras, Mexico (*Chiapas, Oaxaca*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 50,284 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 111,368 sq. km

Size (Max SCL):* male 14.3 cm, female 14.5 cm (Iverson and Berry 2024)

IUCN Red List:* Least Concern (LC) (Forero Medina, Iverson, Hurtado-Gómez, López-Luna, van Dijk, and Rhodin, in press), as part of *Kinosternon scorpioides cruentatum*

CITES:* Appendix II, as *Kinosternon* spp. (2023)

Synonymy:*

Kinosternum mexicanum Le Conte 1854:182

Cinosternum mexicanum, Cinosternon mexicanum, Kinosternon mexicanum, Swanka mexicana, Kinosternon (Kinosternon) mexicanum

Type locality: "Mexico." Restricted to "San Mateo del Mar, Oaxaca, Mexico" [yellow star] by Smith and Taylor (1950a:339, 1950b:23).

Type specimen: ANSP 90, holotype, see Malnate (1971).

Etymology:* Toponym for Mexico.

Kinosternum triliratum Le Conte 1860:6

Cinosternon triliratum, Swanka trilirata, Cinosternum triliratum

Type locality: "Mexico." Restricted to "San Mateo del Mar, Oaxaca, Mexico" [yellow star] by Smith and Taylor (1950a:339, 1950b:23).

Type specimen: Possibly ANSP, apparently lost, not listed by Malnate (1971).


Etymology:* Latin, *tri*, three, and *liratus*, bearing ridges; referring to the three carapacial ridges.

Kinosternon (Kinosternon) oaxacae Berry and Iverson 1980 (37)*


Oaxaca Mud Turtle, *Casquito de Oaxaca*



John B. Iverson / Puerto Angel, Oaxaca, Mexico



John B. Iverson / Puerto Angel, Oaxaca, Mexico



**

(orange dot = trade, introduced, or misidentified; star = type locality)

Distribution: Mexico (*Guerrero, Oaxaca*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 24,005 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 38,821 sq. km

Size (Max SCL): male 17.5 cm, female 15.7 cm (Iverson 1986; Ceballos et al. 2013)

IUCN Red List:* Least Concern (LC) (Macip-Ríos, in press). Previously: TFTSG Provisional Red List: Least Concern (LC) (2018); Data Deficient (DD) (van Dijk and Canseco-Marquez 2007); Near Threatened (NT) (TFTSG *in* Baillie and Groombridge 1996)

CITES:* Appendix II, as *Kinosternon* spp. (2023)

Synonymy:

Kinosternon oaxacae Pritchard 1979:557 (*nomen suppressum*)

Type locality: "vicinity of Pochutla, Oaxaca" [Mexico] [yellow star].

Type specimen: Not designated.

Etymology:* Toponym for Oaxaca, Mexico.

Comment: Name suppressed by ICZN (1985c), see Pritchard and Pronek (1982).

Kinosternon oaxacae Berry and Iverson 1980:313 (*nomen protectum*)

Kinosternon (Kinosternon) oaxacae

Type locality: "11.6 km N. of Pochutla (San Pedro Pochutla), along Mexican Hwy. 175 (ca. 235 m), Oaxaca, Mexico (15º46'N, 96º28'W)" [yellow star].

Type specimen: UCM 48857, holotype.

Etymology:* Toponym for Oaxaca, Mexico.


Comment: Name conserved by ICZN (1985c), see Pritchard and Pronek (1982).

Kinosternon (Kinosternon) scorpioides (Linnaeus 1766) (07:8) (37, 40)*


Scorpion Mud Turtle, *Muçuã, Tapaculo, Jurará*




Frank Deschandol / French Guiana



John B. Iverson / CBFTT / *left*: Colombia / *right*: Cojedes, Venezuela



**



**

(orange dots = probable trade or translocated; stars = type localities)

Distribution: Argentina (*Chaco, Formosa, Jujuy, Salta, Tucumán*), Bolivia, Brazil (*Acre, Alagoas, Amapá, Amazonas, Bahia, Ceará, Goiás, Maranhão, Mato Grosso, Mato Grosso do Sul, Minas Gerais, Pará, Paraiba, Pernambuco, Piauí, Rio Grande do Norte, Rondônia, Sergipe, Tocantins*), Colombia (*Amazonas, Antioquia, Arauca, Atlántico, Bolívar, Caldas, Caquetá, Casanare, Cesar, Chocó, Córdoba, Guainía, Magdalena, Meta, Norte de Santander, Putumayo, Sucre, Vaupés, Vichada*), Ecuador, French Guiana, Guyana, Panama, Paraguay, Peru (*Amazonas, Huánuco, Loreto, Madre de Dios, Ucayali*), Suriname, Trinidad, Venezuela (*Amazonas, Apure, Aragua, Bolívar, Cojedes, Falcón, Guárico, Lara, Monagas, Portuguesa, Sucre, Táchira, Trujillo, Yaracuy, Zulia*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 6,451,732 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 13,294,360 sq. km

Size (Max SCL): male 18.5 cm, female 19.5 cm (Freiberg 1936; Métrailler and Le Gratiet 1996)

CBFTT Account: Berry and Iverson (2011)

IUCN Red List:* Least Concern (LC) (Forero Medina, Iverson, Hurtado-Gómez, López-Luna, van Dijk, and Rhodin, in press). Previously: TFTSG Provisional Red List: Regional (South America): Least Concern (LC) (2011); Least Concern (LC) [Not Listed] (Behler 1996)

CITES:* Appendix II, as *Kinosternon* spp. (2023)

Synonymy:

Testudo scorpioides Linnaeus 1766:352
Emys scorpioides, Chersine scorpioides, Terrapene scorpioidea, Cinosternon scorpioidea, Kinosternon scorpioides, Kinosternon scorpoides, Uronyx scorpioides, Terrapene scorpioides, Cinosternon scorpioides, Clemmys (Cinosternon) scorpioidea, Kinosternum scorpioides, Cinosternum scorpioides, Thyrosternum scorpioides, Swanka scorpiodes, Swanka scorpoides, Swanka scorpioides, Cinosternum scorpioides scorpioides, Kinosternon scorpioides scorpioides, Kinosternon (Kinosternon) scorpioides
Type locality: "Surinami" [Surinam] [yellow star, location estimated].
Type specimen: Possibly UUZM or NRM, but no types located, see Schoepff (1792:xi), not listed by Linné and Thunberg (1780), Thunberg (1828), Lönnberg (1896), Andersson (1900), Holm (1957), or Wallin (2001).
Etymology:* Greek, *skorpios*, scorpion, and *-ides*, similarity; referring to the pronounced horn nail on the tip of the tail, reminiscent of the stinger of a scorpion.
Comment: Description did not cite any sourced references or designated specimens, see also Schoepff (1792:xi).

Testudo tricarinata Retzius *in* Schoepff 1792:9 (senior homonym, not = *Testudo tricarinata* Bory de Saint-Vincent 1804 [= *Geoemyda spengleri*])
Terrapene tricarinata, Clemmys tricarinata
Type locality: Not known. Restricted to "Surinam" [yellow star, location estimated] by Fritz and Havaš (2007:256).
Type specimen: UPSZTY 284 (formerly UUZM 284 and MGA 53), holotype, listed by Thunberg (1810, 1828), Holm (1957), and Wallin (2001), type specimen figured in Schoepff (1792:pl.2).
Etymology:* Latin, *tri-*, three, and *carina*, the keel, *carinata*, keeled; referring to the three carapacial vertebral keels.

Testudo retzii Daudin 1801:174 (*nomen novum*)
Emys retzii, Terrapene retzii
Type locality: Not known. Restricted to "Surinam" [yellow star, location estimated] by Fritz and Havaš (2007:256).
Type specimen: Type specimen: UPSZTY 284 (formerly UUZM 284 and MGA 53), holotype by default, see Thunberg (1828), Holm (1957), and Wallin (2001).
Etymology:* Eponym honoring Anders J. Retzius (1742–1821), Sweden; naturalist at Lund University.
Comment: Unjustified replacement name for *tricarinata*.

Kinosternon longicaudatum Spix 1824:17 (*nomen protectum*)
Cinosternon longicaudatum, Kinosternum longicaudatum, Cinosternum longicaudatum, Thyrosternum longicaudatum, Swanka longicaudata
Type locality: "Brasiliam...ad campis aquosis" [Brazil].
Type specimen: ZSM 2375/0, lectotype, designated by Hoogmoed and Gruber (1983:353), see Franzen and Glaw (2007).
Etymology:* Latin, *longa*, long, wide, and *caudatum*, tailed; referring to the longer tail in males as supposedly representing a separate species.
Comment: Name conserved by ICZN (1989), see Smith et al. (1980b).

Kinosternon brevicaudatum Spix 1824:18
Cinosternon brevicaudatum, Kinosternum brevicaudatum, Cinosternum brevicaudatum
Type locality: "Brasiliam...ad ripam fluminis Solimöens" [Amazonas, Brazil] [= between Tefé and Tabatinga] [green star, location estimated].
Type specimens: ZSM, holotype or syntypes, apparently lost, including type specimen figured (pl.13), see Hoogmoed and Gruber (1983).
Etymology:* Latin, *brevis*, short, and *caudatum*, tailed; referring to the shorter tail in females as supposedly representing a separate species.

Kinosternon shavianum Bell 1825a:302
Cinosternon shavianum, Cinosternum shavianum, Thyrosternum shavianum
Type locality: Not known.
Type specimen: Originally OUM, not located, not listed by Nowak-Kemp and Fritz (2010).
Etymology:* Eponym honoring George Shaw (1751–1813), United Kingdom; naturalist at the British Museum of Natural History, London.

Monoclida retziana Rafinesque 1832:64 (*nomen novum*)
Testudo retziana
Etymology:* Eponym honoring Anders J. Retzius (1742–1821), Sweden; naturalist at Lund University.
Comment: Unjustified replacement name for *tricarinata*.

Cinosternon shawianum Bocourt 1876a:387 (*nomen novum*)
Etymology:* Eponym honoring George Shaw (1751–1813), United Kingdom; naturalist at the British Museum of Natural History, London.
Comment: Unjustified emendation of *shavianum*.

Cinosternum scorpioides integrum brasiliana Siebenrock 1907:579 (unavailable name)
Type locality: "Südamerika" [Brazil].
Type specimen: Possibly NMW, not located, not listed by Tiedemann et al. (1994) or Tiedemann and Grillitsch (1999).
Etymology:* Toponym for Brazil.

Kinosternon scorpioides pachyurum Müller and Hellmich 1936:100
Type locality: "Bolivien...Chaco...Villa Montes" [Bolivia] [blue star].
Type specimens: ZSM 129/1928/1–2, syntypes (2); ZSM 128/1928a–b, also syntypes (2), destroyed during WW II, see Franzen and Glaw (2007).
Etymology:* Greek, *pachys*, thick, coarse, and *oura*, tail, rump; referring to the relatively short, broad tail.


Kinosternon scorpioides seriei Freiberg 1936:169 (07:8)
Type locality: "El Tabacal (Salta)" [Argentina] [pink star].
Type specimen: MACN 1247, holotype.
Etymology:* Eponym honoring Pedro Serié (1865–1951), Argentina; zoologist, herpetologist, and curator at the Museo Argentino de Ciencias Naturales, Buenos Aires.

Kinosternon panamensis Schmidt 1946:5
Type locality: "Panama Railroad, Canal Zone" [Panama] [white star].
Type specimen: USNM 117369 (formerly MCZ 7996), holotype, see Cochran (1961) and Reynolds et al. (2007).
Etymology:* Toponym for Panama.

Kinosternon scorpioides panamense Mertens and Wermuth 1955:338
Etymology:* Toponym for Panama.
Comment: Unjustified emendation of *panamensis*.

Kinosternon scorpioides carajasensis Cunha 1970:1 (07:8)
Type locality: "compartimento da serra dos Carajás (serra Norte) Pará" [Brazil] [gray star].
Type specimen: MPEG 15, holotype.
Etymology:* Toponym for Serra dos Carajás, Brazil.

Kinosternon (Kinosternon) sonoriense (Le Conte 1854) (41)*
Sonora Mud Turtle, *Casquito de Sonora*
(includes 2 subspecies)



**
(subspecies: *sonoriense* = red [light red = extirpated], *longifemorale* = blue; red dots = extirpated, orange dots = questionable)

Distribution:* Mexico (*Baja California [extirpated], Chihuahua, Sinaloa, Sonora*), USA (*Arizona, California [extirpated], Nevada [extirpated], New Mexico*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 186,589 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 467,310 sq. km

Size (Max SCL):* male 20.4 cm, female 17.5 cm (see subspp.)

CBFTT Account:* Stone, Congdon, Stone, Stuart, Iverson, and Rosen (2022)

IUCN Red List:* Vulnerable (VU A4c) (Stone, in press).
Previously: Near Threatened (NT) (van Dijk 2011); Least Concern (LC) [Not Listed] (Behler 1996); Vulnerable (VU) (TFTSG *in* Baillie and Groombridge 1996)

CITES:* Appendix II, as *Kinosternon* spp. (2023)

Kinosternon (Kinosternon) sonoriense sonoriense (Le Conte 1854) (41)*
Desert Mud Turtle, *Casquito Común de Sonora*




William M. Hammond / Blackwater Hole, Peloncillo Mountains, New Mexico





left: John B. Iverson / Chihuahua, Mexico
right: Gerald Kuchling / New Mexico



**
(subspecies: *sonoriense* = red [light red = extirpated], *longifemorale* = purple; red dots = extirpated, orange dot = questionable; stars = type localities)

Distribution:* Mexico (*Baja California [extirpated], Chihuahua, Sinaloa, Sonora*), USA (*Arizona, California [extirpated], Nevada, New Mexico*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 184,500 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 467,310 sq. km

Size (Max SCL):* Northern populations: male 16.9 cm, female 17.5 cm (Iverson 1981; van Loben Sels et al. 1997; Ceballos et al. 2013; Legler and Vogt 2013). Southernmost population (Sinaloa): **Max SCL:** male 20.4 cm, female 13.6 cm [both estimated] (J.B. Iverson, unpubl. data); **Max CCL:** male 23.5 cm, female 15.8 cm (Molina-Serrano et al. 2025)

IUCN Red List:* Vulnerable (VU A4c) (Stone 2025). Previously: TFTSG Provisional Red List: Near Threatened (2022)

CITES:* Appendix II, as *Kinosternon* spp. (2023)

Synonymy:
Kinosternum sonoriense Le Conte 1854:184
Kinosternon sonoriense, *Cinosternum sonoriense*,

Thyrosternum sonoriense, Cinosternon sonoriense, Kinosternon sonoriense sonoriense, Kinosternon (Kinosternon) sonoriense, Kinosternon (Kinosternon) sonoriense sonoriense

Type locality: "Tucson...province of Sonora" [originally Mexico, now Arizona, USA] [yellow star].

Type specimen: Possibly ANSP, apparently lost, not listed by Malnate (1971); Uetz et al. (2019) erroneously list USNM 21710, the holotype for *Kinosternon sonoriense longifemorale*.

Etymology:* Toponym for Sonora, Mexico.

Kinosternum henrici Le Conte 1860:4

Thyrosternum henrici, Cinosternon henrici, Cinosternum henrici, Swanka henrici

Type locality: "New Mexico" [USA]. Data with holotype is "Gila River, New Mexico" [green star, location estimated]; incorrectly restricted to "vicinity of Las Cruces" [New Mexico, USA] [orange star] by Schmidt (1953:91), see Malnate (1971).

Type specimen: ANSP 83, holotype, see Malnate (1971).

Etymology:* Eponym honoring T. Charlton Henry (1825–1877), USA; surgeon with the U.S. Army who collected the type.

Kinosternon (Kinosternon) sonoriense longifemorale Iverson 1981 (41)*


Sonoyta Mud Turtle, *Casquito de Sonoyta*



John B. Iverson / Quitobaquito Springs, Organ Pipe National Monument, Arizona



John B. Iverson / Quitobaquito Springs, Organ Pipe National Monument, Arizona



** (subspecies: *sonoriense* = red [light red = extirpated], *longifemorale* = blue; red dots = extirpated, orange dots = questionable; star = type locality)

Distribution: Mexico (*Sonora*), USA (*Arizona*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 2,089 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 3,177 sq. km

Size (Max SCL): male 14.8 cm, female 16.0 cm (Rosen and Stone 2017 IUCN; M. Bogan and M.A. Grageda García, unpubl. data)

IUCN Red List: Critically Endangered (CR A3c) (Rosen and Stone 2017)

CITES:* Appendix II, as *Kinosternon* spp. (2023)

USA ESA:* Endangered (USFWS 2017b); Species Status Assessment: USFWS (2017a)

Synonymy:

Kinosternon sonoriense longifemorale Iverson 1981:43

Kinosternon (Kinosternon) sonoriense longifemorale

Type locality: "artificial pond fed by springs, Sonoyta, Sonora, Mexico (31°51' N, 112°50' W)" [yellow star].

Type specimen: USNM 21710, holotype, see Reynolds et al. (2007).

Etymology:* Latin, *longus*, long, wide, and *femoralis*, of the femur or thigh; referring to the long plastral interfemoral scute suture.

Kinosternon (Kinosternon) vogti López-Luna, Cupul-Magaña, Escobedo-Galván, González-Hernández, Centenero-Alcala, Rangel-Mendoza, Ramírez-Ramírez, and Cazares-Hernández 2018 (21:48) (37)*


Vallarta Mud Turtle, *Casquito de Vallarta*



Carolina Sánchez Arias / CCB / Puerto Vallarta, Jalisco, Mexico / male



left: Craig B. Stanford / Puerto Vallarta, Jalisco, Mexico / male
right: Carolina Sánchez Arias / Puerto Vallarta, Jalisco, Mexico / male



** (star = type locality)

Distribution: Mexico (*Jalisco, Nayarit*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 263 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 306 sq. km

Size (Max SCL):* male 8.9 cm, female 10.2 cm (López-Luna et al. 2018; T. Butterfield, unpubl. data)

IUCN Red List:* Critically Endangered (CR A4cde) (Cupul-Magaña, Butterfield, Gregory, Iverson, Macip-Ríos, and López-Luna 2022). Previously: TFTSG Provisional Red List: Critically Endangered (CR) (2018)

CITES:* Appendix I (2023)

Synonymy:

Kinosternon vogti López-Luna, Cupul-Magaña, Escobedo-Galván, González-Hernández, Centenero-Alcala,

Rangel-Mendoza, Ramírez-Ramírez, and Cazares-Hernández 2018:4

Kinosternon (Kinosternon) vogti

Type locality: "Puerto Vallarta, Jalisco, Mexico (WSG84, 20°38'24.99"N, 105°13'55.57"W, 10 m elev. masl)" [yellow star].

Type specimen: CNAR 31568 (formerly IBH), holotype.

Etymology:* Eponym honoring Richard C. Vogt (1949–2021), USA / Mexico / Brazil; herpetologist and turtle ecology researcher who worked and lived in USA, Mexico, and Brazil.

Kinosternon (Cryptochelys) Iverson, Le, and Ingram 2013 (14:9, 17:11) (37)*

Central American Mud Turtles

(includes 3 species, 4 taxa)

Cryptochelys Iverson, Le, and Ingram 2013:933 (14:9, 17:11)

Type species: *Cryptochelys leucostomum* [= *Cinosternon leucostomum* Duméril and Bibron *in* Duméril and Duméril 1851], by original designation.


Etymology:* Greek, *krypto*, hidden, and *chelys*, turtle; referring to the hidden, cryptic nature of this clade of turtles not previously recognized as distinct.

Kinosternon (Cryptochelys) angustipons Legler 1965 (14:9, 21:45) (37)*

Narrow-bridged Mud Turtle



Renae Reed / Boca Tapada, Costa Rica



left: John B. Iverson / Limon, Costa Rica
right: Renae Reed / Boca Tapada, Costa Rica



• (star = type locality)

Distribution: Costa Rica, Nicaragua, Panama

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO): 24,179 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 37,554 sq. km

Size (Max SCL): male 11.5 cm, female 12.0 cm (Legler 1965; Acuña-Mesén 1993; Ceballos et al. 2013)

IUCN Red List:* Vulnerable (VU B2ab(ii,iii,iv)) (Rhodin, Villegas Raygoza, Folt, and Mittermeier, in press). Previously: Vulnerable (VU) (TFTSG 1996)

CITES:* Appendix II, as *Kinosternon* spp. (2023)

Synonymy:

Kinosternon angustipons Legler 1965:617

Cryptochelys angustipons, Kinosternon (Cryptochelys) angustipons

Type locality:* "Los Diamantes, Limón Province, Costa Rica...two miles east of Guapiles" [yellow star].

Type specimen: KU 43631, holotype.


Etymology:* Latin, *angustus*, narrow; *pons*, bridge; referring to the narrow plastral bridge.

Kinosternon (Cryptochelys) dunni Schmidt 1947 (14:9, 21:45) (37)*


Dunn's Mud Turtle, *Cabeza de Trozo, Truenito*



German Forero-Medina / CBFTT / Rio Atrato, Chocó, Colombia



German Forero-Medina / Pizarro, Chocó, Colombia



** (star = type locality)

Distribution:* Colombia (*Antioquia*, *Chocó*, *Valle del Cauca*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 26,484 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 52,194 sq. km

Size (Max SCL): male 18.0 cm, female 16.6 cm (Iverson et al. 2012 CBFTT)

CBFTT Account: Iverson, Carr, Castaño-Mora, Galvis-Rizo, Rentería-Moreno, and Forero-Medina (2012)

IUCN Red List:* Endangered (EN B2ab(ii,iii)) (Forero Medina, Rentería-Moreno, Iverson, and Echavarría, in press). Previously: TFTSG Provisional Red List: Vulnerable (VU) (2011); Vulnerable (VU) (TFTSG 1996)

CITES:* Appendix II, as *Kinosternon* spp. (2023)

Synonymy:

Kinosternon dunni Schmidt 1947:109

Cryptochelys dunni, Kinosternon (Cryptochelys) dunni

Type locality: "Pizarro, Choco, Colombia" [yellow star].


Type specimen: FMNH 42804, holotype, see Marx (1958).

Etymology:* Eponym honoring Emmett Reid Dunn (1894–1956), USA; herpetologist at Smith College and curator at the Academy of Natural Sciences, Philadelphia.

Kinosternon (Cryptochelys) leucostomum (Duméril and Bibron *in* Duméril and Duméril 1851) (14:9, 21:45) (37)*

White-lipped Mud Turtle, *Pochitoque, Chechahua*

(includes 2 subspecies)



** (subspecies: *leucostomum* = red, *postinguinale* = blue; overlap = intergrades; orange dots = probable trade or translocated)

Distribution: Belize, Colombia (*Antioquia, Atlántico, Bolívar, Boyacá, Caldas, Cauca, Cesar, Chocó, Córdoba, Cundinamarca, Huila [?], Magdalena, Nariño, Santander, Sucre, Tolima, Valle del Cauca*), Costa Rica, Ecuador, Guatemala, Honduras, Mexico (*Campeche, Chiapas, Oaxaca, Quintana Roo, Tabasco, Veracruz, Yucatán [?]*), Nicaragua, Panama

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 657,730 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 3,179,757 sq. km

Size (Max SCL): male 21.4 cm, female 20.8 cm (see subspp.)

IUCN Red List:* Least Concern (LC) (Páez, Bock, Iverson, Cisneros-Heredia, Macip-Ríos, and Giraldo, in press). Previously: TFTSG Provisional Red List: Regional (South America): Least Concern (LC) (2011); Least Concern (LC) [Not Listed] (Behler 1996)


CITES:* Appendix II, as *Kinosternon* spp. (2023)

Kinosternon (Cryptochelys) leucostomum leucostomum (Duméril and Bibron *in* Duméril and Duméril 1851) (37)*


Northern White-lipped Mud Turtle



John B. Iverson / Belize Distr., Belize



John B. Iverson / Belize Distr., Belize



** (subspecies: *leucostomum* = red, *postinguinale* = blue; overlap = intergrades; orange dots = probable trade or translocated; stars = type localities)

Distribution: Belize, Guatemala, Honduras, Mexico (*Campeche, Chiapas, Oaxaca, Quintana Roo, Tabasco, Veracruz, Yucatán [?]*), Nicaragua

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 339,099 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 901,305 sq. km

Size (Max SCL): male 21.4 cm, female 20.8 cm (Legler and Vogt 2013)

CITES:* Appendix II, as *Kinosternon* spp. (2023)

Synonymy:

Cinosternon leucostomum Duméril and Bibron *in* Duméril and Duméril 1851:17

Kinosternum leucostomum, Kinosternon leucostomum, Cinosternum leucostomum, Thyrosternum leucostomum, Swanka leucostoma, Kinosternon leucostomum leucostomum, Cryptochelys leucostoma, Kinosternon (Cryptochelys) leucostomum, Kinosternon (Cryptochelys) leucostomum leucostomum

Type locality: "N.-Orléans; Mexique; Rio-Sumasinta (Amér. centr.)." Restricted to "Rio Usumacinta, El Peten, Guatemala" [yellow star, location estimated] by Schmidt (1941:488); and to "Cosamaloapam, Veracruz, Mexico" [orange star] by Smith and Taylor (1950a:347, 1950b:26).

Type specimen: MNHN 9087 (formerly 8311), lectotype, designated (as "type") by Stuart (1963:49).

Etymology:* Greek, *lefkos*, white, bright, and *stoma*, mouth, opening;

referring to the brightly colored whitish jaw edges in young animals.

Swanka maculata Gray 1869a:182

Type locality: "Mexico...Papalco Apoia; Vera Paz" [Papalco Apoia = Papaloapan River, Cosamaloapam, Veracruz, Mexico; Vera Paz = Alta Verapaz, Guatemala]. Restricted to "Cosamaloapam, Veracruz, Mexico" [orange star] by Stejneger (1941:457).

Type specimen: NHMUK 1946.1.22.25 (formerly 1862.6.6.1), lectotype, designated as the "type" by Boulenger (1889:43); some of the original syntypes (now paralectotypes) are identified as *K. acutum*, see Stuart (1963), Iverson (1980), Smith and Smith (1980), and Berry and Iverson (2001a).

Etymology:* Latin, *macula*, spot; referring to the spots and blotches on the head, throat, and neck.

Cinosternum brevigulare Günther 1885:17 (senior homonym, not = *Cinosternum brevigulare* Cope 1885 [= *Kinosternon leucostomum postinguinale*])

Type locality: "Mexico, Playa Vicente" [Veracruz] [blue star].

Type specimen: NHMUK 1946.1.22.38 (formerly 1860.6.17.24), holotype.

Etymology:* Latin, *brevis*, short, and *gula*, throat; referring to the small gular scute.

Cinosternum cobanum Günther 1885:18

Cinosternon cobanum

Type locality: "Guatemala, Coban; Cahabon." Restricted to "Cobán, Alta Verapaz, Guatemala" [gree star] by Smith and Taylor (1950a:317, 1950b:25).

Type specimens: NHMUK 1946.1.22.18–19 (formerly 1875.2.26.5 and 1880.11.20.18), syntypes (2), see Stuart (1963).

Etymology:* Toponym for Cobán, Guatemala.

Kinosternon mopanum Neill 1965:117

Type locality: "Waha Leaf Creek, southern Stann Creek District, British Honduras" [Belize] [pink star, location estimated].

Type specimen: MCZ 71635, holotype, apparently lost (Iverson 1976), specimen figured (f.6-7).

Etymology:* Ethnonym for the indigenous Mopan people, native to Belize and Guatemala.

Kinosternon (Cryptochelys) leucostomum postinguinale (Cope 1887) (37)*

Southern White-lipped Mud Turtle



John B. Iverson / Turbo, Antioquia, Colombia



Nathanael Stanek / Puerto Viejo de Sarapiqui, Heredia Prov., Costa Rica



** (subspecies: *leucostomum* = red, *postinguinale* = blue; overlap = intergrades; orange dots = probable trade or translocated; stars = type localities)

Distribution: Colombia (*Antioquia, Atlántico, Bolívar, Boyacá, Caldas, Cauca, Cesar, Chocó, Córdoba, Cundinamarca, Huila [?], Magdalena, Nariño, Santander, Sucre, Tolima, Valle del Cauca*), Costa Rica, Ecuador, Nicaragua, Panama

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 322,848 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 1,312,440 sq. km

Size (Max SCL): male 17.4 cm, female 15.8 cm (Acuña-Mesén 1993; Ceballos et al. 2016)

CITES:* Appendix II, as *Kinosternon* spp. (2023)

Synonymy:

Cinosternum brevigulare Cope 1885:389 (junior homonym, not = *Cinosternum brevigulare* Günther 1885 [= *Kinosternon leucostomum leucostomum*])

Cinosternon brevigulare

Type locality: "Tierra Caliente of Costa Rica at Sipurio, on the east coast" [yellow star, location estimated].

Type specimens: USNM 45582, 51165 (lost), syntypes (2); USNM 19797, listed as a "syntype" by Cochran (1961) and Uetz et al. (2019), is not a type, discussed by Reynolds et al. (2007).

Etymology:* Latin, *brevis*, short, and *gula*, throat; referring to the small gular scute.

Cinosternum postinguinale Cope 1887:23 (*nomen novum*)

Kinosternon postinguinale, Kinosternon leucostomum postinguinale, Kinosternon (Cryptochelys) leucostomum postinguinale

Type locality: "E. coast Costa Rica" [yellow star, location estimated].

Type specimens: USNM 45582, 51165 (lost), syntypes (2) by default; USNM 19797, listed as a "syntype" by Cochran (1961), is not a type, discussed by Reynolds et al. (2007).

Etymology:* Latin, *post*, after, behind, later, and *inguinalis*, belonging to the inguinal scutes; referring to the position of the inguinal scutes in relation to the posterior plastron hinge.

Comment: Justified emendation for *brevigulare*.

Cinosternum spurrelli Boulenger 1913:1030

Kinosternon spurrelli, Kinosternon leucostomum spurrelli

Type locality: "Choco, Colombia...Peña Lisa, Condoto, altitude 300 feet" [green star].

Type specimen: NHMUK 1946.11.12.1 (formerly 1913.11.12.1), holotype.

Etymology:* Eponym honoring Herbert G.F. Spurrell (1877–1918), United Kingdom; physician and zoologist who collected the type.

Kinosternon (Thyrosternum) Agassiz 1857a (14:9, 17:11) (37)*

Nearctic Mud Turtles

(includes 6 species, 7 taxa)

Thyrosternum Agassiz 1857a:418,427

Type species: *Thyrosternum pensilvanica* [= *Testudo pensilvanica* Gmelin 1789 = subjective synonym of *Testudo subrubra* Bonnaterre 1789], by subsequent designation by Lindholm (1929:277).

Platythyra Agassiz 1857a:420,429

Type species: *Platythyra flavescens* Agassiz 1857a, by original monotypy.

Kinosternon (Thyrosternum) baurii (Garman 1891) (37)*


Striped Mud Turtle



Dawn Wilson / CRM 3 / Hillsborough Co., Florida



John B. Iverson / Orange Co., Florida



** (stars = type localities)

Distribution:* USA (*Delaware, Florida, Georgia, Maryland, North Carolina, South Carolina, Virginia*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 374,910 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 887,290 sq. km

Size (Max SCL):* male 11.5 cm, female 13.8 cm (Pritchard 1980; Mitchell 1994; Ewert and Jackson 2005; Wilson et al. 2006; Ceballos et al. 2013; Johnston et al. 2019)

CBFTT Account:* Johnston, Dodd, and Wilson (2021)

IUCN Red List:* Near Threatened (NT A3cd+4cd) (Johnston, Donini, and Dodd 2025). Previously: Least Concern (LC) (van Dijk 2011); Least Concern (LC) [Not Listed] (Behler 1996)

CITES:* Appendix II, as *Kinosternon* spp. (2023)

Synonymy:

Cinosternum baurii Garman 1891:141

Kinosternon baurii, Kinosternon baurii baurii, Kinosternon (Thyrosternum) baurii

Type locality: "the island Key West...Florida..[&]..Cuba." Restricted to "Key West, Monroe County, Florida" [USA] [yellow star] by Uzzell and Schwartz (1955:33).

Type specimens: MCZ 282–87 (now 184720–1, 184723–6), 1558, 1563, 4380, UMMZ 53038 (Peters 1952; formerly MCZ 4718, but formerly MCZ 4379 according to Kluge 1984), FMNH 73481 (Marx 1958; formerly MCZ 4050), syntypes (11); Barbour and Loveridge (1929) listed only MCZ 1563 (2), 4380 (1), 4718 (5), and 4050 (1).

Etymology:* Eponym honoring Georg Baur (1859–1898), Czech Republic / Germany / USA; vertebrate morphologist and herpetologist at Yale University, Clark University, and University of Chicago.

Kinosternon bauri palmarum Stejneger 1925:463

Kinosternon baurii palmarum

Type locality: "Royal Palm State Park, Dade County, Florida" [USA] [green star]. Emended to "Paradise Key, Dade County, Florida" [USA] by Uzzell and Schwartz (1955:34).


Type specimen: USNM 61065, holotype, see Cochran (1961) and Reynolds et al. (2007).

Etymology:* Bionym for palms (Palmae); and Latin, *-arum*; referring to this species' habitat in Florida.


Kinosternon (Thyrosternum) durangoense Iverson 1979b (07:6) (37)*
Durango Mud Turtle, *Casquito de Durango*



John B. Iverson / Pedriceña, Durango, Mexico



Eric V. Goode / San Ignacio, Bolsón de Mapimí, Durango, Mexico



** (orange dots = questionable; star = type locality)

Distribution: Mexico (*Chihuahua, Coahuila, Durango*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 57,452 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 73,146 sq. km

Size (Max SCL): male 19.2 cm, female 14.5 cm (Iverson 1989; Ceballos et al. 2013; Iverson et al. 2018)

IUCN Red List:* Near Threatened (NT A3c) (Woolrich-Piña, Castañeda Gaytán, Rhodin, Iverson, and van Dijk, in press). Previously: TFTSG Provisional Red List: Least Concern (LC) (2018); Data Deficient (DD) (van Dijk 2007)

CITES:* Appendix II, as *Kinosternon* spp. (2023)

Synonymy:

Kinosternon flavescens durangoense Iverson 1979b:219

Kinosternon durangoense, Kinosternon (Thyrosternum) durangoense

Type locality:* "8 km from Ceballos, in Lago de los Palomas, Durango, Mexico" [now Chihuahua] [yellow star].

Type specimen: UF 16180, holotype.

Etymology:* Toponym for Durango, Mexico.


Kinosternon (Thyrosternum) flavescens (Agassiz 1857a) (07:6) (37)*
Yellow Mud Turtle, *Casquito Amarillo*



John B. Iverson / Canadian R., Beaver Co., Oklahoma



John B. Iverson / *left*: Louisa Co., Iowa, *right*: Canadian R., Beaver Co., Oklahoma



** (orange dots = questionable, probably translocated; stars = type localities)

Distribution: Mexico (*Chihuahua, Coahuila, Nuevo León, Tamaulipas, Veracruz*), USA (*Arizona, Colorado, Illinois, Iowa, Kansas, Missouri, Nebraska, New Mexico, Oklahoma, Texas*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 1,337,830 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 2,749,266 sq. km

Size (Max SCL):* male 16.8 cm, female 13.5 cm (Iverson 1989; Ernst and Lovich 2009; Ceballos et al. 2013; Brown et al. 2020)

CBFTT Account: Iverson and Christiansen (2023)

IUCN Red List:* Least Concern (LC) (Iverson 2025). Previously: Least Concern (van Dijk 2011); Least Concern (LC) [Not Listed] (Behler 1996)

CITES:* Appendix II, as *Kinosternon* spp. (2023)

Synonymy:

Platythyra flavescens Agassiz 1857a:430

Cinosternon flavescens, Cinosternum flavescens, Kinosternum flavescens, Kinosternon flavescens, Kinosternon flavescens flavescens, Kinosternon (Thyrosternum) flavescens

Type locality: "Texas, near San Antonio;...lower Rio Grande;...Red River, Arkansas;...Camp Yuma;...Gila River" [USA]. Restricted to "Waco, McLennan County, Texas, USA" [orange star] by Smith and Taylor (1950a:362, 1950b:24), but rejected by Maslin


(1959:22); and to "Rio Blanco, near San Antonio, Texas" [USA] [yellow star] by Schmidt (1953:89) and Iverson (1978:478).
Type specimen: USNM 50, lectotype, designated by Iverson (1978:478), see also Reynolds et al. (2007) and Lieb (2023).
Etymology:* Latin, *flavesco*, gold or yellow-colored; referring to the yellow throat color.


Kinosternon flavescens spooneri Smith 1951:195
Kinosternon spooneri
Type locality: "Henderson County State Forest, 7 miles north of Oquawka, Illinois" [USA] [green star].
Type specimen: INHS 4244, holotype.
Etymology:* Eponym honoring Charles S. Spooner (1885–1963), USA; zoologist and entomologist at Eastern Illinois State University.

Kinosternon (Thyrosternum) steindachneri (Siebenrock 1906b) (12:11, 14:12, 17:12) (37)*

Florida Mud Turtle


John B. Iverson / CRM 3 / CBFTT / Monroe Co., Florida


John B. Iverson / CRM 3 / CBFTT / *left*: Monroe Co., Florida / *right*: Levy Co., Florida



(star = type locality)


Distribution: USA (*Florida*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 87,221 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 111,400 sq. km
Size (Max SCL):* male 11.4 cm, female 11.2 cm (Meshaka et al. 2017 CBFTT; Iverson 2022b)
CBFTT Account: Meshaka, Gibbons, Hughes, Klemens, and Iverson (2017), as part of *Kinosternon subrubrum*
IUCN Red List:* Data Deficient (DD) (Lechowicz 2025). Previously: Least Concern (LC) (van Dijk 2011), as subspecies of *Kinosternon subrubrum*
CITES:* Appendix II, as *Kinosternon* spp. (2023)
Synonymy:
Cinosternum steindachneri Siebenrock 1906b:727
Kinosternon steindachneri, Kinosternon subrubrum steindachneri, Kinosternon (Thyrosternum) steindachneri
Type locality: "Orlando in Florida" [USA] [yellow star].
Type specimens: NMW 23388:1–2, NHMUK 1946.1.22.23–24 (formerly 1922.6.16.3–4), syntypes (4), see Tiedemann and Häupl (1980), Iverson (1992), Tiedemann et al. (1994), and Gemel et al. (2019); Uetz et al. (2019) listed only NHMUK 1946.1.22.23–24.
Etymology:* Eponym honoring Franz Steindachner (1834–1919), Austria; ichthyologist and director at the Naturhistorisches Museum, Vienna.

Kinosternon (Thyrosternum) stejnegeri Hartweg 1938 (07:6, 09:10, 17:13) (37)*

Arizona Mud Turtle


John B. Iverson / nr. Sells, Pima Co., Arizona



John B. Iverson / nr. Sells, Pima Co., Arizona


(orange dots = probably translocated or misidentified; star = type locality)

Distribution: Mexico (*Sonora*), USA (*Arizona*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 36,175 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 69,238 sq. km
Size (Max SCL): male 18.1 cm, female 16.7 cm (Iverson 1989; Ceballos et al. 2013)
IUCN Red List:* Near Threatened (NT A2c+4c) (Jones, Willey, Macip-Ríos, and Lara Resendiz 2025). Previously: Least Concern (LC) (Frost, Hammerson, and Gadsden 2007), as *Kinosternon arizonense*
CITES:* Appendix II, as *Kinosternon* spp. (2023)
Synonymy:

Kinosternon flavescens stejnegeri Hartweg 1938:1 (17:13)
Kinosternon stejnegeri, Kinosternon (Thyrosternum) stejnegeri
Type locality: "Llano, Sonora...approximately midway between Nogales and Hermosillo" [Mexico] [yellow star, location estimated].
Type specimen: UMMZ 72235, holotype, see Peters (1952) and Kluge (1984).
Etymology:* Eponym honoring Leonhard H. Stejneger (1851–1943), Norway / USA; herpetologist, ornithologist, curator at the United States National Museum, Washington, D.C.
Comment: The Arizona Mud Turtle was listed in our checklists prior to TTWG (2017) as *Kinosternon arizonense* † Gilmore 1923, but that taxon is now considered a separate extinct species from the Pliocene–Pleistocene, and no longer included in this checklist, with *K. stejnegeri* (Hartweg 1938) considered the extant species.

Kinosternon (Thyrosternum) subrubrum (Bonnaterre 1789) (09:6, 14:12) (37)*
Eastern Mud Turtle, Common Mud Turtle
(includes 2 subspecies)



**

(subspecies: *subrubrum* = red, *hippocrepis* = blue; overlap = intergrades; orange dots = possibly translocated or misidentified)

Distribution:* USA (*Alabama, Arkansas, Delaware, Florida, Georgia, Illinois, Indiana, Kentucky, Louisiana, Maryland, Mississippi, Missouri, New Jersey, New York, North Carolina, Oklahoma, Pennsylvania, South Carolina, Tennessee, Texas, Virginia, West Virginia*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 1,446,617 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 2,489,041 sq. km

Size (Max SCL): male 11.6 cm, female 12.5 cm (see subspp.)

CBFTT Account: Meshaka, Gibbons, Hughes, Klemens, and Iverson (2017)


IUCN Red List:* Least Concern (LC) (Hughes and Meshaka 2025). Previously: Least Concern (LC) (van Dijk 2011); Least Concern (LC) [Not Listed] (Behler 1996)

CITES:* Appendix II, as *Kinosternon* spp. (2023)


Kinosternon (Thyrosternum) subrubrum subrubrum (Bonnaterre 1789) (09:6) (37)*
Southeastern Mud Turtle



Richard D. Bartlett / CBFTT / Apalachicola National Forest, Florida



Richard D. Bartlett / CBFTT / Apalachicola National Forest, Florida



**

(subspecies: *subrubrum* = red, *hippocrepis* = blue; overlap = intergrades; stars = type localities)

Distribution:* USA (*Alabama, Delaware, Florida, Georgia, Illinois, Indiana, Kentucky, Maryland, Mississippi, New Jersey, New York, North Carolina, Pennsylvania, South Carolina, Tennessee, Virginia, West Virginia*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 817,047 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 1,544,395 sq. km

Size (Max SCL): male 11.6 cm, female 12.5 cm (Meshaka et al. 2017 CBFTT)

CITES:* Appendix II, as *Kinosternon* spp. (2023)

Synonymy:

Testudo subrubra Lacepède 1788:132, synopsis[table] (*nomen suppressum*) (09:6)
Type locality: "Pensylvanie" [Pennsylvania, USA]. Restricted to "vicinity of Philadelphia" [Pennsylvania, USA] [yellow star] by Schmidt (1953:90).
Type specimen: Not located, holotype, originally live, figured in Edwards (1760:pl.287), sent to England in 1757 by W. Bartram.
Etymology:* Latin, *sub-*, below, and *ruber*, red; referring to the red-orange plastron of hatchlings.
Comment: Description cited as sourced from Edwards (1760:pl.287). Name suppressed by ICZN (2005a) as published in a rejected and invalid non-binomial work, see Savage (2003).

Testudo subrubra Bonnaterre 1789:27

Kinosternon subrubrum, Kinosternon subrubrum subrubrum, Kinosternon (Thyrosternum) subrubrum, Kinosternon (Thyrosternum) subrubrum subrubrum

Type locality: "Pensylvanie" [Pennsylvania, USA].

Type specimen: Not located, holotype, originally live, figured in Edwards (1760:pl.287), sent to England in 1757 by W. Bartram.

Etymology:* Latin, *sub-*, below, and *ruber*, red; referring to the red-orange plastron of hatchlings.

Comment: Description cited as sourced from Lacepède (1788) and Edwards (1760).

Testudo pensilvanica Gmelin 1789:1042

Emydes pensilvanica, Kinosternon pensilvanicum, Cinosternum pensilvanicum

Type locality: "Pensilvaniae aquis stagnantibus" [Pennsylvania, USA]. Restricted to "vicinity of Philadelphia" [Pennsylvania, USA] [yellow star] by Schmidt (1953:90).

Type specimen: Not located, holotype, originally live, type specimen figured in Edwards (1760:pl.287) and Seligmann (1773:pl.77), sent to England in 1757 by W. Bartram.

Etymology:* Toponym for Pennsylvania, USA.

Comment: Description cited as sourced from Seligmann (1773:pl. LXXVII), which is the same figure as in Edwards (1760:pl.287).

Emys pensylvanica Schweigger 1812:282 (*nomen novum*)

Terrapene pensylvanica, Terrapene pensyluanica, Cistuda pensylvanica, Sternotherus pensylvanica, Kinosternum pensylvanicum, Cinosternon pensylvanicum, Clemmys (Cinosternon) pensylvanica, Kinosternon pensylvanicum, Cinosternum pensylvanicum

Etymology:* Toponym for Pennsylvania, USA.

Comment: Unjustified emendation of *pensilvanica*.

Kinosternon pennsylvanicum Bell 1825a:304 (*nomen novum*)

Emys (Kinosternon) pennsylvanica, Kinosternum pennsylvanicum, Cinosternon pennsylvanicum, Cinosternum pennsylvanicum, Cistudo pennsylvanica, Terrapene pennsylvanica, Thyrosternum pennsylvanicum

Etymology:* Toponym for Pennsylvania, USA.

Comment: Unjustified emendation of *pensilvanica*.

Kinosternon (Kinosternon) doubledayii Gray 1844:33

Kinosternon doubledayii, Kinosternum doubledayii, Cinosternum doubledayii, Cinosternon doubledayii

Type locality: "California" [in error]. Restricted to "vicinity of Philadelphia" [Pennsylvania, USA] [yellow star] by Schmidt (1953:90).

Type specimen: NHMUK 1947.3.4.67, holotype.

Etymology:* Eponym honoring Edward Doubleday (1810–1849), United Kingdom; entomologist at the British Museum of Natural History, London, who collected the type.

Kinosternon (Kinosternon) oblongum Gray 1844:33

Kinosternon oblongum, Cinosternum oblongum

Type locality: "America." Restricted to "vicinity of Philadelphia" [Pennsylvania, USA] [yellow star] by Schmidt (1953:90).

Type specimens: NHMUK 1947.3.4.65–66, syntypes (2).

Etymology:* Latin, *oblongus*, elongated; referring to the slightly elongated carapace.

Kinosternon punctatum Gray 1856a:198

Cinosternum punctatum

Type locality: "North America." Restricted to "East Florida" [USA] [green star, location estimated] by Gray (1856b:46).

Type specimen: NHMUK 1946.1.22.30, holotype.

Etymology:* Latin, *punctum*, spot; referring to the spots or dots on the head.

Swanka fasciata Gray 1869a:183

Type locality: Not known. Restricted to "vicinity of Philadelphia" [Pennsylvania, USA] [yellow star] by Schmidt (1953:90).

Type specimen: NHMUK 1946.1.22.92, holotype.

Etymology:* Latin, *fascia*, band, ribbon, bundle; referring to the stripes on the head.

Kinosternon (Thyrosternum) subrubrum hippocrepis Gray 1856a (17:14) (37)*


Mississippi Mud Turtle



Richard D. Bartlett / CBFTT / Mississippi or Louisiana



left: John B. Iverson / Texas
right: Richard D. Bartlett / CBFTT / Mississippi or Louisiana



** (subspecies: *subrubrum* = red, *hippocrepis* = blue; overlap = intergrades; orange dots = possibly translocated or misidentified; star = type locality)

Distribution: USA (*Alabama, Arkansas, Florida, Georgia, Louisiana, Mississippi, Missouri, Oklahoma, Tennessee, Texas*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 661,861 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 1,131,025 sq. km

Size (Max SCL): male 10.7 cm, female 11.1 cm (Meshaka et al. 2017 CBFTT)

CITES:* Appendix II, as *Kinosternon* spp. (2023)

Synonymy:

Kinosternon hippocrepis Gray 1856a:198 (17:14)

Cinosternum hippocrepis, Cinosternon hippocrepis, Kinosternon subrubrum hippocrepis, Kinosternon (Thyrosternum) subrubrum hippocrepis

Type locality: "North America; New Orleans" [Louisiana, USA] [yellow star].

Type specimens: NHMUK 1946.1.22.16–17, syntypes (2).

Etymology:* Greek, *hippos*, horse, and *krepis*, shoe; referring to the horseshoe-shaped head stripe pattern.

Kinosternon louisianae Baur 1893c:676

Cinosternum louisianae
Type locality: "New Orleans, La." [Louisiana, USA] [yellow star].
Type specimen: USNM 15527, holotype, see Cochran (1961) and Reynolds et al. (2007).
Etymology:* Toponym for Louisiana, USA.

Sternotherus Bell *in* Gray 1825 (07:9, 17:10)

Musk Turtles
(includes 6 species, 6 taxa)

Sternothaerus Bell 1825a:305 (*partim*, *nomen suppressum*)
Type species: *Sternothaerus leachianus* Bell 1825a [= subjective synonym of *Emys castanea* Schweigger 1812 = *Pelusios castaneus*], by subsequent designation by Bell (1828c:515); not *Sternothaerus odoratus* Bell [= *Testudo odorata* Latreille *in* Sonnini and Latreille 1801], by subsequent incorrect designation by Fitzinger (1843:290).
Etymology:* Greek, *sternon*, chest, sternum, and *thairos*, hinge; referring to the plastral hinge.
Comment: Name suppressed by ICZN (1989), see Smith et al. (1980b).

Sternotherus Bell *in* Gray 1825:211 [Bell 1825b] (*nomen protectum*)
Type species: *Sternotherus odoratus* [= *Testudo odorata* Latreille *in* Sonnini and Latreille 1801], by subsequent designation by Stejneger (1902:237).
Etymology:* Greek, *sternon*, chest, sternum, and *thairos*, hinge; referring to the plastral hinge.
Comment: Name conserved by ICZN (1989), see Smith et al. (1980b).

Aromochelys Gray 1856a:199
Type species: *Aromochelys odorata* [= *Testudo odorata* Latreille *in* Sonnini and Latreille 1801], by subsequent designation by Strauch (1862:38).
Etymology:* Greek, *aroma*, an odor or aroma, and *chelys*, turtle; referring to the pungent musk odor produced by these turtles.

Ozotheca Agassiz 1857a:251,424
Type species: *Ozotheca odorata* [= *Testudo odorata* Latreille *in* Sonnini and Latreille 1801], by subsequent designation by Lindholm (1929:277).
Etymology:* Greek, *osme*, the smell, and *theke*, container, box; referring to the pungent musk odor produced by these turtles.


Goniochelys Agassiz 1857a:420,423
Type species: *Goniochelys triquetra* Agassiz 1857a [= subjective synonym of *Aromochelys carinata* Gray 1856a], by subsequent designation by Lindholm (1929:277).
Etymology:* Greek, *gonia*, angle, and *chelys*, turtle; referring to the carapacial vertebral keel.

Sternotherus carinatus (Gray 1856a)


Razor-backed Musk Turtle, Keeled Musk Turtle



Robert C. Thomson / CBFTT / Pascagoula R., Mississippi



left: James H. Harding / CBFTT / Louisiana
right: Stanley E. Trauth / CBFTT / Arkansas



** (orange dots = questionable, possibly translocations; stars = type localities)


Distribution: USA (*Alabama, Arkansas, Louisiana, Mississippi, Oklahoma, Texas*)
Introduced:* USA (*Florida*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 343,971 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 512,687 sq. km
Size (Max SCL): male 20.9 cm, female 15.5 cm (Iverson 2002; Lindeman 2008 CBFTT)
CBFTT Account: Lindeman (2008)
IUCN Red List:* Least Concern (LC) (Ricardez, Gordon, DeChellis, Brown, and Franklin 2025). Previously: Least Concern (LC) (van Dijk 2011); Least Concern (LC) [Not Listed] (Behler 1996)
CITES:* Appendix II, as *Sternotherus* spp. (2023)
Synonymy:
Aromochelys carinata Gray 1856a:199
Aromochelys carinatum, Aromochelys carinatus, Goniochelys carinata, Cinosternum carinata, Kinosternon carinatum, Sternotherus carinata, Sternotherus carinata carinata, Kinosternon carinata, Sternothaerus carinatus, Sternotherus carinatus
Type locality: "North America, Louisiana" [USA]. Restricted to "vicinity of New Orleans" [Louisiana, USA] [yellow star] by Schmidt (1953:87).
Type specimens: NHMUK 1947.3.4.32, 1947.3.4.64, 1947.3.4.83–84 (formerly 1854.7.1.15, 1855.12.17.5, 1856.12.17.4), syntypes (4).


Etymology:* Latin, *carina*, keel, *carinata*, keeled; referring to the carapacial vertebral keel.

Goniochelys triquetra Agassiz 1857a:420,423
Type locality: "Lake Concordia, in Louisiana" [USA] [green star].
Type specimens: MCZ 15085–86, 46634–35, USNM 2, syntypes (5), see Barbour and Loveridge (1929); USNM 2 lost, see Reynolds et al. (2007).
Etymology:* Latin, *triquetra*, triangular; referring to the triangular shell shape when viewed from the front.

Sternotherus depressus Tinkle and Webb 1955 (07:10, 17:15)
Flattened Musk Turtle


C. Kenneth Dodd, Jr. / CBFTT / TCC / Gurley Creek, Alabama


C. Kenneth Dodd, Jr. / CBFTT / Lost Creek and Sipsey Fork, Alabama


**


(star = type locality)

Distribution: USA (*Alabama*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 12,627 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 14,283 sq. km
Size (Max SCL): male 9.9 cm, female 12.5 cm (Dodd 2008 CBFTT)
CBFTT Account: Dodd (2008)
IUCN Red List:* Critically Endangered (CR A2bce+4bce) (Jenkins and Godwin 2024). Previously: Critically Endangered (CR A2bce+4bce) (van Dijk 2011); Vulnerable (VU) (TFTSG *in* Baillie and Groombridge 1996)
CITES:* Appendix II, as *Sternotherus* spp. (2023)
USA ESA:* Threatened, in the Black Warrior River system upstream from Bankhead Dam (USFWS 1987a)
Synonymy:
Sternotherus depressus Tinkle and Webb 1955:53
Sternothaerus depressus, Sternotherus minor depressus, Kinosternon depressum, Kinosternon depressus
Type locality: "Mulberry Fork of the Black Warrior River, 9 miles east of Jasper, Walker County, Alabama, near the bridge crossing of U.S. highway 78" [USA] [yellow star].
Type specimen: LSUM 109567 (formerly TU 16171), holotype.
Etymology:* Latin, *de-*, down, and *pressus*, pressed; referring to the depressed flattened shell.

Sternotherus intermedius Scott, Glenn, and Rissler 2018 (21:49)
Intermediate Musk Turtle


Peter A. Scott / Econfina River, Florida


Kurt A. Buhlmann / Conecuh National Forest, Alabama


**

(star = type locality)

Distribution: USA (*Alabama, Florida*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 34,646 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 46,519 sq. km
Size (Max SCL): male 11.6 cm, female 10.8 cm (J.B. Iverson and P.A. Scott, unpubl. data)
IUCN Red List:* Data Deficient (DD) (Scott, Iverson, and Brown 2025). Previously: TFTSG Provisional Red List: Least Concern (LC) (2018)
CITES:* Appendix II, as *Sternotherus* spp. (2023)
Synonymy:
Sternotherus intermedius Scott, Glenn, and Rissler 2018:13
Type locality: "Blackwater River ca. 1 mi NW of Milton airfield (now Peter Prince Field), Santa Rosa Co. FL" [Florida, USA] [yellow star].
Type specimen: UAHC 68-814, holotype.
Etymology:* Latin, *intermedius*, intermediate, in between;


referring to the prior perception that this species was a population of *Sternotherus minor* and *S. peltifer* hybrids.

Sternotherus minor (Agassiz 1857a) (21:49)
Loggerhead Musk Turtle


John B. Iverson / Levy Co., Florida



Timothy Walsh / CRM 3 / Marion Co., Florida


**
(orange dots = questionable, possibly translocated; star = type locality)

Distribution: USA (*Alabama, Florida, Georgia*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 168,076 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 204,472 sq. km

Size (Max SCL): male 14.4 cm, female 14.5 cm (Camp 1986; Enge and Foster 1986; Zappalorti and Iverson 2006)

IUCN Red List:* Least Concern (LC) (Brown, Welc, and van Dijk 2025). Previously: Least Concern (LC) (van Dijk 2011); Least Concern (LC) [Not Listed] (Behler 1996)

CITES:* Appendix II, as *Sternotherus* spp. (2023)

Synonymy:

Goniochelys minor Agassiz 1857a:424

Aromochelys minor, Sternotherus minor, Sternotherus carinatus minor, Sternotherus minor minor, Sternothaerus minor minor, Kinosternon minor, Kinosternon minor minor

Type locality: "neighborhood of Mobile;...Columbus, Georgia;...and New Orleans" [USA]. Restricted to "Columbus, Georgia" [USA] [yellow star] by Schmidt (1953:88).

Type specimens: MCZ 1570, 1571(2), 1573, USNM 7111 (formerly MCZ 1572), UMMZ 63520 (formerly MCZ 1572), syntypes (6); MCZ 1573, USNM 7111, and UMMZ 63520 identified as *Sternotherus odoratus* by Tinkle (1958); see Barbour and Loveridge (1929), Peters (1952), Cochran (1961), Kluge (1984), and Reynolds et al. (2007).


Etymology:* Latin, *minor*, small; referring to its smaller size than the other *Goniochelys* described by Agassiz (= *Sternotherus carinatus*).

Sternotherus odoratus (Latreille *in* Sonnini and Latreille 1801) (21:50)
Musk Turtle, Stinkpot, Common Musk Turtle


Peter May / CRM 3 / Volusia Co., Florida



John B. Iverson / Alachua Co., Florida


**
(orange dots = possibly translocated, red dot = erroneous or extirpated; stars = type localities)

Distribution: Canada (*Ontario, Québec*), Mexico (?) (*Chihuahua? [erroneous record or extirpated?]*); USA (*Alabama, Arkansas, Connecticut, Delaware, Florida, Georgia, Illinois, Indiana, Iowa, Kansas, Kentucky, Louisiana, Maine, Maryland, Massachusetts, Michigan, Mississippi, Missouri, New Hampshire, New Jersey, New York, North Carolina, Ohio, Oklahoma, Pennsylvania, Rhode Island, South Carolina, Tennessee, Texas, Vermont, Virginia, West Virginia, Wisconsin*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 2,352,387 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 4,565,464 sq. km

Size (Max SCL): male 13.7 cm, female 15.0 cm (Ewert 2005; Ernst and Lovich 2009)

IUCN Red List:* Least Concern (LC) (Brown, Iverson, and Welc 2025). Previously: Least Concern (LC) (van Dijk 2011); Least Concern (LC) [Not Listed] (Behler 1996)

CITES:* Appendix II, as *Sternotherus* spp. (2023)

Synonymy:

Testudo odorata Latreille *in* Sonnini and Latreille 1801:122 (*nomen protectum*)

Emys odorata, Terrapene odorata, Cistuda odorata, Sternotherus odorata, Sternothaerus odoratus, Kinosternum odoratum, Emys (Kinosternon) odoratum, Kinosternon

odoratum, Didicla odorata, Staurotypus odoratus, Clemmys (Sternothaerus) odorata, Cistudo odorata, Sternotherus odoratus, Aromochelys odorata, Aromochelys odoratum, Cinosternum odoratum, Ozotheca odorata

Type locality: "les eaux dormantes de la Caroline" [USA]. Restricted to "vicinity of Charleston, South Carolina" [USA] [yellow star] by Schmidt (1953:87).

Type specimen: Not located, type specimen figured (pl.opp.112,f.3).

Etymology:* Latin, *odoratus*, to have an odor; referring to the strongly scented musk glands.

Comment: Name conserved by ICZN (1989), see Smith et al. (1980b).

Testudo glutinata Daudin 1801:194

Emys glutinata, Clemmys glutinata

Type locality: "les Etats-Unis d'Amérique" [USA]. Restricted to "vicinity of Lancaster, Pennsylvania" [USA] [green star] by Schmidt (1953:87).

Type specimens: Possibly MNHN, not located, type specimens figured in Schoepff (1895:pl.24,f.B) and Daudin (1801:pl.24,f.4).

Etymology:* Latin, *glutinum*, glue; referring to the poorly developed plastral hinge (as if glued), in comparison to other similar species.

Terrapene boscii Merrem 1820:27

Sternothaerus boscii

Type locality: "America septentrionali." Restricted to "vicinity of Philadelphia" [Pennsylvania, USA] [blue star] by Schmidt (1953:87).

Type specimen: Not known or located.

Etymology:* Eponym honoring Louis Augustin Guillaume Bosc (1759–1828), France / USA; botanist and naturalist who served as French consul to the USA.

Kinosternum guttatum Le Conte 1854:185

Cinosternum guttatum, Aromochelys guttata

Type locality: "Pennsylvania" [USA]. Restricted to "vicinity of Philadelphia" [Pennsylvania, USA] [blue star] by Schmidt (1953:87).

Type specimen: ANSP 63, holotype, see Malnate (1971).

Etymology:* Latin, *guttatus*, spotted or speckled; referring to the spots on the carapace.

Ozotheca tristycha Agassiz 1857a:392,425

Aromochelys tristycha

Type locality: "Osage River, in Missouri, and in Williamson County, in Texas...near San Antonio,...Medina River, in Texas" [USA]. Restricted to "San Antonio" [Texas, USA] [pink star] by Schmidt (1953:87).

Type specimens: MCZ 1574 (4), 1576 (2), 1922 (formerly USNM 70), USNM 64–65, 69–72, 7890, syntypes (14), see Barbour and Loveridge (1929), Cochran (1961), Reynolds et al. (2007), and Lieb (2023).

Etymology:* Greek, *tris*, three, and *stichos*, line, row; referring to the three keels on the carapace of juvenile specimens noted by Agassiz.

Testudo glutinosa Agassiz 1857a:425 (*nomen novum*)

Emys glutinosa

Etymology:* Latin, *glutinum*, glue; referring to the poorly developed plastral hinge (as if glued), in comparison to other similar species.


Comment: Unjustified emendation or error for *glutinata*.

Sternotherus peltifer Smith and Glass 1947 (17:16, 21:49)


Stripe-necked Musk Turtle



C. Kenneth Dodd, Jr. / Buttahatchie R., Marion Co., Alabama



left: Matthew L Niemiller / Hiwassee R., Polk Co., Tennessee
right: C. Kenneth Dodd, Jr. / Buttahatchie R., Marion Co., Alabama



** (orange dots = questionable, possibly translocated; star = type locality)

Distribution: USA (*Alabama, Georgia, Louisiana, Mississippi, North Carolina, Tennessee, Virginia*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 168,042 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 311,673 sq. km

Size (Max SCL):* male 12.6 cm, female 12.6 cm (P.A. Scott, unpubl. data; G.J. Brown, unpubl. data)

IUCN Red List:* Data Deficient (DD) (Brown and Welc 2025). Previously: Least Concern (LC) (van Dijk 2011), as subspecies of *Sternotherus minor*

CITES:* Appendix II, as *Sternotherus* spp. (2023)

Synonymy:

Sternotherus peltifer Smith and Glass 1947:22

Sternotherus carinatus peltifer, Sternotherus minor peltifer, Sternothaerus minor peltifer, Kinosternon minor peltifer

Type locality: "Bassfield, Jefferson Davis County, 30 miles west of Hattiesburg, Miss." [Mississippi, USA] [yellow star].

Type specimen: TCWC 1205, holotype, see Tinkle and Webb (1955).

Etymology:* Greek, *pelte*, a small shield, and *phero*, to bear; referring to the small scutes on the bridge.

Subfamily Staurotypinae Gray 1869a (14:13, 21:51)

Staurotypine Mud Turtles

(includes 2 genera, 3 species, 3 taxa)

Staurotypina Gray 1869a:180

Staurotypinae Siebenrock 1907:531

Staurotypidae Bickham and Carr 1983:925

Claudius Cope 1865

Narrow-bridged Musk Turtles

(includes 1 species, 1 taxon)

Claudius Cope 1865:187

Type species: *Claudius angustatus* Cope 1865, by original monotypy.

Etymology:* Latin, *claudus*, imperfect, defective; referring to the very small and somewhat inadequate plastron.

Staurosternon Duméril *in* Bocourt 1868:122

Type species: *Claudius megalocephalus* Bocourt 1868 [= subjective synonym of *Claudius angustatus* Cope 1865], by original monotypy.

Etymology:* Greek, *stauros*, cross, and *sternon*, chest, sternum; referring to the cruciform plastron.


Claudius angustatus Cope 1865

Narrow-bridged Musk Turtle, *Chopontil*




John B. Iverson / Hattieville, Belize Dist., Belize





left: John R. Polisar / northwest Belize
right: Richard C. Vogt / Lerdo de Tejada, Veracruz, Mexico



** (orange dot = possibly translocated; star = type locality)

Distribution:* Belize, Guatemala, Mexico (*Campeche, Chiapas, Oaxaca, Puebla, Quintana Roo, Tabasco, Veracruz, Yucatán*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 212,668 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 448,706 sq. km

Size (Max SCL): male 16.5 cm, female 15.0 cm (Iverson and Berry 1980; Ceballos et al. 2013; Legler and Vogt 2013)

IUCN Red List:* Endangered (EN A2acd+4acd) (Reyes-Grajales, Reynoso-Rosales, Vásquez-Cruz, Rivera Arroyo, Munscher, and Walde, in press). Previously: Near Threatened (NT) (TFTSG 1996)

CITES:* Appendix II (2023)

Synonymy:

Claudius angustatus Cope 1865:187

Claudius angustatum

Type locality: "Tabasco, Mexico" [yellow star, location estimated]

Type specimen: USNM 6518, holotype (parts of the same specimen formerly USNM 6525), see Cochran (1961) and Reynolds et al. (2007).

Etymology:* Latin, *angustus*, narrow; referring to the narrow plastral bridge.

Claudius megalocephalus Bocourt 1868:122

Type locality: "Mexico." Restricted to "Tabasco, Mexico" [yellow star, location estimated] by Smith and Taylor (1950a:345).

Type specimen: MNHN 1600, holotype, see Smith and Smith (1980:34).

Etymology:* Greek, *megalos*, large, long, and *kefale*, head; referring to the disproportionately large head.

Claudius macrocephalus Gray 1873d:69 (*nomen novum*)

Etymology:* Greek, *makros*, large, long, and *kefale*, head; referring to the disproportionately large head.

Comment: Unjustified emendation or error for *megalocephalus*.

Claudius megacephalus Boulenger 1889:33 (*nomen novum*)

Etymology:* Greek, *megalos*, large, long, and *kefale*, head; referring to the disproportionately large head.

Comment: Unjustified emendation or error for *megalocephalus*.

Claudius agassizii Smith and Taylor 1950a:345 (*nomen nudum*)

Etymology:* Eponym honoring Louis Agassiz (1807–1873), Switzerland / USA; herpetologist at Harvard University who founded the Museum of Comparative Zoology.

Staurotypus Wagler 1830b

Giant Musk Turtles

(includes 2 species, 2 taxa)

Staurotypus Wagler 1830b:137

Type species: *Staurotypus triporcata* [= *Terrapene triporcata* Wiegmann 1828], by original monotypy.

Etymology:* Greek, *stauros*, cross, stake, and *typos*, shape, image, type; referring to the cross-shaped plastron.

Stauremys Gray 1864c:127


Type species: *Staurotypus (Stauremys) salvinii* Gray 1864c, by original monotypy.

Etymology:* Greek, *stauros*, cross, stake, and *emys*, turtle; referring to the cross-shaped plastron.

Staurotypus salvinii Gray 1864c
Pacific Coast Giant Musk Turtle, *Crucilla*



John B. Iverson / Puerto Arista, Chiapas, Mexico



John B. Iverson / Puerto Arista, Chiapas, Mexico



** (stars = type localities)

Distribution: El Salvador, Guatemala, Mexico (*Chiapas, Oaxaca*)
Introduced: USA (*Florida*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 39,342 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 94,961 sq. km
Size (Max SCL): male 18.6 cm, female 20.6 cm (Legler and Vogt 2013); Smith and Smith (1980) and Dean and Bickham (1983) cite ca. 25 cm as maximum, but without documentation.
IUCN Red List:* Endangered (EN A2acd+4acd) (Reyes-Grajales, Walde, Iverson, and Munscher, in press). Previously: Near Threatened (NT) (TFTSG *in* Baillie and Groombridge 1996)
CITES:* Appendix II (2023)
Synonymy:
Staurotypus (Stauremys) salvinii Gray 1864c:127
Stauremys salvinii, Staurotypus salvinii, Staurotypus salvini
Type locality: "Haumanchal, Guatemala." Emended to "Huamuchal, Guatemala" by Boulenger (1889:32); and to "Huamuchal, 14º04'N, 91º34'W, Suchitepequez Province, Guatemala...immediately west of the mouth of Rio Nagua" [yellow star] by Legler and Vogt (2013:99).
Type specimen: NHMUK 1946.1.22.79, holotype, see Stuart (1963).
Etymology:* Eponym honoring Osbert Salvin (1835–1898), United Kingdom; ornithologist, herpetologist, and explorer of Guatemala, who collected the type.
Staurotypus marmoratus Fischer 1872:265
Type locality: "Mexico, Tejas." Restricted to "Santa Efigenia, Oaxaca, Mexico" [green star] by Smith and Taylor (1950a:339, 1950b:27).
Type specimens: Not located, possibly ZIN, type specimen figured (pl.10).
Etymology:* Latin, *marmor*, marble, and *-atus*, provided with; referring to the marbled appearance of the carapace.
Claudius severus Cope 1872:24
Staurotypus (Claudius) severus
Type locality: "Santa Efigenia, on the western side of the Isthmus of Tehuantepec, Mexico" [green star].
Type specimen: USNM 64005, holotype, see Cochran (1961) and Reynolds et al. (2007).
Etymology:* Latin, *severus*, serious, strict, severe; referring to the species being able to bite painfully.
Staurotypus biporcatus Gadow 1905:209 (*nomen nudum*)
Etymology:* Latin, *bi-*, two, and *porcatus*, keeled; referring to an erroneous assumption that the species has two keels.


Staurotypus triporcatus (Wiegmann 1828)
Northern Giant Musk Turtle, *Guao*



Eduardo Reyes-Grajales / Juárez, Chiapas, Mexico



Eduardo Reyes-Grajales / Juárez, Chiapas, Mexico



** (orange dots = probable trade or translocated; stars = type localities)

Distribution:* Belize, Guatemala, Honduras, Mexico (*Campeche, Chiapas, Oaxaca, Puebla, Quintana Roo, Tabasco, Veracruz, Yucatán [?]*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 219,483 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 517,892 sq. km
Size (Max SCL): male 37.8, female 40.2 cm (Legler and Vogt 2013)
IUCN Red List:* Vulnerable (VU A2bcd+4bcd) (Reyes-Grajales, Reynoso-Rosales, Vásquez Cruz, Rivera Arroyo, Munscher, and Walde, in press). Previously: Near Threatened (NT) (TFTSG *in* Baillie and Groombridge 1996)
CITES:* Appendix II (2023)
Synonymy:
Terrapene triporcata Wiegmann 1828:364

Staurotypus triporcata, Staurotypus triporcatus, Staurotypus (Staurotypus) triporcatus, Emys (Kinosternon) triporcata, Kinosternon triporcatum, Clemmys (Staurotypus) triporcata

Type locality: "Rio Alvarado" [Veracruz, Mexico] [yellow star].

Type specimen: ZMB 127, holotype, see Fritz et al. (1994); figured in Gutsche (2016:f.2).

Etymology:* Latin, *tri-*, three, and *porcatus*, ridged; referring to the three carapacial keels.

Claudius pictus Cope 1872:26

Type locality: "Vera Paz" [Guatemala]. Emended to "Alta Verapaz" [Guatemala] by Smith and Taylor (1950a:317, 1950b:27); and to "a tributary of the Rio Polochic...Alta Verapaz, Guatemala" [green star] by Dunn and Stuart (1951:59).

Type specimen: MNHN 1589, holotype, apparently lost, see Iverson (1983); type specimen figured in Duméril and Bocourt (1870:pl.5,f.3).

Etymology:* Latin, *pictus*, painted; referring to the bright colors of this turtle.

Superfamily Testudinoidea Batsch 1788 (21:52) (42)*

Testudinoid Turtles

(includes 4 families, 50 genera, 186 species, 284 taxa)

Testudines Batsch 1788:437

Testudinoidea Fitzinger 1826:5

Family Emydidae Rafinesque 1815 (09:12, 17:17)

Emydid Pond Turtles

(includes 2 subfamilies, 12 genera, 57 species, 91 taxa)

Emidania Rafinesque 1815:75


Emydes Schmid 1819:11

Emydidae Bell 1825a:302

Emydae Swainson 1839:113

Emydides Pictet 1853:446

Emididi Portis 1890:12



Emydidae Species Richness

Subfamily Deirochelyinae Agassiz 1857a (09:12, 17:17)

Deirochelyine Pond Turtles

(includes 6 genera, 42 species, 67 taxa)

Deirochelyoidae Agassiz 1857a:355

Deirochelyinae Gaffney and Meylan 1988:201

Chrysemys Gray 1844 (12:12)

Painted Turtles

(includes 2 species, 4 taxa)

Hydrochelys Wagler 1821:12 (*nomen oblitum*) (12:12)

Type species: *Hydrochelys picta* [= *Testudo picta* Schneider 1783], by original monotypy.

Etymology:* Greek, *hydor*, water, and *chelys*, turtle; referring to the aquatic habitat.

Comment:* Declared a *nomen oblitum* by Gemel and Grillitsch (2008), further confirmed and validated by Frétey (2024).

Chrysemys Gray 1844:27 (*nomen protectum*) (12:12)

Type species: *Emys* (*Chrysemys*) *picta* Schweigger [= *Testudo picta* Schneider 1783], by subsequent designation by Brown (1908:114).

Etymology:* Greek, *chrysos*, gold, and *emys*, turtle; referring to the bright yellowish golden color on the shell, head and neck.

Comment:* Declared a *nomen protectum* by Gemel and Grillitsch (2008), further confirmed and validated by Frétey (2024).


Chrysemys dorsalis Agassiz 1857a (07:11, 10:6, 17:18)

(or *Chrysemys picta dorsalis*)


Southern Painted Turtle



James H. Harding / Reelfoot Lake, Tennessee



left: James H. Harding / Reelfoot Lake, Tennessee
right: Mark Feldman / Louisiana / captive



** (stars = type localities)

Distribution: USA (*Alabama, Arkansas, Illinois, Kentucky, Louisiana, Mississippi, Missouri, Oklahoma, Tennessee, Texas*)

Introduced: USA (*Florida*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 323,135 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 588,056 sq. km

Size (Max SCL): male 11.5 cm, female 15.6 cm (Carr 1952; Ernst and Lovich 2009)

IUCN Red List: Least Concern (LC) (van Dijk 2011), as subspecies of *Chrysemys picta*

TFTSG Provisional Red List:* Near Threatened (NT) (Glorioso, Butterfield, and Munscher, in prep., 2025)

Synonymy:

Chrysemys dorsalis Agassiz 1857a:439,440 (07:11, 10:6)

Clemmys picta dorsalis, Chrysemys cinerea dorsalis, Chrysemys marginata dorsalis, Chrysemys bellii dorsalis,

Chrysemys picta dorsalis

Type locality: "Mississippi and Louisiana...Lake Concordia" [USA]. Restricted to "vicinity of New Orleans" [Louisiana, USA] [orange star] by Schmidt (1953:100), and to "vicinity of Natchez, Mississippi" [USA] [yellow star] by Ernst (1967:133).


Type specimen: MCZ 31960 (formerly 1802), lectotype, designated by Smith and Smith (1980:423); USNM 21, MCZ 1801, paralectotypes, see Barbour and Loveridge (1929), Cochran (1961), and Reynolds et al. (2007); Tiedemann et al. (1994) listed NMW 26218:1–2 as additional syntypes (possible paralectotypes).

Etymology:* Latin, *dorsum*, back; referring to the dorsal carapacial stripe.

Chrysemys picta (Schneider 1783) (07:11, 10:6, 12:13, 17:18) (43)*

Painted Turtle

(includes 3 subspecies)



(subspecies: *picta* = red, *bellii* = blue, *marginata* = green; overlap = intergrades; orange dots = probably introduced or translocated)

Distribution: Canada (*Alberta, British Columbia, Manitoba, New Brunswick, Nova Scotia, Ontario, Québec, Saskatchewan*), Mexico (*Chihuahua*), USA (*Alabama, Arizona, Colorado, Connecticut, Delaware, Georgia, Idaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Maine, Maryland, Massachusetts, Michigan, Minnesota, Missouri, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming*)

Introduced:* Canada (*Alberta, British Columbia, Manitoba, Saskatchewan*), Germany, Indonesia, Philippines, Spain, USA (*Arizona, California, Colorado, Florida, Nevada, Utah, Texas*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 4,569,908 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 8,716,480 sq. km

Size (Max SCL): male 22.5 cm, female 26.6 cm (see subspp.)

IUCN Red List: Least Concern (LC) (van Dijk 2011). Previously: Least Concern [Not Listed] (TFTSG 1996)


TFTSG Provisional Red List:* Least Concern (LC) (Moldowan, in prep., 2025)

Chrysemys picta picta (Schneider 1783) (07:11, 12:13) (43, 44)*


Eastern Painted Turtle



Anders G.J. Rhodin / Acadia National Park, Bar Harbor, Mt. Desert Island, Maine



Anders G.J. Rhodin / Acadia National Park, Pretty Marsh, Mt. Desert Island, Maine



(subspecies: *picta* = red, *bellii* = blue, *marginata* = green; overlap = intergrades; stars = type localities)

Distribution: Canada (*New Brunswick, Nova Scotia, Québec*), USA (*Alabama, Connecticut, Delaware, Georgia, Maine, Maryland, Massachusetts, New Hampshire, New Jersey, New York, North Carolina, Pennsylvania, Rhode Island, South Carolina, Tennessee, Vermont, Virginia*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 765,368 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 1,415,880 sq. km

Size (Max SCL): male 16.0 cm, female 19.0 cm (Rhodin and Mittelhauser 1994; Ultsch et al. 2000; A.G.J. Rhodin, unpubl. data)

Synonymy:

Testudo picta Schneider 1783:348

Emys picta, Hydrochelys picta, Clemmys picta, Terrapene picta, Emys (Chrysemys) picta, Chrysemys picta, Chrysemys picta picta, Pseudemys picta

Type locality: "England" [in error]. Restricted to "Lancaster, Pennsylvania" [USA] [yellow star] by Mittleman (1945:171); and to "vicinity of New York City" [New York, USA] [orange star] by Schmidt


(1953:99) [in error according to Smith and Smith (1980:424)].
Type specimens: UPSZTY 279 (formerly UUZM 279 and MGA 48), possible syntype, see Thunberg (1828), Holm (1957), and Wallin (2001); MZUS s/n, possible syntype, listed as possible "holotype" (Bour, pers. comm. *in* Iverson 1992); MCZ 1764, listed by Uetz et al. (2019) as a syntype, is not apparently a type.
Etymology:* Latin, *picta*, painted; referring to the bright colors of this turtle.
Comment: Original description cited as sourced from Hermann (pers. comm.), whose own description of *Testudo picta* was later published posthumously (Hermann 1804:219).

Testudo cinerea Bonnaterre 1789:25
Emys cinerea, Chrysemys cinerea, Chrysemys cinerea cinerea
Type locality: Not known. Restricted to "vicinity of Philadelphia" [Pennsylvania, USA] [green star] by Schmidt (1953:99).
Type specimen: Not located, type specimen figured in Brown (1776:pl.48,f.1-2).
Etymology:* Latin, *cinerea*, ash gray; referring to the gray color of some specimens, presumably preserved.
Comment: Description cited as sourced from Brown (1776:115,pl.48).

Chrysemys picta bellii (Gray 1830e) (10:7, 17:18, 21:52) (45)*
Western Painted Turtle


John B. Iverson / Oshkosh, Garden Co., Nebraska


John B. Iverson / *left*: nr. Galeana, Chihuahua, Mexico, *right*: Buffalo Co., Nebraska


**
(subspecies: *picta* = red, *bellii* = blue, *marginata* = green; overlap = intergrades; orange dots = probably introduced or translocated; stars = type localities)

Distribution: Canada (*Alberta, British Columbia, Manitoba, Ontario, Saskatchewan*), Mexico (*Chihuahua*), USA (*Arizona, Colorado, Idaho, Illinois, Iowa, Kansas, Michigan, Minnesota, Missouri, Montana, Nebraska, New Mexico, North Dakota, Oklahoma, Oregon, South Dakota, Texas, Utah, Washington, Wisconsin, Wyoming*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 2,732,375 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 5,445,182 sq. km
Size (Max SCL): male 20.8 cm, female 26.6 cm (MacCulloch and Secoy 1983; Rowe 1997; Ernst and Lovich 2009; Ceballos et al. 2013; Marchand et al. 2015)
Synonymy:
Emys bellii Gray 1830e:12 (10:7)
Clemmys (Clemmys) bellii, Emys (Chrysemys) bellii, Chrysemys bellii, Emys bellii, Chrysemys cinerea bellii, Chrysemys marginata bellii, Chrysemys bellii bellii, Chrysemys picta bellii
Type locality:* Not known. Restricted to "West coast of North America; British Columbia" [yellow star, location estimated] by Gray (1873a:147); erroneously to "Manhattan, Kans." [Kansas, USA] [orange star] by Smith and Taylor (1950b:34); and to "Puget Sound, Washington" [USA] [brown star, location estimated] by Schmidt (1953:100).
Type specimens: RCSM s/n, holotype, see Gray (1831d), destroyed during WW II (Iverson 1992); MCZ 1793, listed as "syntype" in MCZ online database, is not apparently a type.
Etymology:* Eponym honoring Thomas Bell (1792–1880), United Kingdom; herpetologist, dental surgeon, and Professor of Zoology at King's College, London.

Emys oregoniensis Harlan 1837:382
Chrysemys oregoniensis, Clemmys oregoniensis, Chrysemys oregonensis
Type locality: "fresh water ponds in the vicinity of the Oregon or Columbia River" [Oregon, USA] [green star, location estimated].
Type specimen: ANSP 165, holotype, see Malnate (1971).
Etymology:* Toponym for Oregon, USA.

Emys originensis Harlan *in* Gray 1844:23 (*nomen novum*)
Etymology:* Toponym for Oregon, USA.
Comment: Unjustified emendation or error for *oregoniensis*.

Chrysemys nuttalii Agassiz 1857a:451 (*nomen nudum*)
Etymology:* Eponym honoring Thomas Nuttal (1786–1859), United Kingdom / USA; naturalist and botanist.

Chrysemys nuttalii Agassiz 1857b:642
Type locality:* "Minesota and westward to the junction of the Yellowstone and Missouri" [Minnesota, USA and North Dakota, USA] [blue stars, syntype locations estimated].
Type specimens: MCZ 1771–78, syntypes (8), see MCZ online database.
Etymology:* Eponym honoring Thomas Nuttal (1786–1859), United Kingdom / USA; naturalist and botanist, who collected the types.


Chrysemys pulchra Gray 1873a:147
Type locality: "North America, Mississippi" [USA]. Restricted to "upper Mississippi River" [USA] [pink star, location estimated] by Schmidt (1953:100).
Type specimens: NHMUK 1946.1.22.26–27, 1946.1.22.93, 1947.3.4.23, 1947.3.4.75–77 (formerly 1844.11.16.5, 1852.9.8.7, 1855.4.12.3, 1855.12.6.9, 1858.9.9.6, 1861.8.21.5–6, 1862.6.5.1), syntypes (7).
Etymology:* Latin, *pulchra*, beautiful; referring to the ornate markings on the head and shell.

Chrysemys treleasei Hurter 1911:235
Type locality: "east side of the Mississippi River, in Madison, St. Clair, and Monroe Counties, Ill." [Illinois, USA] [gray star, one syntype location estimated, partially hidden].
Type specimens: USNM 49427–29, syntypes (3), see Cochran (1961) and Reynolds et al. (2007).
Etymology:* Eponym honoring William Trelease (1857–1945), USA; botanist, director of the Missouri Botanical Garden, and president of the Academy of Science of St. Louis.


Chrysemys picta marginata Agassiz 1857a [(44)*]
Midland Painted Turtle



Scott D. Gillingwater / Long Point, Norfolk Co., Ontario, Canada



left: Matthew G. Keevil / Algonquin Provincial Park, Ontario, Canada
right: Scott D. Gillingwater / Long Point, Norfolk Co., Ontario, Canada



(subspecies: *picta* = red, *bellii* = blue, *marginata* = green; overlap = intergrades; star = type locality)

Distribution: Canada (*Ontario, Québec*), USA (*Illinois, Indiana, Kentucky, Michigan, New York, North Carolina, Ohio, Pennsylvania, Tennessee, Vermont, Virginia, West Virginia*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 1,603,643 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 2,372,893 sq. km

Size (Max SCL): male 18.6 cm, female 19.5 cm (Ernst and Lovich 2009; Dolph 2017)

Synonymy:

Chrysemys marginata Agassiz 1857a:262,439
Clemmys marginata, Chrysemys marginata marginata, Chrysemys bellii marginata, Chrysemys picta marginata
Type locality: "Racine, Wisconsin...Milwaukee, Wisconsin...Flint, Michigan...Ann-Arbor, Michigan...Delphi, Indiana...Burlington, Iowa" [USA]. Restricted to "northern Indiana" [USA] [yellow star, location estimated] by Schmidt (1953:99).
Type specimens: MCZ 1780 (3 specimens), one of them designated as lectotype by Smith and Smith (1980:423); MCZ 1780 (3), 1789–90 (3), 1791 (3), 1796 (9), listed as syntypes by Barbour and Loveridge (1929), including one specimen from 1791 subsequently transferred to UMMZ; MCZ 1789–91, 1796, UMMZ 63519, syntypes (5), listed by Iverson (1992), but UIMNH 41529 (formerly MCZ 1796) also listed as a syntype by Smith et al. (1964); MCZ 1780, 1784–90, 1792–1800, 1899, 182588–90, 182593–606, listed as syntypes in MCZ online database; MCZ 1796, UMMZ 63519, UIMNH 41529, listed as syntypes by Uetz et al. (2019); NMW 26218:1,2 listed as syntypes by Gemel et al. (2019).
Etymology:* Latin, *marginis*, edge or border, and *-atus*, provided with, referring to the bright red markings of the carapacial margin.


Deirochelys Agassiz 1857a

Chicken Turtles
(includes 1 species, 3 taxa)

Deirochelys Agassiz 1857a:252,441
Type species: *Deirochelys reticulata* Schweigger [= *Testudo reticulata* Daudin 1801 = objective synonym of *Testudo reticularia* Latreille *in* Sonnini and Latreille 1801], by original monotypy.
Etymology:* Greek, *deire*, neck, throat, *chelys*, turtle; referring to the relatively long neck.

Hirochelys Beyer 1900:21 (*nomen novum*)
Etymology:* Greek, *deire*, neck, throat, *chelys*, turtle; referring to the relatively long neck.
Comment:* Unjustified emendation or error for *Deirochelys*.

Deirochelys reticularia (Latreille *in* Sonnini and Latreille 1801)
Chicken Turtle
(includes 3 subspecies)



(subspecies: *reticularia* = red, *chrysea* = blue, *miaria* = green; orange dots = questionable, probably translocated)

Distribution: USA (*Alabama, Arkansas, Florida, Georgia, Louisiana, Mississippi, Missouri, North Carolina, Oklahoma, South Carolina, Texas, Virginia*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 720,874 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 2,197,377 sq. km

Size (Max SCL): male 16.5 cm, female 26.0 cm (see subspp.)

CBFTT Account: Buhlmann, Gibbons, and Jackson (2008)

IUCN Red List: Least Concern (LC) [Not Listed] (Behler 1996)

TFTSG Provisional Red List:* Near Threatened (NT) (Hollender, McKnight, and Ligon, in prep., 2025). Previously: Near Threatened (NT) (2011)


Deirochelys reticularia reticularia (Latreille *in* Sonnini and Latreille 1801)
Eastern Chicken Turtle



Kurt A. Buhlmann / Virginia



left: Robert T. Zappalorti / Liberty Co., Florida
right: Kurt A. Buhlmann / CBFTT / Aiken Co., South Carolina



** (subspecies: *reticularia* = red, *chrysea* = blue, *miaria* = green; orange dots = questionable, probably translocated; star = type locality)

Distribution: USA (*Alabama, Florida, Georgia, Louisiana, Mississippi, North Carolina, South Carolina, Virginia*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 308,438 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 662,784 sq. km

Size (Max SCL): male 16.5 cm, female 26.0 cm (Buhlmann et al. 2008 CBFTT; Ernst and Lovich 2009)

Synonymy:

Testudo reticularia Latreille *in* Sonnini and Latreille 1801:124
Emys reticularia, Clemmys reticularia, Deirochelys reticularia, Deirochelys reticularia reticularia
Type locality: "Caroline" [USA]. Restricted to "Charleston" [South Carolina, USA] by Harper (1940:711); to "vicinity of Charleston, South Carolina" [USA] by Schmidt (1953:104); and to "9 miles northwest of Charleston, Charleston County, South Carolina" [USA] by neotype designation by Schwartz (1956a:466) [yellow star].
Type specimen: Originally in MNHN, apparently lost, type specimen figured in Sonnini and Latreille (1801:opp.124,f.1); NCSM 88836 (formerly CHM 54.48.1), neotype, designated by Schwartz (1956a:466); Zug and Schwartz (1971) erroneously listed the neotype as CHM 54.68.1.
Etymology:* Latin, *reticularis*, reticulated or net-like; referring to the carapacial pattern.

Testudo reticulata Bosc *in* Daudin 1801:144 (*nomen novum*)
Emys reticulata, Clemmys (Clemmys) reticulata, Deirochelys reticulata, Hirochelys reticulata, Chrysemys reticulata, Chrysemys reticulatus
Type locality: "Caroline" [USA]. Restricted to "vicinity of Charleston, South Carolina" [USA] [yellow star] by Schmidt (1953:104).
Type specimen: Originally in MNHN, apparently lost, type specimen figured in Sonnini and Latreille (1801:opp.124,f.1); NCSM 88836 (formerly CHM 54.48.1), neotype by default, designated by Schwartz (1956a:466).
Etymology:* Latin, *reticulata*, reticulated or net-like; referring to the carapacial pattern.
Comment: Unjustified replacement name for *reticularia*.


Deirochelys reticularia chrysea Schwartz 1956a
Florida Chicken Turtle



Kurt A. Buhlmann / CBFTT / central Florida



left: Benjamin K. Atkinson / CRM 3 / Hernando Co., Florida
right: Barry Mansell / CRM 3 / Glades Co., Florida



• (subspecies: *reticularia* = red, *chrysea* = blue; star = type locality)

Distribution: USA (*Florida*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 87,533 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 113,879 sq. km

Size (Max SCL): male 16.5 cm, female 25.0 cm (Schwartz 1956a; Ewert et al. 2006)

Synonymy:

Deirochelys reticularia chrysea Schwartz 1956a:476
Type locality: "5.8 miles east of Monroe Station, Collier County, Florida" [USA] [yellow star].
Type specimen: UMMZ 111440, holotype, see Kluge (1984).
Etymology:* Greek, *chryseos*, golden; referring to the plastral color pattern.


Deirochelys reticularia miaria Schwartz 1956a
Western Chicken Turtle



John L. Carr / Ouachita Parish, Louisiana



John L. Carr / Ouachita Parish, Louisiana



(subspecies: *reticularia* = red, *miaria* = green; star = type locality)

Distribution: USA (*Arkansas, Louisiana, Mississippi, Missouri, Oklahoma, Texas*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 324,903 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 610,946 sq. km

Size (Max SCL):* male 16.1 cm, female 23.2 cm (Schwartz 1956a; Bowers et al. 2022)

Synonymy:

Deirochelys reticularia miaria Schwartz 1956a:486
Type locality: "College Station, Brazos County, Texas" [USA] [yellow star].
Type specimen: FMNH 37478, holotype, see Marx (1958) and Lieb (2023).
Etymology:* Greek, *miaria*, stained or defiled; referring to the presence of a plastral pattern.

Graptemys Agassiz 1857a (12:14, 14:14, 17:19, 21:53)
Map Turtles
(includes 13 species, 14 taxa)

Graptemys Agassiz 1857a:252,436
Type species: *Graptemys geographica* [= *Testudo geographica* LeSueur 1817], by subsequent designation by Stejneger and Barbour (1917:117).
Etymology:* Greek, *graptos*, written, drawn, and *emys*, turtle; referring to the striking, complex carapace markings.

Neoclemmys Baur *in* Lindeman 2013:20 (*nomen nudum*)
Etymology:* Greek, *neo*, new, and *klemmys*, turtle.

Megaloclemmys Baur *in* Lindeman 2013:20 (*nomen nudum*)
Etymology:* Greek, *megalos*, big, and *klemmys*, turtle.


Graptemys barbouri Carr and Marchand 1942 (17:19, 17:21, 21:53)
Barbour's Map Turtle



David Dennis / CRM 3 / Florida



left: Matthew Aresco / CRM 3 / Leon Co., Florida / female
right: Robert T. Zappalorti / CRM 3 / Liberty Co., Florida / megacephalic female



(star = type locality)

Distribution: USA (*Alabama, Florida, Georgia*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 23,312 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 63,409 sq. km


Size (Max SCL): male 13.0 cm, female 32.7 cm (Pritchard

1980; Ewert et al. 2006)
IUCN Red List:* Near Threatened (NT A2bcde) (Mays, Smith, Godwin, and Hill 2025). Previously: Vulnerable (VU A2bcde) (van Dijk 2011); Near Threatened (NT) (TFTSG *in* Baillie and Groombridge 1996)
CITES:* Appendix II (2023). Previously: Appendix III (USA), as *Graptemys* spp. (2006)
USA ESA:* Threatened (by Similarity of Appearance) (USFWS 2024b); Species Status Assessment: USFWS (2017c)
Synonymy:
Graptemys barbouri Carr and Marchand 1942:98
Malaclemys barbouri
Type locality: "Chipola River north of Marianna, Jackson County, Florida" [USA] [yellow star].
Type specimen: MCZ 46251, holotype, see Barbour and Loveridge (1946).
Etymology:* Eponym honoring Thomas Barbour (1884–1946), USA; herpetologist and curator at the Museum of Comparative Zoology at Harvard University.

Graptemys caglei Haynes and McKown 1974 (17:19, 17:20, 21:53)
Cagle's Map Turtle, Guadalupe Map Turtle


James H. Harding / Texas / captivity



Viviana Ricardez / Kerr Co., Texas



** (orange dots = misidentified or translocated; star = type locality)

Distribution: USA (*Texas*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 9,015 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 23,473 sq. km
Size (Max SCL): male 12.6 cm, female 21.3 cm (Killebrew and Porter 1989a, 1990; Ernst and Lovich 2009)
IUCN Red List:* Endangered (EN A2c+4c; B2ab(ii,iii)) (Forstner and Mali, in press). Previously: Endangered (EN A2c+4c; B2ab(iii)) (van Dijk 2011); Vulnerable (VU) (TFTSG *in* Baillie and Groombridge 1996)
CITES: Appendix III (USA), as *Graptemys* spp. (2006)
Synonymy:
Graptemys caglei Haynes and McKown 1974:143
Type locality: "Guadalupe River, 8 km NW Cuero, DeWitt Co., Texas" [USA] [yellow star].
Type specimen: TNHC 36061, holotype, see Lieb (2023).
Etymology:* Eponym honoring Fred R. Cagle (1915–1968), USA; herpetologist at Tulane University, Louisiana.

Graptemys ernsti Lovich and McCoy 1992 (17:19, 17:21, 21:53)
Escambia Map Turtle


James C. Godwin / CBFTT / Yellow R., Covington Co., Alabama


James C. Godwin / CBFTT / Conecuh R., Covington Co., Alabama


** (star = type locality)


Distribution: USA (*Alabama, Florida*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 8,895 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 19,461 sq. km
Size (Max SCL): male 13.1 cm, female 28.5 cm (Lovich et al. 2011 CBFTT)
CBFTT Account: Lovich, Godwin, and McCoy (2011)
IUCN Red List:* Near Threatened (NT A2bc) (Godwin, Mays, and Hill 2025). Previously: Near Threatened (NT) (van Dijk 2011); Near Threatened (NT) (TFTSG *in* Baillie and Groombridge 1996)
CITES:* Appendix II (2023). Previously: Appendix III (USA), as *Graptemys* spp. (2006)
USA ESA:* Threatened (by Similarity of Appearance) (USFWS 2024b)
Synonymy:
Graptemys ernsti Lovich and McCoy 1992:300
Graptemys pulchra ernsti
Type locality: "Conecuh River, 1 mile upstream from County Road 4 Bridge, 14 km east of East Brewton, Escambia County, Alabama,

USA" [yellow star].
Type specimen: CM 122408, holotype.
Etymology:* Eponym honoring Carl H. Ernst (1938–2018), USA; herpetologist and turtle researcher at George Mason University, Virginia.

Graptemys flavimaculata Cagle 1954 (14:15, 17:19, 21:53)
Yellow-blotched Map Turtle, Yellow-blotched Sawback


Peter V. Lindeman / Merrill, Mississippi


Will Selman / CBFTT / Pascagoula River, Jackson Co., Mississippi


**
(orange dot = questionable, possibly translocated; star = type locality)


Distribution: USA (*Mississippi*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 10,605 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 16,990 sq. km
Size (Max SCL): male 12.3 cm, female 22.3 cm (Selman and Jones 2011 CBFTT; Ceballos et al. 2013)
CBFTT Account: Selman and Jones (2011)
IUCN Red List: Vulnerable (VU A2bce+4ce) (van Dijk 2011).
Previously: Endangered (EN) (TFTSG *in* Baillie and Groombridge 1996)
TFTSG Provisional Red List:* Vulnerable (VU) (Brown, Pearson, and Selman, in prep., 2025)
CITES: Appendix III (USA), as *Graptemys* spp. (2006)
USA ESA:* Threatened (USFWS 1991a)

Synonymy:
Graptemys flavimaculata Cagle 1954:167
Graptemys oculifera flavimaculata, Malaclemys flavimaculata
Type locality: "Pascagoula River, 13 miles S.W. of Lucedale, George Co., Mississippi" [USA] [yellow star]. Emended to "Pascagoula River at Old Benndale Crossing (T3S, R8W, Sec. 1), George County" [Mississippi, USA] by Cliburn (1971:17).
Type specimen: LSUM 111650 (formerly TU 14798), holotype.
Etymology:* Latin, *flavus*, yellow, and *maculatus*, spotted; referring to the orange spots on the carapacial scutes.

Graptemys geographica (LeSueur 1817) (08:18, 12:14, 17:19, 21:53)
Northern Map Turtle, Common Map Turtle


John B. Iverson / CBFTT / Dewart Lake, Kosciusko Co., Indiana


Grégory Bulté / CBFTT / Lake Opinicon, Ontario, Canada / male, female, female


**
(orange dots = probably translocated or vagrant; stars = type localities)


Distribution:* Canada (*Ontario, Québec*), USA (*Alabama, Arkansas, Delaware, Georgia, Illinois, Indiana, Iowa, Kansas, Kentucky, Louisiana, Maryland, Michigan, Minnesota, Mississippi, Missouri, New Jersey, New York, North Carolina, Ohio, Oklahoma, Pennsylvania, Tennessee, Texas, Vermont, Virginia, West Virginia, Wisconsin*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 726,581 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 2,485,265 sq. km
Size (Max SCL): male 16.0 cm, female 29.2 cm (Vogt et al.

2018 CBFTT)
CBFTT Account: Vogt, Bulté, and Iverson (2018)
IUCN Red List: Least Concern (LC) (van Dijk 2011). Previously: Least Concern (LC) [Not Listed] (Behler 1996)
TFTSG Provisional Red List:* Near Threatened (NT) (Pitt and Brown, in prep., 2025)
CITES: Appendix III (USA), as *Graptemys* spp. (2006)
Synonymy:
Testudo geographica LeSueur 1817:86
Emys geographica, Terrapene geographica, Clemmys (Clemmys) geographica, Clemmys geographica, Graptemys geographica, Malacoclemmys geographica, Malacoclemmys geographicus, Malaclemys geographica, Malaclemys geographicus, Malaclemmys geographicus, Graptemys geographicus
Type locality: "marsh, on the borders of Lake Erie" [USA]. Restricted to "peninsula of Presque Isle and adjacent Presque Isle Bay in Erie County, Pennsylvania" [USA] [yellow star] by Lindeman (2009:97).
Type specimen: MNHN, holotype or syntypes, not located, apparently lost, not listed by Roux-Esteve (1979) or Bonnemains and Bour (1996); type specimen figured (pl.5), designated lectotype by Lindeman (2009:96).
Etymology:* Greek, *gaia, geo-*, the earth, and *grapho*, drawing or writing; referring to the map-like carapacial markings.
Emys lesueurii Gray 1830e:12 (08:18, 10:7)
Emys leseurii, Graptemys lesueurii, Malacoclemmys lesueurii, Malaclemys lesueurii, Malaclemys lesueurii lesueurii
Type locality: "North America."
Type specimen: NHMUK 1982.1297, holotype.
Etymology:* Eponym honoring Charles Alexander LeSueur (1778–1846), France; naturalist, artist, and explorer who collected turtles in the United States.
Emys megacephala Holbrook 1836:51 (senior homonym, not = *Emys megacephalus* Gray 1844 [= *Malaclemys terrapin terrapin*], nor = *Emys megacephala* Gray 1870 [= *Malayemys macrocephala*])
Type locality: "Cumberland river...in the neighbourhood of Nashville, Tennessee" [USA] [green star].
Type specimen: ANSP 255, holotype, listed by Malnate (1971) as *Emys megalocephala*.
Etymology:* Greek, *megas*, great or large, and *kephale*, head; referring to the large head.
Emys macrocephala Agassiz 1857a:436 (*nomen novum*)
Comment: Unjustified emendation or error for *megacephala*.
Etymology:* Greek, *makros*, great or large, and *kephale*, head; referring to the large head.
Emys megalocephala Holbrook *in* Malnate 1971 (*nomen invalidum*)
Etymology:* Greek, *megale*, great or large, and *kephale*, head; referring to the large head.
Comment: Unjustified emendation or error for *megacephala*.

Graptemys gibbonsi Lovich and McCoy 1992 (10:8, 17:19, 21:53)
Pascagoula Map Turtle


Jeffrey E. Lovich / CBFTT / Chickasawhay R., nr. Leakesville, Mississippi / male


Will Selman / CBFTT / Leaf R., Mississippi / hatchling, female


(orange dots = questionable, possibly translocated; star = type locality)

Distribution:* USA (*Alabama*, *Mississippi*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 15,638 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 24,145 sq. km
Size (Max SCL): male 14.1 cm, female 29.5 cm (Lovich et al. 2009 CBFTT)
CBFTT Account: Lovich, Selman, and McCoy (2009)
IUCN Red List:* Endangered (EN A2bce+4ce) (Lindeman 2025). Previously: Endangered (EN A2bce+4ce) (van Dijk 2011); Near Threatened (NT) (TFTSG *in* Baillie and Groombridge 1996)
CITES:* Appendix II (2023). Previously: Appendix III (USA), as *Graptemys* spp. (2006)
USA ESA:* Threatened (by Similarity of Appearance) (USFWS 2024b); Species Status Assessment: USFWS (2021d)
Synonymy:
Graptemys gibbonsi Lovich and McCoy 1992:302
Graptemys pulchra gibbonsi
Type locality: "Chickasawhay River, Leakesville, Greene Co., Mississippi, USA" [yellow star].
Type specimen: CM 94979, holotype.
Etymology:* Eponym honoring J. Whitfield (Whit) Gibbons (1939–), USA; herpetologist and turtle researcher at Savannah River Ecology Laboratory, South Carolina.


Graptemys nigrinoda Cagle 1954 (17:19, 17:22, 21:53)
Black-knobbed Map Turtle, Black-knobbed Sawback



James C. Godwin / CBFTT / Alabama



James C. Godwin / CBFTT / Alabama / male, female, hatchling



** (stars = type localities)

Distribution: USA (*Alabama, Mississippi*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 39547 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 93,268 sq. km

Size (Max SCL): male 12.0 cm, female 22.1 cm (Blankenship et al. 2008 CBFTT; Ceballos et al. 2013)

CBFTT Account: Blankenship, Butterfield, and Godwin (2008)

IUCN Red List:* Least Concern (LC) (Godwin and Selman, in press). Previously: Least Concern (LC) (van Dijk 2011); Near Threatened (NT) (TFTSG *in* Baillie and Groombridge 1996)


CITES: Appendix III (USA), as *Graptemys* spp. (2006)

Synonymy:


Graptemys nigrinoda Cagle 1954:173 (17:22)
Graptemys oculifera nigrinoda, Graptemys nigrinoda nigrinoda, Malaclemys nigrinoda
Type locality: "Black Warrior River, above Lock 9, 17.5 miles SSW of Tuscaloosa, Tuscaloosa County, Alabama" [USA] [yellow star].
Type specimen: LSUM 111649 (formerly TU 14662), holotype.
Etymology:* Latin, *nigra*, black, and *nodus*, knot; referring to the black-knobbed carapace.

Graptemys nigrinoda delticola Folkerts and Mount 1969:677 (17:22)
Type locality: "Hubbard's Landing on Tensaw Lake, 2.6 air miles SW of Latham, Baldwin County, Alabama" [USA] [green star].
Type specimen: UF 26238 (formerly AUM 9229), holotype.
Etymology:* Greek, *delta*, river delta, and Latin, *colo*, to inhabit, referring to the occurrence of the taxon in the lower Mobile River drainage.


Graptemys oculifera (Baur 1890a) (17:19, 21:53) (46)*
Ringed Map Turtle, Ringed Sawback



Robert L. Jones / CCB / Pearl R., Madison Co., Mississippi



Robert L. Jones / CBFTT / Pearl R., Mississippi



** (orange dots = questionable, possibly translocated; stars = type localities)

Distribution: USA (*Louisiana, Mississippi*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 14,751 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 33,988 sq. km

Size (Max SCL): male 10.9 cm, female 21.5 cm (Jones and Selman 2009 CBFTT)

CBFTT Account: Jones and Selman (2009)

IUCN Red List: Vulnerable (VU B2ab(iii)) (van Dijk 2011). Previously: Endangered (EN) (TFTSG *in* Baillie and Groombridge 1996)

TFTSG Provisional Red List:* Endangered (EN) (Selman, Jones, and Jaunsen, in prep., 2025)

CITES: Appendix III (USA), as *Graptemys* spp. (2006)

USA ESA:* Threatened (USFWS 1986)

Synonymy:

Malacoclemmys oculifera Baur 1890a:262
Graptemys oculifera, Malaclemys lesueurii oculifera, Graptemys pseudogeographica oculifera, Graptemys oculifera oculifera, Malaclemys oculifera
Type locality:* "Mandeville, La." [Louisiana, USA] [yellow star].

Restricted to "Pearl River, 26 miles east of Mandeville" [Louisiana, USA] [in error, orange star] by Cagle (1953b:138); discussed by Glorioso et al. (2024).

Type specimen: USNM 15511, lectotype, designated by Lindeman (2013:342), also discussed by Cochran (1961) and Reynolds et al. (2007).

Etymology:* Latin, *oculus*, eye, and Greek, *phero*, to bear; referring to the ring-shaped ocellae on the pleural scutes.

Graptemys ouachitensis Cagle 1953a (12:14, 12:15, 14:16, 17:19, 21:53)
Ouachita Map Turtle



Peter V. Lindeman / CBFTT / Paris Landing State Park, Henry Co., Tennessee





left: Richard C. Vogt / CBFTT / Stoddard, Vernon Co., Mississippi River, Wisconsin / hatchling
right: John B. Iverson / Eufala, Pittsburg Co., Oklahoma



** (orange dots = questionable, possibly translocated; star = type locality)

Distribution: USA (*Alabama, Arkansas, Georgia, Illinois, Indiana, Iowa, Kansas, Kentucky, Louisiana, Minnesota, Mississippi, Missouri, Ohio, Oklahoma, Tennessee, Texas, West Virginia, Wisconsin*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 448,456 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 1,726,489 sq. km

Size (Max SCL): male 16.0 cm, female 26.0 cm (Vogt 2018 CBFTT)

CBFTT Account: Vogt (2018)

IUCN Red List: Least Concern (LC) (van Dijk 2011).
Previously: Least Concern (LC) [Not Listed] (Behler 1996)

TFTSG Provisional Red List:* Least Concern (LC) (Lindeman, in prep., 2025)

CITES: Appendix III (USA), as *Graptemys* spp. (2006)

Synonymy:

Testudo bigibbosa Rafinesque 1818:354 (*nomen nudum*) (21:7)
Type locality: "Ohio river" [USA].
Type specimen: Not located, type specimen figured in manuscript drawing by Rafinesque in 1818, reproduced by Bell and Bauer (2020:f.5).
Etymology:* Latin, *bi-*, twice, and *gibbosa*, humped, strongly arched; referring to two vertebral humps.


Emyda nodosa Rafinesque 1822:3 (*nomen nudum et novum*) (21:7)
Testudo nodosa
Etymology:* Latin, *nodus*, knot, bulge; referring to two vertebral humps.
Comment: Unjustifed replacement name for *bigibbosa*.

Graptemys pseudogeographica ouachitensis Cagle 1953a:10
Malaclemys pseudogeographica ouachitensis, Graptemys ouachitensis, Graptemys ouachitensis ouachitensis
Type locality: "Ouachita River, four miles northeast of Harrisonburg, Louisiana" [USA] [yellow star].
Type specimen: UMMZ 104345, holotype, see Kluge (1984).
Etymology:* Toponym for the Ouachita River, Louisiana, USA.


Graptemys pearlensis Ennen, Lovich, Kreiser, Selman, and Qualls 2010 (10:8, 17:19, 21:53)
Pearl River Map Turtle



Cris Hagen / CCB / Pearl River, Mississippi




Robert L. Jones / CBFTT / Pearl River, nr. Monticello, Mississippi



** (orange dots = possibly translocated; star = type locality)

Distribution: USA (*Louisiana, Mississippi*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 15,441 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 32,489 sq. km
Size (Max SCL): male 12.1 cm, female 29.5 cm (Ennen et al. 2016 CBFTT)
CBFTT Account: Ennen, Lovich, and Jones (2016)
IUCN Red List:* Endangered (EN A2bce+4ce) (Lindeman 2025). Previously: Endangered (EN A1bcde+A4bcde) (van Dijk 2011)
CITES:* Appendix II (2023). Previously: Appendix III (USA), as *Graptemys* spp. (2006)
USA ESA:* Threatened (USFWS 2024b); Species Status Assessment: USFWS (2023b)
Synonymy:
Graptemys pearlensis Ennen, Lovich, Kreiser, Selman, and Qualls 2010:104
Type locality: "Mississippi, Copiah County, Pearl River at State Highway 28, near Georgetown" [USA] [yellow star].
Type specimen: CM 62162, holotype.
Etymology:* Toponym for the Pearl River, Mississippi, USA.

Graptemys pseudogeographica (Gray 1831d) (12:14, 12:15, 17:19, 17:23, 21:53)
False Map Turtle
(includes 2 subspecies)



** (subspecies: *pseudogeographica* = red, *kohnii* = blue; overlap = intergrades; orange dots = introduced or translocated)

Distribution: USA (*Arkansas, Illinois, Indiana, Iowa, Kansas, Kentucky, Louisiana, Minnesota, Mississippi, Missouri, Nebraska, North Dakota, Oklahoma, South Dakota, Tennessee, Texas, Wisconsin*)
Introduced:* Germany, Italy, USA (*non-native range*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 592,401 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 1,982,016 sq. km
Size (Max SCL): male 15.0 cm, female 27.7 cm (see subspp.)
CBFTT Account:* Lindeman and Vogt (2024)
IUCN Red List: Least Concern (LC) (van Dijk 2011). Previously: Least Concern (LC) [Not Listed] (Behler 1996)
TFTSG Provisional Red List:* Least Concern (LC) (Lindeman, in prep., 2025)
CITES: Appendix III (USA), as *Graptemys* spp. (2006)

Graptemys pseudogeographica pseudogeographica (Gray 1831d)
(08:19, 21:53)

Northern False Map Turtle




James H. Harding / Indiana





left: John B. Iverson / Iowa
right: James H. Harding / Indiana



** (subspecies: *pseudogeographica* = red, *kohnii* = blue; overlap = intergrades; orange dots = introduced or translocated, star = type locality)

Distribution: USA (*Illinois, Indiana, Iowa, Kansas, Kentucky, Minnesota, Missouri, Nebraska, North Dakota, South Dakota, Wisconsin*)

Introduced: USA (*Florida, Virginia*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 233,933 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 989,321 sq. km

Size (Max SCL): male 15.0 cm, female 27.7 cm (Vogt 1995; Ceballos et al. 2013)

CBFTT Account:* Lindeman and Vogt (2024)

CITES: Appendix III (USA), as *Graptemys* spp. (2006)

Synonymy:

Emys pseudogeographica Gray 1831d:31

Clemmys pseudogeographica, Graptemys pseudogeographica, Malacoclemmys pseudogeographicus, Malaclemys pseudogeographica, Malaclemys pseudogeographicus, Graptemys pseudogeographicus, Graptemys pseudogeographica pseudogeographica, Malaclemys pseudogeographica pseudogeographica

Type locality: Not designated. Restricted to "Etats-Unis, Indiana, rivière Wabash, entre Mont Vernon et Chaumetown (= Shawneetown)...près du confluent de la Wabash et de l'Ohio" [USA] [yellow star] by Bour and Dubois (1983:45).

Type specimen: MNHN 9147, lectotype, designated by Bour and Dubois (1983:42).

Etymology:* Greek, *pseudo-*, false, *gaia*, *geo-*, the earth, and *grapho*, drawing or writing; referring to the map-like carapacial markings.

Graptemys pseudogeographica kohnii (Baur 1890a) (21:53)

Mississippi Map Turtle




Stanley E. Trauth / Strawberry R., Lawrence Co., Arkansas





left: John B. Iverson / Lake Hamilton, Garland Co., Arkansas
right: James H. Harding / No data / captivity / hatchling



** (subspecies: *pseudogeographica* = red, *kohnii* = blue; overlap = intergrades; orange dots = introduced or translocated; star = type locality)

Distribution: USA (*Arkansas, Kansas, Kentucky, Louisiana, Mississippi, Missouri, Oklahoma, Tennessee, Texas*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 476,062 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 1,375,217 sq. km

Size (Max SCL): male 15.0 cm, female 27.0 cm (R.C. Vogt, unpubl. data)

CBFTT Account:* Lindeman and Vogt (2024)

CITES: Appendix III (USA), as *Graptemys* spp. (2006)

Synonymy:

Malacoclemmys kohnii Baur 1890a:263

Graptemys kohnii, Malaclemys lesueurii kohnii, Graptemys pseudogeographica kohnii, Malaclemys kohnii

Type locality: "Bayou Lafourche, La.; Bayou Teche, St. Martinsville, La." [Louisiana, USA] [yellow star].

Type specimen: TU 16409, lectotype, designated by Lindeman (2013:321).

Etymology:* Eponym honoring Gustave Kohn (1837–1906), American-France; amateur ornithologist and naturalist who collected the types.


Graptemys pulchra Baur 1893c (14:17, 17:19, 21:53)
Alabama Map Turtle



James C. Godwin / CBFTT / Tallapoosa R., Elmore Co., Alabama



James C. Godwin / CBFTT / Tallapoosa R., Elmore Co., Alabama



(orange dot = translocated or vagrant; star = type locality)

Distribution: USA (*Alabama, Georgia, Mississippi, Tennessee*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 47,688 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 128,237 sq. km

Size (Max SCL): male 12.2 cm, female 27.9 cm (Lovich et al. 2014 CBFTT; Coleman 2020)

CBFTT Account: Lovich, Godwin, and McCoy (2014)

IUCN Red List:* Near Threatened (NT A2bcd+4bcd) (Godwin and Brown 2025). Previously: Near Threatened (NT) (van Dijk 2011); Least Concern (LC) (TFTSG 1996)

CITES:* Appendix II (2023). Previously: Appendix III (USA), as *Graptemys* spp. (2006)

USA ESA:* Threatened (by Similarity of Appearance) (USFWS 2024b)

Synonymy:

Graptemys pulchra Baur 1893c:675

Malacoclemmys pulchra, Malaclemys lesueurii pulchra, Malaclemys pulchra, Graptemys pulchra pulchra

Type locality: "Montgomery, Alabama" [USA] [yellow star].

Type specimen: USNM 8808, lectotype, designated by Lovich and McCoy (1992:304), see Reynolds et al. (2007) and Lindeman (2013).

Etymology:* Latin, *pulchra*, beautiful; referring to the ornate markings on the head and shell.

Graptemys alabamensis Baur *in* Lindeman 2013:20 (*nomen nudum*) (14:17)

Etymology:* Toponym for Alabama, USA.


Graptemys grandis Baur *in* Lindeman 2013:20 (*nomen nudum*) (14:17)

Etymology:* Latin, *grandis*, great, significant; possibly referring to the size of the species.


Graptemys sabinensis Cagle 1953a (07:12, 12:15, 14:18, 17:19, 21:53)
Sabine Map Turtle



Peter V. Lindeman / CCB / CBFTT / nr. Estherwood, Acadia Parish, Louisiana



left: Peter V. Lindeman / CBFTT / Calcasieu R., Louisiana
right: Richard C. Vogt / CBFTT / Sabine River, nr. Many, Louisiana



(orange dots = questionable, possibly translocated; star = type locality)

Distribution: USA (*Louisiana, Texas*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 41,168 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 67,820 sq. km

Size (Max SCL): male 10.6 cm, female 15.5 cm (Vogt 2018 CBFTT)

CBFTT Account: Vogt (2018), as subspecies of *Graptemys ouachitensis*

IUCN Red List: Least Concern (LC) (van Dijk 2011), as subspecies of *Graptemys ouachitensis*

TFTSG Provisional Red List:* Least Concern (LC) (Lindeman, in prep., 2025)

CITES: Appendix III (USA), as *Graptemys* spp. (2006)

Synonymy:

Malacoclemmys intermedia Baur *in* Beyer 1900:21 (*nomen nudum*) (14:19)

Graptemys intermedia

Etymology:* Latin, *intermedia*, in-between; referring to the relationship of this species to others.

Graptemys pseudogeographica sabinensis Cagle 1953a:2

Malaclemys pseudogeographica sabinensis, Graptemys ouachitensis sabinensis, Graptemys sabinensis

Type locality: "Sabine River, eight miles southwest of Negreet, Louisiana" [USA] [yellow star].

Type specimen: UMMZ 104351, holotype, see Kluge (1984).


Etymology:* Toponym for the Sabine River, Louisiana, USA.

Graptemys versa Stejneger 1925 (17:19, 21:53)


Texas Map Turtle



Peter V. Lindeman / CBFTT / Live Oak Creek, Gillespie Co., Texas



left: Peter V. Lindeman / CBFTT / South Llano R., East Johnson Fork, Texas
right: Terry Hibbitts / CBFTT / Colorado R., Concho Co., Texas



** (orange dots = questionable, possibly translocated; star = type locality)

Distribution: USA (*Texas*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 35,249 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 81,884 sq. km

Size (Max SCL): male 11.6 cm, female 21.4 cm (Kizirian et al. 1991; Lindeman et al. 2016 CBFTT)

CBFTT Account: Lindeman, Stuart, and Killebrew (2016)

IUCN Red List: Least Concern (LC) (van Dijk 2011). Previously: Near Threatened (NT) (TFTSG *in* Baillie and Groombridge 1996)

TFTSG Provisional Red List:* Least Concern (LC) (Lindeman, in prep., 2025)

CITES: Appendix III (USA), as *Graptemys* spp. (2006)

Synonymy:

Graptemys pseudogeographica versa Stejneger 1925:463

Graptemys versa, Malaclemys versa

Type locality: "Austin, Texas" [USA] [yellow star].

Type specimen: USNM 27473, holotype, see Cochran (1961), Reynolds et al. (2007), and Lieb (2023).

Etymology:* Latin, *versus*, to change; referring to the difference in the head pattern from that of the other subspecies.

Malaclemys Gray 1844

Diamondback Terrapins

(includes 1 species, 7 taxa)

Malaclemys Gray 1844:28

Type species: *Malaclemys concentrica* [= *Testudo concentrica* Shaw 1802 = subjective synonym of *Testudo terrapin* Schoepff 1793a], by original monotypy.

Etymology:* Greek, *malakos*, soft, and *klemmys*, turtle; apparently referring to the ability to retract and hide the head and neck

Maloclemmys Agassiz 1847:222 (*nomen novum*)

Etymology:* Greek, *malakos*, soft, and *klemmys*, turtle.

Comment: Unjustified replacement name or error for *Malaclemys*.

Malacoclemmys Agassiz 1857a:392,437 (*nomen novum*)

Etymology:* Greek, *malakos*, soft, and *klemmys*, turtle.

Comment: Unjustified replacement name or error for *Malaclemys*.

Euchyloclemmys Sclater 1858:292 (*nomen novum*)

Etymology:* Greek, *eu-*, true, good, *chylos*, juices, and *klemmys*, turtle; referring to the culinary appeal and use of the genus.

Euchyloclemys Sclater *in* Gray 1863c:181 (*nomen novum*)

Etymology:* Greek, *eu-*, true, good, *chylos*, juices, and *klemmys*, turtle; referring to the culinary appeal and use of the genus.

Comment: Unjustified replacement name or error for *Euchyloclemmys*.

Malaclemmys Gray 1870c:41 (*nomen novum*)

Etymology:* Greek, *malakos*, soft, and *klemmys*, turtle.

Comment: Unjustified replacement name or error for *Malaclemys*.

Malaclemys terrapin (Schoepff 1793a) (11:5, 17:24, 21:54) (47)*

Diamondback Terrapin

(includes 7 subspecies)



** (subspecies: *terrapin* = red, *centrata* = blue, *littoralis* = orange, *macrospilota* = green, *pileata* = pink, *rhizophorarum* = tourmaline, *tequesta* = purple; orange dots = trade or extralimital)

Distribution: Bermuda, USA (*Alabama, Connecticut, Delaware, Florida, Georgia, Louisiana, Maryland, Massachusetts, Mississippi, New Jersey, New York, North Carolina, Rhode Island, South Carolina, Texas, Virginia*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 124,589 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 2,089,477 sq. km

Size (Max SCL): male 18.0 cm, female 32.0 cm (see subspp.)

IUCN Red List: Vulnerable (VU A4acde) (Roosenburg, Baker, Burke, Dorcas, and Wood 2019). Previously: Near Threatened (NT) (TFTSG 1996)

CITES: Appendix II (2013)

Malaclemys terrapin terrapin (Schoepff 1793a)
Northern Diamondback Terrapin



Joseph C. Mitchell / Fisherman Island, Northampton Co., Virginia



Russell L. Burke / Jamaica Bay, Long Island, New York



(subspecies: *terrapin* = red, *centrata* = blue; orange dot and star = trade and type locality; stars = type localities)

Distribution: USA (*Connecticut, Delaware, Massachusetts, New Jersey, New York, North Carolina, Maryland, Rhode Island, Virginia*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 39,317 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 244,089 sq. km

Size (Max SCL): male 18.0 cm, female 32.0 cm (Hurd et al. 1979; Lovich and Hart 2018; Roosenburg et al. 2019 IUCN)

CITES: Appendix II (2013)

Synonymy:

Testudo terrapin Schoepff 1793a:64
Emys terrapin, Clemmys terrapin, Malaclemys terrapin, Malacoclemmys terrapin, Malacoclemmys terrapen, Malaclemys terrapin terrapin
Type locality: "America septentrionali...in foris Philadelphiae...et... aquis subdulcibus Insulae Longae" [USA]. Restricted to "probably Delaware Bay" [Delaware and New Jersey, USA] [green star, location estimated] by Hay (1905:16); and to "coastal waters of Long Island" [New York, USA] [yellow star, location estimated] by Schmidt (1953:95).
Type specimens: Not located, type specimen figured (pl.15).
Etymology:* Vernacular (Native American, Powhatan), *torope, terrapin, terrapen,* or *turpin,* indigenous general name for a turtle, see Dodd et al. (2016).

Testudo concentrica Shaw 1802:43
Emys concentrica, Malaclemys concentrica, Malaclemmys concentrica, Malaclemmys centrata concentrica, Malaclemys centrata concentrica, Malaclemys terrapin concentrica
Type locality: "North America...sold in the markets at Philadelphia" [Pennsylvania, USA] [orange star]. Restricted to "probably Delaware Bay" [Delaware or New Jersey, USA] [green star, location estimated] by Hay (1905:16).
Type specimens: Not located, type specimen figured (f.9,top).
Etymology:* Latin, *concentrica*, concentric; referring to the characteristic concentric growth rings on the carapace scutes.

Testudo ocellata Link 1807:52
Type locality: "Nord-Amerika." Restricted to "Philadelphia markets" [Pennsylvania, USA] [orange star] by Schmidt (1953:96).
Type specimens: Not located, type specimen figured in Bechstein (1800:pl.4,f.2) and Schoepff (1793a:pl.15).
Etymology:* Latin, *ocellus*, little eye, *ocellata*, having eyespots; referring to the ocellated eye-like markings on the carapace.

Emys concentrica polita Gray 1830e:11 (10:7)
Type locality: Not designated.
Type specimens: OUM 8475, syntype (1), discussed by Nowak-Kemp and Fritz (2010).
Etymology:* Latin, *polita*, polished; referring to polished black appearance of the carapace of this taxon.

Testudo palustris Le Conte 1830:113 (junior homonym and *nomen novum*, not = *Testudo palustris* Gmelin 1789 [= *Trachemys terrapen*])
Emys palustris, Malacoclemmys palustris
Type locality: "New-York to Florida, and even in the West Indies, in salt water" [USA].
Type specimen: Possibly ANSP, apparently lost, not listed by Malnate (1971).
Etymology:* Latin, *palustris*, swamp-dwelling; referring to the habitat in marshes.
Comment: Unjustified replacement name for *centrata*.

Emys macrocephalus Gray 1844:26 (junior homonym, not = *Emys macrocephala* Spix 1824 [= *Peltocephalus dumerilianus*])
Emys macrocephala
Type locality: "America?" Restricted to "Philadelphia markets" [Pennsylvania, USA] [orange star] by Schmidt (1953:96).
Type specimen: NHMUK, possibly lost, see Ernst and Bury (1982).
Etymology:* Greek, *makros*, great or large, and *kephale*, head; referring to the large head.
Comment: Description based on a lost manuscript ("Gray, Desc. Cat. Rept. 13, n. 33" = Gray, J.E. No date. Descriptive Catalogue of Reptiles), where it was originally named *Emys megacephalus*.


Emys megacephalus Gray 1844:26 (junior homonym and *nomen nudum*, not = *Emys megacephala* Holbrook 1836 [= *Graptemys geographica*], nor = *Emys megacephala* Gray 1870 [= *Malayemys macrocephala*])
Etymology:* Greek, *megalos*, great or large, and *kephale*, head; referring to the large head.

Malaclemys terrapin centrata (Latreille *in* Sonnini and Latreille 1801) [(11:5)]

Carolina Diamondback Terrapin



Carla Van Ness / CRM 3 / Duval Co., Florida



** (subspecies: *terrapin* = red, *centrata* = blue, including Bermuda, *tequesta* = purple; star = type locality)

Distribution: Bermuda, USA (*Georgia, Florida, North Carolina, South Carolina*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 17,971 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 105,252 sq. km

Size (Max SCL): male 15.0 cm, female 20.4 cm (Gibbons et al. 2001; Butler 2002)

CITES: Appendix II (2013)

Synonymy:

Testudo centrata Latreille *in* Sonnini and Latreille 1801:145

Emys centrata, Clemmys (Clemmys) centrata, Malaclemmys centrata, Malaclemys centrata, Malaclemys centrata centrata, Malaclemmys terrapin centrata, Malaclemys terrapin centrata

Type locality: “les grands marais de la Caroline” [USA]. Restricted to “neighborhood of Charleston, South Carolina” [USA] [yellow star] by Hay (1905:14).

Type specimen: Not located, type specimen figured in Sonnini and Latreille (1801:pl.opp.124,f.2).

Etymology:* Latin, *centrum*, center; referring to the concentric growth annuli on the carapacial scutes.

Emys concentrica livida Gray 1831d:27

Emys livida

Type locality: “America Boreali.” Restricted to “vicinity of Charleston, South Carolina” [USA] [yellow star] by Schmidt (1953:96).

Type specimen: OUM s/n, holotype, apparently lost, discussed by Nowak-Kemp and Fritz (2010).


Etymology:* Latin, *livida*, bluish or lead-colored; referring to the shell color.

Malaclemys terrapin littoralis (Hay 1905)


Texas Diamondback Terrapin



Aaron S. Baxter / Nueces Bay, Nueces Co., Texas



John L. Carr / Cameron Parish, Louisiana



** (subspecies: *littoralis* = orange, *pileata* = pink; orange dot = extralimital, star = type locality)

Distribution: USA (*Louisiana, Texas*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 14,416 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 47,684 sq. km

Size (Max SCL): male 14.8 cm, female 20.1 cm (Lovich and Hart 2018)

CITES: Appendix II (2013)

Synonymy:

Malaclemmys littoralis Hay 1905:18

Malaclemys centrata littoralis, Malaclemys pileata littoralis, Malaclemys terrapin littoralis

Type locality: “Rockport, Texas” [USA] [yellow star].

Type specimen: USNM 33913, holotype, see Cochran (1961), Reynolds et al. (2007), and Lieb (2023).

Etymology:* Latin, *littoralis*, belonging to the seashore; referring to the coastal estuarine habitat.

Malaclemys terrapin macrospilota (Hay 1905)
Ornate Diamondback Terrapin



George L. Heinrich / St. Marks NWR, Wakulla Co., Florida



left: Richard D. Bartlett / CRM 3 / Lee Co., Florida / hatchling
right: George L. Heinrich / Lee Co., Florida



(subspecies: *centrata* = blue, *macrospilota* = green, *rhizophorarum* = tourmaline, *tequesta* = purple; star = type locality)

Distribution: USA (*Florida*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 17,120 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 74,645 sq. km

Size (Max SCL): female 20.8 cm (B.K. Mealey, unpubl. data)

CITES: Appendix II (2013)


Synonymy:

Malaclemmys macrospilota Hay 1905:16
Malaclemys centrata macrospilota, Malaclemys pileata macrospilota, Malaclemys terrapin macrospilota
Type locality: "Charlotte Harbor, Florida" [USA] [yellow star].
Type specimen: USNM 33917, holotype, see Cochran (1961) and Reynolds et al. (2007).
Etymology:* Greek, *makros*, large, and *spilotos*, soiled or stained; referring to the large yellow spot on each carapacial scute.


Malaclemys terrapin pileata (Wied-Neuwied 1865)
Mississippi Diamondback Terrapin



Andrew T. Coleman / CCB / Bayou Caddy, Hancock Co., Mississippi



left: Jennifer Frey / Jackson Co., Mississippi
right: Grover J. Brown / Jackson Co., Mississippi



(subspecies: *littoralis* = orange, *macrospilota* = green, *pileata* = pink; stars = type localities)

Distribution: USA (*Alabama, Florida, Louisiana, Mississippi*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 27,662 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 77,480 sq. km

Size (Max SCL): male 15.3 cm, female 24.3 cm (Selman et al. 2019; W. Selman, unpubl. data; T.M. Mann and P.S. Floyd, unpubl. data)

CITES: Appendix II (2013)


Synonymy:

Emys pileata Wied-Neuwied 1865:17
Malaclemmys pileata, Malaclemys centrata pileata, Malaclemys pileata pileata, Malaclemys terrapin pileata
Type locality: "Sümpfen mit salzigem Wasser an der Mündung des Mississippi bei New-Orleans" [Louisiana, USA] [orange star]. Emended to "New Orleans, Louisiana" [USA] [yellow star] by Hay (1905:17).
Type specimen: AMNH 916, lectotype, designated as "type?" by Hay (1905:17), and as lectotype by Fritz et al. (1994:167); Uetz et al. (2019) erroneously list AMNH 7064 as holotype.
Etymology:* Latin, *pileatus*, capped, covered with a cap; referring to the black dorsal head surface.


Malaclemys terrapin rhizophorarum (Fowler 1906)
Mangrove Diamondback Terrapin



Brian K. Mealey / CRM 3 / Monroe Co., Florida



left: United States Geological Survey / Mathew Denton / Calusa Keys, Monroe Co., Florida
right: United States Geological Survey / Kristen M. Hart / Everglades, Monroe Co., Florida



** (subspecies: *macrospilota* = green, *rhizophorarum* = tourmaline, *tequesta* = purple; stars = type localities)

Distribution: USA (*Florida*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 3,372 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 17,233 sq. km

Size (Max SCL): male 14.2 cm, female 20.1 cm (Hart and McIvor 2008; Lovich and Hart 2018)

CITES: Appendix II (2013)

Synonymy:


Malaclemys tuberculifera Gray 1844:29 (*nomen oblitum*)
Type locality: "California" [in error]. Restricted to "Philadelphia market" [Pennsylvania, USA] [in error] by Schmidt (1953:96); shown to be from the Florida Keys [Florida, USA] [orange star, location estimated] by Ernst and Hartsell (2000:887), but they did not formally restrict the type locality.
Etymology:* Latin, *tuberculi*, small humps, and *ferre*, to bear, to carry; referring to the vertebral keel having small distinct hump-like projections.
Type specimen: NHMUK 1947.3.5.29, holotype.

Malaclemmys littoralis rhizophorarum Fowler 1906:112
Malaclemmys terrapin rhizophorarum, Malaclemys terrapin rhizophorarum
Type locality: "Boca Grande Key, Florida" [USA] [yellow star].
Type specimen: ANSP 16570, holotype, see Malnate (1971).
Etymology:* Bionym for *Rhizophora*, genus of the Red Mangrove; and Greek, *rhizo*, roots; *phora*, bear; and Latin, *-arum*; referring to this species' habitat in mangrove swamps.

Malaclemys terrapin tequesta Schwartz 1955
Eastern Florida Diamondback Terrapin



Richard D. Bartlett / CRM 3 / Martin Co., Florida



** (subspecies: *centrata* = blue, *macrospilota* = green, *rhizophorarum* = tourmaline, *tequesta* = purple; star = type locality)

Distribution: USA (*Florida*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 6,974 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 25,946 sq. km

Size (Max SCL): male 14.7 cm, female 22.0 cm (Butler et al. 2006; R.A. Seigel, unpubl. data)

CITES: Appendix II (2013)

Synonymy:


Malaclemys terrapin tequesta Schwartz 1955:158
Type locality: "Miami Beach, Dade County, Florida" [USA] [yellow star].
Type specimen: UMMZ 108482, holotype, see Kluge (1984).
Etymology:* Ethnonym honoring the indigenous Tequesta people who previously lived in this area.

Pseudemys Gray 1856a (09:13, 12:16, 14:20)
Cooter Turtles, Cooters
(includes 8 species, 9 taxa)
Pseudemys Gray 1856a:197
Type species: *Pseudemys concinna* [= *Testudo concinna* LeConte 1830], by subsequent designation by Baur (1893a:221).
Etymology:* Greek, *pseudo*, false, untrue, and *emys*, turtle; referring to the view by Gray that this genus is very similar to the genus *Emys*.
Ptychemys Agassiz 1857a:252,431
Type species: *Ptychemys concinna* [= *Testudo concinna* LeConte 1830], by subsequent designation by Brown (1908:114).
Etymology:* Greek, *ptychi*, wrinkle, fold, layer, and *emys*, turtle; referring to the several large scales on the outer edge of the front legs.
Nectemys Agassiz 1857b:642 (*nomen novum*)
Etymology:* Greek, *nektos*, swimming, and *emys*, turtle; referring to the aquatic habitat and swimming behavior.
Comment:* Unjustified provisional replacement name for *Ptychemys*.

Pseudemys alabamensis Baur 1893a (48)*
Alabama Red-bellied Turtle, Alabama Red-bellied Cooter


Robert H. Mount / CCB / CBFTT / Alabama / female


left: Thomas M. Mann / CBFTT / West Pascagoula River, Mississippi / male
right: James L. Dobie / CBFTT / Mississippi / male


**
(orange dots = waifs on offshore islands and translocations; star = type locality)

Distribution: USA (*Alabama, Mississippi*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 4,089 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 9,647 sq. km
Size (Max SCL): male 29.5 cm, female 37.5 cm (Leary et al. 2008 CBFTT)
CBFTT Account: Leary, Dobie, Mann, Floyd, and Nelson (2008)
IUCN Red List:* Endangered (EN A2c+4ce; B1ab(i,ii,iii,iv)) (Godwin and Pearson, in press). Previously: TFTSG Provisional Red List: Endangered (EN) (2011); Endangered (EN B1+2c [ver. 2.3]) (TFTSG 1996)
USA ESA:* Endangered (USFWS 1987b)
Synonymy:
Pseudemys alabamensis Baur 1893a:224
Pseudemys rubriventris alabamensis, Chrysemys (Pseudemys) alabamensis, Chrysemys rubriventris alabamensis
Type locality: "Mobile bay, Ala." [Alabama, USA] [yellow star].
Type specimens: USNM 20966–67, syntypes (2), see Cochran (1961) and Reynolds et al. (2007); USNM 20966 listed as "holotype" by McCoy and Vogt (1985:1).
Etymology:* Toponym for Alabama, USA.
Pseudemys alabamiensis Beyer 1900:20 (*nomen nudum*)
Etymology:* Toponym for Alabama, USA.
Comment:* Unjustified emendation or error for *alabamensis*.

Pseudemys concinna (Le Conte 1830) (09:13, 17:25)
River Cooter
(includes 2 subspecies)



**
(subspecies: *concinna* = red, *suwanniensis* = blue; orange dots = possibly translocated or vagrant)

Distribution: USA (*Alabama, Arkansas, Florida, Georgia, Illinois, Indiana, Kansas, Kentucky, Louisiana, Mississippi, Missouri, North Carolina, Ohio, Oklahoma, Pennsylvania (?), South Carolina, Tennessee, Texas, Virginia, West Virginia*)
Introduced:* Germany, South Korea
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 631,311 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 2,162,022 sq. km
Size (Max SCL):* male 40.2 cm, female 47.6 cm (see subspp.)
CBFTT Account: Ward and Jackson (2008)
IUCN Red List: Least Concern (LC) (van Dijk 2011). Previously: Least Concern (LC) [Not Listed] (Behler 1996)
TFTSG Provisional Red List:* Least Concern (LC) (Johnston and Godwin, in prep., 2025)


Pseudemys concinna concinna (Le Conte 1830) (07:13, 09:13, 10:9, 17:25)
Eastern River Cooter



Peter A. Meylan / CBFTT / CRM 3 / Spring Creek, Decatur Co., Georgia



left: Joseph P. Ward / CBFTT / Flint River, Albany, Madison Co., Alabama
right: Joseph P. Ward / CBFTT / Cahaba River, Bibb Co., Alabama



** (subspecies: *concinna* = red, *suwanniensis* = blue; orange dots = possibly translocated or vagrant; stars = type localities)

Distribution: USA (*Alabama, Arkansas, Florida, Georgia, Illinois, Indiana, Kansas, Kentucky, Louisiana, Mississippi, Missouri, North Carolina, Ohio, Oklahoma, Pennsylvania (?), South Carolina, Tennessee, Texas, Virginia, West Virginia*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 605,401 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 1,864,610 sq. km

Size (Max SCL): male 30.3 cm, female 42.0 cm (Buhlmann and Vaughan 1991; Ward and Jackson 2008 CBFTT)

Synonymy:

Testudo concinna Le Conte 1830:106 (junior homonym, not = *Emys concinna* Cuvier *in* Guérin 1829 (*nomen oblitum*) [= *Trachemys scripta elegans*]) (17:25)
Emys (Testudo) concinna, Terrapene concinna, Clemmys (Clemmys) concinna, Pseudemys concinna, Ptychemys concinna, Chrysemys concinna, Pseudemys floridana concinna, Pseudemys concinna concinna, Chrysemys concinna concinna
Type locality: "rivers of Georgia and Carolina, where the beds are rocky...never...below Augusta on the Savannah, or Columbia on the Congaree" [USA]. Restricted to "vicinity of Columbia, South Carolina" [USA] [yellow star] by Schmidt (1953:101).
Type specimen: MNHN 9172 (with body parts MNHN 2066 of the same specimen), lectotype, designated by Bour (2003:543).
Etymology:* Latin, *concinnus*, pretty, delicate, elegant; referring to the elegant carapacial markings.

Emys annulifera Gray 1830e:12 (10:7)
Trachemys annulifera
Type locality: Not known. Restricted to "Columbia, South Carolina" [USA] [yellow star] by Schmidt (1953:101).
Type specimen: NHMUK 1946.1.22.28, holotype, see Boulenger (1889:84).
Etymology:* Latin, *annulus*, a ring, and *fero*, to bear; referring to the carapacial rings.

Emys hieroglyphica Holbrook 1836:47
Pseudemys hieroglyphica, Ptychemys hieroglyphica, Clemmys hieroglyphica, Chrysemys hieroglyphica, Pseudemys floridana hieroglyphica, Pseudemys concinna hieroglyphica, Chrysemys concinna hieroglyphica
Type locality: "Cumberland river" [Tennessee, USA]. Restricted to "Cumberland River [Nashville, Davidson Co., Tennessee]" [green star] by Ward (1984:38).
Type specimen: ANSP 217, holotype, see Baur (1893a) and Carr (1938b), not listed by Malnate (1971), apparently lost (Iverson 1992).
Etymology:* Greek, *hieros*, sacred, and *glypho*, carve, engrave; referring to the intricate markings on the shell.

Emys mobilensis Holbrook 1838a:53
Ptychemys mobilensis, Clemmys mobilensis, Pseudemys mobilensis, Chrysemys mobilensis, Pseudemys floridana mobilensis, Pseudemys concinna mobilensis, Chrysemys concinna mobilensis
Type locality: "Alabama...in the neighbourhood of Mobile" [USA] [blue star].
Type specimen: ANSP 242, holotype, see Malnate (1971).
Etymology:* Toponym for Mobile, Alabama, USA.

Emys labyrinthica Duméril and Bibron *in* Duméril and Duméril 1851:13
Clemmys labyrinthica, Pseudemys labyrinthica, Chrysemys labyrinthica
Type locality: "Wabash-River (États-Unis)" [Indiana, USA] [pink star, location estimated].
Type specimens: MNHN 9179–81, 9521, syntypes (4), see Roux-Esteve (1979) and Bonnemains and Bour (1996).
Etymology:* Greek, *labyrinthos*, labyrinth; referring to the elaborate markings on the carapace and extremities.

Ptychemys hoyi Agassiz 1857a:433,
Pseudemys concinna hoyi, Pseudemys floridana hoyi, Chrysemys floridana hoyi
Type locality: "south-western Missouri" [USA]. Restricted by neotype designation to "Newton County, Missouri" [USA] [light gray star] by Stejneger (1938:175); invalid restriction to "vicinity of Springfield, Missouri" [USA] [orange star] by Schmidt (1953:101).
Type specimen: USNM 55516, neotype, designated by Stejneger (1938:175), see Cochran (1961) and Reynolds et al. (2007).
Etymology:* Eponym honoring Philo R. Hoy (1816–1892), USA; ornithologist and entomologist who collected the species.

Emys orthonyx Wied-Neuwied 1865:23
Type locality: "südlichen Gewässern bei New-Orleans" [Louisiana, USA] [white star].
Type specimen(s): Not located, syntypes (3), type specimens figured (pl.2,f.5,pl.3,f.1-2).
Etymology:* Greek, *orthos*, straight, upright, correct, intact, and *onyx*, claw, nail; referring to the long, straight claws of the front feet.

Pseudemys vioscana Brimley 1928:66
Type locality: "Lake Des Allemands, La." [Louisiana, USA] [brown star].
Type specimen: USNM 79632 (formerly Brimley no. 115), holotype, see Cochran (1961) and Reynolds et al. (2007).
Etymology:* Eponym honoring Percy Viosca, Jr. (1892–1961), USA; biologist and herpetologist at Tulane University, Louisiana, who collected the types.

Pseudemys elonae Brimley 1928:67
Type locality: "a pond in Guilford County, North Carolina, not far from Elon College, in the Cape Fear drainage" [USA] [purple star].
Type specimen: USNM 79631 (formerly Brimley no. 124), holotype, see Cochran (1961) and Reynolds et al. (2007).
Etymology:* Toponym for Elon College (now Elon University), North Carolina; college originally named for the *elon* (Hebrew for oak tree).

Pseudemys floridana mobiliensis Carr 1937a:1 (*nomen invalidum*)

Emys mobiliensis, Pseudemys mobiliensis, Pseudemys concinna mobiliensis

Etymology:* Toponym for Mobile, Alabama, USA.

Comment: Unjustified emendation or error for *mobilensis*.

Pseudemys concinna metteri Ward 1984:34

Type locality: "Old Fort Cobb, Caddo County, Oklahoma" [USA] [dark gray star].

Type specimen: USNM 7173, holotype, see Reynolds et al. (2007).

Etymology:* Eponym honoring Dean E. Metter (1932–2001), USA; zoologist at the University of Missouri and mentor to the author.

Pseudemys concinna suwanniensis Carr 1937a (07:15, 09:13)


Suwannee Cooter



Dale R. Jackson / CBFTT / CRM 3 / Wakulla River, Wakulla Co., Florida



George L. Heinrich / Rainbow River, Dunnellon, Marion Co., Florida



** (subspecies: *concinna* = red, *suwanniensis* = blue; orange dots = probably introduced or translocated; star = type locality)

Distribution: USA (*Florida, Georgia*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 25,910 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 62,460 sq. km

Size (Max SCL):* male 40.2 cm, female 43.7 cm (Pritchard 1980b; Jackson and Walker 1997; Jackson 2006; Siders et al. 2023; Johnston et al. 2024a)

Synonymy:

Pseudemys floridana suwanniensis Carr 1937a:4

Pseudemys concinna suwanniensis, Chrysemys concinna suwanniensis, Pseudemys suwanniensis

Type locality: "Suwannee River at Manatee Springs, Levy-Dixie County line, Florida" [USA] [yellow star].

Type specimen: UMMZ 81673, holotype, see Peters (1952) and Kluge (1984).

Etymology:* Toponym for the Suwannee River, Florida, USA.

Pseudemys floridana (Le Conte 1830) (07:14, 09:13, 10:9) (49)*


(or *Pseudemys concinna floridana*)

(or *Pseudemys floridana floridana*)


Coastal Plain Cooter, Pond Cooter, Florida Cooter



John Jensen / CRM 3 / Okaloosa Co., Florida



John B. Iverson / *left*: Charlton Co., Georgia / *right*: Marion C., Florida / hatchlings



** (orange dots = questionable, possibly translocated; star = type locality)

Distribution: USA (*Alabama, Georgia, Florida, Mississippi, North Carolina, South Carolina, Virginia*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 299,337 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 471,367 sq. km

Size (Max SCL): male 31.6 cm, female 39.7 cm (Ernst and Lovich 2009; Powell et al. 2016)

IUCN Red List: Least Concern (LC) (van Dijk 2011), as *Pseudemys concinna floridana*. Previously: Least Concern (LC) [Not Listed] (Behler 1996)

Synonymy:

Testudo floridana Le Conte 1830:100

Terrapene floridana, Emys floridana, Clemmys (Clemmys) floridana, Pseudemys floridana, Chrysemys floridana, Pseudemys floridana floridana, Chrysemys floridana floridana, Pseudemys concinna floridana

Type locality: "St. John's river of East Florida" [USA]. Emended to "lower reaches of the St. John's River (Duval County), Florida" [USA] [orange star] by Bour (2003:540).

Type specimens: MNHN 9170, syntype (1), other syntypes apparently lost, discussed by Bour (2003); Uetz et al. (2019) erroneously listed MNHN 91170.

Etymology:* Toponym for Florida, USA.

Comment: The type locality appears to be outside the currently defined native range (see Annotation 47).


Pseudemys gorzugi Ward 1984 (07:16, 12:16, 14:19) (50)*
Rio Grande Cooter, *Jicotea del Rio Bravo*



Charles W. Painter / CBFTT / Black River, Eddy Co., New Mexico



left: Brett Stearns / CBFTT / Devil's River, Val Verde Co., Texas / female
right: Brett Stearns / CBFTT / Black River, Eddy Co., New Mexico



** (light red = extirpated; orange dots = misidentified or translocated; star = type locality)

Distribution: Mexico (*Chihuahua, Coahuila, Nuevo León, Tamaulipas*), USA (*New Mexico, Texas*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 79,877 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 262,558 sq. km

Size (Max SCL): male 28.4 cm, female 40.0 cm (Legler and Vogt 2013; Pierce et al. 2016 CBFTT)

CBFTT Account: Pierce, Stuart, Ward, and Painter (2016)

IUCN Red List:* Vulnerable (VU A2bce) (Mali and Forstner, in press). Previously: Near Threatened (NT) (van Dijk 2011); Near Threatened (NT) (TFTSG *in* Baillie and Groombridge 1996)

USA ESA:* Assessed but Not Listed; Species Status Assessment: USFWS (2021b)


Synonymy:

Pseudemys concinna gorzugi Ward 1984:29
Pseudemys gorzugi
Type locality: "3 1/2 mi. W Jimenez, Río San Diego, Coahuila, México, 850 feet altitude." Emended to "Rio San Diego, 5.6 km W of Jimenez (29º15'N, 100º51'W), Coahuila, Mexico, 260 m elevation" [yellow star] by Legler and Vogt (2013:242).
Type specimen: KU 39986, holotype.
Etymology:* Eponym honoring George R. Zug (1938–), USA; herpetologist and curator at the U.S. National Museum, Washington D.C.


Pseudemys nelsoni Carr 1938a (09:13)
Florida Red-bellied Cooter



Gary Luciano / CBFTT / Big Cypress National Preserve, Collier Co., Florida



Pierson Hill / CBFTT / *left*: Marion Co., *right*: Lake Co., Florida



** (orange dots = questionable, probably translocated; star = type locality)

Distribution: USA (*Florida, Georgia*)

Introduced: British Virgin Islands (*Tortola*), USA (*Texas*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 126,246 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 255,238 sq. km

Size (Max SCL):* male 33.0 cm, female 37.5 cm (Pritchard 1980; Jackson 2010 CBFTT; Siders et al. 2023)

CBFTT Account: Jackson (2010)

IUCN Red List: Least Concern (LC) (van Dijk 2011). Previously: Least Concern (LC) [Not Listed] (Behler 1996)

TFTSG Provisional Red List:* Least Concern (LC) (Munscher, Butterfield, Hauge, Morrison, and Walde, in prep., 2025)

Synonymy:

Pseudemys nelsoni Carr 1938a:307
Pseudemys rubriventris nelsoni, Chrysemys (Pseudemys) nelsoni, Chrysemys nelsoni, Chrysemys rubriventris nelsoni
Type locality: "Fellsmere, Indian River County, Florida" [USA] [yellow star].


Type specimen: MCZ 39888, holotype, see Barbour and Loveridge (1946).
Etymology:* Eponym honoring George Nelson (1876–1962), USA; naturalist and preparator at the Museum of Comparative Zoology, Harvard University, who collected the type.

Pseudemys peninsularis Carr 1938b (07:17, 09:13, 10:9) (49)*
(**or *Pseudemys floridana peninsularis***)
Peninsula Cooter


Richard D. Bartlett / CRM 3 / Lee Co., Florida



left: John B. Iverson / Gainesville, Alachua Co., Florida
right: John B. Iverson / Lake George, Marion Co., Florida / hatchling


**

(orange dot = probably introduced; star = type locality)


Distribution: USA (*Florida*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 98,439 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 114,858 sq. km
Size (Max SCL):* male 35.3 cm, female 38.6 cm (Thomas and Jansen 2006; Siders et al. 2023; Hawthorne and Johnston 2024)
IUCN Red List: Least Concern (LC) (van Dijk 2011), as *Pseudemys peninsularis*
TFTSG Provisional Red List:* Least Concern (LC) (Munscher, Butterfield, Hauge, Morrison, and Walde, in prep., 2025)
Synonymy:
Pseudemys floridana peninsularis Carr 1938b:105
Chrysemys floridana peninsularis, Pseudemys peninsularis
Type locality: "Crystal Springs, Pasco County, Florida" [USA] [yellow star].
Type specimen: MCZ 43849, holotype, see Barbour and Loveridge (1946); erroneously listed as MCZ 42849 by Reynolds et al. (2007).
Etymology:* Latin, *pensilis*, hanging down, *insula*, an island, and *-aris*, pertaining to; referring to the distribution of the species on the Florida peninsula.

Pseudemys rubriventris (Le Conte 1830)
Northern Red-bellied Cooter


James H. Harding / Virginia



C. Kenneth Dodd, Jr. / South Anna River, Hanover Co., Virginia / juvenile


**

(orange dots = probably translocated; stars = type localities)

Distribution: USA (*Delaware, Maryland, Massachusetts, New Jersey, North Carolina, Pennsylvania, Virginia, West Virginia*)
Introduced: South Korea
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 111,065 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 283,425 sq. km
Size (Max SCL):* male 31.2 cm, female 40.0 cm (Graham 1991; Ernst and Lovich 2009)
IUCN Red List: Near Threatened (NT) (van Dijk 2011). Previously: Near Threatened (NT) (TFTSG *in* Baillie and Groombridge 1996)
USA ESA:* Endangered, Massachusetts population only, originally as *Pseudemys rubriventris bangsi* (USFWS 1980); Species Status Assessment: USFWS (2021a)
Synonymy:

Testudo rubriventris Le Conte 1830:101

Terrapene rubriventris, Emys rubriventris, Clemmys (Clemmys) rubriventris, Chrysemys rubriventris, Pseudemys rubriventris, Pseudemys rubiventris, Pseudemys rubriventris rubriventris, Chrysemys rubriventris rubriventris

Type locality: "rivers from New-Jersey to Virginia, chiefly...in such as are rocky; in the Delaware, near Trenton" [USA]. Restricted to "in the Delaware, near Trenton" [New Jersey, USA] [yellow star] by Baur (1893a:224).

Type specimen: MNHN 9208 (with body parts 2069, 2070–71 of the same specimen), lectotype, designated by Bour (2003:541).

Etymology:* Latin, *rubra*, red, and *ventralis*, ventral surface or stomach; referring to the red plastron.

Emys irrigata Bell *in* Duméril and Bibron 1835:276

Emys irrigita

Type locality: "la partie septentrionale du Nouveau-Monde." Restricted to "vicinity of Trenton, New Jersey" [USA] [yellow star] by Schmidt (1953:103).

Type specimens: MNHN 9204, 9206, OUM 8441–42, syntypes (4), discussed by Nowak-Kemp and Fritz (2010).

Etymology:* Latin, *irrigata*, irrigated; referring to the aquatic habitat.

Emys rivulata Gray 1844:22 (junior homonym, not = *Emys rivulata* Valenciennes *in* Bory de Saint-Vincent 1833 [= *Mauremys rivulata*])

Type locality: "N. America." Restricted to "vicinity of Trenton, New Jersey" [USA] [yellow star] by Schmidt (1953:103).

Type specimen: NHMUK 1947.3.5.25, holotype.

Etymology:* Latin, *rivulus*, groove, *rivulata*, having grooves; referring to the rough carapace surface.

Pseudemys rubriventris bangsi Babcock 1937:293

Chrysemys rubriventris bangsi, Pseudemys bangsi

Type locality: "Boot Pond, Plymouth, Massachusetts" [USA]. Corrected to "Gunner's Exchange Pond, Plymouth Co., Massachusetts" [USA] [green star] by Barbour and Loveridge (1946:179).

Type specimen: MCZ 16778, holotype, see Barbour and Loveridge (1946).

Etymology:* Eponym honoring Outram Bangs (1863–1932), USA; mammalogist at the Museum of Comparative Zoology, Harvard University, who discovered this disjunct population.

Pseudemys texana Baur 1893a (12:16)


Texas Cooter



Peter V. Lindeman / South Llano R., Kimble Co., Texas



left: Peter V. Lindeman / South Llano R., Kimble Co., Texas
right: Peter V. Lindeman / Frio R., Real Co., Texas



** (orange dots = probably translocated or misidentified; star = type locality)

Distribution: USA (*Texas*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 102,815 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 203,009 sq. km

Size (Max SCL):* male 29.4 cm, female 34.4 cm (Killebrew and Porter 1989b; Munscher and Franklin 2019; Siders et al. 2023)

IUCN Red List: Least Concern (LC) (van Dijk 2011). Previously: Least Concern (LC) [Not Listed] (Behler 1996)

TFTSG Provisional Red List:* Least Concern (LC) (Gordon, DeChellis, Hammerbach, Ricardez, Franklin, Munscher, and Meik, in prep., 2025)

Synonymy:

Pseudemys texana Baur 1893a:223

Chrysemys texana, Pseudemys floridana texana, Pseudemys concinna texana, Chrysemys concinna texana

Type locality: "San Antonio, Texas" [USA] [yellow star].

Type specimen: ANSP 246, holotype, see Malnate (1971) and Lieb (2023).

Etymology:* Toponym for Texas, USA.

Trachemys Agassiz 1857a (07:18, 09:14, 11:6, 14:21, 14:22, 14:23, 17:26) (51)*

Slider Turtles, Sliders

(includes 17 species, 29 taxa)

Trachemys Agassiz 1857a:252,434

Type species: Originally *Trachemys scabra* [= *Testudo scabra* Linnaeus 1758], by subsequent designation by Brown (1908:114), but its uncertain identity led Lindholm (1929:280) to designate *Trachemys troosti* [= *Emys troosti* Holbrook 1836] as the type species (see also Smith and Smith 1980:434). Rhodin and Carr (2009:14) demonstrated that *Testudo scabra* Linnaeus 1758 was synonymous with *Rhinoclemmys punctularia* [= *Testudo punctularia* Daudin 1801] and declared *T. scabra* a *nomen oblitum*, validating Lindholm's designation of *Trachemys troosti* as type species of *Trachemys*; they also identified the lost holotype of *Trachemys scripta* (*Testudo scripta* Thunberg *in* Schoepff 1792). As a result, Seidel and Ernst (2017:78) suggested that *Trachemys scripta* should be considered the type species for *Trachemys*, but *Trachemys troosti* retains that designation.

Etymology:* Greek, *trachys*, rough, stiff, jagged, prickly, stony, and *emys*, turtle; referring to the rough surfaces on the carapacial scutes.

Comment:* Frétey (2024:74) noted that the designation by Brown (1908) of *Testudo scabra* (sensu Agassiz 1857a) (= *Testudo scabra* Linnaeus 1758) as the type species of *Trachemys* technically may invalidate the declaration by Rhodin and Carr (2009) that the name *Testudo scabra* Linnaeus 1758 is a *nomen oblitum*, since *Testudo scabra* does not meet the requirement of ICZN Article 23.9.1.1 that it has not been used as a valid name after 1899 (being used as such in 1908). Therefore, Frétey (2024) suggested that a petition to the ICZN would need to be submitted to request the use of their plenary powers to suppress the name; otherwise, in his opinion, the name *Callichelys*

with type species *C. ornata* might need to replace the widely used name *Trachemys*. However, the subsequent designation by Lindholm (1929) of *T. troosti* as type species for *Trachemys* seems to have already solved that potential problem.

Callichelys Gray 1863c:179,181
Type species: *Callichelys ornata* [= *Emys ornata* Gray *in* Griffith and Pidgeon 1830], by original designation.
Etymology:* Greek, *kallos*, beauty, and *chelys*, turtle; referring to the fact that the attractive markings on the shell and extremities.


Redamia Gray 1870c:35
Type species: *Redamia olivacea* [= *Emys olivacea* Gray 1856b = subjective synonym of *Pseudemys stejnegeri* Schmidt 1928], by original monotypy.
Etymology:* Anonym, *redamia*, possibly a fabricated name, possibly Vernacular (Old English), or possibly an Eponym honoring someone named Redam.

Trachemys adiutrix Vanzolini 1995 (11:6, 17:27) (51)*
(**or *Trachemys dorbigni adiutrix***)
Maranhao Slider, Carvalho's Slider, *Capininga*, *Cágado*


Richard C. Vogt / Maranhão, Brazil




left: Richard C. Vogt / Maranhão, Brazil
right: Jérôme Maran / Santo Amaro, Maranhão, Brazil


(star = type locality)

Distribution: Brazil (*Maranhão*, *Piauí*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 12,651 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 37,339 sq. km
Size (Max SCL): male 18.6 cm, female 25.1 cm (Batistella 2008; Ernst et al. 2010)
IUCN Red List: Near Threatened (NT B2ab(ii,iii) (Barreto, Sousa Ribeiro, and Bock, in press). Previously: TFTSG Provisional Red List: Near Threatened (NT) (2011); Endangered (EN B1+2c [ver. 2.3]) (TFTSG 1996)

Synonymy:
Trachemys adiutrix Vanzolini 1995:111
Trachemys dorbigni adiutrix, *Trachemys aduitrix*
Type locality: "Brasil: Maranhão: Santo Amaro, 02º33' S, 43º14' W" [Brazil] [yellow star].
Type specimen: MZUSP 3224, holotype.
Etymology:* Latin, *adiutrix*, helper; referring to Maria do Socorro Pinheiro, the author's field assistant and companion. The common name, Carvalho's Slider, honors Antenor Leitão de Carvalho (1910–1985) at Museu Nacional, Rio de Janeiro, who obtained and presented the first known specimen to Vanzolini.

Trachemys callirostris (Gray 1856b) (11:6, 17:26) (51)*
(**or *Trachemys venusta callirostris***)
Colombian Slider, *Jicotea*, *Morrocoy de Agua*
(includes 2 subspecies)


(subspecies: *callirostris* = red, *chichiriviche* = blue; orange dots = trade or translocated)

Distribution:* Colombia (*Antioquia*, *Atlántico*, *Bolívar*, *Cesar*, *Córdoba*, *Cundinamarca*, *La Guajira*, *Magdalena*, *Santander*, *Sucre*), Venezuela (*Carabobo*, *Falcón*, *Trujillo*, *Yaracuy*, *Zulia*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 117,703 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 359,263 sq. km
Size (Max SCL): male 30.0 cm, female 33.0 cm (see subspp.)
CBFTT Account: Bock, Páez, and Daza (2010)
IUCN Red List:* Vulnerable (VU A2bcd+4cde) (Bock, Barros, Guada, Rivas, Hernández, Páez, Rojas-Runjaic, Camargo Siliet, and Cortés-Duque, in press). Previously: TFTSG Provisional Red List: Vulnerable (VU) (2011)
USA ESA:* Endangered, as *Trachemys scripta callirostris* (USFWS 1991b)


Trachemys callirostris callirostris (Gray 1856b) [07:19, 11:6, 17:26]
(or *Trachemys venusta callirostris*)
Colombian Slider



Vivian P. Páez / CBFTT / Colombia



Uwe Fritz / nr. Monteria, Córdoba, Colombia



(subspecies: *callirostris* = red; orange dots = trade or translocated; star = type locality)

Distribution:* Colombia (*Antioquia, Atlántico, Bolívar, Cesar, Córdoba, Cundinamarca, La Guajira, Magdalena, Santander, Sucre*), Venezuela (*Mérida, Trujillo, Zulia*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 109,132 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 255,090 sq. km

Size (Max SCL): male 30.0 cm, female 31.5 cm (Bock et al. 2010 CBFTT; Ceballos et al. 2013)

IUCN Red List:* Vulnerable (VU A2bcd+4cde) (Bock, Barros, Guada, Rivas, Hernández, Páez, Rojas-Runjaic, Camargo Siliet, and Cortés-Duque, in press)

Synonymy:

Emys callirostris Gray 1856b:25

Callichelys callirostris, Pseudemys callirostris, Chrysemys ornata callirostris, Pseudemys scripta callirostris, Pseudemys ornata callirostris, Chrysemys callirostris, Chrysemys scripta callirostris, Trachemys scripta callirostris, Trachemys callirostris, Trachemys ornata callirostris, Trachemys dorbigni callirostris, Trachemys callirostris callirostris, Trachemys venusta callirostris

Type locality:* "America." Restricted to "Unterlauf des Rio Magdalena" [Lower Rio Magdalena, Colombia] [yellow star, location estimated] by Müller (1940:109).


Type specimen: NHMUK 1947.3.4.87 (formerly 1854.10.16.58), holotype.

Etymology:* Greek, *kallos*, beauty, and Latin, *rostrum*, snout, beak; referring to the attractive orange-colored snout region.


Trachemys callirostris chichiriviche (Pritchard and Trebbau 1984) [07:19, 11:6, 17:26]
(or *Trachemys venusta chichiriviche*)
Venezuelan Slider



Hedelvy J. Guada / Jatira-Tacarigua, Falcón, Venezuela



left: Carl H. Ernst / coastal Venezuela
right: Peter C.H. Pritchard / coastal Venezuela



(subspecies: *callirostris* = red, *chichiriviche* = blue; star = type locality)

Distribution: Venezuela (*Carabobo, Falcón, Yaracuy*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 8,571 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 11,467 sq. km

Size (Max SCL): male 24.5 cm, female 33.0 cm (Pritchard and Trebbau 1984; Bock et al. 2010 CBFTT)

IUCN Red List:* Endangered (EN A2cd+4cd) (Bock, Barros, Guada, Rivas, Hernández, Páez, Rojas-Runjaic, Camargo Siliet, and Cortés-Duque, in press)

Synonymy:

Pseudemys scripta chichiriviche Pritchard and Trebbau 1984:191

Trachemys scripta chichiriviche, Trachemys ornata chichiriviche, Trachemys callirostris chichiriviche, Trachemys venusta chichiriviche

Type locality:* "Lago de Tacarigua, Edo. Falcón, Venezuela (68°15' W, 11°4' N)" [= Lago de Tacarigua, Río Tocuyo, Dto. Acosta, Edo. Falcón] [yellow star]. GPS coordinates inaccurate; corrected here to 68°25' W, 11°4' N.


Type specimen: UF 53333, holotype.

Etymology:* Toponym for Cerro de Chichiriviche, a coastal mountain in Falcón, Venezuela.

Trachemys decorata (Barbour and Carr 1940) (14:21)
Hispaniolan Slider



Pablo Feliz / Cabral, Barahona, Dominican Republic / male





left: Harald Artner / Dominican Republic / captivity
right: Pablo Feliz / Algodón, Barahona, Dominican Republic / female



(orange dots = probably introduced; star = type locality)

Distribution: Dominican Republic, Haiti

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 7,110 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 32,841 sq. km

Size (Max SCL): male 21.9 cm, female 34.1 cm (Seidel and Inchaustegui Miranda 1984; Seidel 1988; Ceballos et al. 2013)

IUCN Red List: Vulnerable (VU B1+2c [ver. 2.3]) (TFTSG 1996)

Synonymy:

Pseudemys decorata Barbour and Carr 1940:409


Pseudemys terrapen decorata, Chrysemys (Trachemys) decorata, Chrysemys terrapen decorata, Trachemys decorata, Trachemys stejnegeri decorata

Type locality: "Fond Parisien, Haiti" [yellow star].

Type specimen: MCZ 36862, holotype, see Barbour and Loveridge (1946) and Seidel (1988).

Etymology:* Latin, *decoratus*, adorned, elegant, beautiful; referring to the attractive plastral pattern.

Trachemys decussata (Bell *in* Griffith and Pidgeon 1830) (14:22)
Cuban Slider, *Jamao, Jicotea*
(includes 2 subspecies)



**

(subspecies: *decussata* = red, *angusta* = blue; overlap = intergrades; orange dots = probably introduced or translocated)

Distribution: Cayman Islands [*historical introduction?*], Cuba, Jamaica [*prehistorical introduction?*]

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 112,973 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 370,484 sq. km

Size (Max SCL):* male 27.3 cm, female 45.0 cm (see subspp.)

IUCN Red List: Least Concern (LC) [Not Listed] (Behler 1996)

TFTSG Provisional Red List:* Near Threatened (NT) (Sampedro Marín 2012; Díaz, Fong, Arias, González, Sampedro, and Hurtado, in prep., 2025)

Trachemys decussata decussata (Bell *in* Griffith and Pidgeon 1830) (08:17, 14:22)

Eastern Cuban Slider




Vincenzo Ferri / Cuba / captivity / female





Harald Artner / Cuba / captivity / male



**

(subspecies: *decussata* = red, *angusta* = blue; overlap = intergrades; orange dots = probably introduced or translocated; stars = type localities)

Distribution: Cuba, Jamaica [*prehistorical introduction?*]
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 98,568 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 267,480 sq. km
Size (Max SCL):* male 26.8 cm, female 36.0 cm (Seidel 1988; Thieme 2005)
Synonymy:

Testudo rugosa Shaw 1802:28 (*partim*, junior homonym and *nomen dubium*, not = *Testudo rugosa* Van-Ernest *in* Daudin 1801 [= *Chelonia mydas*])
Emys rugosa, Trachemys rugosa, Clemmys rugosa, Pseudemys rugosa, Pseudemys rugosa rugosa, Pseudemys terrapen rugosa, Chrysemys terrapen rugosa, Trachemys terrapen rugosa
Type locality: Not designated. Restricted to "Rio Jobabo drainage in eastern Cuba" [orange star] by Mittleman (1947:176).
Type specimen: RCSM 990 (formerly LM s/n), holotype, figured in Shaw (1802,pl.4) and Barbour and Carr (1940:pl.1).
Etymology:* Latin, *ruga*, wrinkle, ridge, rugose; referring to the ridged, rugose shell.
Comment:* Name synonymized *in partim* under *Trachemys decussata decussata* and *Trachemys terrapen*.

Emys decussata Bell *in* Griffith and Pidgeon 1830:76 (08:17)
Ptychemys decussata, Clemmys decussata, Pseudemys decussata, Pseudemys decussata decussata, Pseudemys rugosa decussata, Pseudemys terrapen decussata, Chrysemys (Trachemys) decussata, Chrysemys decussata decussata, Chrysemys terrapen decussata, Trachemys decussata, Trachemys decussata decussata
Type locality:* Not designated. Restricted to "North America" by Gray (1830e:11); to "America Boreali" by Gray (1831d:28); to "W. Indies" by Boulenger (1889:80); to "Cuba, exclusive of the drainage systems of the Rio Jobabo and the Caribbean slope of Pinar del Rio Province" by Mittleman (1947:176); to "very likely from Camaguey or Oriente" by Barbour and Carr (1940:402); and hereby to "Camaguey Province, Cuba" [yellow star, location estimated].
Type specimens: OUM 8452–58, NHMUK 1947.3.4.79, syntypes (8), discussed by Nowak-Kemp and Fritz (2010); NHMUK 1947.3.4.79 listed as "holotype" by Seidel (1988), King and Burke (1989), Iverson (1992), and Uetz et al. (2019).
Etymology:* Latin, *decussata*, crossed; referring to the dark markings along the plastron seams.

Emys vermiculata Gray 1844:25
Emys vermicolata
Type locality: "Tropical America." Restricted to "W. Indies" by Gray (1873j:46).
Type specimen: NHMUK s/n, holotype, apparently lost (Seidel 1988; Bour, pers. comm.).
Etymology:* Latin, *vermiculus*, small worm, vermiculated; referring to worm-shaped markings on the plastral marginal scutes.

Emys jamao Duméril 1861b:435 (*nomen nudum*)
Etymology:* Vernacular (Native Cuban), *jamao*, turtle; local indigenous name for the species.

Emys jamao Vilaró 1867a:121
Type locality: Not designated. [Cuba] [no star].
Type specimens: None designated.
Etymology:* Vernacular (Native Cuban), *jamao*, turtle; local indigenous name for the species.


Emys gnatho Vilaró 1867b:204
Type locality: Not designated. [Cuba] [no star].
Type specimens: None designated.
Etymology:* Greek, *gnathos*, jaw; referring to the apparently strongly domed upper jaw.

Pseudemys decussata plana Barbour and Carr 1940:405
Pseudemys terrapen plana, Chrysemys terrapen plana, Trachemys decussata plana
Type locality: "Rio Jobabo, Western Oriente, Cuba" [green star].
Type specimen: MCZ 34134, holotype, see Barbour and Loveridge (1946).
Etymology:* Latin, *plana*, flat, even; referring to the flatter carapace compared to other similar species.

Trachemys decussata angusta (Barbour and Carr 1940) (14:22)
Western Cuban Slider, Cayman Slider


James H. Harding / Grand Cayman Island, Cayman Islands / male


Harald Artner / No data / captivity / female


**
(subspecies: *decussata* = red, *angusta* = blue; overlap = intergrades; orange dots and star = historically introduced *angusta*; stars = type localities)

Distribution: Cayman Islands [*historical introduction?*], Cuba
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 21,973 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 126,068 sq. km
Size (Max SCL):* male 27.3 cm, female 45.0 cm (Barbour and Carr 1940, 1941; L.M. Díaz, unpubl. data)
Synonymy:

Pseudemys decussata angusta Barbour and Carr 1940:402
Pseudemys rugosa angusta, Pseudemys terrapen angusta, Chrysemys terrapen angusta, Trachemys decussata angusta, Trachemys decorata angusta
Type locality: "Taco River, Pinar del Rio, Cuba" [yellow star].
Type specimen: MCZ 34340, holotype, see Barbour and Loveridge (1946).
Etymology:* Latin, *angustus*, narrow; referring to the carapace which is narrower than in the nominate subspecies.

Pseudemys granti Barbour and Carr 1941:59
Pseudemys terrapen granti, Pseudemys decussata granti, Pseudemys stejnegeri granti, Chrysemys terrapen granti, Chrysemys decussata granti, Trachemys decussata granti,

Trachemys granti, Trachemys stejnegeri granti
Type locality: "Grand Cayman" [Cayman Islands] [orange star; possible historical introduction].
Type specimen: MCZ 46045, holotype, see Barbour and Loveridge (1946).
Etymology:* Eponym honoring Major Chapman Grant (1887–1983), USA; herpetologist and grandson of U.S. President Ulysses S. Grant, who collected the species.

Trachemys dorbigni (Duméril and Bibron 1835) (07:20, 11:6, 17:27) (51)*
(**or *Trachemys dorbigni dorbigni***)
D'Orbigny's Slider, Black-bellied Slider, *Tigre d'Água, Tortuga Morrocoyo*


Jérôme Maran / Brazil


Leandro Alcalde / Córdoba Prov., Argentina



** (orange dots = introduced or translocated; stars = type localities)


Distribution:* Argentina (*Buenos Aires, Chaco, Corrientes, Entre Rios, Santa Fe*), Brazil (*Rio Grande do Sul*), Uruguay
Introduced:* Argentina (*Córdoba*), Brazil (*Alagoas, Bahia, Espirito Santo, Goiás, Minas Gerais, Paraná, Rio de Janeiro, Santa Catarina, São Paulo, Sergipe, Tocantins*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 541,619 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 689,976 sq. km
Size (Max SCL):* male 23.0 cm, female 25.5 cm (Seidel 1989; Fagundes et al. 2010)
IUCN Red List:* Near Threatened (NT A2cde+4cde) (Fagundes, Alcalde, Carreira, Fallabrino, Estrades, Florencia David, Famelli, and Lustosa, in press). Previously: TFTSG Provisional Red List: Least Concern (LC) (2011), Least Concern (LC) [Not Listed] (Behler 1996)

Synonymy:
Emys dorbigni Duméril and Bibron 1835:272
Clemmys dorbigni, Pseudemys dorbigni, Chrysemys (Trachemys) dorbigni, Pseudemys scripta dorbigni, Pseudemys dorbigni dorbigni, Chrysemys dorbigni dorbigni, Chrysemys scripta dorbigni, Trachemys scripta dorbigni, Trachemys dorbigni, Trachemys dorbigni dorbigni
Type locality: "Buenos-Ayres" [Buenos Aires, Argentina] [yellow star].
Type specimen: MNHN 9221, holotype, see Seidel (1989).
Etymology:* Eponym honoring Alcide d'Orbigny (1802–1857), France; paleontologist and naturalist who collected the species.
Clemmys (Rhinoclemmys) orbignyi Fitzinger 1835:124 (*nomen novum*)
Emys orbignyi
Type locality: "Buenos-Ayres" [Buenos Aires, Argentina], by default.
Etymology:* Eponym honoring Alcide d'Orbigny (1802–1857), France; paleontologist and naturalist who collected the species.
Comment: Unjustified emendation of *dorbigni*.
Clemmys dorbignyi Boulenger 1886b:424 (*nomen novum*)
Chrysemys dorbignyi, Emys dorbignyi, Pseudemys dorbignyi, Pseudemys dorbignyi dorbignyi, Pseudemys scripta dorbignyi, Chrysemys scripta dorbignyi, Trachemys scripta dorbignyi, Trachemys dorbignyi, Trachemys dorbignyi dorbignyi
Type locality: "Buenos-Ayres" [Buenos Aires, Argentina], by default.
Etymology:* Eponym honoring Alcide d'Orbigny (1802–1857), France; paleontologist and naturalist who collected the species.
Comment: Unjustified emendation of *dorbigni*.
Pseudemys dorbignyi brasiliensis Freiberg 1969:301 (07:20)
Pseudemys dorbigni brasiliensis, Pseudemys scripta brasiliensis, Chrysemys dorbigni brasiliensis, Chrysemys scripta brasiliensis, Trachemys scripta brasiliensis, Trachemys dorbigni brasiliensis, Trachemys dorbignyi brasiliensis
Type locality: "río Guaíba, Porto Alegre, Brasil" [green star].
Type specimen: MACN 23628, holotype.
Etymology:* Toponym for Brazil.

Trachemys gaigeae (Hartweg 1939) (07:18, 21:56, 21:58) (51)*
Big Bend Slider


James N. Stuart / CBFTT / Socorro Co., New Mexico


James N. Stuart / CBFTT / Socorro Co., New Mexico


(star = type locality)

Distribution: Mexico (*Chihuahua, Coahuila*), USA (*New Mexico, Texas*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 39,807 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 195,714 sq. km

Size (Max SCL): male 22.2 cm, female 27.0 cm (Stuart and Ward 2009 CBFTT)

CBFTT Account: Stuart and Ward (2009)

IUCN Red List: Vulnerable (VU A2ce+4ce) (van Dijk 2011). Previously: Vulnerable (VU) (TFTSG *in* Baillie and Groombridge 1996)

TFTSG Provisional Red List:* Vulnerable (VU) (Gordon, DeChellis, Hammerbach, Ricardez, Franklin, in prep., 2025)

Synonymy:

Pseudemys scripta gaigeae Hartweg 1939:1

Pseudemys gaigeae, Chrysemys scripta gaigeae, Chrysemys gaigeae, Trachemys nebulosa gaigeae, Trachemys scripta gaigeae, Trachemys gaigeae, Trachemys ornata gaigeae, Trachemys gaigeae gaigeae

Type locality: "Boquillas, Rio Grande River, Brewster County, Texas" [USA] [yellow star] [now = Rio Grande Village, Big Bend National Park, Texas, USA] (Lieb 2023:250).

Type specimen: UMMZ 66472, holotype, see Peters (1952), Kluge (1984), and Lieb (2023).

Etymology:* Eponym honoring Helen T. Gaige (1890–1976), USA; female herpetologist and curator of the Museum of Zoology at the University of Michigan, who collected the type.

Trachemys grayi (Bocourt 1868) (07:18, 10:10, 11:6, 17:26, 17:28) (52)*
Western Central American Slider, *Jicotea Negra*
(includes 3 subspecies)


**

(subspecies: *grayi* = red, *emolli* = blue, *panamensis* = green)

Distribution: Costa Rica, El Salvador, Guatemala, Honduras, Mexico (*Chiapas, Oaxaca*), Nicaragua, Panama

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 162,881 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 778,168 sq. km

Size (Max SCL): male 29.6 cm, female 54.8 cm (see subspp.)

IUCN Red List:* Least Concern (LC) (Machkour-M'Rabet, Gallardo Alvarez, Lesher-Gordillo, and Bock, in press). Previously: TFTSG Provisional Red List: Data Deficient (DD) (2018)

Trachemys grayi grayi (Bocourt 1868) (07:18, 10:10, 11:6, 17:26, 17:28) (52)*
Gray's Slider, Tehuantepec Slider, *Jicotea de Tehuantepec*


Eduardo Reyes-Grajales / Puerto Arista, Chiapas, Mexico


left: Eduardo Reyes-Grajales / Puerto Arista, Chiapas, Mexico
right: Haitao Shi / Mexico / captivity


•


(subspecies: *grayi* = red, *emolli* = blue; stars = type localities)

Distribution: El Salvador, Guatemala, Mexico (*Chiapas, Oaxaca*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 46,124 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 131,588 sq. km
Size (Max SCL):* male 28.5 cm, female 52.7 cm (Legler 1990; Legler and Vogt 2013; E. Reyes-Grajales, unpubl. data)
IUCN Red List:* Least Concern (LC) (Machkour-M'Rabet, Gallardo Alvarez, Lesher-Gordillo, and Bock, in press).
Synonymy:
Emys grayi Bocourt 1868:121 (senior homonym, not = *Emys grayi* Günther 1869 [= *Mauremys caspica*])
Callichelys grayi, Chrysemys grayi, Pseudemys grayi, Pseudemys ornata grayi, Pseudemys scripta grayi, Chrysemys scripta grayi, Trachemys scripta grayi, Trachemys grayi, Trachemys ornata grayi, Trachemys venusta grayi; Trachemys grayi grayi
Type locality:* "l'embouchure du Nagualate, dans le Pacifique (Guatémala)" [= the mouth of Río Nahualate, Guatemala] [yellow star].
Type specimen: MNHN 9220 (formerly 67.127), holotype, see Stuart (1963), Smith and Smith (1980), and Iverson (1992).
Etymology:* Eponym honoring John Edward Gray (1800–1875), United Kingdom; herpetologist, turtle taxonomist, and curator at the British Museum of Natural History, London.
Callichelys concinna Gray 1873a:148 (17:28)
Type locality:* "San Mateo, Tehuantepec" [= San Mateo del Mar, Tehuantepec, Oaxaca, Mexico] [green star].
Type specimens: NHMUK 1871.2.7.43–44, syntypes (2).
Etymology:* Latin, *concinnus*, pretty, delicate, elegant; referring to the elegant markings.
Emys umbra Bocourt 1876b:26 (*nomen novum*)
Pseudemys umbra, Clemmys umbra, Chrysemys umbra, Pseudemys scripta umbra
Type locality: "l'embouchure du Nagualate, dans le Pacifique (Guatémala)" [= the mouth of Río Nahualate, Guatemala] [yellow star], by default.
Type specimen: MNHN 9220 (formerly 67.127), holotype, by default.
Etymology:* Latin, *umbra*, shadow, darkness; referring to the dark color of some adults.
Comment: Unjustified replacement name for *grayi*.

Trachemys grayi emolli (Legler 1990) (07:18, 11:6, 14:23, 17:26, 21:55) (52)*
Nicaraguan Slider


Haitao Shi / No data / captivity


Charlotte Ducotterd / No data / captivity


**

(subspecies: *grayi* = red, *emolli* = blue, *panamensis* = green; stars = type localities)


Distribution:* Costa Rica, El Salvador, Honduras, Nicaragua, Panama
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 53,469 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 118,587 sq. km
Size (Max SCL): male 29.6 cm, female 54.8 cm (Ernst 2008)
IUCN Red List:* Least Concern (LC) (Machkour-M'Rabet, Gallardo Alvarez, Lesher-Gordillo, and Bock, in press).
Synonymy:
Emys valida Le Conte 1860:7 (*nomen oblitum*) (21:55)
Clemmys valida, Emys (Ptychemys) valida, Trachemys valida
Type locality: "Honduras." Restricted to "Pacific coastal drainages of Honduras" [orange star, location estimated] by McCord et al. (2010:40).
Type specimen: ANSP 216, holotype, listed by Malnate (1971), discussed by McCranie (2018).
Etymology:* Latin, *valida*, strong, healthy, valid, effective; possibly referring to the larger size of this species as compared to other slider turtles.
Pseudemys scripta emolli Legler 1990:91
Trachemys scripta emolli, Trachemys ornata emolli, Trachemys emolli, Trachemys venusta emolli, Trachemys grayi emolli
Type locality: "Río Tepetate, 2.5 km northeast of Granada, Granada Province, Nicaragua" [yellow star].

Type specimen: UU 6728, holotype.
Etymology:* Eponym honoring Edward O. Moll (1939–), USA; herpetologist at Eastern Illinois University and first Chair of the IUCN Freshwater Chelonian Specialist Group, precursor to the IUCN Tortoise and Freshwater Turtle Specialist Group.

Trachemys grayi panamensis McCord, Joseph-Ouni, Hagen, and Blanck 2010 (10:10, 11:6, 17:26) (52)*
Panamanian Slider


© Kai Squires, iNaturalist.org / Anton Valley, Panama


Raúl Fournier / Térraba Sierpe, Costa Rica



**
(subspecies: *emolli* = blue, *panamensis* = green; star = type locality)

Distribution: Costa Rica, Panama
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 63,391 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 160,780 sq. km
Size (Max SCL):* male 23.4 cm, female 32.0 cm (Legler 1990, as "CAPAC" population)
IUCN Red List:* Least Concern (LC) (Machkour-M'Rabet, Gallardo Alvarez, Lesher-Gordillo, and Bock, in press).
Synonymy:
Testudo panama Perry 1810:[unpaginated], pl.33 (*nomen oblitum*) (12:18) (52)*
Type locality: "Panama."
Type specimen: Not located, holotype, originally live, specimen figured (pl.33).
Etymology:* Toponym for Panama.
Comment:* Declared a *nomen oblitum* by Bauer and Petit (2004) and here in our annotation. Previously synonymized under *Trachemys venusta uhrigi* (TTWG 2021).
Trachemys venusta panamensis McCord, Joseph-Ouni, Hagen, and Blanck 2010:46
Trachemys grayi panamensis
Type locality: "Chiva-Chiva Road (trail), 1 km from Gaillané (Gaillard) Highway (Fort Clayton entrance), north of Miraflores Lake, Pacific-side Panama Canal Zone, Panamá Province, Panama" [yellow star].
Type specimen: UF 52511, holotype.
Etymology:* Toponym for Panama.

Trachemys hartwegi (Legler 1990) (07:18, 21:56)
Nazas Slider, Mexican Plateau Slider, *Jicotea del Rio Nazas*


John B. Iverson / CBFTT / Río Nazas, Preza Francisco Zarco, Durango, Mexico


left: John B. Iverson / CBFTT / Río Nazas, Preza Francisco Zarco, Durango, Mexico
right: Peter Paul van Dijk / Río Nazas, Cuidad Lerdo, Durango, Mexico / hatchling


**
(light red = probably extirpated; red dots = probably extirpated; star = type locality)

Distribution:* Mexico (*Coahuila*, *Durango*, *Zacatecas*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 18,397 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 48,397 sq. km
Size (Max SCL): male 14.9 cm, female 30.8 cm (Legler 1990; Stuart and Ward 2009 CBFTT)
CBFTT Account: Stuart and Ward (2009), as a subspecies of *Trachemys gaigeae*.
IUCN Red List:* Vulnerable (VU A2c+4ce) (Villegas Raygoza, Rhodin, and Castañeda Gaytán, in press). Previously: Vulnerable (VU A2ce+4ce) (van Dijk 2011), as subspecies of *Trachemys gaigeae*; Vulnerable (VU) (TFTSG *in* Baillie and Groombridge 1996), as subspecies of *T. gaigeae*
Synonymy:
Pseudemys scripta hartwegi Legler 1990:89
Chrysemys scripta hartwegi, *Trachemys scripta hartwegi*, *Trachemys ornata hartwegi*, *Trachemys nebulosa hartwegi*, *Trachemys gaigeae hartwegi*
Type locality: "Río Nazas, 1.2 km east of Presa Lázaro Cardenas, Durango, Mexico" [yellow star].
Type specimen: UU 3802, holotype.
Etymology:* Eponym honoring Norman E. Hartweg (1904–1964), USA; herpetologist and curator at the Museum of Zoology at the University of Michigan.

Trachemys medemi Vargas-Ramírez, del Valle, Ceballos, and Fritz 2017 (21:57) (51, 53)*

Atrato Slider, *Icotea del Atrato*



Carlos del Valle / Ciénaga de Marriaga, Unguía, Chocó, Colombia



left: Claudia Ceballos / Tulenapa, Carepa, Antioquia, Colombia / juvenile
right: Uwe Fritz / Colombia / adult



**

(orange dot = questionable; star = type locality)

Distribution:* Colombia (*Antioquia, Chocó*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 12,113 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 19,811 sq. km

Size (Max SCL): male 19.8 cm, female 28.1 cm (Ceballos and Brand 2014; Vargas-Ramírez et al. 2017); **(Max CCL):** female 38.5 cm (Ceballos and Brand 2014)

IUCN Red List:* Endangered (EN A4cd) (Vargas-Ramírez, Cuadrado-Ríos, Gaviria-Hernández, and Ceballos, in press). Previously: TFTSG Provisional Red List: Vulnerable (VU) (2017)

Synonymy:

Trachemys medemi Vargas-Ramírez, del Valle, Ceballos, and Fritz 2017:333

Type locality: "Río Sucio, Chocó, Parque Nacional Natural Los Katíos, Colombia" [yellow star].


Type specimen: IAVH R-1606, holotype.

Etymology:* Eponym honoring Federico Medem (Count Friedrich Johann von Medem) (1912–1984), Latvia / Germany / Colombia; nobleman, zoologist, herpetologist, and director of the Instituto Roberto Franco in Villavicencio, Colombia.

Trachemys nebulosa (Van Denburgh 1895) (07:18) (51)*

Baja California Slider

(includes 2 subspecies)



**

(subspecies: *nebulosa* = red, *hiltoni* = blue; orange dots = questionable)

Distribution: Mexico (*Baja California Sur, Sinaloa, Sonora*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 34,618 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 142,770 sq. km


Size (Max SCL): male 33.0 cm, female 37.0 cm (see subspp.)

IUCN Red List:* Vulnerable (VU A2cd+4cd) (Cupul-Magaña, Flores-Guerrero, Nolasco-Luna, and Escobedo-Galván, in press). Previously: TFTSG Provisional Red List: Data Deficient (DD) (2018)


Trachemys nebulosa nebulosa (Van Denburgh 1895) (07:18)
Baja California Slider



James R. Buskirk / Cadeje, Baja California Sur, Mexico



Georg Gassner / La Purísima, Baja California Sur, Mexico



(subspecies: *nebulosa* = red, *hiltoni* = blue; star = type locality)

Distribution: Mexico (*Baja California Sur*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 27,556 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 55,974 sq. km

Size (Max SCL): male 33.0 cm, female 37.0 cm (Seidel 2010)

Synonymy:

Chrysemys nebulosa Van Denburgh 1895:84

Chrysemys ornata nebulosa, Pseudemys ornata nebulosa, Pseudemys nebulosa, Pseudemys scripta nebulosa, Chrysemys scripta nebulosa, Trachemys scripta nebulosa, Trachemys dorbigni nebulosa, Trachemys ornata nebulosa, Trachemys nebulosa, Trachemys nebulosa nebulosa

Type locality: "Mainland abreast of San José Island, Lower California... Los Dolores, L.C." [Baja California Sur, Mexico] [yellow star].

Type specimen: CAS 2244, holotype, see Slevin and Leviton (1956) and Smith and Smith (1980).

Etymology:* Latin, *nebula*, fog, haze, *nebulosa*, smoky gray, foggy, cloudy, nebulous; referring to the more irregular and indistinct ocelli on the carapacial scutes as compared to other similar species.


Trachemys nebulosa hiltoni (Carr 1942) (07:18)
Fuerte Slider, Rio Fuerte Slider, *Jicotea del Fuerte*



Philip C. Rosen / San Miguel Zapotitlán, Rio Fuerte, Sinaloa, Mexico



Philip C. Rosen / San Miguel Zapotitlán, Rio Fuerte, Sinaloa, Mexico



**

(subspecies: *nebulosa* = red, *hiltoni* = blue;
orange dots = questionable; star = type locality)

Distribution: Mexico (*Sinaloa, Sonora*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 7,062 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 11,372 sq. km

Size (Max SCL): male 32.0 cm, female 35.1 cm (Legler 1990; Legler and Vogt 2013)

Synonymy:

Pseudemys scripta hiltoni Carr 1942:1

Pseudemys concinna hiltoni, Chrysemys scripta hiltoni, Chrysemys gaigeae hiltoni, Trachemys scripta hiltoni, Trachemys ornata hiltoni, Trachemys nebulosa hiltoni

Type locality: "Guirocoba about 28 miles southeast of Alamos, Sonora, Mexico, at an elevation of approximately 1,485 feet" [yellow star].

Type specimen: AMNH 63747, holotype.

Etymology:* Eponym honoring John W. Hilton (1904–1983), USA; landscape artist, geologist, and zoologist, who collected the types.

Trachemys ornata (Gray *in* Griffith and Pidgeon 1830) (07:18, 07:19, 10:10, 11:6, 17:26)

Ornate Slider, *Jicotea*



Philip C. Rosen / Río Presídio, Mazatlan, Sinaloa, Mexico





left: Philip C. Rosen / Río Presídio, Mazatlan, Sinaloa, Mexico
right: James R. Buskirk / Presidios, Sinaloa, Mexico



** (orange dots = probably introduced or translocated; star = type locality)

Distribution: Mexico (*Jalisco, Nayarit, Sinaloa*)

Introduced: Mexico (*Guerrero [?], Michoacán [?]*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 19,486 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 48,420 sq. km

Size (Max SCL):* male 35.9 cm, female 43.7 cm (Legler 1990; Legler and Vogt 2013)

IUCN Red List:* Near Threatened (NT A2c+4c) (Cupul-Magaña, Flores-Guerrero, Nolasco-Luna, and Escobedo-Galván, in press). Previously: Vulnerable (VU B1ab(iii)+2ab(iii)) (Frost, Hammerson, and Gadsden 2007)

Synonymy:

Emys ornata Gray *in* Griffith and Pidgeon 1830:76 [Gray 1830c]

Clemmys (Clemmys) ornata, Callichelys ornata, Pseudemys ornata, Chrysemys ornata, Chrysemys ornata ornata, Pseudemys ornata ornata, Pseudemys scripta ornata, Chrysemys scripta ornata, Trachemys scripta ornata, Trachemys ornata, Trachemys ornata ornata

Type locality: Not designated. Restricted to "South America" by Gray (1830e:12); and to "America Meridionali...Mazetland" by Gray (1831d:30). Emended to "Mazatlan" [Mazatlán, Sinaloa, Mexico] [yellow star] by Gray (1856b:24).


Type specimen: NHMUK 1946.1.22.40, lectotype, designated by Legler and Vogt (2013:301).

Etymology:* Latin, *ornatus*, ornate, ornamental; referring to the carapacial color pattern.

Trachemys scripta (Thunberg *in* Schoepff 1792) (09:15, 21:58)

Pond Slider, Common Slider

(includes 3 subspecies)



** (subspecies: *scripta* = red, *elegans* = blue, *troostii* = green; overlap (purple) = intergrades; orange dots = introduced, translocated, and invasive)

Distribution:* Mexico (*Coahuila, Nuevo León, Tamaulipas*), USA (*Alabama, Arkansas, Florida, Georgia, Illinois, Indiana, Iowa, Kansas, Kentucky, Louisiana, Mississippi, Missouri, Nebraska, New Mexico [eastern], North Carolina, Ohio, Oklahoma, South Carolina, Tennessee, Texas, Virginia, West Virginia, Wisconsin*)

Introduced:* Most listings are for *T. s. elegans*: Argentina, Australia, Austria, Bahamas, Bahrain, Bangladesh, Belarus, Belgium, Bermuda, Bhutan, Bosnia and Herzegovina, Bulgaria, Brazil, British Virgin Islands, Cambodia, Canada, Cayman Islands, Chile, China, Colombia, Croatia, Cyprus, Czech Republic, Denmark, Dominican Republic, Ecuador, Egypt, Estonia, Finland, France, French Polynesia, Georgia, Germany, Greece, Guadeloupe, Guam, Guyana, Honduras, Hungary, Indonesia, India, Iran, Ireland, Israel, Italy, Japan, Jordan, Kazakhstan, Kyrgyzstan, Latvia, Lithuania, Luxembourg, Malaysia, Martinique, Mexico (*non-native range*), Micronesia, Moldova, Mongolia, Montenegro, Myanmar, Nepal, Netherlands, Netherlands Antilles, New Zealand, Nicaragua, Northern Mariana Islands, Oman, Pakistan, Palau, Palestine, Panama, Philippines, Poland, Portugal, Puerto Rico, Réunion, Romania, Russia, Saudi Arabia, Serbia, Seychelles, Singapore, Sint Maarten, Slovakia, Slovenia, South Africa, South Korea, Spain, Sri Lanka, Suriname, Sweden, Switzerland, Taiwan, Tajikistan, Thailand, Trinidad, Turkey, Ukraine, United Arab Emirates, United Kingdom, USA (*non-native range*), US Virgin Islands, Uzbekistan, Vietnam

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 2,447,717 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 3,473,334 sq. km

Size (Max SCL): male 26.1 cm, female 32.8 cm (see subspp.)

IUCN Red List: Least Concern (LC) (van Dijk, Harding, and Hammerson 2011). Previously: Near Threatened (NT) (TFTSG *in* Baillie and Groombridge 1996)


Trachemys scripta scripta (Thunberg *in* Schoepff 1792) [(09:15, 21:58)]
Yellow-bellied Slider



Kurt A. Buhlmann / Savannah River Site, Aiken Co., South Carolina



Kurt A. Buhlmann / Savannah River Site, Aiken Co., South Carolina



** (subspecies: *scripta* = red, *elegans* = blue, *troostii* = green; overlap = intergrades; orange dots = introduced, translocated, and invasive; stars = type localities)

Distribution:* USA (*Alabama, Florida, Georgia, Mississippi, North Carolina, South Carolina, Tennessee, Virginia*)

Introduced:* See listing under the species heading above.

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 482,335 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 719,166 sq. km

Size (Max SCL):* male 25.1 cm, female 34.4 cm (Mitchell et al. 2017; Johnston et al. 2024c)

Synonymy:

Testudo scripta Thunberg *in* Schoepff 1792:16 (*nomen protectum*) [(09:15)]

Emys scripta, Trachemys scripta, Chrysemys scripta, Pseudemys scripta, Chrysemys scripta scripta, Chrysemys palustris scripta, Pseudemys scripta scripta, Trachemys scripta scripta

Type locality: Not known. Restricted to "Charleston, South Carolina" [USA] [yellow star] by Schmidt (1953:102).

Type specimen: UPSZTY 7455 (formerly UUZM 7455), holotype, listed by Thunberg (1828), not listed by Lönnberg (1896), Andersson (1900), Holm (1957), or Wallin (2001), rediscovered and designated by Rhodin and Carr (2009:13).

Etymology:* Latin, *scriptus*, written; referring to the markings on the carapace, plastron, and head.

Comment: Name conserved by ICZN (1985b), see Legler et al. (1980).

Testudo serrata Daudin 1801:148 (senior homonym, not = *Testudo serrata* Shaw 1802 [= *Geoemyda spengleri*])

Emys serrata, Terrapene serrata, Clemmys (Clemmys) serrata, Pseudemys serrata

Type locality: "la Caroline" [South Carolina, USA] [green star, location estimated].

Type specimen: Possibly MNHN, not located, type specimen figured (pl.21,f.1-2).

Etymology:* Latin, *serratus*, saw-shaped, serrated; referring to the posterior carapacial margin.

Emys occipitatis Gray *in* Griffith and Pidgeon 1830:75 [Gray 1830c]

Type locality: Not designated.

Type specimen: NHMUK s/n, holotype, apparently lost, redescribed as *Emys vittata* Gray 1830e.

Etymology:* Latin, *occipitatis*, belonging to the occiput, belonging to the back of the head; referring to the conspicuous yellow head stripe behind each eye.

Emys vittata Gray 1830e:11 (*nomen novum*) [(10:7)]

Type locality: "North America?"

Type specimen: NHMUK s/n, holotype, apparently lost, same specimen as holotype of *Emys occipitatis* Gray *in* Griffith and Pidgeon 1830 [Gray 1830c].

Etymology:* Latin, *vitta*, band, ribbon, stripe, *vittata*, striped; referring to the conspicuous yellow head stripe behind each eye.

Comment: Unjustified replacement name for *occipitatis*.


Trachemys scripta elegans (Wied-Neuwied 1839) [17:25, 21:58] [54, 55]*
Red-eared Slider



James C. Godwin / Bon Secour R., Mobile Bay, Alabama



John L. Carr / nr. Monroe, Ouachita Parish, Louisiana



** (subspecies: *scripta* = red, *elegans* = blue, *troostii* = green; overlap = intergrades; orange dots = introduced, translocated, and invasive; stars = type localities)

Distribution: Mexico (*Coahuila, Nuevo León, Tamaulipas*), USA (*Alabama, Arkansas, Florida, Georgia, Illinois, Indiana, Iowa, Kansas, Kentucky, Louisiana, Mississippi, Missouri, Nebraska, New Mexico, Ohio, Oklahoma, Tennessee, Texas, West Virginia, Wisconsin*)

Introduced:* See listing under the species heading above.

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 2,010,329 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 2,695,178 sq. km

Size (Max SCL): male 26.1 cm, female 32.8 cm (Tucker et al. 2006; Brown et al. 2020)

Synonymy:

Testudo chlorops Rafinesque 1818:354 (*nomen nudum*) [21:7]
Type locality: "Kentucky" [USA].
Type specimen: Not located, type specimen figured in manuscript drawing by Rafinesque in 1818, reproduced by Bell and Bauer (2020:f.6).
Etymology:* Greek, *chloros*, green, yellow, pale green, and *ops*, eye, appearance; referring to the greenish shell with yellowish lines.

Emyda semiradiata Rafinesque 1822:3 (*nomen nudum et novum*) [21:7]
Etymology:* Latin, *semi*, half, and *radius*, ray, stick, spoke; referring to the stripes on the carapace extending over half its width.
Comment: Replacement name for *radiata*, a manuscript replacement name for *chlorops* (see below).

Emys concinna Cuvier *in* Guérin 1829:pl.1,f.3 (senior homonym and *nomen oblitum*, not = *Emys concinna* Le Conte 1830 [= *Pseudemys concinna concinna*]) [17:25]
Type locality: Not designated.
Type specimen: Not located, type specimen figured (pl.1,f.3).
Etymology:* Latin, *concinnus*, pretty, delicate, elegant; referring to the elegant markings.

Emys elegans Wied-Neuwied 1839:176(9),213
Trachemys elegans, Clemmys elegans, Pseudemys elegans, Chrysemys elegans, Chrysemys scripta elegans, Chrysemys palustris elegans, Pseudemys troostii elegans, Pseudemys scripta elegans, Trachemys scripta elegans
Type locality: "Gegend von Harmony...in dem Wabasch und Fox-River" [Indiana, USA]. Emended to "in Fox Rivers bei New-Harmony aus einem nebenflusse des Wabasch" [Indiana, USA] [yellow star] by Wied-Neuwied (1865:41).
Type specimen: Not located, type specimen figured in Wied-Neuwied (1865:pl.4).
Etymology:* Latin, *elegans*, elegant, fine, select; referring to the nice markings on the head and limbs.

Emys holbrookii Gray 1844:23
Trachemys holbrookii
Type locality: "N. America. Louisiana" [USA] [green star, location estimated].
Type specimen: NHMUK 1947.3.4.78, holotype.
Etymology:* Eponym honoring John Edwards Holbrook (1794–1871), USA; physician, anatomist, and herpetologist at the Medical College of South Carolina.

Emys sanguinolenta Gray 1856b:26,pl.15,f.1
Type locality: Not designated. Restricted to "N. America" by Gray (1873j:44), and to "Charleston, South Carolina" [USA] [in error] [orange star] by Schmidt (1953:102).
Type specimen: NHMUK 1946.1.22.35 (formerly 1851.3.14.1), holotype.
Etymology:* Latin, *sanguis*, blood, and *-olentus*, full of, *sanguinolentus*, bloody, covered in blood; referring to the conspicuous red stripe behind each eye.

Trachemys lineata Gray 1873a:147
Type locality: "North America." Restricted to "New Harmony, Posey County, Indiana" [USA] [yellow star] by Schmidt (1953:103).
Type specimen: NHMUK 1946.1.22.78 (formerly 1866.6.19.1), holotype.
Etymology:* Latin, *linea*, line, stripe, *lineata*, having lines, striped; referring to the carapacial line markings.


Emys bicolor Berlandier *in* Bour 2017:55 (*nomen nudum*)
Type locality:* Not designated. Restricted to "Tamaulipas, Mexico" [blue star, location estimated] by TTWG (2021:164).
Type specimen: Not located, type specimen figured in manuscript drawing by Berlandier in 1850 (pl.4), reproduced by Bour (2017:f.2).
Etymology:* Latin, *bi*, twice, and *color*, color; referring to the predominantly greenish and yellow coloring of the shell.

Testudo radiata Rafinesque *in* Bell and Bauer 2020:23 (*nomen nudum et novum*) [21:7]
Type locality: "Kentucky" [USA].
Type specimen: Not located, type specimen figured in manuscript drawing by Rafinesque in 1818, reproduced by Bell and Bauer (2020:f.6).
Etymology:* Latin, *radius*, ray, stick, spoke; referring to the stripes on the carapace.
Comment: Manuscript replacement name for *chlorops*.


Trachemys scripta troostii (Holbrook 1836) [21:58]
Cumberland Slider



James H. Harding / Tennessee



James H. Harding / Tennessee / hatchling, adult



**

(subspecies: *scripta* = red, *elegans* = blue, *troostii* = green; overlap = intergrades; orange dots = introduced, translocated, and invasive; star = type locality)

Distribution:* USA (*Alabama, Georgia, Kentucky, North Carolina, Tennessee, Virginia*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 51,992 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 110,580 sq. km

Size (Max SCL): female 21.0 cm (Vetter 2004)

Synonymy:

Emys troostii Holbrook 1836:55

Trachemys troostii, Clemmys troostii, Pseudemys troostii, Chrysemys troostii, Pseudemys scripta troostii, Pseudemys troostii troostii, Chrysemys scripta troostii, Trachemys scripta troostii

Type locality: "Cumberland river" [Tennessee, USA] [yellow star, location estimated].

Type specimen: ANSP 179, holotype, see Malnate (1971); not ANSP 180 as noted by Carr (1937b).

Etymology:* Eponym honoring Gerard Troost (1776–1850), Netherlands / USA; geologist and naturalist and first president of the Academy of Natural Sciences, Philadelphia, who collected the types.


Emys cumberlandensis Holbrook 1840:55

Type locality: "Tennessee...Cumberland river" [USA] [yellow star, location estimated].

Type specimen: Not located, originally live, holotype specimen figured (pl.8).

Etymology:* Toponym for Cumberland River, Tennessee, USA.

Trachemys stejnegeri (Schmidt 1928) [14:22]
Central Antillean Slider
(includes 3 subspecies)



(subspecies: *stejnegeri* = red, *malonei* = purple, *vicina* = blue; orange dots = possibly introduced)

Distribution: Bahamas (*Inagua*), Dominican Republic, Haiti, Puerto Rico

Introduced: Dominica, Guadeloupe

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 38,908 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 123,390 sq. km

Size (Max SCL): male 21.0 cm, female 27.3 cm (see subspp.)

IUCN Red List: Near Threatened (NT) (TFTSG 1996)

TFTSG Provisional Red List: Near Threatened (NT) (2011)


Trachemys stejnegeri stejnegeri (Schmidt 1928)
Puerto Rican Slider



Anders G.J. Rhodin / Fajardo, Puerto Rico



left: Michael E. Seidel / Puerto Rico
right: Anders G.J. Rhodin / Fajardo, Puerto Rico





(subspecies: *stejnegeri* = red; star = type locality)

Distribution: Puerto Rico
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 7,816 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 12,561 sq. km
Size (Max SCL): male 21.0 cm, female 26.7 cm (Seidel 1988; Itescu et al. 2014)
Synonymy:
Emys olivacea Gray 1856b:30 (junior homonym, not = *Emys olivacea* Schweigger 1812 [= *Pelomedusa olivacea*])
Clemmys olivacea, Redamia olivacea, Chrysemys olivacea
Type locality: "N. America?"
Type specimen: NHMUK 1947.3.4.10 (formerly 1843.12.25.17), holotype.
Etymology:* Latin, *oliva*, olive, and *-aceus*, belonging to, having the nature of; referring to the olive-colored carapace.
Pseudemys stejnegeri Schmidt 1928:147
Pseudemys palustris stejnegeri, Pseudemys stejnegeri stejnegeri, Pseudemys terrapen stejnegeri, Pseudemys decussata stejnegeri, Chrysemys decussata stejnegeri, Chrysemys stejnegeri, Chrysemys terrapen stejnegeri, Trachemys stejnegeri, Trachemys stejnegeri stejnegeri
Type locality: "San Juan, Porto Rico" [Puerto Rico, USA] [yellow star].
Type specimen: USNM 25642, holotype, see Cochran (1961), Seidel (1988), and Reynolds et al. (2007).
Etymology:* Eponym honoring Leonhard H. Stejneger (1851–1943), Norway / USA; herpetologist, ornithologist, and curator at the U.S. National Museum, Washington, D.C.

Trachemys stejnegeri malonei (Barbour and Carr 1938)
Inagua Slider


James F. Parham / Inagua, Bahamas / male



James F. Parham / Inagua, Bahamas / *left*: male; *right*: female



(subspecies: *malonei* = purple, *vicina* = blue; orange dot = possibly introduced; star = type locality)

Distribution: Bahamas (*Inagua*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 726 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 1,055 sq. km
Size (Max SCL): male 15.4 cm, female 23.8 cm (Barbour and Carr 1938; J. Rosado, unpubl. data)
USA ESA:* Endangered (USFWS 1991b)
Synonymy:
Pseudemys malonei Barbour and Carr 1938:76
Pseudemys palustris malonei, Pseudemys terrapen malonei, Chrysemys malonei, Chrysemys terrapen malonei, Trachemys stejnegeri malonei, Trachemys malonei
Type locality: "ponds near Northwest Point, Great Inagua Island, B.W.I." (British West Indies) [Bahamas] [yellow star].
Type specimen: MCZ 44338, holotype, see Barbour and Loveridge (1946) and Seidel (1988).
Etymology:* Eponym honoring Japheth Vernon (J.V.) Malone (1886–1973), Bahamas; politician from Abaco, Bahamas, and Commissioner for Inagua, who collected the types.

Trachemys stejnegeri vicina (Barbour and Carr 1940)
Dominican Slider


Peter Paul van Dijk / Vuelta Larga, Dominican Republic


left: Harald Artner / Dominican Republic / captivity
right: Peter Paul van Dijk / Vuelta Larga, Dominican Republic


(subspecies: *vicina* = blue; orange dot = possibly introduced; star = type locality)


Distribution: Dominican Republic, Haiti
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 30,366 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 51,727 sq. km
Size (Max SCL): male 22.2 cm, female 27.3 cm (Seidel and Inchaustegui Miranda 1984; Seidel 1988)
Synonymy:
Pseudemys stejnegeri vicina Barbour and Carr 1940:408
Pseudemys terrapen vicina, Pseudemys decussata vicina, Chrysemys decussata vicina, Chrysemys stejnegeri vicina, Chrysemys terrapen vicina, Trachemys stejnegeri vicina
Type locality: "Sanchez, San Domingo" [Dominican Republic]

[yellow star].
Type specimen: FMNH 5977, holotype, see Marx (1958) and Seidel (1988).
Etymology:* Latin, *vicinus*, related or similar; referring to the close relationship between *vicina* and *stejnegeri*.

Trachemys taylori (Legler 1960) (07:18, 12:17) (51, 55)*
(or *Trachemys venusta taylori*)
Cuatro Cienegas Slider, *Jicotea de Cuatro Ciénegas*


James R. Buskirk / Cuatro Ciénegas, Coahuila, Mexico


John B. Iverson / Cuatro Ciénegas, Coahuila, Mexico



(star = type locality)

Distribution: Mexico (*Coahuila*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 887 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 1,138 sq. km
Size (Max SCL): male 17.9 cm, female 21.8 cm (Legler 1990; Ceballos et al. 2013; Legler and Vogt 2013)
IUCN Red List:* Critically Endangered (CR A2ce+4bce; B1ab (i,ii,iii,iv,v)+2ab(i,ii,iii,iv,v)) (Valenzuela-Ceballos, Stanford, and Castañeda Gaytán, in press). Previously: Endangered (EN A4e; B1ab(iii,v)+2ab(iii,v)) (van Dijk and Flores-Villela 2007)
Synonymy:
Pseudemys scripta taylori Legler 1960:75
Chrysemys scripta taylori, Chrysemys gaigeae taylori, Chrysemys taylori, Trachemys scripta taylori, Trachemys nebulosa taylori, Trachemys ornata taylori, Trachemys taylori, Trachemys venusta taylori
Type locality: "16 km. S Cuatro Ciénegas, Coahuila, México" [yellow star].
Type specimen: KU 46952, holotype, see Duellman and Berg (1962).
Etymology:* Eponym honoring Edward H. Taylor (1889–1978), USA / Philippines; zoologist and herpetologist at the University of Kansas who also worked in the Philippines.

Trachemys terrapen (Bonnaterre 1789) (09:6, 14:22, 21:59)
Jamaican Slider, Bahamian Slider


Kurt A. Buhlmann / CCB / Black River, St. Elizabeth Parish, Jamaica


left: Kurt A. Buhlmann / CCB / Black River, St. Elizabeth Parish, Jamaica
right: Douglas B. Booher / Treasure Beach, St. Elizabeth Parish, Jamaica / melanistic male


(orange dots = introduced and hybrids; stars = type localities)

Distribution: Bahamas (*Cat Island, Eleuthera, Exuma, Long Island, South Andros*) *[prehistorical introductions?]*, Jamaica
Introduced: Bahamas (*Grand Bahama, Great Abaco, New Providence, South Bimini [including hybrids with Trachemys spp.]*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 11,066 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 208,767 sq. km
Size (Max SCL): male 24.3 cm, female 30.0 cm (Tuberville et al. 2005)
IUCN Red List: Vulnerable (VU B1+2c [ver. 2.3]) (TFTSG 1996)
TFTSG Provisional Red List:* Critically Endangered (CR) (Parham and Wilson, in prep., 2025)
USA ESA:* Endangered, population on Cat Island, Bahamas (USFWS 1991b)
Synonymy:
Testudo terrapen Lacepède 1788:129, synopsis[table] (*nomen suppressum*) (09:6)
Type locality: "aux Antilles, & particulièrement à la Jamaïque" [Jamaica] [yellow star, location estimated].

Type specimen: Possibly MNHN, not located, see Seidel (1988).
Etymology:* Vernacular (Native American, Powhatan), *torope*, *terrapin*, *terrapen*, or *turpin*, indigenous general name for a turtle, see Dodd et al. (2016).
Comment: Name suppressed by ICZN (2005a) as published in a rejected and invalid non-binomial work, see Savage (2003).

Testudo terrapen Bonnaterre 1789:30
Pseudemys terrapen, Pseudemys terrapen terrapen, Chrysemys (Trachemys) terrapen, Chrysemys terrapen, Chrysemys terrapen terrapen, Trachemys terrapen, Trachemys terrapen terrapen
Type locality: "La Jamaïque" [Jamaica] [yellow star, location estimated].
Type specimen: Possibly MNHN, not located, see Seidel (1988).
Etymology:* Vernacular (Native American, Powhatan), *torope*, *terrapin*, *terrapen*, or *turpin*, indigenous general name for a turtle, see Dodd et al. (2016).

Testudo palustris Gmelin 1789:1041 (senior homonym, not = *Testudo palustris* Le Conte 1830 [= *Malaclemys terrapin terrapin*])
Trachemys palustris, Pseudemys palustris, Chrysemys scripta palustris, Pseudemys palustris palustris
Type locality: "Jamaicae aquis stagnantibus" [Jamaica] [yellow star, location estimated].
Type specimen: Not known or located, see Seidel (1988); no known figure of type specimen, see also Schoepff (1792:xi).
Etymology:* Latin, *palustris*, swamp-dwelling; referring to its habitat in stagnant waters.
Comment: Description cited as sourced from Browne (1756:465).


Testudo fasciata Suckow 1798:40 (senior homonym, not = *Testudo fasciata* Daudin 1801 [= *Homopus areolatus*])
Type locality: "Amboina, und besonders in Nordamerika zu Carolina" [in error]. Restricted to "Jamaica" [yellow star, location estimated] by Seidel (1988:23).
Type specimen: Not known or located, see Seidel (1988).
Etymology:* Latin, *fascia*, band, ribbon, bundle; referring to the stripes on the head.

Testudo rugosa Shaw 1802:28 (*partim*, junior homonym and *nomen dubium*, not = *Testudo rugosa* Van-Ernest *in* Daudin 1801 [= *Chelonia mydas*])
Emys rugosa, Clemmys rugosa, Chrysemys scripta rugosa, Pseudemys rugosa
Type locality: Not designated. Restricted to "Rio Jobabo drainage in eastern Cuba" [orange star, location estimated] by Mittleman (1947:176).
Type specimen: RCSM 990 (formerly LM s/n), figured in Shaw (1802,pl.4) and Barbour and Carr (1940:pl.1).
Etymology:* Latin, *ruga*, wrinkle, ridge, rugose; referring to the ridged, rugose shell.
Comment:* Name synonymized *in partim* under *Trachemys decussata decussata* and *Trachemys terrapen*.

Emys rugosa livida Gray 1831d:30
Type locality: "America septentrionali?"
Type specimen: Possibly OUM, holotype, apparently lost, not listed by Nowak-Kemp and Fritz (2010).
Etymology:* Latin, *livida*, bluish or lead-colored; referring to the shell color.

Pseudemys felis Barbour 1935:205
Pseudemys palustris felis, Pseudemys terrapen felis, Chrysemys decussata felis, Chrysemys felis, Chrysemys terrapen felis, Trachemys terrapen felis, Trachemys felis
Type locality: "Tea Bay, Cat Island, Bahamas" [green star].
Type specimen: MCZ 38385, holotype, see Barbour and Carr (1940) and Barbour and Loveridge (1946).
Etymology:* Toponym for Cat Island, Bahamas; and Latin, *felis*, cat.

Trachemys venusta (Gray 1856b) (07:18, 10:6, 10:10, 11:6, 12:18, 14:23, 17:26) (51, 52, 55)*
Eastern Central American Slider
(includes 4 subspecies)




** (subspecies: *venusta* = red, *cataspila* = blue, *iversoni* = green, *uhrigi* = brown; orange dots = introduced)

Distribution:* Belize, Guatemala, Honduras, Mexico (*Campeche, Chiapas, Oaxaca, Quintana Roo, San Luis Potosi, Tabasco, Tamaulipas, Veracruz, Yucatán*), Nicaragua
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 478,358 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 1,391,100 sq. km
Size (Max SCL): male 35.0 cm, female 44.0 cm (see subspp.)
IUCN Red List:* Least Concern (LC) (Bock, Cupul-Magaña, Páez, Reyes-Grajales, and van Dijk, in press). Previously: TFTSG Provisional Red List: Data Deficient (DD) (2011, 2018)

Trachemys venusta venusta (Gray 1856b) (07:18, 10:10, 11:6, 12:18, 14:23, 17:26, 21:55)
Central American Slider, *Jicotea*


Eduardo Reyes-Grajales / Juarez, Chiapas, Mexico / female


Eduardo Reyes-Grajales / Juarez, Chiapas, Mexico


**

(subspecies: *venusta* = red, *cataspila* = blue, *iversoni* = green, *uhrigi* = brown; star = type locality)

Distribution: Belize, Guatemala, Mexico (*Campeche, Chiapas, Oaxaca, Quintana Roo, Tabasco, Tamaulipas, Veracruz*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 204,008 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 355,433 sq. km

Size (Max SCL): male 34.1 cm, female 42.4 cm (Legler 1990; Rodrigues et al. 2018)

IUCN Red List:* Least Concern (LC) (Bock, Cupul-Magaña, Páez, Reyes-Grajales, and van Dijk, in press)


Synonymy:


Emys venusta Gray 1856b:24 (12:18)
Callichelys venusta, Pseudemys scripta venusta, Chrysemys scripta venusta, Trachemys scripta venusta, Trachemys ornata venusta, Trachemys venusta, Trachemys venusta venusta
Type locality: "Southern States of America; Honduras." Restricted to "Honduras" by Boulenger (1889:81) and Smith and Smith (1980:495); further restricted to "Belize (British Honduras)" by McCord et al. (2010:40) and to "Belize City, Belize (= British Honduras until 1981)" [yellow star] by Legler and Vogt (2013:266).
Type specimen: NHMUK 1947.3.4.80 (formerly 1845.8.5.26), figured (pl.12A), lectotype, designated by Smith and Smith (1980:495).
Etymology:* Latin, *venusta*, beautiful, graceful, elegant, lovely; referring to the attractive head markings.

Emys (Clemmys) salvini Günther 1885:4
Emys salvini, Emys salvinii, Pseudemys salvini
Type locality: "Guatemala" [no star, location unknown, but probably Petén Department].
Type specimen: NHMUK 1946.1.22.76 (formerly 1862.2.19.5), holotype, figured (pl.2-3).
Etymology:* Eponym honoring Osbert Salvin (1835–1898), United Kingdom; ornithologist, herpetologist, and explorer of Guatemala, who obtained the type.

Trachemys venusta cataspila (Günther 1885) (07:18, 10:10, 11:6, 17:26) (55)*
Huastecan Slider, *Jicotea Huasteca*


James R. Buskirk / Aldama, Tamaulipas, Mexico


left: John B. Iverson / nr. Llerda, Tamaulipas, Mexico
right: James R. Buskirk / Aldama, Tamaulipas, Mexico


**

(subspecies: *venusta* = red, *cataspila* = blue; stars = type localities)

Distribution: Mexico (*San Luis Potosi, Tamaulipas, Veracruz*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 96,417 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 143,716 sq. km

Size (Max SCL): male 31.2 cm, female 34.0 cm (Legler 1990; Legler and Vogt 2013)

IUCN Red List:* Least Concern (LC) (Bock, Cupul-Magaña, Páez, Reyes-Grajales, and van Dijk, in press)

Synonymy:

Emys ventricosa Gray 1856b:28 (*nomen suppressum*)
Pseudemys ventricosa
Type locality: Not known.
Type specimen: NHMUK 1848.7.28.24, holotype, see Smith and Smith (1980).
Etymology:* Latin, *ventricosa*, potbellied, with thick hindquarters; referring to bulges on the sides of the carapace.
Comment: Name suppressed by ICZN (1985b), see Legler et al. (1980).

Emys (Clemmys) cataspila Günther 1885:4 (*nomen protectum*)
Emys cataspila, Pseudemys cataspila, Chrysemys ornata cataspila, Pseudemys scripta cataspila, Pseudemys ornata

cataspila, Chrysemys scripta cataspila, Trachemys scripta cataspila, Trachemys ornata cataspila, Trachemys venusta cataspila, Trachemys cataspila

Type locality: "Mexico." Restricted to "Alvarado, Veracruz, Mexico" [in error] [orange star] by Smith and Taylor (1950a:346, 1950b:32); and to "Tampico, Tamaulipas" [Mexico] [yellow star] by Smith and Smith (1980:486).

Type specimen: NHMUK 1947.3.4.25 (formerly 1851.6.2.4), figured in Gray (1856b:pl.12), lectotype, designated by Legler and Vogt (2013:259).

Etymology:* Greek, *kata*, down, against, and *spilos*, stain, dirt; referring to the dark spot near the lower edge of the second and third costal scutes.


Comment: Name conserved by ICZN (1985b), see Legler et al. (1980).

Trachemys venusta iversoni McCord, Joseph-Ouni, Hagen, and Blanck 2010 (10:10, 11:6, 17:26)


Yucatan Slider



Eduardo Reyes-Grajales / Homún, Yucatán, Mexico



John B. Iverson / *left*: nr. Muná, Yucatán, Mexico, *right*: Cobá, Quintana Roo, Mexico



(subspecies: *venusta* = red, *iversoni* = green, *uhrigi* = brown; star = type locality)

Distribution: Mexico (*Quintana Roo, Yucatán*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO): 58,268 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 70,007 sq. km

Size (Max SCL): male 20.4 cm, female 24.3 cm (Legler and Vogt 2013)

IUCN Red List:* Near Threatened (NT A2cd+4cd) (Bock, Cupul-Magaña, Páez, Reyes-Grajales, and van Dijk, in press)

Synonymy:

Trachemys venusta iversoni McCord, Joseph-Ouni, Hagen, and Blanck 2010:45

Type locality: "Cenote on the north side of the highway, 13.8 km east of Buctzotz, Yucatán, Mexico" [yellow star].

Type specimen: UF 50478, holotype.

Etymology:* Eponym honoring John B. Iverson (1949–), USA; herpetologist and turtle researcher at Earlham College, Indiana.

Trachemys venusta uhrigi McCord, Joseph-Ouni, Hagen, and Blanck 2010 (10:10, 11:6, 14:23, 17:26, 21:55) (52, 55)*


Uhrig's Slider



Dennis W. Uhrig / nr. Mezapa, Honduras / captivity



left: Dennis W. Uhrig / nr. Mezapa, Honduras / captivity
right: Volker Naths and Annett Werner / No data / captivity



** (subspecies: *venusta* = red, *uhrigi* = brown; orange dot = introduced, star = type locality)

Distribution:* Guatemala, Honduras, Nicaragua

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 120,010 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 270,618 sq. km

Size (Max SCL): male 35.0 cm, female 44.0 cm (Moll 1994)

IUCN Red List:* Least Concern (LC) (Bock, Cupul-Magaña, Páez, Reyes-Grajales, and van Dijk, in press)

Synonymy:

Trachemys venusta uhrigi McCord, Joseph-Ouni, Hagen, and Blanck 2010:43


Type locality: "Río Chamelecón drainage 3 km south of San Pedro Sula, northwestern Caribbean coastal Honduras" [yellow star].


Type specimen:* UF 157800, holotype; genotyped by Fritz et al. (2023).

Etymology:* Eponym honoring Dennis W. Uhrig (1952–), USA; amateur herpetologist and turtle breeder.

Trachemys yaquia (Legler and Webb 1970) (07:18)
Yaqui Slider, *Jicotea del Yaqui*


James R. Buskirk / Rio Mayo, below Presa Mocuzari, Sonora, Mexico


Philip C. Rosen / Huásabas, Río Bavisbe, Sonora, Mexico


**

(star = type locality)

Distribution: Mexico (*Sonora*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 29,613 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 87,982 sq. km

Size (Max SCL): male 26.8 cm, female 30.9 cm (Legler 1990; Ceballos et al. 2013; Legler and Vogt 2013)

IUCN Red List:* Data Deficient (DD) (Reyes-Grajales, García-Caballero, and Butterfield, in press). Previously: Vulnerable (VU B1ab(iii)+2ab(iii)) (Frost, Hammerson, and Gadsden 2007)

Synonymy:

Pseudemys scripta yaquia Legler and Webb 1970:158
Chrysemys scripta yaquia, Pseudemys ornata yaquia, Trachemys scripta yaquia, Trachemys dorbigni yaquia, Trachemys ornata yaquia, Trachemys yaquia

Type locality: "Río Mayo, Conicarit, Sonora, México (27º14' N, 109º06' W)" [yellow star].

Type specimen: UU 6030, holotype, see Smith and Smith (1980).

Etymology:* Ethnonym honoring the indigenous Yaqui people who live in this region.

Subfamily Emydinae Rafinesque 1815 (17:17)

Emydine Pond Turtles
(includes 6 genera, 15 species, 24 taxa)
Emidania Rafinesque 1815:75
Emydidae Bell 1825a:302
Emydinae Cope 1870b:123

Actinemys Agassiz 1857a (07:22, 09:16, 10:12, 11:7, 14:24, 17:35)
Western Pond Turtles, Pacific Pond Turtles
(includes 2 species, 2 taxa)
Actinemys Agassiz 1857a:252,444


Type species: *Actinemys marmorata* [= *Emys marmorata* Baird and Girard 1852], by original designation.

Etymology:* Greek, *aktinos*, a ray or beam, and *emys*, turtle; referring to the carapacial rays.

Actinemys marmorata (Baird and Girard 1852) (07:22, 10:15, 17:35)
Northern Pacific Pond Turtle, Northwestern Pond Turtle


David J. Germano / CBFTT / Fresno, San Joaquin Valley, California


R. Bruce Bury / CBFTT / *left*: Klamath Lake region, Oregon, *right*: Shasta Co., California


**

(orange dots = probably translocated; stars = type localities)

Distribution: Canada (*British Columbia*), USA (*California, Nevada, Oregon, Washington*)

Introduced: Australia (*New South Wales*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 236,607 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 488,568 sq. km

Size (Max SCL): male 24.1 cm, female 20.0 cm (Lubcke and Wilson 2007; Bury and Germano 2008 CBFTT; Ernst and Lovich 2009)

CBFTT Account: Bury and Germano (2008)

IUCN Red List: Vulnerable (VU A1cd [ver. 2.3]) (TFTSG 1996)

TFTSG Provisional Red List:* Vulnerable (VU) (Bury, Germano, and Ashton, in prep., 2025). Previously: Vulnerable (VU) (2011)

USA ESA:* Proposed Threatened; final determination not yet decided; Species Status Assessment: USFWS (2023c,d)

Synonymy:

Emys marmorata Baird and Girard 1852:177

Actinemys marmorata, Clemmys marmorata, Geoclemmys marmorata, Chelopus marmoratus, Melanemys marmorata, Clemmys marmorata, Clemmys marmorata marmorata, Actinemys marmorata marmorata, Emys marmorata, Emys marmorata marmorata

Type locality: "Puget Sound" [Washington, USA] [yellow star].
Type specimens: USNM 88, 7594–96, 131830 (formerly 7593), syntypes (5), see Cochran (1961) and Reynolds et al. (2007).
Etymology:* Latin, *marmor*, marble, and *-atus*, provided with; referring to the marbled appearance of the carapace.

Emys nigra Hallowell 1854:91 (senior homonym, not = *Emys nigra* Blyth 1856 [= *Siebenrockiella crassicollis*])

Type locality:* "Posa Creek, Lower California" [Poso Creek, Kern County, California, USA] [green star].
Type specimen: USNM 26, holotype, not figured, apparently lost, see Reynolds et al. (2007).
Etymology:* Latin, *nigra*, black; referring to the shell color.
Comment: Reynolds et al. (2007:14) suggested that this paper was published in 1856, but that was the year that the published numbers in Volume 7 from 1854–55 were bound together and indexed. Hollowell's description was issued in June 1854 in Vol. 7, No. 3.

Clemmys wosnessenskyi Strauch 1862:114

Geoclemmys wosnessenskyi

Type locality:* "Rio Sacramento in Californien" [USA] [nr. Sutter's Fort, Sacramento, California] [blue star].
Type specimen: ZIN 94, holotype, see Strauch (1890).
Etymology:* Eponym honoring Ilya G. Voznesensky (1816–1871), Russia; naturalist and explorer who collected the type in California in 1841 while visiting John Sutter at his fort in "New Helvetia."

Actinemys pallida (Seeliger 1945) (07:22, 10:15, 17:35)

Southern Pacific Pond Turtle, Southwestern Pond Turtle, *Tortuga de Charcos*




Robert H. Goodman, Jr. / Camp Pendleton, San Diego Co., California





left: Kelly Herbinson / Ventura Co., California
right: Dan C. Holland / San Diego Co., California



** (orange dots = probably translocated; star = type locality)

Distribution: Mexico (*Baja California*), USA (*California*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 75,945 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 253,045 sq. km

Size (Max SCL): male 17.9 cm, female 16.4 cm (Germano and Riedle 2015)

CBFTT Account: Bury and Germano (2008) [as part of *A. marmorata*]

IUCN Red List: Vulnerable (VU A1cd [ver. 2.3]) (TFTSG 1996), as part of *Actinemys marmorata*

TFTSG Provisional Red List:* Endangered (EN) (Germano, Bury, and Valdez-Villavicencio, in prep., 2025). Previously: Vulnerable (VU) (2011), as part of *Actinemys marmorata*

USA ESA:* Proposed Threatened; final determination not yet decided; Species Status Assessment: USFWS (2023c,d)

Synonymy:

Clemmys marmorata pallida Seeliger 1945:158

Actinemys marmorata pallida, Emys marmorata pallida, Emys pallida, Actinemys pallida

Type locality: "Lower Coyote Creek, near Alamitos, Orange County, California" [USA] [yellow star].

Type specimen: MVZ 6716, holotype, see Crippen (1962) and Rodríguez-Robles (2003).

Etymology:* Latin, *pallidus*, pale; referring to the light color of the lateral and ventral neck.

Clemmys Ritgen 1828

Spotted Turtles

(includes 1 species, 1 taxon)

Chelopus Rafinesque 1815:75 (*nomen nudum*)

Etymology:* Greek, *chelone*, turtle, and *pus*, foot.

Clemmys Ritgen 1828:270

Type species: *Clemmys punctata* [= *Testudo punctata* Schoepff 1792 = subjective synonym of *Testudo guttata* Schneider 1792], by subsequent designation by Baur (1892:43).

Etymology:* Greek, *klemmys*, turtle, or *kleptein*, to steal, and *emys*, turtle; apparently referrring to the ability to retract and hide the head and neck.

Chelopus Rafinesque 1832:64

Type species: *Chelopus punctatus* [= *Testudo punctata* Schoepff 1792 = subjective synonym of *Testudo guttata* Schneider 1792], by original monotypy.

Etymology:* Greek, *chelone*, turtle, and *pus*, foot.

Nanemys Agassiz 1857a:252,442

Type species: *Nanemys guttata* [= *Testudo guttata* Schneider 1792], by original monotypy.

Etymology:* Greek, *nanos*, dwarf, and *emys*, turtle; referring to the small size of the species in this genus.

Melanemys Shufeldt 1919:157

Type species: *Melanemys guttatus* [= *Testudo guttata* Schneider 1792], by subsequent designation by Dunn (1920:8).

Etymology:* Greek, *melan*, black, dark, and *emys*, turtle.

Clemmys guttata (Schneider 1792) (17:29)


Spotted Turtle



Barry Mansell / CRM 3 / Seminole Co., Florida



left: Richard D. Bartlett / CRM 3 / Clinch Co., Georgia
right: James H. Harding / Ottawa Co., Michigan



** (orange dots = questionable, probably translocated; star = type localities)

Distribution: Canada (*Ontario*), USA (*Connecticut, Delaware, Florida, Georgia, Illinois, Indiana, Maine, Maryland, Massachusetts, Michigan, New Hampshire, New Jersey, New York, North Carolina, Ohio, Pennsylvania, Rhode Island, South Carolina, Vermont, Virginia, West Virginia*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 756,937 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 2,097,291 sq. km

Size (Max SCL): male 12.7 cm, female 12.6 cm (Klemens 1993; Graham 1995); unsexed 13.6 cm (Taylor 1991)

IUCN Red List: Endangered (EN A2cde+4ce) (van Dijk 2011).

Previously: Vulnerable (VU) (TFTSG *in* Baillie and Groombridge 1996)

TFTSG Provisional Red List:* Endangered (EN) (Jones, Willey, Chandler, Howell, Litzgus, Parren, Lipps, Lee, Coury, and Erb, in prep., 2025)

CITES: Appendix II (2013)

Synonymy:

Testudo guttata Schneider 1792:264

Emys guttata, Geoclemys guttata, Nanemys guttata, Clemmys guttata, Clemmys guttatus, Geoclemmys guttata, Chelopus guttatus, Melanemys guttatus

Type locality: Not designated. Restricted to "the vicinity of Philadelphia" [Pennsylvania, USA] [yellow star] by Mittleman (1945:171).

Type specimen: Not located, apparently lost, originally in the collection of Bloch in Dresden.

Etymology:* Latin, *guttatus*, spotted or speckled; referring to the yellow spots on the carapace.

Comment: Description cited as also sourced from Schneider (1789:30) and Seba (1734:pl.80,f.7).

Testudo punctata Schoepff 1792:25 (junior homonym, not = *Testudo punctata* Lacepède 1788 (*nomen suppressum*) = *Testudo punctata* Bonnaterre 1789 [= *Lissemys punctata punctata*])

Emys punctata, Clemmys punctata, Terrapene punctata, Chelopus punctatus

Type locality: "in paludosis Americae septentrionalis... Philadelphiam" [Philadelphia, Pennsylvania, USA] [yellow star].

Type specimen: Not located, type specimen figured (pl.5), labeled *Testudo punctata* Muhlenberg.

Etymology:* Latin, *punctata*, spotted or punctured; referring to the yellow spots on the carapace.

Testudo anonyma Schneider *in* Schoepff 1792:25 (*nomen nudum*)

Etymology:* Greek, *anonymos*, nameless or unknown.

Testudo quarta Hermann *in* Schweigger 1812:434 (*nomen nudum*)

Etymology:* Latin, *quarta*, fourth; referring to the fourth species on a list.
Comment:* Erroneous emendation of "Quarta testudo" listed by Hermann (1804:225).

Geoclemmys sebae Gray 1869a:188
Type locality: Not designated. Restricted to "vicinity of Philadelphia" [Pennsylvania, USA] [yellow star] by Schmidt (1953:91).
Type specimen: NHMUK 1853.11.30.25, possible holotype; erroneously listed by Uetz et al. (2019) as the type of *Testudo guttata* Schneider.
Etymology:* Eponym honoring Albertus Seba (1665–1736), Netherlands; pharmacist and naturalist with a large natural history collection.

Emydoidea Gray 1870c (07:21, 09:16, 10:12, 11:7, 14:24, 17:36)

Blanding's Turtles
(includes 1 species, 1 taxon)

Emydoidea Gray 1870c:19
Type species: *Emydoidea blandingii* [= *Cistuda blandingii* Holbrook 1838b], by original monotypy.
Etymology:* Greek, *emydos*, turtle, and *-oides*, resembling; referring to its similarity with the genus *Emys*.

Neoemys Lindholm 1929:282 (*nomen novum*)
Etymology:* Greek, *neo*, young, new, and *emys*, turtle; referring to its similarity with the genus *Emys*.
Comment:* Unjustified replacement name for *Emydoidea*.

Emydoidea blandingii (Holbrook 1838b) (17:36, 21:60)


Blanding's Turtle



Janet Hostetter / CBFTT / Weaver Dunes, Minnesota



left: Terry E. Graham / CBFTT / Devens, Massachusetts
right: James H. Harding / CBFTT / Michigan



(orange dots = questionable, probably translocated; stars = type localities)

Distribution:* Canada (*Nova Scotia, Ontario, Québec*), USA (*Illinois, Indiana, Iowa, Kansas, Maine, Massachusetts, Michigan, Minnesota, Missouri, Nebraska, New Hampshire, New York, Ohio, Pennsylvania, South Dakota, Wisconsin*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 865,969 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 2,436,571 sq. km

Size (Max SCL): male 28.4 cm, female 23.2 cm (Blasus et al. 2004; Congdon et al. 2008 CBFTT; Ceballos et al. 2013)

CBFTT Account: Congdon, Graham, Herman, Lang, Pappas, and Brecke (2008)

IUCN Red List: Endangered (EN A2cde+4ce) (van Dijk and Rhodin 2011). Previously: Near Threatened (NT) (TFTSG *in* Baillie and Groombridge 1996)

TFTSG Provisional Red List:* Endangered (EN) (Mifsud, McGuire, Congdon, and Harding, in prep., 2025)

CITES: Appendix II (2013)

Synonymy:

Testudo flava Lacepède 1788:135, synopsis[table] (*nomen suppressum*) (09:6)
Type locality: "Amérique..[&]..l'isle de l'Ascension." Restricted to "Amerika" [America = USA] by Wermuth (1956:407).
Type specimen: Possibly MNHN, not located, type specimen figured (pl.opp.135).
Etymology:* Latin, *flava*, yellow, golden yellow; referring to the numerous yellow markings on the carapace and neck.
Comment: Name suppressed by ICZN (1963), see Wermuth (1956) and Mertens and Wermuth (1961), and by ICZN (2005a) as published in a rejected and invalid non-binomial work, see Savage (2003).

Testudo flava Bonnaterre 1789:26 (*nomen oblitum*)
Type locality: "Amérique, l'ile de l'Ascension."
Type specimen: Possibly MNHN, not located, type specimen figured in Lacepède (1788:pl.opp.135).
Etymology:* Latin, *flava*, yellow, golden yellow; referring to the numerous yellow markings on the carapace and neck.

Testudo meleagris Shaw 1793:147 (*nomen suppressum*)
Lutremys meleagris, Emys meleagris
Type locality: "America" [USA].
Type specimen: Not located, type specimen figured (f.144), see Wermuth (1956).
Etymology:* Greek, *meleagris*, guinea-fowl (peacock); referring to the speckled carapace resembling a guinea-fowl.
Comment: Name suppressed by ICZN (1963), see Wermuth (1956) and Mertens and Wermuth (1961).

Cistuda blandingii Holbrook 1838b:35 (*nomen protectum*)
Cistudo blandingii, Emys blandingii, Emydoidea blandingii, Neoemys blandingii
Type locality: "Fox river, a tributary of the Illinois" [Illinois, USA] [yellow star].
Type specimen: ANSP 26123, holotype, listed as *Cistudo blandingii* by Malnate (1971).
Etymology:* Eponym honoring William Blanding (1772–1857), USA; physician and naturalist who first collected the species.
Comment: Name conserved by ICZN (1963), see Wermuth (1956) and Mertens and Wermuth (1961).

Emys twentei † Taylor 1943:250
Type locality: "north side of the Cimarron river, 13 miles southwest of Meade, Meade County, Kansas (Loc. No. 7, XI Ranch)" [USA] [gray star].
Type specimen: KU 6478, holotype, fossil, partial carapace, figured (pl.20), see Vlachos (2018).
Geological age: Pleistocene, high terrace sands.
Etymology:* Eponym honoring John W. Twente (1887–1966), USA; paleontologist at the University of Kansas who collected the type.
Comment: Considered a synonym of *blandingii*, see Preston and McCoy (1971) and Vlachos (2018).

Emys Duméril 1805 (07:21, 09:16, 10:11, 10:12, 11:7, 14:24)

European Pond Turtles

(includes 2 species, 8 taxa)

Emydes Brongniart 1805:27 (*nomen suppressum*)

Etymology:* Greek, *emydos*, turtle.

Comment: Name suppressed by ICZN (1995b), see Webb (1993).

Emys Duméril 1805:76 (*nomen protectum*) (10:11)

Type species: *Emys europaea* Schweigger [= *Testudo europaea* Schneider 1783 = subjective synonym of *Testudo orbicularis* Linnaeus 1758], by subsequent designation by Fitzinger (1843:29).

Comment: Name conserved by ICZN (1995b), see Webb (1993).

Etymology:* Greek, *emys*, turtle.

Hydrone Rafinesque 1814:66

Type species: *Hydrone orbicularis* [= *Testudo orbicularis* Linnaeus 1758], by subsequent designation by Loveridge and Williams (1957:201).

Etymology:* Greek, *hydraino*, to water, to wash, *hydrone*, water dweller.

Emyda Rafinesque 1815:75 (senior homonym and *nomen novum*, not = *Emyda* Gray 1830e [= *Lissemys*])

Etymology:* Greek, *emydos*, turtle.

Lutremys Gray 1844:31

Type species: *Cistudo* (*Lutremys*) *europaea* [= *Testudo europaea* Schneider 1783 = subjective synonym of *Testudo orbicularis* Linnaeus 1758], by original monotypy.

Etymology:* Latin, *lutum*, mud, dirt, clay, and Greek, *emys*, turtle.

Emys orbicularis (Linnaeus 1758) (21:61)

European Pond Turtle

(includes 7 subspecies and several taxonomically unspecified populations of *E. orbicularis* sensu lato)



**

(subspecies: *orbicularis* = red [light red = extirpated], *eiselti* = purple, *galloitalica* = blue, *hellenica* = brown, *ingauna* = pink, *occidentalis* = green, *persica* = tourmaline, taxonomically unspecified populations of *E. orbicularis* sensu lato = gray (Algeria, Tunisia, southern Turkey); overlap = intergrades; orange dots = introduced, reintroduced, translocated, trade, or questionable; red dots = extirpated)

Distribution:* Albania, Algeria, Armenia, Austria, Azerbaijan, Belarus, Belgium (*extirpated*), Bosnia and Herzegovina, Bulgaria, Croatia, Czech Republic (*extirpated, reintroduced*), France (*Continental, Corsica [prehistorical introduction?]*), Georgia, Germany, Greece, Hungary, Iran, Italy (*Continental, Sardinia [prehistorical introduction]*), Kazakhstan, Kosovo [*possibly introduced?*], Latvia, Lithuania, Luxembourg (*extirpated*), Moldova, Montenegro, Morocco, Netherlands (*extirpated, reintroduced*), North Macedonia, Poland, Portugal, Romania, Russia, Serbia, Slovakia, Slovenia, Spain (*Continental*), Switzerland (*extirpated, reintroduced*), Syria, Tunisia, Turkey, Turkmenistan, Ukraine

Introduced:* Denmark, Spain (*Balearic Islands*), United Kingdom

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 4,567,751 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 12,194,183 sq. km

Size (Max SCL): male 21.0 cm, female 23.2 cm (see subspp.)

IUCN Red List: Near Threatened (NT) (TFTSG 1996).

Regional (Europe): Near Threatened (NT) (Luiselli and Vamberger 2024). Previously: Near Threatened (NT) (Cox and Temple 2009); Near Threatened (NT) (van Dijk and Sindaco 2004). Regional: (European Union–EU27): Vulnerable (VU A2bcde) (Cox and Temple 2009)

CITES: Appendix III (Ukraine) (2021)

Emys orbicularis orbicularis (Linnaeus 1758) (09:17, 17:30, 17:31, 21:61)


European Pond Turtle




Uwe Fritz / Dnieper R. Delta, Ukraine



Anders G.J. Rhodin / southeastern Europe, haplotype Ic / captivity / Nordens Ark, Sweden



**



**

(subspecies: *orbicularis* = red [light red = extirpated], *eiselti* = purple, *galloitalica* = blue, *hellenica* = brown, *ingauna* = pink, *occidentalis* = green, *persica* = tourmaline, taxonomically unspecified population of *E. orbicularis* sensu lato = gray (southern Turkey); overlap = intergrades; orange dots = introduced, reintroduced, translocated, trade, or questionable; red dots = extirpated; stars = type localities)

Distribution:* Austria, Belarus, Belgium (*extirpated*), Bosnia and Herzegovina, Bulgaria, Croatia, Czech Republic (*extirpated, reintroduced*), France, Georgia, Germany, Greece, Hungary, Italy, Kazakhstan, Kosovo [*possibly introduced?*],

Latvia, Lithuania, Luxembourg (*extirpated*), Moldova, Netherlands (*extirpated, reintroduced*), North Macedonia, Poland, Romania, Russia, Serbia, Slovakia, Slovenia, Switzerland (*extirpated, reintroduced*), Turkey, Ukraine

Introduced:* Denmark, Spain (*Balearic Islands*), United Kingdom

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 3,618,563 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 8,873,890 sq. km

Size (Max SCL): male 21.0 cm, female 23.2 cm (Fritz 2001; Itescu et al. 2014; Joos et al. 2017)

CITES: Appendix III (Ukraine) (2021)

Synonymy:

Testudo orbicularis Linnaeus 1758:198 (*nomen protectum*)

Hydrone orbicularis, Emys orbicularis, Emys orbicularis orbicularis

Type locality: "meridionalibus Europae." Restricted to "Mecklenburgisch-Pommersche Seenplatte" [Germany] [yellow star] by Fritz (1992:67), and by neotype designation by Fritz (1994:65).

Type specimen: MNM Z4180, neotype, designated by Fritz (1994:65); holotype originally in UUZM, apparently lost, listed by Linné and Thunberg (1780), but not by Thunberg (1828), Lönnberg (1896), Andersson (1900), Holm (1957), or Wallin (2001).

Etymology:* Latin, *orbiculus*, small circle; referring to the subcircular shell.

Comment: Description cited as also sourced from Ray (1693:254). Name conserved by ICZN (1995b), see Webb (1993).

Testudo lutaria Linnaeus 1758:198

Emydes lutaria, Hydrone lutaria, Emys lutaria, Clemmys (Clemmys) lutaria, Cistudo lutaria

Type locality: "Italia, Oriente." Restricted to "Mecklenburgisch-Pommersche Seenplatte" [Germany] [yellow star] by Fritz (1992:67).

Type specimen: UUZM, holotype, apparently lost, listed by Thunberg (1828), but not by Lönnberg (1896), Andersson (1900), Holm (1957), or Wallin (2001).

Etymology:* Latin, *lutum*, mud, dirt, clay; referring to the typically muddy-bottom aquatic habitat.

Comment: Description cited as sourced from a specimen described by Linnaeus (1749a:139, n.23); also based on Worm (1655:315), Grew (1681:38,pl.3,f.3), and Ray (1693:259).

Testudo terrestris Garsault 1764:pl.675 (senior homonym and *nomen oblitum*, not = *Testudo terrestris* Fermin 1765 [= *Chelus fimbriata*], nor = *Testudo terrestris* Forskål 1775 (*nomen protectum*) [= *Testudo (Testudo) graeca terrestris*]) (10:13)

Type locality: Not designated.

Type specimen: Not located, type specimen figured (pl.675).

Etymology:* Latin, *terrestris*, of the earth, terrestrial.

Testudo europaea Schneider 1783:323

Emys europaea, Terrapene europaea, Cistuda europaea, Cistudo europaea, Cistudo (Lutremys) europaea, Lutremys europaea

Type locality: "meisten Ländern von Europa bis in Preussen" [Germany]. Restricted to "Frankfurt an der Oder" [Germany] [green star] by Fritz (1992:67).

Type specimens: Not known or located.

Etymology:* Toponym for Europe.

Testudo pulchella Schoepff 1801:113 (senior homonym, not = *Emys pulchella* Schweigger 1812 [= *Glyptemys insculpta*])

Emys pulchella

Type locality: Not known. Restricted to "Mecklenburgisch-Pommersche Seenplatte" [yellow star] by Fritz (1992:67).

Type specimens: Not located, syntypes (2), type specimens figured (pl.26).

Etymology:* Latin, *pulchra*, beautiful, pretty, and *-ella*, small or smaller; referring to the attractive spot markings on the carapace, head and extremities.

Terrapene europea Bell 1825a:308 (*nomen novum*)

Etymology:* Toponym for Europe.

Comment: Unjustified emendation or error for *europaea*.

Testudo lutraria Gray 1831d:19 (*nomen novum*)

Emys lutraria

Etymology:* Latin, *lutum*, mud, dirt, clay; referring to the typically muddy-bottom aquatic habitat.

Comment: Unjustified emendation or error for *lutaria*.

Emys turfa † Meyer 1835:67

Cistudo lutaria turfa

Type locality: "Torfmooren bei Enkheim unweit Frankfurt" [Germany] [black star].

Type specimens: Possibly SMNS, syntypes, subfossil, carapacial and appendicular elements, not figured, see Kurck (1917).

Geological age: Holocene, peat bog.

Etymology:* Latin, *turfa*, peat; referring to the peat bog where specimens of this taxon were collected.

Clemmys schlotheimii † Fitzinger 1835:127 (*nomen nudum*) (17:30)

Type locality: Not designated. Restricted to "Burgtonna" [Thuringia, Germany] [dark gray star] by Karl and Paust (2014:156).

Type specimen: MNG 1672, lectotype, fossil, small carapacial fragment, figured in Karl and Paust (2014:pl.4,f.4), designated by Karl and Paust (2014:156).

Geological age: Pleistocene, Eemian Period.

Etymology:* Eponym honoring Ernst F. von Schlotheim (1764–1832), Germany; geologist and paleontologist.

Trionyx schlotheimii † Fitzinger 1835:128 (*nomen nudum*) (17:30)

Type locality: Not designated. Restricted to "Burgtonna" [Thuringia, Germany] [dark gray star] by Karl and Paust (2014:156).

Type specimen: MNG 1673, lectotype, fossil, small carapacial fragment, figured in Karl and Paust (2014:pl.4,f.1,3), designated by Karl and Paust (2014:156).

Geological age: Pleistocene, Eemian Period.

Etymology:* Eponym honoring Ernst F. von Schlotheim (1764–1832), Germany; geologist and paleontologist.

Emys lutaria borealis † Nilsson 1841:208

Type locality: "Gräfve af Brågarps pastorat...Skåne..[&]..Götha kanal, Östergöthland...vid Nordskogsvägen nära intill Svartjordshålan" [Gräfve in Brågarp Parish...Scania..[&]..Göta Canal, East Götaland...along Nordskogs Road near Svartjordshålan] [Skåne..[&]..Östergötland, Sweden]. Restricted to "1,5 kilometer Ö om Norsholms jernvägsstation" [1.5 km E of Norsholm's railroad station] [Östergötland, Sweden] by Munthe (1895:163), and to "Norsholm i Kimstads socken...Östergötland" [Norsholm in Kimstad parish... Östergötland] [Sweden] [light gray star] by Kurck (1917:42); erroneously restricted to "Schonen, Südschweden" [Skåne, southern Sweden] [orange star] by Fritz (1992:67).

Type specimens: NRM CU75037, 75039 (formerly R1743–44), syntypes, subfossil, two nearly complete shells and appendicular elements, figured in Dalman (1820:pl.6-7) and Nilsson (1841:pl.3-4), see Kurck (1917) and Sommer et al. (2009).

Geological age: Holocene, Boreal (Atlantic), gravel sediments, ^{14}C age ca. 4,720–5,190 ybp, calendaric age ca. 5,407–5,941 ybp (ca. 3457–3991 BCE), see Sommer et al. (2009).

Etymology:* Latin, *borealis*, northern; referring to the northern geographic location in sub-boreal Sweden where the specimens were collected.

Testudo (Emys) canstadiensis † Plieninger 1847:208 (*nomen nudum*) (17:31)

Emys canstadiensis

Type locality: "Württemberg...Cannstadt" [Germany] [dark brown star].

Type specimen: Possibly SMNS, holotype, fossil, humerus, figured in Jäger (1861:191), not located.

Geological age: Pleistocene, Sinterkalk.

Etymology:* Toponym for Cannstadt [= Cannstatt], Germany.

Emys pulchra Brandt *in* Gray 1873j:22 (*nomen nudum*)

Type locality: "Europe."

Type specimen: NHMUK 1851.12.9.6, holotype, see Gray (1873j).

Etymology:* Latin, *pulchra*, beautiful; referring to the ornate markings on the head and shell.

Cistudo anhaltina † Giebel 1866a:1

Emys anhaltina

Type locality: "Latdorf...Norddeutschland" [Germany]. Emended to "Lattorf vid Bernberg a. d. Saale" [Germany] by Kurck (1917:23); and to "Latdorf bei Bernburg an der Saale, Deutschland" [Germany] [light brown star] by Fritz (1995:227).

Type specimens:* MLUH GeoS.736a–d, syntypes, subfossil, nearly complete carapace and plastron, figured (pl.1-2), see Kurck (1917), Karl and Paust (2024:pl.1).

Geological age: Holocene, Latdorfer Braunkohle.

Etymology:* Toponym for the Duchy of Anhalt [now part of Saxony-Anhalt], Germany.

Emys lutaria taurica Mehnert 1890:537

Type locality: "Ufer des Dnjepr, einige Meilen von seiner Ausmündung" [Dnieper River estuary, Ukraine] [dark blue star].

Type specimens: Possibly MZUS, not located, histological sections figured (pl.20).

Etymology:* Toponym for the Taurida Governorate, Russia [now Ukraine].

Comment: Histological study of embryos and hatchlings of *Emys lutaria* collected in the Taurida Governorate of southern Russia (now Ukraine), not explicitly a description of a new taxon.

Emys europaea sparsa Dürigen 1897:14

Type locality: Not designated. Restricted to "Ungarn" [Hungary] [pink star, location estimated] by Fritz (1992:67).

Type specimen: Not designated, described from the literature, type specimen figured in Sturm (1828:pl.3,f.c), same figure as Schoepff (1792:pl.1), see Fritz (1992).

Etymology:* Latin, *sparsa*, speckled, spotted; referring to the yellowish speckles on the carapace, head and extremities.

Emys europaea punctata Dürigen 1897:15

Type locality: Not designated. Restricted to "Mecklenburg" [Germany] [yellow star] by Fritz (1992:67).

Type specimen: Not designated, described from the literature, type specimen figured in Sturm (1828:pl.3,f.a-b), see Fritz (1992).

Etymology:* Latin, *punctata*, spotted or punctured; referring to the yellow spots on the carapace.

Emys europaea concolor Dürigen 1897:15

Type locality: Not designated. Restricted to "Mark Brandenburg" [Germany] [purple star] by Fritz (1992:67).

Type specimen: Not designated, described from the literature, type specimen figured in Brandt and Ratzeburg (1829:pl.21), see Fritz (1992).

Etymology:* Latin, *con*, the same, and *color*, color; referring to this taxon having a single-colored, dark carapace.

Emys orbicularis aralensis Nikolsky 1915:24

Type locality: "Lac. Aral" [Kazakhstan] [light blue star].

Type specimen: ZIN 70, lectotype, specimen figured in Nikolsky (1899:pl.3), designated as holotype by Fritz (1992:66).

Etymology:* Toponym for the Aral Sea region, Kazakhstan.

Emys orbicularis luteofusca Fritz 1989:145 (09:17)

Type locality: "See-Ebene westlich von Ereğli, Provinz Konya, Türkei" [Turkey] [turquoise star].

Type specimen: SMNS 4615:1, holotype, see Schlüter and Hallermann (1997).

Etymology:* Latin, *luteus*, yellow, and *fuscus*, dark, dusky; referring to the overall dark-colored yellowish brown shell and soft parts.

Emys orbicularis colchica Fritz 1994:61 (09:17)

Emys colchica

Type locality: "Batumi (Batum)" [Georgia] [olive star].

Type specimen: ZIN 9110a, holotype.

Etymology:* Toponym for the prehistorical Colchis region along the east coast of the Black Sea that preceded the nation of Georgia.


Emys orbicularis eiselti Fritz, Baran, Budak, and Amthauer 1998
Eiselt's Pond Turtle, Turkish Pond Turtle



Peter Praschag / Hatay, Antakya, Turkey / female



Peter Praschag / Hatay, Antakya, Turkey / male



(subspecies: *orbicularis* = red, *eiselti* = purple, taxonomically unspecified population of *E. orbicularis* sensu lato = gray; orange dots = introduced, translocated, trade, or questionable; star = type locality)

Distribution: Syria, Turkey

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO): 8,042 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 13,224 sq. km

Size (Max SCL): male 12.9 cm, female 13.1 cm (Fritz et al. 1998b; Fritz 2001)

TFTSG Provisional Red List: Critically Endangered (CR) (2021)

Synonymy:

Emys orbicularis eiselti Fritz, Baran, Budak, and Amthauer 1998b:113

Type locality: "14 km NE of Fevzipaşa (about 450 m above sea level), Vilayet Gaziantep" [Turkey] [yellow star].

Type specimen: NMW 18551:1, holotype, see Gemel et al. (2019).

Etymology:* Eponym honoring Josef Eiselt (1912–2001), Austria; herpetologist who worked in Asia Minor and collected the type.

Emys orbicularis galloitalica Fritz 1995 (10:14, 21:61)
Franco-Italian Pond Turtle




Uwe Fritz / Plaine des Maures, France





left: Romain Levasseur / Provence, France
right: Uwe Fritz / Olmeda, Sardinia, Italy



** (subspecies: *orbicularis* = red, *galloitalica* = blue, *hellenica* = brown, *ingauna* = pink, *occidentalis* = green, taxonomically unspecified populations of *E. orbicularis* sensu lato = gray (Algeria, Tunisia); overlap [purple; dark green] = intergrades; orange dots = introduced, translocated, trade, or questionable; red dots = extirpated; stars = type localities)

Distribution: France (*Continental, Corsica [prehistorical introduction?]*), Italy (*Continental, Sardinia [prehistorical introduction]*), Spain (*Continental*)

Introduced: Spain (*Balearic Islands [possibly prehistorical]*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 171,097 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 800,235 sq. km

Size (Max SCL): male 15.9 cm, female 16.5 cm (Fritz 1995, 2001; Zuffi et al. 2006; M.A.L. Zuffi, unpubl. data; Joos et al. 2017)

Synonymy:

Emys orbicularis (galloitalica) galloitalica Fritz 1995:217
Emys orbicularis galloitalica
Type locality: "5 km östlich Collobrières, Département Var, Südfrankreich" [France] [yellow star].
Type specimen: MNHN 1993.5804, holotype.
Etymology:* Toponym for Gallia [= Gaul = France] and Italy.

Emys orbicularis (galloitalica) capolongoi Fritz 1995:204 (10:14)
Emys orbicularis capolongoi
Type locality: "Olbia (Sardinien)" [Italy] [green star].
Type specimen: SMF 59593, holotype.
Etymology:* Eponym honoring Domenico Capolongo (1938–), Italy; geographer and zoologist at the University of Bari.

Emys orbicularis (galloitalica) lanzai Fritz 1995:211 (10:14)
Emys orbicularis lanzai
Type locality: "Conca-Mündung bei Fontea, unweit Santa Lucia di Porto-Vecchio (Korsika)" [France] [blue star].
Type specimen: MZUF 19095, holotype.
Etymology:* Eponym honoring Benedetto Lanza (1924–2016), Italy; herpetologist and chiropterologist at the University of Florence.

Emys orbicularis hellenica Valenciennes *in* Bory de Saint-Vincent 1833 (12:19, 14:25)
Hellenic Pond Turtle




Andreas Nöllert / Miranje, Croatia / female





Melita Vamberger / Neretva R., Croatia



** (subspecies: *orbicularis* = red, *galloitalica* = blue, *hellenica* = brown, *ingauna* = pink, taxonomically unspecified population of *E. orbicularis* sensu lato = gray (Algeria, Tunisia); overlap [purple] = intergrades; orange dots = introduced, translocated, trade, or questionable; red dots = extirpated; stars = type localities)

Distribution: Albania, Bosnia and Herzegovina, Croatia, Greece, Italy, Montenegro, North Macedonia, Slovenia

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 190,294 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 649,966 sq. km

Size (Max SCL): male 16.2, female 19.5 cm (Joos et al. 2017)

Synonymy:

Emys hellenica Valenciennes *in* Bory de Saint-Vincent 1833:planches, pl.8 (14:25)
Cistuda hellenica, Emys orbicularis hellenica
Type locality: Not designated. Restricted to "plaine de Nisi que baigne le Pamisus au coeur de la Messénie" [Peloponnes, Greece] [yellow star] by Bibron and Bory de Saint-Vincent (1833:61).
Type specimen: MNHN 1943, holotype, see Fritz (1992).
Etymology:* Toponym for Hellas [= Greece].

Emys iberica Valenciennes *in* Bory de Saint-Vincent 1833:planches, pl.9 (14:25)
Type locality: Not designated. Restricted to "Morée, où l'embouchure de l'Eurotas en est remplie, ainsi que le principal ruisseau de l'île de Tine" [Greece] [green star] by Bibron and Bory de Saint-Vincent (1833:61).
Type specimen: Not located, holotype, figured (pl.9).
Etymology:* Toponym for Iberia, a former ancient kingdom in Georgia in the Caucasus, not the Iberian Peninsula (= Spain).

Emys antiquorum Bory de Saint-Vincent 1835:Atlas, pl.9

[corrigenda] (*nomen nudum et novum*) (14:25)
Etymology:* Latin, *antiquorum*, of ancient times.
Comment: Unjustified replacement name for *iberica*.


Emys (Emys) hofmanni Fitzinger 1835:123 (*nomen novum*) (12:19)
Emys orbicularis hoffmanni, *Cistudo hoffmanni*
Etymology:* Eponym honoring A. Hoffmann (1700s–1800s), Austria; naturalist who revised and published Funke's Naturhistorischer Atlas in Vienna in 1825.
Comment: Unjustified replacement name for *hellenica*.

Emys orbicularis atra Werner 1897:15
Type locality: "Dalmatien und Cephallonia" [Croatia and Greece].
Type specimen: Not located, type specimen figured (pl.1,f.2).
Etymology:* Latin, *atra*, black; referring to the color of the shell.

Emys europaea maculosa Dürigen 1897:15
Type locality: Not designated. Restricted to "Dalmatien" [Croatia] [blue star, location estimated] by Fritz (1992:68).
Type specimen: Not designated.
Etymology:* Latin, *macula*, spot, *maculosa*, rich in spots; referring to the carapacial markings.

Emys orbicularis ingauna Jesu, Piombo, Salvidio, Lamagni, Ortale, and Genta 2004 (10:14)
Ligurian Pond Turtle


Pino Piccardo / Albenga, Italy / female


(subspecies: *orbicularis* = red, *galloitalica* = blue, *hellenica* = brown, *ingauna* = pink; overlap [purple] = intergrades; orange dots = introduced, translocated, trade, or questionable; star = type locality)

Distribution: Italy (*Continental*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 1,701 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 2,139 sq. km
Size (Max SCL): male 13.0 cm, female 15.0 cm (Jesu et al. 2004)
Synonymy:

Emys orbicularis ingauna Jesu, Piombo, Salvidio, Lamagni, Ortale, and Genta 2004:139
Emys (Emys) orbicularis ingauna
Type locality: "Peagna (Comune di Ceriale), Provincia di Savona (Regione Liguria, Italia)" [Italy] [yellow star].
Type specimen: MSNG 50650, holotype.
Etymology:* Ethnonym honoring the Ingauni, ancient people that inhabited the Ligurian region during the Iron Age and Roman period.

Emys orbicularis occidentalis Fritz 1993 (17:32, 21:61)
Western Pond Turtle, Spanish Pond Turtle, Magreb Pond Turtle


Uwe Fritz / Ifrane, Morocco


Cesar Ayres / Gándaras de Budiño, Pontevedra, Spain


(subspecies: *orbicularis* = red, *galloitalica* = blue, *occidentalis* = green, taxonomically unspecified populations of *E. orbicularis* sensu lato = gray (Algeria); overlap [dark green] = intergrades; orange dots = introduced, translocated, trade, or questionable; red dots = extirpated; stars = type localities)

Distribution: Morocco, Portugal, Spain
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 452,588 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 807,495 sq. km
Size (Max SCL): male 16.4 cm, female 16.8 cm (Keller et al. 1998; Joos et al. 2017)
Synonymy:

Emys orbicularis occidentalis Fritz 1993:136 (17:32)
Type locality: "Lagune von Medhiya unweit Kenitra, Marokko" [Morocco] [yellow star].
Type specimen: MNHN 1961.0344, holotype.
Etymology:* Latin, *occidens*, the west, *occidentalis*, coming from the west; referring to the westernmost range of this subspecies.


Emys orbicularis fritzjuergenobsti Fritz 1993:132 (09:18, 17:32)
Type locality: "Castellón de la Plana, Spanien" [Spain] [green star].
Type specimen: SMNS 4639:9, holotype, see Schlüter and Hallermann (1997).
Etymology:* Eponym honoring Fritz-Jürgen Obst (1939–2018), Germany; herpetologist and director of the Staatliches Museum für Naturkunde, Dresden.

Emys orbicularis hispanica Fritz, Keller, and Budde 1996:132 (09:18)
Type locality: "Doñana, Huelva" [Spain] [blue star].
Type specimen: EBD 29209, holotype.
Etymology:* Toponym for Hispania, the ancient Roman name for the Iberian Peninsula and Spain.


Emys orbicularis persica Eichwald 1831 (07:23, 07:24, 09:19, 17:33)
Eastern Pond Turtle, Persian Pond Turtle



James F. Parham / CRM 4 / Shahrbijar, Gilan Prov., Iran



left: Luca Luiselli / Iran
right: Asghar Mobaraki / Gilan Prov., Iran



(subspecies: *orbicularis* = red, *persica* = tourmaline; orange dots = introduced, translocated, trade, or questionable; stars = type localities)

Distribution: Armenia, Azerbaijan, Georgia, Iran (*Ardabil, Gilan, Golestan, Mazandaran*), Russia (*Dagestan*), Turkmenistan, Turkey (?)

Introduced: Iran (*Alborz, Teheran*), Turkmenistan

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 143,671 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 488,586 sq. km

Size (Max SCL): male 14.1 cm, female 16.6 cm (Joos et al. 2017)

Synonymy:

Emys europaea persica Eichwald 1831:196 (17:33)
Emys europaea persicae, Emys orbicularis persica
Type locality: "provincia Masanderan" [Mazandaran Province, Caspian Sea, Iran] [yellow star, location estimated].
Type specimen: Not known or located, not figured.
Etymology:* Toponym for Persia [= Iran].

Emys europaea iberica Eichwald 1831:196 (09:19, 17:33)
Emys europaea ibericae, Emys orbicularis iberica
Type locality: Not designated. Restricted to "in Iberiae convallibus paludosis et fluviis, Cyrum amnem petentibus" [in marshy Iberian valleys and rivers, Kura River creeks] [Georgia and Azerbaijan] [green star, location estimated] by Eichwald (1840:47).
Type specimen: Not known or located, not figured.
Etymology:* Toponym for Iberia, a former ancient kingdom in Georgia in the Caucasus, not the Iberian Peninsula (= Spain).

Emys orbicularis orientalis Fritz 1994:72
Type locality: "Bandar-e-Anzali (Enzeli), Prov. Gilan, Iran" [blue star].
Type specimen: NHMUK 1874.11.23.8, holotype.
Etymology:* Latin, *oriens*, the Orient, the East; referring to the easternmost range of this subspecies.

Emys orbicularis kurae Fritz 1994:78 (09:19)
Type locality: "Bank (Bank Promisl) an der Kura-Mündung, Aserbaidshan" [Azerbaijan] [pink star].
Type specimen: NMW 14557:1, holotype, see Gemel et al. (2019).
Etymology:* Toponym for the Kura River, Azerbaijan

***Emys orbicularis* ssp. indet.** (14:26)

Testudo purgottii † Ceselli 1846:24 (*nomen oblitum*) (14:26)
Testudo purgotii
Type locality: "Viterbo...Viterbesi...acque Caje" [Italy].
Type specimen: Not located, fossil, partial carapace, figured (pl.1).
Geological age: Late Pleistocene.
Etymology:* Eponym honoring Sebastiano Purgotti (1799–1879), Italy; chemist, mathematician, and philosopher at the University of Perugia.

Emys major † Portis 1890:16 (*nomen dubium*) (14:26)
Emys maior
Type locality: "Poderaccio sotto Persignano nella Valle Superiore dell'Arno" [Italy].
Type specimen: Possibly IGF, fossil, not figured.
Geological age: Late Pliocene to Early Pleistocene, Villafranchian.
Etymology:* Latin, *major*, greater or larger, referring to the size.

Emys latens † Portis 1890:16 (*nomen dubium*) (14:26)
Type locality: "Colombajolo presso S. Giovanni" [Italy].
Type specimen: Possibly IGF, fossil, not figured.
Geological age: Late Pliocene to Early Pleistocene, Villafranchian.
Etymology:* Latin, *latens*, invisible, hidden; referring to the uncertain affinities of this taxon when described.


Emys tigris Salvator 1897:280 (*nomen nudum*)
Type locality: "Mercadal" [Menorca, Balearic Islands, Spain].
Type specimen: Not designated.
Etymology:* Latin, *tigris*, tiger; presumably referring to the carapacial pattern of radial fine lines.

Emys trinacris Fritz, Fattizzo, Guicking, Tripepi, Pennisi, Lenk, Joger, and Wink 2005 (17:34, 21:62)

Sicilian Pond Turtle, *Testuggine Palustre Siciliana*




Melita Vamberger / CBFTT / Laghetto Gorgo, Sicily, Italy / male





left: Dario Ottonello / CBFTT / Mazara del Vallo, Sicily, Italy
right: Stefania D'Angelo / CBFTT / Gallitello Calatafimi, Sicily, Italy



** (orange dot = probably introduced; star = type locality)

Distribution: Italy (*Sicily*)

Introduced: Italy (*Continental*), Germany

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 23,269 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 30,129 sq. km

Size (Max SCL): male 15.6 cm, female 17.2 cm (Ottonello et al. 2021 CBFTT)

CBFTT Account: Ottonello, D'Angelo, Marrone, Oneto, Spadola, Zuffi, and Fritz (2021)

IUCN Red List:* Data Deficient (DD) (Luiselli, Capula, Di Vittorio, Marrone, and Vamberger 2024). Previously: TFTSG Provisional Red List: Least Concern (LC) (2018); Data Deficient (DD) (van Dijk 2009)

Synonymy:

Emys trinacris Fritz, Fattizzo, Guicking, Tripepi, Pennisi, Lenk, Joger, and Wink 2005a:364

Emys orbicularis trinacris

Type locality: "Lago Gian Fenaro, below the pass of Pizzo Lamiaria approximately 1400 m above sea level, Monte Nebrodi, Sicily" [Italy]. Emended to "Laghetto Gianferraro...14.497241 E, 37.951625 N; Elevation: 1007 m a.s.l." by Marrone et al. (2016:60) [yellow star].

Type specimen: MZUF 11136, holotype.

Etymology:* Toponym for Trinacria [= triangular island], ancient Greek name for Sicily, still used in Italy for Sicily.

Glyptemys Agassiz 1857a (07:21)

Sculpted Turtles

(includes 2 species, 2 taxa)

Glyptemys Agassiz 1857a:252,443

Type species: *Glyptemys inculpta* [= *Testudo insculpta* Le Conte 1830], by original designation.

Etymology:* Greek, *glypto*, to carve, to engrave, and *emys*, turtle; referring to the structured carved-like surface of the carapace scutes.

Calemys Agassiz 1857a:252,443

Type species: *Calemys muhlenbergii* [= *Testudo muhlenbergii* Schoepff 1801], by original designation.

Etymology:* Greek, *kalamos*, reed bed or reeds, and *emys*, freshwater turtle; referring to its preferred habitat.

Glyptemys insculpta (Le Conte 1830) (21:63)


Wood Turtle



Anders G.J. Rhodin / Lunenburg, Worcester Co., Massachusetts



left and center: Anders G.J. Rhodin / Shaftsbury, Bennington Co., Vermont / female
right: Craig B. Stanford / Sussex Co., New Jersey



** (orange dots = questionable, probably translocated; stars = type localities)

Distribution: Canada (*New Brunswick, Nova Scotia, Ontario, Québec*), USA (*Connecticut, Delaware, Iowa, Maine, Maryland, Massachusetts, Michigan, Minnesota, New Hampshire, New Jersey, New York, Pennsylvania, Rhode Island, Vermont, Virginia, West Virginia, Wisconsin*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 793,155 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 1,922,698 sq. km

Size (Max SCL): male 24.4 cm, female 22.5 cm (Lovich et al. 1990; Brooks et al. 1992)

IUCN Red List: Endangered (EN A2cd+4c) (van Dijk and Harding 2011). Previously: Vulnerable (VU) (TFTSG *in* Baillie and Groombridge 1996)

CITES: Appendix II (1992)

Synonymy:

Emys pulchella Schweigger 1812:303 (junior homonym, not = *Testudo pulchella* Schoepff 1801 [= *Emys pulchella*] [= *Emys*

orbicularis orbicularis])
Geoclemys pulchella, Glyptemys pulchella
Type locality:* Unknown.
Type specimens:* Originally MNHN, syntypes (3), possibly lost.
Etymology:* Latin, *pulchra*, beautiful, pretty, and *-ella*, small or smaller; referring to the attractive shell and extremities.

Testudo insculpta Le Conte 1830:112
Clemmys (Clemmys) insculpta, Clemmys insculpta, Emys insculpta, Glyptemys insculpta, Chelopus insculptus, Calemys insculpta
Type locality: "the northern states" [USA]. Restricted to "vicinity of New York City" [New York, USA] [yellow star] by Schmidt (1953:92).
Type specimen: MNHN 9452 (shell) and 6955 (body parts of the same specimen), lectotype, designated by Bour (2003:544).
Etymology:* Latin, *insculptus*, etched or carved; referring to the sculpturing of the carapacial scutes.

Emys speciosa Gray 1830e:10 (10:7)
Type locality: "North America?" Restricted to "America Boreali, New Jersey" [USA] [green star, location estimated] by Gray (1831d:26).
Type specimens: OUM 8489–91, syntypes (3), discussed by Nowak-Kemp and Fritz (2010).
Etymology:* Latin, *speciosa*, beautiful, well formed; referring to the attractive shell and colored extremities.

Emys inscripta Gray 1831d:26 (*nomen novum*)
Etymology:* Latin, *inscripta*, inscribed; referring to the sculpturing of the carapacial scutes.
Comment: Unjustified emendation or error for *insculpta*.

Emys speciosa levigata Gray 1831d:26
Type locality: "America Boreali, New Jersey" [USA] [green star, location estimated].
Type specimen: OUM 8489, holotype, discussed by Nowak-Kemp and Fritz (2010).
Etymology:* Latin, *levigata*, smoothed; referring to this taxon having smoother carapace scutes in contrast to other varieties of *Emys speciosa*.


Glyptemys muhlenbergii (Schoepff 1801) (56)*
Bog Turtle



Brian Zarate / TCC / Sussex Co., New Jersey



left: Peter Paul van Dijk / Maryland
right: Brian Zarate / Sussex Co., New Jersey



** (light red = extirpated; red dots = extirpated, orange dots = questionable historical records; stars = type localities)

Distribution: USA (*Connecticut, Delaware, Georgia, Maryland, Massachusetts, New Jersey, New York, North Carolina, Pennsylvania, South Carolina, Tennessee, Virginia*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 98,998 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 502,790 sq. km
Size (Max SCL): male 11.5 cm, female 11.0 cm (Klemens 1993; Ernst and Lovich 2009; M. Knoerr, unpubl. data)
IUCN Red List: Critically Endangered (CR A2cd+4ce) (van Dijk 2011). Previously: Endangered (EN) (TFTSG *in* Baillie and Groombridge 1996)
TFTSG Provisional Red List:* Critically Endangered (CR) (Munscher, Erb, Knoerr, Pignatelli, Zappalorti, Zarate, and Walde, in prep., 2025)
CITES: Appendix I (1992). Previously: Appendix II (1975)
USA ESA:* Threatened, wherever found, except Georgia, North Carolina, South Carolina, Tennessee, Virginia, or Threatened (by Similarity of Appearance), in Georgia, North Carolina, South Carolina, Tennessee, Virginia (USFWS 1997); Species Status Assessment: USFWS (2022c)
Synonymy:

Testudo muhlenbergii Schoepff 1801:132
Testudo mühlenbergii, Emys muhlenbergii, Emys mühlenbergii, Chersine muhlenbergii, Chersine mühlenbergii, Terrapene muhlenbergii, Clemmys (Clemmys) muhlenbergii, Clemmys muhlenbergii, Geoclemys muhlenbergii, Calemys mühlenbergii, Calemys muhlenbergii, Geoclemmys muhlenbergii, Chelopus muhlenbergii, Melanemys muhlenbergii, Glyptemys muhlenbergii, Glyptemys muehlenbergii
Type locality: "Pensylvaniae rivulis" [USA]. Restricted to "Lancaster, Pennsylvania" [USA] [yellow star] by Stejneger and Barbour (1917:114).
Type specimen: Not located, type specimen figured (pl.31).
Etymology:* Eponym honoring Gotthilf H.E. Mühlenberg (1753–1815), Germany / USA; clergyman and botanist who discovered the species.

Emys biguttata Say 1825:212 (10:16)
Type locality: "United States." Restricted to "vicinity of Philadelphia" [Pennsylvania, USA] [green star] by Schmidt (1953:92).
Type specimen: Originally ANSP, apparently lost, not listed by Malnate (1971).
Etymology:* Latin, *bi-*, two, and *guttata*, spotted; referring to the two prominent head spots.

Emys bipunctata Say *in* Gray 1830e:10 (*nomen novum*)
Etymology:* Latin, *bi-*, two, and *punctata*, spotted or punctured; referring to the two prominent head spots.
Comment: Unjustified emendation or error for *biguttata*.

Emys fusca LeSueur *in* Gray 1831d:25 (*nomen nudum*)

Type locality: Not indicated. Restricted to "Philadelphie" [Philadelphia, Pennsylvania, USA] [yellow star] by Bonnemains and Bour (1996).
Type specimen: MNHN 1502, see Bonnemains and Bour (1996).
Etymology:* Latin, *fusca*, dark brown, dark, gray, black; referring to the carapace color.

Geoclemmys mühlenburgii Gray 1870c:27 (*nomen novum*)
Calemys mühlenburgii
Etymology:* Eponym honoring Gotthilf H.E. Mühlenberg (1753–1815), Germany / USA; clergyman and botanist who discovered the species.
Comment: Unjustified emendation or error for *Geoclemmys mühlenbergii*.

Clemmys nuchalis Dunn 1917:624
Type locality: "side of Yonahlossee Road, about 3 miles from Linville, North Carolina...altitude, 4200 feet" [USA] [blue star].
Type specimen: AMNH 8430, holotype, see Reynolds et al. (2007).
Etymology:* Latin, *nucha*, neck, spinal cord, *nuchalis*, belonging to the neck; referring to the relatively longer nuchal scute in this taxon.

Glyptemys muehlenbergii Frétey 2024:12,85 (*nomen novum*) (56)*
Etymology:* Eponym honoring Gotthilf H.E. Mühlenberg (1753–1815), Germany / USA; clergyman and botanist who discovered the species.
Comment:* Emendation of *Glyptemys muhlenbergii*, here declared unjustified in our annotation.

Terrapene Merrem 1820 (14:27, 21:64)

American Box Turtles
(includes 7 species, 10 taxa)

Didicla Rafinesque 1815:75 (*nomen nudum*)
Etymology:* Greek, *di*, two, and Latin, *claudere*, *clida*, to close, rendered as *dicla*, an anagram of *clida*; referring to the two plastral lobes that close.

Terrapene Merrem 1820:27
Type species: *Terrapene clausa* [= *Testudo clausa* Gmelin 1789 = subjective synonym of *Testudo carolina* Linnaeus 1758], by subsequent designation by Bell (1828c:514).
Etymology:* Vernacular (Native American, Powhatan), *torope*, *terrapin*, *terrapen*, or *turpin*, indigenous general name for a turtle, see Dodd et al. (2016).

Cistuda Fleming 1822:270
Type species: "Box tortoise", by original designation.
Etymology:* Latin, *cista*, box, basket, and *testudo*, turtle; referring to the ability to close the plastral lobes to completely close the shell openings, like a box.

Didicla Rafinesque 1832:64
Type species: *Didicla clausa* [= *Testudo clausa* Gmelin 1789 = subjective synonym of *Testudo carolina* Linnaeus 1758], by original designation.
Etymology:* Greek, *di*, two, and Latin, *claudere*, *clida*, to close, rendered as *dicla*, an anagram of *clida*; referring to the two plastral lobes that close.

Cistudo Duméril and Bibron 1835:207 (*nomen novum*) (10:17)
Etymology:* Latin, *cista*, box, basket, and *testudo*, turtle; referring to the ability to close the plastral lobes to completely close the shell openings, like a box.
Comment:* Unjustified emendation of *Cistuda*.

Pyxidemys Fitzinger 1835:123
Type species: *Pyxidemys clausa* [= *Testudo clausa* Gmelin 1789 = subjective synonym of *Testudo carolina* Linnaeus 1758], by subsequent designation by Fitzinger (1843:29).
Etymology:* Greek, *pyxis*, box, basket, and *emys*, turtle; referring to the ability to close the plastral lobes to completely close the shell openings, like a box.

Emyoides Gray 1844:27
Type species: *Emys* (*Emyoides*) *kinosternoides* [= *Emys kinosternoides* Gray 1830e = subjective synonym of *Testudo carolina* Linnaeus 1758], by original monotypy.
Etymology:* Greek, *emys*, turtle, and *-oides*, resembling; referring to the similarity between this genus and *Emys*.


Onychotria Gray 1849:17
Type species: *Cistudo* (*Onychotria*) *mexicana* Gray 1849, by original monotypy.
Etymology:* Greek, *onyx*, claw, nail, and *tria*, three; referring to the three claws on each of the hind feet.

Pariemys Cope 1895:757
Type species: *Pariemys bauri* [= *Terrapene bauri* Taylor 1895], by original monotypy.
Etymology:* Latin, *paries*, wall, and Greek, *emys*, turtle; referring to the presence of a complete zygomatic arch in the skull.

Toxaspis Cope 1895:757
Type species: *Toxaspis major* [= *Cistudo major* Agassiz 1857a], by original monotypy.
Etymology:* Greek, *toxon*, arch, arrow, and *aspis*, shield; referring to the presence of a complete zygomatic arch in the skull.

Cistudos Herrera 1901:36 (*nomen novum et suppressum*)
Etymology:* Latin, *cista*, box, basket, and *testudo*, turtle; referring to the ability to close the plastral lobes to completely close the shell openings, like a box.
Comment: Unjustified replacement name for *Cistudo*. Name suppressed by ICZN (1922).

Terrapene carolina (Linnaeus 1758) (11:8, 14:27, 17:37, 21:64)
American Box Turtle
(includes 3 subspecies)



** (subspecies: *carolina* = red [light red = extirpated], *bauri* = blue, *major* = green, overlap = intergrades, red dots = extirpated; orange dots = questionable *carolina*, probably translocated or misidentified)

Distribution:* Canada (*Ontario [extirpated]*), USA (*Alabama, Connecticut, Delaware, Florida, Georgia, Illinois, Indiana, Kentucky, Louisiana, Maine, Maryland, Massachusetts, Michigan, Mississippi, New Hampshire, New Jersey, New York, North Carolina, Ohio, Pennsylvania, Rhode Island, South Carolina, Tennessee, Virginia, West Virginia*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 1,632,524 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 3,320,093 sq. km
Size (Max SCL): male 23.5 cm, female 19.8 cm (see subspp.)
CBFTT Account: Kiester and Willey (2015)
IUCN Red List: Vulnerable (VU A2bcde+4bcde) (van Dijk 2011). Previously: Near Threatened (NT) (TFTSG *in* Baillie and Groombridge 1996)
CITES: Appendix II, as *Terrapene* spp. (1995)

Terrapene carolina carolina (Linnaeus 1758) (11:8, 14:27, 17:37, 17:38, 21:64, 21:65)

Eastern Box Turtle, Woodland Box Turtle




Peter Paul van Dijk / CBFTT / Colesville, Montgomery Co., Maryland





left: Michael T. Jones / CBFTT / Massachusetts / male
right: Peter Paul van Dijk / CBFTT / Maryland / male



** (subspecies: *carolina* = red [light red = extirpated], *bauri* = blue, *major* = green; overlap = intergrades, red dots = extirpated; orange dots = questionable *carolina*, probably translocated or misidentified; stars = type localities)

Distribution: Canada (*Ontario [extirpated]*), USA (*Alabama, Connecticut, Delaware, Florida, Georgia, Illinois, Indiana, Kentucky, Maine, Maryland, Massachusetts, Michigan, Mississippi, New Hampshire, New Jersey, New York, North Carolina, Ohio, Pennsylvania, Rhode Island, South Carolina, Tennessee, Virginia, West Virginia*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 1,424,405 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 2,087,572 sq. km

Size (Max SCL): male 17.4 cm, female 19.8 cm (Cook et al. 1972; Klemens 1993; Kiester and Willey 2015 CBFTT)

CITES: Appendix II, as *Terrapene* spp. (1995)

Synonymy:

Testudo carolina Linnaeus 1758:198 (senior homonym, not = *Testudo carolina* Le Conte 1830 [= *Gopherus polyphemus*]) (21:65)

Terrapene carolina, Testudo carolinina, Terrapene carolinensis, Emys (Cistuda) carolinae, Terrapene carolinae, Cistuda carolina, Cistudo carolina, Cistudo carolinensis, Terrapene carolina carolina

Type locality: "Carolina" [USA]. Restricted to "vicinity of Charleston, South Carolina" [USA] [yellow star] by Schmidt (1953:93).

Type specimen: Specimen figured in Edwards (1751:pl.205), holotype; Uetz et al. (2019) erroneously listed MCZ 184564 as a syntype, but that specimen is a syntype of *Cistudo virginea* Agassiz 1857 (= *Terrapene carolina carolina*).

Etymology:* Toponym for the Carolinas [North Carolina and South Carolina], USA.

Comment: Description cited as sourced from Edwards (1751:pl.205).

Testudo carinata Linnaeus 1758:198 (senior homonym and *nomen dubium*, not = *Testudo carinata* Thunberg 1810 [= *Geoemyda spengleri*]) (21:66)

Terrapene carinata, Cistudo carinata

Type locality: "Calidis regionibus." Restricted to "vicinity of Charleston, South Carolina" [USA] [yellow star] by Schmidt (1953:93).

Type specimen: Not known or designated, possibly originally UUZM or NRM, apparently lost, not listed by Linné and Thunberg (1780), nor located by Schoepff (1792:xi); UPSZTY 277 (formerly UUZM 277 and MGA 46), listed by Holm (1957) and Wallin (2001) as the possible type of *T. carinata* Linnaeus 1758, is not the type—it is instead the unpublished type of *T. carinata* Thunberg 1810 (*nomen nudum*) and synonym of *Testudo spengleri* Gmelin 1789 (= *Geoemyda spengleri*).

Etymology:* Latin, *carina*, the keel, *carinata*, keeled; referring to the carapacial vertebral keel.

Comment: Description not based on any cited references or designated specimens.

Testudo brevicaudata Lacepède 1788:169, synopsis[table] (*nomen suppressum*) (09:6)

Testudo brevi-caudata, Testudo brevicauda

Type locality: "Caroline" [South Carolina, USA] [yellow star, location estimated].

Type specimen: Possibly MNHN, not located.

Etymology:* Latin, *brevis*, short, and *caudatus*, tailed; referring to the short tail.

Comment: Name suppressed by ICZN (2005a) as published in a rejected and invalid non-binomial work, see Savage (2003).

Testudo incarcerata Bonnaterre 1789:29

Type locality: "Philadelphie...L'Amérique septentrionale" [Pennsylvania, USA] [green star].

Type specimen: Possibly MNHN, not located.

Etymology:* Latin, *incarcerata*, incarcerated, imprisoned; referring to the ability to close the plastral lobes to completely close the shell openings, like a prison.

Testudo incarceratostriata Bonnaterre 1789:29

Testudo incarcerato-striata

Type locality: "L'Amérique septentrionale." Restricted to "vicinity of Philadelphia" [Pennsylvania, USA] [green star] by Schmidt (1953:93).

Type specimen: Possibly MNHN, not located.

Etymology:* Latin, *incarcerata*, incarcerated, imprisoned, and *striata*, grooved; referring to the ability to close the plastral lobes to completely close the shell openings, like a prison, and the shell is grooved.

Testudo clausa Gmelin 1789:1042

Emydes clausa, Emys clausa, Didicla clausa, Terrapene clausa, Cistudo clausa, Cinosternon clausum, Emys (Pyxidemys) clausa, Pyxidemys clausa, Cinosternum clausum

Type locality: "America septentrionali." Restricted to "vicinity of Philadelphia" [Pennsylvania, USA] [green star] by Schmidt (1953:94).

Type specimen: Not located, type specimen figured in Bloch (1786:pl.1).

Etymology:* Latin, *clausa*, closed; referring to the ability to close the plastral lobes to completely close the shell openings, like a box.

Comment: Description cited as sourced from Bloch (1786:131).

Testudo virgulata Latreille *in* Sonnini and Latreille 1801:100

Emys virgulata, Terrapene virgulata

Type locality: "les grands bois de la Caroline" [USA]. Restricted to "Charleston, South Carolina" [USA] [yellow star] by Schmidt (1953:94).

Type specimen: Possibly MNHN, apparently lost, type specimen figured (pl.opp.100,f.1).

Etymology:* Latin, *virgula*, stick, thin branch, *virgulata*, having small sticks; referring to the pattern of irregularly arranged light lines, rods, spots or dots on the carapace.

Emys schneideri Schweigger 1812:317

Emys (Pyxidemys) schneideri

Type locality: Not known. Restricted to "vicinity of Philadelphia" [Pennsylvania, USA] [green star] by Schmidt (1953:94).

Type specimens: Originally MNHN and MZUS, syntypes (2), not located.

Etymology:* Eponym honoring Johann G. Schneider (1750–1822), Germany; classicist, naturalist, and turtle taxonomist.

Monoclida kentukensis Rafinesque 1822:5 (*nomen suppressum*)

Monoclida kentuckensis

Type locality: "United States...Kentucky" [USA] [orange star, location estimated]

Type specimen: Not known or located.

Etymology:* Toponym for Kentucky, USA.

Comment: Name suppressed by ICZN (1984) as published in a rejected and invalid work, see Smith et al. (1980a).

Didicla erythrops Rafinesque 1822:5 (*nomen nudum*)

Didicula erythrope

Type locality: "United States" [USA]

Type specimen: Not known or located.

Etymology:* Greek, *erythros*, red, and *ops*, eye; referring to the red iris of male box turtles.

Terrapene maculata Bell 1825a:309

Terrapene carolina maculata

Type locality: Not known. Restricted to "vicinity of Philadelphia" [Pennsylvania, USA] [green star] by Schmidt (1953:94).

Type specimen: OUM 8503, holotype, discussed by Nowak-Kemp and Fritz (2010).

Etymology:* Latin, *macula*, spot, *maculata*, having spots; referring to the spots on the shell.

Terrapene nebulosa Bell 1825a:310

Terrapene carolina nebulosa

Type locality: Not known. Restricted to "vicinity of Philadelphia" [Pennsylvania, USA] [green star] by Schmidt (1953:94).

Type specimen: OUM 8506, holotype, discussed by Nowak-Kemp and Fritz (2010).

Etymology:* Latin, *nebula*, fog, haze, *nebulosa*, foggy, cloudy, nebulous; referring to the colors on the shell.

Testudo irregulata Daudin *in* Gray 1830e:7 (*nomen novum*) (10:7)

Etymology:* Latin, *in, ir*, not, and *regula*, ledge, bar, rail, *irregulata*, not having a ledge; referring to the low keel on the carapace.

Comment: Unjustified emendation or error for *Testudo virgulata* Bosc *in* Daudin (1801:204).

Emys (Cistuda) carolinae fusca Gray 1830e:7 (10:7)

Emys carolinae fusca, Cistuda carolinae fusca

Type locality: "North America."

Type specimen: Possibly OUM, not located.

Etymology:* Latin, *fusca*, dark brown, dark, gray, black; referring to the dark carapace color.

Emys kinosternoides Gray 1830e:12 (10:7)

Emys (Emyoides) kinosternoides, Terrapene kinosternoides

Type locality: Not known. Restricted to "vicinity of Philadelphia" [Pennsylvania, USA] [green star] by Schmidt (1953:94).

Type specimen: Originally RCSM, holotype, not located.

Etymology:* Greek, *kino*, to move, and *sternon*, chest, sternum, and *-oides*, similarity; referring to the similarity to the species of the genus *Kinosternon* because of its plastron hinge.

Emys dubia Schweigger *in* Gray 1831d:18 (*nomen nudum*) (10:7)

Etymology:* Latin, *dubia*, doubtful, uncertain; referring to the uncertain identity of this taxon.

Emys cinosternoides Duméril and Bibron 1835:303 (*nomen novum*)

Cistudo carolina cinosternoides, Cistudo cinosternoides, Terrapene cinosternoides

Etymology:* Greek, *kino*, to move, and *sternon*, chest, sternum, and *-oides*, similarity; referring to the similarity to the species of the genus *Kinosternon* because of its plastron hinge.

Comment: Unjustified emendation of *kinosternoides*.

Cistudo pickeringi Holbrook *in* Duméril 1855:199 (*nomen nudum*)

Etymology:* Eponym honoring Charles Pickering (1805–1878), USA; physician and naturalist who collected turtles for Holbrook.

Cistudo virginea Agassiz 1857a:260,445

Type locality: "New England, and westward as far as Michigan, and southward as far as the Carolinas" [USA]. Restricted to "vicinity of Cambridge, Massachusetts" [USA] [brown star] by Schmidt (1953:94) (but invalid restriction); restricted instead to "Springfield, Massachusetts" [USA] [blue star] by Smith and Smith (1980:553).


Type specimens: MCZ 1526–35, 1537–40, 1542–50, 1552, 1554, 1555(1), 1555(2), 1556 (two), 1557, 184563–66, syntypes (34), listed by Barbour and Loveridge (1929), Ernst and McBreen (1991) and MCZ Online database; the restriction of the type locality by Smith and Smith (1980) limits the selection of a potential lectotype to the series 1545–1546 and 1548–50, collected at that locality.

Etymology:* Toponym for Virginia, USA.


Terrapene carolina bauri Taylor 1895 (11:8, 14:27, 17:37, 21:64)
(or *Terrapene bauri*)
Florida Box Turtle



Michael T. Jones / CBFTT / peninsular Florida



left: Michael T. Jones / CBFTT / peninsular Florida / female, male
right: Richard D. Bartlett / CRM 3 / Alachua Co., Florida / hatchling



(subspecies: *carolina* = red, *bauri* = blue, *major* = green, overlap = intergrades; orange dots = questionable *carolina*, probably translocated or misidentified; stars = type localities)

Distribution: USA (*Florida*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 146,860 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 253,327 sq. km

Size (Max SCL): male 19.0 cm, female 15.3 cm (Dodd 1997; Jackson and Brechtel 2006; Kiester and Willey 2015 CBFTT)

CITES: Appendix II, as *Terrapene* spp. (1995)

Synonymy:

Terrapene bauri Taylor 1895:576
Pariemys bauri, Cistudo bauri, Terrapene carolina bauri
Type locality: "Florida" [USA]. Restricted to "Orlando, Florida" [USA] [yellow star] by Schmidt (1953:94).
Type specimen: USNM 8352, holotype, see Cochran (1961) and Reynolds et al. (2007); Uetz et al. (2019) erroneously list MCZ 1508 as syntype, but that specimen is a syntype of *Terrapene carolina major*.
Etymology:* Eponym honoring Georg Baur (1859–1898), Czech Republic / Germany / USA; vertebrate morphologist and herpetologist at Yale University, Clark University, and University of Chicago.

Terrapene formosa † Hay 1916a:57
Type locality: "Ocala, Florida" [USA] [light gray star].
Type specimen: USNM 8825 (formerly FGS 2973), holotype, fossil, posterior carapace, figured (pl.4, f.3), see Vlachos (2018).
Geological age: Late Pleistocene, Ocala limestone.
Etymology:* Latin, *formosa*, beautiful, splendid, well-shaped; referring to the fossil holotype of this taxon as being beautiful.
Comment: Considered a synonym of *carolina* (or *bauri*) by Auffenberg (1958) and Vlachos (2018).

Terrapene innoxia † Hay 1916a:61
Type locality: "Vero, St. Lucie County, Florida" [USA] [dark gray star].
Type specimen: USNM 8824 (formerly FGS 7080), holotype, fossil, complete carapace, figured (pl.6,f.1-4), see Vlachos (2018).
Geological age: Pleistocene.
Etymology:* Latin, *innoxia*, unharmed, harmless, innocent; referring to the shell in the described specimens having no visible bone sutures and very indistinct scute sulci, nearly completely smooth.
Comment: Considered a synonym of *carolina* (or *bauri*) by Auffenberg (1958) and Vlachos (2018).


Terrapene carolina major (Agassiz 1857a) (11:8, 14:27, 17:37, 21:64)
Gulf Coast Box Turtle



Richard D. Bartlett / CRM 3 / Liberty Co., Florida



left: Michael T. Jones / CBFTT / Florida / male
right: Richard D. Bartlett / CRM 3 / Liberty Co., Florida / hatchling



(subspecies: *carolina* = red, *bauri* = blue, *major* = green, overlap = intergrades; orange dots = questionable *carolina*, probably translocated or misidentified; star = type locality)


Distribution:* USA (*Alabama, Florida, Georgia, Louisiana, Mississippi*)


Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 132,076 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 205,038 sq. km

Size (Max SCL): male 23.5 cm, female 18.9 cm (Jackson and Brechtel 2006; Kiester and Willey 2015 CBFTT)
CITES: Appendix II, as *Terrapene* spp. (1995)
Synonymy:
Cistudo major Agassiz 1857a:445
Cistudo carolina major, Terrapene major, Toxaspis major, Terrapene carolina major
Type locality: "Mobile..[&]..Florida. Restricted to "Mobile" [Alabama, USA] [yellow star] by Schmidt (1953:94).
Type specimens: MCZ 1505–10, syntypes (6), listed by Barbour and Loveridge (1929) and Ernst and McBreen (1991), but 1505–08 identified as = *Terrapene carolina bauri* (Rosado, pers. comm. *in* Ernst and McBreen 1991).
Etymology:* Latin, *major*, greater or larger; referring to the larger size of this taxon as compared to other species in the genus.

Terrapene coahuila Schmidt and Owens 1944 (14:27, 21:64) (57)*
Coahuilan Box Turtle, *Tortuga de Cuatro Ciénegas*


Gamaliel Castañeda Gaytán / Cuatro Ciénegas, Coahuila, Mexico


left: Gamaliel Castañeda Gaytán / Cuatro Ciénegas, Coahuila, Mexico
right: Craig B. Stanford / Cuatro Ciénegas, Coahuila, Mexico



(orange dot = uncertain; star = type locality)

Distribution: Mexico (*Coahuila*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 840 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 1,063 sq. km
Size (Max SCL): male 23.0 cm, female 19.9 cm (Howeth and Brown 2011 CBFTT)
CBFTT Account: Howeth and Brown (2011)
IUCN Red List:* Critically Endangered (CR A2bc+4bc; B1ab(i,ii,iii,iv,v)+2ab(i,ii,iii,iv,v)) (Castañeda Gaytán, Stanford, Valenzuela-Ceballos, and Rhodin, in press). Previously: TFTSG Provisional Red List: Critically Endangered (CR) (2020); Endangered (EN A2c+4c, B1ab(i,ii,iii,iv,v)+B2ab(i,ii,iii,iv,v)) (van Dijk, Flores-Villela, and Howeth 2007); Endangered (EN) (TFTSG *in* Baillie and Groombridge 1996)
CITES: Appendix I (1975)
USA ESA:* Endangered (USFWS 1973)
Synonymy:
Terrapene coahuila Schmidt and Owens 1944:101
Terrapene ornata coahuila, Terrapene coahuilae
Type locality: "Cuatro Cienegas, Coahuila" [Mexico] [yellow star, location estimated].
Type specimen: FMNH 41234, holotype, see (Marx 1958); FMNH 55656, a paratype, erroneously listed as holotype by Milstead (1969).
Etymology:* Toponym for Coahuila, Mexico.

Terrapene mexicana (Gray 1849) (07:25, 14:27, 17:37, 21:64)
Mexican Box Turtle, *Tortuga Casquito*



Collette Adams / CBFTT / Mexico


Russell A. Mittermeier / Mexico / captivity / Brownsville Zoo


(orange dots = probable trade; stars = type localities)

Distribution:* Mexico (*Nuevo León [?], Puebla, San Luis Potosi, Tamaulipas, Veracruz*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 76,324 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 102,272 sq. km
Size (Max SCL): male 15.7 cm, female 19.5 cm (Müller 1936; Dodd 2001; Legler and Vogt 2013)
CBFTT Account: Kiester and Willey (2015), as part of *Terrapene carolina*
IUCN Red List:* Vulnerable (VU A2c+4c) (Castañeda Gaytán, García-Velez, Fernández-Badillo, Valencia-Herverth, and Stanford, in press). Previously: Vulnerable (VU A2bcde+4bcde) (van Dijk 2011), as part of *Terrapene carolina*; Near Threatened (NT) (TFTSG 1996)
CITES: Appendix II, as *Terrapene* spp. (1995)
Synonymy:
Cistudo (Onychotria) mexicana Gray 1849:17
Onychotria mexicana, Cistudo mexicana, Cistudo carolina mexicana, Chelopus mexicanus, Terrapene mexicana, Terrapene mexicana mexicana, Terrapene carolina mexicana
Type locality: "Mexico." Restricted to "Tampico, Tamaulipas" [Mexico] [yellow star] by Müller (1936:112).
Type specimens: NHMUK 1947.3.4.4, 1947.3.5.48 (formerly 1848.7.28.29–30) syntypes (2), see Milstead (1969) and Ernst and McBreen (1991); NHMUK 1947.3.4.3 listed as a syntype by Milstead (1969) and Ernst and McBreen (1991).
Etymology:* Toponym for Mexico.
Terrapene goldmani Stejneger 1933:119
Type locality: "Chijol (or Chijoles), southeastern corner of the state of San Luis Potosi, Mexico" [green star].
Type specimen: USNM 46251, holotype, see Cochran (1961) and Reynolds et al. (2007).
Etymology:* Eponym honoring Edward A. Goldman (1873–1946), USA; mammalogist and botanist who collected the type.

Terrapene nelsoni Stejneger 1925 (14:27, 21:64)
Spotted Box Turtle, Sierra Box Turtle, *Caja de Manchas*, *Tortuga de Mochomera*
(includes 2 subspecies)



(subspecies: *nelsoni* = red, *klauberi* = blue)
Distribution: Mexico (*Chihuahua, Jalisco, Nayarit, Sinaloa, Sonora*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 111,936 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 243,918 sq. km
Size (Max SCL): male 15.9 cm, female 14.9 cm (see subspp.)
CBFTT Account: Buskirk and Ponce-Campos (2011)
IUCN Red List:* Near Threatened (NT A2cd+4cd) (Butterfield, Espinosa, Matias-Palafox, Domingúez-Pompa, Monsiváis-Molina, García-Caballero, and Uc-Uc, in press). Previously: TFTSG Provisional Red List: Data Deficient (DD) (2011, 2018); Data Deficient (DD) (TFTSG 1996)
CITES: Appendix II, as *Terrapene* spp. (1995)


Terrapene nelsoni nelsoni Stejneger 1925
Southern Spotted Box Turtle, Southern Sierra Box Turtle



John B. Iverson / CBFTT / Nayarit, Mexico



John B. Iverson / nr. Pedro Pablo, Nayarit, Mexico




(subspecies: *nelsoni* = red; star = type locality)
Distribution: Mexico (*Jalisco, Nayarit, Sinaloa*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 46,468 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 106,938 sq. km
Size (Max SCL): male 14.7 cm, female 14.5 cm (Buskirk and Ponce-Campos 2011 CBFTT; Legler and Vogt 2013)
CITES: Appendix II, as *Terrapene* spp. (1995)
Synonymy:
Terrapene nelsoni Stejneger 1925:463
Terrapene nelsoni nelsoni
Type locality: "Pedro Pablo, Tepic, Mexico; 2500 feet altitude" [yellow star].
Type specimen: USNM 46252, holotype, see Cochran (1961), Reynolds et al. (2007), and Legler and Vogt (2013).

Etymology:* Eponym honoring Edward W. Nelson (1855–1934), USA; naturalist and ethnologist, Chief of the U.S. Biological Survey, who collected the type.


Terrapene nelsoni klauberi Bogert 1943
Northern Spotted Box Turtle, Northern Sierra Box Turtle



Matt Cage / CBFTT / nr. Yecora, Sonora, Mexico



left: Young Cage / CBFTT / nr. Yecora, Sonora, Mexico
right: Michael S. Bogle / CBFTT / Rio Aros, Sonora, Mexico



** (subspecies: *nelsoni* = red, *klauberi* = blue; star = type locality)

Distribution: Mexico (*Chihuahua, Sinaloa, Sonora*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 65,468 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 89,782 sq. km

Size (Max SCL): male 15.9 cm, female 14.9 cm (Bogert and Oliver 1945; Buskirk and Ponce-Campos 2011 CBFTT; Legler and Vogt 2013)

CITES: Appendix II, as *Terrapene* spp. (1995)

Synonymy:

Terrapene klauberi Bogert 1943:2


Terrapene nelsoni klauberi

Type locality: "Rancho Guirocoba, approximately eighteen miles southeast of Alamos, Sonora, Mexico" [yellow star].
Type specimen: AMNH 63751, holotype.
Etymology:* Eponym honoring Laurence M. Klauber (1883–1968), USA; electrical engineer, herpetologist, and authority on rattlesnakes, who inspired the author.


Terrapene ornata (Agassiz 1857a) (12:20, 14:27, 21:64)
Ornate Box Turtle, Western Box Turtle



Jeffrey E. Dawson / Barton Co., Kansas



left: John B. Iverson / Garden Co., Nebraska
right: Craig B. Stanford / Nebraska



** (orange dots = probably translocated; stars = type localities)

Distribution: Mexico (*Chihuahua, Sonora*), USA (*Arizona, Arkansas, Colorado, Illinois, Indiana, Iowa, Kansas, Louisiana, Missouri, Nebraska, New Mexico, Oklahoma, South Dakota, Texas, Wisconsin, Wyoming*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 2,015,079 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 3,109,956 sq. km

Size (Max SCL): male 15.7 cm, female 17.0 cm (Ernst and Lovich 2009; Legler and Vogt 2013)

CBFTT Account:* Hughes, Allender, Bernstein, Iverson, Kolthoff, Martin, Meshaka, and Reed (2024)

IUCN Red List:* Near Threatened (NT A2bcd+4bcd) (Iverson, in press). Previously: Near Threatened (NT) (van Dijk and Hammerson 2011); Near Threatened (NT) (TFTSG *in* Baillie and Groombridge 1996)

CITES: Appendix II, as *Terrapene* spp. (1995)

Synonymy:

Cistudo ornata Agassiz 1857a:392,445

Terrapene ornata, Terrapene ornata ornata, Terrapene carolina ornata

Type locality: "Upper Missouri...and...Iowa" [USA]. Restricted to "Council Bluffs, Pottawatomie County, Iowa, USA" [incorrect] [orange star, partially hidden] by Smith and Taylor (1950a:358, 1950b:36); to "junction of the Platte and Missouri River" [Nebraska, USA] [incorrect] [brown star] by Schmidt (1953:95); and to "Burlington, Des Moines County, Iowa" [USA] [yellow star] by lectotype designation by Smith and Smith (1980:587).
Type specimen: MCZ 1536, lectotype, designated by Smith and Smith (1980:587).
Etymology:* Latin, *ornatus*, ornate, ornamental; referring to the carapacial color pattern.

Terrapene ornata cimarronensis Cragin 1894:37
Type locality: "Red beds country of the Cimarron Basin" [Kansas, USA] [green star, location estimated].
Type specimens: Not known or located.
Etymology:* Toponym for the Cimarron Basin, Colorado.

Terrapene longinsulae † Hay 1908c:166 (12:20)
Terrapene ornata longinsulae
Type locality: "Long Island, Phillips County, Kansas" [USA] [dark gray star].
Type specimen: USNM 5983, holotype, fossil, an almost complete shell, skull, and associated appendicular elements, figured (pl.26) and Joyce et al. (2012:f.3–4), see Milstead (1967), Joyce et al. (2012), and Vlachos (2018).
Geological age: Upper Miocene to possibly Pleistocene.
Etymology:* Toponym for Long Island, Kansas, USA.

Terrapene whitneyi † Hay 1916b:8
Type locality: "Austin, Texas" [USA] [light gray star].
Type specimen: USNM 8617, holotype, fossil, a complete shell, figured (pl.1,f.4-5,pl.2,f.1), see Milstead (1967) and Vlachos (2018).
Geological age: Pleistocene.
Etymology:* Eponym honoring Francis L. Whitney (1878–1962), USA; paleontologist at Texas State University, Austin, who obtained the type.

Terrapene ornata luteola Smith and Ramsey 1952:45 (21:64)
Type locality: "17 miles south of Van Horn, Culberson County, Texas" [USA] [blue star].
Type specimen: TCU 1280, holotype, see Lieb (2023).
Etymology:* Latin, *luteolus*, yellowish; referring to the solid yellow or horn-colored carapace.

Terrapene triunguis (Agassiz 1857a) (11:8, 14:27, 17:37, 21:64)
(or *Terrapene carolina triunguis*)
Three-toed Box Turtle




Michael T. Jones / CBFTT / Missouri





left: John B. Iverson / Logan Co., Oklahoma
right: Elizabeth R. Schwartz / Cole Co., Missouri



** (orange dots = questionable, probably translocated or misidentified; stars = type localities)

Distribution:* USA (*Alabama, Arkansas, Illinois, Iowa, Kansas, Louisiana, Mississippi, Missouri, Oklahoma, Tennessee, Texas*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 1,017,532 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 1,478,267 sq. km

Size (Max SCL): male 14.8 cm, female 15.6 cm (E.R. Schwartz and A.R. Kiester, unpubl. data) [max 17.9 cm (Dodd 2001, citing Conant and Collins 1991)]

CBFTT Account: Kiester and Willey (2015), as part of *Terrapene carolina*

IUCN Red List: Vulnerable (VU A2bcde+4bcde) (van Dijk 2011), as part of *Terrapene carolina*. Previously: Near Threatened (NT) (TFTSG 1996)

CITES: Appendix II, as *Terrapene* spp. (1995)

Synonymy:

Cistudo triunguis Agassiz 1857a:279,445
Cistudo carolina triunguis, Terrapene triunguis, Onychotria triunguis, Terrapene carolina triunguis, Terrapene mexicana triunguis
Type locality:* "Louisiana...Mississippi...New Orleans...Osage River...Georgia" [USA]. Restricted to "New Orleans, Louisiana" [USA] [incorrect, out of range] [orange star] by Schmidt (1953:94), restricted here to "Washington, Adams County, Mississippi, USA" [yellow star], an included locality in the series of syntypes (USNM 22), as discussed by Reynolds et al. (2007:13) and Iverson (2022a:41).
Type specimens:* MCZ 1519 (8 specimens), 1522–25, USNM 22, 7545–46, 86871–72 (formerly MCZ 1519), 131838, 213736, syntypes (19), discussed by Barbour and Loveridge (1929), Cochran (1961), Ernst and McBreen (1991), Reynolds et al. (2007), and Iverson (2022a).
Etymology:* Latin, *tria*, three, and *unguis*, nail, claw; referring to the three claws on each of the hind feet.


Terrapene yucatana (Boulenger 1895b) (07:25, 14:27, 17:37, 21:64)
Yucatan Box Turtle



Michael T. Jones / CBFTT / Yucatán, Mexico



left: Michael T. Jones / CBFTT / Yucatán, Mexico
right: John B. Iverson / Yucatán, Mexico



** (orange dots = probable trade or translocated; star = type locality)

Distribution: Mexico (*Campeche, Quintana Roo, Yucatán*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 61,197 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 99,606 sq. km

Size (Max SCL): male 16.1 cm, female 16.0 cm (Buskirk 1993; Ceballos et al. 2013; M.T. Jones and L.L. Willey, unpubl. data)

CBFTT Account: Kiester and Willey (2015), as part of *Terrapene carolina*

IUCN Red List:* Endangered (EN A2cd+4cd) (Jones, Uc-Uc, Butterfield, Willey, and Díaz-Gamboa, in press). Previously: Vulnerable (VU A2bcde+4bcde) (van Dijk 2011), as part of *Terrapene carolina*; Near Threatened (NT) (TFTSG 1996)

CITES: Appendix II, as *Terrapene* spp. (1995)

Synonymy:

Cistudo yucatana Boulenger 1895b:330
Terrapene yucatana, Terrapene mexicana yucatana, Terrapene carolina yucatana
Type locality: "Mexico...North Yucatan." Restricted to "Chichen Itzá, Yucatán, Mexico" [yellow star] by Smith and Taylor (1950a:351, 1950b:35).
Type specimens: NHMUK 1947.3.5.45–47 (formerly 1894.3.23.2–4), syntypes (3).
Etymology:* Toponym for the Yucatan Peninsula, Mexico.

Family Platysternidae Gray 1869a (07:26)

Asian Big-headed Stream Turtles
(includes 1 genus, 1 species, 3 taxa)
Platysternidae Gray 1869a:208




Platysternidae Species Richness

Platysternon Gray 1831c

Asian Big-headed Turtles
(includes 1 species, 3 taxa)
Platysternon Gray 1831c:106
Type species: *Platysternon megacephalum* Gray 1831c, by original monotypy.
Etymology:* Greek, *platys*, broad, flat, and *sternon*, chest, sternum; referring to the relatively wide, flat plastron.
Platysternum Agassiz 1847:297 (*nomen novum*)
Etymology:* Greek, *platys*, broad, flat, and *sternon*, chest, sternum; referring to the relatively wide, flat plastron.
Comment:* Unjustified emendation of *Platysternon*. Frétey (2024) noted that this publication appeared in 1847, not 1846.

Platysternon megacephalum Gray 1831c (17:39) (58)*
Asian Big-headed Turtle
(includes 3 subspecies)



** (subspecies: *megacephalum* = red, *peguense* = blue, *shiui* = green; taxonomically unspecified lineage of *P. megacephalum* sensu lato = gray; orange dots = probable trade)

Distribution:* Cambodia, China (*Anhui, Fujian, Guangdong, Guangxi, Guizhou, Hainan, Hong Kong, Hubei [?], Hunan, Jiangxi, Yunnan, Zhejian*), Laos, Myanmar, Thailand, Vietnam

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Hist. Indig. Area of Occupancy (AOO):* 1,029,224 sq. km
Est. Hist. Indig. Extent of Occurrence (EOO):* 2,696,573 sq. km

Size (Max SCL): male 25.5 cm, female 21.0 cm (see subspp.)

IUCN Red List: Critically Endangered (CR A2cd) (Fong, Hoang, Li, McCormack, Rao, Timmins, Wang, and van Dijk

2021). Previously: Endangered (EN) (ATTWG 2000); Data Deficient (DD) (TFTSG *in* Baillie and Groombridge 1996)

CITES: Appendix I, as Platysternidae spp. (2013). Previously: Appendix II (2003)


Platysternon megacephalum megacephalum Gray 1831c (07:27, 17:39) (58)*

Chinese Big-headed Turtle




Yik-Hei Sung / Hong Kong, China





left: Torsten Blanck / China / trade
right: Yik-Hei Sung / Hong Kong, China



** (subspecies: *megacephalum* = red, *peguense* = blue, *shiui* = green; taxonomically unspecified lineage of *P. megacephalum* sensu lato = gray; orange dots = probable trade; star = type locality)

Distribution:* China (*Anhui, Fujian, Guangdong, Guangxi, Guizhou, Hong Kong, Hubei [?], Hunan, Jiangxi, Zhejian*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 479,440 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 893,024 sq. km

Size (Max SCL): male 20.1 cm, female 15.3 cm (Ernst et al. 2006b; Itescu et al. 2014; Sung et al. 2014)

IUCN Red List: Critically Endangered (CR A2cd) (Fong et al. 2021). Previously: Endangered (EN) (ATTWG 2000); Data Deficient (DD) (TFTSG 1996)

CITES: Appendix I, as Platysternidae spp. (2013). Previously: Appendix II (2003)

Synonymy:

Platysternon megacephalum Gray 1831c:107
Emys megacephala, Platysternon megacephalus, Platysternon megacephalum megacephalum
Type locality: "Chinâ." Restricted to "S. China" [yellow star, location estimated] by Boulenger (1889:48).
Type specimen: NHMUK 1946.9.7.42, holotype.
Etymology:* Greek, *mega*, large, and *kephale*, head; referring to the large head.

Platysternon megacephalum peguense Gray 1870c (17:39) (58)*

Southeast Asian Big-headed Turtle, Burmese Big-headed Turtle




Peter Paul van Dijk / CRM 2 / TCF / CCB / Phu Luang Wildlife Sanctuary, Loei Prov., Thailand





Peter Paul van Dijk / Phu Luang Wildlife Sanctuary, Loei Prov., Thailand / adult, juvenile



** (subspecies: *megacephalum* = red, *peguense* = blue, *shiui* = green; taxonomically unspecified lineage of *P. megacephalum* sensu lato = gray; orange dots = probable trade; stars = type localities)

Distribution: Cambodia, China (*Yunnan*), Laos, Myanmar, Thailand, Vietnam

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 477,293 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 1,051,949 sq. km

Size (Max SCL): male 25.5 cm, female 21.0 cm (Pipatsawasdikul et al. 2010; Chan-ard et al. 2012)

CITES: Appendix I, as Platysternidae spp. (2013). Previously: Appendix II (2003)

Synonymy:

Platysternon peguense Gray 1870c:70
Platysternon megacephalum peguense
Type locality: "Pegu" [= Bago, Myanmar, out of range] [orange star].
Type specimens: NHMUK 1946.1.22.21–22 (formerly 1868.6.6.2–3), syntypes (2).
Etymology:* Toponym for Pegu, capital of the Mon Kingdom in ancient southern Myanmar.

Platysternon megacephalum vogeli Wermuth 1969:374
Type locality: "Provinz Chiang Mai, Nordwest-Thailand" [green star, location estimated].
Type specimen: SMNS 4573, holotype, see Schlüter and Hallermann (1997).

Etymology:* Eponym honoring Zdeněk Vogel (1913–1986), Czech Republic; writer and amateur herpetologist, who obtained the type.

Platysternon megacephalum tristernalis Schleich and Gruber 1984:68 (17:39)

Type locality: "zwischen Mung Lun und Simao, Ostufer des Mekong-flusses, südliches Yünnan (VR China)" [between Menglun (Menghanzhen) and Simao, Lancang Jiang River, Yunnan] [blue star].

Type specimen: ZSM 317/1980 (not ZSM 319/1980/1 as cited originally), holotype, discussed by Franzen and Glaw (2007).

Etymology:* Latin, *tri-*, three, and Greek, *sternon*, chest; referring to the tripartite supernumerary intergular scute.

Platysternon megacephalum shiui Ernst and McCord 1987 (17:39) (58)*


Vietnamese Big-headed Turtle



Torsten Blanck / Vietnam / trade



Torsten Blanck / Vietnam / trade



** (subspecies: *megacephalum* = red, *peguense* = blue, *shiui* = green; taxonomically unspecified lineage of *P. megacephalum* sensu lato = gray; orange dots = probable trade; star = type locality)

Distribution:* China (*Guangxi*, *Hainan*), Vietnam

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Hist. Indig. Area of Occupancy (AOO):* 47,773 sq. km

Est. Hist. Indig. Extent of Occurrence (EOO):* 124,502 sq. km

Size (Max SCL): male 15.1 cm, female 10.7 cm (Ernst and McCord 1987)

CITES: Appendix I, as Platysternidae spp. (2013). Previously: Appendix II (2003)

Synonymy:

Platysternon megacephalum shiui Ernst and McCord 1987:626

Type locality: "vicinity of Langson, Langson Province, Vietnam (26°50' N, 106°45' E)." GPS coordinates in error, corrected to (21°50' N, 106°45' E) [yellow star] by TTWG (2017:82).

Type specimen: USNM 266160, holotype, see Reynolds et al. (2007).

Etymology:* Eponym honoring Oscar Shiu (1959–), China; animal exporter in Hong Kong who obtained the types.

Family Geoemydidae Theobald 1868a (07:29, 09:20, 12:21) (59)*


Geoemydid Pond and River Turtles

(includes 3 subfamilies, 19 genera, 75 species, 99 taxa)

Geoemydidae Theobald 1868a:vi

Batagurina Gray 1869a:185

Bataguridae Gray 1870c:17



Geoemydidae Species Richness

Subfamily Batagurinae Gray 1869a

Batagurine River Turtles

(includes 8 genera, 20 species, 24 taxa)

Batagurina Gray 1869a:185

Bataguridae Gray 1870c:17

Batagurinae McDowell 1964:239

Batagur Gray 1856b (07:30, 08:9)

River Terrapins

(includes 6 species, 7 taxa)

Tetraonyx Gray 1830e:19 (junior homonym, not = *Tetraonyx* Latreille 1809 [= Coleoptera]) (10:7)

Type species: *Trionyx (Tetraonyx) cuvieri* Gray 1830e [= subjective synonym of *Emys baska* Gray 1830d], by original monotypy.

Etymology:* Greek, *tetra*, four, and *onyx*, claw, nail; referring to the front feet having four claws each.

Tetronyx Lesson 1832:pl.7 (*nomen novum et oblitum*)

Type species: *Tetronyx longicollis* [= *Tetraonyx longicollis* Lesson 1831b = subjective synonym of *Emys baska* Gray 1830d], by original monotypy.

Etymology:* Greek, *tetra*, four, and *onyx*, claw, nail; referring to the front feet having four claws each.

Batagur Gray 1856b:35

Type species: *Batagur baska* [= *Emys baska* Gray 1830d], by subsequent designation by Smith (1931:134).

Etymology:* Anonym, *batagur*, possibly a fabricated name, or possibly a local Vernacular name for these turtles.

Kachuga Gray 1856b:35

Type species: *Batagur (Kachuga) lineata* [= *Emys kachuga* Gray 1831a], by tautonymy. Not *Kachuga trilineata* Gray 1869a [= subjective synonym of *Emys trivittata* Duméril and Bibron 1835], by subsequent erroneous designation by Smith (1931:124).

Etymology:* Vernacular (Hindi), *kachhua*, local indigenous name for a freshwater turtle.

Batagurella Gray 1869a:200

Type species: *Kachuga (Batagurella) peguensis* Gray 1869a [= subjective synonym of *Emys trivittata* Duméril and Bibron 1835], by original monotypy.

Etymology:* Anonym, *batagur*, and *-ella*, small or smaller; possibly a fabricated name, or possibly a local Vernacular name (*batagur*) for these turtles.

Dongoka Gray 1869a:202

Type species: *Dongoka hardwickii* [= *Kachuga (Dongoka) hardwickii* Gray 1869a = subjective synonym of *Emys dhongoka* Gray 1832b], by subsequent designation by Lindholm (1929:278).

Etymology:* Vernacular (Hindi), *dohna*, local indigenous name for the

large river turtles of the genus *Batagur*.

Dhongoka Gray 1870c:57 (*nomen novum*)
Type species: *Dhongoka hardwickii* [= *Kachuga (Dongoka) hardwickii* Gray 1869a = subjective synonym of *Emys dhongoka* Gray 1832b], by subsequent monotypy.
Etymology:* Vernacular (Hindi), *dohna*, local indigenous name for the large river turtles of the genus *Batagur*.
Comment:* Unjustified emendation for *Dongoka*.

Callagur Gray 1870c:53
Type species: *Callagur picta* [= *Batagur picta* Gray 1862b = subjective synonym of *Emys borneoensis* Schlegel and Müller 1845], by original monotypy.
Etymology:* Greek, *kallos*, beauty, and *-agur*, partial name for turtles of the related genus *Batagur*; referring to the beautiful color of the head of males of the species in this genus.

Cantorella Gray 1870c:58
Type species: *Cantorella affinis* [= *Tetraonyx affinis* Cantor 1847], by original monotypy.
Etymology:* Eponym honoring Theodore E. Cantor (1809–1860), Denmark / India; physician and naturalist for the East India Company in Calcutta, who also accompanied British troops in China.

Dhougoka Gray 1873j:52 (*nomen novum*)
Etymology:* Vernacular (Hindi), *dohna*, local indigenous name for the large river turtles of the genus *Batagur*.
Comment:* Probable typographic error for *Dhongoka*.

Cachuga Lydekker 1889:123 (*nomen novum*)
Etymology:* Vernacular (Hindi), *kachhua*, local indigenous name for a freshwater turtle.
Comment:* Unjustified emendation of *Kachuga*.

Batagur affinis (Cantor 1847) (08:9) (60)*
Southern River Terrapin
(includes 2 subspecies)



(subspecies: *affinis* = red, *edwardmolli* = blue)

Distribution: Cambodia, Indonesia (*Sumatra*), Malaysia (*West*), Singapore (*extirpated, reintroduced*), Thailand, Vietnam (*extirpated*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 141,001 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 1,037,543 sq. km

Size (Max SCL): male 50.2 cm, female 62.5 cm (see subspp.)

CBFTT Account: Moll, Platt, Chan, Horne, Platt, Praschag, Chen, and van Dijk (2015)

IUCN Red List: Critically Endangered (CR A2bcd+4bcd) (Horne, Chan, Platt, and Moll 2019). Previously: Critically Endangered (CR A2bcd+4bcd) (Horne, Chan, Platt, and Moll 2016); Critically Endangered (CR), as part of *Batagur baska* (ATTWG 2000); Endangered (EN) as part of *Batagur baska* (TFTSG *in* Baillie and Groombridge 1996).

CITES: Appendix I (1975), originally as part of *B. baska*

Batagur affinis affinis (Cantor 1847) (08:9, 09:21) (60)*
Western Malay River Terrapin



Edward O. Moll / CBFTT / Perak R., Malaysia / female



Edward O. Moll / CBFTT / Perak R., Malaysia / male, hatchlings



(subspecies: *affinis* = red, *edwardmolli* = blue; stars = type localities)

Distribution: Indonesia (*Sumatra*), Malaysia (*West*), Singapore (*extirpated, reintroduced*), Thailand

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 68,241 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 298,357 sq. km

Size (Max SCL): male 50.2 cm, female 54.6 cm (Moll et al. 2015 CBFTT)

CITES: Appendix I (1975), originally as part of *B. baska*

Synonymy:
Tetraonyx affinis Cantor 1847:6
Batagur affinis, Kachuga affinis, Kachuga (Dongoka) affinis, Cantorella affinis, Batagur affinis affinis
Type locality: "sea off Pinang…along the sea-shore of Pinang..[&].. estuaries and rivers on the Peninsula" [Penang, Malaysia (West)] [yellow star].


Type specimen: NHMUK 1947.3.4.31 (formerly 1860.3.19.1450), lectotype, designated by Praschag et al. (2008:66); genotyped by Praschag et al. (2008).
Etymology:* Latin, *affinis*, related or adjacent; referring to its close relationship to *Batagur baska*.

Batagur siebenrocki † Jaekel 1911:76
Type locality: "Pithecanthropus-schichten...Java...Trinil" [Indonesia] [gray star].
Type specimen: ZMB (MNB 199) (formerly Jaekel 1026), holotype, fossil, nearly complete carapace and plastron, figured (pl.14,f.1a-1b), apparently lost or destroyed during World War II (Karl 1987); ZMB (MB R3), neotype, fossil, shell, designated by Karl (1987:38).
Geological age: Pleistocene, Trinil Beds.
Etymology:* Eponym honoring Friedrich Siebenrock (1853–1925), Austria; zoologist, herpetologist, and curator at the Naturhistorisches Museum, Vienna.

Batagur affinis edwardmolli Praschag, Holloway, Georges, Päckert, Hundsdörfer, and Fritz 2009 (09:21) (60)*
Eastern Malay River Terrapin


Eng Heng Chan / CBFTT / TCC / Setiu R., Terengganu, Malaysia / male in breeding color


Edward O. Moll / CBFTT / Terengganu, Malaysia / female, hatchling


(subspecies: *affinis* = red, *edwardmolli* = blue; star = type locality)

Distribution: Cambodia, Malaysia (*West*), Thailand (*extirpated*), Vietnam (*extirpated*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 72,760 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 432,237 sq. km
Size (Max SCL): male 49.6 cm, female 62.5 cm (Moll et al. 2015 CBFTT; E.H. Chen, unpubl. data)
CITES: Appendix I (1975), originally as part of *B. baska*
Synonymy:
Batagur affinis edwardmolli Praschag, Holloway, Georges, Päckert, Hundsdörfer, and Fritz 2009a:64 (09:21)
Type locality: "Sre Ambel River system, Koh Kong Province, Cambodia" [yellow star].
Type specimen: NMW 38903, holotype, see Gemel et al. (2019).
Etymology:* Eponym honoring Edward O. Moll (1939–), USA; herpetologist at Eastern Illinois University and first Chair of the IUCN Freshwater Chelonian Specialist Group, precursor to the IUCN Tortoise and Freshwater Turtle Specialist Group.

Batagur baska (Gray 1830d) (07:31, 08:9, 21:67)
Northern River Terrapin


Peter Praschag / Satkhira Distr., Sundarbans Bangladesh / male in breeding color


left: Rupali Ghosh / TCC 2011, 2018 / Sundarbans Bangladesh / male in breeding color
right: Rick Hudson / India / captivity / Madras Crocodile Bank / female


(orange dots = uncertain or trade; stars = type localities)

Distribution: Bangladesh, India (*Odisha, West Bengal*), Myanmar, Pakistan (?); Thailand (*extirpated?*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 107,265 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 886,629 sq. km
Size (Max SCL): male 49.0 cm, female 60.9 cm (Moll et al. 2009 CBFTT; Platt et al. 2019b)
CBFTT Account: Moll, Platt, Platt, Praschag, and van Dijk (2009)
IUCN Red List: Critically Endangered (CR A2acd+4cd; C1+2a(i)) (Praschag and Singh 2019). Previously: Critically Endangered (CR) (ATTWG 2000); Endangered (EN) (TFTSG 1996)
CITES: Appendix I (1975)

USA ESA:* Endangered (USFWS 1970)

Synonymy:

Emys baska Gray 1830d:pl.75

Testudo baska, Emys batagur baska, Tetronyx baska, Batagur (Batagur) baska, Batagur baska, Tetraonyx baska, Tetraonyx basca, Batagur baska baska

Type locality: "India" [yellow star, location estimated]

Type specimen: NHMUK s/n, holotype, apparently lost, specimen figured (pl.75), see Bour (2009c).

Etymology:* Greek, *baska*, enchanting; presumably referring to the beautiful appearance, or possibly a fabricated name, or possibly a local vernacular name for this species.

Emys batagur Gray 1830e:9 (10:7)

Clemmys (Clemmys) batagur, Tetraonyx batagur, Batagur batagur, Batagur batagur batagur

Type locality: Not designated. Restricted to "India" [yellow star, location estimated] by Gray (1831d:24).

Type specimen: NHMUK, holotype, apparently lost.

Etymology:* Anonym, *batagur*; possibly a fabricated name, or possibly a local Vernacular name for this species.

Trionyx (Tetraonyx) cuvieri Gray 1830e:19 (10:7)

Trionyx cuvierii

Type locality: Not designated. Restricted to "l'Irrawady, fleuve du royaume de Pégu" [Myanmar] [green star, location estimated] by Temminck and Schlegel (1834:43).

Type specimen: MNHN, holotype, not located.

Etymology:* Eponym honoring Baron Georges Cuvier (1769–1832), France; paleontologist, anatomist, and zoologist at the Muséum National d'Histoire Naturelle, Paris.

Tetraonyx longicollis Lesson 1831b:297

Clemmys longicollis, Tetronyx longicollis

Type locality: "le fleuve de l'Irravaddy au Pégou" [Myanmar] [green star, location estimated].

Type specimen: MNHN 7967, holotype, discussed by Bour (2009c).

Etymology:* Latin, *longus*, long, and *collum*, neck; referring to the long neck.

Emys tetraonyx Temminck and Schlegel 1834:43 (*nomen novum*) (10:18)

Etymology:* Greek, *tetra*, four, and *onyx*, claw, nail; referring to the front feet having four claws each.

Comment: Unjustified replacement name for *longicollis*.

Tetraonyx lessonii Duméril and Bibron 1835:338 (*nomen novum*)

Hydraspis (Tetronyx) lessonii

Etymology:* Eponym honoring René Primevère Lesson (1794–1849), France; surgeon, naturalist, ornithologist, and herpetologist.

Comment: Unjustified replacement name for *longicollis*.

Batagur baska ranongensis Nutaphand 1979:181 (07:31)

Batagur ranongensis, Batagur batagur ranongensis

Type locality: "mouth of rivers in Ranong Province" [Thailand] [blue star, location estimated].

Type specimen: Not located, type specimen figured (f.75-76).

Etymology:* Toponym for Ranong Province, Thailand.

Tetronyx unicolor Bibron *in* Bour 2009b:31 (*nomen nudum*)

Etymology:* Latin, *unum*, *uni*, one, single, and *color*, color; probably referring to the specimen examined having a uniform color.

Batagur borneoensis (Schlegel and Müller 1845) (07:30)


Painted Terrapin



Douglas B. Hendrie / TCC / Perak, Malaysia / male in breeding color



left: Gerald Kuchling / Setiu R., Malaysia / female
right: Joko Guntoro / Aceh, Sumatra, Indonesia / female



(stars = type localities)

Distribution: Brunei, Indonesia (*Kalimantan*, *Sumatra*), Malaysia (*East*, *West*), Thailand

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO): 233,763 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 1,657,107 sq. km

Size (Max SCL): male 45.0 cm, female 76.0 cm (de Rooij 1915; Bonin et al. 2006; Ernst et al. 2006b; Itescu et al. 2014)

IUCN Red List: Critically Endangered (CR A2cd) (Shepherd, Horne, Guntoro, and Cota 2021). Previously: Critically Endangered (CR) (ATTWG 2000); Critically Endangered (CR) (TFTSG *in* Baillie and Groombridge 1996)

CITES: Appendix II (1997); zero quota for wild specimens for commercial purposes (2013)

Synonymy:

Emys borneoensis Schlegel and Müller 1845:30

Clemmys borneoensis, Callagur borneoensis, Batagur borneoensis

Type locality: "Borneo" [Kalimantan, Indonesia] [yellow star, location estimated].

Type specimens: RMNH 3296, 6210, syntypes (2), discussed by Hoogmoed et al. (2010); RMNH 6210 listed as "holotype" by Uetz et al. (2019).

Etymology:* Toponym for Borneo, Kalimantan, Indonesia.

Batagur picta Gray 1862b:204

Callagur picta, Tetraonyx pictus, Callagur pictus

Type locality: "Borneo, Sarawak" [East Malaysia] [green star, location estimated].

Type specimen: NHMUK 1947.3.4.19 (formerly 1856.9.19.21),

holotype.
Etymology:* Latin, *picta*, painted; referring to the bright head colors of males of this species.

Clemmys grayi Strauch 1865:88 (*nomen novum*)
Etymology:* Eponym honoring John Edward Gray (1800–1875), United Kingdom; herpetologist, turtle taxonomist, and curator at the British Museum of Natural History, London.
Comment: Unjustified replacement name for *picta*.

Kachuga major Gray 1873c:300
Type locality: "India?" Emended to "India" [in error] by Gray (1873j:51).
Type specimen: NHMUK 1947.3.4.20 (formerly 1863.10.8.11), holotype.
Etymology:* Latin, *major*, greater or larger; referring to the larger size of this species.


Batagur borneensis Hubrecht 1881:47 (*nomen novum*)
Callagur borneensis
Etymology:* Toponym for Borneo, Kalimantan, Indonesia.
Comment: Unjustified emendation or error for *borneoensis*.

Kachuga brookei Bartlett 1895a:29
Type locality: "Borneo" [East Malaysia] [green star, location estimated].
Type specimen: Possibly SMK, holotype, originally live.
Etymology:* Eponym honoring Charles Brooke (1829–1917), United Kingdom / Malaysia; Rajah of Sarawak (now part of Malaysia) and founder of the Sarawak Museum.

Batagur dhongoka (Gray 1832b) (07:30) (59)*
Three-striped Roofed Turtle


Indraneil Das / Bihar, India


Edward O. Moll / India / female


**
(orange dot = questionable; stars = type localities)

Distribution: Bangladesh, India (*Assam, Bihar, Madhya Pradesh, Rajasthan, Uttar Pradesh, West Bengal*), Nepal
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 296,503 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 1,209,803 sq. km
Size (Max SCL): male 26.0 cm, female 48.0 cm (Shrestha 1997; Ceballos et al. 2013)
IUCN Red List: Critically Endangered (CR A2cd+4cd) (Das, Choudhury, Praschag, Ahmed, and Singh 2019). Previously: Endangered (EN) (ATTWG 2000); Near Threatened (NT) (TFTSG *in* Baillie and Groombridge 1996)
CITES: Appendix II (2003)
Synonymy:

Emys dhongoka Gray 1832b:pl.60
Batagur (Kachuga) dhongoka, Batagur dhongoka, Batagur dhougoka, Clemmys dhongoka, Kachuga dhongoka
Type locality: "India" [yellow star, location estimated]. Restricted to "N. India" by Smith (1931:130).
Type specimen: NHMUK, holotype, apparently lost, specimen figured (pl.60).
Etymology:* Vernacular (Hindi), *dohna*, local indigenous name for the large river turtles of the genus *Batagur*.

Emys duvaucelii Duméril and Bibron 1835:334
Batagur duvaucelii, Emys duvancellii
Type locality: "Bengale" [India] [yellow star, location estimated].
Type specimen: MNHN 7987, holotype.
Etymology:* Eponym honoring Alfred Duvaucel (1792–1824), France; naturalist, explorer, and collector for the Muséum National d'Histoire Naturelle, Paris.


Kachuga (Dongoka) hardwickii Gray 1869a:202
Kachuga hardwickii, Dhongoka hardwickii, Dhougoka hardwickii
Type locality: "Nepal" [green star, location estimated].
Type specimen: NHMUK 1947.3.4.59, holotype.
Etymology:* Eponym honoring Thomas Hardwicke (1756–1835), United Kingdom / India; military officer and naturalist who served with the East India Company in India.

Batagur durandi † Lydekker 1885:192 (59)*
Type locality: "Siwalik Hills" [Punjab, India] [gray star, location estimated].
Type specimen:* NHM(P) 39841, holotype, fossil, nearly whole shell, figured (pl.24,f.2), also in Lydekker (1889) and Garbin et al. (2020:f.14).
Geological age: Late Pliocene (Pinjor) to Early Pleistocene (Tatrot).
Etymology:* Eponym honoring Sir Henry Marion Durand (1812–1871), United Kingdom / India; military officer and Lieutenant Governor of Punjab, India, who helped collect the type.
Comment:* Synonymy with *Batagur dhongoka* confirmed by Garbin et al. (2020).


Batagur kachuga (Gray 1831a) (07:30, 21:67) (59)*
Red-crowned Roofed Turtle



Saurav Gawan / Chambal R., Uttar Pradesh, India / male in breeding color



left: Edward O. Moll / Yamuna R., Uttar Pradesh, India / female
right: Edward O. Moll / Chambal R., Uttar Pradesh, India / male in breeding color



** (orange dots = probable trade; stars = type localities)

Distribution: Bangladesh, India (*Bihar, Madhya Pradesh, Punjab, Uttar Pradesh, West Bengal*), Nepal

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 180,161 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 697,885 sq. km

Size (Max SCL): male 29.0 cm, female 56.0 cm (Praschag et al. 2019 IUCN)

IUCN Red List: Critically Endangered (CR A2cd+4cd) (Praschag, Ahmed, Das, and Singh 2019). Previously: Critically Endangered (CR) (ATTWG 2000); Endangered (EN) (TFTSG *in* Baillie and Groombridge 1996)

CITES:* Appendix I (2023). Previously: Appendix II (2003)

Synonymy:

Emys lineata Gray 1830e:9 (*nomen oblitum*) (10:7)
Clemmys (Clemmys) lineata, Batagur (Kachuga) lineata, Batagur lineatus, Kachuga lineata
Type locality: "India" [yellow star, likely near Calcutta].
Type specimen: NHMUK, holotype, apparently lost.
Etymology:* Latin, *linea*, line, stripe, *lineata*, having lines, striped; referring to the line markings on the head and neck of males.
Comment:* Declared a *nomen oblitum* by Rhodin et al. (2008) and TTWG (2021), corroborated and validated by Frétey (2024).

Emys kachuga Gray 1831a:pl.74 (*nomen protectum*)
Batagur kachuga, Kachuga kachuga
Type locality: "India" [yellow star, likely near Calcutta]. Restricted to "N. India" [location imprecise] by Smith (1931:131).
Type specimen: NHMUK s/n, holotype, apparently lost, specimen figured (pl.74).
Etymology:* Vernacular (Hindi), *kachhua*, local indigenous name for a freshwater turtle.
Comment:* Declared and validated as a *nomen protectum* by Frétey (2024).

Batagur ellioti Gray 1862b:264
Batagur elliotti, Clemmys ellioti
Type locality: "Southern India, River Kistna" [Krishna River, Andhra Pradesh, India] [in error] [orange star, location estimated].
Type specimen: NHMUK 1947.3.4.68 (formerly 1855.12.17.15), holotype.
Etymology:* Eponym honoring Sir Walter Elliot (1830–1887), United Kingdom (Scotland) / India; naturalist who served with the East India Company in India.

Kachuga fusca Gray 1870c:56 (*partim*)
Batagur fusca
Type locality:* "India." Restricted to "Pegu..&..Nepaul" [Myanmar and Nepal] [green star, location estimated] by Gray (1873j:51).
Type specimens: NHMUK 1947.3.4.30, 1947.3.4.60 (formerly 1845.1.12.416, 1867.9.28.2), syntypes (2).
Etymology:* Latin, *fusca*, dark brown, dark, gray, black; referring to the carapace color.
Comment:* Syntype 1947.3.4.30 from Nepal = *Batagur kachuga;* syntype 1947.3.4.60 from "Pegu" [Myanmar] = *Batagur trivittata*. Name synonymized *in partim* under *Batagur kachuga* and *Batagur trivittata*.

Batagur bakeri † Lydekker 1885:190 (59)*
Type locality: "Siwalik Hills" [Punjab, India] [gray star, location estimated].
Type specimen:* NHM(P) 39835a, holotype, fossil, nearly whole shell, figured (pl.23,f.2-2a), see Lydekker (1889); see Garbin et al. (2020:f.12).
Geological age: Late Pliocene (Pinjor) to Early Pleistocene (Tatrot).
Etymology:* Eponym honoring Sir William E. Baker (1808–1881), United Kingdom / India; military officer and civil administrator in Sindh, India (now Pakistan), who presented the type.
Comment:* Synonymy with *Batagur lineata* and *B. kachuga* confirmed by Garbin et al. (2020).

Batagur trivittata (Duméril and Bibron 1835) (07:30)
Myanmar Roofed Turtle, Burmese Roofed Turtle




Rick Hudson / CRM 5 / TCC / Myanmar / captivity / Yadanabon Zoo / male in breeding color





left: Brian D. Horne / Myanmar / captivity / male
right: Gerald Kuchling / Myanmar / captivity / female



(stars = type localities)

Distribution: Myanmar

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 107,814 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 335,679 sq. km

Size (Max SCL): male 46.0 cm, female 62.0 cm (Smith 1931; Platt et al. 2019c)

IUCN Red List: Critically Endangered (CR A2bcd; B1ab(i,ii,iii) + 2ab(i,ii,iii); D) (Platt, Horne, and Praschag 2019). Previously: Endangered (EN) (ATTWG 2000); Endangered (EN) (TFTSG *in* Baillie and Groombridge 1996)

CITES: Appendix II (2003); zero quota for wild specimens for commercial purposes (2013)

Synonymy:

Emys trivittata Duméril and Bibron 1835:331
Batagur trivittata, Kachuga trivittata
Type locality: "Bengale" [India or Bangladesh] [in error] [orange star, location estimated].
Type specimens: MNHN 7889, 7892, syntypes (2).
Etymology:* Latin, *tria*, three, and *vitta*, ribbon, band, *trivittata*, having three bands; referring to the three black longitudinal bands on the carapace of males.

Kachuga (Batagurella) peguensis Gray 1869a:200
Kachuga peguensis
Type locality: "India." Restricted to "Pegu" [Myanmar] [green star, location estimated] by Gray (1873j:50).
Type specimen: NHMUK 1947.3.4.86 (formerly 1868.5.11.10), holotype.
Etymology:* Toponym for Pegu (Bago), capital of the Mon Kingdom in ancient southern Myanmar.

Kachuga trilineata Gray 1869a:200
Kachuga (Kachuga) trilineata
Type locality: "India." Restricted to "Pegu" [Myanmar] [green star, location estimated] by Boulenger (1889:56).
Type specimen: NHMUK 1947.3.4.85 (formerly 1867.9.28.4), holotype.
Etymology:* Latin, *tri-*, three, and *linea*, line, stripe, *lineata*, having lines, striped; referring to the three line markings on the carapace of males.

Kachuga fusca Gray 1870c:56 (*partim*)
Batagur fusca
Type locality:* "India." Restricted to "Pegu..&..Nepaul" [Myanmar and Nepal] [green star, location estimated] by Gray (1873j:51).
Type specimens: NHMUK 1947.3.4.30, 1947.3.4.60 (formerly 1845.1.12.416, 1867.9.28.2), syntypes (2).
Etymology:* Latin, *fusca*, dark brown, dark, gray, black; referring to the carapace color.
Comment:* Syntype 1947.3.4.30 from Nepal = *Batagur kachuga;* syntype 1947.3.4.60 from "Pegu" [Myanmar] = *Batagur trivittata*. Name synonymized *in partim* under *Batagur kachuga* and *Batagur trivittata*.

Batagur iravadica Anderson 1879:736
Batagur iravadicus, Clemmys iravadica
Type locality: "Pegu..[&]..Bhamô in Upper Burma..[&].. throughout the Irawady" [Bago and Bhamo, Myanmar] [green star = Bago, blue star = Bhamo].
Type specimens: ZSI 743, syntype (1), others apparently lost, see Das et al. (1998) and Kundu et al. (2018a).
Etymology:* Toponym for the IrrawaddyRiver, Burma [Myanmar].

Geoclemys Gray 1856b

Spotted Pond Turtles

(includes 1 species, 1 taxon)

Geoclemys Gray 1856b:17

Type species: *Geoclemys* - [= *Emys hamiltonii* Gray 1830e], by subsequent designation by Stejneger (1907:496).

Etymology:* Greek, *gaia*, *geo-*, the earth, and *klemmys*, turtle, or *kleptein*, to steal, and *emys*, turtle.

Geoclemmys Cope 1865:186 (*nomen novum*)

Etymology:* Greek, *gaia*, *geo-*, the earth, and *klemmys*, turtle, or *kleptein*, to steal, and *emys*, turtle.

Comment:* Unjustified emendation of *Geoclemys*.

Geoclemys hamiltonii (Gray 1830e) (10:7) (59)*


Spotted Pond Turtle, Black Pond Turtle



Indraneil Das / CBFTT / Kaziranga National Park, Assam, India



left: Chittaranjan Baruah / CBFTT / Assam, India
right: Peter Paul van Dijk / CBFTT / Ganges R., India / captivity / Kukrail Crocodile Center



(stars = type localities)

Distribution: Bangladesh, India (*Assam, Bihar, Jammu, Meghalaya, Punjab, Rajasthan, Uttar Pradesh, West Bengal*), Nepal, Pakistan

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 646,121 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 2,450,116 sq. km

Size (Max SCL): male 39.2 cm, female 40.5 cm (Rashid and Swingland 1997; Das and Bhupathy 2010 CBFTT; Baruah et al. 2012)

CBFTT Account: Das and Bhupathy (2010)

IUCN Red List: Endangered (EN A2cd+4cd) (Praschag, Ahmed, and Singh 2019). Previously: Vulnerable (VU) (ATTWG 2000); Near Threatened (NT) (TFTSG *in* Baillie and Groombridge 1996)

CITES: Appendix I (1975)

USA ESA:* Endangered (USFWS 1976)

Synonymy:

Emys hamiltonii Gray 1830e:9 (10:7)

Clemmys (Clemmys) hamiltonii, Clemmys hamiltonii, Geoclemys hamiltonii, Damonia hamiltonii, Geoclemmys hamiltonii, Geoclemys hamoltoni

Type locality:* "India" [yellow star, likely near Calcutta].

Type specimens: OUM 8477, NHMUK 1947.3.4.41, syntypes (2), 1947.3.4.41 erroneously listed as "holotype" by King and Burke (1989:35) and Uetz et al. (2019), discussed by Nowak-Kemp and Fritz (2010).

Etymology:* Eponym honoring Francis Buchanan-Hamilton (1762–1829), United Kingdom (Scotland) / India / Myanmar; physician and botanist who served with the East India Company in India at Calcutta and Burma (now Myanmar), and collected and figured the type.

Emys guttata Gray 1831b:pl.76 (*nomen novum*)

Type locality:* "India" [yellow star, likely near Calcutta].

Type specimen: NHMUK, possibly 1947.3.4.41, type specimen figured (pl.76), see Smith (1931).

Etymology:* Latin, *guttatus*, spotted or speckled; referring to the yellow spots on the carapace.

Comment:* Unjustified replacement name for *E. hamiltonii*, based on the same figure, see Gray (1831d).

Emys piquotii Lesson 1831a:120

Type locality:* "le Gange" [Ganges River or delta, Bangladesh] [green star, location estimated].

Type specimen: Possibly MNHN 1235, holotype.

Etymology:* Eponym honoring Christophe-Augustin Lamare-Picquot (1785–1873), France; pharmacist, naturalist, and explorer who collected the type.

Emys picquotii Lesson *in* Duméril and Bibron 1835:316 (*nomen novum*)

Etymology:* Eponym honoring Christophe-Augustin Lamare-Picquot (1785–1873), France; pharmacist, naturalist, and explorer who collected the type.

Comment: Unjustified emendation of *piquotii*.

Emys hamiltonoides † Falconer and Cautley *in* Lydekker 1880:21 (*nomen nudum*)

Damonia hamiltonoides

Type locality: "Siwalik Hills" [Punjab, India] [dark gray star, location estimated].

Type specimen: Not known, fossil, shell.

Geological age: Late Pliocene (Pinjor) to Early Pleistocene (Tatrot).

Etymology:* Eponym honoring Francis Buchanan-Hamilton (1762–1829), United Kingdom (Scotland) / India / Myanmar; physician and botanist who served with the East India Company in India and Burma (now Myanmar).

Melanochelys pictus Murray 1884b:107

Type locality: "in the Sind "Doro," in the Kushmore Talooka, Upper Sind" [Pakistan] [blue star].

Type specimen: UKZM s/n, holotype, not located.

Etymology:* Latin, *pictus*, painted; referring to the bright colors of this turtle.

Clemmys palaeindica † Lydekker 1885:178 (59)*

Damonia palaeindica

Type locality: "Siwalik Hills" [Punjab, India] [dark gray star, location estimated].

Type specimens:* NHM(P) 39838, 39840, syntypes, fossil, two nearly complete shells, figured (pl.21,f.1,1a,1b,3,3a), see Lydekker (1889); 39838, lectotype, identified as "holotype" by Garbin et al. (2020:f.22).

Geological age: Late Pliocene (Pinjor) to Early Pleistocene (Tatrot).

Etymology:* Toponym for ancient paleontological India.

Comment:* Synonymy with *Geoclemys hamiltonii* confirmed by Garbin et al. (2020).

Geoclemys sivalensis † Tewari and Badam 1969:555 (59)*

Type locality: "Upper Siwaliks...1 km south-east of Quranwalla (30º46'5", 76º52'50")...6 km northeast of Chandigarh Lake...Panjore, India." [Haryana, India] [light gray star].

Type specimen:* PUM A/665, holotype, fossil, partial anterior carapace, figured (f.2); see Garbin et al. (2020).

Geological age: Lower Pleistocene, Pinjore Stage.

Etymology:* Toponym for the Siwalik Hills, India.
Comment:* Synonymy with *Geoclemys hamiltonii* confirmed by Garbin et al. (2020).

Hardella Gray 1870c


Crowned River Turtles
(includes 1 species, 1 taxon)
Hardella Gray 1870c:58
Type species: *Hardella thurjii* [= *Emys thurjii* Gray 1831d], by subsequent designation as *Hardella thurgi* by Günther (1871:70).
Etymology:* Eponym honoring Thomas Hardwicke (1756–1835), United Kingdom / India; military officer and naturalist who served with the East India Company in India.

Hardella thurjii (Gray 1831d) (07:40) (59)*


Crowned River Turtle, Brahminy River Turtle



Indraneil Das / CBFTT / National Chambal Sanctuary, Madhya Pradesh, India



left: James F. Parham / CBFTT / Khairpur, Sind Province, Pakistan
right: Craig B. Stanford / Sarju R., Uttar Pradesh, India



** (orange dots = probably introduced or trade; stars = type localities)

Distribution: Bangladesh, India (*Assam, Bihar, Madhya Pradesh, Meghalaya, Punjab, Uttar Pradesh, West Bengal*), Nepal, Pakistan

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 534,983 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 2,344,175 sq. km

Size (Max SCL): male 20.5 cm, female 65.3 cm (Rashid and Swingland 1997; Das and Bhupathy 2009a CBFTT; Baruah et al. 2012)

CBFTT Account: Das and Bhupathy (2009a)

IUCN Red List: Endangered (EN A2bcd+4bcd) (Ahmed, Praschag, and Singh 2021). Previously: Vulnerable (VU) (ATTWG 2000); Near Threatened (NT) (TFTSG *in* Baillie and Groombridge 1996)

CITES: Appendix II (2013)

Synonymy:

Emys thuryi Gray 1830e:8 (*nomen oblitum*) (10:7)
Type locality: "Bengal" [Bangladesh or India] [likely near Calcutta, yellow star].
Type specimens: OUM 8433–34, syntypes (2), discussed by Nowak-Kemp and Fritz (2010).
Etymology:* Vernacular (Hindi?), *thurgy*; possibly local name used in India for this species.

Emys thurjii Gray 1831d:22 (*nomen novum*)
Testudo thurjii, Hardella thurjii, Hardella thurjii thurjii
Type locality: "India." Restricted to "Ganges and Brahmaputra systems" [India] [likely near Calcutta, yellow star] as *Hardella thurgii* by McDowell (1964:255).
Type specimens: OUM 8433–34, syntypes (2) by default, discussed by Nowak-Kemp and Fritz (2010).
Etymology:* Vernacular (Hindi?), *thurgy*; possibly local name used in India for this species.
Comment: Accepted emendation of *thuryi*.

Emys thuji Gray 1831e:pl.73 (*nomen novum*)
Etymology:* Vernacular (Hindi?), *thurgy*; possibly local name used in India for this species.
Comment: Unjustified emendation or error for *thurjii*.

Emys flavonigra Lesson 1831a:120
Type locality:* "le Gange" [Ganges River or delta, Bangladesh] [green star, location estimated].
Type specimen: Possibly MNHN, not located.
Etymology:* Latin, *flavo*, yellow, golden yellow, and *nigra*, black, dark-colored; referring to the dark carapace with a yellowish rim in young animals.

Clemmys (Clemmys) thurgii Fitzinger 1835:123 (*nomen novum*)
Clemmys thurgii, Testudo thurgii, Emys thurgii, Emys thurgi, Batagur thurgii, Hardella thurgii, Batagur (Hardella) thurgii
Etymology:* Vernacular (Hindi?), *thurgy*; possibly local name used in India for this species.
Comment: Unjustified emendation or error for *thurjii*.

Kachuga oldhami Gray 1869a:200
Kachuga (Kachuga) oldhami
Type locality: "India." Restricted to "Bengal" [Bangladesh or India] [likely near Calcutta, yellow star] by Boulenger (1889:66).
Type specimen: NHMUK 1947.3.4.73 (formerly 1856.5.6.112), holotype.
Etymology:* Eponym honoring Thomas Oldham (1816–1878), Ireland / India; geologist and naturalist at the Geological Survey of India in Calcutta who collected turtles and mammals for the British Museum. The entry for Oldham in Beolens et al. (2011) is incorrect.

Hardella indi Gray 1870c:58 (07:40)
Hardella thurjii indi
Type locality: "Indus River" [Pakistan] [blue star, location estimated].
Type specimen: NHMUK 1947.3.4.74 (formerly 1869.8.28.1), holotype.
Etymology:* Toponym for the Indus River, Pakistan.

Batagur falconeri † Lydekker 1885:187 (59)*
Hardella falconeri
Type locality: "Siwalik Hills" [Punjab, India] [dark gray star, location estimated].
Type specimen:* NHM(P) 39835, holotype, fossil, shell, figured (pl.23,f.1-1a,pl.24,f.4), see Lydekker (1889); see Garbin et al. (2020:f.19).
Geological age: Late Pliocene (Pinjor) to Early Pleistocene (Tatrot).
Etymology:* Eponym honoring Hugh Falconer (1808–1865), United Kingdom (Scotland) / India; physician, paleontologist, and geologist who worked in India and described several fossil animals, including a giant tortoise.
Comment:* Synonymy with *Hardella thurjii* confirmed by Garbin et al. (2020).

Clemmys watsoni † Lydekker 1886:541 (59)*
Type locality: "Siwaliks of Perim Island, Gulf of Cambay, India" [Gujarat, India] [pink star].

Type specimen:* NHM(P) R.748, holotype, fossil, shell, figured (pl.15), see Lydekker (1889); see Garbin et al. (2020:f.20).
Geological age: Late Pliocene to Early Pleistocene.
Etymology:* Eponym honoring Colonel John W. Watson (1838–1889), United Kingdom / India; military officer in India who obtained the type.
Comment:* Synonymy with *Hardella thurjii* confirmed by Garbin et al. (2020).

Geoemyda pilgrimi † Prasad and Satsangi 1967:536 (59)*
Type locality: "Tatrot beds east of Chakrana (31º32' : 76º41') in the Bilaspur District of Himachal Pradesh" [India] [light gray star].
Type specimen: GSI 18091, holotype, fossil, shell, figured by Das (1994:f.1-2), discussed by Das (1994).
Geological age: Late Pliocene (Pinjor) to Early Pleistocene (Tatrot).
Etymology:* Eponym honoring Henry G.E. Pilgrim (1875–1943), United Kingdom / India; geologist, paleontologist, and superintendent of the Geological Survey of India.
Comment:* Synonymy with *Hardella thurjii* accepted by Garbin et al. (2020).

Malayemys Lindholm 1931 (17:46)

Snail-eating Turtles
(includes 3 species, 3 taxa)

Damonia Gray 1869a:193 (junior homonym, not = *Damonia* Robineau-Desvoidy 1847 [= Diptera])
Type species: *Damonia macrocephala* [= *Geoclemmys macrocephala* Gray 1859], by subsequent designation by Stejneger (1907:496).
Etymology:* Eponym honoring Robert Damon (1814–1889), United Kingdom; naturalist and geologist who traded in natural historical and geological material and corresponded with Gray.

Malayemys Lindholm 1931:30 (*nomen novum*)
Type species: *Malayemys macrocephala* [= *Geoclemmys macrocephala* Gray 1859], in accordance with ICZN Article 67.8, not *Malayemys subtrijuga* [= *Emys subtrijuga* Schlegel and Müller 1845], as by original designation by Lindholm (1931:30).
Etymology:* Toponym for Malaya (Malaysia), and Greek, *emys*, turtle.

Malayemys khoratensis Ihlow, Vamberger, Flecks, Hartmann, Cota, Makchai, Meewattana, Dawson, Kheng, Rödder, and Fritz 2016 (17:46, 17:47) (61)*


Khorat Snail-eating Turtle



Flora Ihlow / Sikhio, Thailand



Flora Ihlow / Thailand



** (stars = type localities)

Distribution: Laos, Thailand

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 120,892 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 158,124 sq. km

Size (Max SCL): male 15.5 cm, female 20.6 cm (Sumontha et al. 2016; Ihlow et al. 2016); **(Max CCL):** male 17.6 cm, female 21.0 cm (Ihlow et al. 2016)

IUCN Red List: Least Concern (LC) (Cota 2021)

CITES: Appendix II (2005), originally as part of *M. subtrijuga*

Synonymy:

Malayemys khoratensis Ihlow, Vamberger, Flecks, Hartmann, Cota, Makchai, Meewattana, Dawson, Kheng, Rödder, and Fritz 2016:16 (17:46, 17:47)
Type locality: "Udon Thani, Udon Thani Province, Thailand (17.36555°N, 102.81427°E)" [yellow star].
Type specimen: THNHM 25816, holotype; genotyped by Ihlow et al. (2016); Uetz et al. (2019) erroneously listed THNHM 25609, the holotype of *Malayemys isan*.
Etymology:* Toponym for the Khorat valley basin of northeastern Thailand, the dry plateau drained by Mekong tributaries.

Malayemys isan Sumontha, Brophy, Kunya, Wiboonatthapol, and Pauwels 2016:2 (17:46, 17:47)
Type locality: "Ban Na Klang (17°14'48.728"N, 102°12'32.479"E), Na Klang Sub-district, Na Klang District, Nong Bua Lamphu Province, northeastern Thailand" [green star].
Type specimen: THNHM 25609, holotype.
Etymology:* Toponym for the Isan region of northeastern Thailand, the dry plateau drained by Mekong tributaries.


Malayemys macrocephala (Gray 1859) (07:43, 17:46)
Malayan Snail-eating Turtle, Rice-field Terrapin



Flora Ihlow / CBFTT / Tap Than, Thailand



Flora Ihlow / CBFTT / *left*: Uthai Thani Prov., Thailand, *right*: Phetchaburi Prov., Thailand



(orange dots = probable trade or introduced; star = type locality)

Distribution:* Malaysia (*West*), Thailand

Possibly Introduced or Native: Indonesia (Sumatra) (prehistorical or historical?)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 199,641 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 642,262 sq. km

Size (Max SCL): male 15.6 cm, female 22.0 cm (Dawson et al. 2018 CBFTT); **(Max CCL):** male 17.8 cm, female 22.5 cm (Ihlow et al. 2016)

CBFTT Account: Dawson, Ihlow, Ettmar, van Dijk, and Thirakhupt (2018)

IUCN Red List: Least Concern (LC) (Cota 2021). Previously: Least Concern (LC) [Not Listed] (Behler 1996), as part of *Malayemys subtrijuga*

CITES: Appendix II (2005)

Synonymy:


Geoclemmys macrocephala Gray 1859:479
Clemmys macrocephala, Emys macrocephala, Damonia macrocephala, Geoclemys macrocephala, Malayemys macrocephala
Type locality: "Siam" [Thailand]. Restricted to "Thanyaburi, Pathum Thani Province, Thailand (Chao Phraya River Basin; approx. 50 km NNE of Bangkok; 14.017 N, 100.733 E)" [yellow star] by Brophy (2004:75).
Type specimen: NHMUK 1947.3.4.52 (formerly 1859.7.8.4), lectotype, designated by Brophy (2004:75).
Etymology:* Greek, *makros*, great or large, and *kephale*, head; referring to the large head.

Damonia megacephala Gray 1870c:44 (*nomen nudum*)
Etymology:* Greek, *mega*, large, and *kephale*, head; referring to the large head.
Comment:* Error for *Damonia macrocephala*.


Malayemys subtrijuga (Schlegel and Müller 1845) (17:46, 17:48) (61)*
Mekong Snail-eating Turtle



Flora Ihlow / Prek Toal, Tonle Sap Lake, Cambodia



Jeffrey E. Dawson / Central Cambodia





(native population = red, possibly introduced population = gray; orange dots = possibly native or introduced or trade; star = type locality)

Distribution:* Cambodia, Indonesia (*Java*), Laos, Thailand, Vietnam

Possibly Introduced or Native: Indonesia (*Java [prehistorical or historical?]*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 268,069 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 1,616,468 sq. km

Size (Max SCL): male 19.9 cm, female 23.7 cm (Platt et al. 2008; Dawson et al. 2020 CBFTT)

CBFTT Account: Dawson, Ihlow, and Platt (2020)

IUCN Red List: Near Threatened (NT A2d) (Horne, McCormack, and Timmins 2021). Previously: Vulnerable (VU) (ATTWG 2000); Least Concern (LC) [Not Listed] (Behler 1996)

CITES: Appendix II (2005)

Synonymy:

Emys subtrijuga Schlegel and Müller 1845:30 (17:46)
Damonia subtrijuga, Geoclemys subtrijuga, Malayemys subtrijuga
Type locality: "Java" [Indonesia] [orange star, location estimated]. Possibly introduced, but perhaps native. (17:46)
Type specimen: RMNH 6082, lectotype, designated by Brophy (2004:74), see also Hubrecht (1881) and Hoogmoed et al. (2010); NHMUK 1947.3.4.53 (formerly 1848.10.31.16) erroneously listed as "holotype" by Iverson (1992:138).
Etymology:* Latin, *sub-*, almost, a little, below, *subtrijuga*, somewhat similar to *trijuga*; referring to the apparent similarity of this species to *Emys trijuga* Schweigger 1812.

Cistudo gibbosa Bleeker 1857b:239 (*nomen nudum*)
Type locality: "Batavia...Java" [Jakarta, Indonesia] [orange star]. Restricted to "Borneo" [Kalimantan, Indonesia, in error, no star] by Gray (1873j:39).
Type specimen: NHMUK 1863.12.4.38, holotype, see Gray (1873j).
Etymology:* Latin, *gibbosa*, humped, strongly arched; referring to the three keels in the form of hump-like elevations on the carapace.

Emys nuchalis Blyth 1863:82
Bellia nuchalis
Type locality: "Java?" [Indonesia] [orange star, location estimated].
Type specimens: ZSI 824–26, syntypes (3), see Das et al. (1998) and Kundu et al. (2018a).
Etymology:* Latin, *nucha*, neck, spinal cord, *nuchalis*, belonging to the neck; referring to the relatively triangular nuchal scute in this taxon.

Emys megacephala Gray 1870c:44 (junior homonym and *nomen nudum*, not = *Emys megacephala* Holbrook 1836 [= *Graptemys geographica*], nor = *Emys megacephalus* Gray 1844 [= *Malaclemys terrapin terrapin*])
Etymology:* Greek, *mega*, large, and *kephale*, head; referring to the large head.

Damonia oblonga Gray 1871d:367
Type locality: "Batavia" [Jakarta, Java, Indonesia] [orange star].
Type specimen: NHMUK 1947.3.5.30 (formerly 1871.4.10.2 or 1871.10.12.2), holotype, see Gray (1873j).
Etymology:* Latin, *oblongus*, elongated; referring to the somewhat elongated carapace.

Morenia Gray 1870c

Indo-Burmese Eyed Turtles
(includes 2 species, 2 taxa)

Morenia Gray 1870c:62
Type species: *Morenia berdmorei* [= *Emys berdmorei* Blyth 1859 = subjective synonym of *Emys ocellata* Duméril and Bibron 1835], by subsequent designation by Lindholm (1929:279).
Etymology:* Eponym apparently honoring Thomas Matthew Berdmore (1811–1859), United Kingdom / Myanmar; naturalist and soldier with the East India Company in Burma (now Myanmar), who obtained the types of *Emys berdmorei* [= *Morenia ocellata*].

Morenia ocellata (Duméril and Bibron 1835)


Burmese Eyed Turtle



Indraneil Das / CBFTT / Yangon, Myanmar



left: Indraneil Das / CBFTT / Yangon, Myanmar
right: George R. Zug / CBFTT / Pyin Oo Lwin, Mandalay Div., Myanmar



** (orange dot = probable trade; stars = type localities)

Distribution:* Myanmar


Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 154,817 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 498,977 sq. km


Size (Max SCL): male 15.0 cm, female 23.9 cm (Platt et al. 2005; Das 2010 CBFTT)

CBFTT Account: Das (2010)
IUCN Red List: Endangered (EN A2cd+4cd) (Praschag, Platt, and Horne 2021). Previously: Vulnerable (VU) (ATTWG 2000); Near Threatened (NT) (TFTSG *in* Baillie and Groombridge 1996)
CITES: Appendix I (1975)
USA ESA:* Endangered (USFWS 1976)
Synonymy:
Emys ocellata Duméril and Bibron 1835:329
Batagur (Kachuga) ocellata, Batagur ocellata, Clemmys ocellata, Morenia ocellata
Type locality: "Bengale" [India or Bangladesh] [in error] [orange star, location estimated]. Restricted to "Irrawaddy (Ayeyarwady) river delta, Myanmar" [yellow star, location estimated] by lectotype designation by Bour (2009b:41).
Type specimen: MNHN 9167, lectotype, designated by Bour (2009b:41), also erroneously listed as MNHN 6167.
Etymology:* Latin, *ocellus*, little eye, *ocellata*, having eyespots; referring to the ocellated eye-like markings on the carapace.
Emys berdmorei Blyth 1859:281
Emys berdmorii, Batagur berdmorii, Batagur berdmorei, Kachuga berdmorei, Kachuga berdmoorei, Kachuga (Dongoka) berdmoorei, Morenia berdmorei
Type locality:* Not stated. Previously clarified as "Schwe Gyen, on the Sitang River, Pegu" [Shwegyin, Sittang River, Bago, Myanmar] [green star] by Blyth (1856:711).
Type specimens: ZSI, syntypes (3), apparently lost, not listed by Das et al. (1998) or Kundu et al. (2018a).
Etymology:* Eponym honoring Thomas Matthew Berdmore (1811–1859), United Kingdom / Myanmar; naturalist and soldier with the East India Company in Burma (now Myanmar), who obtained the types.

Morenia petersi Anderson 1879 (62)*
Indian Eyed Turtle


Shailendra Singh / CBFTT / Ganga R., Uttar Pradesh, India


Shailendra Singh / CBFTT / Ganga R., Uttar Pradesh, India


**
(orange dot = possibly introduced or trade; star = type locality)

Distribution: Bangladesh, India (*Assam, Bihar, Uttarakhand, Uttar Pradesh, West Bengal*), Nepal
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 489,287 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 915,311 sq. km
Size (Max SCL): male 19.4 cm, female 26.0 cm (Rashid and Swingland 1997; Das and Sengupta 2010 CBFTT; Baruah et al. 2012)
CBFTT Account: Das and Sengupta (2010)
IUCN Red List: Endangered (EN A2cd+4cd) (Ahmed and Singh 2021). Previously: Vulnerable (VU) (ATTWG 2000); Near Threatened (NT) (TFTSG *in* Baillie and Groombridge 1996)
CITES: Appendix II (2013)
Synonymy:
Batagur (Morenia) petersi Anderson 1879:761 (62)*
Batagur petersi, Morenia petersi
Type locality:* "Huzurapur in the Jessore District..[&]..Furreedpore [Faridpur]..[&]..Dacca [Dhaka]" [Bangladesh]. Restricted here in our annotation to "Dhaka, Bangladesh" [yellow star].
Type specimen:* ZMB 8865, lectotype, designated "holotype" by Fritz et al. (1994:164); ZSI 155–56, paralectotypes (2), see also Das et al. (1998), Murthy and Das (2009), and Das (2009), who recorded these specimens as "syntypes," also recorded as syntypes but not located by Kundu et al. (2018a).
Etymology:* Eponym honoring Wilhelm K.H. Peters (1815–1883), Germany; zoologist, herpetologist, and curator at the Museum für Naturkunde, Berlin.

Orlitia Gray 1873b
Malaysian Giant Turtles
(includes 1 species, 1 taxon)
Orlitia Gray 1873b:156
Type species: *Orlitia borneensis* Gray 1873b, by original monotypy.
Etymology:* Anonym, *orlitia*, possibly a fabricated name, or possibly an Eponym honoring someone named Orlit.
Heteroclemmys Peters 1874:622
Type species: *Clemmys (Heteroclemmys) gibbera* Peters 1874 [= subjective synonym of *Orlitia borneensis* Gray 1873b], by original monotypy.
Etymology:* Greek, *heteros*, different, other, and *klemmys*, turtle, or *kleptein*, to steal, and *emys*, turtle.
Brookeia Bartlett 1896:113
Type species: *Brookeia baileyi* [= *Hardella baileyi* Bartlett 1895b = subjective synonym of *Orlitia borneensis* Gray 1873b], by original monotypy.
Etymology:* Eponym honoring Sir Charles Brooke (1829–1917), United Kingdom / Malaysia; Rajah of Sarawak (now part of Malaysia) and founder of the Sarawak Museum. The etymology recorded by Frétey (2024) is incorrect in referring to Sir James Brooke (1803–1868), the previous Rajah and uncle to Sir Charles Brooke.
Adelochelys Baur 1896:319
Type species: *Adelochelys crassa* Baur 1896 [= subjective synonym of *Orlitia borneensis* Gray 1873b], by original monotypy.
Etymology:* Greek, *adelos*, concealed or unknown, and *chelys*, turtle.
Liemys Boulenger 1897a:468
Type species: *Liemys inornata* Boulenger 1897a [= subjective synonym of *Orlitia borneensis* Gray 1873b], by original monotypy.
Etymology:* Greek, *lissos*, smooth, and *emys*, turtle; referring to the smooth carapace.


Orlitia borneensis Gray 1873b (12:27)
Malaysian Giant Turtle



Jérôme Maran / Kalimantan, Indonesia [Borneo]



left: Joko Guntoro / Langkat, Sumatra, Indonesia / female
right: Sabine Schoppe / No data / trade



** (stars = type localities)

Distribution: Indonesia (*Kalimantan, Sumatra*), Malaysia (*East, West*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 653,240 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 1,956,123 sq. km

Size (Max SCL):* 80.0 cm (probable SCL) (Lim and Das 1999); **(Max CCL):*** 86.4 cm (Auliya and Chen 2024); 117.0 cm (possible CCL, unverified and questionable, possibly total body length) (Iskandar 2000)

IUCN Red List: Critically Endangered (CR A2acd) (Horne, Das, Hamidy, Kusrini, Guntoro, and As-singkily 2020). Previously: Endangered (EN) (ATTWG 2000); Near Threatened (NT) (TFTSG *in* Baillie and Groombridge 1996)

CITES: Appendix II (2003); zero quota for wild specimens for commercial purposes (2013)

Synonymy:

Cistudo borneensis Bleeker 1857a:473 (*nomen nudum*)
Cistudo borneënsis
Type locality: "Sintang aan de Kapoeas...Borneo" [Sintang, Kapuas River, Kalimantan, Indonesia] [yellow star].
Type specimen:* Not designated, possibly NHMUK 1947.3.4.9 (formerly 1863.12.4.37).
Etymology:* Toponym for Borneo, Kalimantan, Indonesia.

Orlitia borneensis Gray 1873b:157
Bellia borneensis
Type locality:* "Borneo." Restricted to "Sistang...Borneo" [Sintang, Kapuas River, Kalimantan, Indonesia] [yellow star] by Boulenger (1889:100).
Type specimen:* NHMUK 1947.3.4.9 (formerly 1863.12.4.37), holotype, collected and donated by Bleeker.
Etymology:* Toponym for Borneo, Kalimantan, Indonesia.

Clemmys (Heteroclemmys) gibbera Peters 1874:622
Clemmys gibbera, Heteroclemmys gibbera
Type locality: "Pulo Matjan (Borneo)" [Pulau Majang, Kalimantan, Indonesia] [green star].
Type specimen: ZMB 5022, holotype, see Fritz et al. (1994).
Etymology:* Latin, *gibbera*, humped, strongly arched; referring to the convex carapace.

Hardella baileyi Bartlett 1895b:83
Brookeia baileyi
Type locality: "Batang Lupar" [Sarawak, East Malaysia] [blue star].
Type specimen: Possibly SMK, holotype, originally live.
Etymology:* Eponym honoring Demetrius J.S. Bailey (1860–1908), United Kingdom / Malaysia; naval officer and naturalist in Sarawak (now part of Malaysia), who collected the type.

Adelochelys crassa Baur 1896:319
Type locality: Not known.
Type specimen: ZSM 3049/0, holotype, see Franzen and Glaw (2007).
Etymology:* Latin, *crassa*, thick, fat, firm; referring to the relatively strong and solid skull.

Liemys inornata Boulenger 1897a:469
Type locality: "Lobuk Antu district, Borneo" [Lubok Antu, Sarawak, East Malaysia] [pink star].
Type specimens: NHMUK 1947.3.4.28–29 (both formerly 1897.3.4.8), syntypes (2).
Etymology:* Latin, *inornata*, unadorned; referring to the uniformly unpatterned dark carapace and yellowish to brownish plastron.

Batagur signatus † Jaekel 1911:77
Type locality: "Pithecanthropus-schichten...Java...Trinil" [Indonesia] [gray star].
Type specimen: ZMB (MB R5), lectotype, fossil, nuchal, figured (pl.15,f.6), designated as "holotype" by Karl (1987:37).
Geological age: Pleistocene, Trinil Beds.
Etymology:* Latin, *signum*, sign, feature; referring to the pronounced scute furrows on the fossil carapace bone fragments.

Pangshura Gray 1856b (07:48)

Asian Roofed Turtles
(includes 4 species, 7 taxa)


Pangshura Gray 1856b:36
Type species: *Kachuga* (*Pangshura) tecta* [= *Emys tectum* Gray 1830b], by subsequent designation by Lindholm (1929:278).
Etymology:* Vernacular (Bengali), *panshura*, local indigenous name for a turtle.

Cuchoa Gray 1870c:61
Type species: *Cuchoa tentoria* [= *Emys tentoria* Gray 1834a], by subsequent designation by Lindholm (1929:278).
Etymology:* Vernacular (Hindi), *kachhua*, local indigenous name for a freshwater turtle.

Jerdonella Gray 1870c:61
Type species: *Jerdonella sylhetensis* [= *Pangshura sylhetensis* Jerdon 1870], by original monotypy.
Etymology:* Eponym honoring Thomas C. Jerdon (1811–1872), United Kingdom / India; physician, zoologist, and ornithologist with the East India Company in India.

Emia Gray 1870c:61
Type species: *Emia smithii* [= *Batagur smithii* Gray 1863g], by original monotypy.
Etymology:* Anonym, *emia*, possibly a fabricated name, or possibly an Eponym honoring the author's wife, Emma Gray (1787–1876).

Pangshura smithii (Gray 1863g)
Brown Roofed Turtle
(includes 2 subspecies)



(subspecies: *smithii* = red, *pallidipes* = blue; overlap = intergrades; orange dot = possibly introduced)

Distribution: Bangladesh, India (*Assam, Bihar, Punjab, Uttar Pradesh*), Nepal, Pakistan

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 556,637 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 2,686,576 sq. km

Size (Max SCL): male 12.8 cm, female 24.0 cm (see subspp.)

IUCN Red List: Near Threatened (NT A4d) (Ahmed, Praschag, Choudhury, and Singh 2021). Previously: Near Threatened (NT) (ATTWG 2000); Least Concern (LC) [Not Listed] (Behler 1996)

TFTSG Provisional Red List: Near Threatened (NT) (2018)

CITES: Appendix II, as *Pangshura* spp. (2003)


Pangshura smithii smithii (Gray 1863g)
Brown Roofed Turtle



Hermann Schleich / Koshi R., Nepal



Amtyaz Safi / Pakistan



(subspecies: *smithii* = red, *pallidipes* = blue; overlap = intergrades; orange dot = possibly introduced; star = type locality)

Distribution: Bangladesh, India (*Assam, Bihar, Punjab, Uttar Pradesh*), Nepal, Pakistan

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 519,539 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 2,686,576 sq. km

Size (Max SCL): male 12.8 cm, female 24.0 cm (Rashid and Swingland 1997; Ernst et al. 2006b; Itescu et al. 2014; Baruah et al. 2016)

Synonymy:

Batagur smithii Gray 1863g:253

Pangshura smithii, Clemmys smithii, Emia smithii, Kachuga smithii, Kachuga smithii smithii, Pangshura smithii smithii

Type locality: "North-western India: Punjab; River Chenab" [Punjab, Pakistan] [yellow star, location estimated].


Type specimens: NHMUK 1947.3.4.69–70 (formerly 1863.2.21.87, 1863.6.5.3)., syntypes (2).

Etymology:* Eponym honoring Sir Andrew Smith (1797–1872), United Kingdom (Scotland) / South Africa / India; physician, administrator, and zoologist, who worked primarily in South Africa, and obtained the types in India.


Pangshura smithii pallidipes (Moll 1987)
Pale-footed Roofed Turtle



Joyee Chan / India / trade



Peter Praschag / India



(subspecies: *smithii* = red, *pallidipes* = blue; overlap = intergrades; orange dot = possibly introduced; star = type locality)

Distribution: India (*Bihar, Uttar Pradesh*), Nepal

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 41,447 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 75,803 sq. km

Size (Max SCL): male 10.8 cm, female 22.3 cm (Baruah et al. 2016)

Synonymy:

Kachuga smithii pallidipes Moll 1987:8

Pangshura smithii pallidipes

Type locality: "Gandak River, Bherihari Wildlife Sanctuary, Bettiah (West Champaran) District, Bihar" [India] [yellow star].

Type specimen: FMNH 224177, holotype.

Etymology:* Latin, *pallidus*, pale, and *pes*, foot; referring to the pale colored feet.

Pangshura sylhetensis Jerdon 1870

Assam Roofed Turtle, Sylhet Roofed Turtle




Indraneil Das / CBFTT / Kulsi River, Kamrup District, Assam, India





left: Indraneil Das / CBFTT / Kulsi River, Kamrup District, Assam, India
right: Peter Praschag / India



** (star = type locality)

Distribution: Bangladesh, Bhutan (?), India (*Arunachal Pradesh, Assam, Meghalaya, Mizoram, Nagaland*), Myanmar (?)

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 144,452 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 346,432 sq. km

Size (Max SCL): male 9.8 cm, female 21.0 cm (Das et al. 2010 CBFTT; Baruah et al. 2012, 2016)

CBFTT Account: Das, Sengupta, and Praschag (2010)

IUCN Red List: Critically Endangered (A2cd+4cd) (Praschag, Das, Ahmed, and Singh 2021). Previously: Endangered (EN) (ATTWG 2000); Data Deficient (DD) (TFTSG *in* Baillie and Groombridge 1996)

CITES: Appendix II, as *Pangshura* spp. (2003)

Synonymy:

Pangshura sylhetensis Jerdon 1870:69

Jerdonella sylhetensis, Kachuga sylhetensis

Type locality: "the stream that runs from the Terria Ghat at the foot of the Khasi Hills" [originally Assam, India; now nr. Bholaganj, Sylhet District, Bangladesh]. Emended to "Sylhet, a stream at the foot of the Khasi Hills" by Boulenger (1889:58) [yellow star].

Type specimens: NHMUK 1947.3.4.22, 1947.3.4.62–63 (formerly 1870.11.29.51–52), syntypes (3), see Das (2009).

Etymology:* Toponym for Sylhet District, India (now Bangladesh).

Pangshura tecta (Gray 1830b)

Indian Roofed Turtle




Peter Praschag / India





left: Peter Praschag / India
right: Joyee Chan / Bangladesh / trade



** (orange dots = possible translocations; stars = type localities)

Distribution:* Bangladesh, India (*Arunachal Pradesh, Assam, Bihar, Gujarat, Haryana, Jammu and Kashmir, Madhya Pradesh, Meghalaya, Odisha, Punjab, Rajasthan, Uttar Pradesh, Uttarakhand, West Bengal*), Nepal, Pakistan

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 920,058 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 2,783,217 sq. km

Size (Max SCL): male 27.6 cm, female 33.9 cm (Agarwal et al. 1986)

IUCN Red List: Vulnerable (VU A4d) (Ahmed, Praschag, and Singh 2021). Previously: Least Concern (LC) (ATTWG 2000); Least Concern (LC) [Not Listed] (Behler 1996)

CITES: Appendix I (1975)

USA ESA:* Endangered, as *Kachuga tecta tecta* (USFWS 1976)

Synonymy:

Emys tectum Gray 1830b:pl.72

Emys tecta, Clemmys (Clemmys) tecta, Clemmys tecta, Batagur (Pangshura) tecta, Batagur tecta, Clemmys

tectum, Pangshura tecta, Pangshura tectum, Kachuga tectum, Kachuga (Pangshura) tecta, Kachuga tectum tectum, Kachuga tecta, Kachuga tecta tecta

Type locality: "India." [yellow star, likely near Calcutta]

Type specimens: OUM 8430–31, syntypes (2), discussed by Nowak-Kemp and Fritz (2010).

Etymology:* Latin, *tectum*, roof; referring to the roof-shaped carapace.

Testudo dura Hamilton *in* Gray 1831d:23 (*nomen nudum*)

Etymology:* Latin, *dura*, hard; referring to the hard shell.

Testudo katuya Hamilton *in* Gray 1831d:23 (*nomen nudum*)

Etymology:* Vernacular (Bengali?), *katuya*; possibly a local indigenous name for the species.

Testudo pangshure Hamilton *in* Gray 1831d:23 (*nomen nudum*)

Testudo pangshura

Etymology:* Vernacular (Bengali), *panshura*; local indigenous name for a turtle.

Testudo khagraskata Hamilton *in* Gray 1831d:23 (*nomen nudum*)

Etymology:* Vernacular (Bengali?), *khagraskata*; possibly a local indigenous name for the species.

Emys trigibbosa Lesson 1831a:121

Type locality: "le Gange" [Bangladesh] [Ganges River or delta, Bangladesh] [green star, location estimated].

Type specimen: Possibly MNHN, holotype, not located.

Etymology:* Latin, *tria*, three, and *gibbosa*, humped; referring to the three anterior vertebral humps or spines.

Emys namadicus † Theobald 1860:295 (*nomen nudum*)

Emys nomadicus, Emys namadica

Type locality: "near Moar Domur...Nerbudda Valley" [Madhya Pradesh, India] [gray star, location estimated].

Type specimen: whole fossil shell, not located.

Geological age: Tertiary.

Etymology:* Toponym for the Narmada River in the Nerbudda Valley, India.

Pangshura dura Gray 1869a:205

Type locality: Not designated.

Type specimen: NHMUK s/n, holotype, not located.

Etymology:* Latin, *dura*, hard; referring to the hard shell.

Pangshura ventricosa Gray 1870c:60

Type locality: "India." Restricted to "Assam" [India] [blue star, location estimated] by Boulenger (1889:59).


Type specimen: NHMUK 1947.3.4.71 (formerly 1870.11.29.55), holotype.

Etymology:* Latin, *ventricosa*, potbellied; referring to the bulging, deep shell.

Pangshura tentoria (Gray 1834a)

Indian Tent Turtle

(includes 3 subspecies)



** (subspecies: *tentoria* = red, *circumdata* = blue, *flaviventer* = green; overlap = intergrades; orange dots = probable trade or questionable)

Distribution: Bangladesh, India (*Andhra Pradesh, Assam, Bihar, Chhattisgarh, Maharashtra, Odisha, Rajasthan, Telangana, Uttar Pradesh, Uttarakhand, West Bengal*), Nepal

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 569,009 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 1,892,769 sq. km

Size (Max SCL): male 17.9 cm, female 27.1 cm (see subspp.)

IUCN Red List: Least Concern (LC) (Choudhury, Das, Praschag, and Singh 2021). Previously: Least Concern (LC) (ATTWG 2000); Least Concern (LC) [Not Listed] (Behler 1996)

CITES: Appendix II, as *Pangshura* spp. (2003)

Pangshura tentoria tentoria (Gray 1834a) (07:49)


Indian Tent Turtle



Peter Paul van Dijk / India / captivity / Madras Crocodile Bank Trust



Peter Praschag / *left*: Mahanadi R. Odisha, India, *right*: Godavari R., Odisha, India



** (subspecies: *tentoria* = red, *flaviventer* = green; stars = type localities)

Distribution: India (*Andhra Pradesh, Chhattisgarh, Maharashtra, Odisha*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 144,814 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 449,403 sq. km

Size (Max SCL): male 14.9 cm, female 22.8 cm (Baruah et al. 2016)

Synonymy:

Emys tentoria Gray 1834a:54

Batagur (Pangshura) tentoria, Batagur tentoria, Clemmys tentoria, Pangshura tentoria, Pangshura tentori, Cuchoa tentoria, Kachuga tectum tentoria, Kachuga tecta tentoria, Kachuga tentoria tentoria, Pangshura tentoria tentoria

Type locality: "Indiae Orientalis regione Dukhun" [Deccan Plateau, India] [yellow star, location estimated and imprecise]. Emended to "Deccan" [Deccan Plateau, India] by Boulenger (1889:59); restricted to "Dhond, Poona Dist." [= Daund, Pune District, Maharashtra, India; in error] [orange star] by Smith (1931:128).

Type specimen: NHMUK 1947.3.4.72, holotype.

Etymology:* Latin, *tentoria*, tent; referring to the strongly arched carapace shape reminiscent of a tent.

Pangshura tentorium Gray 1869a:205 (*nomen novum*)
Comment: Unjustified emendation or error for *tentoria*.
Etymology:* Latin, *tentoria*, tent; referring to the strongly arched carapace shape reminiscent of a tent.


Emys (Pangshura) tectum intermedia Blanford 1870:339
Kachuga intermedia, Kachuga tectum intermedia
Type locality: "Chappa and Korba in Biláspúr, on the Hasdo River, a tributary of the upper Mahanaddi...above Sambhalpúr" [Chhattisgarh, India] [green star].
Type specimens: ZSI 122–23, syntypes (2), see Das et al. (1998), not located by Kundu et al. (2018a).
Etymology:* Latin, *intermedia*, in-between; referring to the relationship of this species to others.

Pangshura leithii Gray 1870c:60
Type locality: "River Poonah" [Maharashtra, India; in error, possible trade] [brown star, location estimated].
Type specimen: NHMUK 1947.3.4.61, holotype.
Etymology:* Eponym honoring Andrew H. Leith (1827–1882), United Kingdom / India; physician, naturalist, and malacologist who worked in India on the Bombay Sanitary Commission and collected the type.

Pangshura tentoria circumdata (Mertens 1969)
Pink-ringed Tent Turtle


Nikhil Whitaker / Uttar Pradesh, India / juvenile


Peter Paul van Dijk / Uttar Pradesh, India / captive / Kukrail Breeding Center


**

(subspecies: *tentoria* = red, *circumdata* = blue, *flaviventer* = green; overlap = intergrades; star = type locality)

Distribution:* India (*Bihar, Haryana, Madhya Pradesh, Rajasthan, Uttar Pradesh*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 145,844 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 332,721 sq. km

Size (Max SCL): male 8.8 cm, female 27.1 cm (Rai et al. 2002)


Synonymy:

Kachuga tecta circumdata Mertens 1969a:24
Kachuga tentoria circumdata, Pangshura tentoria circumdata
Type locality: "Meerut, Indien" [Uttar Pradesh, India] [yellow star].
Type specimen: SMF 52793, holotype, see Das (2009).
Etymology:* Latin, *circumdata*, surrounded, enclosed; referring to the reddish-pink ring that surrounds the carapace.

Pangshura tentoria flaviventer Günther 1864 (07:49)
Yellow-bellied Tent Turtle


Hermann Schleich / Koshi R., Nepal


Peter Praschag / India


**

(subspecies: *tentoria* = red, *circumdata* = blue, *flaviventer* = green; overlap = intergrades; orange dot = probable trade or questionable; star = type locality)

Distribution: Bangladesh, India (*Assam, Bihar, Uttar Pradesh, West Bengal*), Nepal

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 296,494 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 776,609 sq. km

Size (Max SCL): male 17.9 cm, female 20.4 cm (Baruah et al. 2012)

Synonymy:

Pangshura flaviventer Günther 1864:35
Kachuga tecta flaviventer, Pangshura flaviventus, Kachuga tentoria flaviventer, Pangshura tentoria flaviventer
Type locality: "Bengal" [India] [yellow star, location estimated, likely nr. Calcutta]. Emended to "India" by Boulenger (1889:59).
Type specimen: NHMUK 1947.3.4.82, holotype, see Das (2009).
Etymology:* Latin, *flavus*, yellow, *venter*, belly, abdomen; referring to the uniformly yellowish or cream-colored plastron.

Cuchoa flaviventris Gray 1870c:61 (*nomen novum*)
Etymology:* Latin, *flavus*, yellow, *venter*, belly, abdomen; referring to the uniformly yellowish or cream-colored plastron.
Comment: Unjustified emendation or error for *flaviventer*.

Siebenrockiella Lindholm 1929

Asian Marsh Turtles

(includes 2 species, 2 taxa)

Bellia Gray 1869a:197 (junior homonym, not = *Bellia* Milne-Edwards 1848 [= Crustacea] or *Bellia* Bate 1851 [= Crustacea] or *Bellia* Tutt 1902 [= Lepidoptera])

Type species: *Bellia crassicollis* [= *Emys crassicollis* Gray 1830e], by original monotypy.

Etymology:* Eponym honoring Thomas Bell (1792–1880), United Kingdom; herpetologist, dental surgeon, and Professor of Zoology at King's College, London.

Siebenrockiella Lindholm 1929:280 (*nomen novum*)

Type species: *Siebenrockiella crassicollis* [= *Emys crassicollis* Gray 1830e], by original designation.

Etymology:* Eponym honoring Friedrich Siebenrock (1853–1925), Austria; zoologist, herpetologist, and curator at the Naturhistorisches Museum, Vienna.

Panyaenemys Diesmos, Parham, Stuart, and Brown 2005:38

Type species: *Siebenrockiella (Panyaenemys) leytensis* [= *Heosemys leytensis* Taylor 1920], by original designation.


Etymology:* Vernacular (Pala'wan), *panya-en*, enchanted, and Greek, *emys*, turtle; referring to indigenous tradition of this turtle being a favorite of forest spirits on Palawan.

Siebenrockiella crassicollis (Gray 1830e) [(10:7)]


Black Marsh Turtle



Edward O. Moll / Malaysia



left: Sabine Schoppe / No data / trade
right: Jérôme Maran / No data / captivity



(orange dot = probable trade; stars = type localities)

Distribution:* Cambodia, Indonesia (*Java*, *Kalimantan*, *Sumatra*), Laos (?), Malaysia (*East [?]*, *West*), Myanmar, Singapore, Thailand, Vietnam

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 907,850 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 3,036,820 sq. km

Size (Max SCL): male 20.3 cm, female 20.0 cm (Le 2007; Itescu et al. 2014)

IUCN Red List: Endangered (EN A2cd) (Horne, Kusrini, Hamidy, Platt, Guntoro, and Cota 2021). Previously: Vulnerable (VU) (ATTWG 2000); Least Concern (LC) [Not Listed] (Behler 1996)

CITES: Appendix II (2003)

Synonymy:

Emys crassicollis Gray 1830e:8 [(10:7)]

Clemmys (Clemmys) crassicollis, *Clemmys crassicollis*, *Bellia crassicollis*, *Orlitia crassicollis*, *Siebenrockiella crassicollis*

Type locality: "Sumatra" [Indonesia] [yellow star, location estimated].

Type specimens: OUM 8479–81, syntypes (3), see Nowak-Kemp and Fritz (2010); NHMUK 1947.3.5.36, erroneously listed as "holotype" by King and Burke (1989) and Iverson (1992), is not a type.

Etymology:* Latin, *crassum*, thick, fat, firm, and *collum*, neck; referring to the relatively broad head and neck.

Emys nigra Blyth 1856:713 (junior homonym, not = *Emys nigra* Hallowell 1854 [= *Emys* or *Actinemys marmorata*])

Type locality: "valley of the Tenasserim" [Myanmar] [green star, location estimated].

Type specimens: Possibly ZSI, syntypes, apparently lost, not listed by Das et al. (1998) or Kundu et al. (2018a).

Etymology:* Latin, *nigra*, black; referring to the shell color.

Emys crassilabris Gray *in* Theobald 1876:10 (*nomen novum*)

Bellia crassilabris

Etymology:* Latin, *crassum*, thick, fat, firm, and *labrum*, lip; referring to a relatively thick lip.

Comment: Error for *crassicollis* (corrected in Errata insert).

Pangshura cochinchinensis Tirant 1884:159

Kachuga cochinchinensis

Type locality: "Cochinchine" [Vietnam] [blue star, location estimated].

Type specimens: Not known or located.

Etymology:* Toponym for Cochinchina (now southern Vietnam).

Siebenrockiella leytensis (Taylor 1920) (07:50)
Palawan Forest Turtle, Philippine Forest Turtle




Rafe M. Brown / TCC / CBFTT / Palawan, Philippines





left: Sabine Schoppe / CBFTT / Palawan, Philippines
right: Emerson Sy / CBFTT / Palawan, Philippines



(orange dots = probable trade; stars = type localites)

Distribution: Philippines (*Palawan [not Leyte]*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 4,231 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 8,983 sq. km

Size (Max SCL): male 29.9 cm, female 21.2 cm (Diesmos et al. 2012 CBFTT)

CBFTT Account: Diesmos, Buskirk, Schoppe, Diesmos, Sy, and Brown (2012)

IUCN Red List: Critically Endangered (CR A4abcd) (Schoppe, Diesmos, and Diesmos 2021). Previously: Critically Endangered (CR) (ATTWG 2000); Endangered (EN) (TFTSG *in* Baillie and Groombridge 1996)

CITES: Appendix II (2003)

Synonymy:

Heosemys leytensis Taylor 1920:131
Geoemyda leytensis, Siebenrockiella leytensis, Siebenrockiella (Panyaenemys) leytensis
Type locality: "Cabalian, southern Leyte, P. I." [Philippines] [in error] [orange star]. Restricted to "northern Palawan in the Province of Palawan, Philippines" [yellow star, location estimated] by Diesmos et al. (2012:3).
Type specimens: University of the Philippines s/n, syntypes (2), moved to the Bureau of Science in Manila, destroyed during WW II; CAS 60930, neotype, designated by Buskirk (1989:226).
Etymology:* Toponym for Leyte Island, Philippines.

Subfamily Geoemydinae Theobald 1868a (12:21)

Geoemydine Pond Turtles
(includes 10 genera, 46 species, 61 taxa)
Geoemydidae Theobald 1868a:vi
Geoemydina Gray 1869a:184
Geoemydinae Gaffney and Meylan 1988:207


Cuora Gray 1856a (07:32, 12:22, 17:40)

Asian Box Turtles
(includes 17 species, 25 taxa)
Cuora Gray 1856a:198
Type species: *Cuora amboinensis* [= *Testudo amboinensis* Riche *in* Daudin 1801], by subsequent designation by Stejneger (1907:503).
Etymology:* Vernacular (Malayan), *kura-kura*, freshwater turtle.
Cistoclemmys Gray 1863e:175 (63)*
Type species: *Cistoclemmys flavomarginata* Gray 1863e, by original monotypy.
Etymology:* Latin, *cista*, a box, and Greek, *klemmys*, turtle, or *kleptein*, to steal, and *emys*, turtle; referring to its ability to retract and hide its head and neck.
Pyxidea Gray 1863e:175
Type species: *Pyxidea mouhotii* [= *Cyclemys mouhotii* Gray 1862a], by original monotypy.
Etymology:* Greek, *pyxis*, box, and *-idea*, similar; referring to the box-like appearance of these turtles.
Pyxiclemmys Gray 1863e:176
Type species: *Cuora* (*Pyxiclemmys*) *trifasciata* [= *Sternothaerus trifasciatus* Bell 1825a], by original monotypy.
Etymology:* Greek, *pyxis*, a box, and *klemmys*, turtle, or *kleptein*, to steal, and *emys*, turtle; referring to its ability to retract and hide its head and neck.
Cystoclemmys Gray 1870c:20 (*nomen novum*)
Comment:* Unjustified emendation of *Cistoclemmys*.

"Cuora amboinensis" Species Complex

Southeast Asian Box Turtles
(includes 5 species, 7 taxa)

Cuora amboinensis (Riche *in* Daudin 1801) (12:23, 17:41) (64)*
East Indonesian Box Turtle, Amboina Box Turtle
(includes 2 subspecies)



(subspecies: *amboinensis* = red, *aurantiae* = blue; overlap = intergrades)

Distribution:* Indonesia (*Moluccas, Sulawesi*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 219,356 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 986,360 sq. km

Size (Max SCL):* male 19.2 cm, female 20.8 cm (see subspp.)

CBFTT Account: Schoppe and Das (2011) as part of *Cuora amboinensis* (sensu lato)

IUCN Red List: Endangered (EN A2d) (Cota, Hoang, Horne, Kusrini, McCormack, Platt, Schoppe, and Shepherd 2020) as part of *Cuora amboinensis* (sensu lato). Previously: Vulnerable

(VU) (ATTWG 2000); Near Threatened (NT) (TFTSG *in* Baillie and Groombridge 1996)

CITES: Appendix II, as *Cuora* spp. (2000)

Cuora amboinensis amboinensis (Riche *in* Daudin 1801) (12:23) (64)*


East Indonesian Box Turtle, Moluccas Box Turtle



Sabine Schoppe / CBFTT / Sulawesi, Indonesia



left: Sabine Schoppe; *right*: Torsten Blanck / Sulawesi, Indonesia



** (subspecies: *amboinensis* = red, *aurantiae* = blue, overlap = intergrades; stars = type localities)

Distribution:* Indonesia (*Moluccas*, *Sulawesi*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 169,123 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 986,360 sq. km

Size (Max SCL):* male 19.2 cm, female 20.8 cm (Blanck et al. 2023; T. Blanck, unpubl. data)

TFTSG Provisional Red List: Endangered (EN) (2025)

CITES: Appendix II, as *Cuora* spp. (2000)

Synonymy:

Testudo melanocephala Van-Ernest *in* Daudin 1801:128 (*nomen oblitum*)

Emys melanocephala, *Clemmys (Clemmys) melanocephala*

Type locality: "une des îles Moluques ou Philippines" [Moluccas, Indonesia, or Philippines]. Restricted to "Batavia [= Djakarta, Java] [in error] und die Molukken oder die Philippinen" by Rummler and Fritz (1991:38), and to "Moluccas" [Indonesia] [yellow star, location estimated, possibly Ambon] by Fritz and Havaš (2007:214).

Type specimen: Possibly MNHN, not known, apparently lost (Bour, pers. comm.).

Etymology:* Greek, *melanos*, black, dark, and *kefale*, head; referring to the dark-colored top of the head.

Testudo amboinensis Riche *in* Daudin 1801:309

Emys amboinensis, *Terrapene amboinensis*, *Kinosternon amboinense*, *Cistuda amboinensis*, *Cuora amboinensis*, *Cistudo amboinensis*, *Cyclemys amboinensis*, *Cuora amboinensis amboinensis*

Type locality: "Amboine" [Ambon, Moluccas, Indonesia] [yellow star].

Type specimen: Not known, lost at sea en route to MNHN (Bour, pers. comm. *in* Rummler and Fritz 1991). MHNG 748.011, neotype, designated by Blanck et al. (2023:23).

Etymology:* Toponym for Ambon Island, Moluccas, Indonesia.

Emys melanogaster Bleeker *in* Gray 1864a:12 (*nomen nudum*)

Type locality: "Batchian and Boero" [Bacan and Buru, Moluccas, Indonesia] [green stars]. Restricted to "Borneo" [in error] by Gray (1873j:21).

Type specimen: NHMUK 1863.12.4.121, holotype, see Gray (1873j).

Etymology:* Greek, *melanos*, black, dark, and *gaster*, belly, stomach; referring to the dark-colored plastron.

Emys hypselonotus Bleeker *in* Gray 1864a:12 (*nomen nudum*)

Emys hypsilonotus

Type locality: "Batchian and Boero" [Bacan and Buru, Moluccas, Indonesia] [green stars]. Restricted to "Borneo" [in error] by Gray (1873j:22).

Type specimen: NHMUK 1863.12.4.139, holotype, see Gray (1873j).

Etymology:* Greek, *ypsilon*, high, steep, and *notos*, the back; referring to the relatively high-arched carapace.

Emys gastrotaenia Bleeker *in* Gray 1873j:21 (*nomen nudum*)

Type locality: "Borneo." Locality likely in error, probably "Batchian and Boero" [Bacan and Buru, Moluccas, Indonesia] [green stars].

Type specimen: NHMUK 1863.12.4.48, holotype, see Gray (1873j).

Etymology:* Greek, *gastros*, belly, stomach, and *tainia*, bandage, band; possibly referring to the dark mark on the plastron.

Cuora amboinensis aurantiae Blanck, Gaillard, Protiva, Wheatley, Shi, Liu, Ray, and Anders 2023 (64)*
Palu Box Turtle, Sulawesi Box Turtle




Torsten Blanck / Sulawesi, Indonesia





Torsten Blanck / Sulawesi, Indonesia



**

(subspecies: *amboinensis* = red, *aurantiae* = blue, overlap = intergrades; star = type locality)

Distribution:* Indonesia (*Sulawesi*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 76,808 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 194,664 sq. km

Size (Max SCL):* male 15.5 cm, female 20.1 cm (Blanck et al. 2023; T. Blanck, unpubl. data)


TFTSG Provisional Red List: Critically Endangered (CR) (2025)

CITES: Appendix II, as *Cuora* spp. (2000)

Synonymy:*

Cuora amboinensis aurantiae Blanck, Gaillard, Protiva, Wheatley, Shi, Liu, Ray, and Anders 2023:26
Type locality: "Palu, Sulawesi, Indonesia" [yellow star].
Type specimen: NHMW 41494:1, holotype.
Etymology:* Latin, *aurantius*, orange; referring to the orange head pattern of this taxon.

Cuora couro (Leschenault de la Tour *in* Schweigger 1812) (64)*
Southeast Asian Box Turtle
(includes 2 subspecies)



**

(subspecies: *couro* = red, *kamaroma* = blue; orange dots = probable trade or introduced)

Distribution:* Brunei, Cambodia, India (*Nicobar Islands*), Indonesia (*Java, Kalimantan, Lesser Sundas, Sumatra, Timor*), Laos, Malaysia (*East, West*), Myanmar, Philippines (*Palawan, Sulu Archipelago*), Singapore, Thailand, Timor-Leste, Vietnam

Introduced:* Philippines (*Luzon, Mindoro*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 1,804,560 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 8,312,367 sq. km

Size (Max SCL):* male 21.9 cm, female 27.8 cm (see subspp.)

CBFTT Account: Schoppe and Das (2011) as part of *Cuora amboinensis*


IUCN Red List: Endangered (EN A2d) (Cota, Hoang, Horne, Kusrini, McCormack, Platt, Schoppe, and Shepherd 2020) as part of *Cuora amboinensis*. Previously: Vulnerable (VU) (ATTWG 2000); Near Threatened (NT) (TFTSG 1996)

CITES: Appendix II, as *Cuora* spp. (2000)

Cuora couro couro (Leschenault de la Tour *in* Schweigger 1812) [(64)*]
Indo-Malayan Box Turtle


Torsten Blanck / nr. Jakarta, Java, Indonesia


Sabine Schoppe / CBFTT / Indonesia


**

(subspecies: *couro* = red, *kamaroma* = blue; star = type locality)

Distribution:* Brunei, Indonesia (*Java, Kalimantan, Lesser Sundas, Sumatra, Timor*), Malaysia (*East, West*), Philippines (*Palawan, Sulu Archipelago*), Singapore, Thailand, Timor-Leste

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 1,286,966 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 6,391,521 sq. km

Size (Max SCL):* male 21.3 cm, female 25.0 cm (Lim and Das 1999; Schoppe and Das 2011 CBFTT)

TFTSG Provisional Red List: Vulnerable (VU) (2025)

CITES: Appendix II, as *Cuora* spp. (2000)

Synonymy:

Emys couro Leschenault de la Tour *in* Schweigger 1812:315
Terrapene couro, Cuora amboinensis couro
Type locality: "Java" [Indonesia] [yellow star, location estimated, possibly Jakarta].
Type specimen: MNHN 7931, holotype, see Rummler and Fritz (1991) and Ceríaco and Bour (2012).
Etymology:* Vernacular (Malayan), *kura-kura*, freshwater turtle.
Comment:* The original author's name was incorrectly given by Schweigger as Lechenault, but his name was actually Leschenault de la Tour, and has therefore been corrected (see Blanck et al. 2023).


Terrapene bicolor Bell 1826:485
Type locality: "Americâ septentrionali" [in error].
Type specimens: OUM, syntypes (2), one originally live, apparently lost, not located by Nowak-Kemp and Fritz (2010), type specimen figured (pl.16).
Etymology:* Latin, *bi*, twice, and *color*, color; referring to the predominantly blackish and yellowish coloring of the shell.

Emys (Cistuda) amboinensis leveriana Gray 1830e:7 [(10:7)]
Cistuda amboinensis leveriana
Type locality: "Java and Penang" [Indonesia and West Malaysia].
Type specimens: NHMUK, holotype, not located.
Etymology:* Eponym for Sir Ashton Lever (1729–1788), United Kingdom; naturalist collector and founder of the Leverian Museum in London that housed many specimens from the circumglobal voyages of James Cook.

Cuora couro kamaroma Rummler and Fritz 1991 [(12:23)] [(64)*]
Southeast Asian Box Turtle


Torsten Blanck / Khao Yai National Park, Nahkon Rachasima, Thailand


Torsten Blanck / Khao Yai National Park, Nahkon Rachasima, Thailand


**

(subspecies: *couro* = red, *kamaroma* = blue, orange dots = probable trade; star = type locality)

Distribution:* Cambodia, India (*Nicobar Islands*), Laos, Myanmar, Thailand, Vietnam

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 517,594 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 1,743,998 sq. km

Size (Max SCL):* male 21.9 cm, female 27.8 cm (Blanck et al. 2023; T. Blanck, unpubl. data)

TFTSG Provisional Red List: Vulnerable (VU) (2025)

CITES: Appendix II, as *Cuora* spp. (2000)


Synonymy:

Cuora amboinensis kamaroma Rummler and Fritz 1991:39
Type locality: "circa 50 km nördlich von Bangkok, Thailand" [yellow star].
Type specimen: ZMH R-277, holotype, see Hallermann (1998).
Etymology:* Greek, *kamaroma*, a dome or bulge; referring to the high vaulted carapace.

Cuora lineata McCord and Philippen 1998 (17:41) (64, 66)*
Myanmar Box Turtle, Burmese Box Turtle




Hans-Dieter Philippen / CBFTT / Myanmar / captivity





Hans-Dieter Philippen / Myanmar / captivity



** (star = type locality)

Distribution:* India (*Manipur*), Myanmar

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 65,179 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 144,583 sq. km

Size (Max SCL):* male 22.1 cm, female 25.7 cm (Blanck et al. 2023; T. Blanck, unpubl. data)

TFTSG Provisional Red List: Endangered (EN) (2025)

CITES: Appendix II, as *Cuora* spp. (2000)


Synonymy:

Cuora amboinensis lineata McCord and Philippen 1998:54
Type locality: "Myitkyina, Kachin Province, Myanmar (Burma)" [yellow star].
Type specimen: USNM 122189, holotype, see Reynolds et al. (2007).
Etymology:* Latin, *linea*, line, stripe, *lineata*, having lines, striped; referring to the mid-dorsal carapacial light stripe and the lateral costal stripes.


Cuora philippinensis Blanck, Gaillard, Protiva, Wheatley, Shi, Liu, Ray, and Anders 2023 (64)*
Philippine Box Turtle



Torsten Blanck / Visayas, Philippines



left: Alice Schumacher / Holotype, NHMW 29470:2 / Negros, Philippines
right: Torsten Blanck / Visayas, Philippines



** (star = type locality)

Distribution:* Philippines

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 286,029 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 685,546 sq. km

Size (Max SCL):* male 16.7 cm, female 17.3 cm (Blanck et al. 2023; T. Blanck, unpubl. data)

TFTSG Provisional Red List: Vulnerable (VU) (2025)

CITES: Appendix II, as *Cuora* spp. (2000)


Synonymy:*

Cuora philippinensis Blanck, Gaillard, Protiva, Wheatley, Shi, Liu, Ray, and Anders 2023:40
Type locality: "Negros, Philippines" [yellow star, location estimated].
Type specimen: NHMW 29470:2, holotype.
Etymology:* Toponym for the Philippines.


Cuora praschagi Blanck, Gaillard, Protiva, Wheatley, Shi, Liu, Ray, and Anders 2023 [64]*
Assam Box Turtle



Torsten Blanck / Assam, India



Torsten Blanck / Assam, India




** (star = type locality)

Distribution:* Bangladesh, Bhutan, India (*Arunachal Pradesh, Assam, Manipur, Meghalaya, Nagaland, Tripura*), Myanmar
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 149,274 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 456,213 sq. km
Size (Max SCL):* male 23.3 cm, female 25.0 cm (Baruah et al. 2012; Blanck et al. 2023)
TFTSG Provisional Red List: Vulnerable (VU) (2025)
CITES: Appendix II, as *Cuora* spp. (2000)
Synonymy:*
Cuora praschagi Blanck, Gaillard, Protiva, Wheatley, Shi, Liu, Ray, and Anders 2023:38
Type locality: "Gelabill River, Jorhat District, Assam, India" [= Bhogdoi River] [yellow star].
Type specimen: MHNG 1557.15, holotype.
Etymology:* Eponym honoring Peter Praschag (1974–), Austria; turtle biologist who first recognized the species as distinct.

Other *Cuora* ("*Cistoclemmys*" Species Complex)

East Asian Box Turtles
(includes 12 species, 18 taxa)

Cuora aurocapitata Luo and Zong 1988 (12:22, 17:42)
Yellow-headed Box Turtle
(includes 2 subspecies)



(subspecies: *aurocapitata* = red, *dabieshani* = purple; orange dots = probable trade)

Distribution: China (*Anhui, Henan, Hubei, Zhejiang*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 29,679 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 57,919 sq. km
Size (Max SCL): male 13.5 cm, female 19.5 cm (see subspp.)
IUCN Red List: Critically Endangered (CR A1d+2d [ver. 2.3]) (ATTWG 2000). Previously: Data Deficient (DD) (TFTSG *in* Baillie and Groombridge 1996)
TFTSG Provisional Red List: Critically Endangered (CR) (2011)
CITES: Appendix II, as *Cuora* spp. (2000); zero quota for wild specimens for commercial purposes (2013)

Cuora aurocapitata aurocapitata Luo and Zong 1988 (12:22, 17:42)
Eastern Yellow-headed Box Turtle




Torsten Blanck / Jing Co., Anhui, China





left: John B. Iverson / Anhui, China / trade
right: Gerald Kuchling / TCC / Anhui, China / captivity



(subspecies: *aurocapitata* = red, *dabieshani* = purple; orange dots = probable trade; star = type locality)

Distribution: China (*Anhui, Zhejiang*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 17,640 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 21,137 sq. km

Size (Max SCL): male 12.5 cm, female 16.0 cm (Blanck et al. 2017)

CITES: Appendix II, as *Cuora* spp. (2000); zero quota for wild specimens for commercial purposes (2013)

Synonymy:

Cuora aurocapitata Luo and Zong 1988:13
Cuora pani aurocapitata, Pyxiclemmys aurocapitata, Pyxiclemmys pani aurocapitata, Cuora aurocapitata aurocapitata
Type locality: "Nanling County, Anhui" [China] [yellow star, location estimated].
Type specimen: SNHM 87012, holotype.
Etymology:* Latin, *aurum*, gold, and *capitatus*, with a head; referring to the golden yellow color on the head.

Cuora aurocapitata dabieshani Blanck, Protiva, Zhou, Li, Crow, and Tiedemann 2017 (17:42)
Western Yellow-headed Box Turtle




Torsten Blanck / Anhui, China / captivity





Torsten Blanck / Anhui, China / captivity



(subspecies: *aurocapitata* = red, *dabieshani* = purple; orange dots = probable trade; star = type locality)

Distribution: China (*Anhui, Henan, Hubei*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 12,039 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 15,922 sq. km

Size (Max SCL): male 13.5 cm, female 19.5 cm (Blanck et al. 2017)

CITES: Appendix II, as *Cuora* spp. (2000); zero quota for wild specimens for commercial purposes (2013)


Synonymy:

Cuora aurocapitata dabieshani Blanck, Protiva, Zhou, Li, Crow, and Tiedemann 2017:382
Type locality: "Anhui province, China" [yellow star, location estimated].
Type specimen: NMW 32987:2, holotype, see Gemel et al. (2019).
Etymology:* Toponym for Dabie Shan Mountains, Anhui, China.
Comment: The type specimen, donated by Weissinger, noted by Gemel et al. (2019) to originally have been obtained from an animal importer in Austria.


Cuora bourreti Obst and Reimann 1994 (07:35, 09:22, 12:22, 21:68) (65)*
Bourret's Box Turtle



Torsten Blanck / TCC / Vietnam



Thong Van Pham / Bach Ma, Vietnam



** (stars = type localities)

Distribution: Laos, Vietnam
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 36,815 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 60,101 sq. km
Size (Max SCL): male 19.4 cm, female 20.8 cm (C. Hagen, unpubl. data)
IUCN Red List: Critically Endangered (CR A2bd+4bd) (McCormack and Stuart 2020): Previously: Critically Endangered (CR A2bd+4bd) (McCormack and Stuart 2016); Critically Endangered (CR), as part of *Cuora galbinifrons* (ATTWG 2000)
CITES: Appendix I (2019). Previously: Appendix II, as *Cuora* spp. (2000)
Synonymy:
Cuora galbinifrons serrata Iverson and McCord 1992b:434 (*partim*, hybrid) (07:33)
Type locality:* "100 km east of Tungfang at Tainhfien in central Hainan Island, China" Emended to "100 km east of Dongfang at Tan Hsien (= Danzhou), Hainan"] [orange star, location estimated] by Shi et al. 2005:377.
Type specimen: UF 81791, holotype.
Etymology:* Latin, *serratus*, saw-shaped, serrated; referring to the posterior carapacial margin.
Comment:* Type specimen likely acquired from the trade; type locality likely a trade location. Apparent hybrid of *Cuora bourreti*, *Cuora galbinifrons*, and *Cuora mouhotii mouhotii*.
Cuora galbinifrons bourreti Obst and Reimann 1994:135
Cistoclemmys galbinifrons bourreti, Cuora bourreti, Cistoclemmys bourreti
Type locality: "Linh-Cam (heute Ha-Tinh), in Mittel-Vietnam." Emended to "Bach Ma, Thua Thien Hue Province, Central Vietnam" [yellow star] by Fritz et al. (2002:71).
Type specimen:* MNHN 1948.37 (formerly VNUH T.60), holotype, see also Bour (2005a); possibly lost according to Uetz et al. (2019), but since relocated.
Etymology:* Eponym honoring René L. Bourret (1884–1957), France / Vietnam; naturalist and herpetologist who worked in French Indochina (now Vietnam).

Cuora cyclornata Blanck, McCord, and Le 2006a (07:36, 09:23, 12:22, 17:40, 17:42)
Vietnamese Three-striped Box Turtle
(includes 3 subspecies)



(subspecies: *cyclornata* = red, *annamitica* = purple, *meieri* = blue; orange dots = trade)


Distribution: China (*Guangxi*), Laos, Vietnam
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 124,545 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 337,326 sq. km
Size (Max SCL): male 26.0 cm, female 35.0 cm (see subspp.)
IUCN Red List: Critically Endangered (CR A1d+2d [ver. 2.3]) (ATTWG 2000), as part of *Cuora trifasciata*. Previously: Endangered (EN) (TFTSG 1996), as part of *C. trifasciata*.
TFTSG Provisional Red List: Critically Endangered (CR) (2016)
CITES: Appendix II, as *Cuora* spp. (2000); zero quota for wild specimens for commercial purposes (2013)

Cuora cyclornata cyclornata Blanck, McCord, and Le 2006a (07:36, 09:23, 12:22, 17:40, 17:42)

Southern Vietnamese Three-striped Box Turtle



Torsten Blanck / Vietnam / captivity



(subspecies: *cyclornata* = red, *annamitica* = purple; star = type locality)

Distribution: Laos, Vietnam

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 42,187 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 91,066 sq. km

Size (Max SCL): male 26.0 cm, female 35.0 cm (Blanck et al. 2017)

CITES: Appendix II, as *Cuora* spp. (2000); zero quota for wild specimens for commercial purposes (2013)

Synonymy:

Cuora cyclornata cyclornata Blanck, McCord, and Le 2006a:58 (07:36, 09:23, 12:22)


Type locality: "Phong Nha Ke Bang Nat. Res., Quang Binh Province, central Vietnam" [yellow star].
Type specimen: ZFMK 71348, holotype; genotyped by Blanck et al. (2006a) and Tiedemann et al. (2014).
Etymology:* Greek, *kyklos*, circular, and Latin, *ornata*, adorned, ornate, decorated; referring to the rounded carapace and attractive color.

Cuora cyclornata annamitica Blanck, Protiva, Zhou, Li, Crow, and Tiedemann 2017 (17:42)

Central Vietnamese Three-striped Box Turtle



Torsten Blanck / Vietnam / captivity



(subspecies: *cyclornata* = red, *annamitica* = purple, *meieri* = blue; star = type locality)

Distribution: Vietnam

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 30,501 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 59,098 sq. km

Size (Max SCL): male 26.0 cm, female 35.0 cm (Blanck et al. 2017)

CITES: Appendix II, as *Cuora* spp. (2000); zero quota for wild specimens for commercial purposes (2013)

Synonymy:

Cuora cyclornata annamitica Blanck, Protiva, Zhou, Li, Crow, and Tiedemann 2017:378


Type locality: "Vietnam, Nghe An Province, Tan Ky district, near Ky Son Village" [yellow star].
Type specimen: VUM R.Em.04, holotype.
Etymology:* Toponym for Annam (= Vietnam).

Cuora cyclornata meieri Blanck, McCord, and Le 2006a (07:36, 09:23, 12:22, 17:40, 17:42)

Northern Vietnamese Three-striped Box Turtle



Torsten Blanck / Vietnam / captivity



(subspecies: *annamitica* = purple, *meieri* = blue; orange dot = trade; star = type locality)

Distribution: China (*Guangxi*), Vietnam

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 51,857 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 76,694 sq. km

Size (Max SCL):* male 24.0 cm, female 32.2 cm (Blanck et al. 2017; T. Blanck, unpubl. data)

CITES: Appendix II, as *Cuora* spp. (2000); zero quota for wild specimens for commercial purposes (2013)

Synonymy:

Cuora cyclornata meieri Blanck, McCord, and Le 2006a:73 (07:36, 09:23, 12:22)

Type locality: "Tam Dao, Provinz Vinh Phuc, Nord-Vietnam" [yellow star].

Type specimen: MHNT CHE 1992-11, holotype; genotyped by Blanck et al. (2006a).

Etymology:* Eponym honoring Elmar Meier (1957–), Germany; turtle breeder and conservationist focused on the genus *Cuora*.

Cuora flavomarginata (Gray 1863e) (08:21, 11:9, 12:22, 17:43) (63)*

Yellow-margined Box Turtle

(includes 2 subspecies)



(subspecies: *flavomarginata* = red, *evelynae* = purple (Ryukyus); orange dots = introduced or trade)

Distribution: China (*Anhui, Fujian, Henan, Hubei, Hunan, Jiangsu, Jiangxi, Sichuan, Zhejiang*), Japan (*Ryukyu Archipelago*), Taiwan

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 383,782 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 1,279,192 sq. km

Size (Max SCL): male 18.9 cm, female 19.0 cm (see subspp.)

CBFTT Account: Ota, Yasukawa, Fu, and Chen (2009)

IUCN Red List: Endangered (EN A1cd+2cd [ver. 2.3]) (ATTWG 2000). Previously: Vulnerable (VU) (TFTSG *in* Baillie and Groombridge 1996)

TFTSG Provisional Red List: Critically Endangered (CR) (2011)

CITES: Appendix II, as *Cuora* spp. (2000); zero quota for wild specimens for commercial purposes (2013)

Cuora flavomarginata flavomarginata (Gray 1863e) [(63)*]
Yellow-margined Box Turtle


Tien-Hsi Chen / CBFTT / Nantou, Taiwan


Tien-Hsi Chen / CBFTT / Keelung, Taiwan


(subspecies: *flavomarginata* = red; orange dots = introduced or trade; stars = type localities)

Distribution: China (*Anhui, Fujian, Henan, Hubei, Hunan, Jiangsu, Jiangxi, Zhejiang*), Taiwan

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 383,114 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 1,171,064 sq. km

Size (Max SCL): male 18.9 cm, female 19.0 cm (McCord and Iverson 1991; Ota et al. 2009 CBFTT)

CITES: Appendix II, as *Cuora* spp. (2000); zero quota for wild specimens for commercial purposes (2013)

Synonymy:

Cistoclemmys flavomarginata Gray 1863e:175 [(63)*]
Cuora flavomarginata, Terrapene flavomarginata, Cyclemys flavomarginata, Cyclemys flavomarginata flavomarginata, Cuora flavomarginata flavomarginata, Geoemyda flavomarginata, Cistoclemmys flavomarginatus, Cistoclemmys flavomarginata flavomarginata
Type locality: "China; Formosa...district of Tamsuy, N.W. Formosa" [Taiwan]. Restricted to "Tamsui, Formosa" [Taiwan] [yellow star] by Stejneger (1907:503).
Type specimens: NHMUK 1947.3.5.50, 1947.3.5.68 (formerly 1863.2.23.8), syntypes (2).
Etymology:* Latin, *flavus*, yellow, *marginis*, edge or border, and *-atus*, provided with; referring to the color of the marginal scutes.

Cystoclemmys flavomarginata Gray 1870c:20
Comment:* Unjustified emendation of *Cistoclemmys flavomarginata*.


Cyclemys flavomarginata sinensis Hsü 1930:3 [(07:34, 08:21)]
Cuora flavomarginata sinensis, Cistoclemmys flavomarginata sinensis
Type locality: "Künshan Island, Tungting Lake, Central China." [Junshan Dao, Dongting Lake, Hunan, China] [green star].
Type specimens: MBLSSC 1174–75, syntypes (2).
Etymology:* Toponym for China.

Terrapene culturalia † Yeh 1961:59
Emydoidea culturalia
Type locality: "Dawenkou, Taian, Shantung" [Shandong, China] [gray star].
Type specimen: IVPP V.2520, holotype, subfossil, nearly whole shell, figured (f.1,pl.1).
Geological age: Holocene, Neolithic, Lung-shan Period (Ying Dynasty), subfossil.
Etymology:* Latin, *culturalia*, belonging to culture; referring to the apparent cultural use of this species by Neolithic people.

Cuora flavomarginata evelynae Ernst and Lovich 1990 [(08:21, 11:9)]
Ryukyu Yellow-margined Box Turtle


Iriomote Wildlife Conservation Center / CBFTT / Iriomotejima, Japan


left: Yuichirou Yasukawa / CBFTT / Iriomotejima, Japan
right: Jarmo Perälä / CBFTT / Iriomotejima, Japan


(subspecies: *flavomarginata* = red, *evelynae* = purple; orange dots = introduced or trade; star = type locality)

Distribution: Japan (*Ryukyu Archipelago*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 668 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 2,529 sq. km

Size (Max SCL): male 15.9 cm, female 17.0 cm (Ernst and Lovich 1990; Itescu et al. 2014)

CITES: Appendix II, as *Cuora* spp. (2000); zero quota for wild specimens for commercial purposes (2013)


Synonymy:

Cuora evelynae Ernst and Lovich 1990:31
Cuora flavomarginata evelynae, Cistoclemmys flavomarginata evelynae
Type locality: "Ishigaki Shima, Ryukyu Islands, Japan" [yellow star].
Type specimen: CAS 26113, holotype.
Etymology:* Eponym honoring Evelyn M. Ernst (1948–), USA; female educator and parasitologist and the senior author's wife.

Cuora galbinifrons Bourret 1940 (07:35, 09:22, 12:22, 12:24, 21:68) (65)*
Indochinese Box Turtle


Douglas B. Hendrie / TCF / TCC / Vietnam


left: Archive of the Museum of Zoology Senckenberg Dresden / Tam Dao, Vietnam
right: Edgar Lehr / Vietnam



(stars = type localities)


Distribution: China (*Guangxi*, *Hainan*), Laos, Vietnam
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 127,569 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 296,925 sq. km
Size (Max SCL): male 18.6 cm, female 20.4 cm (Itescu et al. 2014; J.B. Iverson, unpubl. data)
IUCN Red List: Critically Endangered (CR A2bd+4bd) (Li, McCormack, Rao, Shi, Stuart, and Wang 2020). Previously: Critically Endangered (CR A2bd+4bd) (McCormack, Shi, and Stuart 2016); Critically Endangered (CR) (ATTWG 2000); Near Threatened (NT) (TFTSG *in* Baillie and Groombridge 1996)
CITES:* Appendix I (2023). Previously: Appendix II, as *Cuora* spp. (2000); zero quota for wild specimens for commercial purposes (2013)
Synonymy:
Cuora galbinifrons Bourret 1940:11
Cistoclemmys galbinifrons, Cuora galbinifrons galbinifrons, Cistoclemmys galbinifrons galbinifrons
Type locality: "Tam-Dao..[&]..Bach-Ma (Annam)..[&]..Linh-Cam (Ha-Tinh, Nord Annam)" [Vietnam]. Restricted to "Tam-Dao, Nord-Vietnam" [yellow star] by Obst and Reimann (1994:135).
Type specimen: MNHN 1948.36 (formerly VNUH T.54), lectotype, designated by Obst and Reimann (1994:135), see also Bour (2005a); Uetz et al. (2019) erroneously listed MNHN 1948.36 and 1948.37 as syntypes.
Etymology:* Latin, *galbinum*, green-yellow, and *frons*, forehead, the front; referring to the color on top of the head.
Cyclemus flavomarginata hainanensis Li 1958:234
Cuora hainanensis, Cyclemys flavomarginata hainanensis, Cistoclemmys hainanensis, Cuora flavomarginata hainanensis, Cistoclemmys flavomarginata hainanensis, Cuora galbinifrons hainanensis, Cyclemys flavomarginatus hainanensis, Cistoclemmys galbinifrons hainanensis
Type locality: "Dali village, Mt. Diaoluo, Linshui County, Hainan Island, China" [in Chinese] [green star].
Type specimen: FU 200, holotype, CIB 64III6110 (formerly SBRI), allotype, see Koshikawa (1982).
Etymology:* Toponym for Hainan, Chaina.
Cuora galbinifrons serrata Iverson and McCord 1992b:434 (*partim*, hybrid) (07:33)
Type locality:* "100 km east of Tungfang at Tainhfien in central Hainan Island, China" Emended to "100 km east of Dongfang at Tan Hsien (= Danzhou), Hainan"] [orange star, location estimated] by Shi et al. 2005:377.
Type specimen: UF 81791, holotype.
Etymology:* Latin, *serratus*, saw-shaped, serrated; referring to the posterior carapacial margin.
Comment:* Type specimen likely acquired from the trade; type locality likely a trade location. Apparent hybrid of *Cuora bourreti*, *Cuora galbinifrons*, and *Cuora mouhotii mouhotii*

Cuora mccordi Ernst 1988 (12:22)
McCord's Box Turtle




Torsten Blanck / TCC / Guangxi, China / captivity





left: John B. Iverson / Guangxi, China / trade
right: Torsten Blanck / Guangxi, China / captivity



(red = possible range; orange dots = trade; stars = type localities)

Distribution: China (*Guangxi*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 25,056 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 34,964 sq. km

Size (Max SCL): male 18.4 cm, female 22.5 cm (Ceballos et al. 2013; T. Blanck, unpubl. data)

IUCN Red List: Critically Endangered (CR A1d+2d [ver. 2.3]) (ATTWG 2000). Previously: Data Deficient (DD) (TFTSG *in* Baillie and Groombridge 1996)


TFTSG Provisional Red List: Critically Endangered (CR) (2011)

CITES: Appendix II, as *Cuora* spp. (2000); zero quota for wild specimens for commercial purposes (2013)

Synonymy:

Cuora mccordi Ernst 1988:466
Cistoclemmys mccordi
Type locality: "highland near Paise, Guangxi Province, China (23º54'N, 106º37'E)" [in error; orange star]. Emended to "Yunnan Province, west of Paise, Guangxi Province, China" [in error, location estimated, yellow star] by McCord and Iverson (1991:414).
Type specimen: USNM 281850, holotype, see Reynolds et al. (2007).
Etymology:* Eponym honoring William P. McCord (1950–), USA; veterinarian and herpetologist, who obtained specimens of this species from the trade and first recognized them as distinct.
Comment:* Type specimen likely acquired from the trade; type locality likely a trade location.

Cuora mouhotii (Gray 1862a) (07:32, 12:22)
Keeled Box Turtle
(includes 2 subspecies)



** (subspecies: *mouhotii* = red, *obsti* = blue; overlap = intergrades, orange dots = probable trade)

Distribution: Bangladesh, Bhutan, China (*Guangxi*, *Hainan*, *Yunnan*), India (*Arunachal Pradesh*, *Assam*, *Manipur*, *Meghalaya*, *Mizoram*, *Nagaland*), Laos, Myanmar, Thailand (?), Vietnam

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 580,625 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 2,317,440 sq. km

Size (Max SCL): male 25.7 cm, female 23.7 cm (see subspp.)

CBFTT Account: Das, McCormack, van Dijk, Hoang, and Struijk (2016)


IUCN Red List: Endangered (EN A2cd) (Ahmed, Horne, Li, Platt, Rahman, and Wang 2020). Previously: Endangered (EN) (ATTWG 2000); Least Concern (LC) [Not Listed] (Behler 1996)

CITES: Appendix II, as *Cuora* spp. (2000); zero quota for wild specimens for commercial purposes (2013)

Cuora mouhotii mouhotii (Gray 1862a)
Northern Keeled Box Turtle



Indraneil Das / CBFTT / China / captivity



Timothy E.M. McCormack / CBFTT / Cuc Phuong National Park, Vietnam / female, hatchling



(subspecies: *mouhotii* = red, *obsti* = blue; overlap = intergrades, orange dots = probable trade; stars = type localities)

Distribution: Bangladesh, Bhutan, China (*Guangxi*, *Hainan*, *Yunnan*), India (*Arunachal Pradesh*, *Assam*, *Manipur*, *Meghalaya*, *Mizoram*, *Nagaland*), Laos, Myanmar, Thailand (?), Vietnam

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 527,129 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 1,819,248 sq. km

Size (Max SCL): male 25.7 cm, female 23.7 cm (Das et al. 2016 CBFTT; T. Blanck, unpubl. data)

CITES: Appendix II, as *Cuora* spp. (2000); zero quota for wild specimens for commercial purposes (2013)

Synonymy:

Cyclemys mouhotii Gray 1862a:157

Pyxidea mouhotii, Cyclemys mohouti, Pyxidea mouboti, Emys mouhotii, Geoemyda mouhotii, Pyxidea mouhotii mouhotii, Cuora mouhotii, Cuora mouhotii mouhotii

Type locality: "Lao Mountains, in Siam." Restricted to "Luang Prabang, Laos (19º54' N, 102º8' O)" [yellow star] by lectotype designation by Fritz et al. (1998:40).

Type specimen: NHMUK 1947.3.4.27, lectotype, designated by Fritz et al. (1998a:40), discussed by Das (2009); Uetz et al. (2019) erroneously listed NHMUK 1947.3.4.27, 1947.3.4.48–49, and 1947.3.4.64–67 as syntypes.

Etymology:* Eponym honoring Henri Mouhot (1826–1861), France; naturalist and explorer in Thailand and Laos who collected the types.

Cuora galbinifrons serrata Iverson and McCord 1992b:434 (*partim*, hybrid) (07:33)

Type locality:* "100 km east of Tungfang at Tainhfien in central Hainan Island, China" Emended to "100 km east of Dongfang at Tan Hsien (= Danzhou), Hainan"] [orange star, location estimated] by Shi et al. 2005:377.

Type specimen: UF 81791, holotype.


Etymology:* Latin, *serratus*, saw-shaped, serrated; referring to the posterior carapacial margin.

Comment:* Type specimen likely acquired from the trade; type locality likely a trade location. Apparent hybrid of *Cuora bourreti*, *Cuora galbinifrons*, and *Cuora mouhotii mouhotii*


Cuora mouhotii obsti (Fritz, Andreas, and Lehr 1998a)
Southern Keeled Box Turtle



Torsten Blanck / Phu Yen Prov., Vietnam / male



left: Nguyen Thanh Luan / CBFTT / Quang Tri, Vietnam / trade / female
right: Richard Struijk / CBFTT / southern Vietnam / hatchling



(subspecies: *mouhotii* = red, *obsti* = blue; overlap = intergrades; orange dots = probable trade; star = type locality)

Distribution: Laos, Vietnam

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 63,806 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 179,784 sq. km

Size (Max SCL): male 18.0 cm, female 16.0 cm (Fritz et al. 1998a)

CITES: Appendix II, as *Cuora* spp. (2000); zero quota for wild specimens for commercial purposes (2013)

Synonymy:

Pyxidea mouhotii obsti Fritz, Andreas, and Lehr 1998a:35

Cuora mouhotii obsti

Type locality: "Umgebung von Phú Lôc, Annam (Vietnam), 16º16' N, 107º56' O)" [yellow star].

Type specimen: MTD 31458, holotype.

Etymology:* Eponym honoring Fritz-Jürgen Obst (1939–2018), Germany; herpetologist and director of the Staatliches Museum für Tierkunde, Dresden.

Cuora pani Song 1984 (12:22) (67)*
Pan's Box Turtle



Torsten Blanck / TCC / China / captivity





left: John B. Iverson / China / trade
right: Torsten Blanck / China / captivity



** (orange dots = probable trade; stars = type localities)

Distribution: China (*Henan, Hubei, Shaanxi, Sichuan*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 77,051 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 179,341 sq. km

Size (Max SCL): male 14.5 cm, female 19.5 cm (Song 1984; Ernst and McCord 1987; T. Blanck, unpubl. data)

IUCN Red List: Critically Endangered (CR A1d+2d [ver. 2.3]) (ATTWG 2000). Previously: Data Deficient (DD) (TFTSG *in* Baillie and Groombridge 1996)

TFTSG Provisional Red List: Critically Endangered (CR) (2011)

CITES: Appendix II, as *Cuora* spp. (2000); zero quota for wild specimens for commercial purposes (2013)

Synonymy:

Cuora pani Song 1984:330
Cuora pani pani, Pyxiclemmys pani pani
Type locality: "Xujiaba (alt. 420 m) of Pingli County in Shaanxi Province" [China] [yellow star].
Type specimen:* SIZ 80170, holotype, figured (f.1) and in Song (2001:f.1).
Etymology:* Eponym honoring Liu Pan (1900s), China; herpetologist at Shanghai Museum of Natural History, Shanghai.

Cuora chriskarannarum Ernst and McCord 1987:624
Type locality: "Ta Lau Shan, Yunnan Province, China (23º30' N, 102º25' E)" [in error?] [orange star].
Type specimen: USNM 266162, holotype, see Reynolds et al. (2007).
Etymology:* Eponym honoring Christine, Karen, and Anne McCord (1978–, 1979–, and 1950–), USA; the junior author's two daughters and wife.
Comment:* Type specimen likely acquired from the trade; type locality likely a trade location.

Cuora picturata Lehr, Fritz, and Obst 1998 (07:35, 09:22, 12:22, 21:68) (65)*
Southern Vietnam Box Turtle



Edgar Lehr / Vietnam / trade





Edgar Lehr / Vietnam / trade



• (orange dot = trade; star = type locality)

Distribution: Vietnam

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 3,541 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 5,774 sq. km

Size (Max SCL): male 18.9 cm, female 19.2 cm (Itescu et al. 2014; T. Blanck, unpubl. data)


IUCN Red List: Critically Endangered (CR A2bd+4bd) (McCormack, Stuart, Blanck 2020). Previously: Critically Endangered (CR A2bd+4bd) (McCormack, Stuart, Blanck 2016); Critically Endangered (CR), as part of *Cuora galbinifrons* (ATTWG 2000)

CITES: Appendix I (2019). Previously: Appendix II, as *Cuora* spp. (2000)

Synonymy:

Cuora galbinifrons picturata Lehr, Fritz, and Obst 1998:7
Cistoclemmys galbinifrons picturata, Cuora picturata, Cistoclemmys picturata
Type locality: "südliches Annam (Vietnam)" [yellow star, location estimated].
Type specimen: MTD 34565, holotype.
Etymology:* Latin, *picturata*, embroidered; referring to the soft parts having a fine web-like pattern similar to bead-work.

Cuora trifasciata (Bell 1825a) (07:36, 09:23, 12:22, 17:40, 17:42)
Chinese Three-striped Box Turtle, Golden Coin Turtle
(includes 2 subspecies)



(subspecies: *trifasciata* = red, *luteocephala* = purple; orange dots = trade)

Distribution: China (*Fujian, Guangdong, Guangxi, Hainan, Hong Kong, Macau*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 241,670 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 452,311 sq. km

Size (Max SCL): male 23.0 cm, female 26.0 cm (see subspp.)

IUCN Red List: Critically Endangered (CR A2d) (Fong, Hoang, Horne, Li, McCormack, Rao, Timmins, and Wang 2020). Previously: Critically Endangered (CR) (ATTWG 2000); Endangered (EN) (TFTSG *in* Baillie and Groombridge 1996)


CITES: Appendix II, as *Cuora* spp. (2000); zero quota for wild specimens for commercial purposes (2013)

Cuora trifasciata trifasciata (Bell 1825a) (07:36, 09:23, 12:22, 17:40, 17:42)
Chinese Three-striped Box Turtle




Paul Crow / TCF / TCC / Hong Kong, China / captivity





left: Eric V. Goode / China / captivity
right: Michael Lau / Hong Kong, China



(subspecies: *trifasciata* = red, *luteocephala* = purple; orange dots = trade; stars = type localities)

Distribution: China (*Fujian, Guangdong, Guangxi, Hong Kong, Macau*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 225,935 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 302,216 sq. km

Size (Max SCL): male 21.0 cm, female 23.0 cm (Blanck et al. 2017)

CITES: Appendix II, as *Cuora* spp. (2000); zero quota for wild specimens for commercial purposes (2013)

Synonymy:

Sternothaerus trifasciatus Bell 1825a:305
Emys (Cistuda) trifasciata, Emys trifasciata, Cistuda trifasciata, Cistudo trifasciata, Cuora trifasciata, Pyxidemys trifasciata, Terrapene trifasciata, Cyclemys trifasciata, Cuora (Pyxiclemmys) trifasciata, Pyxiclemmys trifasciata, Cuora trifasciata trifasciata
Type locality: Not known. Restricted to "Luofo Shan Mountains, Guangdong, China" [yellow star] by Blanck et al. (2006a:40).
Type specimen: OUM 8557, holotype, discussed by Blanck et al. (2006a) and Nowak-Kemp and Fritz (2010); genotyped by Blanck et al. (2006a) and Tiedemann et al. (2014).
Etymology:* Latin, *tri-*, three, and *fasciatus*, banded; referring to the three longitudinal, dark stripes on the carapace.

Mauremys iversoni Pritchard and McCord 1991:140 (*partim*, hybrid) (07:33)
Type locality: "People's Republic of China: Fujian Province: vicinity of Nanping (26°38' N, 118°10' E)" [in error?] [orange star].
Type specimen: UF 71865, holotype.
Etymology:* Eponym honoring John B. Iverson (1949–), USA; herpetologist and turtle researcher at Earlham College, Indiana.
Comment:* Type specimen likely acquired from the trade; type locality likely a trade location. Apparent hybrid of *Cuora trifasciata trifasciata* and *Mauremys mutica mutica*. Previously listed as Data Deficient (DD) on the IUCN Red List (TFTSG *in* Baillie and Groombridge 1996).

Clemmys guangxiensis Qin 1992:60 (*partim*, hybrid) (14:29)
Type locality: "Nanning, Guangxi" [China] [in error?] [brown star].
Type specimen: SNHM 88701, holotype.
Etymology:* Toponym for Guangxi, China.
Comment:* Type specimen likely acquired from the trade; type locality likely a trade location. Apparent hybrid of *Cuora trifasciata trifasciata*, *Mauremys mutica mutica*, and possibly *Mauremys annamensis*.

Cuora trifasciata luteocephala Blanck, Protiva, Zhou, Li, Crow, and Tiedemann 2017 (17:42)

Hainan Three-striped Box Turtle




Torsten Blanck / Hainan, China / captivity





Scott D. Gillingwater / Hainan, China / captivity / adult female, hatchling



(subspecies: *trifasciata* = red, *luteocephala* = purple; orange dots = trade; stars = type localities)

Distribution: China (*Hainan*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO): 15,734 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 19,035 sq. km

Size (Max SCL): male 23.0 cm, female 26.0 cm (Blanck et al. 2017)

CITES: Appendix II, as *Cuora* spp. (2000); zero quota for wild specimens for commercial purposes (2013)

Synonymy:

Sacalia pseudocellata Iverson and McCord 1992a:426 (*partim*, hybrid) (07:33)

Type locality: "between Tungfang [19º03' N, 108º56' E] and Kancheng [18º51' N, 108º37' E; ca. 48 km from Tungfang], western Hainan Island, China" [= Dongfang and Gancheng] [orange star].

Type specimen: UF 81505, holotype.

Etymology:* Greek, *pseudo*, false, and Latin, *ocellata*, eyed; referring to the lack of black eye spots on the neck.

Comment:* Type specimen likely acquired from the trade; type locality likely a trade location. Apparent hybrid of *Cuora trifasciata luteocephala* and *Sacalia quadriocellata*. Previously listed as Data Deficient (DD) on the IUCN Red List (TFTSG *in* Baillie and Groombridge 1996).

Ocadia philippeni McCord and Iverson 1992:13 (*partim*, hybrid) (07:33)

Type locality: "near Kancheng [18º51'N, 108º37'E; = 48 km from Tungfang (19º03'N, 108º56'E)], western Hainan Island, China" [= Gancheng and Dongfang] [brown star].

Type specimen: UF 80766, holotype.

Etymology:* Eponym honoring Hans-Dieter Philippen (1957–2016), Germany; herpetologist and turtle researcher.

Comment:* Type specimen likely acquired from the trade; type locality likely a trade location. Apparent hybrid of *Cuora trifasciata luteocephala* and *Mauremys sinensis*. Previously listed as Data Deficient (DD) on the IUCN Red List (TFTSG *in* Baillie and Groombridge 1996).

Cuora trifasciata luteocephala Blanck, Protiva, Zhou, Li, Crow, and Tiedemann 2017:379

Type locality: "China, Hainan Province, Dan County, 300 m elevation" [= Danzhou] [yellow star].

Type specimen: CIB 64III5279, holotype.


Etymology:* Latin, *lutea*, yellow, and Greek, *kefale*, head; referring to the golden yellow head.

Cuora yunnanensis (Boulenger 1906) (07:37, 12:22)

Yunnan Box Turtle




Ting Zhou, William P. McCord, Torsten Blanck / TCC / Yunnan, China / captivity





left: John B. Iverson / Yunnan, China / holotype [NHMUK, London]
right: Haitao Shi / Yunnan, China / captivity



(red = possible range; dots = possible native or trade; star = type locality)

Distribution: China (*Sichuan*, *Yunnan*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO): 31,947 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 61,978 sq. km

Size (Max SCL): male 15.2 cm, female 19.0 cm (Blanck et al. 2006b; T. Blanck, unpubl. data)

IUCN Red List: Critically Endangered (CR B2ab(ii,iii,v), D) (van Dijk, Blanck, and Lau 2010). Previously: Extinct (EX) (ATTWG 2000); Data Deficient (DD) (TFTSG *in* Baillie and Groombridge 1996)

CITES: Appendix II, as *Cuora* spp. (2000); zero quota for wild specimens for commercial purposes (2013)

Synonymy:


Cyclemys yunnanensis Boulenger 1906a:567

Cuora yunnanensis, *Pyxiclemmys yunnanensis*

Type locality: "Yunnan fu..[&].. Tongchuan fu" [Kunming Shi.. [&].. Dongchuan Shi, Yunnan, China]; restricted to "vicinity of Zhongping (Huize) City (26°42'N, 103°30'E) in Huize County, northeastern Yunnan Province, China (= Tongchuan Fu)" [yellow star] by Blanck et al. (2006b:31).
Type specimen: NHMUK 1946.1.22.97 (formerly 1905.5.30.34), lectotype, designated by Blanck (2005:11).
Etymology:* Toponym for Yunnan, China.

Cuora zhoui Zhao *in* Zhao, Zhou, and Ye 1990 (12:22)
Zhou's Box Turtle


Torsten Blanck / TCC / Guangxi, China? / captivity


John B. Iverson / Guangxi, China? / captivity


(red = possible range; yellow dot = possible native; orange dots = trade; stars = type localities)


Distribution: China (*Guangxi [?], Yunnan [?]*), Vietnam
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 12,983 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 15,593 sq. km
Size (Max SCL): male 17.0, female 22.3 cm (T. Blanck, unpubl. data)
CBFTT Account:* Blanck, Iverson, Zhou, Meier, and McCord (2024)
IUCN Red List: Critically Endangered (CR A1d+2d [ver. 2.3]) (ATTWG 2000). Previously: Data Deficient (DD) (TFTSG *in* Baillie and Groombridge 1996)
TFTSG Provisional Red List: Critically Endangered (CR) (2011)
CITES: Appendix II, as *Cuora* spp. (2000); zero quota for wild specimens for commercial purposes (2013)
Synonymy:
Cuora zhoui Zhao *in* Zhao, Zhou, and Ye 1990:213
Pyxiclemmys zhoui
Type locality: "market of Nanning, Guangxi Zhuang Autonomous Region" [China] [orange star, trade].
Type specimen:* CIB 119803, holotype; previously NTM(P) 9001, holotype; genotyped by Tiedemann et al. (2014).
Etymology:* Eponym honoring Jiufa Zhou (1938–), China; economist, naturalist, and founder of the Nanjing Turtle Museum.
Comment:* Type specimen acquired from the trade; type locality a trade location.
Cuora pallidicephala McCord and Iverson 1991:414
Type locality: "Wuting [= Wuding: 25º26'N, 102º21'W] or Yuanmow [25º41'N, 101º54'W], Yunnan Province, China" [brown stars, trade].
Type specimen: UF 77230, holotype.
Etymology:* Latin, *pallidus*, pale, and Greek, *kefale*, head; referring to the pale head markings.
Comment:* Type specimen acquired from the trade; type locality a trade location.

Cyclemys Bell 1834 (07:38, 08:6, 09:24)
Southeast Asian Leaf Turtles
(includes 7 species, 7 taxa)
Cyclemys Bell 1834:17
Type species: *Cyclemys orbiculata* Bell 1834 [= subjective synonym of *Emys dentata* Gray 1831d], by original designation.
Etymology:* Greek, *kyklos*, circular, and *emys*, turtle; referring to the rounded oval carapace.
Cyclemis Tirant 1884:156 (*nomen novum*)
Etymology:* Greek, *kyklos*, circular, and *emys*, turtle; referring to the rounded oval carapace.
Comment:* Unjustifed emendation or error for *Cyclemys*.

Cyclemys atripons Iverson and McCord 1997b (17:44)
Western Black-bridged Leaf Turtle


Flora Ihlow / Trat, Thailand-Cambodia border


Flora Ihlow / Trat, Thailand-Cambodia border



(orange dots = hybrids with *C. oldhamii*; star = type locality)


Distribution: Cambodia, Thailand, Vietnam
Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 37,284 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 55,743 sq. km
Size (Max SCL): male 19.5 cm, female 23.6 cm (Fritz et al. 2008b; J.B. Iverson, unpubl. data)
IUCN Red List: Endangered (EN A2cd+4cd) (Timmins, Hoang, and McCormack 2021)
CITES: Appendix II, as *Cyclemys* spp. (2013)
Synonymy:
Cyclemys atripons Iverson and McCord 1997b:632
Cyclemys atripons atripons, Cyclemys pulchristriata atripons
Type locality: "Thailand, Krat [= Trat], Kao [= Mt.] Kuap (= Khao Kuap)" [yellow star].
Type specimen: USNM 81865, holotype, see Reynolds et al. (2007); genotyped by Stuart and Fritz (2008).
Etymology:* Latin, *atra*, black, and *pons*, bridge; referring to the black markings on the plastral bridge.

Cyclemys dentata (Gray 1831d) (08:7)
Asian Leaf Turtle


Hynek Prokop / Binunsalian, Palawan, Philippines


left: Hynek Prokop / Palawan, Philippines
right: Maren Gaulke / Kota Pinang, Sumatra, Indonesia



** (blue dots = genetically confirmed; orange dot = erroneous record; stars = type localities)


Distribution: Brunei, Indonesia (*Bali, Java, Kalimantan, Sumatra*), Malaysia (*East, West*), Philippines (*Palawan, Sulu Archipelago*), Singapore, Thailand
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 1,333,998 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 4,433,372 sq. km
Size (Max SCL): male 19.6, female 21.0 cm (Fritz et al. 2008b; J.B. Iverson, unpubl. data)
IUCN Red List: Near Threatened (NT A3cd) (As-singkily, Guntoro, Kusrini, and Schoppe 2021). Previously: Near Threatened (NT) (ATTWG 2000); Least Concern (LC) [Not Listed] (Behler 1996)
CITES: Appendix II, as *Cyclemys* spp. (2013)
Synonymy:
Emys hasseltii Boie *in* Fitzinger 1826:45 (*nomen nudum*)
Clemmys (Clemmys) hasseltii
Type locality: "Asia, Insula Java" [Indonesia] [yellow star, location estimated].
Type specimen: Not known or located.
Etymology:* Eponym honoring Johan Conrad van Hasselt (1797–1823), Netherlands / Indonesia; physician and zoologist who worked on Java in the Dutch East Indies (now Indonesia).
Emys dhor Gray 1830e:8 (*nomen oblitum*) (10:7)
Cyclemys dhor
Type locality: "India." Restricted to "Bengal..[and]..Java" by Gray (1831d:20); and to "Java" [Indonesia] [yellow star, location estimated] by Fritz et al. (1997:188).
Type specimen: NHMUK 1946.1.22.62 (formerly 28.5.12.1), lectotype, designated by Fritz et al. (1997:194), discussed by Nowak-Kemp and Fritz (2010); genotyped by Stuart and Fritz (2008); Uetz et al. (2019) erroneously listed NHMUK 1946.1.22.62 and 1946.1.22.63 as syntypes.
Etymology:* Vernacular (Hindi?), *dhor*, possibly a designation for various regional indigenous tribes in India, or possibly a local indigenous name for the species.
Emys dentata Gray 1831d:errata:btw.78-79 (*nomen novum*)
Cistudo (Cyclemys) dentata, Cistudo dentata, Cyclemys dentata, Cyclemys dentata dentata
Type locality: "Bengal..[and]..Java." Restricted to "Java" [Indonesia] [yellow star, location estimated] by Smith (1931:80) and by Fritz et al. (1997:188).
Type specimen: NHMUK 1946.1.22.62 (formerly 1828.5.12.1), lectotype, designated by Fritz et al. (1997:194), discussed by Nowak-Kemp and Fritz (2010); genotyped by Stuart and Fritz (2008).
Etymology:* Latin, *dens*, tooth, *dentata*, toothed, serrated; referring to the serrated rear marginal scutes in juveniles.
Comment: Accepted replacement name for *dhor*.
Cyclemys orbiculata Bell 1834:17
Emys orbiculata, Emys (Cyclemys) orbiculata, Cistudo orbiculata
Type locality: "Indiâ." Restricted to "Java" [Indonesia] [yellow star, location estimated] by Fritz et al. (1997:188).
Type specimens: OUM 8512–13, 8867, syntypes (3); NHMUK 1946.1.22.62 erroneously designated neotype by Fritz et al. (1997:188), discussed by Nowak-Kemp and Fritz (2010); OUM 8512–13 genotyped by Stuart and Fritz (2008).
Etymology:* Latin, *orbiculus*, small circle, *orbiculata*, circular; referring to the relatively circular shell.
Cistudo diardii Duméril and Bibron 1835:227 (*nomen novum*)
Emys diardii
Type locality:* "Bengale et l'ile de Java." Restricted here to "Java" [Indonesia] [yellow star, location estimated].
Type specimens: MNHN 9104–08, syntypes (5).
Comment: Unjustified replacement name for *dhor*.
Etymology:* Eponym honoring Pierre-Médard Diard (1794–1863), France / Indonesia; naturalist, explorer, and collector who founded the Buitenzorg Botanical Gardens in Java.
Cyclemys ovata Gray 1863e:178
Type locality: "Sarawak" [East Malaysia] [green star, location estimated].
Type specimen: NHMUK 1863.6.21.1, holotype; genotyped by Stuart and Fritz (2008).
Etymology:* Latin, *ovata*, egg-shaped, oval; referring to the rounded oval carapace.
Cyclemys bellii Gray 1863e:179
Type locality: Not known.

Type specimen: OUM 8513, holotype, discussed by Nowak-Kemp and Fritz (2010); OUM 8867 erroneously listed as holotype by Das (2005); OUM 8513 genotyped by Stuart and Fritz (2008).
Etymology:* Eponym honoring Thomas Bell (1792–1880), United Kingdom; herpetologist, dental surgeon, and Professor of Zoology at King's College, London.

Cyclemys enigmatica Fritz, Guicking, Auer, Sommer, Wink, and Hundsdörfer 2008 (08:6)
Enigmatic Leaf Turtle


Maren Gaulke / Rantau Prapat, Sumatra, Indonesia



Maren Gaulke / Rantau Prapat, Sumatra, Indonesia



** (blue dots = genetically confirmed; star = type locality)

Distribution: Malaysia (*West*), Singapore, Indonesia (*Java*, *Kalimantan*, *Sumatra*), Thailand
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 920,084 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 2,440,116 sq. km
Size (Max SCL): female 23.5 cm (Fritz et al. 2008b)
IUCN Red List: Near Threatened (NT A3cd) (As-singkily, Guntoro, Kusrini, and Hamidy 2021)
CITES: Appendix II, as *Cyclemys* spp. (2013)
Synonymy:
Cyclemys enigmatica Fritz, Guicking, Auer, Sommer, Wink, and Hundsdörfer 2008b:381
Type locality: "Padang, Sumatra" [Indonesia] [yellow star].
Type specimen: NMW 9811, holotype, see Gemel et al. (2019).
Etymology:* Greek, *enigma*, riddle, mystery, *enigmatica*, enigmatic; referring to confusion about the identity of this cryptic species.

Cyclemys fusca Fritz, Guicking, Auer, Sommer, Wink, and Hundsdörfer 2008 (08:6)
Myanmar Brown Leaf Turtle


Peter Paul van Dijk / Myanmar / trade


Peter Paul van Dijk / Myanmar / trade


** (orange dots = translocated; star = type locality)

Distribution: India (*Nagaland*, *Manipur*), Myanmar
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 141,788 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 308,421 sq. km
Size (Max SCL): male 22.5 cm, female 26.9 cm (J.B. Iverson, unpubl. data)
IUCN Red List: Least Concern (LC) (Platt, Praschag, and Horne 2021)
CITES: Appendix II, as *Cyclemys* spp. (2013)
Synonymy:
Cyclemys fusca Fritz, Guicking, Auer, Sommer, Wink, and Hundsdörfer 2008b:383
Type locality: "Kachin State, Myanmar" [yellow star, location estimated].
Type specimen: MTD 42578, holotype.
Etymology:* Latin, *fusca*, dark brown, dark, gray, black; referring to the carapace color.


Cyclemys gemeli Fritz, Guicking, Auer, Sommer, Wink, and Hundsdörfer 2008 (08:6)
Assam Leaf Turtle



Peter Praschag / Dimapur, Nagaland, India



Peter Praschag / Dimapur, Nagaland, India



(star = type locality)

Distribution: Bangladesh, Bhutan, India (*Arunachal Pradesh, Assam, Bihar, Manipur, Meghalaya, Mizoram, Nagaland, Uttar Pradesh*), Myanmar, Nepal

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 273,105 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 974,256 sq. km

Size (Max SCL): male 19.0 cm, female 23.2 cm (Praschag et al. 2009b; Baruah et al. 2012)

IUCN Red List: Near Threatened (NT A4c) (Praschag and Ahmed 2021)

CITES: Appendix II, as *Cyclemys* spp. (2013)

Synonymy:

Cyclemys gemeli Fritz, Guicking, Auer, Sommer, Wink, and Hundsdörfer 2008b:384

Type locality: "street from Tezpur to Arunachal Pradesh, 5 km to border of Arunachal Pradesh, Jia Bhoroli River Region, Assam, India" [yellow star].

Type specimen: NMW 37153, holotype, see Gemel et al. (2019); genotyped by Fritz et al. (2008b).


Etymology:* Eponym honoring Richard Gemel (1948–), Austria; herpetologist who collected the type.

Cyclemys oldhamii Gray 1863e (08:8, 17:44)
Southeast Asian Leaf Turtle




Thomas Ziegler / NW of Ky Thuong, Vietnam





Peter Praschag / No data / captive



(orange dots = probable trade; stars = type localities)

Distribution:* Cambodia, China (*Yunnan*), Laos, Myanmar, Thailand, Vietnam

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 634,619 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 1,416,910 sq. km

Size (Max SCL): male 24.1 cm, female 26.0 cm (Fritz et al. 2008b; Seateun et al. 2019)

IUCN Red List: Endangered (EN A2cd+4cd) (Timmins, Hoang, and McCormack 2021)

CITES: Appendix II, as *Cyclemys* spp. (2013)

Synonymy:

Cyclemys oldhamii Gray 1863e:178

Cyclemis oldhami

Type locality:* "Mergui..[&]..Siam." Restricted to "Mergui" [Myanmar] [= Myeik, Tanintharyi Region] [yellow star] by Smith (1931:81) and by Fritz et al. (1997:196).

Type specimen: NHMUK 1947.3.5.63 (formerly 1856.5.6.1), lectotype, designated by Fritz et al. (1997:201); genotyped by Stuart and Fritz (2008).

Etymology:* Eponym honoring Thomas Oldham (1816–1878), Ireland / India; geologist and naturalist at the Geological Survey of

India who collected turtles and mammals for the British Museum. The entry for Oldham in Beolens et al. (2011) is incorrect.

Cyclemys dhor shanensis Annandale 1918:67

Cyclemys shanensis, Cyclemys shanensis shanensis

Type locality:* "Fort Stedman on the Inlé Lake, altitude 3,000 feet..[&].. He-Ho plain 800 feet higher" [Myanmar]. Restricted to "Fort Stedman (3000 Fuß Höhe) am Inlé-See, Birma" [Myanmar] by lectotype designation by Fritz et al. (1997:196) [green star].

Type specimen: ZSI 18594, lectotype, designated by Fritz et al. (1997:196); ZSI 18593–94 listed as syntypes by Das et al. (1998) and Kundu et al. (2018a).

Etymology:* Toponym for Shan State, Burma (= Myanmar).

Geoemyda tcheponensis Bourret 1939:7

Cyclemys tcheponensis, Cyclemys dentata tcheponensis, Cyclemys shanensis tcheponensis

Type locality:* "Haute Sé-Bang-Hien, (centre de la Chaîne annamitique)" [Upper Banghiang River, Seponh District, Laos] [blue star].

Type specimen:* CRES T.43 (formerly VNUH T.43), holotype, previously assumed to be lost (Fritz et al. 1997; TTWG 2017, 2021, Iverson 2022a) but actually rediscovered by Fritz et al. (1999) in a previously overlooked publication.

Etymology:* Toponym for Tchépone District in French Indochina (= Seponh District, Savannakhet Province, Laos).

Cyclemys tiannanensis Kou 1989:193

Type locality: "Nanliang, Mengla County, Xishuangbanna of Yunnan Province, alt. 620 m" [China] [pink star].

Type specimen: YU RT8311002, holotype.

Etymology:* Toponym for Tiannan [in old Pinyin] (= Diannan [in new Pinyin] = Southern Yunnan, China).


Comment: Previously recorded as probable market specimen, according to Fritz et al. (1997), but confirmed as native occurrence (Wang et al. 2022; K. Wang, unpubl. data).

Cyclemys pulchristriata Fritz, Gaulke, and Lehr 1997 (17:45)


Eastern Black-bridged Leaf Turtle



Edgar Lehr / Vietnam / trade



left: Thong Van Pham / Vietnam
right: Edgar Lehr / Vietnam / trade



** (star = type locality)

Distribution: Cambodia, Vietnam

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 70,362 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 112,295 sq. km

Size (Max SCL): male 17. 5 cm, female 22.7 cm (Fritz et al. 2008b)

IUCN Red List: Endangered (EN A2cd+4cd) (Timmins, Hoang, and McCormack 2021)

CITES: Appendix II, as *Cyclemys* spp. (2013)

Synonymy:

Cyclemys pulchristriata Fritz, Gaulke, and Lehr 1997:203

Cyclemys atripons pulchristriata

Type locality: "Phuc-Son, Annam" [Quang Nam, Vietnam] [yellow star].

Type specimen: NMW 29525:4, holotype, see Gemel et al. (2019); genotyped by Stuart and Fritz (2008).

Etymology:* Latin, *pulchra*, beautiful, and *striata*, striped; referring to the striking coloration compared to other *Cyclemys* species.

Geoemyda Gray 1834b (07:39)

East Asian Leaf Turtles

(includes 2 species, 2 taxa)

Geoemyda Gray 1834b:100 (*nomen protectum*)

Type species: *Geoemyda spengleri* [= *Testudo spengleri* Gmelin 1789], by original designation.

Etymology:* Greek, *gaia*, *geo-*, the earth or land, and *emydos*, turtle.

Comment: Name conserved by ICZN (1985a), see Smith et al. (1980c).

Geoemys Bonaparte 1836:6 (*nomen novum*)

Etymology:* Greek, *gaia*, *geo-*. the earth or land, and *emys*, turtle.

Comment:* Invalid emendation or error for *Geoemyda*.

Geomyda Blyth 1856:714 (*nomen novum*)

Etymology:* Greek, *gaia*, *geo-*, the earth or land, and *emydos*, turtle.

Comment:* Invalid emendation or error for *Geoemyda*.

Nicoria Gray 1856b:17

Type species: *Nicoria spengleri* [= *Testudo spengleri* Gmelin 1789], by original monotypy.

Etymology:* Mythonym for *nicor*, an Old English name for a water monster or water goblin.


Geoemyda japonica Fan 1931
Ryukyu Black-breasted Leaf Turtle



Maximilian Maurer and Simon Rouot / Okinawajima, Japan



Maximilian Maurer and Simon Rouot / Okinawajima, Japan



(orange dots = introduced; star = type locality)

Distribution: Japan (*Ryukyu Archipelago*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 1,613 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 8,653 sq. km

Size (Max SCL):* male 16.2 cm, female 17.0 cm (Vetter and van Dijk 2006; Itescu et al. 2014; Rouot and Maurer 2023; H. Ota, unpubl. data)

CBFTT Account: Yasukawa and Ota (2008)

IUCN Red List: Endangered (EN A1ce, B1+2c [ver. 2.3]) (ATTWG 2000). Previously: Endangered (EN) (TFTSG *in* Baillie and Groombridge 1996)

CITES: Appendix II (2013)

Synonymy:

Geoemyda spengleri japonica Fan 1931:148

Geoemyda japonica, Geoemyda japonicus

Type locality: "Japan and other Pacific Islands." Restricted to "Nawa (?Naha), Okinawajima Is., Japan" [yellow star] by Yasukawa et al. (1992:149).

Type specimen: USNM 34053, holotype, listed by Yasukawa et al. (1992), but not listed by Cochran (1961), Reynolds et al. (2007), or Uetz et al. (2019).

Etymology:* Toponym for Japan.


Geoemyda spengleri (Gmelin 1789) [(09:25)]
Black-breasted Leaf Turtle



Flora Ihlow / Tam Dao National Park, Vietnam



left: Nobuhiro Kawazoe / CBFTT / No data / captivity
right: Thong Van Pham / Vietnam



(orange dots = probable trade; star = type locality)

Distribution:* China (*Guangdong, Guangxi, Guizhou [?], Hainan, Hunan [?], Jiangxi, Yunnan [?]*), Laos, Vietnam

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 246,243 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 816,570 sq. km

Size (Max SCL): male 10.9 cm, female 12.8 cm (Fan 1931; Pham et al. 2018; T. Pham, unpubl. data)

CBFTT Account: Yasukawa and Ota (2010)

IUCN Red List: Endangered (EN A2cd+4cd) (Fong, Hoang, Li, McCormack, Rao, Timmins, and Wang 2020). Previously: Endangered (EN) (ATTWG 2000); Least Concern (LC) [Not Listed] (Behler 1996)

CITES: Appendix II (2013). Previously: Appendix III (China) (2005)

Synonymy:

Testudo spengleri Gmelin 1789:1043 (*nomen protectum*)

Emys spengleri, Geoemyda spengleri, Clemmys (Clemmys) spengleri, Clemmys spengleri, Nicoria spengleri, Geoemyda spengleri spengleri

Type locality: Not designated. Originally indicated as "vermuthlich... Ostindien" [no star] by Walbaum (1785:129); translated to "probably...the East Indies" by Pope (1935:36).

Type specimen: Possibly MNUL s/n, destroyed in World War II; type specimen figured in Walbaum (1785:pl.3).

Etymology:* Eponym honoring Lorenz Spengler (1720–1807), Switzerland / Denmark; educator and malacologist, and owner of the type in his private museum.
Comment: Description cited as sourced from Walbaum (1785:122,pl.3). Name conserved by ICZN (1985a), see Smith et al. (1980c).

Testudo serrata Shaw 1802:51 (junior homonym, not = *Testudo serrata* Daudin 1801 [= *Trachemys scripta scripta*])
Type locality: Not designated.
Type specimen: RCSM s/n (formerly LM s/n), holotype, see Gray (1831d), apparently lost, possibly destroyed during World War II, type specimen figured (pl.9,left).
Etymology:* Latin, *serratus*, saw-shaped, serrated; referring to the posterior carapacial margin.

Testudo tricarinata Bory de Saint-Vincent 1804:308 (junior homonym, not = *Testudo tricarinata* Retzius *in* Schoepff 1792 [= *Kinosternon scorpioides scorpioides*]) (21:66)
Type locality: "Bourbon...mare d'Arzule" [Réunion] [in error, trade].
Type specimen: Not located, type specimen figured (pl.37,f.1).
Etymology:* Latin, *tri-*, three, and *carina*, the keel, *carinata*, keeled; referring to the three carapacial vertebral keels.

Testudo carinata Thunberg 1810:6 (junior homonym and *nomen nudum et novum*, not = *Testudo carinata* Linnaeus 1758 [= *Terrapene carolina carolina*]) (21:66)
Type locality: Not designated.
Type specimen: UPSZTY 277 (formerly UUZM 277 and MGA 46), holotype, listed by Thunberg (1810, 1828), Holm (1957), and Wallin (2001), is a *Geoemyda spengleri* and not the type of *T. carinata* Linnaeus 1758.
Etymology:* Latin, *carina*, the keel, *carinata*, keeled; referring to the carapacial vertebral keel.
Comment: Probable unjustified replacement name for *Testudo tricarinata* Bory de Saint-Vincent.

Geoemyda spengleri sinensis Fan 1931:146
Type locality: "Loshiang and Kutchen" [Luoxiang and Guchen, Guangxi, China] [green star].
Type specimens: SYS 1202, 8001–68, syntypes (69), possibly lost; RMNH 5887, listed as paratype by Hoogmoed et al. (2010); Uetz et al. (2019) erroneously listed no types as designated.
Etymology:* Toponym for China.

Heosemys Stejneger 1902

Asian Forest Turtles
(includes 4 species, 4 taxa)

Heosemys Stejneger 1902:238
Type species: *Heosemys spinosa* [= *Emys spinosa* Gray 1831d], by original designation.
Etymology:* Greek, *eos*, east, morning, dawn, and *emys*, turtle; referring to the eastern hemisphere distribution of the genus.


Hieremys Smith 1916:50
Type species: *Hieremys annandalii* [= *Cyclemys annandalii* Boulenger 1903a], by original monotypy.
Etymology:* Greek, *hieros*, sacred, holy, and *emys*, turtle; referring to the type species being revered as a sacred animal in its countries of origin and kept in temple ponds.

Heosemys annandalii (Boulenger 1903a) (07:41)
Yellow-headed Temple Turtle


Sitha Som / TL / Tonle Sap, Cambodia



left: Peter Paul van Dijk / Thailand / captivity, Samut Prakan Crocodile Farm and Zoo / male
right: Craig B. Stanford / Thailand / captivity / Bangkok Zoo


(star = type locality)

Distribution: Cambodia, Laos, Malaysia (*West*), Thailand, Vietnam
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 289,524 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 855,655 sq. km
Size (Max SCL): male 50.0 cm, female 42.0 cm (Cox et al. 1998; Platt et al. 2008; Ceballos et al. 2013)
IUCN Red List: Critically Endangered (CR A2cd+4cd) (Cota, Horne, McCormack, and Timmins 2021). Previously: Endangered (EN) (ATTWG 2000); Vulnerable (VU) (TFTSG *in* Baillie and Groombridge 1996)
CITES: Appendix II (2003); zero quota for wild specimens for commercial purposes (2013)
Synonymy:

Cyclemys annandalii Boulenger 1903a:142 (07:41)
Hieremys annandalii, *Heosemys annandalii*
Type locality: "Kampong Jalor" [= Yala, Yala Prov., Thailand] [yellow star].
Type specimens: NHMUK 1946.1.22.67–68, 1947.3.5.62 (formerly 1903.4.13.2–4), syntypes (3).
Etymology:* Eponym honoring Nelson Annandale (1876–1924), United Kingdom (Scotland) / India; naturalist and herpetologist at the Indian Museum in Calcutta, who co-founded the Zoological Survey of India.


Hieremys annandalei Smith 1916:50 (*nomen novum*)
Cyclemys annandalei
Etymology:* Eponym honoring Nelson Annandale (1876–1924), United Kingdom (Scotland) / India; naturalist and herpetologist at the Indian Museum in Calcutta, who co-founded the Zoological Survey of India.
Comment: Unjustified emendation of *annandalii*.

Heosemys depressa (Anderson 1875)
Arakan Forest Turtle


Brian D. Horne / TCC / Myanmar



John B. Iverson / No data / captivity


**
(orange dot = probable trade; stars = type localities)

Distribution: Bangladesh, Myanmar
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 43,864 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 63,487 sq. km
Size (Max SCL): male 29.6 cm, female 25.9 cm (Platt et al. 2010; Ceballos et al. 2013)
IUCN Red List: Critically Endangered (CR A2cd+4cd) (Platt, Rahman, Horne, and Praschag 2020). Previously: Critically Endangered (CR) (ATTWG 2000); Critically Endangered (CR) (TFTSG *in* Baillie and Groombridge 1996)
CITES: Appendix II (2003); zero quota for wild specimens for commercial purposes (2013)
Synonymy:
Geoemyda depressa Anderson 1875:284
Geoëmyda depressa, Heosemys depressa
Type locality: "Arakan" [Myanmar]. Restricted to "hilly region in the neighbourhood of Akyab in Arracan" [nr. Sittwe, Myanmar] [yellow star] by Anderson (1879:722).
Type specimens: ZSI 751, syntype, a skull, other syntypes apparently lost, including specimen figured in Anderson (1879:pl.55-56), see Iverson and McCord (1997a), Das et al. (1998), and Kundu et al. (2018a); NHMUK 1887.3.30.1 may be a syntype (Iverson 1992); ZMB 8869, recorded as "holotype" by Fritz et al. (1994:165), is not a type, see Iverson and McCord (1997a).
Etymology:* Latin, *de-*, down, and *pressus*, pressed; referring to the depressed flattened shell.
Geomyda arakana Theobald 1876:vii
Geoemyda arakana
Type locality: "Akyab" [Sittwe, Myanmar] [green star].
Type specimen: NHMUK 1947.3.4.28, 1947.3.5.69 (formerly 1888.6.18.1, 1890.1.28.9) (separate parts of same specimen), holotype, see Iverson and McCord (1997a).
Etymology:* Toponym for the Arakan region, Burma (=Myanmar).

Heosemys grandis (Gray 1860d)
Giant Asian Pond Turtle


Jérôme Maran / Vietnam



Sabine Schoppe / No data / trade


**
(orange dots = probable trade; star = type locality)

Distribution: Cambodia, Laos, Malaysia (*West*), Myanmar, Thailand, Vietnam
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 575,784 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 1,800,090 sq. km
Size (Max SCL):* male 48.0 cm, female 45.0 cm (Cox et al. 1998; Rudolphi and Weser 2000; Bonin et al. 2006; Iverson and McCord 2006; Le 2007)

IUCN Red List: Critically Endangered (CR A2cd+4cd) (Cota, Horne, and Shepherd 2021). Previously: Vulnerable (VU) (ATTWG 2000); Near Threatened (NT) (TFTSG *in* Baillie and Groombridge 1996)

CITES: Appendix II (2003)

Synonymy:


Emys siamensis Gray *in* Günther 1860:114 *(nomen nudum)*
Etymology:* Toponym for Siam (= Thailand).

Geoemyda grandis Gray 1860d:218
Geomyda grandis, Clemmys grandis, Heosemys grandis
Type locality:* "Cambojia" [Cambodia] [possibly nr. Angkor Wat] [yellow star].
Type specimens:* NHMUK 1947.3.4.7 and 1947.3.4.55–56 (formerly 1860.8.28.1–2 and 1861.6.1.7), syntypes (3), collected by Henri Mouhot; Uetz et al. (2019) erroneously listed NHMUK 1867.12.30.66 as a possible syntype.
Etymology:* Latin, *grandis*, great, significant, large; referring to the somewhat larger size of this species.

Heosemys spinosa (Gray 1830e) (12:25)
Spiny Turtle


Peter Paul van Dijk / Perlis State Park, Perlis, Malaysia / adult


John B. Iverson / captivity / juvenile


**
(star = type locality)

Distribution: Brunei, Indonesia (*Sumatra, Kalimantan*), Malaysia (*East, West*), Myanmar, Philippines (*Sulu Archipelago [Tawi-Tawi]*), Singapore, Thailand

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 931,054 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 3,298,125 sq. km

Size (Max SCL):* male 29.9 cm, female 20.1 cm (Goetz 2007; Pierlioni 2019)

IUCN Red List: Endangered (EN A2cd) (Cota, Guntoro, Horne, Kusrini, Krishnasamy, and Shepherd 2021).
Previously: Endangered (EN) (ATTWG 2000); Vulnerable (VU) (TFTSG *in* Baillie and Groombridge 1996)

CITES: Appendix II (2003)

Synonymy:

Emys spinosae Bell *in* Gray 1830a:pl.77 (*nomen oblitum*)
Type locality: "Penang" [West Malaysia] [yellow star].
Type specimens: OUM 8517, NHMUK 1862.8.28.1, syntypes (2), discussed by Bourret (1941c) and Nowak-Kemp and Fritz (2010).
Etymology:* Latin, *spina*, thorn, *spinosa*, thorny, full of thorns, prickly; referring to the numerous marginal spinous processes on the carapace of juveniles of this species.

Emys spinosa Gray 1830e:8 (*nomen novum*)
Geoemyda spinosa, Geomyda spinosa, Clemmys (Clemmys) spinosa, Clemmys spinosa, Heosemys spinosa
Type locality: "Penang" [West Malaysia] [yellow star].
Type specimens: OUM 8517, NHMUK 1862.8.28.1, syntypes (2) by default, discussed by Bourret (1941c) and Nowak-Kemp and Fritz (2010).
Etymology:* Latin, *spina*, thorn, *spinosa*, thorny, full of thorns, prickly; referring to the numerous marginal spinous processes on the carapace of juveniles of this species.
Comment: Accepted emendation of *spinosae*.

Leucocephalon McCord, Iverson, Spinks, and Shaffer 2000

Sulawesi Forest Turtles

(includes 1 species, 1 taxon)

Leucocephalon McCord, Iverson, Spinks, and Shaffer 2000:21

Type species: *Leucocephalon yuwonoi* McCord, Iverson, and Boeadi 1995, by original designation.


Etymology:* Greek, *lefkos*, white, bright, *kefale*, head; referring to the white head of most males of this genus.

Leucocephalon yuwonoi (McCord, Iverson, and Boeadi 1995) (07:42)


Sulawesi Forest Turtle



Cris Hagen / CBFTT / TCC / Sulawesi, Indonesia / male



Cris Hagen / CBFTT / Sulawesi, Indonesia / male



(orange dots = trade; star = type locality)

Distribution: Indonesia (*Sulawesi*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 30,718 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 65,701 sq. km

Size (Max SCL): male 27.8 cm, female 24.0 cm (Ives et al. 2008; Hagen et al. 2009 CBFTT)

CBFTT Account: Hagen, Platt, and Innis (2009)

IUCN Red List: Critically Endangered (CR A2bcd) (Stanford and Hamidy 2021). Previously: Critically Endangered (CR) (ATTWG 2000); Data Deficient (DD) (TFTSG *in* Baillie and Groombridge 1996)

CITES: Appendix II (2003)

Synonymy:

Geoemyda yuwonoi McCord, Iverson, and Boeadi 1995:311
Heosemys yuwonoi, *Leucocephalon yuwonoi*

Type locality: "near Gorontalo [0°33' N, 123°05' E] on the Minahassa Peninsula, northern Sulawesi [Celebes], Indonesia" [yellow star].

Type specimen: MZB 10295, holotype; MZB 10296, UF 97333–5, paratypes (4); MZB specimens (2) possibly lost (McCord, pers. comm. to Iverson).

Etymology:* Eponym honoring Frank Yuwono (1958–), Indonesia; animal exporter who obtained the types.

Mauremys Gray 1869b (07:44, 09:26)

Eurasian Stripe-necked Turtles

(includes 9 species, 11 taxa)

Mauremys Gray 1869b:500

Type species: *Mauremys fuliginosa* [= *Emys fuliginosus* Gray 1860c = subjective synonym of *Emys leprosa* Schoepff *in* Schweigger 1812], by subsequent designation by Lindholm (1929:281).

Etymology:* Greek, *mavros*, dark, black, and *emys*, turtle; referring to the dark carapace color of the species originally described in this genus.

Ocadia Gray 1870c:35

Type species: *Ocadia sinensis* [= *Emys sinensis* Gray 1834a], by original monotypy.

Etymology:* Anonym, *ocadia*, possibly a fabricated name, or possibly an Eponym honoring someone named Ocad, or whose name started with Ocad or O'Cad.

Emmenia Gray 1870c:38

Type species: *Emmenia grayi* [= *Emys grayi* Günther 1869 = subjective synonym of *Mauremys caspica siebenrocki* Wischuf and Fritz *in* Fritz and Wischuf 1997 = subjective synonym of *Testudo caspica* Gmelin 1774], by original monotypy.

Etymology:* Anonym, *emmenia*, possibly a fabricated name, or possibly an Eponym honoring the author's wife, Emma Gray (1787–1876), and/or possibly Greek, *emmeno*, to be faithful or to cleave to, possibly referring to his wife.

Eryma Gray 1870c:44 (junior homonym, not = *Eryma* Meyer 1840 [= Crustacea †] or *Eryma* Albers 1854 [= Gastropoda] or *Eryma* Förster 1868 [= Hymenoptera])

Type species: *Eryma laticeps* [= *Emys laticeps* Gray 1854a = subjective synonym of *Emys leprosa* Schoepff *in* Schweigger 1812], by original monotypy.

Etymology:* Greek, *erymnos*, defensive, reinforced, strong; referring to the strong jaws and claws of this taxon.

Cathaiemys Lindholm 1931:29

Type species: *Cathaiemys mutica* [= *Emys mutica* Cantor 1842b], by original designation.

Etymology:* Toponym for Cathay (= China) and Greek, *emys*, turtle.

Pseudocadia Lindholm 1931:30

Type species: *Pseudocadia anyangensis* [= *Testudo anyangensis* † Ping 1930], by original designation.

Etymology:* Greek, *pseudo*, false, untrue, and *ocadia*; referring to the similarity of this genus to *Ocadia*.

Chinemys Smith 1931:116

Type species: *Chinemys reevesi* [= *Emys reevesii* Gray 1831d)], by original monotypy.

Etymology:* Toponym for China and Greek, *emys*, turtle.

Annamemys Bourret 1940:15

Type species: *Annamemys merkleni* Bourret 1940 [= subjective synonym of *Cyclemys annamensis* Siebenrock 1903a], by original monotypy.

Etymology:* Toponym for Annam (= Vietnam) and Greek, *emys*, turtle.


Mauremys annamensis (Siebenrock 1903a) (07:44, 14:29, 17:49, 21:69) (68)*
Vietnamese Pond Turtle, Annam Pond Turtle



Jeffrey E. Dawson / CBFTT / Vietnam / captivity



Jeffrey E. Dawson / CBFTT / Vietnam / captivity



** (stars = type localities)

Distribution: Vietnam

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 20,576 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 44,027 sq. km

Size (Max SCL): male 23.2 cm, female 28.5 cm (McCormack et al. 2014 CBFTT)

CBFTT Account: McCormack, Dawson, Hendrie, Ewert, Iverson, Hatcher, and Goode (2014)

IUCN Red List: Critically Endangered (CR A2bcd+4bcd; B2ab(i,ii,iii,iv,v); D) (McCormack, van Dijk, Roberton, and Dawson 2020). Previously: Critically Endangered (CR) (ATTWG 2000), Least Concern (LC) [Not Listed] (Behler 1996)

CITES: Appendix I (2019). Previously: Appendix II (2003)

Synonymy:

Cyclemys annamensis Siebenrock 1903a:341
Cuora (Cyclemys) annamensis, Cuora annamensis, Annamemys annamensis, Mauremys annamensis, Cathaiemys annamensis
Type locality: "Annam (Phuc-Son)" [= Phuoc Son, Quang Nam, Vietnam (15°33'00"N; 108°04'00"E) [yellow star]; see Le et al. 2004].
Type specimen: NMW 23394, holotype, see Tiedemann and Häupl (1980), Tiedemann et al. (1994), and Gemel et al. (2019).
Etymology:* Toponym for Annam (= Vietnam).

Annamemys merkleni Bourret 1940:15
Annamemys mekleni
Type locality: "Annam...Fai-Fo" [= Hoi An, Quang Nam, Vietnam] [green star].
Type specimen: CAS-SU 9142 (formerly VNUH Y.20), lectotype, designated by Savage (1953:471).
Etymology:* Eponym honoring Monsieur Merklen (given name not discovered) (1900s), France / Vietnam; official and naturalist working for the Service Forestier de l'Indochine in Annam, French Indochina (now part of Vietnam), who collected the types.

Clemmys guangxiensis Qin 1992:60 (*partim*, hybrid) (14:29)
Mauremys guangxiensis
Type locality: "Nanning, Guangxi" [China]. [in error?] [orange star].
Type specimen: SNHM 88701, holotype.
Etymology:* Toponym for Guangxi, China.
Comment:* Type specimen likely acquired from the trade; type locality likely a trade location. Apparent hybrid of *Cuora trifasciata trifasciata*, *Mauremys mutica mutica*, and possibly *Mauremys annamensis*.

Ocadia glyphistoma McCord and Iverson 1994:53 (*partim*, hybrid) (07:33)
Type locality: "near the Vietnam border southwest of Nanning, Guangxi Province, China." [in error?] [brown star, location estimated].
Type specimen: UF 84818, holotype.
Etymology:* Greek, *glyphis*, carved or notched, and *stoma*, mouth; referring to the distinctly notched upper jaw.
Comment:* Type specimen likely acquired from the trade; type locality likely a trade location. Apparent hybrid of *Mauremys sinensis* and *Mauremys annamensis*. Previously listed as Data Deficient (DD) on the IUCN Red List (TFTSG *in* Baillie and Groombridge 1996).


Mauremys caspica (Gmelin 1774) (09:27, 12:26, 17:50)
Caspian Turtle, Caspian Terrapin, West Asian Stripe-necked Terrapin



Asghar Mobaraki / Maharloo Lake, Shiraz, Fars, Iran



Anders G.J. Rhodin / captivity / Sharjah Zoo, United Arab Emirates



** (orange dots = possibly introduced or questionable; stars = type localities)

Distribution: Armenia, Azerbaijan, Bahrain, Georgia, Iran (*Ardabil, Bushehr, Chahar Mahal Va Bakhtiari, East Azarbaijan, Fars, Gilan, Golestan, Ilam, Isfahan, Kermanshah, Khuzestan, Kordestan, Lorestan, Mazandaran, West Azarbaijan*), Iraq, Russia (*Dagestan*), Saudi Arabia (*Eastern*), Syria, Turkey, Turkmenistan

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 742,006 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 2,731,757 sq. km

Size (Max SCL): male 24.0 cm, female 24.0 cm (Fritz and Wischuf 1997; Yazarloo et al. 2017)

CBFTT Account:* Vamberger, Busack, Ayaz, Fritz, and Mazanaeva (2024)

IUCN Red List: Least Concern (LC) [Not Listed] (Behler 1996)

TFTSG Provisional Red List: Least Concern (LC) (2011)

Synonymy:

Testudo caspica Gmelin 1774:59,pl.X
Emys caspica, Clemmys caspica, Clemmys (Clemmys) caspica, Terrapene caspica, Clemmys caspica caspica, Mauremys caspica, Mauremys caspica caspica
Type locality: "Schamachie...bei Bach Pusahat" [= Pirsagat (39º53'N, 49º25'E), Shemakha, Azerbaijan] [yellow star]. Emended to "Hircaniae aquis dulcibus" by Gmelin (1789:1042); and to "les bords de la mer Caspienne...dans les eaux douces de l'Hircanie" [Azerbaijan] by Daudin (1801:124).
Type specimen: Not located, specimen figured (pl.10-11), lectotype, designated by Fritz and Wischuf (1997:237).
Etymology:* Toponym for the Caspian Sea region.

Testudo ecaudata Daudin 1801:125 (senior homonym and *nomen novum*, not = *Testudo ecaudata* Pallas 1814 [= *Testudo (Testudo) graeca buxtoni*])
Type locality: "les bords de la mer Caspienne...dans les eaux douces de l'Hircanie" [Azerbaijan] [yellow star].
Type specimen: Not located, specimen figured in Gmelin (1774:pl.10-11), by default, lectotype, designated by Fritz and Wischuf (1997:237).
Etymology:* Latin, *e-*, without, and *caudata*, having a tail, tailed; referring to the apparent lack of a tail on the examined specimen.
Comment: Unjustified replacement name for *caspica*.

Emys caspia Rüppell *in* Gray 1830e:9 (*nomen novum*)
Testudo caspia, Clemmys caspia, Clemmys caspia caspia
Type locality: "Caspian Sea."
Etymology:* Toponym for the Caspian Sea region.
Comment: Unjustified emendation or error for *caspica*.

Emys grayi Günther 1869:504 (junior homonym, not = *Emys grayi* Bocourt 1868 [= *Trachemys grayi grayi*])
Emmenia grayi
Type locality: "Bussora" [= Basrah (30º30'N, 47º50'E), Iraq] [green star].
Type specimen: NHMUK 1947.4.3.47 (formerly 1869.8.28.4), holotype, see Fritz and Wischuf (1997).
Etymology:* Eponym honoring John Edward Gray (1800–1875), United Kingdom; herpetologist, turtle taxonomist, and curator at the British Museum of Natural History, London.

Mauremys caspica siebenrocki Wischuf *in* Maran 1996:17 (*nomen nudum*)
Etymology:* Eponym honoring Friedrich Siebenrock (1853–1925), Austria; zoologist, herpetologist, and curator at the Naturhistorisches Museum, Vienna.

Mauremys caspica schiras Wischuf *in* Maran 1996:17 (*nomen nudum*)
Etymology:* Toponym for Schiras (= Shiraz), Iran.

Mauremys caspica ventrimaculata Wischuf and Fritz 1996:114
Type locality: "Tang-e Karam (Tang-i-Kerim), 70 Meilen östlich von Schiraz, Iran (29º 03' N, 53º 38' E)" [blue star].
Type specimen: NHMUK 1874.11.23.4, holotype.
Etymology:* Latin, *venter*, belly or abdomen, and *maculata*, spotted, speckled, stippled; referring to the plastral pattern.

Mauremys caspica siebenrocki Wischuf and Fritz *in* Fritz and Wischuf 1997:240
Type locality: "Kerbela, Irak" [Iraq] [pink star].
Type specimen: MTD 27101, holotype.
Etymology:* Eponym honoring Friedrich Siebenrock (1853–1925), Austria; zoologist, herpetologist, and curator at the Naturhistorisches Museum, Vienna.


Mauremys japonica (Temminck and Schlegel 1838) (10:18, 14:30, 21:70)
Japanese Pond Turtle, *Ishigame*



Hiromi Tada / Tokyo, Honshu, Japan



Yuichirou Yasukawa / CBFTT / *left*: Shiga, Honshu, Japan, *right*: Kyoto, Honshu, Japan



**

(orange dots = probably introduced; star = type locality)

Distribution: Japan (*Honshu, Kyushu, Shikoku*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 153,544 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 495,533 sq. km

Size (Max SCL):* male 17.4 cm, female 21.0 cm (Iverson and McCord 1994; Schilde 2004; Ceballos et al. 2013; Itescu et al. 2014)

CBFTT Account: Yasukawa, Yabe, and Ota (2008)

IUCN Red List: Near Threatened (NT) (ATTWG 2000). Previously: Least Concern (LC) [Not Listed] (Behler 1996)

CITES: Appendix II (2013)

Synonymy:


Emys palustris var. *Japon* Temminck and Schlegel 1834:pls.8-9 (invalid vernacular name) (14:30)

Emys vulgaris var. *japonaise* Temminck and Schlegel 1834:54 (invalid vernacular name) (14:30)

Emys vulgaris japonica Temminck and Schlegel 1838:139 (10:18, 14:30)
Emys japonica, Emys caspica japonica, Clemmys japonica, Mauremys japonica, Ocadia japonica
Type locality:* "Japon" [Japan] [probably nr. Dajima, Nagasaki, Kyushu] [yellow star].
Type specimens:* RMNH 3331–34, 6142, syntypes (6), collected by Philipp Franz von Siebold, discussed by Hoogmoed et al. (2010); MNHN 1954 erroneously listed as a syntype by King and Burke (1989), Iverson (1992), and Uetz et al. (2019).
Etymology:* Toponym for Japan.

Testudo margaritifera Schlegel *in* Gray 1856b:11 (*nomen nudum*)
Etymology:* Latin, *margarita*, pearl, and *ferre*, to bear or carry; possibly referring to carapacial filamentous algae growth suggestive of pearls.

Mauremys leprosa (Schoepff *in* Schweigger 1812) (07:45,17:51)
Mediterranean Pond Turtle, Spanish Terrapin, Mediterranean Stripe-necked Terrapin
(includes 2 subspecies)



**

(subspecies: *leprosa* = red, *saharica* = blue, overlap = intergrades; orange dots = possibly prehistorically introduced or traded or questionable localities)

Distribution: Algeria, France, Libya, Mauritania (*prehistorical introduction?*), Mali (*prehistorical introduction?*), Morocco, Niger (*prehistorical introduction?*), Portugal, Spain (*Continental*), Tunisia

Introduced: France, Italy, Spain (*Balearic Islands*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 957,571 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 6,551,749 sq. km

Size (Max SCL): male 21.0 cm, female 24.0 cm (see subspp.)

CBFTT Account: Bertolero and Busack (2017)


IUCN Red List:* Least Concern (LC) [Not Listed] (Behler 1996). Regional (Europe): Near Threatened (NT) (Luiselli 2024). Previously: Vulnerable (VU A2ac+3c) (van Dijk, Mateo Miras, Cheylan, Joger, Sá-Sousa, and Pérez-Mellado 2004; Cox and Temple 2009). Regional (European Union–EU27): Vulnerable (VU A2ac+A3c) (Cox and Temple 2009)

TFTSG Provisional Red List: Vulnerable (VU) (2011)


Mauremys leprosa leprosa (Schoepff *in* Schweigger 1812) [07:45,17:51]
Mediterranean Pond Turtle, Mediterranean Stripe-necked Terrapin



Albert Bertolero / CBFTT / Flix, Catalonia, Spain



Albert Bertolero / CBFTT / Flix, Catalonia, Spain



(subspecies: *leprosa* = red, *saharica* = blue, overlap = intergrades; orange dots = probably introduced; stars = type localities)

Distribution: France, Morocco, Portugal, Spain (*Continental*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 515,727 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 969,097 sq. km

Size (Max SCL): male 21.0 cm, female 24.0 cm (Bertolero and Busack 2017 CBFTT)

Synonymy:

Emys leprosa Schoepff *in* Schweigger 1812:298
Clemmys (Clemmys) leprosa, Clemmys leprosa, Emys caspica leprosa, Clemmys caspica leprosa, Mauremys caspica leprosa, Mauremys leprosa, Mauremys leprosa leprosa
Type locality: Not known. Restricted to "Südspanien" [southern Spain] [yellow star, location estimated] by Mertens and Müller (1928:22).
Type specimen: MNHN 1934, holotype, see Bour and Maran (1999) and Ceríaco and Bour (2012); not MNHN 7936, listed as "holotype" (Bour, pers. comm. *in* Iverson 1992).
Etymology:* Latin, *leprosa*, leprous; referring to the holotype with pathological tubercles on the carapace.

Emys lutescens Schweigger 1812:302
Clemmys (Clemmys) lutescens, Clemmys lutescens
Type locality: Not known. Restricted to "southern Spain" [yellow star, location estimated] by Fritz and Havaš (2007:230).
Type specimen: MNHN s/n, holotype, apparently lost, see Bour and Maran (1999).
Etymology:* Latin, *lutescens*, yellowish, golden yellowish; referring to the somewhat yellowish colored carapace.

Emys marmorea Spix 1824:13
Clemmys marmorea
Type locality: "Brasil" [Brazil, in error]. Restricted to "environs de Gibraltar" [green star] by Temminck and Schlegel (1834:53), and to "southern Spain" [yellow star, location estimated] by Fritz and Havaš (2007:230).
Type specimen: ZSM, holotype or syntypes, not known, apparently lost, type specimen figured (pl.10), not listed by Hoogmoed and Gruber (1983) or Franzen and Glaw (2007).
Etymology:* Latin, *marmor*, marble; referring to the marbled appearance of the carapace.

Clemmys sigriz Michahelles 1829:1296
Terrapene sigriz, Emys sigriz, Clemmys caspica sigriz
Type locality: "paludibus Hispaniae meridionalis" [marshes of southern Spain] [yellow star, location estimated].
Type specimens: Not known or located, syntypes, originally live, see Bour and Maran (1999).
Etymology:* Eponym honoring Adalbert von Sigriz (1800s), Germany; physician and explorer who obtained turtles identified as *Clemmys caspica* at Ragusa (Sicily) on the Adriatic Sea, used by the author as comparison for his description of Spanish *C. sigriz*.

Emys vulgaris Gray 1830e:9 [10:7]
Type locality: "South of Europe. Spain."
Type specimens: Specimen figured in Gray (1831d:pl.4,top), lectotype, designated by Fritz and Wischuf (1997:249); original syntypes (20+) in OUM, NHMUK, and RMNH, discussed by Fritz and Wischuf (1997), Hoogmoed et al. (2010), and Nowak-Kemp and Fritz (2010); NHMUK 1988.79, erroneously listed as "holotype" by Bour and Maran (1999), also discussed by Fritz and Wischuf (1997).
Etymology:* Latin, *vulgaris*, common or usual; referring to the apparent commonness of the species.

Emys sigritzii Michahelles *in* Gray 1831d:24 (*nomen novum*)
Etymology:* Eponym honoring Adalbert von Sigriz (1800s), Germany; physician and explorer who obtained turtles identified as *Clemmys caspica* at Ragusa (Sicily) on the Adriatic Sea, used by the author as comparison for his description of Spanish *C. sigriz*.
Comment: Unjustified emendation of *sigriz*.

Emys laticeps Gray 1854a:134
Clemmys laticeps, Eryma laticeps
Type locality: "West Africa, River Gambia" [Gambia, in error]. Emended to "Tétouan, Morocco" [blue star] by Fritz and Havaš (2007:230) without explanation.
Type specimens: NHMUK 1947.3.4.43–44, 1947.3.4.46, 1947.3.5.38–39 (formerly 1848.8.22.6–9, 1848.8.22.[s/n]), syntypes (5), see Bour and Maran (1999).
Etymology:* Latin, *latus*, broad, and *-ceps*, head; referring to the short, broad head.

Emys fuliginosus Gray 1860c:232
Emys fuliginosa, Clemmys fuliginosa, Mauremys fuliginosa
Type locality: "Egypt?...North Africa?" Emended to "southern Spain" [yellow star, location estimated] by Fritz and Havaš (2007:230) without explanation.
Type specimen: NHMUK 1947.3.4.45, holotype, see Bour and Maran (1999).
Etymology:* Latin, *fuligo*, soot, fog, *fuliginosus*, soot-colored; referring to the fact generally dark shell color.

Mauremys laniaria Gray 1869b:499
Emys (Mauremys) laniaria, Emys lamaria
Type locality: Not known. Restricted to "southern Spain" [yellow star, location estimated] by Fritz and Havaš (2007:230).
Type specimens: NHMUK 1946.1.22.52–53 (formerly 1869.12.21.3–4), syntypes (2), listed by Bour and Maran (1999).
Etymology:* Latin, *laniaria*, to tear apart, to tear into pieces; referring to the ravenous carnivorous eating habit of the live type specimen.

Emys flavipes Gray 1869c:643
Type locality: Not known. Restricted to "southern Spain" [yellow star, location estimated] by Fritz and Havaš (2007:230).
Type specimens: NHMUK 1946.1.22.31, 1946.1.22.43, and s/n (formerly 1859.5.11.2, 1861.10.1.4, 1870.12.22.7), syntypes (3), see Bour and Maran (1999).
Etymology:* Latin, *flava*, yellow, golden yellow, and *pes*, foot; referring to the forelegs with bright yellowish-orange undersides.

Emys marginata Gray 1873j:36 (*nomen nudum*)
Type locality: "W. Africa."
Type specimen: NHMUK, holotype, not located.
Etymology:* Latin, *marginis*, edge or border, and *-atus*, provided with, referring to the expanded carapacial margin.

Mauremys leprosa atlantica Schleich 1996a:32 (07:45)
Type locality: "Larache, Nord-Marokko" [Morocco] [white star].
Type specimen: ZSM 35/1996, holotype, see Franzen and Glaw (2007).
Etymology:* Toponym for the Atlantic Ocean.

Mauremys leprosa marokkensis Schleich 1996a:40 (07:45)
Type locality: "ca. 2 km NE Tahanaoute / Kreuzung S 501, wenige km vor Aguelmouss; S-Marrakech, N 31º 24' 54.0, W 7º 49' 44.4, Zentral-Morokko" [Morocco] [brown star].
Type specimen: ZSM 79/1994, holotype, see Franzen and Glaw (2007).
Etymology:* Toponym for Morocco.

Mauremys leprosa wernerkaestlei Schleich 1996a:44 (07:45)
Type locality: "Wasserlauf im Schnittpunkt von Oued Serou, Oued Oum de Oumbia. Oued Oum er Rbia, S-Khenifra, N 32º 44' 55.1, W 5º 41' 10.7, ca. 90 m NN, Zentral-Morokko" [Morocco] [pink star].
Type specimen: ZSM 81/1994, holotype, see Franzen and Glaw (2007).
Etymology:* Eponym honoring Werner Kästle (1926–2019), Germany; herpetologist.


Mauremys leprosa saharica Schleich 1996 (07:45,17:51) (69)*
Saharan Pond Turtle



Andreas Nöllert / CBFTT / Bou-Jerif, Morocco



left: Andreas Nöllert / Oued Noun, Morocco
right: Khaled Merabet / Tlemcen National Park, Ras Oued El Garsa, M'sila, Algeria



(subspecies: *leprosa* = red, *saharica* = blue, overlap = intergrades; orange dots = possibly prehistorically introduced or traded or questionable localities; stars = type localities)

Distribution: Algeria, Libya, Mauritania (*prehistorical introduction?*), Mali (*prehistorical introduction?*), Morocco, Niger (*prehistorical introduction?*), Tunisia

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 507,796 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 5,343,917 sq. km

Size (Max SCL): male 17.5 cm, female 22.0 cm (Bertolero and Busack 2017 CBFTT)

Synonymy:

Emys fraseri Gray 1869c:643 (*nomen oblitum*) (69)*
Type locality: "North Africa." Restricted to "Lake Tetzara [= Fetzara], Algiers" [Algeria] [orange star] by Gray (1873a:146).
Type specimens: NHMUK 1946.1.22.33–34 (formerly 1846.11.4.21–22), syntypes (2), see Bour and Maran (1999).
Etymology:* Eponym honoring Louis Fraser (1810–1866), United Kingdom; naturalist and curator of the Museum of the Zoological Society of London, who collected the types.
Comment: Declared a *nomen oblitum* here in our annotation.

Mauremys leprosa saharica Schleich 1996a:45 (*nomen protectum*) (07:45) (69)*
Type locality: "Fort Bou Jerif, Goulmime, Südwest-Marokko" [Morocco] [yellow star].
Type specimen: ZSM 200/1993, holotype, see Franzen and Glaw (2007).
Etymology:* Toponym for the Sahara Desert.
Comment: Declared a *nomen protectum* here in our annotation.

Mauremys leprosa erhardi Schleich 1996a:36 (07:45) (69)*
Type locality:* "NNE Taza...N 35º 25' 32.8, W 02º 52' 11.5...Nordost Marokko" [Morocco]; geographic coordinates inaccurate, emended here to N 34º 25' 32.8, W 03º 52' 11.5 [green star].
Type specimen: ZSM 13/1996, holotype, see Franzen and Glaw (2007).
Etymology:* Eponym honoring Andreas Erhard (1960–), Germany; historian, conservationist, and naturalist.

Mauremys leprosa zizi Schleich 1996a:48 (07:45) (69)*
Type locality: "Oued Ziz, Meski bis Erfoud, Süd-Marokko" [Morocco] [blue star].
Type specimen: ZSM 15/1996, holotype, see Franzen and Glaw (2007).
Etymology:* Toponym for the Oued Ziz river, Morocco.

Mauremys leprosa vanmeerhaeghei Bour and Maran 1999:42 (07:45) (69)*

Type locality: "Mare amont dans l'oasis de Sidi El Mehdaoui (29º30 N, 8º00 W)...Province de Tata (ca. 30 km de Tata), Maroc...affluent de...l'oued Draa" [Morocco] [pink star].
Type specimen: MNHN 1996.8204, holotype.
Etymology:* Eponym honoring Bernard Vanmeerhaeghe (1950–1995), France; chemist and amateur herpetologist.

Mauremys mutica (Cantor 1842b) (17:49, 21:69) (68)*
Yellow Pond Turtle
(includes 2 subspecies)



(subspecies: *mutica* = red; *kami* = blue [Ryukyus]; orange dots = trade or questionable)

Distribution:* China (*Anhui, Fujian, Guangdong, Guangxi, Hainan, Hubei, Jiangsu, Zhejiang*), Japan (*Ryukyu Archipelago*), Laos (?), Taiwan, Vietnam
Introduced: Japan (*mainland*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 425,278 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 1,981,730 sq. km
Size (Max SCL):* male 22.5 cm, female 19.5 cm (see subspp.)
IUCN Red List: Critically Endangered (CR A2cd+4cd) (Fong, Hoang, Li, McCormack, Rao, and Wang 2021). Previously: Endangered (EN) (ATTWG 2000); Least Concern (LC) [Not Listed] (Behler 1996)
CITES: Appendix II (2003)

Mauremys mutica mutica (Cantor 1842b) (68)*
Yellow Pond Turtle



Tien-Hsi Chen / Taiwan



Peter Paul van Dijk / Hainan, China / trade



(subspecies: *mutica* = red; orange dots = trade or questionable; stars = type localities)


Distribution:* China (*Anhui, Fujian, Guangdong, Guangxi, Hainan, Hubei, Jiangsu, Zhejiang*), Laos (?), Taiwan, Vietnam
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 424,573 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 1,856,296 sq. km
Size (Max SCL):* male 22.5 cm, female 19.5 cm (Yasukawa et al. 1996; Schilde 2004; Ceballos et al. 2013)
CITES: Appendix II (2003)
Synonymy:
Emys muticus Cantor 1842b:482
Emys mutica, Clemmys mutica, Damonia mutica, Geoclemys mutica, Cathaiemys mutica, Mauremys mutica, Mauremys mutica mutica, Cathaiemys mutica mutica
Type locality: "Chusan...Island" [Zhoushan Island, Zhejiang, China] [yellow star]. Cited as "Canton" [in error] by many authors, corrected by Iverson and McCord (1989:23).
Type specimen: NHMUK 1947.3.5.34, holotype.
Etymology:* Latin, *muticus*, smooth; referring to the smooth carapace surface.
Clemmys schmackeri Boettger 1894:129
Type locality: "China, wahrscheinlich Hainan" [green star, location estimated].
Type specimen: SMF 7584, holotype, see Mertens (1967b).
Etymology:* Eponym honoring Bernhard Schmacker (1852–1896), Germany / China; malacologist who worked in China and collected the type.
Annamemys grochovskiae Dao 1957:1214
Mauremys grochovskiae
Type locality:* "лес Винь-Линь" (in Russian) [= "forest Vinh-Linh", Quang Tri Province, central Vietnam] [blue star].
Type specimen: Not located, type specimen figured (f.a-d).
Etymology:* Eponym honoring Inne Grochowski (1900s), Soviet Union / Vietnam; female herpetologist who first collected the species.
Comment:* Dao is the family name of the Vietnamese author, Dao Van Tien, not Tien or Van Tien, as sometimes recorded.
Mauremys iversoni Pritchard and McCord 1991:140 (*partim*, hybrid) (07:33)
Type locality: "People's Republic of China: Fujian Province: vicinity of Nanping (26º38' N, 118º10' E)" [in error?] [orange star].
Type specimen: UF 71865, holotype.
Etymology:* Eponym honoring John B. Iverson (1949–), USA; herpetologist and turtle researcher at Earlham College, Indiana.
Comment:* Type specimen likely acquired from the trade; type locality likely a trade location. Apparent hybrid of *Cuora trifasciata trifasciata* and *Mauremys mutica mutica*. Previously listed as Data Deficient (DD) on the IUCN Red List (TFTSG *in* Baillie and Groombridge 1996).
Clemmys guangxiensis Qin 1992:60 (*partim*, hybrid) (14:29)
Mauremys guangxiensis
Type locality: "Nanning, Guangxi" [China] [in error?] [brown star].


Type specimen: SNHM 88701, holotype.
Etymology:* Toponym for Guangxi, China.
Comment:* Type specimen likely acquired from the trade; type locality likely a trade location. Apparent hybrid of *Cuora trifasciata trifasciata*, *Mauremys mutica mutica*, and possibly *Mauremys annamensis*.

Mauremys pritchardi McCord 1998:555 (*partim*, hybrid) (07:33)
Type locality: "Lashio, Myanmar (97°14' E, 22°56' N)" [in error?] [gray star].
Type specimen: UF 105422, holotype.
Etymology:* Eponym honoring Peter C.H. Pritchard (1943–2020), United Kingdom (Northern Ireland) / USA; herpetologist and turtle researcher who founded Chelonian Research Institute, Florida.
Comment:* Type specimen likely acquired from the trade; type locality likely a trade location. Apparent hybrid of *Mauremys mutica mutica* and *Mauremys reevesii*.

Mauremys mutica kami Yasukawa, Ota, and Iverson 1996
Ryukyu Yellow Pond Turtle, *Kami*


John B. Iverson / Ryukyus, Japan


John B. Iverson / Ryukyus, Japan




(subspecies: *mutica* = red; *kami* = blue;
orange dots = possible introductions, star = type locality)

Distribution: Japan (*Ryukyu Archipelago*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 705 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 4,910 sq. km
Size (Max SCL): male 18.9 cm, female 17.6 cm (Yasukawa et al. 1996)
CITES: Appendix II (2003)
Synonymy:
Mauremys mutica kami Yasukawa, Ota, and Iverson 1996:311
Cathaiemys mutica kami
Type locality: "Okawa, Ishigakijima Is., the Yaeyama Group, Ryukyu Archipelago, Japan" [yellow star].
Type specimen: KUZ 19541, holotype.
Etymology:* Vernacular (Japanese), local indigenous name for the turtle in the Yaeyama Group, southern Ryukyus.

Mauremys nigricans (Gray 1834a) (70)*
Chinese Red-necked Turtle, Red-necked Pond Turtle


Ben Anders / CBFTT / China / trade


left: Ben Anders / CBFTT / China / captivity
right: Cris Hagen / China / captivity


(orange dots = possible trade or questionable; stars = type localities)

Distribution: China (*Fujian [?], Guangdong, Guangxi, Hainan [?]*), Vietnam (?)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 133,188 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 236,336 sq. km
Size (Max SCL): male 19.7 cm, female 29.8 cm (Fang 1934; Iverson and McCord 1989; Anders and Iverson 2012 CBFTT; Ceballos et al. 2013)
CBFTT Account: Anders and Iverson (2012)
IUCN Red List: Endangered (EN A1d+2d [ver. 2.3]) (ATTWG 2000). Previously: Data Deficient (DD) (TFTSG *in* Baillie and Groombridge 1996)
TFTSG Provisional Red List: Critically Endangered (CR) (2011)
CITES: Appendix II (2013). Previously: Appendix III (China) (2005)
Synonymy:
Emys nigricans Gray 1834a:53
Emys nigracans, Clemmys nigricans, Damonia nigricans, Chinemys nigricans, Mauremys nigricans
Type locality: "Chinâ prope Canton" [Guangzhou, Guangdong] [yellow star].
Type specimen: NHMUK 1947.3.5.35, holotype.
Etymology:* Latin, *nigricans*, to become black, blackish; referring to the dark carapace color.
Damonia crassiceps Gray 1870c:43 (17:48) (70)*
Type locality: "China." Emended to "likely...Guangdong Province, in the vicinity of Guangzhou (formerly Canton)" [yellow star, location estimated] by Dawson et al. (2025:449).
Type specimen: Specimen figured in Hardwicke (1835:f.19-21), iconotype, by original designation; also in Dawson et al. (2025:f.1).
Etymology:* Latin, *crassum*, thick, fat, and *caput* (*ceps*), head;

referring to the relatively large, broad head.

Geoclemys kwangtungensis Pope 1934:1

Clemmys kuangtungensis, Chinemys kwangtungensis

Type locality: "Lofaoshan, Kwangtung, 300–400 m. altitude" [Mt. Luofu, Guangdong, China] [green star].

Type specimen: ZMB 34955, holotype, see Fritz et al. (1994).

Etymology:* Toponym for Kwangtung (= Guangdong), China.

Geoclemys palaeannamitica † Bourret 1941a:10

Geoclemmys palaeannamitica, Chinemys palaeannamitica

Type locality: "la grotte de Dong-Giao, appelée 'Thung Gianh'... à moins de 1 km de la voie ferrée, vers le N.W., au S.W. de la station de Dong-Giao, tout près de la frontière entre Annam et Tonkin" [Vietnam]. Emended here to "Thung Gianh cave, southwest of Ga Dong Giao rail station, northwest of Bim Son, Thanh Hoa, Vietnam (20.134ºN, 105.851ºE)" [gray star].

Type specimen: MNHN 1948.42 (formerly VNUH T.88), holotype, subfossil, partial skull, figured, see Bour (1980b).

Geological age: Holocene, Neolithic.

Etymology:* Toponym for Annam, an old name for part of Vietnam, and Greek, *palaeo* (*palaios*), ancient, old.


Comment:* Analyzed by Bour (1980b) and synonymized by Pritchard (1994).

Mauremys reevesii (Gray 1831d) (07:46, 17:52, 21:71)

Reeves' Turtle, Chinese Three-keeled Pond Turtle



Nobuhiro Kawazoe / CBFTT / Honshu, Japan / melanistic male



Nobuhiro Kawazoe / CBFTT / Honshu, Japan





(native populations = red, presumed introduced populations = gray; orange dots = presumed trade and/or introduced; stars = type localities)

Distribution: China (*Anhui, Fujian, Gansu, Guangdong, Guizhou, Henan, Hong Kong, Hubei, Hunan, Jiangsu, Jiangxi, Macau, Shangdong, Shanxi, Shaanxi, Sichuan, Zhejiang*), Japan (*prehistorical introduction*), North Korea, South Korea, Taiwan (*prehistorical introduction*)

Introduced: Indonesia (*Timor*), Palau, Philippines, Timor-Leste

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 907,262 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 4,277,816 sq. km

Size (Max SCL): male 23.6 cm, female 30.0 cm (Lovich et al. 2011 CBFTT; J.B. Iverson, unpubl. data)

CBFTT Account: Lovich, Yasukawa, and Ota (2011)

IUCN Red List: Endangered (EN A2bcd+4bcd) (van Dijk 2011). Previously: Endangered (EN) (ATTWG 2000); Least Concern (LC) [Not Listed] (Behler 1996)

CITES: Appendix III (China) (2005)

Synonymy:

Emys reevesii Gray 1831d:73

Clemmys (Clemmys) reevesii, Clemmys reevesii, Geoclemys reevesii, Geoclemmys reevesii, Damonia reevesii, Geoclemys reevesii reevesii, Chinemys reevesi, Chinemys reevesii, Mauremys reevesii

Type locality: "China" [likely nr. Canton (= Guangzhou, Guangdong)] [yellow star, location estimated].

Type specimens: NHMUK 1947.3.5.31–32, syntypes (2).

Etymology:* Eponym honoring John R. Reeves, Jr. (1804–1877), United Kingdom / China; naturalist who worked for the East India Company in China and collected turtles, likely near Canton. The entry for Reeves in Beolens et al. (2011) is incorrect, referring instead to the father.

Emys vulgaris picta Schlegel 1844:127

Type locality: "Japan."

Type specimens: RMNH 3330A–B, MNHN 1954, syntypes (3), discussed by Hoogmoed et al. (2010).

Etymology:* Latin, *picta*, painted; referring to the bright colors of this turtle.

Emys japonica Duméril and Bibron *in* Duméril and Duméril 1851:8 (*nomen novum*)

Type locality: "Japon" [Japan].

Type specimens: RMNH 3330A–B, MNHN 1954, syntypes (3), by default, discussed by Hoogmoed et al. (2010), not MNHN 4096.

Etymology:* Toponym for Japan.

Comment: Unjustified replacement name for *picta*.

Damonia unicolor Gray 1873e:78

Clemmys unicolor, Damonia reevesii unicolor, Geoclemys reevesii unicolor

Type locality: "China...Shanghai" [green star].

Type specimens: NHMUK 1946.1.22.46–47 and 1947.3.4.11 (formerly 1873.6.21.4, 1876.6.12.1–2), syntypes (3).

Etymology:* Latin, *unum*, *uni*, one, single, and *color*, color;

probably referring to the specimen examined having a uniform color.

Geoclemys grangeri Schmidt 1925:1

Geoclemys reevesii grangeri, Chinemys grangeri

Type locality: "Yenchingkao, Wanhsien, Szechwan, 1500 feet altitude" [Yanjinggou, Sichuan, China] [blue star].

Type specimen: AMNH 23481, holotype.

Etymology:* Eponym honoring Walter W. Granger (1872–1941), USA; paleontologist, curator at the American Museum of Natural History, New York, and collector of the type.

Geoclemys paracaretta Chang 1929:1

Geoclemys papacaretta

Type locality: "Fuchow" [Fuzhou Shi, Fujian, China] [pink star].

Type specimen: MBLSSC 134, holotype.

Etymology:* Greek, *para-*, beside, near, and Vernacular (Old French), *caret*, sea turtle; referring to the similarity to *Caretta*.

Chinemys megalocephala Fang 1934:158

Mauremys megalocephala, Chinemys megacephala

Type locality: "hill-sides of the vicinity of Nanking city" [Nanjing Shi, Jiangsu, China] [brown star].

Type specimen: Possibly MMNHN, holotype, originally live, figured (f.6).

Etymology:* Greek, *megalos*, great or large, and *kefale*, head; referring to the large head.

Chinemys macrocephala Bourret 1941c:140 (*nomen invalidum*)

Etymology:* Greek, *makros*, great or large, and *kefale*, head; referring to the large head.

Comment: Unjustified replacement name for *megalocephala*.

Chinemys pani † Tao 1985:45

Type locality: "Tsochen, Tsai-liao-chi, Tainan, Taiwan" [Zuojhen, Taiwan] [gray star].

Type specimen:* Private collection of Chang-Wu Pan, Tainan, Taiwan, holotype, fossil, nearly whole shell, figured (f.2-4), likely lost; NTUM R0001, cast of the holotype, likely lost; CJSHS 911001, NTM I02023, I02039, replica casts of the holotype (see Liaw and Tsai 2022).

Geological age:* Middle to Late Pleistocene, Chiting.

Etymology:* Eponym honoring Chang-Wu Pan (1932–2015), Taiwan; amateur paleontologist, fossil collector, and owner of the type.

Comment:* Synonymization with *Mauremys reevesii* confirmed by Liaw and Tsai (2022), who also noted that the type specimen was probably obtained from the sea bottom of the Taiwan Strait between Taiwan and the Penghu islands.

Mauremys pritchardi McCord 1998:555 (*partim*, hybrid) (07:33)

Type locality: "Lashio, Myanmar (97°14' E, 22°56' N)" [in error?] [orange star].

Type specimen: UF 105422, holotype.

Etymology:* Eponym honoring Peter C.H. Pritchard (1943–2020), United Kingdom (Northern Ireland) / USA; herpetologist and turtle researcher who founded Chelonian Research Institute, Florida.


Comment:* Type specimen likely acquired from the trade; type locality likely a trade location. Apparent hybrid of *Mauremys mutica mutica* and *Mauremys reevesii*.

Mauremys rivulata (Valenciennes *in* Bory de Saint-Vincent 1833)
(14:25, 17:50, 17:53)


Western Caspian Turtle, Balkan Terrapin



Apostolis Trichas / CBFTT / Crete, Greece



left: Georgia Mantziou / CBFTT / Almyros River, Irakleio, Crete, Greece
right: Apostolis Trichas / CBFTT / Crete, Greece



(orange dots = uncertain or introduced; stars = type localities)

Distribution:* Albania, Bosnia and Herzegovina, Bulgaria, Croatia, Cyprus (*possible historical introduction*), Greece, Israel, Jordan, Lebanon, Montenegro, North Macedonia, Palestine (*West Bank*), Syria, Turkey

Introduced: Latvia

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 330,174 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 1,439,834 sq. km

Size (Max SCL): male 21.5 cm, female 24.4 cm (Ayaz and Budak 2008; Mantziou and Rifai 2014 CBFTT)

CBFTT Account: Mantziou and Rifai (2014)

IUCN Red List:* Not Evaluated (NE). Regional (Europe): Least Concern (LC) (Luiselli and Vamberger 2024). Previously: Least Concern (LC) (Cox and Temple 2009); Least Concern (LC) (van Dijk, Lymberakis, Mousa Disi, Ajtic, Tok, Ugurtas, Sevinç, and Haxhiu 2004). Regional (European Union–EU27): Least Concern (LC) (Cox and Temple 2009)

TFTSG Provisional Red List: Least Concern (LC) (2011)

Synonymy:

Emys pannonica Michahelles *in* Bonaparte 1831:87 (*nomen nudum*)

Etymology:* Toponym for Pannonia, an historical province of the Roman Empire in eastern Europe south and west of the Danube River.

Emys rivulata Valenciennes *in* Bory de Saint-Vincent 1833:planches,pl.9 (senior homonym, not = *Emys rivulata* Gray 1844 [= *Pseudemys rubriventris*]) (14:25)

Clemmys caspica rivulata, Mauremys caspica rivulata, Mauremys rivulata, Mauremys rivulata rivulata, Emmenia rivulata

Type locality: Not designated. Restricted to "Morée...Modon" [Methoni, Greece] [yellow star] by Bory de Saint-Vincent (1836:110), and emended to "Umgebung von Modon, Morea, Griechenland" [Greece] by Mertens and Müller (1928:22).

Type specimen: MNHN 1930, holotype, see Fritz and Wischuf (1997).

Etymology:* Latin, *rivulus*, groove, *rivulata*, having grooves; referring to the carapace surface.

Emys tristrami Gray 1869a:190

Emys tristram, Mauremys rivulata tristrami

Type locality: "Holy Land" [Israel or Jordan]. Restricted to "Jabook River" [Zarqa River, Jordan] [green star, location estimated] by Gray (1873j:35); and to "Yarmuk River" [Israel or Jordan] [orange star, location estimated] by Boulenger (1889:104).

Type specimen: NHMUK 1947.3.5.37 (formerly 1864.8.23.156), holotype, see Fritz and Wischuf (1997).

Etymology:* Eponym honoring Henry B. Tristram (1822–1906), United Kingdom; clergyman, Bible scholar, and ornithologist, who collected the type.

Emys caspica arabica Gray 1870c:36

Emys arabica

Type locality: "Arabia Petraea?" Restricted to "Mt. Carmel" [Israel] [blue star] by Gray (1873j:34).

Type specimen: NHMUK 1946.1.22.48 (formerly 1864.8.25.51), holotype, see Fritz and Wischuf (1997).

Etymology:* Toponym for the Arabian Peninsula.

Emys pannonica Gray 1870c:36

Type locality:* "Xantos" [Xanthos, Turkey] [pink star].

Type specimen: NHMUK 1946.1.22.37, lectotype, designated by Fritz and Wischuf (1997:244).

Etymology:* Toponym for Pannonia, an historical province of the Roman Empire in eastern Europe south and west of the Danube River.

Clemmys caspica orientalis Bedriaga 1881:335

Type locality:* "Umgebung von Athen und von Nauplia, ferner auf den Inseln Milo, Mykonos, Syra, Tinos, Siphnos und Seriphos" [Greece]. Restricted here to "Athens, Greece" [brown star].

Type specimens: Bedriaga private collection, syntypes (2), apparently lost, see Fritz and Wischuf (1997).

Etymology:* Latin, *oriens*, the East, *orientalis*, coming from the East; referring to the provenance of the species from eastern Greece.

Clemmys caspica obsoleta Schreiber 1912:946

Type locality:* "südlichen Dalmatien...Bocche di Cattaro..[&].. Halbinsel Sabbioncello..[&]..Menge" [Croatia]. Restricted to "Bocche di Cattaro, Süd-Dalmatien" [Montenegro] [white star] by Wermuth and Mertens (1961:60).

Type specimens: Schreiber private collection, syntypes, apparently lost, see Fritz and Wischuf (1997).

Etymology:* Latin, *obsoleta*, worn, obsolete; referring to the apparent ontogenetic loss of stripes on the carapace in this taxon.

Clemmys caspica cretica Mertens 1946:115

Mauremys caspica cretica, Mauremys rivulata cretica

Type locality: "Rapaniana, Kreta" [Crete, Greece] [gray star].

Type specimen: SMF 36396, holotype, see Mertens (1967b) and Fritz and Wischuf (1997).


Etymology:* Toponym for the island of Crete, Greece.

Mauremys sinensis (Gray 1834a) (07:44)

Chinese Stripe-necked Turtle



John B. Iverson / No data / trade



John B. Iverson / No data / trade



** (orange dots = trade or introduced; stars = type localities)

Distribution:* China (*Fujian, Guangdong, Guangxi [?], Hainan*), Taiwan, Vietnam

Introduced:* Italy, Micronesia (*Guam, Pohnpei*), Réunion, Seychelles, South Korea, Spain, USA (*Florida*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 239,282 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 1,358,157 sq. km

Size (Max SCL):* male 20.0 cm, female 30.0 cm (McCord and Iverson 1994; Schilde 2004)

IUCN Red List: Critically Endangered (CR A2cd+4cd) (Li, Rao, and Wang 2021). Previously: Endangered (EN) (ATTWG 2000); Near Threatened (NT) (TFTSG *in* Baillie and Groombridge 1996)

CITES: Appendix III (China) (2005)

Synonymy:

Emys sinensis Gray 1834a:53

Graptemys sinensis, Clemmys sinensis, Ocadia sinensis, Ocadia sinensis sinensis, Mauremys sinensis

Type locality: "Chinâ." Restricted to "China; Canton" [Guangzhou, Guangdong] [yellow star] by Gray (1856b:21).

Type specimen: NHMUK 1947.3.5.26, holotype.

Etymology:* Toponym for China.

Emys bennettii Gray 1844:21

Clemmys bennettii

Type locality: "North America?" [in error]. Restricted to "China?" by Gray (1856b:22).

Type specimens: NHMUK 1947.3.4.24 and 1947.3.5.27 (formerly 1854.5.4.10, 1854.5.4.19), syntypes (2).

Etymology:* Eponym honoring Edward T. Bennett (1797–1836), United Kingdom; physician, zoologist, and secretary of the Zoological Society of London, who obtained the types.

Testudo anyangensis † Ping 1930:217

Ocadia anyangensis, Pseudocadia anyangensis, Mauremys anyangensis

Type locality: "ancient ruins in Annyang Hsien, Honan Province" [Anyang Shi, Henan, China] [dark gray star].

Type specimen: Probably NZMC s/n, holotype, not located, subfossil, whole shell, figured (f.1-3).

Geological age: Holocene, Neolithic.

Etymology:* Toponym for Annyang Hsien (= Anyang Shi), Henan, China.

Comment:* Synonymized by Pritchard (1994).

Ocadia sinensis changwui † Tao 1988:233

Type locality: "Penghu Channel in the Taiwan Strait" [Taiwan] [light gray star].

Type specimen: Private collection of Chang-Wu Pan, Tainan, Taiwan, holotype, fossil, partial carapace fragment, figured (f.1-2); NTUM F0076, cast of the holotype.

Geological age: Late Pleistocene.

Etymology:* Eponym honoring Chang-Wu Pan (1932–2015), Taiwan; fossil collector and owner of the type.

Ocadia philippeni McCord and Iverson 1992:13 (*partim*, hybrid) (07:33)

Type locality: "near Kancheng [18º51'N, 108º37'E; = 48 km from Tungfang (19º03'N, 108º56'E)], western Hainan Island, China" [in error?] [orange star].

Type specimen: UF 80766, holotype.

Etymology:* Eponym honoring Hans-Dieter Philippen (1957–2016), Germany; herpetologist and turtle researcher.

Comment:* Type specimen likely acquired from the trade; type locality likely a trade location. Apparent hybrid of *Cuora trifasciata luteocephala* and *Mauremys sinensis*. Previously listed as Data Deficient (DD) on the IUCN Red List (TFTSG *in* Baillie and Groombridge 1996).

Ocadia glyphistoma McCord and Iverson 1994:53 (*partim*, hybrid) (07:33)

Type locality: "near the Vietnam border southwest of Nanning, Guangxi Province, China" [in error?] [brown star].

Type specimen: UF 84818, holotype.

Etymology:* Greek, *glyphis*, carved or notched, and *stoma*, mouth; referring to the distinctly notched upper jaw.

Comment:* Type specimen likely acquired from the trade; type locality likely a trade location. Apparent hybrid of *Mauremys sinensis* and *Mauremys annamensis*. Previously listed as Data Deficient (DD) on the IUCN Red List (TFTSG *in* Baillie and Groombridge 1996).

Melanochelys Gray 1869a

Indian Black Turtles

(includes 2 species, 7 taxa)

Melanochelys Gray 1869a:187

Type species: *Melanochelys trijuga* [= *Emys trijuga* Schweigger 1812], by original monotypy.

Etymology:* Greek, *melanos*, black, dark, and *chelys*, turtle.

Chaibassia Theobald 1876:6

Type species: *Geomyda* (*Chaibassia) tricarinata* [= *Geomyda tricarinata* Blyth 1856], by original monotypy.

Etymology:* Toponym for Chaibasa, Jharkhand, India.

Melanochelys tricarinata (Blyth 1856) (59)*


Tricarinate Hill Turtle, Three-keeled Land Turtle



Indraneil Das / CBFTT / Dehradun, Uttarakhand, India



Indraneil Das / CBFTT / Dehradun, Uttarakhand, India



(orange dots = possible trade or uncertain; stars = type localities)

Distribution: Bangladesh, Bhutan, India (*Arunachal Pradesh, Assam, Bihar, Jharkhand, Uttar Pradesh, Uttarakhand, West Bengal*), Nepal

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 298,787 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 1,233,467 sq. km

Size (Max SCL): male 19.6 cm, female 18.5 cm (Baruah et al. 2012; Ceballos et al. 2013)

CBFTT Account: Das (2009)

IUCN Red List: Endangered (EN A4cd) (Horne, Praschag, Choudhury, and Singh 2020). Previously: Vulnerable (VU) (ATTWG 2000); Vulnerable (VU) (TFTSG *in* Baillie and Groombridge 1996)

CITES: Appendix I (1975)

USA ESA:* Endangered (USFWS 1976)

Synonymy:

Geomyda tricarinata Blyth 1856:714

Geomyda (Chaibassia) tricarinata, Geoemyda tricarinata,

Chaibassia tricarinata, Nicoria tricarinata, Melanochleys tricarinata

Type locality: "Central India (Chaibása)." Restricted to "Chybassa, in the District of Singhbhum, Chota Nagpur, Bengal" [Chaibasa, Jharkhand, India] [yellow star] by Anderson (1879:717).

Type specimen: ZSI 816, holotype, see Das et al. (1998), Das (2009), and Kundu et al. (2018a).

Etymology:* Latin, *tri-*, three, and *carina*, keel, *carinata*, keeled; referring to the three carapacial vertebral keels.

Geoemyda carinata Blyth *in* Jerdon 1870:69 (*nomen novum*)

Etymology:* Latin, *carina*, keel, *carinata*, keeled; referring to the carapacial vertebral keel.

Comment: Unjustified replacement name for *tricarinata*.


Chaibassia theobaldi Anderson 1879:718

Type locality: "Bishnath Plain, near Tezpur, in Assam, close to the Brahmaputra" [Assam, India] [green star].

Type specimens: ZSI 188–89, syntypes (2), see Das et al. (1998), 189 not located by Kundu et al. (2018a).

Etymology:* Eponym honoring William J. Theobald (1829–1908), United Kingdom / India; naturalist, herpetologist, and deputy superintendent of the Geological Survey of India.

Melanochelys trijuga (Schweigger 1812) [59, 71]*
Indian Black Turtle
(includes 6 subspecies)



** (subspecies: *trijuga* = red, *coronata* = blue, *edeniana* = green, *indopeninsularis* = pink, *parkeri* = brown, *thermalis* = tourmaline; orange dots = introduced or trade)

Distribution:* Bangladesh, India (*Andhra Pradesh, Assam, Bihar, Goa, Gujarat, Karnataka, Kerala, Maharashtra, Meghalaya, Mizoram, Nagaland, Odisha, Tamil Nadu, Telangana, Tripura, Uttar Pradesh, West Bengal*), Myanmar, Nepal, Sri Lanka, Thailand

Introduced: British Indian Ocean Territory (*Chagos Archipelago*), Maldives

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 1,511,818 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 5,163,214 sq. km

Size (Max SCL): male 38.3 cm, female 26.2 cm (see subspp.)

CBFTT Account: Das and Bhupathy (2009b)


IUCN Red List: Least Concern (LC) (Ahmed, Praschag, de Silva, Das, Singh, and de Silva 2020). Previously: Near Threatened (NT) (ATTWG 2000); Data Deficient (DD) (TFTSG *in* Baillie and Groombridge 1996)

CITES: Appendix II (2013)


Melanochelys trijuga trijuga (Schweigger 1812) [21:72] [71]*
Indian Black Turtle



S. Jayakumar / CBFTT / Anaikatti Hills, Coimbatore, Western Ghats, Tamil Nadu, India



S. Jayakumar / CBFTT / Anaikatti Hills, Coimbatore, Western Ghats, Tamil Nadu, India



** (subspecies: *trijuga* = red, *coronata* = blue, *indopeninsularis* = pink, *thermalis* = tourmaline; stars = type localities)

Distribution:* India (*Andhra Pradesh, Goa, Gujarat, Karnataka, Maharashtra, Tamil Nadu, Telangana*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 581,201 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 645,301 sq. km

Size (Max SCL): No data located; **(Max CCL):** male 25.0 cm, female 22.0 cm (Mukherjee et al. 2006; Das and Bhupathy 2009b CBFTT)

CITES: Appendix II (2013)

Synonymy:

Emys trijuga Schweigger 1812:310 [21:72]

Clemmys (Clemmys) trijuga, Clemmys trijuga, Melanochelys trijuga, Nicoria trijuga, Geoemyda trijuga, Geoemyda trijuga trijuga, Melanochelys trijuga trijuga

Type locality: "insula Java" [Indonesia] [no star] [21:72].

Type specimen: MNHN, holotype, apparently lost, see Das (2009); specimen figured by Oppel *in* Bour and Schmidtler (2014:f.9,pl.A,B), identified as a *Malayemys subtrijuga*.

Etymology:* Latin, *tria*, three, and *jugum*, yoke; referring to the three longitudinal keels on the carapace.

Emys herrmanni Schweigger 1812:311 [21:72]

Emys hermanni, Clemmys (Clemmys) hermanni

Type locality: Not known. Restricted to "India" by Gray (1873j:30).

Type specimens: Possibly in MZUS, syntype specimen figured by Duvernoy (1838:pl.5), reproduced by Bour and Schmidtler (2014:f.9,C).

Etymology:* Eponym honoring Johann Hermann (or Herrmann) (1738–1800), France; zoologist at the University of Strasbourg in Alsace, who first recognized this species as distinct.

Emys belangeri Lesson 1831b:291 (21:72) (71)*
Type locality:* "Bengale..[&]..Carnate" [India]. Restricted to "Bengal" [India] [in error; likely nr. Calcutta, location estimated; orange star] by Smith (1931:97); restricted in our annotation 67 to "Carnate (= Carnatic region), nr. Pondicherry, Tamil Nadu, India]" [yellow star].
Type specimen:* MNHN, holotype, not located, specimen figured (pl.1), reproduced by Bour and Schmidtler (2014:f.9,E), designated lectotype in our annotation 67.
Etymology:* Eponym honoring Charles P. Bélanger (1805–1881), France; botanist and mycologist who collected the type.
Comment: Type collected by Bélanger while stationed in the French port of Pondicherry in the Carnate (Carnatic) region of southeast India.


Emys trijuga maderaspatana Anderson 1879:729
Type locality: "Madras" [Chennai, Tamil Nadu, India] [green star].
Type specimens: ZSI 831, 1008–9, syntypes (3), see Das et al. (1998) and Kundu et al. (2018a), listed as *Emys trijuga madraspatana*.
Etymology:* Toponym for Madrasapattinam (= Chennai), Tamil Nadu, India; long form of the name "Madras" previously used for this city.


Emys trijuga madraspatana Annandale 1913:67 (*nomen novum*)
Geoemyda trijuga madraspatana
Etymology:* Toponym for Madrasapattinam (= Chennai), Tamil Nadu, India; long form of the name "Madras" previously used for this city.
Comment: Unjustified emendation of *maderaspatana*.

Geoemyda trijuga plumbea Annandale 1915a:192
Type locality: "Coorg on the east side of the Western Ghats considerably east and a little south of the Madras district" [= Coorg Province, nr. Mercara (= Madikeri), Karnataka, India] [blue star; location estimated].
Type specimens: ZSI 17712, 17715, syntypes (2), see Das et al. (1998); 17715 not located by Kundu et al. (2018a).
Etymology:* Latin, *plumbum*, lead, *plumbea*, made of lead, leaden; referring to the lead-gray color of the soft parts.

Melanochelys trijuga coronata (Anderson 1879)
Cochin Black Turtle


Subramanian Bhupathy / CBFTT / Kumarakom, Kerala, India


Václav Gvoždík / Mudumalai, Tamil Nadu, India


**
(subspecies: *trijuga* = red, *coronata* = blue, *parkeri* = brown, *thermalis* = tourmaline; overlap = intergrades; star = type locality)

Distribution: India (*Kerala*, *Tamil Nadu*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 49,640 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 82,630 sq. km

Size (Max SCL): No data located; **(Max CCL):** male 26.0 cm, female 23.0 cm (Mukherjee et al. 2006; Das and Bhupathy 2009b CBFTT)

CITES: Appendix II (2013)

Synonymy:

Emys trijuga coronata Anderson 1879:729
Nicoria trijuga coronata, *Geoemyda trijuga coronata*, *Melanochelys trijuga coronata*
Type locality: "Southern India...Travancore" [nr. Trivandrum (= Thiruvananthapuram), Kerala, India] [yellow star, location estimated].
Type specimen: ZSI 1012, holotype, see Das et al. (1998), Das (2009), and Kundu et al. (2018a).
Etymology:* Latin, *corona*, crown, *coronata*, having a crown; referring to the diamond-shaped pattern on top of the head.

Melanochelys trijuga edeniana Theobald 1876 (07:47)
Burmese Black Turtle, Myanmar Black turtle




Rick Hudson / Myanmar / captivity / Yadanabon Zoo





left: Indraneil Das / CBFTT / Myanmar
right: Cris Hagen / CBFTT / Myanmar




(subspecies: *edeniana* = green, *indopeninsularis* = pink; orange dot = probable trade; stars = type localities)

Distribution: Myanmar, Thailand
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 204,721 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 421,674 sq. km
Size (Max SCL): 28.0 cm (Das and Bhupathy 2009b CBFTT)
CITES: Appendix II (2013)
Synonymy:
Melanochelys edeniana Theobald 1876:12
Nicoria trijuga edeniana, Geoemyda trijuga edeniana, Melanochelys trijuga edeniana, Emys trijuga edeniana, Melanochelys edeniana edeniana
Type locality: "Inhabits Arakan, Pegu, and Tenasserim [Provinces]. A female taken near Tounghu" [= Taungoo, Myanmar] [yellow star].
Type specimens: ZSI 830, 1010–11, 1018, 1097, 1369, 1371, syntypes (7), see Das et al. (1998) and Kundu et al. (2018a).
Etymology:* Eponym honoring Sir Ashley Eden (1831–1887), United Kingdom / India / Myanmar; official and diplomat in British India and first civil governor of Burma (now Myanmar).
Emys trijuga burmana Anderson 1879:723
Type locality: "Bhamô..[&]..Moulmein..[&]..Khyouk-Phyoo in Arracan" [Myanmar]. Restricted to "Bhamo, Burma" [Myanmar] [green star] by Smith (1931:98).
Type specimens: ZSI 830, syntype (1), others apparently lost, see Das et al. (1998) and Kundu et al. (2018a).
Etymology:* Toponym for Burma (= Myanmar).
Geoemyda trijuga wiroti Reimann *in* Nutaphand 1979:177
Melanochelys trijuga wiroti, Melanochelys edeniana wiroti
Type locality: "Thai-Burmese border area (Tak and Mae Hong Son Provinces)" [Thailand] [blue star, location estimated].
Type specimen: Not located, possibly in ZFMK, type specimen figured (f.66-67).
Etymology:* Eponym honoring Wirot Nutaphand (or Nutphand) (1932–2005), Thailand; airforce officer, naturalist, and herpetologist.

Melanochelys trijuga indopeninsularis (Annandale 1913) (71)*
Bengal Black Turtle



Bhaba Amatya / CBFTT / Rupa Lake, Pokhara, Nepal



(subspecies: *trijuga* = red, *edeniana* = green, *indopeninsularis* = pink; star = type locality)

Distribution:* Bangladesh, India (*Assam, Bihar, Jharkhand, Meghalaya, Mizoram, Nagaland, Odisha, Tripura, Uttar Pradesh, West Bengal*), Nepal
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 600,281 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 1,291,299 sq. km
Size (Max SCL): male 34.2 cm, female 26.2 cm (Farkas and Csorba 1999; Das and Bhupathy 2009b CBFTT)
CITES: Appendix II (2013)
Synonymy:
Geoemyda indopeninsularis Annandale 1913:71
Geoemyda trijuga indopeninsularis, Melanochelys trijuga indopeninsularis, Melanochelys edeniana indopeninsularis
Type locality: "Singhbhum district of Chota Nagpur..[&]..Dharwar district in...Bombay Presidency...at Devikop, 26 miles south of... Dharwar" [India]. Restricted to "Singhbhum District, Chota Nagpur" [Jharkhand, India] [yellow star, location estimated] by Smith (1931:99).
Type specimens: ZSI 17098, 17100, syntypes (2), see Das et al. (1998), Das (2009), and Kundu et al. (2018a).
Etymology:* Toponym for India, and Latin, *pensilis*, hanging down, *insula*, an island, and *-aris*, pertaining to; referring to the distribution of the species on the Indian Peninsula.


Melanochelys trijuga parkeri Deraniyagala 1939 (17:54)
Parker's Black Turtle



Indraneil Das / CBFTT / Sri Lanka



Anslem de Silva / Sri Lanka



** (subspecies: *parkeri* = brown, *thermalis* = tourmaline; stars = type localities)

Distribution: Sri Lanka

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 12,728 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 17,080 sq. km

Size (Max SCL): 38.3 cm (Deraniyagala 1939; Das and Bhupathy 2009b CBFTT)

CITES: Appendix II (2013)

Synonymy:


Melanochelys trijuga parkeri Deraniyagala 1939:269
Geoemyda trijuga parkeri
Type locality: "Polonnaruva, N.-C.P., ...Ceylon" [Sri Lanka] [yellow star].
Type specimen: NHMUK 1947.3.6.14 (formerly 1937.10.3.1), holotype.
Etymology:* Eponym honoring Hampton W. Parker (1897–1968), United Kingdom; herpetologist at the British Museum of Natural History, London, who first recognized the taxon as distinctive.

Melanochelys trijuga sinhaleyus † Deraniyagala 1953:4
Geoemyda trijuga sinhaleya, Melanochelys sinhaleyus
Type locality: "near Ratnapura...Ceylon" [Sri Lanka] [gray star].
Type specimens: RNM F80, F93, syntypes, fossil, carapacial bones, nuchal figured (f.2).
Etymology:* Toponym for Sinhala [Sanskrit name for Ceylon = Sri Lanka].
Geological age: Late Pleistocene, Ratnapura Beds.
Comment:* This taxon was described from fragmentary carapacial material showing slight differences from modern *Melanochelys trijuga parkeri*. Its taxonomic status is uncertain; we listed it as a possibly distinct species in TEWG (2015), pending further analysis, but now provisionally synonymize it here.


Melanochelys trijuga thermalis (Lesson 1830)
Sri Lanka Black Turtle



Peter Paul van Dijk / CBFTT / nr. Bentota, Sri Lanka



left: Peter Paul van Dijk / CBFTT / nr. Bentota, Sri Lanka
right: Anslem de Silva / Sri Lanka



** (subspecies: *trijuga* = red, *coronata* = blue, *parkeri* = brown, *thermalis* = tourmaline; overlap = intergrades; star = type locality)

Distribution: India (*Tamil Nadu*), Sri Lanka

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 65,164 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 119,495 sq. km

Size (Max SCL):* male 25.6 cm, female 22.9 cm (Das and Bhupathy 2009b CBFTT; de Silva et al. 2022)

CITES: Appendix II (2013)

Synonymy:

Emys thermalis Lesson 1830:86
Clemmys thermalis, Nicoria trijuga thermalis, Geoemyda trijuga thermalis, Melanochelys trijuga thermalis
Type locality: "eaux thermales de Cannia, près Trinquemalé à Ceylan," [hot springs of Kanniya, nr. Trincomalee, Sri Lanka] [yellow star].
Type specimen: ZSI, not located, apparently lost; holotype specimen figured (f.29), see Das (2009) and Bour and Schmidtler (2014:f.9;D).
Etymology:* Greek, *therme*, warm source, warmth, heat, *thermalis*, belonging to a warm spring; referring to provenance of this species from thermal springs.

Emys sebae Gray 1831d:75
Geoclemys sebae, Melanochelys sebae, Emys trijuga sebae
Type locality: Not designated. Restricted to "Ceylon" [Sri Lanka] [no star] by Gray (1844:19).
Type specimen: Specimen figured in Seba (1734:pl.79,f.1-2), holotype, by original designation.

Etymology:* Eponym honoring Albertus Seba (1665–1736), Netherlands; pharmacist and naturalist with a large natural history collection.

Emys seba Gray 1844:19 (*nomen novum*)

Geoclemys seba, Geoclemmys seba

Etymology:* Eponym honoring Albertus Seba (1665–1736), Netherlands; pharmacist and naturalist with a large natural history collection.

Comment: Unjustified emendation of *Emys sebae*.

Notochelys Gray 1863e

Flat-shelled Turtles

(includes 1 species, 1 taxon)

Notochelys Gray 1863e:177 (senior homonym, not = *Notochelys* Owen 1882 [= Testudines: Protostegidae †])

Type species: *Notochelys platynota* [= *Emys platynota* Gray 1834a], by original monotypy.

Etymology:* Greek, *notos*, the back, and *chelys*, turtle.

Notochelys platynota (Gray 1834a)


Malayan Flat-shelled Turtle



Sabine Schoppe / Malaysia / trade



Sabine Schoppe / Malaysia / trade



(orange dots = probable trade; stars = type localities)

Distribution: Brunei, Indonesia (*Java*, *Kalimantan*, *Sumatra*), Malaysia (*East*, *West*), Myanmar, Singapore, Thailand

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 688,017 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 3,522,719 sq. km

Size (Max SCL): male 28.2 cm, female 33.0 cm (Brophy and Ernst 2004)

IUCN Red List: Vulnerable (VU A2cd) (Kusrini, Hamidy, Guntoro, Cota, and Schoppe 2021). Previously: Vulnerable (VU) (ATTWG 2000); Data Deficient (DD) (TFTSG *in* Baillie and Groombridge 1996)

CITES: Appendix II (2005)

Synonymy:

Emys platynota Gray 1834a:54

Emys platynotha, Cyclemys platynota, Notochelys platynota

Type locality: "Indiâ Orientali." Restricted to "Sumatra" [Indonesia] [yellow star, location estimated] by Gray (1835:pl.57).

Type specimen: NHMUK 1947.3.4.6, holotype.

Etymology:* Greek, *platys*, flat, broad, or wide, and *notos*, back; referring to the flat-topped carapace.

Cistudo bankanensis Bleeker *in* Gray 1864a:12

Type locality: "Banka Island" [Bangka Island, Sumatra, Indonesia] [green star, location estimated].

Type specimen: NHMUK 1946.1.22.64 (formerly 1863.12.4.120), see Gray (1873j).

Etymology:* Toponym for Banka Island (= Bangka Island, Sumatra, Indonesia).

Cyclemys giebelii Hubrecht 1881:45

Type locality: "Borneo" [Kalimantan, Indonesia] [blue star, location estimated, possibly nr. Pontianak].

Type specimens: RMNH 3348, syntype (1), others apparently lost, discussed by Hoogmoed et al. (2010).

Etymology:* Eponym honoring Christoph G.A. Giebel (1820–1881), Germany; zoologist and paleontologist at the University of Halle.

Sacalia Gray 1870c

Asian Eyed Turtles

(includes 2 species, 2 taxa)

Sacalia Gray 1870c:35

Type species: *Sacalia bealii* [= *Emys bealii* Gray 1834a = *Cistuda bealei* Gray 1831d], by original monotypy.


Etymology:* Anonym, *sacalia*, possibly a fabricated name, or possibly an Eponym honoring someone named Sacal or Sakal, or starting with Sacal- or Sakal-.

Sacalia bealei (Gray 1831d) [(72)*]


Beale's Eyed Turtle



John B. Iverson / No data / trade



John B. Iverson / No data / trade



** (star = type locality)

Distribution: China (*Fujian*, *Guangdong*, *Hong Kong*, *Jiangxi*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 273,959 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 664,481 sq. km

Size (Max SCL): male 14.1 cm, female 15.6 cm (Pope 1935; Iverson and McCord 1992; Lin et al. 2018b)

IUCN Red List: Endangered (EN A1d+2d [ver. 2.3]) (ATTWG 2000). Previously: Vulnerable (VU) (TFTSG *in* Baillie and Groombridge 1996)

TFTSG Provisional Red List: Critically Endangered (CR) (2011)

CITES: Appendix II (2013). Previously: Appendix III (China) (2005)

Synonymy:

Cistuda bealei Gray 1831d:71

Emys (Pyxidemys) bealei, *Emys bealei*, *Clemmys bealei*, *Cistudo bealei*, *Sacalia bealei*, *Mauremys bealei*, *Sacalia bealei bealei*

Type locality: "China" [yellow star, possibly near Macao, location estimated].

Type specimens: NHMUK 1947.3.4.33, 1947.3.4.42, syntypes (2).

Etymology:* Eponym honoring Thomas Beale (1775–1842), United Kingdom (Scotland) / China; naturalist and merchant who lived and worked in China in Macao and obtained the type.

Emys bealii Gray 1834a:54 (*nomen novum*)

Clemmys bealii, *Sacalia bealii*, *Cistudo bealii*, *Clemmys bealii bealii*

Etymology:* Eponym honoring Thomas Beale (1775–1842), United Kingdom (Scotland) / China; naturalist and merchant who lived and worked in China and obtained the type.


Comment: Unjustified emendation or error for *bealei*.

Sacalia quadriocellata (Siebenrock 1903a) [(08:24, 21:73) (72)*]


Four-eyed Turtle



Haitao Shi / Guangdong, China



John B. Iverson / No data / trade



** (orange dot = probable trade; stars = type localities)

Distribution: China (*Guangdong*, *Guangxi*, *Hainan*), Laos, Vietnam

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 304,229 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 952,792 sq. km

Size (Max SCL): male 14.3 cm, female 15.2 cm (Iverson and McCord 1992; de Bruin and Artner 1999; Ceballos et al. 2013)

IUCN Red List: Critically Endangered (CR A2cd+4cd) (Fong, Hoang, Li, McCormack, Rao, Shi, and Wang 2021).

Previously: Endangered (EN) (ATTWG 2000); Vulnerable (VU) (TFTSG *in* Baillie and Groombridge 1996)

CITES: Appendix II (2013). Previously: Appendix III (China) (2005)

Synonymy:

Clemmys bealii quadriocellata Siebenrock 1903a:336 (senior homonym, not = *Clemmys bealii quadriocellata* Li 1958)

Clemmys quadriocellata, Clemmys quadriocellata, Clemmys bealei quadriocellata, Sacalia quadriocellata, Sacalia bealei quadriocellata, Sacalia quadriocellata quadriocellata

Type locality: "Annam" [Vietnam]. Restricted to "Annam, Phuc-Son" [= Phuoc Son, Quang Nam, Vietnam (15°33'00"N; 108°04'00"E); see Le et al. 2004] [yellow star] by Siebenrock (1909a:482).

Type specimen: NMW 23393, holotype, see Tiedemann and Häupl (1980), Tiedemann et al. (1994), and Gemel et al. (2019).

Etymology:* Latin, *quattuor*, four, and *ocellata*, having eye spots, ocellated; referring to the four ocellated markings on the head.

Clemmys bealii quadriocellata Li 1958:235 (junior homonym, not = *Clemmys bealii quadriocellata* Siebenrock 1903a)

Type locality: "small stream of Dali village of Mt. Diaoluo, Lingshui Co., Hainan Island, China" [green star] [in Chinese].

Type specimen: Not located, holotype specimen figured (f.2).

Etymology:* Latin, *quattuor*, four, and *ocellata*, having eye spots, ocellated; referring to the four ocellated markings on the head.

Clemmys beali insulensis Adler 1962:135 (*nomen novum*) (08:24, 21:73) (72)*

Clemmys bealii insulensis, Sacalia insulensis

Etymology:* Latin, *insula*, an island; referring to the distribution of the species on Hainan Island.

Comment: Justified replacement name for *Clemmys bealii quadriocellata* Li 1958.

Sacalia pseudocellata Iverson and McCord 1992a:426 (*partim*, hybrid) (07:33)

Type locality: "between Tungfang [19º03'N, 108º56'E] and Kancheng [18º51'N, 108º37'E; ca. 48 km from Tungfang], western Hainan Island, China" [orange star] [in error?].

Type specimen: UF 81505, holotype.

Etymology:* Greek, *pseudo*, false, and Latin, *ocellata*, eyed; referring to the lack of black eye spots on the neck.

Comment:* Type specimen likely acquired from the trade; type locality likely a trade location. Apparent hybrid of *Cuora trifasciata luteocephala* and *Sacalia quadriocellata*. Previously listed as Data Deficient (DD) on the IUCN Red List (TFTSG *in* Baillie and Groombridge 1996).

Sacalia quadriocellata insularis Artner 2003:xviii (*nomen invalidum*)

Etymology:* Latin, *insula*, an island, and *-aris*, pertaining to; referring to the distribution of the species on Hainan Island.

Comment: Unjustified emendation or error for *insulensis*.

Vijayachelys Praschag, Schmidt, Fritzsch, Müller, Gemel, and Fritz 2006 (07:51)

Cane Turtles

(includes 1 species, 1 taxon)

Vijayachelys Praschag, Schmidt, Fritzsch, Müller, Gemel, and Fritz 2006:156

Type species: *Vijayachelys silvatica* [= *Geoemyda silvatica* Henderson 1912], by original designation.


Etymology:* Eponym honoring Jagannathan Vijaya (1959–1987), India; female herpetologist who studied the turtles of this genus, and Greek, *chelys*, turtle.

Vijayachelys silvatica (Henderson 1912)

Cochin Forest Cane Turtle




Veerappan Deepak / CBFTT / Anaimalai Hills, Western Ghats, Tamil Nadu, India / male





left: Peter Praschag / CBFTT / Tamil Nadu, India / male
right: Veerappan Deepak / CBFTT / Anaimalai Hills, Tamil Nadu, India / female



** (orange dots = probable trade; star = type locality)

Distribution: India (*Karnataka, Kerala, Tamil Nadu*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 27,364 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 60,369 sq. km

Size (Max SCL): male 13.4 cm, female 17.0 cm (Whitaker and Vijaya 2009; Deepak et al. 2014 CBFTT)

CBFTT Account: Deepak, Praschag, and Vasudevan (2014)

IUCN Red List: Endangered (EN C1) (Choudhury and Praschag 2021). Previously: Endangered (EN) (ATTWG 2000); Endangered (EN) (TFTSG *in* Baillie and Groombridge 1996)

CITES: Appendix II (2013)

Synonymy:

Geoemyda silvatica Henderson 1912:217

Heosemys silvatica, Vijayachelys silvatica

Type locality: "South India...near Kavalai in the Cochin State Forests... an elevation of about 1500 feet" [Kerala, India] [yellow star].

Type specimen: ZSI 17117, holotype, see Das et al. (1998) and Das (2009), specimen not located by Kundu et al. (2018a).

Etymology:* Latin, *silvatica*, belonging to the forest; referring to the forest-dwelling habitat of this species.

Subfamily Rhinoclemmydinae Gray 1873j (12:21) (73)*

Neotropical Wood Turtles

(includes 9 species, 14 taxa)

Rhinoclemmyina Gray 1873j:27

Rhinoclemminae Le and McCord 2008:763

Rhinoclemmydinae van Dijk, Iverson, Shaffer, Bour, and Rhodin [Turtle Taxonomy Working Group] 2012:274

Rhinoclemmyinae Frétey 2024:67

Rhinoclemmys Fitzinger 1835 (09:20, 12:21, 14:31)

Neotropical Wood Turtles

(includes 9 species, 14 taxa)

Chemelys Rafinesque 1815:75 (*nomen nudum*)

Etymology:* Greek, *chelys*, turtle, and *emys*, turtle; possibly a fabricated combination of the two names as a play on words by Rafinesque.

Chemelys Rafinesque 1832:64 (*nomen suppressum*)

Type species: *Chemelys verrucosa* [= *Testudo verrucosa* Suckow 1798 = subjective synonym of *Testudo punctularia* Daudin 1801], by original monotypy.

Etymology:* Greek, *chelys*, turtle, and *emys*, turtle; possibly a fabricated combination of the two names as a play on words by Rafinesque.

Comment: Name suppressed by ICZN (1985a), see Smith et al. (1980c).

Rhinoclemmys Fitzinger 1835:115 (*nomen protectum*)

Type species: *Geoemyda (Rhinoclemmys) dorsata* [= *Testudo dorsata* Schoepff 1801 = subjective synonym of *Testudo punctularia* Daudin 1801], by subsequent designation as *Emys dorsata* sensu Schweigger 1812 by Lindholm (1929:283). Genus established as *Clemmys (Rhinoclemmys)* without a type species.

Etymology:* Greek, *rhinos*, nose, and *klemmys*, turtle, or *kleptein*, to steal, and *emys*, turtle; referring to the prominent nose and ability to retract and hide head and neck.

Comment: Name conserved by ICZN (1985a), see Smith et al. (1980c).

Rhinoclemys Gray 1863c:182 (*nomen novum*)

Etymology:* Greek, *rhinos*, nose, and *klemmys*, turtle, or *kleptein*, to steal, and *emys*, turtle; referring to the prominent nose and ability to retract and hide head and neck.

Callopsis Gray 1863c:183

Type species: *Rhinoclemys* (*Callopsis) annulata* [= *Geoclemmys annulata* Gray 1860b], by original monotypy.

Etymology:* Greek, *kallos*, beauty, and *opsis*, appearance, resemblance; referring to the attractive stripes on the head and neck.

Rhinoclemmys annulata (Gray 1860b)


Brown Wood Turtle, *Montañero, Bambera*



John L. Carr / Isla Palma, Valle del Cauca, Colombia



John L. Carr / Isla Palma, Valle del Cauca, Colombia



(orange dots = probable trade; stars = type localities)

Distribution: Colombia (*Antioquia, Cauca, Chocó, Córdoba, Nariño, Valle del Cauca*), Costa Rica, Ecuador, Honduras, Nicaragua, Panama

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 302,983 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 1,365,514 sq. km

Size (Max SCL): male 20.2 cm, female 22.6 cm (Ernst 1978; Giraldo et al. 2012)

CBFTT Account:* Carr, Moll, Giraldo, and Garcés-Restrepo (2023)

IUCN Red List:* Near Threatened (NT A2c+4c) (Carr and Giraldo, in press). Previously: TFTSG Provisional Red List: Data Deficient (DD) (2011, 2018); Near Threatened (NT) (TFTSG 1996)

CITES:* Appendix II, as *Rhinoclemmys* spp. (2023)

Synonymy:

Geoclemmys annulata Gray 1860b:231

Clemmys annulata, Rhinoclemys annulata, Rhinoclemys (Callopsis) annulata, Chelopus annulatus, Rhinoclemmys annulata, Nicoria annulata, Geoemyda annulata, Callopsis annulata

Type locality: “Esmeraldas, ...western coast of Ecuador” [yellow star].

Type specimens: NHMUK 1946.1.22.56, 1947.3.5.58–59 (formerly (formerly 1860.4.22.23–25), syntypes (3).

Etymology:* Latin, *annulatus*, furnished or ornamented with a ring; referring to the growth rings on the carapacial scutes.

Chelopus gabbii Cope 1875a:153 (21:6)

Emys gabbii, Nicoria gabbii, Geoemyda gabbii, Rhinoclemmys gabbii

Type locality: “Costa Rica.” Restricted to “Sipurio, Costa Rica” [green star] by Dunn (1930:32).

Type specimen: USNM 45905, holotype, see Cochran (1961) and Reynolds et al. (2007).

Etymology:* Eponym honoring William M. Gabb (1839–1878), USA / Costa Rica; paleontologist and geographer who worked in Costa Rica and collected the type.

Testudo mercatoria Vaillant 1911:47 (*nomen nudum*)

Type locality: “Amérique centrale.”

Type specimen: Not known or located.

Etymology:* Latin, *mercatoria*, commercial; referring to the provenance of the examined specimen from a commercial animal dealer.


Rhinoclemmys areolata (Duméril and Bibron *in* Duméril and Duméril 1851)
Furrowed Wood Turtle, *Mojina*



John B. Iverson / CBFTT / nr. Paamul, Quintana Roo, Mexico



left: Eduardo Reyes-Grajales / Palenque, Chiapas, Mexico
right: James H. Harding / CBFTT / captivity



** (orange dots = probable trade; stars = type localities)

Distribution:* Belize, Guatemala, Honduras (?), Mexico (*Campeche, Chiapas, Oaxaca, Quintana Roo, Tabasco, Veracruz, Yucatán*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 289,973 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 525,944 sq. km

Size (Max SCL):* male 18.8 cm, female 24.0 cm (Ernst 1978; Legler and Vogt 2013; Charruau et al. 2014; Charruau and Morales-Garduza 2022)

CBFTT Account: Vogt, Platt, and Rainwater (2009)

IUCN Red List:* Least Concern (LC) (Butterfield, Monsiváis-Molina, Díaz-Gamboa, Willey, Jones, and Platt, in press). Previously: Near Threatened (NT) (van Dijk, Lee, Calderón Mandujano, Flores-Villela, and López-Luna 2007); Least Concern (LC) [Not Listed] (Behler 1996)

CITES:* Appendix II, as *Rhinoclemmys* spp. (2023)

Synonymy:

Emys areolata Duméril and Bibron *in* Duméril and Duméril 1851:10

Malaclemmys concentrica areolata, Clemmys areolata, Malaclemys concentrica areolata, Chelopus areolatus, Nicoria punctularia areolata, Geoemyda punctularia areolata, Geoemyda areolata, Rhinoclemmys areolata, Callopsis areolata

Type locality: "Province du Petén (Amér. centr.)." Restricted to "La Libertad, El Petén, Guatemala" [orange star] by Smith and Taylor (1950a:318; 1950b:30); and to "Flores...El Petén, La Libertad... Guatemala" [yellow star] by Dunn and Stuart (1951:60).


Type specimen: MNHN 9424, holotype, see Ernst (1978, 1980a) and Bour (2007b); MNHN 8310, listed as "type" by Stuart (1963), is not a type.

Etymology:* Latin, *areola*, small open space; referring to the pattern on the carapacial scutes.


Rhinoclemmys diademata (Mertens 1954)
Maracaibo Wood Turtle, *Galapago Negro*



Carlos A. Galvis-Rizo / Colombia / captivity



Thomas S.B. Akre / Catatumbo, Zulia, Venezuela



** (orange dots = probably introduced or trade; stars = type localities)

Distribution: Colombia (*Norte de Santander*), Venezuela (*Mérida, Táchira, Trujillo, Zulia*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 38,711 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 58,878 sq. km

Size (Max SCL): male 18.7 cm, female 28.5 cm (Pritchard and Trebbau 1984; Rivas et al. 2007)

IUCN Red List:* Endangered (EN A2cde+4cde) (Morales-Betancourt, Lasso, and Rojas-Runjaic 2024). Previously: TFTSG Provisional Red List: Vulnerable (VU) (2011); Least Concern (LC) [Not Listed] (Behler 1996)

CITES:* Appendix II, as *Rhinoclemmys* spp. (2023)

Synonymy:

Geoemyda punctularia diademata Mertens 1954:4

Callopsis punctularia diademata, Rhinoclemmys punctularia diademata, Rhinoclemmys diademata

Type locality: "Maracay, Venezuela" [orange star, in error]. Considered erroneous by Pritchard (1979:182); and considered in error for "Maracaibo" [Venezuela] [yellow star] by Pritchard and Trebbau (1984:172).

Type specimen: SMF 48141, holotype, see Mertens (1967b).

Etymology:* Greek, *diadima*, a diadem; referring to the distinctive yellow, diadem-like figure behind the eyes on top of the head.

Rhinoclemmys funerea (Cope 1875a) (21:6)
Black Wood Turtle, *Jicotea Negra*



Manuel Merchán Fornelino / Costa Rica





Nathanael Stanek / Guapiles, Limón Prov., Costa Rica



** (orange dots = probable trade or translocated or misidentified; star = type locality)

Distribution: Costa Rica, Honduras, Nicaragua, Panama

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 71,823 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 198,690 sq. km

Size (Max SCL):* male 37.5 cm, female 35.5 cm (Ernst 1980b; Savage 2002; Folt 2020; R. Acuña-Mesén and D. Moll, unpubl. data)

IUCN Red List:* Least Concern (LC) (Folt, van Dijk, Rhodin, and Such, in press). Previously: Near Threatened (NT) (TFTSG 1996)

CITES:* Appendix II, as *Rhinoclemmys* spp. (2023)

Synonymy:

Chelopus funereus Cope 1875a:154 (21:6)
Emys funerea, Geoemyda funerea, Geoemyda punctularia funerea, Rhinoclemmys funerea, Callopsis funerea
Type locality: "Costa Rica...Limon" [Puerto Limon] [yellow star].
Type specimens: USNM 45900–01, 46134–35, syntypes (4), see Cochran (1961) and Reynolds et al. (2007); Uetz et al. (2019) erroneously listed USNM 45000.
Etymology:* Latin, *funeris*, death, corpse, funeral; referring to the black carapace.

Geoemyda costaricensis Kanberg 1930:162
Type locality: "Costa Rica" [no star].
Type specimen: ZMB 36496, holotype, previously listed as not known (Peters, pers. comm. *in* Ernst 1978, 1980b), identified by Fritz et al. (1994).
Etymology:* Toponym for Costa Rica.

Rhinoclemmys melanosterna (Gray 1861b) (14:31)
Colombian Wood Turtle, *Palmera*




Uwe Fritz / El Silencio, Antioquia, Colombia





John B. Iverson / Turbo, Antioquia, Colombia / juvenile



** (orange dots = probable trade or translocated; stars = type localities)


Distribution: Colombia (*Antioquia, Atlántico, Bolívar, Boyacá, Caldas, Cauca, Cesar, Chocó, Córdoba, Cundinamarca, La Guajira, Magdalena, Nariño, Santander, Sucre, Valle del Cauca*), Ecuador, Panama

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 187,077 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 540,304 sq. km
Size (Max SCL): male 27.3 cm, female 30.4 cm (V.P. Páez and B. Bock, unpubl. data)
IUCN Red List:* Near Threatened (NT A2c+4cd) (Páez, Carr, Bock, Garcés-Restrepo, Galvis-Rizo, Giraldo, and Echeverri-García, in press). Previously: TFTSG Provisional Red List: Least Concern (LC) (2011); Least Concern (LC) [Not Listed] (Behler 1996)
CITES:* Appendix II, as *Rhinoclemmys* spp. (2023)
Synonymy:
Geoclemmys melanosterna Gray 1861b:205
Clemmys melanosterna, Rhinoclemmys melanosterna, Rhinoclemys melanosterna, Nicoria punctularia melanosternum, Geoemyda punctularia melanosternum, Geoemyda punctularia melanosterna, Callopsis punctularia melanosterna, Rhinoclemmys punctularia melanosterna
Type locality: "Gulf of Darien: Cherunha" [Colombia]. Restricted to "River Buonaventura [orange star]..[and]..Chirambira, Gulf of Darien [yellow star]" [Colombia] by Boulenger (1889:124); to "Chirambira bei Buenaventura, Columbien" [Colombia] [yellow star] by Mertens and Wermuth (1955: 352), and to "Punta Chirambira en el Delta del Río San Juan...al norte de Buenaventura en la Costa del Pacífico, Chocó" [Colombia] [yellow star] by Medem (1958:21).
Type specimens: NHMUK 1947.3.4.8, 1947.3.5.51 (formerly 1861.11.22.3), syntypes (2).
Etymology:* Greek, *melas*, *melano*, black, and *sternon*, chest; referring to the black plastron in most specimens.

Rhinoclemmys nasuta (Boulenger 1902a)
Large-nosed Wood Turtle, *Sabaletera*


José Vicente Rueda-Almonacid / CBFTT / Isla Palma, Valle del Cauca, Colombia


left: José Vicente Rueda-Almonacid / CBFTT / Isla Palma, Valle del Cauca, Colombia
right: Alan Giraldo / CBFTT / Isla Palma, Valle del Cauca, Colombia


**
(star = type locality)

Distribution:* Colombia (*Antioquia*, *Cauca*, *Chocó*, *Nariño*, *Valle del Cauca*), Ecuador
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 83,244 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 167,866 sq. km
Size (Max SCL): male 19.6 cm, female 22.8 cm (Carr and Giraldo 2009 CBFTT)
CBFTT Account: Carr and Giraldo (2009)
IUCN Red List:* Vulnerable (VU A2bcd+4bcd) (Carr, Garcés-Restrepo, and Giraldo, in press). Previously: TFTSG Provisional Red List: Near Threatened (NT) (2010); Near Threatened (NT) (TFTSG 1996)
CITES:* Appendix II, as *Rhinoclemmys* spp. (2023)
Synonymy:
Nicoria nasuta Boulenger 1902a:53
Geoemyda nasuta, Geoemyda punctularia nasuta, Callopsis punctularia nasuta, Rhinoclemmys punctularia nasuta, Callopsis nasuta, Rhinoclemmys nasuta
Type locality:* "North-western Ecuador...Bulún, 160 feet, and from the Rio Durango, 350 feet." Emended to "San Francisco de Pulún..[&].. Río Bogota...1º3'N, 78º38'W" [Ecuador] [yellow star] by Carr and Almendariz (1990:92).
Type specimens: NHMUK 1947.3.5.54–57 (formerly 1902.5.27.10–12, 1902.5.27.18), syntypes (4).
Etymology:* Latin, *nasuta*, having a big nose; referring to the prominent snout.

Rhinoclemmys pulcherrima (Gray 1856b)
Painted Wood Turtle, *Sabanera, Casco Rojo*
(includes 4 subspecies)



(subspecies: *pulcherrima* = red, *incisa* = blue, *manni* = green, *rogerbarbouri* = pink; orange dots = probably trade or introduced or translocated)

Distribution: Costa Rica, El Salvador, Guatemala, Honduras, Mexico (*Chiapas, Colima, Guerrero, Jalisco, Nayarit, Oaxaca, Sinaloa, Sonora*), Nicaragua

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 259,172 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 1,553,099 sq. km

Size (Max SCL): male 18.1 cm, female 20.7 cm (see subspp.)

IUCN Red List:* Near Threatened (NT A2cd+4cd) (Butterfield, Iverson, and Cupul-Magaña, in press). Previously: Least Concern (LC) [Not Listed] (Behler 1996)

CITES:* Appendix II, as *Rhinoclemmys* spp. (2023)

Rhinoclemmys pulcherrima pulcherrima (Gray 1856b)
Guerrero Wood Turtle



John L. Carr / Guerrero, Mexico / captivity





John L. Carr / Guerrero, Mexico / captivity



(subspecies: *pulcherrima* = red, *incisa* = blue, *rogerbarbouri* = pink; orange dots = probably trade or translocated; stars = type localities)

Distribution: Mexico (*Guerrero, Oaxaca*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 31,699 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 51,837 sq. km

Size (Max SCL): male 16.0 cm, female 16.6 cm (Ernst 1978)

CITES:* Appendix II, as *Rhinoclemmys* spp. (2023)

Synonymy:

Emys pulcherrimus Gray 1856b:25

Clemmys pulcherrima, Callichelys pulcherrima, Emys pulcherrima, Rhinoclemmys pulcherrima, Chelopus pulcherrima, Pseudemys pulcherrima, Chelopus pulcherrimus, Nicoria punctularia pulcherrima, Geoemyda punctularia pulcherrima, Geoemyda pulcherrima pulcherrima, Rhinoclemmys pulcherrima pulcherrima, Callopsis pulcherrima pulcherrima

Type locality: "Mexico." Restricted to "Presidio de Mazatlán, Sinaloa, Mexico" [in error; orange star] by Smith and Taylor (1950a:343, 1950b:30); and to "vicinity of San Marcos, Guerrero, Mexico" [yellow star] by Ernst (1978:125).

Type specimen: NHMUK 1947.3.5.52 (formerly 1851.8.25.24), holotype.

Etymology:* Latin, *pulcherrimus*, most beautiful; referring to the ornate coloration of the shell.

Rhinoclemmys pulcherrima incisa (Bocourt 1868)
Incised Wood Turtle



John B. Iverson / Honduras



John B. Iverson / Honduras



(subspecies: *pulcherrima* = red, *incisa* = blue, *manni* = green; orange dots = probably trade or translocated; star = type locality)

Distribution: El Salvador, Guatemala, Honduras, Nicaragua, Mexico (*Chiapas, Oaxaca*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 130,183 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 279,008 sq. km

Size (Max SCL): male 18.1 cm, female 20.6 cm (Ernst 1978; Legler and Vogt 2013)

CITES:* Appendix II, as *Rhinoclemmys* spp. (2023)

Synonymy:

Emys incisa Bocourt 1868:121

Chelopus incisa, Chelopus incisus, Glyptemys incisa, Nicoria punctularia incisa, Clemmys incisa, Chrysemys incisa, Geoemyda punctularia incisa, Rhinoclemmys incisa, Geoemyda pulcherrima incisa, Rhinoclemmys pulcherrima incisa, Callopsis pulcherrima incisa

Type locality:* "l'Union, un des ports du Salvador" [El Salvador] [yellow star]. Emended to "la Union, un des ports du Salvador sur le Pacifique; elle a été trouvée sur la montagne de Conchavoua" [Conchagua, El Salvador] by Duméril and Bocourt (1870:12).

Type specimen:* MNHN 9131, holotype, specimen figured in Duméril and Bocourt (1870:pl.1,f.1–3,pl.2,f.1–2); see also Ernst (1978) and Bour (2007b); MNHN 6217, listed as "type" by Stuart (1963), is not a type.

Etymology:* Latin, *incisus*, to cut into; evidently referring to the serrated shell margin, based on the original description.

Rhinoclemmys frontalis Gray 1873a:144

Type locality: "Tropical America."

Type specimen: NHMUK 1947.3.5.53 (formerly 1865.3.31.4), holotype.

Etymology:* Latin, *frontis*, forehead, front, *frontalis*, belonging to the forehead; referring to the noticeable striped pattern on the head.

Rhinoclemmys bocourti Gray 1873f:111

Type locality: "Central America."


Type specimen: Possibly MNHN, holotype, apparently lost, specimen figured in Duméril and Bocourt (1870:pl.7,f.1,1a–b).

Etymology:* Eponym honoring Marie Firmin Bocourt (1819–1904), France; naturalist and artist who collected reptiles in Central America.


Rhinoclemmys pulcherrima manni (Dunn 1930)
Central American Wood Turtle



John B. Iverson / nr. Canas, Guanacaste Prov., Costa Rica



John B. Iverson / nr. Canas, Guanacaste Prov., Costa Rica



(subspecies: *incisa* = blue, *manni* = green; orange dots = probably trade or translocated; star = type locality)

Distribution: Costa Rica, Nicaragua

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 25,134 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 49,910 sq. km

Size (Max SCL): male 16.6 cm, female 20.7 cm (Merchan Fornelino 2003)

CITES:* Appendix II, as *Rhinoclemmys* spp. (2023)

Synonymy:

Geoemyda manni Dunn 1930:33

Geoemyda pulcherrima manni, Geoemyda punctularia manni, Callopsis pulcherrima manni, Rhinoclemmys pulcherrima manni


Type locality: "San Jose, Costa Rica" [yellow star].

Type specimen: MCZ 29097, holotype.
Etymology:* Eponym honoring William M. Mann (1886–1960), USA; entomologist and director of the U.S. National Zoological Park, Washington, D.C.


Rhinoclemmys pulcherrima rogerbarbouri (Ernst 1978)
Western Mexican Wood Turtle



John B. Iverson / nr. Colima, Colima, Mexico



left: John B. Iverson / nr. Colima, Colima, Mexico
right: Anders G.J. Rhodin / Monte Mojino Reserve, Alamos, Sonora, Mexico



** (subspecies: *pulcherrima* = red, *rogerbarbouri* = pink; orange dots = probably trade or translocated; star = type locality)

Distribution:* Mexico (*Chihuahua, Colima, Jalisco, Nayarit, Sinaloa, Sonora*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 73,256 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 153,172 sq. km
Size (Max SCL): male 17.6 cm, female 20.2 cm (Ernst 1978)
CITES:* Appendix II, as *Rhinoclemmys* spp. (2023)
Synonymy:
Callopsis pulcherrima rogerbarbouri Ernst 1978:127
Rhinoclemmys pulcherrima rogerbarbouri
Type locality:* "Guirocoba, Sonora, Mexico" [= Huirocoba, 28 km SE of Alamos, Sonora] [yellow star].
Type specimen: AMNH 63760, holotype.
Etymology:* Eponym honoring Roger W. Barbour (1919–1993), USA; vertebrate zoologist and herpetologist at the University of Kentucky.

Rhinoclemmys punctularia (Daudin 1801) [08:12, 09:28]
Spot-legged Turtle
(includes 2 subspecies)



** (subspecies: *punctularia* = red, *flammigera* = blue; orange dots = probably introduced)

Distribution: Brazil (*Amapá, Amazonas, Bahia, Espirito Santo, Maranhão, Pará, Piauí, Roraima, Tocantins*), French Guiana, Guyana, Suriname, Trinidad and Tobago, Venezuela (*Amazonas, Bolívar, Delta Amacuro, Monagas*)
Introduced: Brazil (*Rio de Janeiro*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 2,213,334 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 5,992,970 sq. km
Size (Max SCL): male 20.1 cm, female 24.8 cm (see subspp.)
IUCN Red List:* Near Threatened (NT A2cd+4cd) (Brito, Valadão, Cunha, Oliveira, Hernández, Ferrara, Andrade, and Carr 2025). Previously: TFTSG Provisional Red List: Least Concern (LC) (2011); Least Concern (LC) [Not Listed] (Behler 1996)
CITES:* Appendix II, as *Rhinoclemmys* spp. (2023)

Rhinoclemmys punctularia punctularia (Daudin 1801) (08:12, 09:28, 21:74)
Eastern Spot-legged Turtle, *Perema, Aperema*



Jérôme Maran / French Guiana



John L. Carr / Brazil



(subspecies: *punctularia* = red, *flammigera* = blue;
orange dots = probably introduced; stars = type localities)

Distribution: Brazil (*Amapá, Amazonas, Espirito Santo, Maranhão, Pará, Piauí, Roraima, Tocantins*), French Guiana, Guyana, Suriname, Trinidad and Tobago, Venezuela (*Bolívar, Delta Amacuro, Monagas*)

Introduced: Brazil (*Rio de Janeiro*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 2,188,265 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 5,735,675 sq. km

Size (Max SCL): male 20.1 cm, female 24.8 cm (Ernst 1978; Pritchard and Trebbau 1984; Wariss et al. 2012)

CITES:* Appendix II, as *Rhinoclemmys* spp. (2023)

Synonymy:

Testudo amboinensis Linnaeus 1754:50 (invalid pre-1758 name)
Type locality: "India."
Type specimen: NRM 31491 (formerly MAF 89), type; erroneously included in composite syntype series of *Testudo geometrica* Linnaeus 1758; NRM 1 (formerly MDG s/n), not the type, erroneously interpreted as the type by Andersson (1900), identified as = *Hydraspis gibba* (= *Mesoclemmys gibba*) by Andersson (1900), see also Wallin (1977) and Bour (2005d).
Etymology:* Toponym for Ambon, Molukkas, Indonesia [in error].
Comment: Unavailable pre-1758 binomial name. Description cited as also sourced from Seba (1734:126,pl.79,f.1–2) (= *Melanochelys trijuga*).

Testudo scabra Linnaeus 1758:198 (senior homonym and *nomen oblitum*, not = *Testudo scabra* Retzius *in* Schoepff 1792 [= *Pelomedusa galeata*], nor = *Testudo scabra* Latreille *in* Sonnini and Latreille 1801 [= *Lissemys punctata punctata*]) (08:12) (74)*
Emys scabra, Rhinoclemmys scabra , Rhinoclemys scabra
Type locality: "Indiis" [West Indies]. Restricted to "India orientali, Carolina" [in error] by Linnaeus (1766:351); and to "Cayenne, French Guiana" [yellow star] by Rhodin and Carr (2009:11).
Type specimen: UPSZTY 129 (formerly UUZM 129), holotype, identified and described by Rhodin and Carr (2009).
Etymology:* Latin, *scabra*, rough, shabby, mangy, unclean; referring to the rough carapacial surface.

Testudo verrucosa Walbaum 1782:116 (*nomen novum* and unavailable name)
Etymology:* Latin, *verruca*, wart, *verrucosa*, full of warts, warty; referring to the rough carapacial surface.
Comment: Unjustified replacement name for *Testudo scabra* Linnaeus, see Rhodin and Carr (2009). Unavailable name from a non-binomial work, see Wermuth (1956).

Testudo verrucosa Suckow 1798:40 (junior homonym, not = *Testudo (ferox) verrucosa* Schoepff 1795 [= *Apalone ferox*])
Chemelys verrucosa
Type locality: Not known.
Type specimen: Not known or located.
Etymology:* Latin, *verruca*, wart, *verrucosa*, full of warts, warty; referring to the rough carapacial surface.

Testudo punctularia Daudin 1801:249("349") (*nomen protectum*)
Emys punctularia, Chersine punctularia, Clemmys (Clemmys) punctularia, Clemmys punctularia, Chelopus punctularius, Nicoria punctularia, Geoemyda punctularia, Geoemyda punctularia punctularia, Rhinoclemmys punctularia, Rhinoclemmys punctularia punctularia, Callopsis punctularia punctularia
Type locality: "Amérique méridionale...sur-tout...la Guiane..[&].. Cayenne" [French Guiana] [yellow star].
Type specimen: MNHN 9130, holotype, see Fretey et al. (1977) and Bour (2007b).
Etymology:* Latin, *punctulatus*, spotted; referring to the head spots.
Comment: Name conserved by ICZN (1963), see Wermuth (1956) and Mertens and Wermuth (1961).

Testudo dorsata Schoepff 1801:136 (*nomen novum et suppressum*)
Emys dorsata, Clemmys dorsata
Type locality: Not known.
Type specimen: Possibly ZMUS, not located, type specimen apparently figured on unpublished plate cited by Schoepff (1801:pl.34).
Etymology:* Latin, *dorsum*, the back, referring to the carapacial keel.
Comment: Unjustified replacement name for *Testudo scabra* Linnaeus. Description cited as sourced from Hermann (pers. comm.), whose own description of *Testudo dorsata* was published posthumously (Hermann 1804:219). Name suppressed by ICZN (1963), see Wermuth (1956) and Mertens and Wermuth (1961).

Testudo alacacca Stedman *in* Schweigger 1812:429 (*nomen nudum)*
Type locality: "Surinam" [green star, location estimated].
Type specimen: Not known or located.
Etymology:* Vernacular (Native Surinamese), *alacacca*, local indigenous name for the species.

Emys dorsualis Spix 1824:11
Emys dorsalis
Type locality: "juxta flumen Solimoens" [Amazonas, Brazil] [= along Rio Solimões between Tefé and Tabatinga, possibly near Tefé] [blue star, location estimated].
Type specimen: ZSM 2424/0, lectotype, designated by Hoogmoed and Gruber (1983:350), see Franzen and Glaw (2007).

Etymology:* Latin, *dorsum*, the back, referring to the carapacial keel.

Rhinoclemys bellii Gray 1863c:183

Rhinoclemmys bellii, Rhinoclemmys scabra bellii

Type locality: "Tropical America."

Type specimen: OUM 8510, holotype, see Nowak-Kemp and Fritz (2010).

Etymology:* Eponym honoring Thomas Bell (1792–1880), United Kingdom; herpetologist, dental surgeon, and Professor of Zoology at King's College, London.

Geoclemys callocephalus Gray 1863h:254

Clemmys callocephala, Geoclemmys callocephala, Geoclemmys callocephalus, Rhinoclemmys callocephala, Rhinoclemys callocephala

Type locality: Not known. Restricted to "S. America" by Boulenger (1889:124).

Type specimen: NHMUK 1947.3.5.50 (formerly 1853.11.30.3), holotype.

Etymology:* Greek, *kallos*, beauty, and *kefale*, head; referring to the attractive head markings.

Rhinoclemmys lunata Gray 1873a:144

Geoemyda punctularia lunata, Callopsis punctularia lunata, Rhinoclemys lunata, Rhinoclemmys punctularia lunata

Type locality: Not designated. Restricted to "Dutch Guiana" [Suriname] [green star, location estimated] by Boulenger (1889:124).

Type specimens: NHMUK 1946.1.22.69–70 (formerly 1866.8.14.227), syntypes (2).

Etymology:* Latin, *luna*, moon, *lunata*, crescent-shaped, curved; referring to the crescent-shaped marking on the head.

Rhinoclemmys ventricosa Gray 1873a:145

Rhinoclemys ventricosa

Type locality: "Tropical America." Restricted to "Dutch Guiana" [Suriname] [green star, location estimated] by Boulenger (1889:124).

Type specimen: NHMUK 1947.3.5.49 (formerly 1867.4.2.156), holotype.


Etymology:* Latin, *ventricosa*, potbellied; referring to the bulging, deep shell.

Rhinoclemmys punctularia flammigera Paolillo O. 1985 (08:25, 09:28)

Upper Orinoco Spot-legged Turtle



Fernando J.M. Rojas-Runjaic / nr. Orinoco-Ventuari confluence, Venezuela



** (subspecies: *punctularia* = red, *flammigera* = blue; star = type locality)

Distribution: Venezuela (*Amazonas*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 25,125 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 40,085 sq. km

Size (Max SCL): male 19.3 cm, female 22.5 cm (Pritchard and Trebbau 1984; Paolillo O. 1985)

CITES:* Appendix II, as *Rhinoclemmys* spp. (2023)

Synonymy:

Rhinoclemmys punctularia flammigera Paolillo O. 1985:294


Rhinoclemmys flammigera

Type locality: "Caño Maica, 10 km SE of Carmelitas, Territorio Federal Amazonas, Venezuela (4º4' N, 66º31' W)" [yellow star].

Type specimen: EBRG 1467, holotype.

Etymology:* Latin, *flamma*, flame, fire, and *gerere*, wear, carry, *flammigera*, to wear flames; referring to the flame-like shaped radially arranged markings on the head.

Rhinoclemmys rubida (Cope 1870a)
Mexican Spotted Wood Turtle
(includes 2 subspecies)



(subspecies: *rubida* = red, *perixantha* = blue;
orange dots = probably translocated)

Distribution: Mexico (*Chiapas, Colima, Guerrero, Jalisco, México, Michoacán, Oaxaca*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 74,137 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 284,927 sq. km

Size (Max SCL):* male 23.0 cm, female 17.9 cm (see subspp.)

IUCN Red List:* Vulnerable (VU A2c+4c) (Holcomb and Butterfield, in press). Previously: Near Threatened (NT) (van Dijk, Canseco-Marquez, and Muñoz 2007); Vulnerable (VU) (TFTSG *in* Baillie and Groombridge 1996)

CITES:* Appendix II, as *Rhinoclemmys* spp. (2023)


Rhinoclemmys rubida rubida (Cope 1870a)
Oaxaca Wood Turtle



Michael Redmer / Mexico / captivity



Eduardo Reyes-Grajales / Arriaga, Chiapas, Mexico



(subspecies: *rubida* = red; orange dots = probably translocated;
stars = type localities)

Distribution: Mexico (*Chiapas, Oaxaca*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 32,083 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 86,202 sq. km

Size (Max SCL):* male 23.0 cm, female 17.9 cm (Ernst 1978; Legler and Vogt 2013)

CITES:* Appendix II, as *Rhinoclemmys* spp. (2023)

Synonymy:


Chelopus rubidus Cope 1870a:148
Geoclemmys rubida, Emys rubida, Nicoria rubida, Clemmys rubida, Geoemyda rubida, Geoemyda rubida rubida, Rhinoclemmys rubida, Rhinoclemmys rubida rubida, Callopsis rubida, Callopsis rubida rubida, Chelopus rubidus rubidus
Type locality: "Tuchitan Tehuantepec, Mexico" [Juchitán, Oaxaca] [yellow star].
Type specimens: ANSP 285, 337–341, USNM 45612–14, syntypes (9), see Cochran (1961), Malnate (1971), and Reynolds et al. (2007).
Etymology:* Latin, *rubidus*, reddish; referring to the light red skin markings.

Rhinoclemmys mexicana Gray 1870b:659
Chelopus mexicana, Chelopus mexicanus, Emys mexicana
Type locality: "Mexico; San Juan del Rio" [Oaxaca] [green star].
Type specimens: NHMUK 1947.3.4.34 and 1947.3.5.60–61 (formerly 1870.6.20.1, 1871.2.7.45–46), syntypes (3).
Etymology:* Toponym for Mexico.


Rhinoclemmys rubida perixantha (Mosimann and Rabb 1953)
Colima Wood Turtle



John B. Iverson / nr. Colima, Colima, Mexico



John B. Iverson / Colima, Mexico




(subspecies: *perixantha* = blue; orange dot = probably translocated;
star = type locality)

Distribution: Mexico (*Colima, Guerrero, Jalisco, México, Michoacán*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 42,054 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 103,220 sq. km
Size (Max SCL): male 14.1 cm, female 15.6 cm (Ernst 1978)
CITES:* Appendix II, as *Rhinoclemmys* spp. (2023)
Synonymy:
Geoemyda rubida perixantha Mosimann and Rabb 1953:1
Rhinoclemmys rubida perixantha, Callopsis rubida perixantha, Chelopus rubidus perixanthus
Type locality: "8 kilometers south of Tecoman, Colima, Mexico" [yellow star].
Type specimen: UMMZ 80336, holotype, see Kluge (1984).
Etymology:* Greek, *peri-*, around, and *xanthos*, yellow; referring to the yellowish marginals.

Family Testudinidae Batsch 1788 (21:3)

Tortoises
(includes 3 subfamilies, 18 genera, 53 species, 91 taxa)
Testudines Batsch 1788:437
Testudia Rafinesque 1814:66
Tortudines Schmid 1819:10
Testudinidae Gray 1825:210
Tylopodae Wagler 1828:861
Dysmydae Ritgen 1828:270
Tylopodes Burmeister 1837:732
Baenodactyli Mayer 1849:198
Testudinides Pictet 1853:442
Testudinina Agassiz 1857a:356
Testudinidi Portis 1890:12



Testudinidae Species Richness

Subfamily Manouriinae Gray 1869a (21:3)

Manourine Tortoises
(includes 1 genus, 2 species, 3 taxa)
Manourina Gray 1869a:167
Manouriana Gray 1873j:17
Manouriina Baur 1888b:597
Manouriinae Rhodin, Iverson, Bour, Fritz, Georges, Shaffer, and van Dijk [Turtle Taxonomy Working Group] 2021:251

Manouria Gray 1854a

Indochinese Tortoises
(includes 2 species, 3 taxa)
Manouria Gray 1854a:133
Type species: *Manouria fusca* Gray 1854a [= subjective synonym of *Testudo emys* Schlegel and Müller 1840], by original monotypy.
Etymology:* Anonym, *manouria*, possibly a fabricated name, or possibly an Eponym honoring someone named Manour; or possibly Greek, *manos*, thin, and *ouria*, tail, but unlikely, since the specimens described were shells without soft parts.
Teleopus Le Conte 1854:187
Type species: *Teleopus luxatus* Le Conte 1854 [= subjective synonym of *Testudo emys* Schlegel and Müller 1840], by monotypy.
Etymology:* Greek, *telos*, end, and *pus*, foot; referring to the front and back feet having five claws each.
Scapia Gray 1869a:167
Type species: *Testudo (Scapia) falconeri* Gray 1869 [= subjective synonym of *Testudo emys* Schlegel and Müller 1840 or *Testudo phayrei* Blyth 1854], by monotypy.
Etymology:* Anonym, *scapia*, possibly a fabricated name, or possibly Latin, *scapus*, roller, trunk, shaft; possibly referring to the massive, cylindrical body shape, or possibly an Eponym honoring someone named Scap or Skap, or starting with Scap- or Skap-.

Manouria emys (Schlegel and Müller 1840) (21:75)

Asian Giant Tortoise
(includes 2 subspecies)



** (subspecies: *emys* = red, *phayrei* = blue; orange dots = probable trade)
Distribution: Bangladesh, Brunei, India (*Assam, Meghalaya, Mizoram, Nagaland*), Indonesia (*Kalimantan, Sumatra*), Malaysia (*East, West*), Myanmar, Singapore (*extirpated*), Thailand
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 571,019 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 5,393,869 sq. km
Size (Max SCL): male 60.0 cm, female 58.0 cm (see subspp.)
CBFTT Account: Stanford, Wanchai, Schaffer, Schaffer, and Thirakhupt (2015)
IUCN Red List: Critically Endangered (CR A2cd+4cd) (Choudhury, Cota, McCormack, Platt, Das, Ahmed, Timmins, Rahman, and Singh 2019). Previously: Endangered (EN) (ATTWG 2000); Vulnerable (VU) (TFTSG *in* Baillie and Groombridge 1996)
CITES: Appendix II, as Testudinidae spp. (1977). Previously: Appendix II (1975)

Manouria emys emys (Schlegel and Müller 1840) [21:75]
Asian Brown Giant Tortoise




Chuck Schaffer / CBFTT / Tabin Wildlife Reserve, Sabah, Malaysia (East)





left: Chuck Schaffer / CBFTT / No data / captivity
right: Mark Auliya / CBFTT / West Malaysia



** (subspecies: *emys* = red, *phayrei* = blue; orange dots = probable trade; stars = type localities)

Distribution: Brunei, Indonesia (*Kalimantan, Sumatra*), Malaysia (*East, West*), Singapore (*extirpated*), Thailand

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 416,022 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 2,439,884 sq. km

Size (Max SCL): male 56.0 cm (de Rooij 1915; Lim and Das 1999)

CITES: Appendix II, as Testudinidae spp. (1977). Previously: Appendix II (1975)

Synonymy:

Testudo emys Schlegel and Müller 1840:pl.4
Manouria emys, Manouria emys emys, Geochelone emys, Geochelone emys emys, Testudo emys emys
Type locality: "Sumatra" [Indonesia]. Restricted to "versant sud du Gunung (= Mont) Singgalang, dans la grande vallée de la rivière Anai, et à l'est de Padang, Sumatera Barat, Sumatra, Indonésie" [Indonesia] [yellow star] by Bour (1998:36).
Type specimen: RMNH 3808, lectotype, designated by Hoogmoed and Crumly (1984:251), discussed by Hoogmoed et al. (2010).
Etymology:* Greek, *emys*, turtle; referring to its apparent similarity to freshwater turtles.

Testudo emydoides Duméril and Bibron *in* Duméril and Duméril 1851:4 (*nomen novum*)
Manouria emydoides
Etymology:* Greek, *emydos*, turtle, and *-oides*, resembling; referring to its apparent similarity to freshwater turtles.
Comment: Unjustified replacement name for *emys*.

Manouria fusca Gray 1854a:134
Type locality: "Singapore" [probable trade, orange star]. Restricted to "Pinang" [Penang, Malaysia] [green star, location estimated] by Gray (1856:16).
Type specimen: NHMUK 1947.3.4.57, holotype.
Etymology:* Latin, *fusca*, dark brown, dark, gray, black; referring to the carapace color.

Teleopus luxatus Le Conte 1854:187
Manouria luxata
Type locality: "Java" [Indonesia] [probable trade, brown star, location estimated].
Type specimen: ANSP, holotype, see Baur (1892), apparently lost, not listed by Malnate (1971).
Etymology:* Latin, *luxatus*, dislocated; referring to the triangular pectoral shields that are separated at the plastral midline.

Manouria emys phayrei (Blyth 1854) [17:66, 21:75]
Burmese Black Giant Tortoise




Craig B. Stanford / CBFTT / Kaeng Krachan National Park, Thailand





left: Chuck Schaffer / CBFTT / No data / captivity
right: Craig B. Stanford / CBFTT / Kaeng Krachan National Park, Thailand



(subspecies: *emys* = red, *phayrei* = blue; orange dots = probable trade; stars = type localities)

Distribution: Bangladesh, India (*Assam, Meghalaya, Mizoram, Nagaland*), Myanmar, Thailand

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO): 154,997 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 767,031 sq. km

Size (Max SCL): male 60.0 cm, female 58.0 cm (Nutaphand 1979; Høybye-Mortensen 2004; Stanford et al. 2015 CBFTT)

CITES: Appendix II, as Testudinidae spp. (1977). Previously: Appendix II (1975)

Synonymy:

Testudo phayrei Blyth 1854:639 (17:66)
Scapia phayrei, Manouria emys phayrei
Type locality: "Arakan; Tenasserim Provinces...Burma" [Myanmar]. Restricted to "Arakan" [nr. Sittwe, Myanmar] [yellow star, location estimated] by Anderson (1872:132).
Type specimen: ZSI 813, lectotype, designated "type" by Anderson (1871:426,f.1–2), discussed by Bour (1998), see Das et al. (1998), Das (2009), and Kundu et al. (2018a).
Etymology:* Eponym honoring Sir Arthur Purves Phayre (1812–1885), United Kingdom / Myanmar; army officer and eventual Commissioner in Burma (now Myanmar) who collected the type while a Captain serving in Arakan.


Testudo (Scapia) falconeri Gray 1869a:169
Testudo falconeri, Scapia falconeri
Type locality: "India?" [no star].
Type specimen: Formerly NHMUK s/n, holotype, figured (f.1); returned to donor (Gray 1873j:18).
Etymology:* Eponym honoring Hugh Falconer (1808–1865), United Kingdom (Scotland) / India; physician, paleontologist, and geologist who worked in India and described several fossil animals, including a giant tortoise.

Testudo nutapundi Reimann and Nutaphand *in* Nutaphand 1979:193
Geochelone nutapundi, Manouria emys nutapundi, Geochelone emys nutapundi, Manouria nutapundi
Type locality: "Northern Thailand (Tak Province) and western Central Region (Kanchanaburi Province); Assam, Burma". Restricted to "Nord-Thailand, Tak-Provinz" [green star, location estimated] by Obst (1983:253).
Type specimen: Not located, specimen figured (f.105-106), but not considered the holotype by Obst (1983).
Etymology:* Eponym honoring Wirot Nutaphand (or Nutphand) (1932–2005), Thailand; airforce officer, naturalist, and herpetologist.

Manouria impressa (Günther 1882)
Impressed Tortoise


David Emmett / TL / Cardamom Mts., Cambodia


Peter Paul van Dijk / Loei Prov., Thailand


(orange dots = probable trade; stars = type localities)

Distribution: Cambodia, China (*Yunnan*), India (*Arunachal Pradesh, Assam [?]*), Laos, Malaysia (*West*), Myanmar, Thailand, Vietnam

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 471,131 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 2,641,658 sq. km

Size (Max SCL): male 29.3 cm, female 35.9 cm (Chan-ard et al. 1996; Mital et al. 2020)

IUCN Red List: Endangered (EN A2cd) (Cota, Li, Platt, Rao, Stanford, Wang, and Wanchai 2021). Previously: Vulnerable (VU) (ATTWG 2000); Vulnerable (VU) (TFTSG *in* Baillie and Groombridge 1996)

CITES: Appendix II, as Testudinidae spp. (1977). Previously: Appendix II (1975)

Synonymy:

Geoemyda impressa Günther 1882:343
Testudo impressa, Geochelone impressa, Manouria impressa
Type locality: "Siam" [Thailand] [orange star, location estimated, probable trade specimen].
Type specimen: NHMUK 1947.3.5.7 (formerly 1882.4.20.1), holotype, see Bour (1998).
Etymology:* Latin, *impressa*, pressed in, impressed; referring to slightly concave indented pleural scutes.

Geoemyda latinuchalis Vaillant 1894:68
Testudo latinuchalis
Type locality: "la rivière Noire, Tonkin" [Black River (Song Da), Vietnam] [green star, location estimated].
Type specimen: MNHN 1892.266, lectotype, designated by Bour (1998:30).
Etymology:* Latin, *latus*, broad, wide, and *nuchalis*, belonging to the neck; referring to the nuchal scute being significantly wider than long.

Testudo pseudemys Boulenger 1903a:144
Type locality: "Batang Padang district, South Perak (1,000 feet to 2,000 feet)" [Malaysia] [blue star].
Type specimen: NHMUK 1947.3.5.6 (formerly 1903.4.13.6), lectotype, designated by Bour (1998:31).
Etymology:* Greek, *pseudo*, false, untrue, and *emys*, turtle; referring to the fact that this taxon, while similar to *Manouria emys*, is not the same.

Subfamily Xerobatinae Gray 1874 (21:3)

North American Tortoises

(includes 1 genus, 6 species, 6 taxa)

Xerobatina Gray 1874:723

Xerobatinae Gaffney and Meylan 1988:210

Gopherus Rafinesque 1832 (17:67, 21:76)

Gopherine Tortoises

(includes 6 species, 6 taxa)

Gopherus Rafinesque 1815:74 (*nomen nudum*)

Etymology:* Vernacular (English), *gopher*, anglicized from Vernacular (Old French), *gaufre*, local name for the species, first used by Bartram (1791).

Gopherus Rafinesque 1832:64

Type species: *Gopherus polyphemus* [= *Testudo polyphemus* Daudin 1801], by original designation.

Etymology:* Vernacular (English), *gopher*, anglicized from Vernacular (Old French), *gaufre*, local name for the species, first used by Bartram (1791).

Xerobates Agassiz 1857a:252,446

Type species: *Xerobates berlandieri* Agassiz 1857a, by subsequent designation by Brown (1908:115).

Etymology:* Greek, *xeros*, dry, arid, and *baino*, walk, run, to wander; referring to the dry habitats in which the species lives and walks.

Bysmachelys Johnston 1937:439

Type species: *Bysmachelys canyonensis* † Johnston 1937 [= subjective synonym of *Testudo pertenuis* † Cope 1892b (see TEWG 2015)], by original monotypy.

Etymology:* Greek, *bysma*, plug, dowel, and *chelys*, turtle; referring to the short, broad head.

Scaptochelys Bramble 1982:852

Type species: *Scaptochelys agassizii* [= *Xerobates agassizii* Cooper 1861], by original designation.


Etymology:* Greek, *skaptos*, digger, and *chelys*, turtle; referring to the burrowing behavior of the included species.

Gopherus agassizii (Cooper 1861) (10:22, 11:11)


Mojave Desert Tortoise, Agassiz's Desert Tortoise



Bev Steveson / CCB / CBFTT / Mojave Desert, California



U.S. Geological Survey c/o Kristin H. Berry / CBFTT / Mojave Desert, California



** (orange dots = probably introduced or translocated; star = type locality)

Distribution: USA (*Arizona*, *California*, *Nevada*, *Utah*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 120,488 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 185,471 sq. km

Size (Max SCL):* male 33.0 cm, female 37.4 cm (Berry and Murphy 2019 CBFTT); gigantic captive: 49.3 cm (Jackson et al. 1980)

CBFTT Account: Berry and Murphy (2019)

IUCN Red List: Critically Endangered (CR A2abce+4abce) (Berry, Allison, McLuckie, Vaughn, and Murphy 2021). Previously: Vulnerable (VU) (TFTSG 1996)

CITES: Appendix II, as Testudinidae spp. (1977). Previously: Appendix II, as *Gopherus* spp. (1975)

USA ESA:* Threatened, wherever found, except Arizona south and east of Colorado River, and Mexico; or Threatened (by Similarity of Appearance) in Arizona south and east of Colorado River, and Mexico, when found outside of Mexico or said range in Arizona (USFWS 1990)

Synonymy:

Xerobates agassizii Cooper 1861:120 (10:22)

Testudo agassizii, *Gopherus agassizii*, *Gopherus polyphemus agassizii*, *Geochelone agassizii*, *Scaptochelys*

agassizii

Type locality: "mountains of California, near Fort Mojave" [USA]. Restricted to "California, San Bernardino County; Mountains of California, near Fort Mojave; Soda Valley (very approximately 35°6' N, 116°6' W)" [yellow star] by Murphy et al. (2011:51).

Type specimen: USNM 7888, lectotype, designated and genotyped by Murphy et al. (2011:51), see also Cochran (1961) and Reynolds et al. (2007).

Etymology:* Eponym honoring Louis Agassiz (1807–1873), Switzerland / USA; herpetologist at Harvard University who founded the Museum of Comparative Zoology.

Xerobates lepidocephalus Ottley and Velázques Solis 1989:497 (11:11)

Type locality: "western base of the Sierra San Vicente, approximately 1 km N of the Buena Mujer Dam, Baja California Sur, Mexico" [ca. 24.1 N, -110.2 W] [in error, introduced] [outside of map, not shown].

Type specimen: BYU 39706, holotype; genotyped by Murphy et al. (2011).

Etymology:* Greek, *lepidos*, scaly, and *kefala*, head; referring to the thickened raised scales on the head.


Comment:* Shown by Murphy et al. (2011) to have the same mitochondrial DNA as *G. agassizii* from the Mojave Desert region.

Gopherus berlandieri (Agassiz 1857a) (14:34, 21:77)


Texas Tortoise, Berlandier's Tortoise, *Galapago Tamaulipas*



Larry Ditto / CCB / Texas



left: Larry Ditto / Texas
right: John B. Iverson / northeast Mexico



** (orange dots = probably introduced or translocated; stars = type localities)

Distribution: Mexico (*Coahuila, Nuevo León, San Luis Potosi, Tamaulipas, Veracruz*), USA (*Texas*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 287,918 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 408,465 sq. km

Size (Max SCL): male 23.8 cm, female 21.0 cm (Rose and Judd 2014)

IUCN Red List:* Near Threatened (NT A2c+4c) (Mali and Forstner, in press). Previously: TFTSG Provisional Red List: Near Threatened (NT) (2011); Least Concern (LC) (TFTSG 1996)

CITES: Appendix II, as Testudinidae spp. (1977). Previously: Appendix II, as *Gopherus* spp. (1975)

Synonymy:

Testudo tuberculata Berlandier 1850:287 (junior homonym and *nomen oblitum*, not = *Testudo tuberculata* Pennant *in* Schoepff 1801 [= *Dermochelys coriacea*]) (14:34, 21:77)

Testudo tuberculatu

Type locality: "Tamaulipas" [Mexico]. Restricted to "los llanos de Tamaulipas" [Mexico] [orange star, location estimated, likely nr. Matamoros] by Bour (2017:53).

Type specimen: Not located, specimen figured in manuscript drawing by Berlandier in 1850 (pl.5), reproduced and designated lectotype by Bour (2017:f.3).

Etymology:* Latin, *tuberculum*, small hump, nodule; referring to the humplike skin nodules on the head and neck.

Testudo bicolor Berlandier 1850:287 (*nomen oblitum*) (14:34, 21:77)

Type locality: "Tamaulipas" [Mexico] [orange star, location estimated, likely nr. Matamoros].

Type specimen: Not known or located.

Etymology:* Latin, *bi*, twice, and *color*, color; referring to the predominantly blackish and yellowish coloring of the shell.

Xerobates berlandieri Agassiz 1857a:392,447

Testudo berlandieri, Xerobates gopher berlandieri, Gopherus berlandieri, Gopherus polyphemus berlandieri, Scaptochelys berlandieri

Type locality: "southern Texas [USA] and Mexico." Restricted to "Lower Rio Grande, Texas" [USA] by Stejneger and Barbour (1917:121); and to "Brownsville, Cameron County, Texas, USA" [yellow star] by Smith and Taylor (1950a:361).

Type specimen: USNM 60(2), lectotype, designated by Bour (2017:61,f.8), also discussed by Cochran (1961), Reynolds et al. (2007), and Lieb (2023).

Etymology:* Eponym honoring Jean Louis Berlandier (1803–1851),

Belgium / Mexico; naturalist who worked in Mexico and collected some of the types.

Testudo minima Berlandier *in* Bour 2017:58 (*nomen nudum*)

Etymology:* Latin, *minima*, smallest, very small; referring to the relatively small size of this taxon.

Comment: Original manuscript name for *Testudo bicolor* Berlandier 1850, never published.

Gopherus evgoodei Edwards, Karl, Vaughn, Rosen, Meléndez Torres, and Murphy 2016 (17:68)


Sinaloan Thornscrub Tortoise, Goode's Thornscrub Tortoise, *Tortuga Sinaloense de Matorral, Tortuga de Monte Sinaloense*



Eric V. Goode / Monte Mojino Reserve, Alamos, Sonora, Mexico



Eric V. Goode / Monte Mojino Reserve, Alamos, Sonora, Mexico



** (star = type locality)

Distribution: Mexico (*Chihuahua, Sinaloa, Sonora*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 42,108 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 61,033 sq. km

Size (Max SCL): male 25.5 cm, female 26.1 cm (A. Karl, unpubl. data)

IUCN Red List: Vulnerable (VU A4ce) (Edwards, Karl, Rosen, Vaughn, Melendez Torres, Kiester, and Goode 2018)

CITES: Appendix II, as Testudinidae spp. (1977)

Synonymy:

Gopherus evgoodei Edwards, Karl, Vaughn, Rosen, Meléndez Torres, and Murphy 2016:140

Type locality: "Alamos (approximate location 27°02' N, 108°55' W, elevation 433 m), Sonora, Mexico" [yellow star].

Type specimen: AMNH 64160, holotype.


Etymology:* Eponym honoring Eric V. Goode (1957–), USA; hotelier, herpetologist, and conservationist who founded Turtle Conservancy and supported land purchase to protect this species as it was being described.

Gopherus flavomarginatus Legler 1959 (12:32, 21:76)


Bolson Tortoise, Mexican Giant Tortoise, *La Tortuga del Bolsón, Tortuga Grande, Tortuga Llanero*



Eric V. Goode / TCC / Bolson de Mapimí, Durango, Mexico



Eric V. Goode / Bolson de Mapimí, Durango, Mexico



** (orange dots = introduced, translocated, or possible relictual populations; stars = type localities)

Distribution: Mexico (*Chihuahua, Coahuila, Durango*)

Introduced: USA (*Arizona, New Mexico*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 19,375 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 29,309 sq. km

Size (Max SCL): male 35.6 cm, female 39.1 cm (Legler and Vogt 2013; McCoy et al. 2014; C. Wiese and S. Hillard, Turner Endangered Species Fund, unpubl. data); **(Max CCL):** ca. 46.0 cm [estimated].


IUCN Red List: Critically Endangered (CR A4bcd) (Kiester, Palomo-Ramos, Ríos-Arana, and Goode 2018). Previously: Vulnerable (VU A1cd) (van Dijk and Flores-Villela 2007); Vulnerable (VU) (TFTSG *in* Baillie and Groombridge 1996)

IUCN Green List:* Largely Depleted (LD) (van Dijk and Castañeda Gaytán 2024)
CITES: Appendix I (1979). Previously: Appendix II, as Testudinidae spp. (1977); Appendix II, as *Gopherus* spp. (1975)
USA ESA:* Endangered (USFWS 1979)
Synonymy:
Gopherus flavomarginatus Legler 1959:337
Gopherus polyphemus flavomarginatus, Gopherus flavomarginata
Type locality: "30 to 40 miles from Lerdo, Durango, Mexico" [yellow star, location estimated].
Type specimen: USNM 61253, holotype, see Cochran (1961) and Reynolds et al. (2007).
Etymology:* Latin, *flavus*, golden or yellow, and *marginatus*, border; referring to the color of the marginal scutes.
Gopherus huecoensis † Strain 1966:24 (21:52)
Type locality: "Madden Arroyo, locality B.E.G. 40240, NW1/4NW1/4 Sec. 19, T. 7, Blk. 73, Hudspeth County, Texas" [USA] [gray star].
Type specimen: BEG 40240-27, holotype, fossil, plastron and appendicular bones, figured (pl.2-6), see also Vlachos (2018).
Geological age: Lower Pleistocene, Fort Hancock Formation, Late Blancan, Piacenzian–Gelasian; see Vlachos (2018).
Etymology:* Toponym for the Hueco Bolson, Texas, USA.

Gopherus morafkai Murphy, Berry, Edwards, Leviton, Lathrop, and Riedle 2011 (11:11)
Sonoran Desert Tortoise, Morafka's Desert Tortoise



Peter Paul van Dijk / Pima Co., Arizona



left: Roy Averill-Murray / Arizona
right: Peter Paul van Dijk / Pima Co., Arizona


**
(orange dots = probably introduced or tanslocated; star = type locality)
Distribution: Mexico (*Sonora*), USA (*Arizona*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 205,771 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 284,624 sq. km
Size (Max SCL): male 31.4 cm, female 32.0 cm (Averill-Murray et al. 2002; Arizona Game and Fish Dept., unpubl. data)
IUCN Red List:* Vulnerable (VU A2abce+4abce) (Averill-Murray, Rosen, Jones, Jones, Lara-Resendiz, Edwards, Karl, and Berry 2023). Previously: TFTSG Provisional Red List: Vulnerable (VU) (2011, 2020)
CITES: Appendix II, as Testudinidae spp. (1977). Previously: Appendix II, as *Gopherus* spp. (1975)
USA ESA:* Assessed but Not Listed; Species Status Assessment: USFWS (2021f)
Synonymy:
Gopherus morafkai Murphy, Berry, Edwards, Leviton, Lathrop, and Riedle 2011:53 (11:11)
Xerobates morafkai
Type locality: "Tucson (approximate location 32°7' N, 110°56' W, elevation 948 m), Pima County, Arizona, U.S.A." [yellow star].
Type specimen: CAS 33867, holotype.
Etymology:* Eponym honoring David J. Morafka (1946–2004), USA; zoologist and herpetologist at California State University.

Gopherus polyphemus (Daudin 1801) (12:33, 21:78)
Gopher Tortoise


Kevin Main / CCB / Lake Wales Ridge, Florida


left: David Dennis / CRM 3 / Florida
right: John B. Iverson / nr. Orange Springs, Marion Co., Florida


**
(star = type localities)
Distribution: USA (*Florida, Georgia, South Carolina, Alabama, Mississippi, Louisiana*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 271,213 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 568,895 sq. km
Size (Max SCL): male 44.0 cm, female 42.5 cm; **(Max CCL):** male 54.5 cm (Mushinsky et al. 2006; Mrykalo et al. 2016; Bohlman et al. 2019)
IUCN Red List:* Critically Endangered (CR A2bcde+3b+4bce) (Stemle and Howell, in press). Previously: TFTSG Provisional Red List: Endangered (EN) (2011); Vulnerable (VU A1acde [ver. 2.3]) (TFTSG 1996)
CITES: Appendix II, as Testudinidae spp. (1977). Previously: Appendix II, as *Gopherus* spp. (1975)
USA ESA:* Threatened, in Western Distinct Population Segment (Louisiana, Mississippi, western Alabama) (USFWS 1987c); Species Status Assessment: USFWS (2021e)
Synonymy:
Testudo polyphaemus Bartram 1791:18 (*nomen nudum*)
Etymology:* Mythonym for *Polyphemus*, the Greek mythological cyclops from Homer's Odyssey, who lived in a cave, from Greek, *poly*, many, and *pheme*, voice; referring to the habitat of the species being a cave-like burrow.
Testudo polyphemus Daudin 1801:256
Emys polyphemus, Gopherus polyphemus, Xerobates polyphemus, Gopherus polyphemus polyphemus
Type locality: "l'Amérique septentrionale, principalement sur les bords de la rivière Savannah et près de l'Alatamaha" [USA]. Restricted to "vicinity of Savannah, Georgia" [USA] [yellow star] by Schmidt (1953:104).
Type specimen: Not known or located.
Etymology:* Mythonym for *Polyphemus*, the Greek mythological cyclops from Homer's Odyssey, who lived in a cave, from Greek, *poly*, many, and *pheme*, voice; referring to the habitat of the species being a cave-like burrow.
Testudo depressa Guérin 1829:pl.1,f.1
Type locality: Not designated. Restricted to "vicinity of Savannah, Georgia" [USA] [yellow star] by Schmidt (1953:105).
Type specimens: MNHN 9252 and 9269, syntypes (2), identified by Bour (2003).
Etymology:* Latin, *de-*, down, and *pressus*, pressed; referring to the depressed flattened shell.
Testudo carolina Le Conte 1830:97 (junior homonym and *nomen novum*, not = *Testudo carolina* Linnaeus 1758 [= *Terrapene carolina carolina*])
Xerobates carolinus, Gopherus carolinus
Type locality: "pine forests of Georgia and Florida, ...never found north of Savannah river" [USA]; restricted by default to "vicinity of Savannah, Georgia" [USA] [yellow star] by Schmidt (1953:104).
Type specimens: MNHN 9252 and 9269, syntypes (2), identified by Bour (2003).
Etymology:* Toponym for the Carolina region (North Carolina, South Carolina), USA.
Comment: Unjustified replacement name for *polyphemus*.
Testudo gopher Bartram *in* Gray 1844:4
Xerobates gopher
Type locality: Not designated. Restricted to "vicinity of Savannah, Georgia" [USA] [yellow star] by Schmidt (1953:105).
Type specimens: NHMUK 1839.10.18.30 and 1970.1916, syntypes (2).
Etymology:* Vernacular (English), *gopher*, anglicized from Vernacular (Old French), *gaufre*, local name for the species, first used by Bartram (1791).

***Gopherus*, sp. indet.** (12:34)
Testudo australis Girard 1858:470 (*nomen dubium et oblitum*) (12:34)
Gopherus australis
Type locality: "Bay of Islands, New Zealand" [in error, trade specimen; no star shown].
Type specimen: Probably USNM, apparently lost, see TTWG (2012; Annotation 12:34).
Etymology:* Latin, *austral*, the south, *australis*, from the south; referring to the provenance of the specimen from the southern hemisphere.
Comment: Possibly either *Gopherus agassizii* or *Gopherus polyphemus*.

Subfamily Testudininae Batsch 1788 (21:3)


Testudine Tortoises
(includes 16 genera, 45 species, 82 taxa)
Testudines Batsch 1788:437
Testudinina Gray 1869a:167
Testudininae Siebenrock 1909a:508

Aldabrachelys Loveridge and Williams 1957 (07:52, 07:53, 08:13, 09:29, 09:30, 12:28, 14:32, 17:55) (75)*

Indian Ocean Giant Tortoises
(includes 1 species, 4 taxa)
Megalochelys Fitzinger 1843:29 (junior homonym, not = *Megalochelys* Falconer and Cautley 1837 [= Testudinidae †])
Type species: *Testudo (Megalochelys) gigantea* [= *Testudo gigantea* Schweigger 1812], by original designation.
Etymology:* Greek, *megalos*, big, large, and *chelys*, turtle; referring to the large body size.
Aldabrachelys Loveridge and Williams 1957:225
Type species: *Geochelone* (*Aldabrachelys*) *gigantea* [= *Testudo gigantea* Schweigger 1812], by original designation.
Etymology:* Toponym for Aldabra Atoll, Seychelles, and Greek, *chelys*, turtle.
Dipsochelys Bour 1982a:117
Type species: *Dipsochelys elephantina* [= *Testudo elephantina* Duméril and Bibron 1835 = subjective synonym of *Testudo gigantea* Schweigger 1812], by original designation.
Etymology:* Greek, *dipso*, thirsty, and *chelys*, turtle; referring to the ability to drink or inhale water through its nostrils.

Aldabrachelys gigantea (Schweigger 1812) (07:54, 07:55, 08:13, 09:29, 09:30, 11:10, 12:28, 14:32, 17:56) (75, 76)*

Aldabra Giant Tortoise
(includes 4 subspecies)



(subspecies: *gigantea* = red range with yellow dot, *daudinii* = red dot, *arnoldi* and *hololissa* = orange dots [probable])
Distribution: Seychelles (*Aldabra*, *Granitic Islands*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 438 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 66,120 sq. km
Size (Max SCL): male 138.0 cm, female 114.0 cm (see subspp.)
IUCN Red List: Vulnerable (VU D2 [ver. 2.3]) (TFTSG 1996)
TFTSG Provisional Red List: Endangered (EN) (2019)
CITES: Appendix II, as Testudinidae spp. (1977). Previously: Appendix II, as *Geochelone* spp. (1975)

Aldabrachelys gigantea gigantea (Schweigger 1812) (07:54, 08:13, 09:29, 09:30, 11:10, 17:57)

Aldabra Giant Tortoise




Bernard Devaux / Aldabra, Seychelles





left: Peter Paul van Dijk / Aldabra, Seychelles / captivity / Singapore Zoo
right: Bernard Devaux / Aldabra, Seychelles



(subspecies: *gigantea* = red; stars = type localities)

Distribution: Seychelles (*Aldabra*)

Introduced: Mauritius (*Aigrettes, Rodrigues, Round*), Seychelles (*Assomption, Alphonse, Astove, Cerf, Cosmolédo, Cousin, Cousine, Curieuse, D'Arros, Desroches, Farquhar, Frégate, Grande Soeur, Moyenne, North, Rémire, Silhouette*), Tanzania (*Zanzibar [Changuu]*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 155 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 406 sq. km

Size (Max SCL): male 123.0 cm, female 97.0 cm; **(Max CCL):** male 160.0 cm, female 131.0 (Gerlach and Canning 1998; J. Gerlach, unpubl. data)

TFTSG Provisional Red List: Endangered (EN) (2019)

CITES: Appendix II, as Testudinidae spp. (1977). Previously: Appendix II, as *Geochelone* spp. (1975)

Synonymy:

Testudo gigantea Schweigger 1812:327 (*partim*, senior homonym and *nomen protectum*, misidentified type, not *Testudo gigantea* † (Grandidier 1868) [= *Aldabrachelys grandidieri* †]) (08:13, 09:29, 14:32)
Geochelone (Chelonoidis) gigantea, Geochelone gigantea, Testudo gigantea gigantea, Geochelone gigantea gigantea, Aldabrachelys gigantea, Megalochelys gigantea, Megalochelys gigantea gigantea, Scapia gigantea, Dipsochelys gigantea, Aldabrachelys gigantea gigantea, Dipsochelys giganteus
Type locality: "Brasilia" [Brazil]. Restricted to "Dune Patates, South Island, Aldabra Atoll, Republic of Seychelles" [-9.42 S, 46.40 E] [yellow star] by neotype designation by Frazier (2006:278).
Type specimen: USNM 269962, neotype, designated by Frazier (2006:278); MNHN 9554, the original holotype (= *Chelonoidis denticulatus*), rediscovered and designated by Bour (2006c:19), see also Ceríaco and Bour (2012) and Bour (2013a), but rejected and set aside by ICZN (2013b) in favor of the neotype designation.
Etymology:* Greek, *gigas*, giant, huge, *gigantea*, gigantic; referring to the very large body size.
Comment: Name conserved by ICZN (2013b), see Frazier (2008, 2009), Zug et al. (2009), Bour et al. (2009), and Takahashi et al. (2009). Name synonymized *in partim* under *Aldabrachelys gigantea gigantea* and *Chelonoidis denticulatus*.

Testudo dussumieri Schlegel *in* Gray 1830e:3 (*nomen nudum*)
Etymology:* Eponym honoring Jean-Jacques Dussumier (1792–1883), France; shipowner in the merchant marine and naturalist collector.

Testudo dussumieri Gray 1831d:9 (*nomen suppressum*) (09:29, 14:32)
Dipsochelys dussumieri, Geochelone dussumieri, Aldabrachelys dussumieri, Dipsochelys dussumieri dussumieri
Type locality: "Insula Mauritiana, Insula Aldebra" [Mauritius and Aldabra, Seychelles]. Restricted to "Aldabra (N.W. of Madagascar)" [yellow star] by Hubrecht (1881:44), and to "Insula Aldebra" [Aldabra, Seychelles] by Bour (1984b:291).
Type specimen: RMNH 3231, lectotype, designated by Bour (2006c:21), discussed by Hoogmoed et al. (2010) and Frazier and Matyot (2010), who concluded that its provenance was most likely the granitic Seychelles rather than Aldabra; genotyped by Austin et al. (2003).
Etymology:* Eponym honoring Jean-Jacques Dussumier (1792–1883), France; shipowner in the merchant marine and naturalist collector.
Comment: Name suppressed by ICZN (2013b), see Frazier (2008, 2009), Zug et al. (2009), Bour et al. (2009), and Takahashi et al. (2009).

Testudo elephantina Duméril and Bibron 1835:110
Testudo gigantea elephantina, Geochelone elephantina, Geochelone gigantea elephantina, Dipsochelys elephantina, Aldabrachelys elephantina, Dipsochelys elephantina elephantina
Type locality: "la plupart des îles qui sont situées dans le Canal de Mosambique, telle que Anjouan, Aldebra, les Comores, d'où on l'apporte fréquemment à Bourbon et à Maurice." Restricted to "Aldabra" [Seychelles] by Boulenger (1889:168); to "North Aldabra" by Rothschild (1915:418); and to "Ile Malabar, Aldabra Atoll, Republic of Seychelles" [green star] by Bour (1984b:291).
Type specimen:* MNHN 7874, lectotype, designated by Rothschild (1915:425) per Bour (1985:54); genotyped by Austin et al. (2003); MNHN 1942, 9314, 9562, 2002.160, paralectotypes, genotyped by et al. (2023).
Etymology:* Greek, *elefas*, elephant, and Latin, *-ina*, similarity of form or texture, *elephantina*, elephant-like; referring to the very large size.

Testudo ponderosa Günther 1877:35
Aldabrachelys ponderosa
Type locality: Not known. Restricted to "Aldabra" [Seychelles] [yellow star] by Bour (1984b:292).
Type specimen:* NHMUK 1947.3.4.89 (formerly 1864.12.20.27), lectotype, designated by Bour (1985:54), genotyped by Kehlmaier et al. (2023); NHMUK 1947.3.4.94 (formerly 1876.1.4.1), paralectotype, discussed by Gerlach and Canning (1998); genotyped by Austin et al. (2003).
Etymology:* Latin, *ponderosa*, heavy; referring to the heavy weight of the species.

Testudo sumeirei Sauzier 1892:398
Geochelone sumeirei, Dipsochelys sumeirei, Megalochelys sumeirei, Dipsochelys elephantina sumeirei, Aldabrachelys sumeirei, Dipsochelys dussumieri sumeirei, Testudo sumieri
Type locality: "probable...de Maurice..[&]..la Réunion..[&]..des

Séchelles." Restricted to "Seychelles" by Rothschild (1899:360); to "Seychelles Islands?" by Auffenberg (1974:144), and to "central Seychelles" [introduced; no star shown] by Bour (1984b:292).
Type specimen:* NHMUK 1947.3.4.1 (formerly 1929.12.27.1), holotype, discussed by Gerlach and Canning (1998); genotyped by Austin et al. (2003) and Kehlmaier et al. (2023).
Etymology:* Eponym honoring Benoit Camille Sumeire (1850–1920), Réunion / Mauritius; businessman and port station director who provided photos of the type.


Testudo gouffei Rothschild 1906:753
Testudo goufféi, Geochelone gouffei, Geochelone (Aldabrachelys) gouffei, Geochelone (Aldabrachelys) gigantea gouffei, Geochelone gigantea gouffei, Megalochelys gouffei, Aldabrachelys gouffei
Type locality: "Therese Island, St. Anne's Channel, Seychelles Islands." Emended to "more likely...from Juan de Novo or Farquhar Island" [Seychelles] [introduced; no star shown] by Rothschild (1915:427).
Type specimen:* NHMUK 1949.1.4.66, holotype; genotyped by Austin et al. (2003) and Kehlmaier et al. (2023).
Etymology:* Eponym honoring Alphonse A. Gouffé (1845–1931), United Kingdom; ship and insurance broker who obtained the type.

Aldabrachelys gigantea arnoldi (Bour 1982a) (07:54, 09:30, 11:10) (75)*
Arnold's Giant Tortoise


John Pemberton / CBFTT / Seychelles




Justin Gerlach / CBFTT / Seychelles


(subspecies: *arnoldi* = blue; orange dot and star [probable]; star = possible type locality)

Distribution: Seychelles (*Mahé?, North? [all extirpated]*)
Introduced: Seychelles (*North, Cousine, Frégate, Silhouette*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 162 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 492 sq. km
Size (Max SCL): male 96.0 cm, female 88.0 cm; **(Max CCL):** male 124.0 cm, female 111.0 cm (Gerlach and Canning 1998; J. Gerlach, unpubl. data)
CBFTT Account: Gerlach (2009)
TFTSG Provisional Red List: Critically Endangered (CR) (2019)
CITES: Appendix II, as Testudinidae spp. (1977). Previously: Appendix II, as *Geochelone* spp. (1975)
Synonymy:
Dipsochelys arnoldi Bour 1982a:121
Testudo arnoldi, Aldabrachelys arnoldi, Aldabrachelys gigantea arnoldi, Dipsochelys dussumieri arnoldi
Type locality: "Seychelles...probablement...les îles granitiques" [orange star, location estimated, possibly North Island].
Type specimen:* MNHN 9564, holotype, genotyped by Kehlmaier et al. (2023); NHMUK 1874.2.6.5, paratype, genotyped by Austin et al. (2003) and Kehlmaier et al. (2023); MNHN 7872, paratype, genotyped by Kehlmaier et al. (2023).
Etymology:* Eponym honoring E. Nicholas Arnold (1940–2023), United Kingdom; herpetologist at the British Museum of Natural History, London.

Aldabrachelys gigantea daudinii (Duméril and Bibron 1835) (09:30) (75)*
Daudin's Giant Tortoise
(Extinct, ca. 1850)


Roger Bour / Seychelles [MNHN, Paris]


(subspecies: *daudinii* = purple; red star [probable, extinct] = possible type locality)

Distribution: Seychelles (*Mahé? [extinct]*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 159 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 286 sq. km
Size (Max SCL): male 83.0 cm; **(Max CCL):** male 98.0 cm (Gerlach and Canning 1998)
TFTSG Provisional Red List: Extinct (EX) (2011)
CITES: Appendix II, as Testudinidae spp. (1977). Previously: Appendix II, as *Geochelone* spp. (1975)
Synonymy:
Testudo daudinii Duméril and Bibron 1835:123
Testudo gigantea daudinii, Geochelone gigantea daudinii, Dipsochelys daudinii, Geochelone daudinii, Aldabrachelys daudinii, Aldabrachelys gigantea daudinii, Dipsochelys dussumieri daudinii

Type locality: “Indes orientales.” Restricted to “les îles Seychelles granitiques” [red star = location estimated, possibly Mahé] by Bour (1985:58).

Type specimen:* MNHN 11819 (shell) and 7640 (its scutes) (now MNHN-ZA-AC-1884-549), holotype, discussed by Bour (1985) and Gerlach and Canning (1998); genotyped by Austin et al. (2003) and Kehlmaier et al. (2023).; Uetz et al. (2019) erroneously listed only MNHN 7640.

Etymology:* Eponym honoring François Marie Daudin (1774–1804), France; zoologist, ornithologist, and herpetologist who described many species of birds and reptiles.


Aldabrachelys gigantea hololissa (Günther 1877) (09:30, 11:10) (75)*
Seychelles Giant Tortoise



Justin Gerlach / CBFTT / Seychelles



Justin Gerlach / CBFTT / Seychelles



(subspecies: *hololissa* = green; orange dots and star [probable]; star = possible type locality)

Distribution: Seychelles (*Cerf [?], Cousine [?], Frégate [?], La Digue [?], Mahé [?], Praslin [?], Round [?], Silhouette [?] [all extirpated]*)

Introduced: Seychelles (*Cerf, Cousine, Round*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 229 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 2,803 sq. km

Size (Max SCL): male 138.0 cm, female 114.0 cm; **(Max CCL):** male 175.0 cm, female 147.0 cm (Gerlach and Canning 1998; J. Gerlach, unpubl. data)

CBFTT Account: Gerlach (2011)

TFTSG Provisional Red List: Critically Endangered (CR) (2019)

CITES: Appendix II, as Testudinidae spp. (1977). Previously: Appendix II, as *Geochelone* spp. (1975)

Synonymy:

Testudo hololissa Günther 1877:39
Dipsochelys hololissa, Geochelone hololissa, Aldabrachelys hololissa, Aldabrachelys gigantea hololissa, Dipsochelys dussumieri hololissa

Type locality: “Seychelle Islands...[probably] originally imported from the Aldabra group.” Restricted to “des îles Seychelles” [orange star, location estimated, possibly Praslin] by Sauzier (1899:142).

Type specimen:* UMZ R3796, neotype, designated by Gerlach and Canning (1998:12); RCSM 1021, lectotype, figured by Günther (1877:pl.7), designated by Rothschild (1915:426), destroyed during World War II (Bour 1985); RCSM 1020, paralectotype, also destroyed during WW II (Bour 1985); NHMUK 1888.3.20.1, paralectotype, labeled “type” by Boulenger (1889), however, identified by Gerlach and Canning (1998) as a *Dipsochelys dussumieri* [= *Aldabrachelys gigantea gigantea*] and removed from the type series, genotyped by Austin et al. (2003) and Kehlmaier et al. (2023) as a purported *A. g. hololissa*; Uetz et al. (2019) erroneously listed only NHMUK 1888.3.20.1 as syntype.

Etymology:* Greek, *holos*, whole, and *lissos*, smooth; referring to the shells of the originally examined specimens being perfectly smooth.

Dipsochelys resurrecta Gerlach and Canning 1996:133 (*nomen nudum*)

Etymology:* Latin, *resurrecta*, to be resurrected; referring to the apparent rediscovery of a taxon previously thought to be extinct.

Astrochelys Gray 1873j (07:52) (75)*

Madagascan Tortoises

(includes 2 species, 2 taxa)

Astrochelys Gray 1873j:4

Type species: *Testudo* (*Astrochelys*) *radiata* [= *Testudo radiata* Shaw 1802], by original monotypy.

Etymology:* Greek, *asteros*, a star, and *chelys*, turtle; referring to the starburst like markings on the carapace.

Asterochelys Gray 1874:724 (*nomen novum*)

Etymology:* Greek, *asteros*, a star, and *chelys*, turtle; referring to the starburst like markings on the carapace.

Comment:* Unjustified emendation of *Astrochelys*.

Angonoka Le, Raxworthy, McCord, and Mertz 2006:528 (09:31)

Type species: *Angonoka yniphora* [= *Testudo yniphora* Vaillant 1885a], by original designation.

Etymology:* Vernacular (Malagasy), *angonoka*, local indigenous name for the species.


Comment: Misspelled as *Anganoka* by Frétey (2024:8).

Astrochelys radiata (Shaw 1802)


Radiated Tortoise, *Sokake*



Anders G.J. Rhodin / TCC / CRM 6 / Cap Sainte Marie Special Reserve, Madagascar



Anders G.J. Rhodin / nr. Tsihombe, Cap Sainte Marie, Madagascar



** (star = type locality)

Distribution: Madagascar

Introduced: Mauritius (*Rodrigues*, *Round*), Réunion

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 28,082 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 56,574 sq. km

Size (Max SCL): male 39.5 cm, female 35.6 cm (Pedrono 2008; Leuteritz and Ganz 2013; Pedrono and Smith 2013); longterm captive: female 48.0 cm (N.A.T. Rakotoarisoa and R. Hudson, unpubl. data)

IUCN Red List: Critically Endangered (CR A4d, E) (Leuteritz and Rioux Paquette 2008). Previously: Vulnerable (VU) (TFTSG *in* Baillie and Groombridge 1996)

CITES: Appendix I (1975)

USA ESA:* Endangered, as *Geochelone radiata* (USFWS 1970)

Synonymy:

Testudo coui Daudin 1801:271 (*nomen oblitum*)

Type locality: Not known.

Type specimen:* Possibly MNHN, not located; type figured (pl.26).

Etymology:* Vernacular (Old French), *coui*, decorated calabash bowl often used by natives for eating and drinking; referring to the hemispherical shape of the tortoise shell with its striations.

Testudo radiata Shaw 1802:22

Psammobates radiatus, Asterochelys radiata, Testudo radiata radiata, Geochelone radiata, Astrochelys radiata

Type locality: "Madagascar." Restricted to "Soalara (Baie de Saint-Augustin), sud-ouest de Madagascar" [yellow star] by Bour (1979:152).

Type specimen: NHMUK 1947.3.5.15, holotype.

Etymology:* Latin, *radius*, ray, spoke, radius, *radiata*, radiated; referring to the characteristic radiated shell pattern.

Testudo madagascariensis Schweigger 1812:457 (*nomen nudum*)

Etymology:* Toponym for Madagascar.

Testudo desertorum Grandidier 1869:257

Type locality: "Madagascar."

Type specimen: Not known or located.

Etymology:* Latin, *desertorum*, coming from the desert; referring to the very arid habitat of this taxon.

Testudo hypselonota Bourret 1941b:9

Type locality: "provenant d'un Chinois de Cholon qui l'avait...achetée au marché...il n'est pas certain qu'elle ait été trouvée en Cochinchine" [Vietnam] [trade specimen, no star shown]. Restricted to "Cholon?...Indochina" by Wermuth and Mertens (1961:213) [in error]. Shown to be identical to *Testudo radiata* from Madagascar by Auffenberg (1963a:465); type locality restricted to "durch den Tierhandel...von Madagascar nach Indochina" by Wermuth (1965:285).


Type specimen: MNHN 1948.41 (formerly VNUH T.85), holotype, see Auffenberg (1963a) and Wermuth (1965).

Etymology:* Greek, *ypelos*, high, steep, and *notos*, the back; referring to the highly domed carapace.

Astrochelys yniphora (Vaillant 1885a) (07:56, 09:31, 21:79)
Ploughshare Tortoise, Plowshare Tortoise, *Angonoka*


Anders G.J. Rhodin / CCB / Baly Bay National Park, Madagascar


Anders G.J. Rhodin / Madagascar / captivity / Antananarivo



(orange dot = trade; star = type locality)


Distribution: Madagascar
Introduced: Mauritius (*Rodrigues*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 884 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 2,886 sq. km
Size (Max SCL): male 51.9 cm, female 42.6 cm (Pedrono and Markwell 2001; Pedrono 2008; Pedrono and Smith 2013; Pedrono and Clausen 2018)
IUCN Red List: Critically Endangered (CR A4ad, B2ab(v), C1,E) (Leuteritz and Pedrono 2008). Previously: Endangered (EN) (TFTSG *in* Baillie and Groombridge 1996)
CITES: Appendix I (1975)
USA ESA:* Endangered, as *Geochelone yniphora* (USFWS 1976)
Synonymy:
Testudo yniphora Vaillant 1885a:441
Testudo radiata yniphora, Asterochelys yniphora, Geochelone yniphora, Geochelone (Asterochelys) yniphora, Astrochelys yniphora, Angonoka yniphora, Anganoka yniphora
Type locality: "un îlot situé au nord-nord-est de...grande Comore ...d'une localité située vers Aldabra" [in error; no star]. Restricted to "cap d'Amparafaka (Baie de Baly), nord-ouest de Madagascar" [yellow star] by Bour (1979:152).
Type specimen: MNHN 1885.499, holotype, apparently lost, specimen figured (pl.13-14), see Bour (2007a).
Etymology:* Greek, *ynio*, ploughshare, and *phora*, carrying; referring to the distinctive ploughshare-like projection from the anterior plastron.
Testudo hyniphora Vaillant *in* Vaillant and Grandidier 1910:40 (*nomen novum*)
Etymology:* Greek, *ynio*, ploughshare, and *phora*, carrying; referring to the distinctive ploughshare-like projection from the anterior plastron.
Comment: Unjustified emendation of *yniphora*.

Centrochelys Gray 1872c (07:52) (25)*
African Spurred Tortoises
(includes 1 species, 1 taxon)
Centrochelys Gray 1872c:5
Type species: *Peltastes (Centrochelys) sulcatus* [= *Testudo sulcata* Miller 1779], by original monotypy.
Etymology:* Greek, *kentron*, spike, spur, and *chelys*, turtle; referring to the large, conical, spur-like horn scales on the back of the thighs.

Centrochelys sulcata (Miller 1779) (12:29)
African Spurred Tortoise, Grooved Tortoise, Sahel Tortoise, *Tortue Sillonnée*


Tomas Diagne / Katane, Senegal / male


left and center: Tomas Diagne / Senegal / subadult
right: Håkan Pohlstrand / Kafta Sheraro National Park, Ethiopia / female


(orange dots = probable trade or translocated; stars = type localities)

Distribution:* Algeria, Benin, Burkina Faso, Cameroon, Central African Republic, Chad, Djibouti (*extirpated*), Eritrea, Ethiopia, Mali, Mauritania, Niger, Nigeria, Saudi Arabia, Senegal, Sudan, Togo (?), Yemen

Introduced: Costa Rica, Ethiopia, Gambia, Somalia, Spain, USA (*Hawaii*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 3,416,077 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 6,524,249 sq. km

Size (Max SCL): male 86.0 cm, female 57.8 cm (Lambert 1993; Ernst et al 2006b; Itescu et al. 2014; Ardjima et al. 2020); **(Max CCL):** male 101.0 cm, female 67.0 cm (L. Luiselli and F. Petrozzi, unpubl. data)

CBFTT Account: Petrozzi, Hema, Demaya, Benansio, Eniang, Diagne, Segniagbeto, and Luiselli (2020)

IUCN Red List: Endangered (EN A4bcd) (Petrozzi, Luiselli, Hema, Diagne, Segniagbeto, Eniang, Leuteritz, and Rhodin 2021). Previously: Vulnerable (VU) (TFTSG 1996)

CITES: Appendix II, as Testudinidae spp. (1977); zero annual export quota for specimens removed from the wild and traded for primarily commercial purposes (2000). Previously: Appendix II, as *Geochelone* spp. (1975)

Synonymy:

Testudo sulcata Miller 1779:pl.26
Geochelone (Geochelone) sulcata, Geochelone sulcata, Peltastes sulcatus, Centrochelys sulcatus, Centrochelys sulcata
Type locality: "India occidentali" [in error]. Restricted to "West Indies" [in error] [no star] by Miller and Shaw (1796:54).
Type specimen: Not located, type specimen figured (pl.26), discussed by Bour (2013b).
Etymology:* Latin, *sulcus*, groove or furrow, and *-ata*, having; referring to the furrowed carapace.

Testudo calcarata Schneider 1784:317 (*nomen novum*)
Chersine calcarata
Type locality: "Westindien." [in error] [no star].
Type specimen: Figured in Miller (1779:pl.26), by default.
Etymology:* Latin, *calcaratus*, furnished with a spur; referring to the large thigh spurs.
Comment: Unjustified replacement name for *sulcata*.

Testudo radiata senegalensis Gray 1831d:11
Geochelone senegalensis, Geochelone sulcata senegalensis
Type locality: "Senegal" [orange star, probable trade, location estimated].
Type specimen: MNHN, holotype, not located.
Etymology:* Toponym for Senegal.

Testudo schoepfii Rüppell *in* Gray 1873j:13 (junior homonym and *nomen nudum*, not *Testudo schoepfii* Fitzinger 1826 [= *Kinixys erosa*]).
Type locality:* "Abyssinia" [Eritrea] [green star, location estimated].
Type specimen: NHMUK, holotype, not located, see Gray (1873j).
Etymology:* Eponym honoring Johann David Schoepff (or Schöpf) (1752–1800), Germany; physician, botanist, herpetologist, and turtle taxonomist.

Geochelone sulcata sudanensis Ballasina, Vandepitte, Mochi, and Fenwick 2006:111 (*nomen nudum*)
Etymology:* Toponym for Sudan.

Chelonoidis Fitzinger 1835 (07:52, 17:58)

South American Tortoises, Neotropical Tortoises
(includes 4 species, 18 taxa)

Chelonoidis Fitzinger 1835:112
Type species: *Testudo (Chelonoidis) boiei* [= *Testudo boiei* Wagler 1830a = subjective synonym of *Testudo carbonaria* Spix 1824], by subsequent designation by Fitzinger (1843:29). Genus established as *Geochelone (Chelonoidis)* without a type species.
Etymology:* Greek, *chelone*, turtle, and *-ides*, similarity; referring to its similarity to the genus *Geochelone*.

Gopher Gray 1870a:190
Type species: *Testudo (Gopher) chilensis* Gray 1870a, by original monotypy.
Etymology:* Vernacular (English), *gopher*, anglicized from Vernacular (Old French), *gaufre*.

Elephantopus Gray 1874:724 (junior homonym, not = *Elephantopus* Agassiz 1847 [= Siphonophora])
Type species: *Elephantopus planiceps* [= *Testudo planiceps* Gray 1854b = *Chelonoidis porteri*], by original monotypy.
Etymology:* Greek, *elefas*, elephant, and *pus*, foot, leg; referring to the elephant-like feet.

Pampatestudo Lindholm 1929:285 (*nomen novum*)
Type species: *Testudo (Pampatestudo) chilensis* [= *Testudo (Gopher) chilensis* Gray 1870a], by original monotypy.
Etymology:* Toponym for the Pampas region, Argentina, and Latin, *testudo*, tortoise.

Monachelys Williams 1952:547
Type species: *Testudo (Monachelys) monensis* † Williams 1952, by original designation.
Etymology:* Toponym for Mona Island, Puerto Rico, and Greek, *chelys*, turtle.

Darwintestudo Antenbrink-Vetter and Vetter 1998:4
Type species: *Darwintestudo hoodensis* [= *Testudo hoodensis* Van Denburgh 1907], by original designation.
Etymology:* Eponym honoring Charles Darwin (1809–1882), United Kingdom; naturalist and proponent of the theory of evolution by natural selection who studied tortoises in the Galapagos Islands, and Latin, *testudo*, tortoise.

Chelonoidis carbonarius (Spix 1824) (10:19, 10:20, 14:33, 17:58, 21:80) (77)*

Red-footed Tortoise, *Jabuti-Vermelho*, *Morrocoy Pata Roja*, *Jabuti-Piranga*




Cassiano Zaparoli / Fazenda Caiman, Mato Grosso do Sul, Brazil



left: Eric V. Goode / captivity / El Impenetrable National Park, Chaco, Argentina
right: Cassiano Zaparoli / Fazenda Caiman, Mato Grosso do Sul, Brazil


**


**


**

(orange dots = questionable localities, probably introduced or trade or translocated; star = likely type locality)

Distribution:* Argentina (*Chaco, Formosa, Salta*), Bolivia (*Beni, Cochabamba, La Paz, Pando, Santa Cruz*), Brazil (*Alagoas, Amazonas, Bahia, Ceará, Maranhão, Mato Grosso, Mato Grosso do Sul, Pará, Paraíba, Pernambuco, Piauí, Rondônia, Roraima, Sergipe*), Colombia (*Antioquia, Arauca, Atlántico, Bolívar, Boyacá, Caldas, Caquetá, Casanare, Cauca, Cesar, Chocó, Córdoba, Cundinamarca, Guainía, La Guajira, Magdalena, Meta, Santander, Sucre, Tolima, Vichada*), French Guiana, Guyana, Panama (*Darién, San Blas*), Paraguay, Suriname, Venezuela (*Apure, Barinas, Bolívar, Carabobo, Cojedes, Falcón, Guárico, Mérida, Miranda, Portuguesa, Sucre, Yaracuy, Zulia*)

Introduced (prehistorical and historical):* Anguilla, Antigua, Argentina (*Misiones*), Barbuda, Barbados, Brazil (*Acre, Espirito Santo, Minas Gerais, Rio de Janeiro, Santa Catarina, São Paulo*), British Virgin Islands, Colombia (*Amazonas, Providencia, San Andrés, Valle del Cauca*), Cuba, Dominica, Ecuador, Grenada, Guadeloupe, Jamaica, Martinique, Montserrat, Netherlands Antilles, Nicaragua (*Maíz Grande*), Panama, Peru (*San Martín*), Saint-Barthélemy, Saint Kitts and Nevis, Saint Lucia, Saint Martin, Saint Vincent and the Grenadines, Trinidad, U.S. Virgin Islands (*St. Croix, St. John, St. Thomas*), Venezuela (*Isla Margarita, Los Tostigos*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 5,323,930 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 11,429,957 sq. km

Size (Max SCL):* male 59.3 cm, female 44.2 (Pritchard 1980b; Pritchard and Trebbau 1984; Vinke and Vinke 1999; Vinke et al. 2008)

IUCN Red List:* Endangered (EN A2cd+4cde) (Ceballos, Vinke, Vinke, Rojano Bolaño, Morcatty, Escalona, and Gallego-García, in press). Previously: TFTSG Provisional Red List: Vulnerable (VU) (2011); Least Concern (LC) [Not Listed] (Behler 1996)

CITES: Appendix II, as Testudinidae spp. (1977). Previously: Appendix II, as *Geochelone* spp. (1975)

Synonymy:

Testudo carbonaria Spix 1824:22
Testudo tabulata carbonaria, Geochelone carbonaria, Chelonoidis carbonaria, Geochelone carbonaria carbonaria, Chelonoidis carbonaria carbonaria, Chelonoidis carbonarius
Type locality:* "flumen Amazonum" [Amazonas or Pará, Brazil]. Clarified as the Rio Amazonas between the mouth of the Tajapuru (01º02'S, 51º02'W) and the mouth of the Negro (03º08'S, 59º55'W) by Vanzolini (1981:xxvi), likely the vicinity of Parintins, Amazonas, Brazil [yellow star].
Type specimen: ZSM, holotype or syntypes, not known, apparently lost; specimen figured (pl.16), lectotype, designated by Hoogmoed and Gruber (1983:354), also see Franzen and Glaw (2007).
Etymology:* Latin, *carbo*, coal, *carbonaria*, coal-like; referring to the dark carapace color.
Comment:* Spix collected extensively while staying in Parintins on the Rio Amazonas (Vanzolini 1981), probably including the type specimen.

Testudo boiei Wagler 1830a:pl.13 (14:33)
Geochelone (Chelonoidis) boiei, Geochelone boiei, Chelonoidis boiei
Type locality: Not known.
Type specimen: ZSM, holotype or syntypes, not known, apparently lost; specimen figured (pl.13), lectotype, designated by Hoogmoed and Gruber (1983:354), also see Franzen and Glaw (2007).
Etymology:* Eponym honoring Heinrich Boie (1794–1827), Germany / Netherlands; naturalist and herpetologist who worked at the Rijksmuseum van Natuurlijke Historie, Leiden.

Testudo hercules truncata Gray 1830e:3 (10:7, 10:20)
Type locality: "South America."
Type specimen: OUM 8461, holotype, discussed by Nowak-Kemp and Fritz (2010).
Etymology:* Latin, *truncata*, truncated, blunt, capped; referring to the conically raised and truncated carapace scutes of the holotype.


Geochelone atlantica † López-Jurado, Mateo, and García-Márquez 1998 (21:80)
Centrochelys atlantica †
Type locality: "cráter de Pedra Lume...isla de Sal (Archipiélago de Cabo Verde)" [Cape Verde] [error; no star]. However, the type specimen's genotype closely aligns with extant specimens of *Chelonoidis carbonarius* from Paraguay (Kehlmaier et al. 2021b).
Type specimen: DBULPGC 17, holotype, femur, genotyped by Kehlmaier et al. (2021b).
Etymology:* Toponym for the Atlantic Ocean.
Comment: Originally described from a femur and two peripheral bones as alleged Quaternary fossils embedded in chalky tufa-like matrix from Cape Verde, but shown by Kehlmaier et al. (2021b) to

be from a single individual of *Chelonoidis carbonarius*, likely from Paraguay, that was alive at least from 1962 to 1974. How the type specimen bones became embedded in the matrix is unknown.


Chelonoidis chilensis (Gray 1870a) (07:57, 10:21, 12:30, 17:59, 21:81)
Chaco Tortoise, Pampas Tortoise, Argentine Tortoise, *Tortuga Chaqueña*



Maurice Rodrigues / TTN / Monte Desert, nr. Neuquén, Neuquén, Argentina



left: Leandro Alcalde / Baldecitos, San Juan Prov., Argentina
right: Thomas and Sabine Vinke / Boquerón Dept., Paraguay



** (orange dots = probable trade or translocated; stars = type localities)

Distribution:* Argentina (*Buenos Aires, Catamarca, Chaco, Chubut, Córdoba, Formosa, La Pampa, La Rioja, Mendoza, Neuquén, Rio Negro, Salta, San Juan, San Luis, Santa Fe, Santiago del Estero, Tucumán*), Bolivia (*Chuquisaca, Santa Cruz, Tarija*), Paraguay

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 1,118,096 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 1,786,245 sq. km

Size (Max SCL): male 23.9 cm, female 43.3 cm (Freiberg 1973; Fritz et al. 2012a)

IUCN Red List:* Endangered (EN A2cd+4cde) (Kubisch, Alcalde, and Waller, in press). Previously: TFTSG Provisional Red List: Vulnerable (VU) (2011); Vulnerable (VU A1cd [ver. 2.3]) (TFTSG 1996), as *Geochelone chilensis*

CITES: Appendix II, as Testudinidae spp. (1977). Previously: Appendix II, as *Geochelone* spp. (1975)

Synonymy:

Testudo (Gopher) chilensis Gray 1870a:190
Testudo chilensis, Geochelone chilensis, Geochelone chilensis chilensis, Chelonoidis chilensis, Chelonoidis chilensis chilensis
Type locality: "Chili" [Chile; in error], see Sclater (1870:470). Emended to "Chili...N. Patagonia...Mendoza and the Pampas..[&]..Monte Video and Buenos Ayres" by Gray (1870d:707); and restricted to "Mendoza" [Argentina] [yellow star] by Boulenger (1889:159).
Type specimens: NHMUK 1947.3.5.8–9 (both formerly 1870.12.18.2), syntypes (2), possibly lost (McCarthy, pers. comm. *in* Vinke et al. 2008).
Etymology:* Toponym for Chile.

Testudo argentina Sclater 1870:471 (*nomen novum*)
Comment: Unjustified replacement name for *chilensis*.
Etymology:* Toponym for Argentina.

Geochelone donosobarrosi Freiberg 1973:83 (12:30)
Geochelone chilensis donosobarrosi, Chelonoidis chilensis donosobarrosi, Chelonoidis donosobarrosi
Type locality: "San Antonio, Río Negro" [Argentina] [green star].
Type specimen: USNM 192961, holotype, see Reynolds et al. (2007).
Etymology:* Eponym honoring Roberto Donoso-Barros (1922–1975), Chile; physician, biologist, and herpetologist at the Universidad de Concepción.

Geochelone petersi Freiberg 1973:86 (12:30)
Chelonoidis chilensis petersi, Chelonoidis petersi, Geochelone chilensis petersi
Type locality: "Kishka, La Banda, Santiago del Estero" [Argentina] [blue star].
Type specimen: USNM 192959, holotype, see Reynolds et al. (2007).
Etymology:* Eponym honoring James A. Peters (1922–1972), USA; zoologist, herpetologist, and curator at the Smithsonian U.S. National Museum, Washington, D.C.

Chelonoidis denticulatus (Linnaeus 1766) (10:19, 17:58)
Yellow-footed Tortoise, *Jabuti, Jabuti-Tinga, Morrocoy Pata Amarilla*



Carlos Alberto Jimenez / Madre de Dios, Peru



left: Guta Agostini / Amazonas, Brazil / captivity
right: Frank Deschandol / French Guiana



(light red = extirpated; red dots = extirpated, orange dots = probably trade or translocated; stars = type localities)

Distribution: Bolivia (*Beni, La Paz, Pando, Santa Cruz*), Brazil (*Acre, Amapá, Amazonas, Bahia, Espírito Santo, Maranhão, Mato Grosso, Mato Grosso do Sul, Pará, Rio de Janeiro, Rondônia, Roraima, Tocantins*), Colombia (*Amazonas, Arauca, Caquetá, Casanare, Guainía, Guaviare, Meta, Putumayo, Vaupés, Vichada*), Ecuador, French Guiana, Guyana, Peru (*Cusco, Loreto, Madre de Dios, Pasco, Ucayali*), Suriname, Trinidad, Venezuela (*Amazonas, Bolívar, Delta Amacuro, Monagas*)

Introduced:* Dominica

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 5,799,636 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 10,304,369 sq. km

Size (Max SCL):* male 82.0 cm, female 75.9 cm (Pritchard and Trebbau 1984; Sajdak and Molina 1991)

IUCN Red List:* Endangered (EN A2bcd+4bcd) (Tavares, Morcatty, Rojas-Runjaic, Rojano Bolaño, Fagundes, and Valadão, in press). Previously: TFTSG Provisional Red List: Near Threatened (NT) (2011); Vulnerable (VU A1cd+2cd [ver. 2.3]) (TFTSG 1996)

CITES: Appendix II, as Testudinidae spp. (1977). Previously: Appendix II, as *Geochelone* spp. (1975)

Synonymy:

Testudo denticulata Linnaeus 1766:352
Chersine denticulata, Geochelone (Geochelone) denticulata, Geochelone denticulata, Chelonoidis denticulata, Chelonoidis denticulatus
Type locality: "Virginia" [USA, in error] [no star].
Type specimens: UUZM s/n, NRM s/n (formerly MDG 21), syntypes (2), discussed by Schoepff (1801:119) and Andersson (1900:25), both apparently lost, listed by Thunberg (1828), but not by Holm (1957) or Wallin (2001), UUZM s/n figured in Schoepff (1801:pl.28,f.1).
Etymology:* Latin, *denticulus*, small tooth, *denticulata*, serrated, denticulated; referring to serrated carapacial margin in juveniles.
Comment: Description cited as sourced from "Mus. De Geer" (MDG).

Testudo tabulata Walbaum 1782:122 (unavailable name)
Type locality: Not designated.
Type specimens: UPSZTY 281–83 (formerly UUZM 281–83 and MGA 50–52), possible syntypes (3), see Wallin (2001).
Etymology:* Latin, *tabula*, panel, board, *tabulata*, paneled; referring to the shell having the appearance of wood paneling.
Comment: Unavailable name from a non-binomial work, see Wermuth (1956).

Testudo tessellata Stobaeus *in* Schneider 1792:262
Chersine tessellata
Type locality: "America australi" [South America].
Type specimen: Not known or located.
Etymology:* Latin, *tessela*, little cube, mosaic stone, *tessellata*, having cubes; referring to the brightly colored, angular scute areolae.
Comment: Description cited as sourcing Schneider (1783:362) and the unnamed species described in Gmelin (1789:1045,n.33), which in turn cited Stobaeus (1730:59).

Testudo stobaeana Gmelin *in* Schoepff 1793a:48 (*nomen nudum*)
Etymology:* Eponym honoring Kilian Stobaeus (1690–1742), Sweden; physician, paleontologist, and naturalist.

Testudo tabulata Walbaum *in* Schoepff 1793a:56
Chersine tabulata, Geochelone (Chelonoidis) tabulata, Geochelone tabulata, Chelonoidis tabulata
Type locality: "Africa australi?" [South Africa, in error].
Type specimens: UPSZTY 281–83 (formerly UUZM 281–83 and MGA 50–52), possible syntypes (3), listed by Thunberg (1828) and Wallin (2001), type specimens figured in Schoepff (1793:pl.12,f.2,pl.13-14).
Etymology:* Latin, *tabula*, panel, board, *tabulata*, paneled; referring to the shell having the appearance of wood paneling.

Testudo gigantea Schweigger 1812:327 (*partim*, misidentified type, senior homonym, not *Testudo gigantea* † (Grandidier 1868) [= *Aldabrachelys grandidieri* †]) (08:13, 09:29)
Type locality: "Brasilia" [Brazil]. Restricted to "Dune Patates, South Island, Aldabra Atoll, Republic of Seychelles" by neotype designation by Frazier (2006:278).
Type specimen: USNM 269962, neotype (= *Aldabrachelys gigantea*), designated by Frazier (2006:278); MNHN 9554, the original holotype (= *Chelonoidis denticulata*), rediscovered and designated by Bour (2006c:19), see also Ceríaco and Bour (2012) and Bour (2013a).
Etymology:* Greek, *gigas*, giant, huge, *gigantea*, gigantic; referring to the very large body size.
Comment: Original holotype and name rejected and set aside by ICZN (2013b) in favor of the neotype designation. Name synonymized *in partim* under *Aldabrachelys gigantea gigantea* and *Chelonoidis denticulatus*.

Testudo terrestris brasiliensis Seba *in* Schweigger 1812:445 (*nomen nudum*)
Testudo brasiliensis
Etymology:* Toponym for Brazil.

Testudo terrestris americana Stobaeus *in* Schweigger 1812:445 (*nomen novum*)
Etymology:* Toponym for the Americas (South America).

Testudo tabulata cayennensis Schweigger 1812:445 (*nomen nudum*)
Testudo terrestris cayennensis
Etymology:* Toponym for Cayenne, French Guiana.

Testudo terrestris surinamensis Stedman *in* Schweigger 1812:445 (*nomen nudum*)
Etymology:* Toponym for Suriname.

Testudo hercules Spix 1824:20
Type locality:* "sylvis ad flumen Solimoens" [Amazonas, Brazil] [= between Tefé and Tabatinga, possibly near São Paulo de Olivença] [green star, location estimated].
Type specimen: ZSM, holotype or syntypes, not known, apparently lost, including type specimen figured (pl.14), see Hoogmoed and Gruber (1983).
Etymology:* Mythonym for *Hercules*, the Greek mythological strong man; referring to the large strong shell.

Testudo sculpta Spix 1824:21 (senior homonym, not *Testudo sculpta* Brandt *in* Gray 1856b [= *Chersina angulata*]).
Type locality:* "sylvis juxta flumen Amazonum" [Amazonas or Pará, Brazil]. Clarified as the Rio Amazonas between the mouth of the Tajapuru (01º02'S, 51º02'W) and the mouth of the Negro (03º08'S, 59º55'W) by Vanzolini (1981:xxvi), likely the vicinity of Parintins, Amazonas [blue star].

Type specimen: ZSM 2753/0/1, lectotype, designated by Hoogmoed and Gruber (1983:354), see Franzen and Glaw (2007).
Etymology:* Latin, *sculpta*, carved, sculpted, chiseled, engraved; referring to the clearly pronounced growth rings on the carapace scutes.
Comment:* Spix collected extensively while staying in Parintins on the Rio Amazonas (Vanzolini 1981), probably including the type specimen.


Testudo cagado Spix 1824:23
Type locality:* "campis et nemoribus campestribus Bahiae" [Bahia, Brazil] [possibly in the Ilhéus region of Bahia near Almada on the Rio Itaípe (14º40'S, 39º23'W), as per Vanzolini (1981)] [pink star, location estimated].
Type specimen: ZSM, holotype or syntypes, not known, apparently lost, including type specimen figured (pl.17), see Hoogmoed and Gruber (1983).
Etymology:* Vernacular (Portuguese), *cágado*, vernacular name for turtles or tortoises in Brazil.
Comment:* Spix collected extensively at Almada on the Rio Itaípe (Vanzolini 1981:xiv), probably including the type specimen.

Testudo planata Gmelin *in* Gray 1831d:9 (*nomen nudum*)
Etymology:* Latin, *planata*, flattened; probably referring to the slightly flattened carapace.

Testudo foveolata Schinz 1833:40 (*nomen nudum*)
Etymology:* Latin, *foveola*, small round pit, *foveolata*, having small round pits; probably referring to small shell depressions.

Chelonoidis niger (Quoy and Gaimard 1824b) (09:32, 12:31, 17:58, 17:60, 21:82) (78, 79, 80)*

Galápagos Giant Tortoise
(includes 15 subspecies, of which one is unnamed)



**

Distribution: Ecuador (*Galápagos: Española [Hood]; Fernandina [Narborough] (possibly extinct); Floreana [Charles] (extinct); Isabela [Albemarle]; Pinta [Abingdon] (extinct); Pinzón [Duncan]; San Cristóbal [Chatham]; Santa Cruz [Indefatigable]; Santa Fé [Barrington] (extinct); Santiago [San Salvador; James]*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 3,043 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 34,131 sq. km
Size (Max SCL): male 135.8 cm, female 122.0 cm; **(Max CCL):** male 199.7 cm, female 197.2 cm; 95th Percentile CCL: male 151.0 cm, female 118.0 cm (see subspp.) (21:83)
IUCN Red List: Vulnerable (VU A2c, B1+2c [ver. 2.3]) (TFTSG 1996), as *Chelonoidis nigra*, but see all more recently updated subspecies assessments
CITES: Appendix I (1975)
USA ESA:* Endangered, as *Geochelone nigra* (= *elephantopus*) (USFWS 1970)


Chelonoidis niger niger (Quoy and Gaimard 1824b) (07:58, 09:33, 12:31, 17:58, 17:62, 21:84)

Floreana Giant Tortoise, Charles Island Giant Tortoise

(Extinct, ca. 1850)



Peter C.H. Pritchard / CRM 1 / Floreana [NHMUK, London]



(stars = type localities)

Distribution: Ecuador (*Galápagos: Floreana [Charles] [extinct]*)

Introduced: Ecuador (*Galápagos: Isabela [Albemarle]) (hybrids with C. becki*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 78 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 86 sq. km

Size (Max SCL): male ca. 96.0 cm [estimated]; **(Max CCL):** male ca. 105.0 cm (Broome 1929)

IUCN Red List: Extinct (EX) (van Dijk, Rhodin, Cayot, and Caccone 2017), as *Chelonoidis niger*. Previously: Extinct (EX) (TFTSG 1996), as *Chelonoidis nigra nigra*

CITES: Appendix I, as *Chelonoidis niger* (1975)

Synonymy:

Testudo californiana Quoy and Gaimard 1824a:90 (*nomen oblitum*) (17:62)

Testudo californica

Type locality: "La Californie...donnée vivante aux îles Sandwich" [California, USA; in error]. Erroneously given as "Sandwich Islands" [Hawaii, USA] [no star] by Wermuth and Mertens (1961, 1977).

Type specimen: MNHN 9550, holotype, discussed by Pritchard (1996) and Olson (2015).

Etymology:* Toponym for California, USA.

Testudo nigra Quoy and Gaimard 1824b:174 (17:62, 21:84) (*nomen novum*)

Chelonoidis nigra, Geochelone nigra, Geochelone nigra nigra, Chelonoidis nigra nigra, Geochelone elephantopus nigra, Chelonoidis niger, Chelonoidis niger niger

Type locality: "la Californie...donnée vivante, aux îles Sandwich" [California, USA; in error]. Restricted arbitrarily to "Cerro Azul d'Isabela" [Galápagos, Ecuador] [in error, orange star] by Bour *in* David (1994:83); and to "Charles Island (Santa María or Floreana)" [Galápagos, Ecuador] [red star] by Pritchard (1996:49).

Type specimen: MNHN 9550, holotype, discussed by Pritchard (1996) and Olson (2015).

Etymology:* Latin, *nigra*, black; referring to the shell color.

Comment: Unjustified replacement name for *californiana*.

Testudo elephantopus Harlan 1826:284 (09:33, 17:62, 21:84) (*nomen dubium*)

Testudo elephantopus elephantopus, Geochelone elephantopus, Geochelone elephantopus elephantopus, Chelonoidis elephantopus, Chelonoidis elephantopus elephantopus, Geochelone nigra elephantopus

Type locality: "Gallapagos Islands" [Galápagos, Ecuador]. Restricted to "South Albemarle" [Isabela, Galápagos, Ecuador] [orange star, location estimated] by Pritchard (1967:168); restricted to "Charles Island (Floreana, Santa Maria), Galapagos" [Ecuador] [red star] by Olson and Humphrey (2017:114).

Type specimens: ANSP, holotype or syntype, apparently lost, not listed by Malnate (1971), discussed by Olson and Humphrey (2017); MCZ 11063, possible syntype, designated by Garman (1917:292), see also Barbour and Loveridge (1929:350).

Etymology:* Greek, *elefas*, elephant, and *pus*, foot, leg; referring to the elephant-like feet.

Testudo galapagoensis Baur 1889:1044

Testudo elephantopus galapagoensis, Geochelone elephantopus galapagoensis, Chelonoidis galapagoensis, Chelonoidis elephantopus galapagoensis, Geochelone (Chelonoidis) nigra galapagoensis, Geochelone nigra galapagoensis, Chelonoidis nigra galapagoensis

Type locality: "Charles Island" [Floreana, Galápagos, Ecuador] [red star].


Type specimen: MCZ 11064, lectotype, designated as the "type" by Garman (1917:289) and the "holotype" by Barbour and Loveridge (1929); Olson (2017) suggested that MCZ 11069 and 11070 should be regarded as "syntypes" (paralectotypes); MCZ 11070 genotyped by Poulakakis et al. (2008).

Etymology:* Toponym for the Galapagos Islands, Ecuador.

Chelonoidis niger abingdonii (Günther 1877) (09:34, 12:31)
Pinta Giant Tortoise, Abingdon Island Giant Tortoise
(Extinct, 2012)



Anders G.J. Rhodin / TCC / Pinta / male / Lonesome George, CDRS, Santa Cruz



(stars = type localities)


Distribution: Ecuador (*Galápagos: Pinta [Abingdon] [extinct]*)
Introduced: Ecuador (*Galápagos: Isabela [Albemarle]) (hybrids with C. becki*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 15 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 19 sq. km
Size (Max SCL): male 99.1 cm, female 70.0 cm [estimated]; **(Max CCL):** male 106.0 cm, female ca. 85.0 cm [estimated] (Pritchard 2005; Chiari 2020) (21:83)
IUCN Red List: Extinct (EX) (Cayot, Gibbs, Tapia, and Caccone 2016), as *Chelonoidis abingdonii*. Previously: Extinct in the Wild (EW) (TFTSG 1996)
CITES: Appendix I, as *Chelonoidis niger* (1975)
Synonymy:
Testudo ephippium Günther 1874:422 (*nomen nudum*)
Etymology:* Greek, *ephippion*, saddle blanket, horse blanket; referring to the saddle-shaped carapace.
Testudo ephippium Günther 1875a:271 (*partim*, previously misidentified type)
Type locality: "Charles Island" [Floreana, Galápagos, Ecuador] [in error, orange star]. Restricted to "Duncan" [Pinzón, Galápagos, Ecuador] by Van Denburgh (1914:259) [in error, brown star], and to "Pinta (Abingdon)?" [Galápagos, Ecuador] [red star] by Garman (1917:pl.39).
Type specimen: NMSZ 1932.027.001–012 (shell and skeletal parts of same specimen) (formerly RSM 1822.058), holotype, see Baur (1889), Garman (1917), Herman et al. (1990), Pritchard (1996), and Olson (2017); NHMUK 1874.6.1.6, listed as "syntype" in the NHMUK catalogue, is not a type.
Etymology:* Greek, *ephippion*, saddle blanket, horse blanket; referring to the saddle-shaped carapace.
Comment:* Name synonymized *in partim* under *Chelonoidis niger abingdonii* and *Chelonoidis niger duncanensis*.
Testudo abingdonii Günther 1877:85
Testudo elephantopus abingdonii, Geochelone abingdonii, Geochelone elephantopus abingdonii, Chelonoidis abingdonii, Chelonoidis elephantopus abingdonii, Geochelone (Chelonoidis) nigra abingdonii, Geochelone nigra abingdonii, Chelonoidis nigra abingdonii, Chelonoidis elephantopus abingdonii, Chelonoidis niger abingdonii
Type locality: "Abingdon Island" [Pinta, Galápagos, Ecuador] [red star].
Type specimens: NHMUK 1947.3.4.39, 1947.3.4.95–96 (formerly 1876.6.21.38–40), syntypes (3).
Etymology:* Eponym honoring the first Earl of Abingdon (James Bertie [1653–1699]), United Kingdom; nobleman in whose honor Abingdon Island was also named.

Chelonoidis niger becki (Rothschild 1901) (12:31, 17:60)

Volcán Wolf Giant Tortoise, Wolf Volcano Giant Tortoise



Paul M. Gibbons / Volcán Wolf, Isabela / non-hybrid



(stars = type localities)

Distribution: Ecuador (*Galápagos: Isabela [Albemarle]*)

Invasives: Hybrids with *Chelonoidis abingdonii* and *C. niger*

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 234 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 271 sq. km

Size (Max SCL): male ca. 105.0 cm, female ca. 83.0 cm [both estimated]; **(Max CCL):** male 140.0 cm, female 110.4 cm; 95th Percentile CCL: male128.0 cm, female 87.0 cm (Galapagos Conservancy and Galapagos National Park Directorate, unpubl. data; Chiari 2020) (21:83)

IUCN Red List: Vulnerable (VU A1bde) (Caccone, Cayot, Gibbs, and Tapia 2017), as *Chelonoidis becki*. Previously: Vulnerable (VU) (TFTSG 1996), as *Chelonoidis nigra becki*

CITES: Appendix I, as *Chelonoidis niger* (1975)

Synonymy:

Testudo becki Rothschild 1901:372

Geochelone becki, Geochelone elephantopus becki, Chelonoidis becki, Chelonoidis elephantopus becki, Geochelone (Chelonoidis) nigra becki, Geochelone nigra becki, Chelonoidis nigra becki, Chelonoidis niger becki

Type locality: “Cape Berkeley, northern point of Albemarle Island, Galapagos Archipelago” [Isabela, Galápagos, Ecuador] [possibly in error, orange star]. Emended to “Bank’s Bay, Albemarle” [Isabela, Galápagos, Ecuador] [yellow star] by Van Denburgh (1914:259).

Type specimen: NHMUK 1949.1.3.87, holotype.


Etymology:* Eponym honoring Rollo H. Beck (1870–1950), USA; ornithologist and explorer who collected the type.

Chelonoidis niger chathamensis (Van Denburgh 1907) (07:59, 12:31, 17:61) (79)*

San Cristóbal Giant Tortoise, Chatham Island Giant Tortoise



Washington Tapia / San Cristóbal



**

(star = type locality)

Distribution: Ecuador (*Galápagos: San Cristóbal [Chatham]*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 282 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 368 sq. km

Size (Max SCL): male ca. 95.0 cm, female ca. 79.0 cm [both estimated]; **(Max CCL):** male 124.7 cm, female 103.5 cm; 95th Percentile CCL: male 110.0 cm, female 85.0 cm (Galapagos Conservancy and Galapagos National Park Directorate, unpubl. data; Chiari 2020) (21:83)

IUCN Red List: Endangered (EN A1abde) (Caccone, Cayot, Gibbs, and Tapia 2017), as *Chelonoidis chathamensis*. Previously: Vulnerable (VU) (TFTSG 1996), as *Chelonoidis nigra chathamensis*

CITES: Appendix I, as *Chelonoidis niger* (1975)

Synonymy:

Testudo chathamensis Van Denburgh 1907:4

Testudo elephantopus chathamensis, Geochelone chathamensis, Geochelone elephantopus chathamensis, Chelonoidis chathamensis, Chelonoidis elephantopus chathamensis, Geochelone (Chelonoidis) nigra chathamensis, Geochelone nigra chathamensis, Chelonoidis nigra chathamensis, Chelonoidis niger chathamensis

Type locality: “Chatham Island, Galapagos Archipelago” [San Cristóbal, Galápagos, Ecuador] [yellow star].


Type specimen:* CAS 8127, holotype, see Slevin and Leviton (1956) and Pritchard (1996), genotyped by Jensen et al. (2022a); CAS 8133, paratype, genotyped by Poulakakis et al. (2012) and Jensen et al. (2022a).

Etymology:* Toponym for Chatham Island (San Cristóbal), Galapagos, Ecuador.

Chelonoidis niger darwini (Van Denburgh 1907) [12:31, 17:60]
Santiago Giant Tortoise, James Island Giant Tortoise



Tui De Roy/Roving Tortoise Photos / Santiago



(star = type localities)

Distribution: Ecuador (*Galápagos: Santiago [San Salvador] [James]*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 173 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 183 sq. km

Size (Max SCL): male ca. 107.0 cm, female ca. 95.0 cm [both estimated]; **(Max CCL):** male 150.2 cm, female 134.4 cm; 95th Percentile CCL: male 134.0 cm, female 95.0 cm (Galapagos Conservancy and Galapagos National Park Directorate, unpubl. data; Chiari 2020) [21:83]

IUCN Red List: Critically Endangered (CR A1bde) (Cayot, Gibbs, Tapia, and Caccone 2016), as *Chelonoidis darwini*. Previously: Endangered (EN) (TFTSG 1996), as *Chelonoidis nigra darwini*

CITES: Appendix I, as *Chelonoidis niger* (1975)

Synonymy:

Testudo schweiggeri Fitzinger 1826:44 (*nomen nudum*)
Etymology:* Eponym honoring Augustus F. Schweigger (1783–1821), Germany; naturalist, herpetologist, and turtle taxonomist.


Geochelone (Geochelone) schweiggeri Fitzinger 1835:122 (junior homonym and *nomen dubium et oblitum*, not = *Testudo schweiggeri* Gray *in* Duméril and Bibron 1835 [= *Testudo schweigeri* Gray 1830e] [= *Cylindraspis triserrata*])
Geochelone schweiggeri
Type locality: Not designated. Restricted to "America: Insel St. Jacob der Gallopagen" [= Santiago, Galápagos, Ecuador] [yellow star, location estimated] by Fitzinger (1853:110).
Type specimen: Originally live in MSW, see Fitzinger (1853), apparently lost, possibly in NMW, not located or listed by Tiedemann and Häupl (1980) or Tiedemann et al. (1994).
Etymology:* Eponym honoring Augustus F. Schweigger (1783–1821), Germany; naturalist, herpetologist, and turtle taxonomist.

Testudo darwini Van Denburgh 1907:4
Testudo elephantopus darwini, Geochelone darwini, Geochelone elephantopus darwini, Chelonoidis darwini, Chelonoidis elephantopus darwini, Geochelone (Chelonoidis) nigra darwini, Geochelone nigra darwini, Chelonoidis nigra darwini, Chelonoidis niger darwini
Type locality: "James Island, Galapagos Archipelago" [Santiago, Galápagos, Ecuador] [yellow star].
Type specimen: CAS 8108, holotype, see Slevin and Leviton (1956).
Etymology:* Eponym honoring Charles Darwin (1809–1882), United Kingdom; naturalist and proponent of the theory of evolution by natural selection who studied tortoises in the Galapagos Islands.

Chelonoidis niger donfaustoi Poulakakis, Edwards, and Caccone *in* Poulakakis, Edwards, Chiari, Garrick, Russello, Benavides, Watkins-Colwell, Glaberman, Tapia, Gibbs, Cayot, and Caccone 2015 [17:61]
Eastern Santa Cruz Giant Tortoise, Cerro Fatal Giant Tortoise, Don Fausto's Giant Tortoise



Peter Paul van Dijk / Cerro Fatal, Santa Cruz



(star = type locality)

Distribution: Ecuador (*Galápagos: Santa Cruz [Indefatigable]*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 89 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 90 sq. km

Size (Max SCL): male ca. 112.0 cm, female ca. 87.0 cm [both estimated]; **(Max CCL):** male 172.6.0 cm, female 133.0 cm; 95th Percentile CCL: male 141.0 cm, female 114.0 cm (Galapagos Conservancy and Galapagos National Park Directorate, unpubl. data; Chiari 2020) [21:83]

IUCN Red List: Critically Endangered (CR A2abce; B1ab(ii)) (Cayot, Gibbs, Tapia, and Caccone 2017), as *Chelonoidis donfaustoi*

CITES: Appendix I, as *Chelonoidis niger* (1975)

Synonymy:

Chelonoidis donfaustoi Poulakakis, Edwards, and Caccone *in* Poulakakis, Edwards, Chiari, Garrick, Russello, Benavides, Watkins-Colwell, Glaberman, Tapia, Gibbs, Cayot, and Caccone 2015:12
Chelonoidis nigra donfaustoi, Chelonoidis niger donfaustoi
Type locality: "Cerro Fatal in Santa Cruz" [Galápagos, Ecuador]

[yellow star].
Type specimen: UWZM 32700, holotype; genotyped by Poulakakis et al. (2015).
Etymology:* Eponym honoring Don Fausto Llerena Sánchez (1940–), Ecuador; park ranger at Galápagos National Park responsible for the care of the giant tortoises at the Charles Darwin Station.

Chelonoidis niger duncanensis (Pritchard 1996) (07:60, 12:31)
Pinzón Giant Tortoise, Duncan Island Giant Tortoise



Peter C.H. Pritchard / CRM 1 / Pinzón / male / Onan


(orange dot = introduced; stars = type localities)

Distribution: Ecuador (*Galápagos: Pinzón [Duncan]*)
Introduced: Ecuador (*Galápagos: Santa Cruz [Indefatigable]*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 10 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 11 sq. km
Size (Max SCL): male ca. 85.0 cm, female ca. 80.0 cm [both estimated]; **(Max CCL):** male 101.2 cm, female 94.6 cm; 95th Percentile CCL: male 94.0 cm, female 80.0 cm (Galapagos Conservancy and Galapagos National Park Directorate, unpubl. data; Chiari 2020) (21:83)
IUCN Red List: Vulnerable (VU A1abde; D1+2) (Cayot, Gibbs, Tapia, and Caccone 2017), as *Chelonoidis duncanensis*. Previously: Extinct in the Wild (EW) (TFTSG 1996), as *Chelonoidis nigra duncanensis*
CITES: Appendix I, as *Chelonoidis niger* (1975)
Synonymy:
Testudo ephippium Günther 1875a:271 (*partim*, previously misidentified type)
Testudo elephantopus ephippium, Geochelone elephantopus ephippium, Geochelone ephippium, Chelonoidis ephippium, Chelonoidis nigra ephippium, Chelonoidis elephantopus ephippium, Geochelone (Chelonoidis) nigra ephippium, Geochelone nigra ephippium
Type locality: "Charles Island" [Floreana, Galápagos, Ecuador] [in error, orange star]. Restricted to "Duncan" [Pinzón, Galápagos, Ecuador] by Van Denburgh (1914:259) [in error, brown star, partly hidden], and to "Pinta (Abingdon)?" [Pinta, Galápagos, Ecuador] [red star] by Garman (1917:pl.39).
Type specimen: NMSZ 1932.027.001–012 (shell and skeletal parts of same specimen) (formerly RSM 1822.058), holotype, see Baur (1889), Garman (1917), Herman et al. (1990), Pritchard (1996), and Olson (2017); NHMUK 1874.6.1.6, listed as "syntype" in the NHMUK catalogue, is not a type.
Etymology:* Greek, *ephippion*, saddle blanket, horse blanket; referring to the saddle-shaped carapace.
Comment:* Name synonymized *in partim* under *Chelonoidis niger abingdonii* and *Chelonoidis niger duncanensis*.
Testudo duncanensis Garman 1917:269 (*nomen nudum*)
Type locality: "Pinzon (Duncan)" [Galápagos, Ecuador] [yellow star].
Type specimen: MCZ 11068, labeled by Garman (unpubl.) as "type of Baur's *Testudo duncanensis*."
Etymology:* Toponym for Duncan Island (Pinzón), Galapagos, Ecuador.
Geochelone nigra duncanensis Pritchard 1996:47
Chelonoidis nigra duncanensis, Chelonoidis duncanensis, Chelonoidis niger duncanensis
Type locality: "Duncan Island" [Pinzón, Galápagos, Ecuador] [yellow star].
Type specimen: MCZ 11068, holotype, designated lectotype of *Testudo duncanensis* Garman 1917 (*nomen nudum*), by Pritchard (1996:47); specimen figured in Garman (1917:pl.41).
Etymology:* Toponym for Duncan Island (Pinzón), Galapagos, Ecuador.

Chelonoidis niger guntheri (Baur 1889) (07:64, 08:14, 09:36, 12:31, 17:60)
Sierra Negra Giant Tortoise



Tui De Roy/Roving Tortoise Photos / Sierra Negra, Isabela / captivity


(stars = type localities)

Distribution: Ecuador (*Galápagos: Isabela [Albemarle]*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 499 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 672 sq. km
Size (Max SCL): male ca. 94.0 cm, female ca. 77.0 cm [both estimated]; **(Max CCL):** male 136.5 cm, female 111.4 cm; 95th Percentile CCL: male 129.0 ccm, female 99.0 cm (Galapagos Conservancy and Galapagos National Park Directorate, unpubl. data; Chiari 2020) (21:83)
IUCN Red List: Critically Endangered (CR A2bde) (Cayot, Gibbs, Tapia, and Caccone 2018), as *Chelonoidis guntheri*. Previously: Endangered (EN) (TFTSG 1996), as *Chelonoidis nigra guentheri*
CITES: Appendix I, as *Chelonoidis niger* (1975)
Synonymy:
Testudo guntheri Baur 1889:1044 (senior homonym, not = *Testudo guntheri* Gadow *in* Van Denburgh 1914 [= *Cylindraspis triserrata*]) (07:64, 09:36)
Testudo güntheri, Geochelone elephantopus güntheri, Geochelone güntheri, Geochelone (Chelonoidis) nigra güntheri, Geochelone nigra güntheri, Chelonoidis elephantopus güntheri, Geochelone elephantopus guntheri, Geochelone guntheri, Geochelone (Chelonoidis) nigra guntheri, Geochelone nigra guntheri, Chelonoidis elephantopus guntheri, Chelonoidis guntheri, Chelonoidis niger guntheri
Type locality: "Galapagos." Restricted to "Vilamil, Albemarle" [Isabela, Galápagos, Ecuador] [yellow star] by Van Denburgh (1914:259).
Type specimen: OUM 8656, lectotype, designated by Pritchard (1996:48), figured in Günther (1877:pl.30,f.A); Nowak-Kemp and Fritz (2010) listed this specimen as one of several syntypes.
Etymology:* Eponym honoring Albert C.L.G. Günther (1830–1914), Germany / United Kingdom; zoologist, herpetologist, ichthyologist, and curator at the British Museum of Natural History, London.
Testudo wallacei Rothschild 1902:619 (12:31)
Testudo elephantopus wallacei, Geochelone elephantopus wallacei, Geochelone wallacei, Chelonoidis elephantopus wallacei, Chelonoidis nigra wallacei, Geochelone nigra wallacei
Type locality: "Chatham Island" [San Cristóbal, Galápagos, Ecuador] [in error, orange star] by supposition. Restricted to "Jervis" [Rábida, Galápagos, Ecuador] [in error, brown star] by Van Denburgh (1914:259).
Type specimen: NHMUK 1949.1.4.43, holotype.
Etymology:* Eponym honoring Alfred Russell Wallace (1823–1913), United Kingdom; naturalist who, with Charles Darwin, championed the idea of evolution by natural selection.
Comment:* Species description based on a single specimen of unknown origin, considered by Van Denburgh (1914) to possibly be from Jervis (Rábida), where tortoises may have been introduced from southern Isabela.
Geochelone elephantopus guentheri Baur *in* Pritchard 1971b:50 (junior homonym and *nomen novum*, not = *Testudo guentheri* Gadow 1894 [= *Cylindraspis triserrata*]) (07:64, 09:36)
Testudo guentheri, Geochelone guentheri, Chelonoidis guentheri, Chelonoidis elephantopus guentheri, Geochelone nigra guentheri, Chelonoidis nigra guentheri
Etymology:* Eponym honoring Albert C.L.G. Günther (1830–1914), Germany / United Kingdom; zoologist, herpetologist, ichthyologist, and curator at the British Museum of Natural History, London.
Comment: Unjustified emendation of *güntheri* (= *guntheri*).

Chelonoidis niger hoodensis (Van Denburgh 1907) (12:31)
Española Giant Tortoise, Hood Island Giant Tortoise


Peter C.H. Pritchard / TCC / Española / captivity / CDRS, Santa Cruz


(orange dot = introduced; star = type locality)

Distribution: Ecuador (*Galápagos: Española [Hood]*)
Introduced: Ecuador (*Galápagos: Santa Fé [Barrington]*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 21 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 22 sq. km
Size (Max SCL): male ca. 97.0 cm, female ca. 78.0 cm [both estimated]; **(Max CCL):** male 119.2 cm, female 96.0 cm;

95th Percentile CCL: male 91.0 cm, female 76.0 cm (Galapagos Conservancy and Galapagos National Park Directorate, unpubl. data; Chiari 2020) (21:83)

IUCN Red List: Critically Endangered (CR A1bde) (Cayot, Gibbs, Tapia, and Caccone 2017), as *Chelonoidis hoodensis*. Previously: Critically Endangered (CR) (TFTSG 1996), as *Chelonoidis nigra hoodensis*

CITES: Appendix I, as *Chelonoidis niger* (1975)

Synonymy:

Testudo hoodensis Van Denburgh 1907:3

Testudo elephantopus hoodensis, Geochelone elephantopus hoodensis, Geochelone hoodensis, Chelonoidis hoodensis, Chelonoidis elephantopus hoodensis, Geochelone (Chelonoidis) nigra hoodensis, Geochelone nigra hoodensis, Chelonoidis nigra hoodensis, Darwintestudo hoodensis, Chelonoidis niger hoodensis

Type locality: "Hood Island, Galapagos Archipelago" [Española, Galápagos, Ecuador] [yellow star].

Type specimen: CAS 8121, holotype, see Slevin and Leviton (1956).


Etymology:* Toponym for Hood Island (Española), Galapagos, Ecuador.

Chelonoidis niger microphyes (Günther 1875a) (07:64, 08:14, 09:36, 12:31)

Volcán Darwin Giant Tortoise, Darwin Volcano Giant Tortoise, Tagus Cove Giant Tortoise



Anders G.J. Rhodin / Volcán Darwin, nr. Tagus Cove, Isabela



(stars = type localities)

Distribution: Ecuador (*Galápagos: Isabela [Albemarle]*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 111 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 121 sq. km

Size (Max SCL): male ca. 73.8 cm, female ca. 64.5 cm [both estimated]; **(Max CCL):** male 121.0 cm, female 104.0 cm; 95th Percentile CCL: male 120.0 cm, female 103.0 cm (Galapagos Conservancy and Galapagos National Park Directorate, unpubl. data; Chiari 2020) (21:83)

IUCN Red List: Endangered (EN A1bde) (Cayot, Gibbs, Tapia, and Caccone 2018), as *Chelonoidis microphyes*. Previously: Vulnerable (VU) (TFTSG 1996), as *Chelonoidis nigra microphyes*

CITES: Appendix I, as *Chelonoidis niger* (1975)

Synonymy:

Testudo microphyes Günther 1874:422 (*nomen nudum*)

Etymology:* Greek, *mikros*, small, and *fysis*, body, *microphyes*, having a small body; referring to the relatively small size of this taxon.

Testudo microphyes Günther 1875a:275 (08:14)

Geochelone elephantopus microphyes, Geochelone microphyes, Chelonoidis microphyes, Chelonoidis elephantopus microphyes, Geochelone (Chelonoidis) nigra microphyes, Geochelone nigra microphyes, Chelonoidis nigra microphyes, Chelonoidis niger microphyes

Type locality: "Hood's Island" [Española, Galápagos, Ecuador] by supposition [in error, orange star]. Emended to "Tagus Cove, about 4 miles inland...northern Albemarle Island" [Isabela, Galápagos, Ecuador] [yellow star] by Günther (1877:78).

Type specimen: NHMUK 1947.3.4.88 (formerly 1875.12.29.1), holotype.

Etymology:* Greek, *mikros*, small, and *fysis*, body, *microphyes*, having a small body; referring to the relatively small size of this taxon.

Testudo macrophyes Garman 1917:273

Type locality: "Santa Isabela Island (Albemarle) near Tagus Cove" [Isabela, Galápagos, Ecuador] [yellow star].

Type specimens: NHMUK 1876.6.21.41, 1947.3.4.91–3 (formerly 1876.6.21.42, 1880.12.20.1, 1886.3.5.1), syntypes (4).

Etymology:* Greek, *makros*, large, and *fysis*, body, *macrophyes*, having a large body; referring to the relatively large size of this taxon.

Chelonoidis niger phantasticus (Van Denburgh 1907) (07:62, 12:31, 17:58, 17:63, 21:85) (80)*


Fernandina Giant Tortoise, Narborough Island Giant Tortoise



Peter C.H. Pritchard / CRM 1 / Fernandina / male [CAS, San Francisco]



©Galápagos Conservancy / Fernandina / female




** (star = type locality)

Distribution: Ecuador (*Galápagos: Fernandina [Narborough]*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 109 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 142 sq. km
Size (Max SCL): male 88.0 cm, female ca. 45.0 cm [estimated]; **(Max CCL):** male 106.7 cm, female 59.5 cm (Van Denburgh 1914; Giant Tortoise Restoration Initiative, unpubl. data)
IUCN Red List: Critically Endangered (Possibly Extinct) (CR(PE) D) (Rhodin, Gibbs, Cayot, Kiester, and Tapia 2017), as *Chelonoidis phantasticus*. Previously: Critically Endangered (Possibly Extinct) (CR(PE)) (TFTSG Provisional Red List 2017); Extinct (EX) (TFTSG Provisional Red List 2011)
TFTSG Provisional Red List: Critically Endangered (CR) (2021)
CITES: Appendix I, as *Chelonoidis niger* (1975)
Synonymy:
Testudo phantasticus Van Denburgh 1907:4
Testudo phantastica, Testudo elephantopus phantastica, Geochelone elephantopus phantastica, Geochelone phantastica, Chelonoidis phantastica, Geochelone phantasticus, Chelonoidis elephantopus phantastica, Geochelone (Chelonoidis) nigra phantastica, Geochelone nigra phantastica, Chelonoidis nigra phantastica, Chelonoidis phantasticus, Chelonoidis niger phantasticus
Type locality: "Narborough Island, Galapagos Archipelago" [Fernandina, Galápagos, Ecuador] [yellow star].
Type specimen: CAS 8101, holotype, see Slevin and Leviton (1956) and Pritchard (1996); genotyped by Poulakakis et al. (2012).
Etymology:* Greek, *fantasia*, fantasy, *phantasticus*, fantastic; referring to the unusually prominent saddle-backed shell with enlarged and raised anterior shell opening in the single original male specimen collected.

Chelonoidis niger porteri (Rothschild 1903) (07:63, 09:35, 12:31, 17:62, 21:86)
Western Santa Cruz Giant Tortoise, Indefatigable Island Giant Tortoise


Russell A. Mittermeier / El Chato, Santa Cruz


(star = type localities]


Distribution: Ecuador (*Galápagos: Santa Cruz [Indefatigable]*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 306 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 319 sq. km
Size (Max SCL): male 135.8 cm, female ca. 122.0 cm (Pritchard 1996:37; White 2015) [female estimated]; **(Max CCL):** male 191.7 cm, female 197.2 cm; 95th Percentile CCL: male 151.0 cm, female 118.0 cm (Galapagos Conservancy and Galapagos National Park Directorate, unpubl. data; Chiari 2020) (21:83)
IUCN Red List: Critically Endangered (CR A2bde) (Cayot, Gibbs, Tapia, and Caccone 2017), as *Chelonoidis porteri*. Previously: Endangered (EN) (TFTSG 1996), as *Chelonoidis nigra porteri*
CITES: Appendix I, as *Chelonoidis niger* (1975)
Synonymy:
Testudo nigrita Duméril and Bibron 1835:80 (senior homonym and *nomen dubium*, not = *Testudo nigrita* Tamayo 1962 [= *Chelonia mydas*]) (17:62, 21:86)
Testudo elephantopus nigrita, Geochelone nigrita, Chelonoidis nigrita, Geochelone elephantopus nigrita, Chelonoidis elephantopus nigrita, Geochelone (Chelonoidis) nigra nigrita, Geochelone nigra nigrita, Chelonoidis nigra nigrita
Type locality: Not known. Restricted to "Insel Indefatigable" [Santa Cruz, Galápagos, Ecuador] [yellow star, location estimated] by Mertens and Wermuth (1955:376).
Type specimen: NHMUK 1949.1.4.37, lectotype, designated as "type" by Günther (1875a:268), discussed by Olson (2015).
Etymology:* Latin, *nigrita*, blackish; referring to relatively dark overall color of the shell.
Testudo planiceps Gray 1854b:12 (junior homonym and *nomen dubium*, not = *Testudo planiceps* Schneider 1792 [= *Platemys platycephala platycephala*]) (21:86)
Geochelone planiceps, Chelonoidis planiceps
Type locality: "Galapagos Islands" [Galápagos, Ecuador] [no star].
Type specimen: NHMUK 1947.3.5.3 (formerly 1848.4.6.3), holotype.
Etymology:* Latin, *planis*, flat, plane, and *ceps*, head; referring to the flattened head.
Testudo porteri Rothschild 1903:119
Geochelone elephantopus porteri, Geochelone porteri, Geochelone nigra porteri, Chelonoidis elephantopus porteri, Chelonoidis nigra porteri, Chelonoidis porteri, Chelonoidis niger porteri
Type locality: "Indefatigable Island, Galapagos group" [Santa Cruz, Galápagos, Ecuador] [yellow star, location estimated].
Type specimen: NHMUK 1949.1.4.38 and 1974.2464 (shell and bones from same specimen), holotype; genotyped by Poulakakis et al. (2015).
Etymology:* Eponym honoring Captain David Porter (1780–1843), USA; naval commander, the first to explore Santa Cruz, first named Porter's Island, then Indefatigable, who first noted differences among the tortoises from various Galapagos Islands (Porter 1815).

Chelonoidis niger vandenburghi (DeSola 1930) (07:64, 08:14, 09:36, 12:31, 17:60)

Volcán Alcedo Giant Tortoise, Alcedo Volcano Giant Tortoise



Peter C.H. Pritchard / CRM 1 / TCF / Volcán Alcedo, Isabela



(stars = type localities]

Distribution: Ecuador (*Galápagos: Isabela [Albemarle]*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 317 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 318 sq. km

Size (Max SCL): male 129.5 cm, female 90.2 cm [female estimated]; **(Max CCL):** male 199.7 cm, female 145.5 cm; 95th Percentile CCL: male 148.0 cm, female 107.0 cm (Pritchard 1971a; Galapagos Conservancy and Galapagos National Park Directorate, unpubl. data; Chiari 2020) (21:83)

IUCN Red List: Vulnerable (VU A1bde) (Cayot, Gibbs, Tapia, and Caccone 2018), as *Chelonoidis vandenburghi*. Previously: Vulnerable (VU) (TFTSG 1996), as *Chelonoidis nigra vandenburghi*

CITES: Appendix I, as *Chelonoidis niger* (1975)

Synonymy:

Testudo vandenburghi DeSola 1930:80

Geochelone vandenburghi, Geochelone elephantopus vandenburghi, Chelonoidis vandenburghi, Chelonoidis elephantopus vandenburghi, Geochelone (Chelonoidis) nigra vandenburghi, Geochelone nigra vandenburghi, Chelonoidis nigra vandenburghi, Chelonoidis niger vandenburghi

Type locality: "mid-Albemarle Island...forty miles from Villamil...at the coast on the southern border of Perry Isthmus...Cowley Mountain nearly north, while behind us...to the south...Villamil Mountain" [Isabela, Galápagos, Ecuador] [orange star]. Emended to "Cowley Mountain...the first mountain north of Villamil Mountain" [= Volcán Alcedo, Isabela, Galápagos, Ecuador] [yellow star] by DeSola (1930:80).

Type specimen: CAS 8141, holotype, see Slevin and Leviton (1956).


Etymology:* Eponym honoring John Van Denburgh (1872–1924), USA; physician, herpetologist, and curator at the California Academy of Sciences, who studied Galapagos tortoises.

Chelonoidis niger vicina (Günther 1875a) (07:64, 08:14, 09:36, 12:31, 17:60)

Cerro Azul Giant Tortoise, Iguana Cove Giant Tortoise



Vincenzo Ferri / Cerro Azul, Isabela / captivity



(star = type locality)

Distribution: Ecuador (*Galápagos: Isabela [Albemarle]*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 357 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 402 sq. km

Size (Max SCL): male 84.2 cm, female 71.8 cm [both estimated]; **(Max CCL):** male 138.0 cm, female 115.8 cm; 95th Percentile CCL: male 123.0 cm, female 98.0 cm (Galapagos Conservancy and Galapagos National Park Directorate, unpubl. data; Chiari 2020) (21:83)

IUCN Red List: Endangered (EN A2bde) (Cayot, Gibbs, Tapia, and Caccone 2018), as *Chelonoidis vicina*. Previously: Endangered (EN) (TFTSG 1996), as *Chelonoidis nigra vicina*

CITES: Appendix I, as *Chelonoidis niger* (1975)

Synonymy:

Testudo vicina Günther 1874:422 (*nomen nudum*)

Etymology:* Latin, *vicinus*, related or similar; referring to the close relationship between *vicina* and *microphyes*.

Testudo vicina Günther 1875a:277

Geochelone vicina, Geochelone (Chelonoidis) nigra vicina, Geochelone nigra vicina, Geochelone elephantopus vicina, Chelonoidis elephantopus vicina, Chelonoidis nigra vicina, Chelonoidis vicina, Chelonoidis niger vicina

Type locality: "Galapagos" [Galápagos, Ecuador]. Restricted to "South Albemarle" [Isabela, Galápagos, Ecuador] by Günther (1877:73); and to "Iguana Cove, Albemarle" [Isabela, Galápagos, Ecuador] [yellow star] by Van Denburgh (1914:259).

Type specimen: NHMUK 1947.3.4.90 (formerly 1874.7.15.1), holotype.

Etymology:* Latin, *vicinus*, related or similar; referring to the close relationship between *vicina* and *microphyes*.

***Chelonoidis niger* unnamed subspecies** (21:87)

Santa Fé Giant Tortoise, Barrington Island Giant Tortoise

(Extinct, ca. 1890 [?])



Rebekah Kim / Plate 123, Fig. 2, Van Denburgh 1914 / Santa Fé
© California Academy of Sciences



(red dot = extinct, orange dot = possible original provenance)

Distribution: Ecuador (*Galápagos: Santa Fé [Barrington] [?] [extinct]; possibly San Cristóbal [Chatham] [?] [extinct]*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 18 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 19 sq. km

Size (Max SCL): unknown; no shells known

IUCN Red List: Not Evaluated (NE)

TFTSG Provisional Red List: Extinct (EX) (2021)

Synonymy:

Chelonoidis sp. (Santa Fe) Poulakakis et al. (2015:123)

Specimens: CAS 8142–54, appendicular bones without associated skulls or shells, collected on Santa Fé (Barrington Island) in 1906.

Comment:* Mitochondrial DNA of specimens CAS 8143, 8145–6, and 8148 genotyped by Poulakakis et al. (2012) and shown to be distinct and separate from a clade containing *abingdonii*, *hoodensis*, *chathamensis*, and *donfaustoi* (the latter still unnamed at that time). The original provenance of this taxon is questionable (Pritchard 1996:65), possibly southwestern San Cristóbal, but it has been referred to as an unnamed extinct taxon from Santa Fé by many since 2012 (see Caccone 2020; Gibbs et al. 2020, TTWG 2021). Subsequent analysis of the same specimens by Jensen et al. (2022a) demonstrated that this taxon is also separate and distinct from an extinct lineage of *C. n. chathamensis* on central San Cristóbal.

***Chelonoidis niger*, subsp. indet.**

Testudo clivosa Garman 1917:283 (*nomen dubium*)

Geochelone clivosa, Chelonoidis clivosa

Type locality: "Mascarenes" by supposition [in error]. Emended to "?Galapagos Is." [Galápagos, Ecuador] by Barbour and Loveridge (1929:350).

Type specimen: MCZ 11075, holotype, see Barbour and Loveridge (1929).

Etymology:* Latin, *clivosa*, steep, hilly, sloping; referring to the arched carapace.

Testudo typica Garman 1917:285 (*nomen dubium*)

Geochelone typica, Chelonoidis typica

Type locality: "Galapagos" [Galápagos, Ecuador].

Type specimen: MCZ 11072, holotype, see Barbour and Loveridge (1929).

Etymology:* Greek, *typos*, image, pattern, type, *typica*, typical; referring to the appearance of this taxon as typical of Galapagos tortoises.

Chersina Gray 1830e (10:7) (25)*

Bowsprit Tortoises

(includes 1 species, 1 taxon)

Chersina Gray 1830e:5 (10:7)

Type species: *Testudo (Chersina) angulata* Duméril *in* Schweigger 1812], by original monotypy.

Etymology:* Greek, *chersos*, tortoise.

Goniochersus Lindholm 1929:285

Type species: *Testudo (Goniochersus) angulata* [= *Testudo angulata* Duméril *in* Schweigger 1812], by original designation.

Etymology:* Greek, *gonia*, angle, and *chersos*, tortoise; referring to the type species *angulata*.

Comment:* Unjustified replacement name for *Chersina*.

Neotestudo Hewitt 1931:504

Type species: *Neotestudo angulata* [= *Testudo angulata* Duméril *in* Schweigger 1812], by original monotypy.

Etymology:* Greek, *neo*, new, and Latin, *testudo*, tortoise.


Comment:* Unjustified replacement name for *Chersina*.

Chersina angulata (Duméril *in* Schweigger 1812) (09:37, 21:88)


Angulate Tortoise, South African Bowsprit Tortoise



Margaretha D. Hofmeyr / CBFTT / West Coast National Park, South Africa



Margaretha D. Hofmeyr / CBFTT / Clanwilliam, South Africa



(orange dots = probable trade; star = type localities)

Distribution: Namibia, South Africa

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 259,049 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 629,956 sq. km

Size (Max SCL): male 35.1 cm, female 21.6 cm (Hofmeyr 2009 CBFTT; Ceballos et al. 2013)

CBFTT Account: Hofmeyr (2009)

IUCN Red List: Least Concern (LC) (Hofmeyr and Keswick 2018). Previously: Least Concern (LC) [Not Listed] (Behler 1996)

CITES: Appendix II, as Testudinidae spp. (1977). Previously: Appendix II, as *Chersina* spp. (1975)

Synonymy:

Testudo angulata Duméril *in* Schweigger 1812:321
Testudo (Chersina) angulata, Chersina angulata, Testudo (Goniochersus) angulata, Goniochersus angulata, Goniochersus angulatus, Neotestudo angulata, Chersine angulata
Type locality: Not known.
Type specimen: MNHN 4087, lectotype, designated by Bour (2008e:32), see also Ceríaco and Bour (2012); MNHN 4067 erroneously referred to as lectotype by Bour (2008e:f.5) (as a typo).
Etymology:* Latin, *angulus*, angle or corner; referring to the angled gular projection.

Testudo bellii Gray 1828:2
Type locality: "Cape of Good Hope" [South Africa] [green star].
Type specimen: NHMUK 2005.1631, holotype, discussed by Bour (2008e), also erroneously referred to as 2005.1641 by Bour (2008e).
Etymology:* Eponym honoring Thomas Bell (1792–1880), United Kingdom; herpetologist, dental surgeon, and Professor of Zoology at King's College, London.

Chersina angulata pallida Gray 1831d:69
Type locality: Not known.
Type specimen: Possibly ZMH, holotype, not located.
Etymology:* Latin, *pallida*, pale; referring to the light background color of the carapace.

Testudo flavofusca Gray 1844:11 (*nomen nudum*)
Etymology:* Latin, *flavus*, yellow, and *fusca*, dark, gray; referring to the colors of the shell.

Testudo sculpta Brandt *in* Gray 1856b:12 (junior homonym and *nomen nudum*, not *Testudo sculpta* Spix 1824 [= *Chelonoidis denticulatus*]).
Type locality: "Cape of Good Hope" [South Africa] [green star].
Type specimen: NHMUK 1850.12.12, see Gray (1873j).
Etymology:* Latin, *sculpta*, carved, sculpted; referring to the clearly pronounced growth rings on the carapace scutes.

Chersobius Fitzinger 1835 (07:67, 10:23, 17:64, 21:89) (25)*

Padloper Tortoises, Padlopers
(includes 3 species, 3 taxa)

Chersobius Fitzinger 1835:112
Type species: *Testudo (Chersobius) signatus* Walbaum [= *Testudo signata* Gmelin 1789], by subsequent designation by Lindholm (1929:284).
Etymology:* Greek, *chersos*, tortoise.

Pseudomopus Hewitt 1931:496
Type species: *Pseudomopus signatus* Walbaum [= *Testudo signata* Gmelin 1789], by original designation.
Etymology:* Greek, *pseudo*, false, untrue, and Greek, *homopus*, even feet; referring to the difference of this genus from *Homopus*.

Chersobius boulengeri (Duerden 1906) (17:64, 21:89)


Karoo Dwarf Tortoise, Karoo Padloper



Victor J.T. Loehr / Northern Cape Prov., South Africa



Victor J.T. Loehr / Northern Cape Prov., South Africa



(stars = type localities)

Distribution: South Africa

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 111,235 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 163,211 sq. km

Size (Max SCL): male 10.0 cm, female 11.0 cm (Boycott and Bourquin 2000)

IUCN Red List: Endangered (EN A4ace) (Hofmeyr, Loehr, Baard, and Juvik 2018). Previously: Least Concern (LC) [Not Listed] (Behler 1996), as *Homopus boulengeri*

CITES: Appendix II, as Testudinidae spp. (1977), as *Homopus boulengeri*. Previously: Appendix II (1975)

Synonymy:


Homopus boulengeri Duerden 1906:406
Pseudomopus boulengeri, Chersobius boulengeri
Type locality: "South Africa—Districts of Willowmore, Aberdeen, and Beaufort West" [yellow stars].

Type specimen: NHMUK 1946.1.23.4 (formerly1906.6.21.1), holotype.
Etymology:* Eponym honoring George Albert Boulenger (1858–1937), Belgium / United Kingdom; herpetologist and curator at the British Museum of Natural History, London.

Chersobius signatus (Gmelin 1789) (10:24, 17:64, 21:89)
Speckled Dwarf Tortoise, Speckled Tortoise, Speckled Padloper


Victor J.T. Loehr / Northern Cape Prov., South Africa


Victor J.T. Loehr / Northern Cape Prov., South Africa


(stars = type localities)

Distribution: South Africa
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 39,014 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 75,392 sq. km
Size (Max SCL): male 9.6 cm, female 11.0 cm (Loehr et al. 2006; Ceballos et al. 2013)
IUCN Red List: Endangered (EN A4ace) (Hofmeyr, Loehr, and Baard 2018). Previously: Near Threatened (NT) (Branch *in* Baillie and Groombridge 1996)
CITES: Appendix II, as Testudinidae spp. (1977), as *Homopus signatus*. Previously: Appendix II (1975)
Synonymy:
Testudo signata Walbaum 1782:120 (unavailable name)
Type locality: Not known.
Type specimen: Not located, type specimen figured (f.1-2.).
Etymology:* Latin, *signum*, the sign or feature; referring to the features that appeared to differentiate the examined specimens from other species.
Comment: Unavailable name from a non-binomial work, see Wermuth (1956).
Testudo signata Gmelin 1789:1043
Chersine signata, Homopus signatus, Pseudomopus signatus, Pseudomopus signatus signatus, Chersobius signatus, Homopus signata, Homopus signatus signatus
Type locality: "Virginia" [USA, in error]. Restricted to "vicinity of Springbok, Cape Province, South Africa" [yellow star] by Bour (1988:3).
Type specimens: Possibly NRM s/n (formerly MDG s/n), apparently lost, not listed by Andersson (1900), specimen figured in Walbaum (1782:f.1-2) designated lectotype by Bour (1988:1).
Etymology:* Latin, *signum*, the sign or feature; referring to the features that appeared to differentiate the examined specimens from other species.
Comment: Description cited as sourced from Walbaum (1782:120).
Testudo cafra Daudin 1801:291 (10:24)
Homopus signatus cafra, Homopus signatus cafer, Homopus cafer
Type locality: "Afrique...la Cafrerie" [South Africa]. Restricted to "Drainage of the Olifants River, Cape Province, South Africa" [green star, location estimated] by Bour (1988:3).
Type specimen: MNHN 7924, holotype, see Bour (1988).
Etymology:* Toponym for Cafrerie (= Kaffraria), an historical region of South Africa.
Testudo juvencella Daudin 1802:380 (10:25)
Type locality: "Afrique" [Africa].
Type specimen: Possibly MNHN, not located.
Etymology:* Latin, *juven*, young, and *-ella*, small or smaller; referring to this taxon being smaller than *Testudo cafra* Daudin 1801.
Pseudomopus signatus peersi Hewitt 1935:345
Homopus signatus peersi, Chersobius peersi
Type locality: "Klaver District, C.P. [Cape Province], near Van Rhynsdorp" [South Africa] [blue star].
Type specimens: Possibly PEM or AMG, syntypes (2), figured (pl.36).
Etymology:* Eponym honoring Bertram Peers (1903–1939), South Africa; amateur archeologist and naturalist who collected the species.

Chersobius solus (Branch 2007) (07:67, 17:64, 21:89)
Nama Tortoise, Nama Padloper




Maurice Rodrigues / TCC / Namibia





left: Alfred Schleicher / Namibia / male
right: Uwe Fritz / Namibia



(star = type locality)

Distribution: Namibia

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 22,377 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 30,614 sq. km

Size (Max SCL): male 9.6 cm, female 11.4 cm (Schleicher 2004; Branch 2007)

IUCN Red List: Vulnerable (VU C2a) (Branch 2018). Previously: Vulnerable (VU) (Branch *in* Baillie and Groombridge 1996), as *Homopus bergeri*

TFTSG Provisional Red List: Endangered (EN) (2013)

CITES: Appendix II, as Testudinidae spp. (1977), as *Homopus solus*. Previously: Appendix II (1975), as *Homopus bergeri*

Synonymy:

Homopus solos Devaux 2003:40 (*nomen nudum*)
Etymology:* Latin, *solus*, alone, unique, lonely, and *sol*, the sun; referring to the isolated range of this taxon and in allusion to the sun in the Namib Desert.

Homopus solus Branch 2007:11
Chersobius solus
Type locality: "vicinity of Aus, Luderitz District, Namibia" [yellow star].
Type specimen: PEM 8754, holotype.
Etymology:* Latin, *solus*, alone, unique, lonely, and *sol*, the sun; referring to the isolated range of this taxon and in allusion to the sun in the Namib Desert.

Cylindraspis Fitzinger 1835 (17:55, 21:90) (75)*
Mascarene Giant Tortoises
(Extinct, ca. 1840)
(includes 5 species, 5 taxa)

Chelonura Rafinesque 1815:74 (*nomen nudum*)
Etymology:* Greek, *chelone*, turtle, and *oura*, tail.


Chelonura Rafinesque 1832:64 (junior homonym, not = *Chelonura* Fleming 1822 [= *Chelydra*])
Type species: *Chelonura indica* [= *Testudo indica* Schneider 1783], by original designation.
Etymology:* Greek, *chelone*, turtle, and *oura*, tail.

Cylindraspis Fitzinger 1835:112
Type species: *Chersina* (*Cylindraspis) vosmaeri* Fitzinger 1826 [= *Testudo indica vosmaeri* Suckow 1798], by subsequent designation by Fitzinger (1843:29). Genus established as *Geochelone (Cylindraspis)* without a type species.
Etymology:* Latin, *kylindros*, roller, cylinder, and Greek, *aspis*, shield; referring to the cylindrical shell.

Cylindraspis indica (Schneider 1783) (07:65)
Reunion Giant Tortoise
(Extinct, ca. 1840)



Roger Bour / Réunion [MNHN, Paris]



(star = type locality)

Distribution: Réunion [*extinct*]

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 2,536 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 2,700 sq. km

Size (Max SCL): estimated: male ca. 117 cm, female ca. 50 cm (Bour 1985; Bour et al. 2014); measured: male 75.0 cm, female 36.5 cm [*C. borbonica*] (Bour 1985)

IUCN Red List: Extinct (EX) (WCMC 1996)

Synonymy:

Testudo indica Schneider 1783:355
Chelonura indica, Cylindrapis indica, Megalochelys indica, Geochelone indica, Cylindraspis indica
Type locality: "Ostindien". Restricted to "Côte de Coromandel" [Coromandel Coast, India] [no star] by Bour (1985:35) as per Perrault (1676:193), and to "Réunion" [red star] by Austin and Arnold (2001:2517).
Type specimen: MNHN 7819, holotype, discussed by Bour (1985, 2004c) and Gerlach (2004); genotyped by Austin and Arnold (2001) and Austin et al. (2002).
Etymology:* Toponym for India.

Testudo tabulata africana Schweigger 1812:322 (09:38)
Type locality: "Africa" by inference [no star].
Type specimen: MNHN 9374, holotype, see Bour (1985, 2008) and Ceríaco and Bour (2012); genotyped by Austin et al. (2002).
Etymology:* Toponym for Africa.

Chersine retusa Merrem 1820:29 (*nomen novum*)
Testudo retusa
Etymology:* Latin, *retusus*, blunted or rounded; referring to the truncated anterior plastron.
Comment: Unjustified replacement name for *indica*.

Testudo perraultii Duméril and Bibron 1835:126 (*nomen novum*)
Geochelone (Cylindraspis) perraultii, Testudo indica perraultii
Type locality: "Indes Orientales" [no star].
Type specimen: MNHN 9374, holotype, see Bour (1985, 2004c, 2008); genotyped by Austin et al. (2002).
Etymology:* Eponym honoring Claude Perrault (1613–1688), France; architect, physician, and anatomist, and the original describer of the species in the pre-Linnaean literature (Perrault 1676).
Comment: Unjustified replacement name for *indica*.


Testudo graii Duméril and Bibron 1835:135 (*nomen novum*)
Geochelone graii, Cylindraspis graii
Type locality: "Afrique?" [Africa] [no star].
Type specimen: MNHN 9374, holotype, see Bour (1985, 2008); genotyped by Austin et al. (2002).
Etymology:* Eponym honoring John Edward Gray (1800–1875), United Kingdom; herpetologist, turtle taxonomist, and curator at the British Museum of Natural History, London.
Comment: Unjustified replacement name for *Testudo tabulata africana* Schweigger 1812.

Chersina grayi Strauch 1865:36 (*nomen novum*)
Geochelone grayi, Geochelone (Cylindraspis) grayi
Type locality: "Afrika" [Africa] [no star]
Type specimen: MNHN 9374, holotype by default, see Bour (1985, 2008); genotyped by Austin et al. (2002).
Etymology:* Eponym honoring John Edward Gray (1800–1875), United Kingdom; herpetologist, turtle taxonomist, and curator at the British Museum of Natural History, London.
Comment: Unjustified emendation of *graii*.

Cylindraspis borbonica Bour 1978:492
Type locality: "Réunion" [red star]
Type specimen: Specimen figured in Petit (1737:pl.7,f.3,5-6), holotype, by original designation, discussed also by Gerlach (2004).
Etymology:* Toponym for Bourbon Island (old name for Réunion).

Cylindraspis inepta (Günther 1873a)
Mauritius Giant Domed Tortoise
(Extinct, ca. 1735**)**


Roger Bour / Mare aux Songes, Mauritius [NHMUK, London]


(star = type locality)

Distribution: Mauritius (*Mauritius [extinct]*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 1,882 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 2,090 sq. km
Size (Max SCL): estimated: male ca. 104 cm (Bour et al. 2014); measured: female 76.0 cm (Bour 1985)
IUCN Red List: Extinct (EX) (WCMC 1996)
Synonymy:


Testudo neraudii Gray 1831d:14 (*nomen oblitum*)
Type locality: "Isle of France" [Mauritius] [red star]. Restricted to "Quatre Cocos, Flacq, Maurice" [Mauritius] [red star] by Bour (1985:37).
Type specimens: MNHN 8383, 8385, syntypes (2).
Etymology:* Eponym honoring Jules Néraud (1795–1855), France; botanist who collected bone fragments of this species on Mauritius.

Testudo inepta Günther 1873a:397
Geochelone inepta, Cylindraspis inepta
Type locality: "Mauritius." Restricted to "La Mare aux Songes...near Mahe'bourg...Mauritius" [red star] by Günther (1875b:297).
Type specimens: NHMUK 39927, 39931, 39937–38, 39943–44, 39963, 1877.11.13.1, syntypes (8), see Bour (1985) and Gerlach (2004).
Etymology:* Latin, *inepta*, awkward, stupid, inept; referring to the type specimens of this taxon being found together with a skeleton of the extinct Dodo (*Didus ineptus*, a synonym of *Raphus cucullatus*).


Testudo boutonii Günther 1875b:297
Type locality: "La Mare aux Songes...near Mahe'bourg...Mauritius" [red star].
Type specimens: NHMUK 39936, 39943, syntypes (2); see Bour (1985).
Etymology:* Eponym honoring Louis S. Bouton (1800–1878), France / Mauritius; botanist and secretary of the Natural History Society of Mauritius.

Testudo sauzieri Gadow 1894:315
Geochelone sauzieri
Type locality: "Mare aux Songes, in Mauritius" [red star].
Type specimen: NHMUK 4676, holotype, see Bour (1985).
Etymology:* Eponym honoring Théodore Sauzier (1829–1904), France / Réunion / Mauritius; naturalist and paleontologist who lived and worked on Réunion and Mauritius.

Cylindraspis peltastes (Duméril and Bibron 1835) [75]*
Rodrigues Domed Tortoise
(Extinct, ca. 1800**)**



Roger Bour / Mauritius [NHMUK, London]



(star = type locality)

Distribution: Mauritius (*Rodrigues [extinct]*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 113 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 135 sq. km

Size (Max SCL): measured: male 46.0 cm, female 41.0 cm (Bour 1985; Bour et al. 2014)

IUCN Red List: Extinct (EX) (WCMC 1996)

Synonymy:

Testudo peltastes Duméril and Bibron 1835:138
Geochelone peltastes, Geochelone (Cylindraspis) peltastes, Cylindraspis peltastes
Type locality: Not known. Restricted to "Rodriguez" [Rodrigues] [red star] by Günther (1877:53).
Type specimen: MNHN 7831, holotype, discussed by Bour (1985) and Gerlach (2004); genotyped by Austin et al. (2002).
Etymology:* Greek, *peltastes*, lightly armed foot soldiers in ancient Greece who wore light shields (*pelte*) made of twigs and leather; referring to the relatively thin shell bones in this taxon.


Cylindraspis triserrata (Günther 1873a)
Mauritius Giant Flat-shelled Tortoise
(Extinct, ca. 1735**)**



Trustees of the Natural History Museum, London / Mauritius [holotype *T. schweigeri*]



Trustees of the Natural History Museum, London / Mauritius [holotype *T. schweigeri*]



(star = type locality)

Distribution: Mauritius (*Mauritius [extinct]*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 1,882 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 2,090 sq. km

Size (Max SCL): estimated: male ca. 75 cm (Bour 1985)

IUCN Red List: Extinct (EX) (WCMC 1996)

Synonymy:

Testudo schweigeri Gray 1830e:3 (*nomen oblitum*) [10:7]
Cylindraspis schweigeri
Type locality: Not known. Restricted to "Mauritius" [red star] by Austin and Arnold (2001:2517).
Type specimen:* NHMUK 1947.3.5.5 (formerly 1876.10.28.4), holotype, figured in Boulenger (1889:156,pl.2-3, as "*Testudo schweiggeri* Gray 1831"), see also Gray (1831d:10), discussed by Bour (1985) and Bour et al. (2014); genotyped by Austin and Arnold (2001) and Kehlmaier et al. (2023).
Etymology:* Eponym honoring Augustus F. Schweigger (1783–1821), Germany; naturalist, herpetologist, and turtle taxonomist.

Testudo schweiggeri Gray *in* Duméril and Bibron 1835:108 (senior homonym and *nomen novum et oblitum*, not = *Geochelone (Geochelone) schweiggeri* Fitzinger 1835 [= *Chelonoidis darwini*])
Etymology:* Eponym honoring Augustus F. Schweigger (1783–1821), Germany; naturalist, herpetologist, and turtle taxonomist.
Comment: Unjustified emendation of *schweigeri*.

Testudo triserrata Günther 1873a:397
Geochelone triserrata, Cylindraspis triserrata
Type locality: "Mauritius." Restricted to "La Mare aux Songes...near Mahe'bourg...Mauritius" [red star] by Günther (1875b:297).
Type specimens: NHMUK 39928–29, 39932–35, 39939–42, 39945–47, 39964, 1947.3.5.5, syntypes (15), see Bour (1985) and Gerlach (2004).
Etymology:* Latin, *tria*, three, and *serrata*, serrated, sawn, jagged; referring to the three serrated horn ridges in the lower jaw.

Testudo leptocnemis Günther 1875b:297
Geochelone leptocnemis, Cylindraspis leptocnemis
Type locality: "district of Flacq..[&]..La Mare aux Songes...near Mahe'bourg...Mauritius." Restricted to "District of Flacq" [Mauritius] [red star] by Günther (1877:14).
Type specimen: NHMUK 1876.11.4.17, lectotype, designated and genotyped by Austin and Arnold (2001).
Etymology:* Greek, *leptos*, narrow, thin, slender, and *knemis*, shin, lower leg; referring to the relatively thin leg bones in this taxon.

Testudo microtympanum Boulenger 1891:4
Geochelone microtympanum
Type locality: "Mauritius" [red star] by supposition.
Type specimen: RCSM s/n, holotype, apparently lost or destroyed during World War II, see Bour (1985) and Bour et al. (2014).
Etymology:* Greek, *mikros*, small, and *tympanon*, tympanic cavity, ear drum; referring to the relatively small tympanic cavity.


Testudo guentheri Gadow 1894:320 (senior homonym, not = *Geochelone elephantopus guentheri* Baur *in* Pritchard [= *Chelonoidis guntheri*])
Type locality: "Mare aux Songes...Mauritius" [red star].
Type specimens: NHMUK 4138–40, MNHN 1943.3, syntypes (4), see Bour (1985).
Etymology:* Eponym honoring Albert C.L.G. Günther (1830–1914), Germany / United Kingdom; zoologist, herpetologist, ichthyologist, and curator at the British Museum of Natural History, London.


Testudo guntheri Gadow *in* Van Denburgh 1914:257 (junior homonym and *nomen novum*, not = *Testudo guntheri* Baur 1889 [= *Chelonoidis guntheri*])
Testudo güntheri
Etymology:* Eponym honoring Albert C.L.G. Günther (1830–1914), Germany / United Kingdom; zoologist, herpetologist, ichthyologist, and curator at the British Museum of Natural History, London.
Comment: Unjustified replacement name for *Testudo guentheri* Gadow.

Testudo gadowi Van Denburgh 1914:257 (*nomen novum*)
Geochelone (Megalochelys) gadowi, Geochelone (Cylindraspis) gadowi
Type locality: "Mare aux Songes...Mauritius" [red star].
Type specimens: NHMUK 4138–40, MNHN 1943.3, syntypes (4) by default, see Bour (1985).
Etymology:* Eponym honoring Hans F. Gadow (1855–1928), Germany / United Kingdom; zoologist, ornithologist, and curator at Cambridge University.
Comment: Justified replacement name for *Testudo guentheri* Gadow.

Cylindraspis vosmaeri (Suckow 1798) (07:66, 21:91) (75)*
Rodrigues Giant Saddleback Tortoise
(Extinct, ca. 1800)


Roger Bour / Rodrigues [Ste. Geneviève]


Roger Bour / Rodrigues


(star = type locality)

Distribution: Mauritius (*Rodrigues [extinct]*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 113 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 135 sq. km
Size (Max SCL): estimated: male ca. 109 cm, female ca. 80 cm (Taffortet 1726; Milne-Edwards 1875; Bour 1985; Bour et al. 2014); measured: male 88.0 cm, female 68.0 cm (Bour 1985; Bour et al. 2014)
IUCN Red List: Extinct (EX) (WCMC 1996)
Synonymy:
Testudo indica. Vosmaeri Schoepff 1795:103 (unavailable name) (21:91)
Type locality: "Promontorio Bonae Spei" [Cape of Good Hope, South Africa] [in error] [no star].
Type specimen: RMNH 6001, type specimen figured (pl.22.B,f.[2-3]), see Hoogmoed et al. (2010) and Bour et al. (2014).
Etymology:* Eponym honoring Arnout Vosmaer (1720–1799), Netherlands; naturalist and curator of a private natural history collection.
Comment: Name written to mean "*Testudo indica* sensu Vosmaer."

Testudo indica vosmaeri Suckow 1798:57

Testudo vosmaeri, Geochelone (Cylindraspis) vosmaeri, Chersina (Cylindraspis) vosmaeri, Geochelone vosmaeri, Cylindraspis vosmaeri

Type locality: Not given. Restricted to "Rodriguez" [Rodrigues] [red star] by Günther (1877:53); and to "Cap de Bonne-Espérance" [Cape of Good Hope, South Africa] [in error] by Vaillant (1893:269).

Type specimen: RMNH 6001, lectotype, designated by Bour (1985:40), discussed by Gerlach (2004) and Hoogmoed et al. (2010); genotyped by Austin et al. (2002); NMW 1461, listed by Tiedemann and Häupl (1980), Tiedemann et al. (1994), and Gemel et al. (2019) as a "syntype" of "*Testudo vosmaeri* Fitzinger 1826," is a paralectotype, genotyped by Kehlmaier et al. (2019b).

Etymology:* Eponym honoring Arnout Vosmaer (1720–1799), Netherlands; naturalist and curator of a private natural history collection.

Testudo rotunda Latreille *in* Sonnini and Latreille 1801:107 (75)*

Chersine rotunda, Geochelone (Geochelone) rotunda

Type locality: Not known. Restricted to "Mascarenes...Rodrigues" [red star] by Bour (2005e:25).

Type specimens:* MNHN 1991, 1991A, syntypes (2), see Pritchard (1996), Gerlach (2004), and Bour (2005e), MNHN 1991 genotyped by Kehlmaier et al. (2023).

Etymology:* Latin, *rotunda*, round; referring to the rounded carapace in the hatchling type specimens.

Testudo rodericensis Günther 1873a:397

Type locality: "Rodriguez" [Rodrigues] [red star].

Type specimens: NHMUK 1876.11.4.1–9, syntypes (10), see Bour (1985).

Etymology:* Toponym for Rodrigues Island, Mauritius.

Testudo commersoni Vaillant 1898:138

Geochelone commersoni, Cylindraspis commersoni

Type locality: "Rodrigue" [Rodrigues] [red star].

Type specimen: Not located, type specimen figured (f.1-4), see Bour (1985).

Etymology:* Eponym honoring Philibert Commerson (1727–1773), France; botanist and explorer who studied the type.

Geochelone Fitzinger 1835 (07:52)

Asian Star Tortoises

(includes 2 species, 3 taxa)

Geochelone Fitzinger 1835:112


Type species: *Geochelone stellata* [= *Testudo stellata* Schweigger 1812 = objective synonym of *Testudo elegans* Schoepff 1795], by subsequent designation by Fitzinger (1843:29).

Etymology:* Greek, *gaia, geo-*, the earth, and *chelone*, turtle.

Geochelone elegans (Schoepff 1795) (17:65, 21:92) (81, 82)*

Indian Star Tortoise

(includes 2 subspecies)



** (subspecies: *elegans* = red, *actinodes* = blue; orange dots = probable trade)

Distribution: India (*Andhra Pradesh, Chhattisgarh, Gujarat, Karnataka, Kerala, Madhya Pradesh, Odesha, Rajasthan, Tamil Nadu, Telangana*), Pakistan, Sri Lanka

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 599,206 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 2,593,830 sq. km

Size (Max SCL): male 25.7 cm, female 46.7 cm (see subspp.)

CBFTT Account: D'Cruze, Mookerjee, Vyas, Macdonald, and de Silva (2018)

IUCN Red List: Vulnerable (VU A4cd) (Choudhury, de Silva, and Shepherd 2020). Previously: Vulnerable (VU A4cd) (D'Cruze, Choudhury, and Mookerjee 2016); Least Concern (LC) (ATTWG 2000); Least Concern (LC) [Not Listed] (Behler 1996)

CITES: Appendix I (2019). Previously: Appendix II, as Testudinidae spp. (1977); Appendix II, as *Geochelone* spp. (1975)


Geochelone elegans elegans (Schoepff 1795) (17:65, 21:92) (81, 82)*
Southern Indian Star Tortoise



Peter Paul van Dijk / Tamil Nadu, India



Brian D. Horne / captivity, Madras Crocodile Bank Trust, Tamil Nadu, India



** (subspecies: *elegans* = red, *actinodes* = blue; orange dots = probable trade; stars = type localities)

Distribution:* India (*Andhra Pradesh, Chhattisgarh, Karnataka, Kerala, Odesha, Tamil Nadu, Telangana*), Sri Lanka

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 321,011 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 1,183,260 sq. km

Size (Max SCL):* India: male 25.7 cm, female 32.2 cm (Frazier *in* Moll 1989); Sri Lanka: female 46.7 cm [estimated] (de Silva et al. 2017); **(Max CCL):** female 57.5 cm (de Silva et al. 2017; D'Cruze et al. 2018 CBFTT)

CBFTT Account: D'Cruze, Mookerjee, Vyas, Macdonald, and de Silva (2018)

CITES: Appendix I (2019). Previously: Appendix II, as Testudinidae spp. (1977); Appendix II, as *Geochelone* spp. (1975)

Synonymy:

Testudo elegans Schoepff 1795:111 (17:65)
Chersine elegans, Peltastes stellatus elegans, Geochelone elegans, Geochelone elegans elegans
Type locality: "India orientali" [eastern India] [yellow star; location estimated; probably in the vicinity of Pulicat, former Dutch Coromandel governorate, Tamil Nadu].
Type specimen: Not located, possibly lost, type specimen figured (pl.25); however, Schoepff (1801:113) stated that the figured specimen was from Amsterdam and others were in the "Museis Hagae Comitis et Harlemi" (possibly the Teylers Museum in Haarlem, Netherlands).
Etymology:* Latin, *elegans*, elegant, fine, select; referring to the elegant radiating markings on the shell.

Testudo stellata Schweigger 1812:325 (*nomen novum*) (17:65) (82)*
Geochelone (Geochelone) stellata, Geochelone stellata, Peltastes stellatus, Geochelone elegans stellata
Type locality: "India orientali" [eastern India] [yellow star; location estimated; probably in the vicinity of Pulicat, former Dutch Coromandel governorate, Tamil Nadu].
Type specimen: Not located, possibly lost, type specimen figured in Schoepff (1795:pl.25); however, Schoepff (1801:113) stated that the figured specimen was from Amsterdam and others were in the "Museis Hagae Comitis et Harlemi" (possibly the Teylers Museum in Haarlem, Netherlands). Schweigger noted that he had seen Schoepff's specimen in the natural history museum in Berlin.
Etymology:* Latin, *stella*, star, and *-ata*, having; referring to the star-like radiating pattern on the shell.
Comment: Unjustified replacement name for *elegans* (see Annotation 17:65).

Peltastes stellatus maura Gray 1870b:654 (81)*
Testudo stellata maura
Type locality: Not known. Restricted to "India" by Gray (1873j:8).
Type specimen: NHMUK, holotype, see Gray (1873j), possibly lost.
Etymology:* Greek, *mavra*, black; referring to the predominantly black carapace.

Peltastes stellatus sebae Gray 1870b:654 (81)*
Peltastes stellatus seba
Type locality: Not known. Restricted to "India" by Gray (1873j:8).
Type specimen: NHMUK 1843.2.29.6, holotype, see Gray (1873j), not located, possibly lost.
Etymology:* Eponym honoring Albertus Seba (1665–1736), Netherlands; pharmacist and naturalist with a large natural history collection.

Testudo megalopus Blyth 1854:640 (17:66)
Type locality: Not known. Restricted to "Calcutta (in West Bengal State, eastern India)" [Kolkata, West Bengal, India] [orange star; trade specimen] by Das et al. (1998:127).
Type specimen: ZSI 792, holotype, see Das et al. (1998), not located by Kundu et al. (2018a).
Etymology:* Greek, *megalos*, large, long, and *pus*, foot; referring to the size of the feet in this taxon.

Geochelone elegans actinodes (Bell 1828a) (17:65, 21:92) (82)*
Northwestern Indian Star Tortoise




Raju Vyas / CBFTT / Gir Wildlife Sanctuary and National Park, Gujarat, India





Neil D'Cruze / Gujarat, India



** (subspecies: *elegans* = red, *actinodes* = blue; orange dots = probable trade)

Distribution:* India (*Gujarat, Madhya Pradesh, Rajasthan*), Pakistan

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 277,195 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 342,292 sq. km

Size (Max SCL):* female 41.0 cm (Vyas 2011); **(Max CCL):** female 50.5 cm (Vyas 2011)

CBFTT Account: D'Cruze, Mookerjee, Vyas, Macdonald, and de Silva (2018)

CITES: Appendix I (2019). Previously: Appendix II, as Testudinidae spp. (1977); Appendix II, as *Geochelone* spp. (1975)

Synonymy:*

Testudo actinodes Bell 1828a:419 (82)*
Type locality: "Africâ" [in error] [no star].
Type specimens: OUM 8555–56, 8558–60, syntypes (5), discussed by Nowak-Kemp and Fritz (2010).
Etymology:* Greek, *aktinos*, a ray or beam, and *-oides*, resembling; referring to the ray-like carapacial markings.

Testudo actinoides Bell *in* Gray 1844:7 (*nomen novum*)
Peltastes stellatus actinoides
Etymology:* Greek, *aktinos*, a ray or beam, and *-oides*, resembling; referring to the ray-like carapacial markings.
Comment: Unjustified emendation for *actinodes*.

Geochelone platynota (Blyth 1863)
Burmese Star Tortoise, Myanmar Star Tortoise




Kalyar Platt / CBFTT / Lawkanandar Wildlife Sanctury, Bagan, Myanmar





Kalyar Platt / CBFTT / Lawkanandar Wildlife Sanctury, Bagan, Myanmar



(orange dots = trade; star = type locality)

Distribution: Myanmar

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 85,181 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 100,164 sq. km

Size (Max SCL): male 25.0 cm, female 45.5 cm (Platt et al. 2011 CBFTT; Platt et al. 2019a)

CBFTT Account: Platt, Thanda Swe, Win Ko Ko, Platt, Khin Myo Myo, Rainwater, and Emmett (2011)

IUCN Red List: Critically Endangered (CR A1cd) (Praschag, Platt, and Horne 2020). Previously: Critically Endangered (CR) (ATTWG 2000); Critically Endangered (CR) (TFTSG *in* Baillie and Groombridge 1996)

IUCN Green List:* Critically Depleted (CD) (Horne, Platt, and Bennett 2024)

CITES: Appendix I (2013). Previously: Appendix II, as Testudinidae spp. (1977); Appendix II, as *Geochelone* spp. (1975)

Synonymy:

Testudo platynotus Blyth 1863:83
Peltastes platynotus, Testudo platynota, Geochelone platynota, Geochelone elegans platynota
Type locality: "Lower Pegu..[&]..Valley of the Irawadi" [Myanmar]. Restricted to "Irrawaddy Valley" [Myanmar] [yellow star, location estimated] by Smith (1931:140).
Type specimens: ZSI 787–89, syntypes (3), see Das et al. (1998) and Kundu et al. (2018a); Uetz et al. (2019) erroneously listed syntypes as not located.
Etymology:* Greek, *platys*, flat, even, broad, wide, and *notos*, the back; referring to the slightly flattened carapace as compared to *G. elegans*.

Homopus Duméril and Bibron 1834 (07:67, 10:23, 17:64, 21:89) (25)*

Cape Tortoises

(includes 2 species, 2 taxa)

Homopus Duméril and Bibron 1834:357 (10:23

Type species: *Homopus areolatus* [*Tortue Aréolée* Schoepff = *Testudo areolata* Thunberg 1787], by subsequent designation by Duméril and Bibron (1835:7).

Etymology:* Greek, *homos*, equal, similar, corresponding, agreeing, and *pus*, foot; referring to the front and rear feet all having four claws each.

Homopus areolatus (Thunberg 1787) (17:64, 21:89, 21:93)

Parrot-beaked Dwarf Tortoise, Parrot-beaked Tortoise, Common Padloper




Gerald Kuchling / Elandsberg, South Africa





left: Gerald Kuchling / Elandsberg, South Africa
right: Craig B. Stanford / South Africa



(star = type locality)

Distribution: South Africa

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO): 85,707 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 240,965 sq. km

Size (Max SCL): male 10.0 cm, female 12.0 cm (Boycott and Bourquin 2000; Ceballos et al. 2013)

IUCN Red List: Least Concern (LC) (Hofmeyr and Keswick 2018). Previously: Least Concern (LC) [Not Listed] (Behler 1996)

CITES: Appendix II, as Testudinidae spp. (1977). Previously: Appendix II (1975)

Synonymy:

Testudo areolata Thunberg 1787:180

Chersine areolata, Homopus areolatus, Homopus areolata

Type locality: "Indien" [India] [in error, no star].

Type specimen: UPSZTY 298 (formerly UUZM 298 and ZIUS 298), holotype, see Thunberg (1828) and Iverson (1992).

Etymology:* Latin, *areola*, small space, small field; referring to the defined areas in the middle of the carapacial scutes framed by raised growth rings.

Testudo minuta Thunberg 1788:206 (*nomen nudum*)

Etymology:* Latin, *minuta*, very small, tiny; referring to the very small body size.

Testudo miniata Lacepède 1788:166, synopsis[table] (*nomen suppressum*) (09:6)

Type locality: "Cap de Bonne-espérance" [South Africa] [green star].

Type specimen: Possibly MNHN, not located.

Etymology:* Latin, *miniata*, vermilion; referring to the orange-red prefrontal scales on top of the head of males during the breeding season.

Comment: Name suppressed by ICZN (2005a) as published in a rejected and invalid non-binomial work, see Savage (2003).

Testudo fasciata Daudin 1801:294 (junior homonym, not = *Testudo fasciata* Suckow 1798 [= *Trachemys terrapen*])

Chersine fasciata

Type locality: "l'île de Ceilan" [Sri Lanka] [in error].

Type specimen: Possibly MNHN, not located.

Etymology:* Latin, *fascia*, band, ribbon, bundle; referring to the stripes on the carapace.

Testudo africana Hermann 1804:218

Type locality: "Africa" by inference.

Type specimen: Possibly ZMUS, not located.

Etymology:* Toponym for Africa.

Chersine tetradactyla Merrem 1820:32

Type locality: "India orientali" [in error].

Type specimen: Not known or located.

Etymology:* Greek, *tetra*, four, and *daktylos*, toe, finger; referring to the four claws on all front and hind feet.

Comment: Description cited as sourced from Linnaeus (1745:[33],n.23 and 1749a:139,n.23["354"]) and questionably Worm (1655:317).

Testudo areolata pallida Gray 1831d:13

Type locality: Not designated. Restricted to "Süd-Afrika" [South Africa] [no star] by Wermuth and Mertens (1961:175).

Type specimen: Possibly in OUM, originally live, not located, not listed by Nowak-Kemp and Fritz (2010).

Etymology:* Latin, *pallidus*, pale; referring to the light background color of the carapace.


Homopus femoralis Boulenger 1888a (17:64, 21:89)
Greater Padloper, Greater Dwarf Tortoise



Victor J.T. Loehr / Karoo, Northern Cape Prov., South Africa



Victor J.T. Loehr / Karoo, Northern Cape Prov., South Africa



(star = type locality)

Distribution: South Africa

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 212,535 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 337,901 sq. km

Size (Max SCL): male 13.7 cm, female 16.8 cm (Boycott and Bourquin 2000)

IUCN Red List: Least Concern (LC) (Hofmeyr and Keswick 2018). Previously: Least Concern (LC) [Not Listed] (Behler 1996)

CITES: Appendix II, as Testudinidae spp. (1977). Previously: Appendix II (1975)

Synonymy:

Homopus femoralis Boulenger 1888a:251
Testudo femoralis
Type locality: "South Africa...Cradock" [yellow star].
Type specimens: NHMUK 1946.1.22.54 (formerly 1888.12.28.1), syntype, two others not located.
Etymology:* Latin, *femur*, thigh, *femoralis*, belonging to the thigh; referring to the large conical tubercle on each thigh.


Indotestudo Lindholm 1929

Asian Elongated Tortoises
(includes 3 species, 3 taxa)
Indotestudo Lindholm 1929:285
Type species: *Testudo (Indotestudo) elongata* [= *Testudo elongata* Blyth 1854], by original designation.
Etymology:* Toponym for India, and Latin, *testudo*, turtle.


Indotestudo elongata (Blyth 1854) (17:66) (83)*
Elongated Tortoise, Yellow-headed Tortoise



Flora Ihlow / CBFTT / Doi Phu Nang National Park, Phayao Prov., Thailand



Flora Ihlow / CBFTT / Cuc Phuong National Park, Vietnam / male



(orange dots = probable trade or introduced or temples; stars = type localities)

Distribution: Bangladesh, Bhutan, Cambodia, China (?) (*Guangxi [?], Yunnan [?]*), India (*Assam, Bihar, Jharkhand, Meghalaya, Mizoram, Odisha, Sikkim, Tripura, Uttarakhand, Uttar Pradesh, West Bengal*), Laos, Malaysia (*West*), Myanmar, Nepal, Thailand, Vietnam

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 1,308,716 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 4,802,341 sq. km

Size (Max SCL): male 38.0 cm, female 31.0 cm (Senneke 2006; Ceballos et al. 2013; Ihlow et al. 2016 CBFTT; Khin Myo Myo and Platt 2016)

CBFTT Account: Ihlow, Dawson, Hartmann, and Som (2016)

IUCN Red List: Critically Endangered (CR A2cd) (Rahman, Platt, Das, Choudhury, Ahmed, Cota, McCormack, Timmins, and Singh 2019). Previously: Endangered (EN) (ATTWG 2000); Vulnerable (VU) (TFTSG *in* Baillie and Groombridge 1996)

CITES: Appendix II, as Testudinidae spp. (1977). Previously: Appendix II (1975)

Synonymy:

Testudo elongata Blyth 1854:639 (17:66)

Peltastes elongatus, Indotestudo elongata, Geochelone elongata, Indotestudo elongata elongata, Geochelone elongata elongata

Type locality: "Arakan" [Myanmar] [yellow star, location estimated].

Type specimens: ZSI 796, 798–800, syntypes (4), see Das et al. (1998), Das (2009), and Kundu et al. (2018a).

Etymology:* Latin, *elongata*, stretched; referring to the elongated carapace.

Testudo parallelus Annandale 1913:76

Type locality: "Chaibassa (Singhbhum) district, Chota Nagpur" [India] [green star].

Type specimen: ZSI 11379, holotype, see Das et al. (1998).

Etymology:* Latin, *parallelus*, parallel; referring to the approximately parallel sides of the carapace.

Indotestudo forstenii (Schlegel and Müller 1845) (17:69)


Sulawesi Tortoise, Forsten's Tortoise



Jérôme Maran / Sulawesi, Indonesia



left: Sabine Schoppe / Sulawesi, Indonesia
right: Jérôme Maran / Sulawesi, Indonesia



(stars = type localities)

Distribution: Indonesia (*Sulawesi*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 43,814 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 116,439 sq. km

Size (Max SCL): male 30.9 cm, female 25.4 cm (Ernst et al. 2006b; Ives et al. 2008; Itescu et al. 2014)

IUCN Red List: Critically Endangered (CR A4cd) (Kusrini, As-singkily, Light, and Stanford 2021). Previously: Endangered (EN) (ATTWG 2000); Vulnerable (VU) (TFTSG *in* Baillie and Groombridge 1996)

CITES: Appendix II, as Testudinidae spp. (1977). Previously: Appendix II (1975)

Synonymy:

Testudo forstenii Schlegel and Müller 1845:30 (17:69)

Peltastes forstenii, Geochelone forstenii, Indotestudo forstenii, Indotestudo elongata forstenii, Geochelone elongata forstenii

Type locality: "Gilolo...Indischen Archipel" [Halmahera, Moluccas, Indonesia] [orange star, probably in error]; restricted to "near Gorontalo, Sulawesi, Indonesia" [yellow star] by TTWG (2017:230).

Type specimen: RMNH 3811, holotype, identified and discussed by Hoogmoed and Crumly (1984) and Hoogmoed et al. (2010); Uetz et al. (2019) erroneously listed this specimen as a lectotype.

Etymology:* Eponym honoring Eltio A. Forsten (1811–1843), Netherlands / Indonesia; naturalist and botanist who explored and collected in the Dutch East Indies (now Indonesia).


Indotestudo travancorica (Boulenger 1907) (07:68)
Travancore Tortoise



Veerappan Deepak / CBFTT / Anaimalai Hills, Western Ghats, Tamil Nadu, India



Veerappan Deepak / CBFTT / Anaimalai Hills, Western Ghats, Tamil Nadu, India



**

(star = type locality)

Distribution: India (*Karnataka, Kerala, Tamil Nadu*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 41,526 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 92,013 sq. km

Size (Max SCL): male 33.1 cm, female 29.5 cm (Deepak et al. 2011 CBFTT; Ceballos et al. 2013)

CBFTT Account: Deepak, Ramesh, Bhupathy, and Vasudevan (2011)

IUCN Red List: Vulnerable (VU A1cd [ver. 2.3]) (ATTWG 2000)

TFTSG Provisional Red List: Endangered (EN) (2011)

CITES: Appendix II, as Testudinidae spp. (1977). Previously: Appendix II (1975)

Synonymy:

Testudo travancorica Boulenger 1907:560
Geochelone travancorica, Indotestudo travancorica, Indotestudo elongata travancorica, Geochelone elongata travancorica
Type locality: "near Trivandrum..[&]..Travancore hills between 500 and 1,000 feet altitude" [India] [yellow star].
Type specimens: NHMUK 1946.1.22.80–81 (formerly 1906.7.18.6–7), and 1947.3.5.4 (formerly 1933.11.11.1), syntypes (3), see Das (2009).
Etymology:* Toponym for Travancore (former kingdom now part of Kerala), India.

Kinixys Bell 1827 (12:35) (25)*

Hinge-back Tortoises
(includes 8 species, 9 taxa)

Kinixys Bell 1827:398
Type species: *Kinixys homeana* Bell 1827, by subsequent designation by Bell (1828c:514).
Etymology:* Greek, *kino*, move, motion, and *ixys*, hip, waist; referring to the movable hinge in the posterior part of the carapace.

Kinyxis Gray 1830e:6 (*nomen novum*)
Etymology:* Greek, *kino*, move, motion, and *ixys*, hip, waist; referring to the movable hinge in the posterior part of the carapace.
Comment:* Unjustified emendation of *Kinixys*.

Cinixys Wagler 1830b:138 (*nomen novum*)
Etymology:* Greek, *kino*, move, motion, and *ixys*, hip, waist; referring to the movable hinge in the posterior part of the carapace.
Comment:* Unjustified emendation of *Kinixys*.

Cinothorax Fitzinger 1835:108
Type species: *Cinixys (Cinothorax) bellianus* [= *Testudo (Kinyxis) belliana* Gray 1830e], by subsequent designation by Fitzinger (1843:29).
Etymology:* Greek, *kino*, move, motion, and *thorax*, shell, torso; referring to the movable hinge in the posterior part of the carapace.

Kinixis Hallowell 1839:161 (*nomen novum*)
Etymology:* Greek, *kino*, move, motion, and *ixys*, hip, waist; referring to the movable hinge in the posterior part of the carapace.
Comment:* Unjustified emendation of *Kinixys*.

Cinyxis Peters 1866:887 (*nomen novum*)
Etymology:* Greek, *kino*, move, motion, and *ixys*, hip, waist; referring to the movable hinge in the posterior part of the carapace.
Comment:* Unjustified emendation of *Kinixys*.


Kinothorax Gray 1873j:16 (*nomen novum*)
Etymology:* Greek, *kino*, move, motion, and *thorax*, shell, torso; referring to the movable hinge in the posterior part of the carapace.
Comment:* Unjustified emendation of *Cinothorax*.

Madakinixys Vuillemin 1972b:169
Type species: *Madakinixys domerguei* Vuillemin 1972b, by original monotypy.
Etymology:* Toponym for Madagascar, and Greek, *kinixys*; referring to the similarity to the genus *Kinixys*.


Kinixys belliana (Gray 1830e) (07:69, 08:11, 10:7, 12:35)
Bell's Hinge-back Tortoise



Vincenzo Ferri / Entebbe, Uganda



left: Russell A. Mittermeier / northern Serengeti, Tanzania
right: Phil Allman / Queen Elizabeth National Park, Uganda



(orange dots = possible trade or transport or misidentified; stars = type localities)

Distribution: Angola, Congo (DRC), Congo (ROC), Eritrea, Ethiopia, Kenya, Rwanda, Somalia, South Sudan, Sudan, Tanzania, Uganda

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 2,268,595 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 7,015,110 sq. km

Size (Max SCL): 23.0 cm (Mifsud and Stapleton 2014)

IUCN Red List: Least Concern (LC) [Not Listed] (Behler 1996)

TFTSG Provisional Red List: Vulnerable (VU) (2013)

CITES: Appendix II, as Testudinidae spp. (1977). Previously: Appendix II, as *Kinixys* spp. (1975)

Synonymy:

Testudo (Kinyxis) belliana Gray 1830e:6 (10:7)
Kinixys belliana, Cinixys (Cinothorax) belliana, Cinixys belliana, Cinothorax bellianus, Kinixys belliana belliana
Type locality: Not known.
Type specimen: NHMUK 1979.919, holotype, see Bour (2006d).
Etymology:* Eponym honoring Thomas Bell (1792–1880), United Kingdom; herpetologist, dental surgeon, and Professor of Zoology at King's College, London.


Kinixys schoensis Rüppell 1845:226
Kinixys belliana schoensis
Type locality: "Schoa, südlich von Abyssinien" [Shewa, Ethiopia] [green star, location estimated].
Type specimen: SMF 7719, holotype, see Mertens (1967b) and Bour (2006d).
Etymology:* Toponym for Schoa (now Shewa) historical region and former kingdom in Abyssinia (now Ethiopia).

Kinixys belliana mertensi Laurent 1956:27
Type locality: "Dika, Uele...le nord du Congo" [Democratic Republic of Congo (DRC)] [blue star].
Type specimen: MRAC 4417, holotype.
Etymology:* Eponym honoring Robert F.W. Mertens (1894–1975), Russia / Germany; herpetologist and director of the Senckenberg Museum, Frankfurt.


Kinixys erosa (Schweigger 1812) (12:35)
Forest Hinge-back Tortoise, Serrated Hinge-back Tortoise



Tomas Diagne / Oyo State, Nigeria



left: Laurent Chirio / Kinero Malibé, Gabon
right: Tomas Diagne / Oyo State, Nigeria



** (orange dots = probable trade; star = type locality)

Distribution: Angola, Benin, Cameroon, Central African Republic, Congo (DRC), Congo (ROC), Equatorial Guinea, Gabon, Ghana, Guinea, Ivory Coast (Côte d'Ivoire), Liberia, Nigeria, Sierra Leone, South Sudan, Togo, Uganda

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 2,920,544 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 5,563,008 sq. km

Size (Max SCL): male 40.0 cm, female 29.9 cm (Maran 2006c; Luiselli and Diagne 2014 CBFTT)

CBFTT Account: Luiselli and Diagne (2014)

IUCN Red List:* Endangered (EN A2bcd+4bcd) (Luiselli, Segniagbeto, Behangana, McGovern, Diagne, Eniang, and Petrozzi, in press). Previously: TFTSG Provisional Red List: Endangered (EN) (2013); Data Deficient (DD) (TFTSG 1996)

CITES: Appendix II, as Testudinidae spp. (1977). Previously: Appendix II, as *Kinixys* spp. (1975)

Synonymy:

Testudo erosa Schweigger 1812:321
Kinixys erosa, Cinixys erosa, Kinixis erosa, Kinixys belliana erosa

Type locality: "America septentrionali" [North America] [in error] [no star].
Type specimen: Not located, specimen figured in Shaw (1802:f.13), lectotype, designated by Bour (2006d:8).
Etymology:* Latin, *erosa*, eaten, eaten away, eroded; referring to the strongly serrated front and rear carapace edges.

Testudo schoepfii Fitzinger 1826:44 (senior homonym and *nomen nudum*, not *Testudo schoepfii* Rüppell *in* Gray 1873j [= *Centrochelys sulcata*])
Etymology:* Eponym honoring Johann David Schoepff (or Schöpf) (1752–1800), Germany; physician, botanist, herpetologist, and turtle taxonomist.

Kinixys castanea Bell 1827:398
Cinixys (Cinixys) castanea, Cinixys castanea, Kinixis castanea
Type locality: "Africa." [no star]
Type specimens: OUM 8519–21, syntypes (3), discussed by Nowak-Kemp and Fritz (2010). NHMUK 1947.3.5.71 also listed as a syntype in NHMUK catalogue.
Etymology:* Greek, *kastano*, brown; referring to the brown carapace color.

Kinixis denticulata Hallowell 1839:161
Kinixys denticulata
Type locality: "Liberia...banks of the St. Paul and Mesurado rivers" [green star].
Type specimen: Not located, type specimen figured (pl.8-9).
Etymology:* Latin, *denticulus*, small tooth, *denticulata*, serrated, denticulated; referring to serrated carapacial margin.

Kinixys homeana Bell 1827 (12:35)


Home's Hinge-back Tortoise



Tomas Diagne / CBFTT / Cameroon



left: Luca Luiselli / Port Harcourt, Nigeria
right: Pearson McGovern / Ivory Coast (Côte d'Ivoire)



** (orange dots = questionable or trade; star = type locality)

Distribution: Benin, Cameroon, Equatorial Guinea, Ghana, Guinea (?), Ivory Coast (Côte d'Ivoire), Liberia, Nigeria, Togo

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 565,797 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 1,459,772 sq. km

Size (Max SCL): male 25.1 cm, female 25.8 cm (Luiselli and Diagne 2013 CBFTT; Segniagbeto et al. 2015)

CBFTT Account: Luiselli and Diagne (2013)

IUCN Red List: Critically Endangered (CR A2bcd+4bcd) (Luiselli, Agyekumhene, Akani, Allman, Diagne, Eniang, Mifsud, Petrozzi, and Segniagbeto 2021). Previously: Vulnerable (VU A2cd) (Luiselli, Politano, and Lea 2006); Data Deficient (DD) (TFTSG *in* Baillie and Groombridge 1996)

CITES: Appendix II, as Testudinidae spp. (1977). Previously: Appendix II, as *Kinixys* spp. (1975)

Synonymy:


Kinixys homeana Bell 1827:400
Testudo homeana, Cinixys homeana, Cinixys (Cinothorax) homeana, Testudo (Kinyxis) homeana, Kinixys belliana homeana
Type locality: "Africâ occidentali...Sierra Leone". Emended to "Cape Coast, Ashantee" [Ghana] by Boulenger (1889:143) and to "Cape Coast, Ashanti, i.e. Gold Coast" [Ghana] [yellow star, location estimated] by Loveridge and Williams (1957:400).
Type specimens: OUM 8522, NHMUK a,b (2), syntypes (3), discussed by Nowak-Kemp and Fritz (2010). Two unnumbered NHMUK specimens (a and b, listed by Boulenger 1889) identified as syntypes by Bour (pers. comm.).
Etymology:* Eponym honoring Sir Everard Home (1756–1832), United Kingdom; physician, anatomist, and naturalist who presented the types to the British Museum of Natural History.

Kinixys lobatsiana (Power 1927) (12:35)


Lobatse Hinged Tortoise, Lobatse Hinge-back Tortoise



Flora Ihlow / Lapalala Reserve, Limpopo Prov., South Africa



Victor J.T. Loehr / Limpopo Prov., South Africa





(orange dots = possibly introduced or uncertain; star = type locality)

Distribution: Botswana, South Africa
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 114,837 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 147,733 sq. km
Size (Max SCL): male 17.0 cm, female 20.0 cm (Boycott and Bourquin 2000; Ceballos et al. 2013)
IUCN Red List: Vulnerable (VU A4cde) (Hofmeyr and Boycott 2018)
CITES: Appendix II, as Testudinidae spp. (1977). Previously: Appendix II, as *Kinixys* spp. (1975)
Synonymy:
Cinixys lobatsiana Power 1927:410
Kinixys lobatsiana, Kinixys belliana lobatsiana
Type locality: "Lobatsi [Bechuanaland Protectorate]" [Botswana] [yellow star].
Type specimens: NHMUK 1947.3.5.72 (formerly 1931.12.14.1) and MMK 217, 221, and 224, syntypes (4); Uetz et al. (2019) erroneously listed NHMUK 1979.919 as the holotype, but that specimen is the holotype of *Kinixys belliana*.
Etymology:* Toponym for Lobatsi, Bechuanaland (now Botswana).

Kinixys natalensis Hewitt 1935 (12:35)
Natal Hinge-back Tortoise, KwaZulu-Natal Hinge-back Tortoise


James Harvey / Lebombo Mountains, South Africa



James Harvey / Lebombo Mountains, South Africa



(star = type locality)

Distribution: Eswatini (Swaziland), Mozambique, South Africa
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 70,751 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 121,704 sq. km
Size (Max SCL): male 13.0 cm, female 17.2 cm (Bourquin 1992; Boycott and Bourquin 2000; Ceballos et al. 2013)
IUCN Red List: Vulnerable (VU A4c) (Hofmeyr and Boycott 2018). Previously: Near Threatened (NT) (TFTSG 1996)
CITES: Appendix II, as Testudinidae spp. (1977). Previously: Appendix II, as *Kinixys* spp. (1975)
Synonymy:
Kinixys natalensis Hewitt 1935:353
Kinixys belliana natalensis
Type locality: "Jameson Drift, Tugela River, and...Dimane stream, near Jameson Drift" [KwaZulu-Natal] [South Africa]. Restricted to "Dimane stream, near Jameson Drift, Tugela Valley, Natal" [South Africa] [yellow star] by lectotype designation by Broadley (1981b:206).
Type specimens: PEM s/n (formerly AMG s/n), lectotype, designated by Broadley (1981b:206,f.7).
Etymology:* Toponym for Natal (now KwaZulu-Natal), South Africa.

Kinixys nogueyi (Lataste 1886) (08:11, 09:39, 12:35)
Western Hinge-back Tortoise


Tomas Diagne / Dene, Senegal


Pearson McGovern / Senegal


(stars = type localities)

Distribution: Benin, Burkina Faso, Cameroon, Central African Republic, Chad, Congo (DRC), Gambia, Ghana, Guinea, Guinea-Bissau, Ivory Coast (Côte d'Ivoire), Mali, Mauritania (?), Niger, Nigeria, Senegal, Sierra Leone, South Sudan, Togo
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 2,154,510 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 4,194,819 sq. km
Size (Max SCL): male 28.3 cm, female 28.7 cm (L. Luiselli and G.H. Segniagbeto, unpubl. data)
IUCN Red List: Not Evaluated (NE)
TFTSG Provisional Red List: Vulnerable (VU) (2013)
CITES: Appendix II, as Testudinidae spp. (1977). Previously: Appendix II, as *Kinixys* spp. (1975)
Synonymy:
Homopus nogueyi Lataste 1886:286
Cinixys nogueyi, Cinixys belliana nogueyi, Kinixys


nogueyi, *Kinixys belliana nogueyi*
Type locality: "Médine (Haut-Sénégal)" [= Médine, Mali] [yellow star].
Type specimens: NHMUK 1946.1.22.46, 1947.3.5.75 (formerly 1920.1.20.3240–1, syntypes (2).
Etymology:* Eponym honoring Gustave Noguey (1800s), France; naturalist of the Société Linnéenne de Bordeaux and a traveling companion to the author.

Cinixys dorri Lataste 1888:164
Type locality: "Haut-Sénégal" [Mali]. Restricted to "Bakel, Haut-Sénégal" [= Bakel, Senegal] [green star] by Mertens and Wermuth (1955:372).
Type specimens: NHMUK 1946.1.22.94 (formerly 1920.1.20.3709), syntype, plus at least two others, not located.
Etymology:* Eponym honoring Emile Dorr (1857–1907), France / Senegal; naval officer and naturalist stationed in Senegal who collected the types.

Kinixys spekii Gray 1863d (12:35)
Speke's Hinge-back Tortoise, Speke's Hinged Tortoise


Richard C. Boycott / Dinedor Farm, Eswatini (Swaziland)


left: Flora Ihlow / Big Bend region, Eswatini (Swaziland)
right: Victor J.T. Loehr / nr. Lephalale, Limpopo Prov., South Africa


**
(orange dots = possible trade or introduced or misidentified; stars = type localities)

Distribution: Angola, Botswana, Burundi, Congo (DRC), Eswatini (Swaziland), Kenya, Malawi, Mozambique, Namibia, Rwanda, South Africa, Tanzania, Zambia, Zimbabwe
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 2,522,956 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 5,561,920 sq. km
Size (Max SCL): male 18.1 cm, female 21.0 cm (Boycott and Bourquin 2000; Boycott 2001)
CBFTT Account:* Hailey, Broadley, Boycott, and Spawls (2021)
IUCN Red List: Not Evaluated (NE)
TFTSG Provisional Red List: Vulnerable (VU) (2013)
CITES: Appendix II, as Testudinidae spp. (1977). Previously: Appendix II, as *Kinixys* spp. (1975)
Synonymy:

Kinixys spekii Gray 1863d:381
Cinixys spekii, *Kinixys belliana spekii*
Type locality: "Central Africa". Restricted to "Tanganyika Territory east of the lakes" [Tanzania] and estimated as being near "Simbo a few miles north of Tabora (Unyanyembe), Unyamwezi, Tanganyika Territory" [Tanzania] [yellow star] by Loveridge (1936:218).
Type specimen: NHMUK 1947.3.5.74 (formerly 1863.8.11.8), holotype; Uetz et al. (2019) erroneously listed NHMUK 1936.5.3.117 as holotype.
Etymology:* Eponym honoring John H. Speke (1827–1864), United Kingdom; explorer and first European to reach Lake Victoria, the source of the Nile River, and who collected the type.

Homopus darlingi Boulenger 1902b:15
Kinixys darlingi, *Kinixys belliana darlingi*
Type locality: "Mashonaland...Rhodesia...district about Salisbury...at Mazoë and between Umtali and Marandellas" [Zimbabwe] [green star]. Emended to "Mashonaland" [Zimbabwe] by Mertens and Wermuth (1955:372) and to "Salisbury District, Mashonaland, Southern Rhodesia" [Zimbabwe] by Fritz and Havaš (2007:286).
Type specimen: NHMUK 1946.1.22.60 (formerly 1902.2.12.1), holotype.
Etymology:* Eponym honoring James J.F. Darling (1859–1929), Ireland; gold prospector, medical officer, and naturalist who collected the type.

Testudo procterae Loveridge 1923:928
Malacochersus procterae
Type locality: "Ikikuyu, Dodoma Dist., Tanganyika Territory" [Tanzania] [Ikuyu, Tanzania] [blue star]. Emended to "Ikikuyu, south of Gulwe, Tanganyika Territory" [Tanzania] by Fritz and Havaš (2007:286).
Type specimen: NHMUK 1946.1.22.59 (formerly 1923.10.9.102), holotype.
Etymology:* Eponym honoring Joan B. Procter (1897–1931), United Kingdom; female zoologist, herpetologist, and curator at the British Museum of Natural History, London.

Kinixys australis Hewitt 1931:477
Kinixys australis australis, *Kinixys belliana australis*
Type locality: "White River, Eastern Transvaal" [South Africa] [pink star].
Type specimen: AMG 78C, lectotype, designated by Broadley (1981b:212).
Etymology:* Latin, *auster*, the south, *australis*, from the south; referring to the provenance of the taxon from the south.

Kinixys jordani Hewitt 1931:482
Type locality: "Isoka, N. Rhodesia" [Zambia] [brown star].
Type specimens: Apparently AMG s/n, syntypes (2), specimens figured (pl.37,f.7-9).
Etymology:* Eponym honoring E. Knowles Jordan (1800s–1900s), United Kingdom / Zambia; political official and naturalist in Northern Rhodesia (now Zambia) who collected the types.

Kinixys youngi Hewitt 1931:486
Type locality: "Nyasaland...near Livingstonia, on the shore of the lake" [Malawi] [light gray star].
Type specimens: Apparently AMG s/n, syntypes (2), specimens figured (pl.37,f.10-11).
Etymology:* Eponym honoring Reverend William Paulin Young (1886–1969), United Kingdom / South Africa; clergyman and naturalist who presented the types.

Kinixys australis mababiensis FitzSimons 1932:37
Kinixys belliana mababiensis
Type locality: "Tsotsoroga Pan, Mababe Flats...Kalahari" [Botswana] [dark gray star].

Type specimen: TMP 14689, holotype.
Etymology:* Toponym for the Mababe Depression, Botswana.

Kinixys zombensis Hewitt 1931 (12:35)
Southeastern Hinge-back Tortoise
(includes 2 subspecies)



(subspecies: *zombensis* = red, *domerguei* = orange dots)

Distribution: Eswatini (Swaziland) (?), Kenya, Malawi, Mozambique, South Africa, Tanzania, Madagascar (*possible prehistorical introduction*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 610,369 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 2,494,004 sq. km

Size (Max SCL): male 23.0 cm, female 23.0 cm (see subspp.)

IUCN Red List: Not Evaluated (NE)


TFTSG Provisional Red List: Vulnerable (VU) (2013)

CITES: Appendix II, as Testudinidae spp. (1977). Previously: Appendix II, as *Kinixys* spp. (1975)

Kinixys zombensis zombensis Hewitt 1931 (12:35)
Southeastern Hinge-back Tortoise




Flora Ihlow / Bonamanzi Game Reserve, KwaZulu Natal, South Africa





Flora Ihlow / KwaZulu Natal, South Africa



(subspecies: *zombensis* = red; stars = type localities)

Distribution: Eswatini (Swaziland) (?), Kenya, Malawi, Mozambique, South Africa, Tanzania

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 609,534 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 1,400,516 sq. km

Size (Max SCL): male 20.6 cm, female 21.7 cm (Mifsud and Stapleton 2014)

CITES: Appendix II, as Testudinidae spp. (1977). Previously: Appendix II, as *Chersina* spp. (1975)

Synonymy:


Kinixys belliana zombensis Hewitt 1931:469
Kinixys zombensis, *Kinixys zombensis zombensis*
Type locality: "Zomba, Nyassaland" [Malawi] [yellow star].
Type specimen: PEM 14961 (formerly AMG 121), holotype, see Uetz et al. (2019).
Etymology:* Toponym for Zomba, Nyasaland (now Malawi).

Kinixys belliana zuluensis Hewitt 1931:471
Kinixys zuluensis
Type locality: "Richards Bay, Zululand" [South Africa] [green star].
Type specimen: NM 1203, lectotype, designated as the type by Broadley (1981b:210,f.10).
Etymology:* Toponym for Zululand, South Africa.

Kinixys zombensis domerguei (Vuillemin 1972b) [12:35]
Madagascan Hinge-back Tortoise


Thomas E.J. Leuteritz / CRM 6 / Nosy Faly, Madagascar


Flora Ihlow / Madagascar


(subspecies: *domerguei* = purple; orange dots; star = type locality)

Distribution: Madagascar (*possible prehistorical introduction*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 835 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 1,520 sq. km

Size (Max SCL): male 23.0 cm, female 23.0 cm (Pedrono 2008)

CITES: Appendix II, as Testudinidae spp. (1977). Previously: Appendix II, as *Chersina* spp. (1975)

Synonymy:
Madakinixys domerguei Vuillemin 1972b:169
Kinixys belliana domerguei, Kinixys zombensis domerguei

Type locality: "Madagascar." Restricted to "canton d'Antsakoamanondro, N.W. de Madagascar" [orange star] by Bour (1985:60).

Type specimen: MNHN 1993.0275 (formerly A.275), holotype, see Bour (2006e).

Etymology:* Eponym honoring Charles A. Domergue (1914–2008), France / Madagascar; naturalist, ornithologist, and herpetologist who lived and worked in Madagascar most of his life.

Malacochersus Lindholm 1929


Pancake Tortoises
(includes 1 species, 1 taxon)
Malacochersus Lindholm 1929:285


Type species: *Testudo (Malacochersus) tornieri* [= *Testudo tornieri* Siebenrock 1903b], by original designation.

Etymology:* Greek, *malakos* = soft, tender, and *chersos*, tortoise; referring to the soft, relatively flexible shell.

Malacochersus tornieri (Siebenrock 1903b)
Pancake Tortoise


Fabian Schmidt / CBFTT / Ruaha National Park, Tanzania


Fabian Schmidt / CBFTT / Ruaha National Park, Tanzania


(orange dots = probable trade or erroneous records; stars = type localities)

Distribution: Kenya, Tanzania, Zambia

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 121,863 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 624,474 sq. km

Size (Max SCL): male 17.0 cm, female 17.8 cm (Mwaya et al. 2018 CBFTT)

CBFTT Account: Mwaya, Moll, Malonza, and Ngwava (2018)

IUCN Red List: Critically Endangered (CR A4abcd) (Mwaya, Malonza, Ngwava, Moll, Schmidt, and Rhodin 2019). Previously: Vulnerable (VU) (TFTSG 1996)

CITES: Appendix I (2019). Previously: Appendix II, as

Testudinidae spp. (1977); Appendix II, as *Kinixys* spp. (1975)

Synonymy:

Testudo tornieri Siebenrock 1903b:443

Testudo (Malacochersus) tornieri, Malacochersus tornieri

Type locality: "Bussisia am Viktoria Nyanza" [Tanzania] [yellow star]. Emended to "Busisi, s. end of Lake Victoria, Tanganyika Territory" [Tanzania] by Loveridge and Williams (1957:286).

Type specimen: ZMB 11748, holotype, figured (f.1-3), destroyed during World War II, see Fritz et al. (1994); Iverson (1992) and Uetz et al. (2019) erroneously listed ZMB 11740 as holotype.

Etymology:* Eponym honoring Gustav Tornier (1859–1938), Germany; zoologist, herpetologist, and curator at Humboldt University, Berlin.

Testudo loveridgii Boulenger 1920:263

Malacochersus loveridgii

Type locality: "Afrique orientale à Dodoma, entre Kilossa et Tabora" [Tanzania] [green star]. Emended to "Dodoma, Ugogo, Tanganyika Territory" [Tanzania] by Loveridge and Williams (1957:287).

Type specimens: NHMUK 1946.1.22.50–51 (formerly 1920.5.4.1–2) and MCZ 39956–57, syntypes (4), see Barbour and Loveridge (1946).

Etymology:* Eponym honoring Arthur Loveridge (1891–1980), United Kingdom / Kenya / USA; herpetologist and curator at first the Nairobi Museum and then the Museum of Comparative Zoology, Harvard University.

Psammobates Fitzinger 1835 [25]*

South African Star Tortoises

(includes 3 species, 5 taxa)

Psammobates Fitzinger 1835:113

Type species: *Psammobates geometricus* Fitzinger [= *Testudo geometrica* Linnaeus 1758], by subsequent designation by Fitzinger (1843:29).

Etymology:* Greek, *psammos*, sand, dune, and *baino*, walk, run, to wander; referring to the dry habitats where the genus occurs.

Chersinella Gray 1870c:8

Type species: *Chersinella geometrica* [= *Testudo geometrica* Linnaeus 1758], by subsequent designation by Hewitt (1933:259).

Etymology:* Greek, *chersos*, tortoise, and Latin, *-ella*, small.

Comment: Genus established as *Peltastes (Chersinella)* without a type species. Lindholm (1929:286) previously designated *Testudo graeca* Linnaeus 1758 as type species, but it was not originally included in *Chersinella* by Gray (1870c), so therefore has no validity as type.

Psammobates geometricus (Linnaeus 1758)


Geometric Tortoise



Eric V. Goode / South African Geometric Tortoise Reserve, South Africa



Eric V. Goode / South African Geometric Tortoise Reserve, South Africa



(stars = type localities)

Distribution: South Africa

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO): 6,261 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 12,260 sq. km

Size (Max SCL): male 12.3 cm, female 16.5 cm (Baard 1995; Ernst et al. 2006b; Itescu et al. 2014)

IUCN Red List: Critically Endangered (CR A4ace) (Hofmeyr and Baard 2018). Previously: Critically Endangered (CR) (Hofmeyr and Baard 2015); Endangered (EN) (TFTSG *in* Baillie and Groombridge 1996)

CITES: Appendix I (1975)

USA ESA:* Endangered (USFWS 1976)

Synonymy:

Testudo geometrica Linnaeus 1758:199

Chersine geometrica, Hydrone geometrica, Psammobates geometricus, Peltastes geometricus, Peltastes geometrica, Peltastes (Chersinella) geometrica, Chersinella geometrica, Psammobates geometrica, Geochelone geometrica

Type locality: "Asia." Restricted to "southwestern Cape Province, South Africa" [yellow star, location estimated] by Baard (1991:9).

Type specimen: UPSZTY 20 (formerly UUZM 20), syntype, recorded as type by Wallin (1977, 2001), but identified as a *Testudo elegans* Schoepff (= *Geochelone elegans*); specimen figured in Piso (1658:105:f.1) (part of the composite syntype series), designated lectotype by Hoogmoed and Crumly (1984:255).

Etymology:* Greek, *gaia, geo-*, the earth, and *metriki*, metrics; referring to the geometrical rayed pattern of the carapace.

Comment: Description cited as sourced from Linnaeus (1749a:139, n.24), Worm (1655:317), Piso (1658:pl.105), Grew (1681:36,pl.3,f.1–2), Ray (1693:259), Seba (1734: pl.80,f.8), and Linnaeus (1754:50), but this last entry is in error, as this specimen at NRM = *Testudo amboinensis* Linnaeus 1754 (= *Rhinoclemmys punctularia*; A.G.J. Rhodin and E. Åhlander, unpubl. data) and was previously erroneously identified as a *Mesoclemmys gibba* (Andersson 1900; Wallin 1977).

Testudo luteola Daudin 1801:277

Type locality: Not known.

Type specimen: Possibly MNHN, not located, type specimen figured (pl.25,f.3).

Etymology:* Latin, *luteolus*, yellowish; referring to the yellowish color of the carapace in the juvenile type specimen.

Peltastes geographicus Gray 1869a:173 (*nomen novum*)

Testudo geographica

Etymology:* Greek, *gaia, geo-*, the earth, and *grapho*, drawing or writing; referring to the rayed pattern of the carapace.

Comment: Erroneous misprint for *geometrica*, see Gray (1870c:9).

Testudo strauchi Lidth de Jeude 1893:312

Chersinella strauchi

Type locality: "Cape of Good Hope" [South Africa] [probable trade specimen, orange star].

Type specimen: RMNH 6011, holotype, discussed by Hoogmoed and Crumly (1984) and Hoogmoed et al. (2010).

Etymology:* Eponym honoring Alexander A. Strauch (1832–1893), Russia; herpetologist and director of the Zoological Museum of the Imperial Academy of Sciences at St. Petersburg.

Psammobates oculifer (Kuhl 1820)


Serrated Tent Tortoise, Kalahari Tent Tortoise



William R. Branch / Botswana



Victor J.T. Loehr / Northern Cape Prov., South Africa



(orange dots = trade or possible misidentified; stars = type localities)

Distribution: Botswana, Namibia, South Africa, Zimbabwe

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO): 1,078,671 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 1,477,015 sq. km

Size (Max SCL): male 11.8 cm, female 14.7 cm (Boycott and Bourquin 2000; Itescu et al. 2014)

IUCN Red List: Least Concern (LC) [Not Listed] (Behler 1996)

TFTSG Provisional Red List: Least Concern (LC) (2013)

CITES: Appendix II, as Testudinidae spp. (1977). Previously: Appendix II (1975)

Synonymy:

Testudo oculifera Kuhl 1820:77

Emys oculifera, Clemmys oculifera, Chersinella oculifera, Psammobates oculifera, Psammobates oculifer, Psammobates oculiferus

Type locality: "Cap" [Cape of Good Hope, South Africa] [in error] [orange star].

Type specimen: ZMB 223, holotype, apparently lost, destroyed during World War II, see Fritz et al. (1994).

Etymology:* Latin, *oculus*, eye, and *ferre*, to bear, to carry; referring to the stripe pattern that forms an eye spot pattern in the area between adjacent carapacial scutes.

Emys occilifera Kuhl *in* Gray 1830e:9 (*nomen novum*)


Etymology:* Latin, *oculus*, eye, and *ferre*, to bear, to carry; referring to the stripe pattern that forms an eye spot pattern in the area between

adjacent carapacial scutes.
Comment: Unjustified emendation or error for *oculifera*.

Emys kuhlii Gray 1831d:73 (*nomen dubium*)
Type locality: Not known.
Type specimen: ZMB s/n, holotype, apparently lost.
Etymology:* Eponym honoring Heinrich Kuhl (1797–1821), Germany / Netherlands / Indonesia; zoologist and herpetologist who worked in the Netherlands and on Java in the Dutch East Indies (now Indonesia).

Testudo semiserrata Smith 1839a:Reptilia,pl.6
Peltastes semiserratus
Type locality: "districts between Latakoo and the Tropic of Capricorn" [= north of Dithakong, in North-West Province, South Africa] [green star, location estimated].
Type specimens: NMSZ 1859.013.1729 (formerly RSM) and NHMUK 1947.3.5.13 (formerly 1863.2.21.44), syntypes (2); NMSZ 1859.013.1729 listed as "holotype" by Herman et al. (1990).
Etymology:* Latin, *semi*, half, and *serrata*, serrated, jagged; referring to the serrated anterior and posterior shell margins, but only weakly serrated on the sides.

Psammobates tentorius (Bell 1828a) (17:61, 21:94)
Tent Tortoise
(includes 3 subspecies)



(subspecies: *tentorius* = red, *trimeni* = blue, *verroxii* = green; overlap [brown] = intergrades; orange dots = probable trade or introduced)

Distribution: Namibia, South Africa
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 548,201 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 742,135 sq. km
Size (Max SCL): male 12.0 cm, female 14.5 cm (see subspp.)
IUCN Red List: Least Concern (LC) (Hofmeyr, Leuteritz, and Baard 2018). Previously: Least Concern (LC) [Not Listed] (Behler 1996)
CITES: Appendix II, as Testudinidae spp. (1977). Previously: Appendix II (1975)


Psammobates tentorius tentorius (Bell 1828a) (21:94)
Southern Tent Tortoise, Common Tent Tortoise



Victor J.T. Loehr / nr. Prince Albert, Western Cape Prov., South Africa



Victor J.T. Loehr / *left*: Northern Cape Prov., *right*: Western Cape Prov., South Africa



(subspecies: *tentorius* = red, *trimeni* = blue, *verroxii* = green; overlap [brown] = intergrades; orange dot = probable trade or translocated; stars = type localities)

Distribution: South Africa
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 174,265 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 283,555 sq. km
Size (Max SCL): male 10.0 cm, female 13.1 cm (Boycott and Bourquin 2000)
TFTSG Provisional Red List: Least Concern (LC) (2017)
CITES: Appendix II, as Testudinidae spp. (1977). Previously: Appendix II, as *Chersina* spp. (1975)
Synonymy:
Testudo tentoria Bell 1828a:420
Testudo geometrica tentoria, Peltastes tentorius, Peltastes (Chersinella) tentoria, Chersinella tentoria, Chersinella tentoria tentoria, Psammobates tentoria, Psammobates tentoria tentoria, Psammobates tentorius, Psammobates tentorius tentorius, Testudo tentoria tentoria
Type locality: "Africâ?" [no star].
Type specimen: OUM 8570, holotype, discussed by Nowak-Kemp and Fritz (2010).
Etymology:* Latin, *tentoria*, tent; referring to the tent-like raised vertebral and pleural scutes.

Testudo geometrica nigriventris Gray 1856b:8
Type locality: "South Africa" [no star].
Type specimens: NHMUK s/n, not located, syntypes (6).
Etymology:* Latin, *niger*, black, dark-colored, and *venter*, belly,

abdomen; referring to the large dark plastral figure.

Chersinella tentoria albanica Hewitt 1933b:266

Psammobates tentoria albanica

Type locality: "neighbourhood of farm Mayfair, Albany District" [Eastern Cape Province, South Africa] [green star, location estimated].

Type specimens: Possibly PEM or AMG, syntypes (4), figured (pl.14,f.10-15).

Etymology:* Toponym for Albany District, South Africa.

Chersinella tentoria tentorioides Hewitt 1933b:268

Psammobates tentoria tentorioides

Type locality: "Bowden Hall, Middelburg district, C.P." [Eastern Cape Province, South Africa] [blue star].

Type specimen: Possibly PEM or AMG, holotype, figured (pl.14,f.16).

Etymology:* Latin, *tentoria*, tent, and *-oides*, similar; referring to the tent-like raised vertebral and pleural scutes similar to those in *Testudo tentoria*.

Chersinella tentoria piscatella Hewitt 1933b:269

Psammobates tentoria piscatella

Type locality: "Little Fish River, Somerset East district" [Eastern Cape Province, South Africa] [pink star].

Type specimen: Possibly PEM, holotype, figured (pl.14,f.17-18).

Etymology:* Toponym (Latin, *pisces*, fish, and *-ella*, small) for the Little Fish River, South Africa.

Chersinella tentoria subsulcata Hewitt 1933b:270

Type locality: "farm 'Brighton' near Steytlerville" [Eastern Cape Province, South Africa] [light gray star].

Type specimen: Possibly PEM or AMG, holotype.

Etymology:* Latin, *sub-*, below, under, *sulcus*, groove or furrow, and *-ata*, having; referring to the furrows on the carapace.

Chersinella tentoria karuica Hewitt 1933b:272

Psammobates tentoria karuica

Type locality: "farm 'Drogekloof' near Klaarstroom, Prince Albert district" [Western Cape Province, South Africa] [dark gray star].

Type specimens: Possibly PEM or AMG, syntypes (9), figured (pl.14,f.19-24,pl.15,f.25-28).

Etymology:* Toponym for the Karoo (previously spelled Karru), a semi-desert natural region of South Africa.

Chersinella tentoria duerdeni Hewitt 1933b:279

Psammobates tentoria duerdeni

Type locality: "Graaff Reinet, C.P." [Eastern Cape Province, South Africa] [black star].

Type specimen: Possibly PEM or AMG, holotype, figured (pl.15,f.29).

Etymology:* Eponym honoring James E. Duerden (1865–1937), United Kingdom / South Africa; zoologist, herpetologist, and turtle researcher in South Africa.

Chersinella tentoria lativittata Hewitt 1933b:281

Type locality: "Willowmore" [Eastern Cape Province, South Africa] [white star].

Type specimens: Possibly PEM or AMG, syntypes (2), figured (pl.15,f.32-34).

Etymology:* Latin, *latum*, broad, wide, extended, and *vitta*, band, stripe; referring to the relatively broad stripes on the carapace.

Chersinella tentoria karuella Hewitt 1933b:283

Type locality: "Uniondale, C.P." [Western Cape Province, South Africa] [purple star].

Type specimen: Possibly PEM or AMG, holotype, figured (pl.15,f.35).

Etymology:* Toponym (Vernacular, *karru*, and Latin, *-ella*, small) for the Little Karoo (previously spelled Karru), a semi-desert natural region of South Africa.

Chersinella tentoria hexensis Hewitt 1933b:286

Type locality: "Hex River, Worcester district" [Western Cape Province, South Africa] [brown star].

Type specimens: Possibly PEM or AMG, syntypes (4), figured (pl.15,f.39-40).

Etymology:* Toponym for the Hex River, Western Cape Province, South Africa.

Psammobates tentorius trimeni (Boulenger 1886a) [21:94]


Western Tent Tortoise



William R. Branch / Groenrivier, South Africa / male



left: Victor J.T. Loehr / Northern Cape Prov., South Africa
right: William R. Branch / Groenrivier, Northern Cape Prov., South Africa / male



(subspecies: *tentorius* = red, *trimeni* = blue, *verroxii* = green; overlap [brown] = intergrades; star = type locality)

Distribution: Namibia, South Africa

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO): 32,253 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 46,987 sq. km

Size (Max SCL): male 10.0 cm, female 14.5 cm (Boycott and Bourquin 2000)

TFTSG Provisional Red List: Endangered (EN) (2017)

CITES: Appendix II, as Testudinidae spp. (1977). Previously: Appendix II, as *Chersina* spp. (1975)

Synonymy:

Testudo trimeni Boulenger 1886a:541

Chersinella trimeni, Psammobates trimeni, Psammobates tentorius trimeni, Testudo tentoria trimeni, Psammobates tentoria trimeni

Type locality: "Mouth of the Orange River" [Little Namaqualand, Northern Cape Province, South Africa] [yellow star].


Type specimens: NHMUK 1947.3.5.10–12, syntypes (3).

Etymology:* Eponym honoring Roland Trimen (1840–1916), United Kingdom / South Africa; entomologist who worked in South Africa and became director of the South African Museum.


Psammobates tentorius verroxii (Smith 1839b) [(21:94)]
Northern Tent Tortoise



Victor J.T. Loehr / nr. Beaufort West, Northern Cape Prov., South Africa



Victor J.T. Loehr / Northern Cape Prov., South Africa



(subspecies: *tentorius* = red, *trimeni* = blue, *verroxii* = green; overlap [brown] = intergrades; orange dot = probable trade; stars = type localities)

Distribution: Namibia, South Africa

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 367,871 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 531,092 sq. km

Size (Max SCL): male 12.0 cm, female 14.0 cm (Boycott and Bourquin 2000)

TFTSG Provisional Red List: Least Concern (LC) (2017)

CITES: Appendix II, as Testudinidae spp. (1977). Previously: Appendix II, as *Chersina* spp. (1975)

Synonymy:

Testudo verroxii Smith 1839b:Reptilia,pl.8
Peltastes verroxii, Chersinella verroxii, Chersinella verroxii verroxii, Psammobates tentorius verroxii, Testudo tentoria verroxii, Psammobates tentoria verroxii, Testudo tentorius verroxii
Type locality: "South Africa, near the sources of the Garriep or Orange River." Restricted to "north of Aliwal North, between the Orange and Caledon Rivers...roughly 260 miles east of Niekirk's Hope" [Free State Province, South Africa] [yellow star] by Power (1932:466).
Type specimen: NMSZ 1859.013.1680 (formerly RSM), holotype, see FitzSimons (1937); Herman et al. (1990) erroneously recorded it as 1959.013.1680 (a typo).
Etymology:* Eponym honoring Jules P. Verreaux (1807–1873), France / South Africa; botanist, ornithologist, and naturalist collector and trader in South Africa, who helped Smith found the South African Museum, Cape Town.

Peltastes verreauxii Gray 1870b:656 (*nomen novum*)
Testudo verreauxii, Psammobates verreauxii
Etymology:* Eponym honoring Jules P. Verreaux (1807–1873), France / South Africa; botanist, ornithologist, and naturalist collector and trader in South Africa, who helped Smith found the South African Museum, Cape Town.
Comment: Unjustified emendation of *verroxii*.

Testudo fiski Boulenger 1886a:542
Testudo tentoria fiski, Chersinella fiski, Chersinella fiski fiski, Psammobates fiski, Psammobates fiski fiski
Type locality: "De Aar, not far from Hopetown" [Northern Cape Province, South Africa] [green star].
Type specimen: NHMUK 1946.1.22.65 (formerly 1886.8.30.1), holotype.
Etymology:* Eponym honoring Reverend George H.R. Fisk (1829–1911), United Kingdom / South Africa; clergyman and naturalist who worked in South Africa in the 1870s and collected the types.

Testudo smithi Boulenger 1886a:542
Chersinella verroxii smithi, Testudo smithi smithi, Testudo verroxii smithi
Type locality: "S. Africa" [South Africa] [no star].
Type specimen: NHMUK 1947.3.4.2 (formerly 1847.3.5.20), holotype.
Etymology:* Eponym honoring Sir Andrew Smith (1797–1872), United Kingdom (Scotland) / South Africa; physician, administrator, and zoologist, who worked primarily in South Africa.

Testudo seimundi Boulenger 1903b:216
Chersinella fiski seimundi
Type locality: "3 miles east of Deelfontein" [Northern Cape Province, South Africa] [pink star].
Type specimen: NHMUK 1946.1.22.58 (formerly 1903.4.27.1), holotype.
Etymology:* Eponym honoring Elbert C.H. Seimund (1878–1942), United Kingdom / South Africa; military medical technician and taxidermist, worked for the British Museum in South Africa and collected the type.

Testudo boettgeri Siebenrock 1904a:194 (junior homonym, not = *Testudo graeca boettgeri* Mojsisovics 1889 [= *Testudo (Chersine) hermanni boettgeri*])
Chersinella verroxii boettgeri, Chersinella boettgeri
Type locality: "Groß-Namaland in Südwestafrika" [Namaland, Karas Region, Namibia] [light gray star, location estimated].
Type specimen: SMF 7793, holotype, see Mertens (1967b).
Etymology:* Eponym honoring Oskar Boettger (or Böttger) (1844–1910), Germany; paleontologist, herpetologist, and curator at the Senckenberg Museum, Frankfurt.

Homopus bergeri Lindholm 1906:348 [(07:67)]
Testudo bergeri, Chersinella verroxii bergeri, Testudo smithi bergeri, Testudo verroxii bergeri
Type locality: "nach Gibeon in Deutsch-Südwestafrika" [Hardap Region, Namibia] [dark gray star].
Type specimen:* MWNH 711, holotype, discussed by Karl and Tichy (1999) and Branch (2007).
Etymology:* Eponym honoring Carl Berger (1871–1962), Germany / Namibia; missionary in South West Africa (now Namibia), who collected the types. The entry for Berger in Beolens et al. (2011) is incorrect.

Testudo oscarboettgeri Lindholm 1929:295 (*nomen novum*)
Type locality: "Groß-Namaland in Südwestafrika" [Namaland, Karas Region, Namibia] [light gray star].
Type specimen: SMF 7793, holotype, by default, see Mertens (1967b).
Etymology:* Eponym honoring Oskar Boettger (or Böttger) (1844–1910), Germany; paleontologist, herpetologist, and curator at the Senckenberg Museum, Frankfurt.
Comment: Justified emendation for *Testudo boettgeri* Siebenrock 1904a.

Chersinella schonlandi Hewitt 1934:303

Type locality: "Namaqualand, C.P." [nr. Okiep, Little Namaqualand, Northern Cape Province, South Africa] [black star, location estimated].
Type specimen: Possibly PEM or AMG, holotype, figured (pl.16,f.46).
Etymology:* Eponym honoring Selmar Schonland (or Schönland) (1860–1940), Germany / South Africa; botanist and curator at the Albany Museum, Grahamstown.

Chersinella fiski cronwrighti Hewitt 1934:317
Psammobates fiski cronwrighti
Type locality: "Hanover, C.P." [Northern Cape Province, South Africa] [white star].
Type specimen: Possibly PEM or AMG, holotype, figured (pl.16,f.58).
Etymology:* Eponym honoring Samuel C. Cronwright (1863–1936), South Africa; farmer, biographer, and naturalist who collected and donated several tortoises to the Albany Museum, Grahamstown.

Chersinella fiski orangensis Hewitt 1934:319
Type locality: "between Philipstown and Petrusville district" [Northern Cape Province, South Africa] [purple star].
Type specimen: Possibly PEM or AMG, holotype, figured (pl.16,f.59-60).
Etymology:* Toponym for the Orange River, Northern Cape, South Africa.

Chersinella fiski colesbergensis Hewitt 1934:321
Psammobates fiski colesbergensis
Type locality: "Colesberg" [Northern Cape Province, South Africa] [light brown star].
Type specimen: Possibly PEM or AMG, holotype, figured (pl.16,f.61).
Etymology:* Toponym for Colesberg, Northern Cape, South Africa.

Chersinella fiski grica Hewitt 1934:323
Type locality: "Marydale, ... midway between Prieska and Kenhardt" [Northern Cape Province, South Africa] [dark brown star].
Type specimens: Possibly PEM or AMG, syntypes (2), figured (pl.16,f.63,pl.17,f.67).
Etymology:* Ethnonym honoring the Griqua, semi-nomadic, Afrikaans-speaking people in the area of the former province of Griqualand West of the Cape Colony, now Northern Cape, South Africa.

Chersinella fiski gricoides Hewitt 1934:326
Type locality: "Niekerks Hope, about midway between Griquatown and Prieska, C.P." [Niekerkshoop, Northern Cape Province, South Africa] [light blue star].
Type specimens: Possibly PEM or AMG, syntypes (2), figured (pl.17,f.72-75).
Etymology:* Ethnonym honoring the Griqua, semi-nomadic, Afrikaans-speaking people in the area of the former province of Griqualand West of the Cape Colony, now Northern Cape, South Africa, and Latin, *-oides*, similar.

Chersinella verroxii amasensis Hewitt 1934:333
Type locality:* "Ukamas district" [Karas Region, Namibia]; erroneously emended to "Ukamas District, Kap-Provinz" [South Africa] by Wermuth and Mertens (1961:228); hereby corrected to "Ukamas district (ca. 28.03S, 19.73E), Karas Region (Great Namaqualand), Namibia" [dark blue star].
Type specimens: Possibly PEM or AMG, syntypes (2).
Etymology:* Toponym for Amas (= Ukamas), Great Namaqualand, former South West Africa (now Namibia).

Psammobates depressa FitzSimons 1938:154
Type locality: "8 miles west of Aus, Great Namaqualand" [Karas Region, Namibia] [orange star].
Type specimen: TMP 17821, holotype.
Etymology:* Latin, *de-*, down, and *pressus*, pressed; referring to the depressed flattened shell.

Pyxis Bell 1827 [84]*

Spider Tortoises
(includes 2 species, 4 taxa)

Pyxis Bell 1827:395 [84]*
Type species: *Pyxis arachnoides* Bell 1827, by original monotypy.
Etymology:* Greek, *pyxis*, box; referring to the box-like appearance of these turtles.
Comment:* Previously considered an apparent junior homonym of *Pyxis* Chemnitz 1784 (= Brachiopoda and Bivalvia).


Acinixys Siebenrock 1902b:12
Type species: *Acinixys planicauda* [= *Testudo planicauda* Grandidier 1867], by original monotypy.
Etymology:* Greek, *a-*, without, and *kineo*, to move, and *ixys*, the small of the back; referring to the non-kinetic carapace.

Bellemys Williams 1950a:512 (*nomen novum*) [84]*
Etymology:* Eponym honoring Thomas Bell (1792–1880), United Kingdom; herpetologist, dental surgeon, and Professor of Zoology at King's College, London.
Comment:* Initially considered a valid replacement name for *Pyxis* Bell 1827, but determined to be invalid.

Pyxoides Vuillemin and Domergue 1972:193
Type species: *Pyxoides brygooi* Vuillemin and Domergue 1972, by original monotypy.
Etymology:* Greek, *pyxis*, box, and *-oides*, resemblance; referring to the box-like shell.

Pyxis arachnoides Bell 1827

Spider Tortoise, *Kapila*
(includes 3 subspecies)



(subspecies: *arachnoides* = red, *brygooi* = blue, *oblonga* = green; overlap [purple] = intergrades)

Distribution: Madagascar
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 18,559 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 77,591 sq. km
Size (Max SCL): male 14.4 cm, female 15.7 cm (see subspp.)
IUCN Red List: Critically Endangered (CR A4cd; E) (Leuteritz and Walker 2020). Previously: Critically Endangered (CR) (Leuteritz and Walker 2014); Critically Endangered (CR) (Leuteritz and Walker 2008); Vulnerable (VU) (TFTSG *in* Baillie and Groombridge 1996)
CITES: Appendix I (2005). Previously: Appendix II, as Testudinidae spp. (1977); Appendix II, as *Pyxis* spp. (1975)


Pyxis arachnoides arachnoides Bell 1827 (07:70)
Spider Tortoise, Common Spider Tortoise



Michael Ogle / Anakao, Madagascar



left: Anders G.J. Rhodin / Madagascar / captivity
right: Michael Ogle / Anakao, Madagascar



(subspecies: *arachnoides* = red, *brygooi* = blue, *oblonga* = green; overlap [purple] = intergrades; stars = type localities)

Distribution: Madagascar

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 7,088 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 15,907 sq. km

Size (Max SCL): male 14.4 cm, female 15.7 cm (Pedrono 2008; Pedrono and Smith 2013; R. Walker, unpubl. data)

CITES: Appendix I (2005). Previously: Appendix II, as Testudinidae spp. (1977); Appendix II, as *Pyxis* spp. (1975)

Synonymy:

Pyxis arachnoides Bell 1827:395
Testudo (Pyxis) arachnoides, Testudo arachnoides, Pyxis arachnoidea, Bellemys arachnoides, Pyxis arachnoides arachnoides
Type locality: Not known. Restricted to "Soalara (Baie de Saint-Augustin), sud-ouest de Madagascar" [yellow star] by Bour (1979:153).
Type specimen: OUM 1092, lectotype, designated by Bour (1979:153), discussed by Nowak-Kemp and Fritz (2010).
Etymology:* Greek, *arachne*, a spider, and *-oides*, resembling; referring to the spidernet-like reticulations on the carapace.


Testudo (Pyxis) aranoides Gray 1830e:6 (*nomen novum*)
Pyxis aranoides
Etymology:* Latin, *aranea*, a spider, and Greek, *-oides*, resembling, referring to the spidernet-like reticulations on the carapace.
Comment: Unjustified emendation or error for *arachnoides*.

Pyxis madagascariensis Lesson 1831a:120
Type locality: "Madagascar" [green star, location estimated].
Type specimen: Possibly MNHN, holotype, not located.
Etymology:* Toponym for Madagascar.


Pyxis arachnoides brygooi (Vuillemin and Domergue 1972)
Northern Spider Tortoise



Roger Bour / Salary, nr. Forêt des Mikea, Madagascar



Roger Bour / Salary, nr. Forêt des Mikea, Madagascar



(subspecies: *arachnoides* = red, *brygooi* = blue; overlap [purple] = intergrades; star = type locality)

Distribution: Madagascar

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 5,464 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 7,544 sq. km

Size (Max SCL): male 12.3 cm, female 12.8 cm (R. Walker, unpubl. data)

CITES: Appendix I (2005). Previously: Appendix II, as Testudinidae spp. (1977); Appendix II, as *Pyxis* spp. (1975)


Synonymy:

Pyxoides brygooi Vuillemin and Domergue 1972:193
Pyxis arachnoides brygooi
Type locality: "la côte Sud-Ouest, entre Morombe et Tuléar, aux alentours du lac Ihotry, dans la forêt des Mikea" [Madagascar]. Restricted to "Ampanonga (N.-W. Lac Ihotry), sud-ouest de Madagascar" [yellow star] by Bour (1979:153).
Type specimen: MNHN A.277, holotype, see Bour (1981).
Etymology:* Eponym honoring Édouard-Raoul Brygoo (1920–2016), France / Madagascar; physician and herpetologist who worked in Madagascar.

Pyxis arachnoides oblonga Gray 1869a
Southern Spider Tortoise



Anders G.J. Rhodin / TCC / CRM 6 / Berenty, Madagascar



Anders G.J. Rhodin / Berenty, Madagascar



(subspecies: *arachnoides* = red, *oblonga* = green; star = type localities)

Distribution: Madagascar

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 6,849 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 12,603 sq. km

Size (Max SCL): male 11.8 cm, female 15.4 cm (R. Walker, unpubl. data)

CITES: Appendix I (2005). Previously: Appendix II, as Testudinidae spp. (1977); Appendix II, as *Pyxis* spp. (1975)

Synonymy:


Pyxis arachnoidea oblonga Gray 1869a:173
Pyxis arachnoides oblonga
Type locality: Not known. Restricted to "Asia" by Gray (1873j:14); to "Madagascar; Mauritius" by Boulenger (1889:145); and to "Cap Sainte-Marie, province de Tuléar (Toliara), sud du Madagascar" [yellow star] by Bour (1982b:30).
Type specimen: NHMUK 1861.3.20.31, holotype, see Bour (1982b).
Etymology:* Latin, *oblongus*, elongated; referring to the somewhat elongated carapace.

Pyxis arachnoides matzi Bour 1979:143
Type locality: "Cap Sainte-Marie (Province de Tuléar), extrême sud du Madagascar" [yellow star]
Type specimen: MNHN 1978.24, holotype, see Bour (1981).
Etymology:* Eponym honoring Gilbert Matz (1934–), France; biologist and herpetologist at the University of Angers.


Pyxis planicauda (Grandidier 1867)
Flat-tailed Tortoise, Flat-shelled Spider Tortoise, *Kapidolo*



Miguel Pedrono / CRM 6 / Kirindy Forest, Madagascar



Anders G.J. Rhodin / captivity / Ampijoroa, Madagascar



(star = type localities)

Distribution: Madagascar

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 5,374 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 6,605 sq. km

Size (Max SCL): male 14.1 cm, female 15.9 cm (Bloxam and Hayes 1991; Pedrono and Smith 2013)

IUCN Red List: Critically Endangered (CR A4acd) (Leuteritz, Randriamahazo, and Lewis 2008). Previously: Endangered (TFTSG *in* Baillie and Groombridge 1996)

CITES: Appendix I (2003). Previously: Appendix II, as Testudinidae spp. (1977); Appendix II, as *Pyxis* spp. (1975)

Synonymy:

Testudo planicauda Grandidier 1867:223
Acinixys planicauda, Pyxis (Acinixys) planicauda, Pyxis planicauda
Type locality: "Mouroundava...cote sud-ouest de Madagascar" [Morondava] [yellow star].
Type specimen: MNHN 9373 (formerly 1867.44), holotype; NMW 23479, listed as "syntype" by Tiedemann and Häupl (1980),

Tiedemann et al. (1994), and Gemel et al. (2019), is not a type; discussed by Bour (2005c).
Etymology:* Latin, *plana*, flat, even, and *cauda*, tail; referring to the strongly flattened tail.

Testudo morondavaensis Vuillemin 1972a:127
Type locality: "Morondava (côte sud-ouest de Madagascar)" [yellow star].
Type specimen: MNHN A.276, holotype, discussed by Bour (2005c).
Etymology:* Toponym for Morondava, Madagascar.

Stigmochelys Gray 1873j (07:52, 10:26) (25)*

Leopard Tortoises
(includes 1 species, 1 taxon)

Stigmochelys Gray 1873j:5
Type species: *Testudo (Stigmochelys) pardalis* Bell 1828a, by original monotypy.
Etymology:* Greek, *stigma*, spot, point, mark, and *chelys*, turtle; referring to the conspicuous carapacial spot and line pattern.

Megachersine Hewitt 1933b:257
Type species: *Megachersine pardalis* [= *Testudo pardalis* Bell 1828a], by original designation.
Etymology:* Greek, *megas*, big, and *chersos*, tortoise; referring to the large size.

Stigmochelys pardalis (Bell 1828a) (07:71, 10:27, 21:95)

Leopard Tortoise




Victor J.T. Loehr / nr. Beaufort West, Northern Cape, South Africa







left: Flora Ihlow / South Africa
right: Ronald Orenstein / Addo Elephant Park, South Africa



**



**



**

(orange dots = probably introduced or translocated; stars = type localities)

Distribution: Angola, Botswana, Burundi, Djibouti (?) (*extirpated?*), Eswatini (Swaziland), Ethiopia, Kenya, Malawi, Mozambique, Namibia, Rwanda, Somalia, South Africa, South Sudan, Tanzania, Uganda, Zambia, Zimbabwe

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 4,450,736 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 11,956,566 sq. km

Size (Max SCL): male 65.5 cm, female 75.0 cm (Lambert 1995; Lambert et al. 1998; Boycott and Bourquin 2000)

IUCN Red List: Least Concern (LC) (Baker, Kabigumila, Leuteritz, Hofmeyr, and Ngwava 2015). Previously: Least Concern (LC) [Not Listed] (Behler 1996)

CITES: Appendix II, as Testudinidae spp. (1977). Previously: Appendix II (1975)

Synonymy:

Testudo pardalis Bell 1828a:420

Geochelone (Geochelone) pardalis, Geochelone pardalis, Testudo (Stigmochelys) pardalis, Megachersine pardalis, Testudo pardalis pardalis, Geochelone pardalis pardalis, Stigmochelys pardalis, Centrochelys pardalis, Centrochelys pardalis pardalis, Stigmochelys pardalis pardalis, Psammobates pardalis

Type locality: "Promont. Bonae Spei" [Cape of Good Hope, South Africa] [yellow star].

Type specimen: OUM s/n, originally live, holotype figured (pl.25), apparently lost, not listed by Nowak-Kemp and Fritz (2010).

Etymology:* Latin, *pardalis*, panther; referring to the leopard-like pattern on the shell.

Testudo biguttata Cuvier 1829:10 (*nomen nudum*)

Etymology:* Latin, *bi-*, two, and *guttata*, spotted; referring to the carapacial spots.

Testudo armata Boie *in* Gray 1830e:4 (*nomen nudum*)

Etymology:* Latin, *armata*, armed, armored, protected; referring to the strong protective shell and scaly armored front legs.

Testudo bipunctata Gray 1830e:4 (*nomen nudum*)

Etymology:* Latin, *bi-*, two, and *punctata*, spotted or punctured; referring to the carapacial spots.

Testudo pardalis babcocki Loveridge 1935:4 (10:27)

Geochelone pardalis babcocki, Geochelone babcocki, Stigmochelys pardalis babcocki

Type locality: "western slopes of Mount Debasien, Karamoja, Uganda at 5,500 feet" [green star].

Type specimen: MCZ 40003, holotype, see Barbour and Loveridge (1946).

Etymology:* Eponym honoring Harold L. Babcock (1886–1953), USA; herpetologist and curator at the Museum of the Boston Society of Natural History.

Testudo Linnaeus 1758 (07:72, 17:56, 17:70)

Palearctic Tortoises

(includes 3 subgenera, 5 species, 20 taxa)

Testudo Linnaeus 1758:197

Type species: *Testudo graeca* Linnaeus 1758 [= subjective synonym of *Testudo ibera* Pallas 1814], by subsequent designation by Fitzinger (1843:29) and Lindholm (1929:284); not *Testudo graeca* Auct. [= subjective synonym of *Testudo hermanni* Gmelin 1789] by earlier designation by Bell (1828c:514).

Etymology:* Latin, *testudo*, turtle or tortoise.

Chersine Merrem 1820:29 (09:40)

Type species: "*T. graeca* »L.« auct. = *T. hermanni* Gmelin" [= *Testudo hermanni* Gmelin 1789], by subsequent designation by Lindholm (1929:286).

Etymology:* Greek, *chersos*, tortoise.

Chersini Merrem *in* Gray 1825:210 (*nomen novum*)

Etymology:* Greek, *chersos*, tortoise.

Comment: Unjustified emendation of *Chersine*.

Chersus Wagler 1830b:138

Type species: *Chersus marginatus* [*Testudo marginata* Schoepff 1793a], by monotypy.

Etymology:* Greek, *chersos*, tortoise.

Comment: Unjustified emendation of *Chersine*.

Cherseus Gray 1856b:10 (*nomen novum*)

Etymology:* Greek, *chersos*, tortoise.

Comment: Unjustified emendation of *Chersus*.

Peltastes Gray 1869a:167,171 (junior homonym, not = *Peltastes* Illiger 1807 [= Hymenoptera] or *Peltastes* Agassiz 1838 [= Echinodermata] or *Peltastes* Fischer-Waldheim 1839 [= Orthoptera])

Type species: *Testudo graeca* Linnaeus 1758 [= *Peltastes graecus* sensu Gray 1869] [= subjective synonym of *Testudo ibera* Pallas 1814], by subsequent designation by Lindholm (1929:286).

Etymology:* Greek, *peltastes*, lightly armed foot soldiers in ancient Greece who wore light shields (*pelte*) made of twigs and leather; referring to the relatively thin shell bones in this genus.

Testudinella Gray 1870b:658 (junior homonym, not = *Testudinella* Bory de Saint-Vincent 1822 [= Rotatoria])

Type species: *Testudinella horsfieldii* [= *Testudo horsfieldii* Gray 1844], by original monotypy.

Etymology:* Latin, *testudo*, turtle, and *-ella*, small or smaller; referring to the relatively smaller size of the type species as compared to other tortoises.

Peltonia Gray 1872c:4 (*nomen novum*)

Etymology:* Greek, *peltastes*, lightly armed foot soldiers in ancient Greece who wore light shields (*pelte*) made of twigs and leather; referring to the relatively thin shell bones in this genus.

Comment: Unjustified replacement name for *Peltastes*.

Medaestia Wussow 1916:170 (09:40)

Type species: *Medaestia graeca* sensu Wussow 1916 [= subjective synonym of *Testudo hermanni* Gmelin 1789] by subsequent designation by Mertens (1949:232).

Etymology:* Anonym, *medaestia*, possibly a fabricated name, but possibly a Toponym for the Eastern Mediterranean region (*meda-* and *est*, east).

Comment: Genus referred to as *Testudo (Medaestia)* without indicating a type species, but noted as including *Medaestia graeca, M. ibera, M. leithii* [=*Testudo kleinmanni*], and *M. horsfieldii*.

Pseudotestudo Loveridge and Williams 1957:166, 276

Type species: *Testudo (Pseudotestudo) kleinmanni* [= *Testudo kleinmanni* Lortet 1883], by original designation.

Etymology:* Greek, *pseudo*, false, untrue, and Latin, *testudo*, turtle; referring to the difference of this genus from the genus *Testudo*.

Agrionemys Khosatzky and Młynarski 1966:123 (*nomen novum*)

Type species: *Agrionemys horsfieldi* [= *Testudo horsfieldii* Gray 1844], by original designation.

Etymology:* Greek, *agrios*, wild, as in living in a field, and *emys*, turtle; referring to the grassland habitat.

Comment: Justified replacement name for *Testudinella*.

Protestudo Chkhikvadze 1970:245

Type species: *Protestudo bessarabica* [= *Testudo bessarabica* † Riabinin 1918], by original designation.

Etymology:* Greek, *protos*, earliest, the first, and Latin, *testudo*, turtle; referring to this genus the ancestor of the genus *Testudo*.

Furculachelys Highfield 1990:1

Type species: *Furculachelys nabeulensis* Highfield 1990 [= *Testudo graeca nabeulensis*], by subsequent designation by Fritz and Havaš (2007:295).

Etymology:* Latin, *furcula*, forked, and Greek, *chelys*, turtle; referring to the forked suprapygal.

Eurotestudo Lapparent de Broin, Bour, Parham, and Perälä 2006a:803

Type species: *Eurotestudo hermanni* [= *Testudo hermanni* Gmelin 1789], by original designation.

Etymology:* Toponym for Europe, and Latin, *testudo*, tortoise.

Testudo (Testudo) Linnaeus 1758 (07:72, 17:70)

Eurasian Tortoises, Mediterranean Tortoises
(includes 3 species, 13 taxa)

Testudo Linnaeus 1758:197
- Type species: *Testudo graeca* Linnaeus 1758 [= subjective synonym of *Testudo ibera* Pallas 1814], by subsequent designation by Fitzinger (1843:29) and Lindholm (1929:284); not *Testudo graeca* Auct. [= subjective synonym of *Testudo hermanni* Gmelin 1789] by earlier designation by Bell (1828c:514).
- Etymology:* Latin, *testudo*, turtle or tortoise.

Chersus Wagler 1830b:138
- Type species: *Chersus marginatus* [*Testudo marginata* Schoepff 1793a], by monotypy.
- Etymology:* Greek, *chersos*, tortoise.

Cherseus Gray 1856b:10 (*nomen novum*)
- Etymology:* Greek, *chersos*, tortoise.
- Comment: Unjustified emendation of *Chersus*.

Peltastes Gray 1869a:167,171 (junior homonym, not = *Peltastes* Illiger 1807 [= Hymenoptera] or *Peltastes* Agassiz 1838 [= Echinodermata] or *Peltastes* Fischer-Waldheim 1839 [= Orthoptera])
- Type species: *Testudo graeca* Linnaeus 1758 [= *Peltastes graecus* sensu Gray 1869] [= subjective synonym of *Testudo ibera* Pallas 1814], by subsequent designation by Lindholm (1929:286).
- Etymology:* Greek, *peltastes*, lightly armed foot soldiers in ancient Greece who wore light shields (*pelte*) made of twigs and leather; referring to the relatively thin shell bones in this genus.

Peltonia Gray 1872c:4 (*nomen novum*)
- Etymology:* Greek, *peltastes*, lightly armed foot soldiers in ancient Greece who wore light shields (*pelte*) made of twigs and leather; referring to the relatively thin shell bones in this genus.
- Comment: Unjustified replacement name for *Peltastes*.


Pseudotestudo Loveridge and Williams 1957:166, 276
- Type species: *Testudo (Pseudotestudo) kleinmanni* [= *Testudo kleinmanni* Lortet 1883], by original designation.
- Etymology:* Greek, *pseudo*, false, untrue, and Latin, *testudo*, turtle; referring to the difference of this genus from the genus *Testudo*.


Protestudo Chkhikvadze 1970:245
- Type species: *Protestudo bessarabica* [= *Testudo bessarabica* † Riabinin 1918], by original designation.
- Etymology:* Greek, *protos*, earliest, the first, and Latin, *testudo*, turtle; referring to this genus the ancestor of the genus *Testudo*.


Furculachelys Highfield 1990:1
- Type species: *Furculachelys nabeulensis* Highfield 1990 [= *Testudo graeca nabeulensis*], by subsequent designation by Fritz and Havaš (2007:295).
- Etymology:* Latin, *furcula*, forked, and Greek, *chelys*, turtle; referring to the forked suprapygal.

Testudo (Testudo) graeca Linnaeus 1758 (07:73, 11:12, 12:36, 14:35, 17:71, 17:72, 21:96) (85, 86)*

Spur-thighed Tortoise, Greek Tortoise, Moorish Tortoise
(includes 11 subspecies)


1 **


2 **


3 **

(subspecies [map #]: *graeca* = red [1], *armeniaca* = purple [3], *buxtoni* = dark blue [3], *cyrenaica* = orange [2], *ibera* = green [2,3], *marokkensis* = tourmaline [1], *nabeulensis* = gray [1], *terrestris* = brown [2,3], *tripolitania* = light blue [1], *whitei* = pink [1], *zarudnyi* = olive [3]; overlap = intergrades [2,3]; orange dots = uncertain, trade, or introduced)


Distribution:* Algeria, Armenia, Azerbaijan, Bulgaria, Georgia, Greece, Iran (*Alborz, Ardabil, Chahar Mahal Va Bakhtiari, East Azarbaijan, Fars, Gilan, Hamedan, Hormozgan, Ilam, Isfahan, Kerman, Kermanshah, Khorasan Razavi, Kohgiluyeh Va Boyer Ahmad, Kordestan, Lorestan, Markazi, Qazvin, Qom, Sistan Va Baluchestan, South Khorasan, Tehran, West Azarbaijan, Yazd, Zanjan*), Iraq, Israel, Jordan, Kosovo, Lebanon, Libya, Moldova, Morocco, North Macedonia, Palestine (*Gaza, West Bank*), Romania, Russia (*Dagestan, Krasnodar*), Serbia, Spain, Syria, Tunisia, Turkey (*Asian, European*), Turkmenistan (?)

Introduced:* Albania, Cyprus, Egypt, France, Greece (*Crete*), Italy (*Continental*, *Sardinia [prehistorical]*, *Sicily*), Malta (?), Spain (*Continental*, *Balearic Islands*), Ukraine
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 2,013,047 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 9,669,271 sq. km
Size (Max SCL):* male 46.0 cm, female 31.5 cm (see subspp.)
CBFTT Account:* (Western Subspecies Clade): Escoriza, Díaz-Paniagua, Andreu, and Ben Hassine (2022)
CBFTT Account:* (Eastern Subspecies Clades): Türkozan, Javanbakht, Mazanaeva, Meiri, Kornilev, Tzoras, Popgeorgiev, Shanas, and Escoriza (2023)
IUCN Red List:* Vulnerable (VU A1cd [ver. 2.3]) (TFTSG 1996). Regional (Europe): Near Threatened (NT) (Luiselli 2024). Previously: Vulnerable (VU A2bcde+4bcde) (Cox and Temple 2009); Vulnerable (VU A2bcde+4bcde) (van Dijk, Corti, Mellado, and Cheylan 2004). Regional (European Union–EU27): Vulnerable (VU A2bcde+A4bcde) (Cox and Temple 2009)
CITES: Appendix II, as Testudinidae spp. (1977). Previously: Appendix II, as *Testudo* spp. (1975)

Testudo (Testudo) graeca graeca Linnaeus 1758 (09:41, 21:96)
Souss Valley Tortoise


Uwe Fritz / foothills of Jbel Amsittene, Morocco


Andreas Nöllert / east of Taroudannt, Morocco / male


**

(subspecies: *graeca* = red, *marokkensis* = tourmaline, *whitei* = pink; orange dots = uncertain, trade, or introduced; stars = type localities)

Distribution: Morocco
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 66,771 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 136,365 sq. km
Size (Max SCL): male 19.5 cm, female 24.9 cm (Pieh 2001)
CBFTT Account:* (Western Subspecies Clade): Escoriza, Díaz-Paniagua, Andreu, and Ben Hassine (2022)
CITES: Appendix II, as Testudinidae spp. (1977). Previously: Appendix II, as *Testudo* spp. (1975)

Synonymy:

Testudo graeca Linnaeus 1758:198 (21:96)
Testudo graeca graeca, Peltastes graecus
Type locality: "Africa." Restricted to "Santa Cruz in Barbarie" by Houttuyn (1764:56); incorrectly emended to "Santa-Cruz (alte spanische Festung bei Oran in der Algérie)" [Algeria] [orange star] by Strauch (1862:67); correctly restricted to "Agadir in Morocco" [yellow star] by Schweiger and Gemel (2020:33).
Type specimen: Specimen figured in Edwards (1751:pl.204), holotype, designated and discussed by Bour (1987a); Thunberg (1828) listed a specimen in the UUZM acquired in the early 1800s from the MGA, possibly formerly in the MFA, but unlikely a type.
Etymology:* Toponym for Greece, or possibly Latin for Greek Tortoise (despite being described as coming from Africa), or possibly referring to the shell pattern reminiscent of Greek mosaic, or that the species was mentioned by Ancient Greek authors.
Comment: Description cited as sourced from Edwards (1751:pl.204). Type locality corrected by Schweiger and Gemel (2020), necessitating redefining the distributional range of the subspecies *graeca* from northeast Morocco through central Algeria to southwest Morocco.

Testudo pusilla Linnaeus 1758:199 (senior homonym, not = *Testudo pusilla* † Bergounioux 1936)
Chersine pusilla
Type locality:* "India." Restricted to "Cap. b. spei" [Cape of Good Hope, South Africa] by Linnaeus (1766:353); restricted to "Santa-Cruz (alte spanische Festung bei Oran in der Algérie)" [Algeria] [in error, orange star] by Strauch (1862:67) and "Santa Cruz, Oran" [Algeria] by Wermuth (1956:402); further restricted to "Agadir, Morocco" [yellow star] by TTWG (2021:298).
Type specimens: UUZM s/n, possible syntype, apparently lost, listed by Linné and Thunberg (1780), but not by Thunberg (1828), Lönnberg (1896), Andersson (1900), Holm (1957), or Wallin (2001).
Etymology:* Latin, *pusilla*, small, tiny, dwarfish; referring to the small size as compared to other tortoise species.
Comment: Description cited as sourced from Grew (1681:38,pl.3,f.3), Worm (1655:315), and Ray (1693:259), but cited by Linnaeus (1766:353) as also sourced from Edwards (1751:pl.204).

Testudo mauritanica Duméril and Bibron 1835:44 (21:96)
Testudo graeca mauritanica, Chersus mauritanicus, Peltastes mauritanicus, Testudo ibera mauritanica
Type locality:* "Mauritanie..[&]..les côtes occidentales de la mer Caspienne..[&]..les environs d'Alger." Restricted to "Umgebung von Algier" [Algeria] [in error, gray star] by Mertens and Wermuth (1955:378); further restricted to "Agadir, Morocco" [yellow star] by TTWG (2021:298).
Type specimens: Specimen figured in Edwards (1751:pl.204), lectotype, designated by Bour (1987a:111); USNM 10980 (formerly MNHN 22), originally a syntype, now a paralectotype, discussed by Cochran (1961) and Reynolds et al. (2007); MNHN 0.1937, invalid neotype, designated by Schweiger and Gemel (2020:36).
Etymology:* Toponym for Mauritania.


Testudo graeca soussensis Pieh 2001:211 (21:96)
Testudo soussensis, Testudo (Testudo) graeca soussensis
Type locality:* "Umgebung von Agadir 30°28'N, 9°55'W, Sousstal, Südwest-Marokko" [Morocco], geographic coordinates corrected here to 30°28'N, 9°35'W [green star, largely hidden by yellow star].
Type specimen: MTD 33842, holotype.
Etymology:* Toponym for the Souss Valley, Morocco.

Testudo (Testudo) graeca armeniaca Chkhikvadze and Bakradze 1991 (11:12, 21:97, 21:98)


Armenian Tortoise, Araxes Tortoise



Marine Arakelyan / Vanand region, Araxes valley, Armenia



Viacheslav Chkhikvadze / Dagestan, Russia / "*Testudo dagestanica*"



(subspecies: *armeniaca* = purple, *buxtoni* = dark blue, *ibera* = green, *terrestris* = brown; overlap = intergrades; orange dots = uncertain, trade, or introduced; stars = type localities)

Distribution: Armenia, Azerbaijan, Georgia, Iran (*Ardabil, East Azarbaijan, West Azarbaijan*), Russia (*Dagestan*), Turkey

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 111,897 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 253,884 sq. km

Size (Max SCL):* male 24.9 cm, female 27.1 cm (Pieh et al. 2002; Arakelyan et al. 2018; H. Javanbakht, unpubl. data; Türkozan et al. CBFTT 2023)

CBFTT Account:* (Eastern Subspecies Clades): Türkozan, Javanbakht, Mazanaeva, Meiri, Kornilev, Tzoras, Popgeorgiev, Shanas, and Escoriza (2023)

CITES: Appendix II, as Testudinidae spp. (1977). Previously: Appendix II, as *Testudo* spp. (1975)

Synonymy:

Testudo graeca pallasi Chkhikvadze 1989:67 (*nomen nudum*)
Etymology:* Eponym honoring Peter Simon Pallas (1741–1811), Germany / Russia; zoologist and naturalist at the Imperial Academy of Sciences in St. Petersburg.

Testudo graeca armeniaca Chkhikvadze 1989:67 (*nomen nudum*)
Etymology:* Toponym for Armenia.

Testudo graeca armeniaca Chkhikvadze and Bakradze 1991:60
Testudo armeniaca, Testudo terrestris armeniaca
Type locality: "Мегри, ЮВ Армянской ССР" [Megri, SE Armenian SSR] [Armenia] [yellow star].
Type specimen: GNM 13.3.007 (formerly ANGSSR), holotype.
Etymology:* Toponym for Armenia.

Testudo graeca pallasi Chkhikvadze and Bakradze 2002:276
Testudo pallasi, Testudo marginata pallasi
Type locality: "Дагестан, окрестности села Гиляры-Даг" [Dagestan, near the village of Gilyary-Dag]. Emended to "Gilyar, near Gilyary-Dag mountain, Magaramkent District, Dagestan, Russia (41.558499 N, 48.257204 E)" [green star] by TTWG (2017:152).
Type specimen: GNM NQ13-3-5 (formerly IPANG), holotype.
Etymology:* Eponym honoring Peter Simon Pallas (1741–1811), Germany / Russia; zoologist and naturalist at the Imperial Academy of Sciences in St. Petersburg.

Testudo dagestanica Chkhikvadze, Mazanaeva, and Shammakov 2011:337 (11:12)
Testudo graeca dagestanica
Type locality: "Папас (Южный Дагестан)" [Papas (Southern Dagestan)] [Russia] [blue star].
Type specimen: GNM 13.11.1, holotype.
Etymology:* Toponym for Dagestan, Russia.

Testudo (Testudo) graeca buxtoni Boulenger 1921 (10:28, 21:97, 21:98) (85)*


Zagros Tortoise, Buxton's Tortoise



Pavel Široký / Dehlili, Kermanshah Prov., Iran



Pavel Široký / *left*: Nahoj, Isfahan Prov., Iran, *right*: Siyahdare, Kermanshah Prov., Iran



(subspecies: *armeniaca* = purple, *buxtoni* = dark blue, *ibera* = green, *terrestris* = brown, *zarudnyi* = olive; overlap = intergrades; orange dots = uncertain, trade, or introduced; stars = type localities)

Distribution: Iran (*Alborz, Ardabil, Chahar Mahal Va*

Bakhtiari, East Azarbaijan, Fars, Gilan, Hamedan, Ilam, Isfahan, Kermanshah, Kordestan, Lorestan, Markazi, Qazvin, Tehran, West Azarbaijan, Zanjan), Turkey

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 405,933 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 739,181 sq. km

Size (Max SCL):* male 46.0 cm, female 30.0 cm (Perälä 2002c; Türkozan et al. 2010; Rezazadeh et al. 2014)

CBFTT Account:* (Eastern Subspecies Clades): Türkozan, Javanbakht, Mazanaeva, Meiri, Kornilev, Tzoras, Popgeorgiev, Shanas, and Escoriza (2023)

CITES: Appendix II, as Testudinidae spp. (1977). Previously: Appendix II, as *Testudo* spp. (1975)

Synonymy:


Testudo ecaudata Pallas 1814:19 (junior homonym and *nomen dubium*, not = *Testudo ecaudata* Daudin 1801 [= *Mauremys caspica*]) (10:28)
Type locality: "nemorosis Persiae mari caspio conterminis" [forests of Persia along the Caspian Sea] [Iran] [orange star, location estimated].
Type specimen: Not located; type specimen figured in an originally unpublished drawing by Pallas (1814:pl.3), reproduced in Darevsky and Mertens (1973:101), misidentified there as a *Psammobates tentorius*, but identified as a juvenile *Testudo graeca* ssp. by TTWG (2017:244).
Etymology:* Latin, *e-*, without, and *caudata*, having a tail, tailed; referring to the apparent lack of a tail on the examined specimen.


Testudo buxtoni Boulenger 1921:251
Testudo terrestris buxtoni, Testudo ibera buxtoni, Testudo graeca buxtoni
Type locality: "Manjil, between Resht and Kasuin, South Coast of the Caspian Sea, on a hill-side about 7,000–7,500 feet...northern Persia" [Iran] [yellow star].
Type specimen: NHMUK 1947.3.5.16 (formerly 1920.8.6.1), holotype.
Etymology:* Eponym honoring Patrick A. Buxton (1892–1955), United Kingdom / Iran; physician and entomologist who worked with the Royal Army Medical Corps in Persia (now Iran) and collected the type.

Testudo perses Perälä 2002c:81
Testudo graeca perses, Testudo ibera perses
Type locality: "vicinity of Lalabad village, some 25 mi NW Kermānshāh, Kermānshāhān province, W Iran...34°27'N 46°50'E" [green star].
Type specimen: FMNH 130820, holotype.
Etymology:* Toponym for Persia (now Iran).

Testudo (Testudo) graeca cyrenaica Pieh and Perälä 2002
Cyrenaican Tortoise, Cyrenaican Spur-thighed Tortoise


Christoph Schneider and Willi Schneider / Qasr Libya, Libya


Christoph Schneider and Willi Schneider / *left*: Ras al Hilal, Libya, *right*: Taknis, Libya


(subspecies: *cyrenaica* = orange, *ibera* = green, *terrestris* = brown, *tripolitania* = light blue; overlap = intergrades; orange dots = uncertain, trade, or introduced; star = type locality)

Distribution: Libya

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 23,138 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 44,624 sq. km

Size (Max SCL): male 19.3 cm, female 18.1 cm (Pieh and Perälä 2002)

CBFTT Account:* (Western Subspecies Clade): Escoriza, Díaz-Paniagua, Andreu, and Ben Hassine (2022)

CITES: Appendix II, as Testudinidae spp. (1977). Previously: Appendix II, as *Testudo* spp. (1975)

Synonymy:


Testudo graeca cyrenaica Pieh and Perälä 2002:8
Testudo cyrenaica
Type locality: "Derna 32°46'N, 22°39'E (= Darnah, Cyrenaika Ostlibyen" [Libya] [yellow star].
Type specimen: MTDK 31880, holotype.
Etymology:* Toponym for the Cyrenaica region of Libya.

Testudo (Testudo) graeca ibera Pallas 1814 (11:12, 21:97, 21:98)
Anatolian Tortoise, Greek Tortoise, Asia Minor Tortoise



Alexander A. Inozemtsev / CCB / TCF / nr. Novorossiysk, Krasnodar, Russia





Chris Leone / Ankara, Turkey / captivity



** (subspecies: *armeniaca* = purple, *buxtoni* = dark blue, *ibera* = green, *terrestris* = brown; overlap = intergrades; orange dots = uncertain, trade, or introduced; stars = type localities)

Distribution: Armenia, Azerbaijan, Bulgaria, Georgia, Greece, Kosovo, Moldova, North Macedonia, Romania, Russia (*Krasnodar*), Serbia, Turkey

Introduced:* Cyprus, Greece (*Crete*), Ukraine

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 550,805 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 1,663,554 sq. km

Size (Max SCL): male 38.9 cm, female 31.5 cm (Beshkov 1997; Auer and Herz 2006)

CBFTT Account:* (Eastern Subspecies Clades): Türkozan, Javanbakht, Mazanaeva, Meiri, Kornilev, Tzoras, Popgeorgiev, Shanas, and Escoriza (2023)

IUCN Red List: The currently synonymized taxon *Testudo graeca nikolskii* listed as Critically Endangered (CR A1abcde+2bcde [ver. 2.3]) (ERASG 1996)

CITES: Appendix II, as Testudinidae spp. (1977). Previously: Appendix II, as *Testudo* spp. (1975)

Synonymy:

Testudo ibera Pallas 1814:18
Chersus iberus, Cherseus iberus, Medaestia ibera, Testudo graeca ibera, Testudo ibera ibera, Testudo terrestris ibera
Type locality: "Caucasi meridionalibus..&..convallibus orae meridionalis montosae Chersonesi tauricae" [southern Caucasus ..&.. steep mountainous southern coast of Crimea]. Restricted to "Iberia, haud pro cul a Tiflisio" [Tbilisi, Georgia] by Eichwald (1831:196); to "das Gebiet des mittleren Kura-Tales im Kaukasus" [Georgia] by Mertens (1946:113); and to "Tbilissi (Tiflis), Géorgie, URSS" [Georgia] [yellow star] by Bour (1987a:112).
Type specimen: Not located; type specimen figured in an originally unpublished drawing by Pallas (1814:pl.2,f.2-3), discussed by Strauch (1862) and reproduced in Darevsky and Mertens (1973:100) and Bour (1987a:113), designated as lectotype by Bour (1987a:112).
Etymology:* Toponym for Iberia, a former ancient kingdom in Georgia in the Caucasus, not the Iberian Peninsula (= Spain).

Testudo georgicana Güldenstädt *in* Pallas 1814:18 (*nomen nudum*)
Etymology:* Toponym for Georgia in the Caucasus.

Testudo iberia Blyth 1854:642 (*nomen novum*) (17:66)
Etymology:* Toponym for Iberia, a former ancient kingdom in Georgia in the Caucasus, not the Iberian Peninsula (= Spain).
Comment: Unjustified emendation or error for *ibera*.

Testudo ibera bicaudalis Venzmer 1920:289
Type locality: "cilicischen Taurus in der kleinasiatischen Türkei" [Taurus Mts., Cilicia, Turkey] [green star, location estimated].
Type specimens: Not located, type specimen figured (f.1-2).
Etymology:* Latin, *bi-*, two, and *caudalis*, caudal scutes; referring to the two supracaudal marginal scutes.

Testudo ibera racovitzai Călinescu 1931:169
Type locality: "Turtucaia (jud. Durostor)" [Romania] [now Tutrakan, Silistra, Bulgaria] [blue star].
Type specimens: Not located, type specimen figured (f.6-7).
Etymology:* Eponym honoring Emil G. Racovitza (or Racoviță) (1868–1947), Romania; biologist, zoologist, and president of the Romanian Academy.


Testudo graeca nikolskii Chkhikvadze and Tuniyev 1986:618
Testudo ibera nikolskii, Testudo terrestris nikolskii, Testudo nikolskii
Type locality: "Поселок Небуг Туапсинского района (Краснодарский край)" [Nebug Settlement, Tuapse Co., Krasnodar District] [Russia] [pink star].
Type specimen: GNM 13.3.008 (formerly ANGSSR), holotype.
Etymology:* Eponym honoring Alexander M. Nikolsky (1858–1942), Russia / Soviet Union; naturalist and herpetologist at the Zoological Museum of the Academy of Sciences, St. Petersburg.

Testudo graeca pontica Khosatzky 1987:58 (*nomen nudum*)
Etymology:* Toponym for the Pontus region of northeastern Turkey, along the southeastern shores of the Black Sea.


Testudo (Testudo) graeca marokkensis Pieh and Perälä 2004 (09:42)
Moroccan Tortoise



Uwe Fritz / Sidi Bou Othmane, Morocco



Uwe Fritz / Sidi Bou Othmane, Morocco



**

(subspecies: *graeca* = red, *marokkensis* = tourmaline, *whitei* = pink; orange dot = uncertain, trade, or introduced; stars = type localities)

Distribution: Morocco

Introduced: Spain

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 69,188 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 99,260 sq. km

Size (Max SCL): male 17.2 cm, female 21.4 cm (Carretero et al. 2005; Ben Kaddour et al. 2008; Bonnet et al. 2010)

CBFTT Account:* (Western Subspecies Clade): Escoriza, Díaz-Paniagua, Andreu, and Ben Hassine (2022)

CITES: Appendix II, as Testudinidae spp. (1977). Previously: Appendix II, as *Testudo* spp. (1975)

Synonymy:


Testudo graeca marokkensis Pieh and Perälä 2004:22 (09:42)
Type locality:* "Tarmilete (33°23'N 6°04'W)" [= Tarmilate, Morocco] [yellow star].
Type specimen: SMNS 7602, holotype, see Bour (2004d).
Etymology:* Toponym for Morocco.

Testudo graeca lamberti Pieh and Perälä 2004:33 (09:42)
Type locality: "22 km nördlich von Tetuan (= Tétouan; Tétuan 35°34'N 5°22'W)" [Morocco] [green star].
Type specimen: NHMUK 1974.661, holotype, see Bour (2004d).
Etymology:* Eponym honoring Michael R.K. Lambert (1941–2004), United Kingdom; entomologist and herpetologist at the Natural Resources Institute, University of Greenwich.


Testudo (Testudo) graeca nabeulensis (Highfield 1990) (86)*
Tunisian Tortoise, Nabeul Tortoise



Tomáš Mazuch / Bou-Hedma National Park, Tunisia



Tomáš Mazuch / Bou-Hedma National Park, Tunisia



**

(subspecies: *cyrenaica* = orange, *nabeulensis* = gray, *tripolitania* = light blue, *whitei* = pink; orange dots = uncertain, trade, or introduced; star = type locality)

Distribution:* Algeria, Tunisia

Introduced:* Italy (*Sardinia*, *Sicily*), Malta

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 122,657 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 182,213 sq. km

Size (Max SCL):* male 13.7 cm, female 16.5 cm (Highfield and Martin 1989c; A. Highfield, unpubl. data, https://www.tortoisetrust.org/articles/tunisian.html)

CBFTT Account:* (Western Subspecies Clade): Escoriza, Díaz-Paniagua, Andreu, and Ben Hassine (2022)

CITES: Appendix II, as Testudinidae spp. (1977). Previously: Appendix II, as *Testudo* spp. (1975)

Synonymy:

Furculachelys nabeulensis Highfield 1990:32
Testudo nabeulensis, *Testudo graeca nabeulensis*
Type locality: "forested area in the region of Nabeul, Tunisia." Restricted to "Waldgebiet 7–8 km nordwestlich von Nabeul Richtung Grombalia (Tunesien)" [Tunisia] [yellow star] by Pieh and Perälä (2004:42).
Type specimen: TT s/n, not located, holotype figured (p.51).
Etymology:* Toponym for Nabeul, Tunisia.

Testudo graeca sarda Ballasina 1995:166 (*nomen nudum*)
Etymology:* Toponym for Sardinia, Italy.

Testudo graeca sardinia van der Kuyl, Ballasina, Dekker, Maas, Willemsen, and Goudsmit 2002:180 (*nomen nudum*)
Etymology:* Toponym for Sardinia, Italy.


Testudo (Testudo) graeca terrestris Forskål 1775 (10:29, 17:72)
Levantine Tortoise, Mesopotamian Tortoise



Norbert Halasz / Aleppo, Syria / captivity



Andreas Nöllert / Jabal an Nusayriyah, Latakia, Syria / captivity



** (subspecies: *armeniaca* = purple, *buxtoni* = dark blue, *ibera* = green, *terrestris* = brown; overlap = intergrades; orange dots = uncertain, trade, or introduced; stars = type localities)

Distribution: Iraq, Israel, Jordan, Lebanon, Palestine (*Gaza*, *West Bank*), Syria, Turkey

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 258,462 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 975,500 sq. km

Size (Max SCL):* male 28.0 cm, female 30.8 cm (Türkozan et al. 2010; Crucitti and Emiliani 2012)

CBFTT Account:* (Eastern Subspecies Clades): Türkozan, Javanbakht, Mazanaeva, Meiri, Kornilev, Tzoras, Popgeorgiev, Shanas, and Escoriza (2023)

CITES: Appendix II, as Testudinidae spp. (1977). Previously: Appendix II, as *Testudo* spp. (1975)

Synonymy:

Testudo terrestris Forskål 1775:viii,12 (junior homonym and *nomen protectum*, not = *Testudo terrestris* Garsault 1764 (*nomen oblitum*) [= *Emys orbicularis orbicularis*], nor = *Testudo terrestris* Fermin 1765 [= *Chelus fimbriata* (*nomen protectum*)]) (10:13)
Testudo graeca terrestris, Testudo terrestris terrestris, Testudo ibera terrestris
Type locality: "Lohajae..&..Kahirae..&..Aleppo..&..Libanon" [Al Luhayyah..&..Cairo..&..Aleppo..&..Mount Lebanon]. Restricted to "Arabien" [Saudi Arabia] by Wermuth (1956:402), to "Libanon-Gebirge, Israel" by Wermuth (1958:152), to "environs d'Alep (= Halab), Syrie" [Syria] by Bour (1989:14), and to "Aleppo (Alep, Halab; 36°12' N, 37°09' E), Syria (Syrian Arab Republic)" [yellow star] by Perälä and Bour (2004:117) by neotype designation.
Type specimen: Not designated, original syntypes apparently lost (Webb, pers. comm. *in* Iverson 1992); NMW 18674:2, neotype, designated by Perälä and Bour (2004:117), see Gemel et al. (2019).
Etymology:* Latin, *terrestris*, of the earth, terrestrial; referring to its terrestrial habitat.
Comment: Name conserved by ICZN (1963), see Wermuth (1956) and Mertens and Wermuth (1961).

Testudo zolhafa Forskål *in* Gray 1830e:5 (*nomen nudum*)
Etymology:* Vernacular (Arabic), *zolhafa*, *sulhufaa*, local indigenous name for the species.

Testudo floweri Bodenheimer 1935:197 (17:72)
Testudo graeca floweri, Testudo terrestris floweri, Testudo ibera floweri
Type locality: "the Negeb...Palestine" [Israel]. Restricted to "Negev, Palestine (environs de Gaza, Israël)" [Israel] [green star] by Bour (1989:14).
Type specimens: Not known or located.
Etymology:* Eponym honoring Stanley S. Flower (1871–1946), United Kingdom / Egypt; army officer, zoologist, and director of the Zoological Gardens at Giza, Egypt.

Testudo graeca anamurensis Weissinger 1987:14
Testudo ibera anamurensis, Testudo terrestris anamurensis, Testudo anamurensis
Type locality: "Strand von Anamurum, 7 km westlich von Anamur, SW–Küste der Türkei" [Turkey] [blue star].
Type specimen: NMW 30795:1, holotype, see Tiedemann et al. (1994) and Gemel et al. (2019).
Etymology:* Toponym for Anamur, Turkey.


Testudo antakyensis Perälä 1996:23;Appendix:16
Testudo graeca antakyensis, Testudo terrestris antakyensis, Testudo ibera antakyensis
Type locality:* "Antakyasta itään jäävä vuoristo Etelä-Turkissa"; "mountains to the east of Antakya in southern Turkey" [pink star, location estimated].
Type specimen: NHMUK 1998.70, holotype.
Etymology:* Toponym for Antakya, Turkey.

Testudo (Testudo) graeca tripolitania Poterala, Rattay, Johnson, Buijlayyil, Kiraz, and Ajaj 2024 (86)*


Tripolitanian Tortoise



Willi Schneider / near Tarhunah, Libya



Willi Schneider / near Gharyan, Libya



(subspecies: *cyrenaica* = orange, *nabeulensis* = gray, *tripolitania* = light blue, *whitei* = pink; orange dots = uncertain, trade, or introduced; star = type locality)

Distribution:* Libya

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 21,037 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 25,948 sq. km

Size (Max SCL):* male 14.3 cm, female 15.2 cm (Poterala et al. 2024)

CITES: Appendix II, as Testudinidae spp. (1977). Previously: Appendix II, as *Testudo* spp. (1975)

Synonymy:*

Testudo flavominimaralis Highfield and Martin 1989c:[9] (*nomen dubium*)

Testudo graeca flavominimaralis

Type locality: "North Africa." Restricted to "Libya" by Highfield (1990:41) [no star].

Type specimen: TT 89/07/AI, holotype, originally live, not located, specimen figured (pl.1).

Etymology:* Latin, *flavo*, yellow, golden yellow, and *minima*, smallest, very small; referring to the yellow carapace color and small body size.

Comment: Declared a *nomen dubium* by Pieh and Perälä (2002).

Testudo graeca tripolitania Poterala, Rattay, Johnson, Buijlayyil, Kiraz, and Ajaj 2024:117 (86)*

Type locality:* "Northwestern Libya." [yellow star, location estimated].

Type specimen:* FMNH 289176, originally US-I1-008, holotype, originally live, specimen figured (f.2).

Etymology:* Toponym for the Tripolitania region, Libya.

Testudo (Testudo) graeca whitei Bennett *in* White 1836 (21:96)


Mediterranean Spur-thighed Tortoise, Moorish Tortoise



Uwe Fritz / Murcia, Spain



Uwe Fritz / Murcia, Spain



(subspecies: *graeca* = red, *marokkensis* = tourmaline, *nabeulensis* = gray, *whitei* = pink; orange dots = uncertain, trade, or introduced; star = type locality)

Distribution: Algeria, Morocco, Spain

Introduced: Spain (*Balearic Islands*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 145,426 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 535,025 sq. km

Size (Max SCL): male 24.0 cm, female 28.0 cm (Highfield and Martin 1989b)

CBFTT Account:* (Western Subspecies Clade): Escoriza, Díaz-Paniagua, Andreu, and Ben Hassine (2022)

CITES: Appendix II, as Testudinidae spp. (1977). Previously: Appendix II, as *Testudo* spp. (1975)

Synonymy:

Testudo whitei Bennett *in* White 1836:361 (09:41)

Peltastes marginatus whitei, Testudo marginata whitei, Furculachelys whitei, Testudo graeca whitei

Type locality: Not known. Restricted to "S. Europe" by Gray (1873j:11) and to "Algiers and its environs, Algeria" [yellow star] by Highfield and Martin (1989b:21).

Type specimen: NHMUK 1853.4.17.1, holotype, discussed by Gray (1870b, 1870c, 1873j) and Highfield and Martin (1989b).

Etymology:* Eponym honoring Reverend Gilbert White (1720–1793), United Kingdom; naturalist and popular writer who obtained the type.

Comment: Type locality for the subspecies *graeca* corrected by Schweiger and Gemel (2020), necessitating redefining its distributional range to southwest Morocco and resurrecting *whitei* as the name for the subspecies from northeast Morocco through central Algeria.


Testudo (Testudo) graeca zarudnyi Nikolsky 1896 [(85)*]
Kerman Tortoise, Iranian Tortoise



Asghar Mobaraki / Saravan, Sistan Va Baluchestan, Iran / captivity



Asghar Mobaraki / Saravan, Sistan Va Baluchestan, Iran / captivity



** (subspecies: *buxtoni* = dark blue, *zarudnyi* = olive, overlap = intergrades; orange dots = uncertain, trade, or introduced; star = type locality)

Distribution:* Iran (*Fars, Hormozgan, Isfahan, Kerman, Khorasan Razavi, Sistan Va Baluchestan, South Khorasan, Yazd*), Turkmenistan (?)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 371,136 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 601,717 sq. km

Size (Max SCL):* male 24.7 cm, female 28.2 cm (Highfield and Martin 1989a; H. Javanbakht, unpubl. data; Türkozan et al. CBFTT 2023)

CBFTT Account:* (Eastern Subspecies Clades): Türkozan, Javanbakht, Mazanaeva, Meiri, Kornilev, Tzoras, Popgeorgiev, Shanas, and Escoriza (2023)

CITES: Appendix II, as Testudinidae spp. (1977). Previously: Appendix II, as *Testudo* spp. (1975)

Synonymy:

Testudo zarudnyi Nikolsky 1896:369

Testudo graeca zarudnyi, Testudo ibera zarudnyi, Testudo terrestris zarudnyi

Type locality: "Persia orientali" [Iran]. Restricted to "montibus provinciae Birdschan in Persiae orientali" [Iran] by Nikolsky (1897:308); and to "Birjand in Khorāsan province, NE Iran (32º53' N 59º03' E)" [yellow star] by Perälä (2002c:84).

Type specimen: ZIN 8738, holotype, erroneously designated lectotype by Perälä (2002c:84), discussed and genotyped by Parham et al. (2012).

Etymology:* Eponym honoring Nikolai A. Zarudny (1859–1919), Russia; zoologist, herpetologist, and explorer.


Testudo (Testudo) kleinmanni Lortet 1883 [(07:74, 17:73)]
Egyptian Tortoise



Basem Rabia / Zaranik Protected Area, Sinai, Egypt



Basem Rabia / Zaranik Protected Area, Sinai, Egypt / hatchling, adult



** (orange dots = probable trade; stars = type localities)

Distribution: Egypt, Israel, Libya, Palestine (*extirpated*) (*Gaza, West Bank*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 104,750 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 483,564 sq. km

Size (Max SCL): male 10.6 cm, female 14.4 cm (Farkas et al. 1997; Perälä 2001; Itescu et al. 2014)

IUCN Red List: Critically Endangered (CR A2abcd+3d) (Perälä 2003a). Previously: Endangered (EN) (TFTSG *in* Baillie and Groombridge 1996); the currently synonymized taxon *Testudo werneri* was previously listed separately as Critically Endangered (CR A2abcde+3de) (Perälä 2003b).

CITES: Appendix I (1995). Previously: Appendix II, as Testudinidae spp. (1977); Appendix II, as *Testudo* spp. (1975)

USA ESA:* Threatened (USFWS 2023a); Species Status Assessment: USFWS (2022a)

Synonymy:

Testudo leithii Günther 1869:502 (junior homonym, not = *Testudo leithii* † Carter 1852)

Peltastes leithii, Medaestia leithii

Type locality: "Sindh" [Pakistan] [in error] [no star].

Type specimen: NHMUK 1947.3.4.35 (formerly 1869.8.28.5), holotype.

Etymology:* Eponym honoring Andrew H. Leith (1827–1882), United Kingdom / India; physician, naturalist, and malacologist who worked in India on the Bombay Sanitary Commission and collected the type.

Testudo kleinmanni Lortet 1883:188

Pseudotestudo kleinmanni, Testudo kleinmanni kleinmanni

Type locality: "la basse Égypte, surtout dans les environs d'Alexandrie" [Egypt] [yellow star].

Type specimen: MHNL 42000414, lectotype, designated by Perälä (2001:579); SMF 7810, erroneously designated "lectotype" by Mertens (1967b:52), is not a type.

Etymology:* Eponym honoring Édouard Kleinmann (1832–1901), France / Egypt; banker, naturalist, and director of Crédit Lyonnais in Egypt, who collected the types.

Testudo werneri Perälä 2001:570 (07:74)

Testudo kleinmanni werneri

Type locality: "Northern Negev desert, Israel (14 km south of Be'er Sheva)" [green star].

Type specimen: HUJ 949, holotype.

Etymology:* Eponym honoring Yehudah L. Werner (1931–), Germany / Israel; zoologist and herpetologist at Hebrew University, Jerusalem.

Testudo (Testudo) marginata Schoepff 1793a (07:75, 11:12, 11:13)

Marginated Tortoise




Andreas Nöllert / Marathea, Peloponnese, Greece



Andreas Nöllert / Marathea, Peloponnese, Greece / *left*: female, *right*: male



**



**

(orange dots = probable trade or introduced or translocated; stars = type localities)

Distribution: Albania, Greece

Introduced:* Bulgaria, Cyprus, Greece (*Crete and various Ionian and Aegean islands*), Italy (*Continental*, *Sardinia [prehistorical]*), North Macedonia, Spain

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 64,496 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 135,660 sq. km

Size (Max SCL):* male 47.5 cm, female 42.0 cm (Bour 1996; Coutard 2006; Bringsøe et al. 2001; Escoriza et al. 2023 CBFTT); **(Max CCL):** female 50.0 cm (Ducotterd 1997)

CBFTT Account:* Escoriza, Tzoras, and Bringsøe (2023)

IUCN Red List:* Least Concern (LC) (Luiselli 2024). Previously: Least Concern (LC) (van Dijk, Lymberakis, and Böhme 2004); Least Concern (LC) [Not Listed] (Behler 1996)

CITES: Appendix II, as Testudinidae spp. (1977). Previously: Appendix II, as *Testudo* spp. (1975)

Synonymy:

Testudo tabulata campanulata Walbaum 1782:124 (unavailable name)

Etymology:* Latin, *campanula*, small bell, and *-atus*, provided with; referring to the flared posterior carapacial margin.

Comment: Unavailable name from a non-binomial work, see Wermuth (1956).

Testudo marginata Schoepff 1793a:52

Chersine marginata, Chersus marginatus, Peltastes marginatus, Peltastes marginata, Testudo marginata marginata

Type locality: Not known. Restricted to "Morea oggi Pelopponneso, Grecia" [Peloponnese, Greece] by Bruno (1986:238) [invalid restriction], and to "Greece, probably province of Attica, Stereá Eláda" [yellow star, location estimated] by lectotype designation by Bour (1987a:111).

Type specimen: MZUS 143, specimen figured in Schoepff (1792:pl.11-12,f.1) and Bour (1987a:f.20-22), designated lectotype by Bour (1987a:118); Thunberg (1828) listed a UUZM specimen from MGA and Wallin (2001) listed UUZM 278 (formerly MGA 47) as a type, possibly a paralectotype.

Etymology:* Latin, *marginis*, edge or border, and *-atus*, provided with, referring to the expanded carapacial margin.

Testudo graja Hermann *in* Schoepff 1793a:52

Type locality: Not designated. Restricted to "Griechenland" [Greece] by Mertens and Wermuth (1955:380).

Type specimen: MZUS 143, lectotype, designated by Bour (1987a:118), figured in Schoepff (1792:pl.11-12,f.1).

Etymology:* Toponym for Greece (Latin, *graia* = Greece).

Comment: Schoepff cited the author as Hermann, whose own description of *Testudo graja* was published posthumously (Hermann 1804:219).

Testudo campanulata Walbaum *in* Gray 1830e:4 (*nomen nudum*)

Etymology:* Latin, *campanula*, small bell, and *-atus*, provided with;

referring to the flared posterior carapacial margin.

Testudo graji Gray 1830e:4 (*nomen novum*)
Etymology:* Toponym for Greece (Latin, *graia* = Greece).
Comment: Unjustified emendation or error for *graja*.

Testudo campanulata Walbaum *in* Strauch 1862:65
Type locality: "Griechenland" [Greece].
Type specimens: Possibly ZIN, syntypes (2), not located.
Etymology:* Latin, *campanula*, small bell, and *-atus*, provided with; referring to the flared posterior carapacial margin.

Peltastes marginatus melas Gray 1870c:10
Type locality: "Greece."
Type specimen: NHMUK 1846.6.15.65, holotype.
Etymology:* Greek, *melas*, black; referring to the generally black carapace.

Testudo nemoralis Schreiber 1875:557
Type locality: "Griechenland und Süditalien" [Greece and Italy]. Restricted to "Griechenland" [Greece] by Mertens and Wermuth (1955:380).
Type specimens: Not located, type specimen figured (f.118-119).
Etymology:* Latin, *nemoralis*, woody, sylvan, coming from the forest; referring to the habitat.

Testudo marginata cretensis † Bachmayer, Brinkerink, and Symeonidis 1975:111
Type locality: "Gerani-Höhle an der Nordküste der Insel Kreta in der Nähe von Rethymnon" [Gerani Cave, Crete, Greece] [gray star].
Type specimen:* AMPG 3/1974, holotype, fossil, nearly whole shell, figured (pl.18-20); see also Vlachos (2022).
Geological age: Pleistocene, Neogenkalk [Neogene lime].
Etymology:* Toponym for Crete, Greece.

Testudo marginata sarda Mayer 1992:95
Type locality: "Olbia...Sardinia" [Italy] [orange star].
Type specimens: None designated.
Etymology:* Toponym for Sardinia, Italy.

Testudo marginata weissingeri Trutnau 1994:238 (*nomen suppressum* petitioned)
Type locality:* "Peleponnes" [Peloponnese, Greece].
Type specimen:* None designated.
Etymology:* Eponym honoring Heinz Weissinger (1945–1993), Austria; amateur herpetologist.
Comment:* Name mentioned prematurely and inadvertently with very brief characterization in a hobbyist book, prior to the delayed publication of the full scientific description of this taxon by Bour (1996). The ICZN has been petitioned to suppress this earlier name (Escoriza et al., in review).

Testudo weissingeri Bour 1996:30
Testudo marginata weissingeri
Type locality: "Kardamili, Messénie, Grèce" [Greece] [green star].
Type specimen: MNHN 1989.3317, holotype.
Etymology:* Eponym honoring Heinz Weissinger (1945–1993), Austria; amateur herpetologist.


Testudo (Agrionemys) Khosatzky and Młynarski 1966
(07:72, 09:40, 17:70)

Central Asian Tortoises
(includes 1 species, 5 taxa)

Testudinella Gray 1870c:12 (junior homonym, not = *Testudinella* Bory de Saint-Vincent 1822 [= Rotatoria])
Type species: *Testudinella horsfieldii* [= *Testudo horsfieldii* Gray 1844], by original monotypy.
Etymology:* Latin, *testudo*, turtle, and *-ella*, small or smaller.

Agrionemys Khosatzky and Młynarski 1966:123 (*nomen novum*)
Type species: *Agrionemys horsfieldi* [= *Testudo horsfieldii* Gray 1844], by original designation.
Etymology:* Greek, *agrios*, wild, as in living in a field, and *emys*, turtle; referring to the grassland habitat.
Comment: Justified replacement name for *Testudinella*.

Testudo (Agrionemys) horsfieldii Gray 1844 (07:77, 08:15, 09:43, 10:30, 14:38)
Central Asian Tortoise, Steppe Tortoise, Horsfield's Tortoise
(includes 5 subspecies and one taxonomically unspecified population of *T. (A.) horsfieldii* sensu lato)




** (subspecies: *horsfieldii* = red, *bogdanovi* = pink, *kazachstanica* = blue, *kuznetzovi* = brown, *rustamovi* = green, taxonomically unspecified population of *T. (A.) horsfieldii* sensu lato = gray, overlap [purple] = intergrades; orange dots = probable trade or translocated)

Distribution: Afghanistan, China (*Xinjiang*), Iran (*Golestan, Khorasan Razavi, North Khorasan, Semnan, South Khorasan*), Kazakhstan, Kyrgyzstan, Pakistan, Tajikistan, Turkmenistan, Uzbekistan
Introduced:* Azerbaijan, Latvia, Mongolia
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 2,122,553 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 4,029,696 sq. km
Size (Max SCL): male 19.0 cm, female 28.6 cm (see subspp.)
IUCN Red List: Vulnerable (VU A2d [ver. 2.3]) (TFTSG 1996)
CITES: Appendix II, as Testudinidae spp. (1977). Previously: Appendix II, as *Testudo* spp. (1975)

Testudo (Agrionemys) horsfieldii horsfieldii Gray 1844 (10:30)
Central Asian Tortoise, Steppe Tortoise, Horsfield's Tortoise



Salman Baloch / iNaturalist.org / Kalat, Baluchistan, Pakistan



**

(subspecies: *horsfieldii* = red, *rustamovi* = green, taxonomically unspecified population of *T. (A.) horsfieldii* sensu lato = gray; orange dots = probable trade or translocated; overlap [purple] = intergrades; stars = type localities)

Distribution: Afghanistan, Pakistan

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 96,572 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 191,321 sq. km

Size (Max SCL): female 21.1 cm (Annandale 1906)

CITES: Appendix II, as Testudinidae spp. (1977). Previously: Appendix II, as *Testudo* spp. (1975)

Synonymy:

Testudo horsfieldii Gray 1844:7

Testudinella horsfieldii, Homopus horsfieldii, Testudo (Homopus) horsfieldii, Medaestia horsfieldii, Agrionemys horsfieldii, Testudo horsfieldi, Testudo horsfieldii horsfieldii, Agrionemys horsfieldii horsfieldii

Type locality:* "India, Affghanistan [p.7]...Cabul [p.vi]." Restricted to "Afghanistan" by Günther (1864:7), and to "Kabul" [Afghanistan] [yellow star] by Iverson (1992:289).

Type specimen: NHMUK 1947.3.4.3, holotype.

Etymology:* Eponym honoring Thomas Horsfield (1773–1859), USA / Indonesia / India; physician and naturalist who worked in the Dutch East Indies (now Indonesia) and India and obtained the type.

Homopus burnesii Blyth 1854:642 (17:66)

Type locality: "Afghanistan" [yellow star, location estimated].

Type specimen: ZSI 793, holotype, see Das et al. (1998) and Kundu et al. (2018a).

Etymology:* Eponym honoring Sir Alexander Burnes (1805–1841), United Kingdom / Afghanistan; explorer, military officer, and diplomat in Afghanistan who obtained the type.

Testudo baluchiorum Annandale 1906:75 (10:28)

Agrionemys horsfieldii baluchiorum

Type locality: "Baluchistan" [Pakistan] [green star, location estimated].

Type specimen: ZSI 11420, holotype, see Das et al. (1998) and Kundu et al. (2018a).

Etymology:* Toponym for Baluchistan (= Balochistan), Pakistan.

Testudo hardwickei Cotton 1918:114 (*nomen nudum*)

Testudo hardwickii


Type locality: "Wano" [Wana, Khyber Pakhtunkhwa Prov., Pakistan] [blue star].

Etymology:* Eponym honoring Thomas Hardwicke (1756–1835), United Kingdom / India; military officer and naturalist who served with the East India Company in India.

Testudo (Agrionemys) horsfieldii bogdanovi Chkhikvadze *in* Chkhikvadze, Brushko, and Kubykin 2008 (10:30)
Fergana Valley Steppe Tortoise



Vyacheslav Yusupov / iNaturalist.org / Namangan, Uzbekistan



**

(subspecies: *horsfieldii* = red, *bogdanovi* = pink, *kazachstanica* = blue, *kuznetzovi* = brown, *rustamovi* = green, taxonomically unspecified population of *T. (A.) horsfieldii* sensu lato = gray, overlap [purple] = intergrades; stars = type localities)

Distribution: Kyrgyzstan, Tajikistan, Turkmenistan, Uzbekistan

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 158,196 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 258,595 sq. km

Size (Max SCL): male 12.4 cm, female 18.2 cm (Gnetneva 2020)

CITES: Appendix II, as Testudinidae spp. (1977). Previously: Appendix II, as *Testudo* spp. (1975)

Synonymy:

Agrionemys bogdanovi Chkhikvadze *in* Chkhikvadze, Brushko, and Kubykin 2008:100

Testudo horsfieldii bogdanovi, Agrionemys horsfieldii bogdanovi, Agrionemys bogdanovi

Type locality: "Узбекистан (Окрестности городов Бухара, Самарканд, Карши), Восточный Туркменистан (окрестности Чарджоу) и Киргизстан (Чуйская долина и окрестности города Ош)" [Uzbekistan (vicinity of the cities of Bukhara, Samarkand, Karshi), East Turkmenistan (near Chardzhou) and Kyrgyzstan (Chuy valley and vicinity of Osh)] [orange stars]. Restricted to "Ферганская долина" [Ferghana Valley] [Uzbekistan] [yellow star, location estimated] by Chkhikvadze et al. (2009:49).

Type specimen: GNM 13.4.60 (formerly GSM), holotype.


Etymology:* Eponym honoring Oleg P. Bogdanov (1925–2007), Soviet Union / Uzbekistan; herpetologist who collected the type.

Testudo (Agrionemys) horsfieldii kazachstanica Chkhikvadze 1988 (09:43, 10:30, 14:38)


Kazakhstan Steppe Tortoise



Haitao Shi / Huocheng Nature Reserve, Xinjiang, China



Haitao Shi / Huocheng Nature Reserve, Xinjiang, China



** (subspecies: *bogdanovi* = pink, *kazachstanica* = blue, *kuznetzovi* = brown, *rustamovi* = green, taxonomically unspecified population of *T. (A.) horsfieldii* sensu lato = gray, overlap [purple] = intergrades; orange dots = probable trade or translocated; stars = type localities)

Distribution: China (*Xinjiang*), Kazakhstan, Kyrgyzstan, Turkmenistan, Uzbekistan

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 1,213,129 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 2,180,040 sq. km

Size (Max SCL): male 18.9 cm, female 28.6 cm (Yakovleva 1961; Gnetneva 2020)

CITES: Appendix II, as Testudinidae spp. (1977). Previously: Appendix II, as *Testudo* spp. (1975)

Synonymy:

Agrionemys horsfieldi kazachstanica Chkhikvadze 1988:110

Testudo horsfieldii kazachstanica, Agrionemys horsfieldii kazachstanica, Agrionemys kazachstanica

Type locality: "Южное Прибалхашье, поселок Каратал" [southern Pribalkhashye (= Balkash), Karatal village]" [Kazakhstan] [yellow star, location estimated].

Type specimen: GNM 13.4.1, holotype.

Etymology:* Toponym for Kazakhstan.

Agrionemys kazachstanica terbishi Chkhikvadze 2009:60 (10:30, 14:38)

Testudo horsfieldii terbishi, Agrionemys horsfieldii terbishi


Type locality: "Монголия, г. Кобдо" [Mongolia, nr. Khovd] [orange star] [probable trade specimen].

Type specimen: PIK s/n, holotype, figured (p.60); apparently lost (Ansorge et al. 2012).

Etymology:* Eponym honoring Khayankhyarvaa Terbish (1952–), Mongolia; herpetologist at the National University of Mongolia who collected the type.

Testudo (Agrionemys) horsfieldii kuznetzovi Chkhikvadze, Ataev, Shammakov, and Zatoka *in* Chkhikvadze, Ataev, and Shammakov 2009 (10:30)

Turkmenistan Steppe Tortoise



** (subspecies: *bogdanovi* = pink, *kazachstanica* = blue, *kuznetzovi* = brown, *rustamovi* = green, taxonomically unspecified population of *T. (A.) horsfieldii* sensu lato = gray, overlap [purple] = intergrades; orange dot = probable trade or translocated; star = type locality)

Distribution: Turkmenistan, Uzbekistan

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 87,497 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 126,593 sq. km

Size (Max SCL): male 18.2 cm, female 22.4 cm (Gnetneva 2020)

CITES: Appendix II, as Testudinidae spp. (1977). Previously: Appendix II, as *Testudo* spp. (1975)

Synonymy:

Agrionemys kazachstanica kuznetzovi Chkhikvadze, Ataev, Shammakov, and Zatoka *in* Chkhikvadze, Ataev, and Shammakov 2009:52

Testudo horsfieldii kuznetzovi, Agrionemys horsfieldii kuznetzovi, Agrionemys kuznetzovi, Agrionemys kuznetsovi

Type locality: "впадины Акчакая (Северный Туркменистан, к западу от г. Дашогуз – бывш. Ташауз)" [Akchakai depression (North Turkmenistan, west of the city of Dashoguz – prev. Tashauz)] [yellow star].

Type specimen: GNM s/n, holotype, figured (f.4-6).


Etymology:* Eponym honoring Valentin V. Kuznetsov (1923–2006), Soviet Union / Kazakhstan; paleoherpetologist at the Institute of Zoology of Academy of Sciences of Kazakhstan.

Testudo (Agrionemys) horsfieldii rustamovi Chkhikvadze, Amiranashvili, and Ataev 1990 (09:43, 10:30)

Kopet-Dag Steppe Tortoise



Andreas Nöllert / Kara-Kum Desert, N. of Ashgabat, Turkmenistan



**

(subspecies: *horsfieldii* = red, *bogdanovi* = pink, *kazachstanica* = blue, *kuznetzovi* = brown, *rustamovi* = green, taxonomically unspecified population of *T. (A.) horsfieldii* sensu lato = gray, overlap [purple] = intergrades; orange dots = probable trade or translocated; star = type locality)

Distribution: Afghanistan, Iran (*Golestan, Khorasan Razavi, North Khorasan, Semnan*), Turkmenistan

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 550,888 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 825,160 sq. km

Size (Max SCL): male 23.4 cm, female 23.9 cm (Chkhikvadze et al. 2010)

CITES: Appendix II, as Testudinidae spp. (1977). Previously: Appendix II, as *Testudo* spp. (1975)

Synonymy:

Agrionemys horsfieldi rustamovi Chkhikvadze, Amiranashvili, and Ataev 1990:73

Agrionemys horsfieldii rustamovi, Testudo horsfieldii rustamovi, Agrionemys rustamovi

Type locality: "село Мадау (Кизыл-Атрекский район), Юго-Западный Туркменистан" [Madau village (Kizyl-Atrek region), south-western Turkmenistan] [yellow star].

Type specimen: GNM 13.4.88, holotype.

Etymology:* Eponym honoring Anver K. Rustamov (1917–2005), Soviet Union / Turkmenistan; zoologist, ornithologist, and herpetologist who worked in Turkmenistan.

Testudo (Chersine) Merrem 1820 (07:72, 09:40, 17:70)

European Tortoises

(includes 1 species, 2 taxa)

Chersine Merrem 1820:29 (09:40)

Type species: "*T. graeca* »L.« auct. = *T. hermanni* Gmelin" [= *Testudo hermanni* Gmelin 1789], by subsequent designation by Lindholm (1929:286).

Etymology:* Greek, *chersos*, tortoise.

Chersini Merrem *in* Gray 1825:210 (*nomen novum*)

Etymology:* Greek, *chersos*, tortoise.

Comment: Unjustified emendation of *Chersine*.

Medaestia Wussow 1916:170 (09:40)

Type species: *Medaestia graeca* sensu Wussow 1916 [= subjective synonym of *Testudo hermanni* Gmelin 1789] by subsequent designation by Mertens (1949:232). Genus established as *Testudo (Medaestia)* without a type species.

Etymology:* Anonym, *medaestia*, possibly a fabricated name, but possibly a Toponym for the Eastern Mediterranean region (*meda-* and *est*, east).

Comment: Genus referred to as *Testudo (Medaestia)* without indicating a type species, but noted as including *Medaestia graeca, M. ibera, M. leithii* [=*Testudo kleinmanni*], and *M. horsfieldii*.

Eurotestudo Lapparent de Broin, Bour, Parham, and Perälä 2006a:803


Type species: *Eurotestudo hermanni* [= *Testudo hermanni* Gmelin 1789], by original designation.

Etymology:* Toponym for Europe, and Latin, *testudo*, tortoise.

Testudo (Chersine) hermanni Gmelin 1789 (14:36, 14:37, 17:74, 21:99) (87)*

Hermann's Tortoise

(includes 2 subspecies)



**

(subspecies: *hermanni* = red, *boettgeri* = blue; orange dots = probably introduced or trade)

Distribution: Albania, Bosnia and Herzegovina, Bulgaria, Croatia, France (*Continental, Corsica*), Greece, Italy (*Continental, Sardinia [prehistorical introduction], Sicily*), Kosovo, Montenegro, North Macedonia, Romania, Serbia, Slovenia, Spain (*Continental, Balearic Islands [prehistorical introduction]*), Turkey (*European*)

Introduced: France, Malta (?), Spain (*Balearic Islands [prehistorical introduction]*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 383,025 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 1,926,808 sq. km

Size (Max SCL): male 31.4 cm, female 35.7 cm (see subspp.)

CBFTT Account: Bertolero, Cheylan, Hailey, Livoreil, and Willemsen (2011)

IUCN Red List:* Vulnerable (VU A2acde+4acde) (Luiselli 2024). Previously: Near Threatened (NT) (van Dijk, Corti, Mellado, and Cheylan 2004); Near Threatened (NT) (TFTSG *in* Baillie and Groombridge 1996)

CITES: Appendix II, as Testudinidae spp. (1977). Previously: Appendix II, as *Testudo* spp. (1975)

Testudo (Chersine) hermanni hermanni Gmelin 1789 (07:76, 14:36, 21:99, 21:100) (87)*

Western Hermann's Tortoise




Francesco Luigi Leonetti / Strongoli, Calabria, Italy





Albert Bertolero / CBFTT / Minorca, Spain



** (subspecies: *hermanni* = red, *boettgeri* = blue; orange dots = probably introduced or trade; stars = type localities)

Distribution: France (*Continental, Corsica*), Italy (*Continental, Sicily*), Spain (*Continental*)

Introduced: Italy (*Sardinia [prehistorical introduction]*), Spain (*Balearic Islands [prehistorical introduction]*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 136,705 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 900,797 sq. km

Size (Max SCL): male 19.6 cm, female 20.0 cm (Cheylan 2001)

IUCN Red List: Endangered (EN B1+2abcde [ver. 2.3]) (ERASG 1996)

CITES: Appendix II, as Testudinidae spp. (1977). Previously: Appendix II, as *Testudo* spp. (1975)

Synonymy:

Testudo hermanni Gmelin 1789:1041

Testudo hermanni hermanni, Protestudo hermanni, Agrionemys hermanni, Eurotestudo hermanni, Chersine hermanni, Chersine hermanni hermanni

Type locality: Not designated. Restricted to "Collobrières, massif de Maures, Var, France" [yellow star] by Bour (1987a:116).

Type specimen: MZUS 121, holotype, figured in Schoepff (1792:pl.8) and Bour (1987a:f.6-7), discussed by Bour (1987a, 2001).

Etymology:* Eponym honoring Johann Hermann (or Herrmann) (1738–1800), France; zoologist at the University of Strasbourg in Alsace, who first recognized this species as distinct.

Comment: Description cited as sourced from Schneider (1783:348), who cited notes by Hermann (pers. comm.).

Testudo graeca bettai Lataste 1881:396

Type locality:* Not known, but hypothesized as "une provenance plus occidentale que les îles Baléares, européenne (peninsule ibérique)" [orange star, location estimated].

Type specimen: NHMUK 1947.3.4.54 (formerly 1920.1.20.915), holotype, see Bour (1987a).

Etymology:* Eponym honoring Édoardo de Betta (1822–1896), Italy; herpetologist and politician who provided comparative material from Venice for the description of the new species.

Testudo hermanni robertmertensi Wermuth 1952:162

Type locality: "Gebirge nördlich von Arta, Mallorca, Balearen" [mountains north of Arta, Mallorca, Balearic Islands] [Spain] [green star].

Type specimen: SMF 37468, holotype, see Mertens (1967b).

Etymology:* Eponym honoring Robert F.W. Mertens (1894–1975), Russia / Germany; herpetologist and director of the Senckenberg Museum, Frankfurt.

Testudo (Chersine) hermanni boettgeri Mojsisovics 1889 (21:99)

Eastern Hermann's Tortoise




Adrian Hailey / CBFTT / Greece





Andreas Nöllert/ Marathea, Peloponnese, Greece



(subspecies: *hermanni* = red, *boettgeri* = blue; orange dots = probably introduced or trade; stars = type localities)

Distribution: Albania, Bosnia and Herzegovina, Bulgaria, Croatia, Greece, Italy (*Continental*), Kosovo, Montenegro, North Macedonia, Romania, Serbia, Slovenia, Turkey (*European*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 225,031 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 893,752 sq. km

Size (Max SCL): male 31.4 cm, female 35.7 cm (Beshkov 1997; Cheylan 2001)

CITES: Appendix II, as Testudinidae spp. (1977). Previously: Appendix II, as *Testudo* spp. (1975)

Synonymy:

Testudo graeca boettgeri Mojsisovics 1889:242 (senior homonym, not = *Testudo boettgeri* Siebenrock 1904a [= *Psammobates tentorius verroxii*])

Testudo hermanni boettgeri, Testudo boettgeri, Testudo boettgeri boettgeri, Eurotestudo boettgeri, Chersine hermanni boettgeri

Type locality: "Orsova...Cernathal...Süd-Ungarn" [Romania]. Restricted to "Orsova, Banat" [Romania] [yellow star] by Boettger (1893:11).

Type specimen: SMF 7836, lectotype, designated as the type by Boettger (1893:11), listed as "holotype" by Mertens (1967b), see Bour (1987a); specimen figured in Bour (1987a:f.12-14).

Etymology:* Eponym honoring Oskar Boettger (or Böttger) (1844–1910), Germany; paleontologist, herpetologist, and curator at the Senckenberg Museum, Frankfurt.

Testudo graeca hercegovinensis Werner 1899:818

Testudo hercegovinensis, Testudo boettgeri hercegovinensis, Testudo hermanni hercegovinensis, Eurotestudo hercegovinensis

Type locality: "Trebinje...Hercegovina" [Bosnia and Herzegovina] [green star].

Type specimen: NMW 1222, lectotype, designated by Bour (1987a:116), see Tiedemann et al. (1994) and Gemel et al. (2019).

Etymology:* Toponym for Herzegovina.

Testudo enriquesi Parenzan 1932:1160

Type locality: "Conca di Elbassan, nella vallata dello Skumbi, in Albania Centrale" [Elbassan basin, in the Skumbi valley, in Central Albania] [blue star].

Type specimen: Not located, holotype figured (pl.1-4), see Bour (1987a).

Etymology:* Eponym honoring Paolo Enriques (1878–1932), Italy; zoologist at the University of Padua who mentored the author.

Infraorder Trionychia Zittel 1889 (21:3)

Softshell Hidden-necked Turtles

(includes 2 families, 14 genera, 36 species, 48 taxa)

Potamites Duméril and Bibron 1834:353

Diacostoidea Baur 1887:99

Trionychia Zittel 1889:513

Chilotae Baur 1890c:536

Superfamily Trionychoidea Gray 1825

Softshell Hidden-necked Turtles

(includes 2 families, 14 genera, 36 species, 48 taxa)

Trionicidae Gray 1825:212

Trionychoidea Fitzinger 1826:5

Trionychia Hümmel 1929:362

Family Carettochelyidae Boulenger 1887a

Pig-nosed Turtles

(includes 1 genus, 1 species, 1 taxon)

Carettochelydidae Boulenger 1887a:171

Carettochelydes Baur 1891a:190

Carettochelyidae Baur 1891c:637



Carettochelyidae Species Richness

Carettochelys Ramsay 1886

Pig-nosed Turtles

(includes 1 species, 1 taxon)

Carettochelys Ramsay 1886:158

Type species: *Carettochelys insculpta* Ramsay 1886, by original monotypy.

Etymology:* Vernacular (Old French), *caret*, *Caretta*, sea turtle, and Greek, *chelys*, turtle; referring to the genus as representing an apparent link between river turtles and sea turtles.


Carettochelys insculpta Ramsay 1886 (07:78, 08:10) (88)*
Pig-nosed Turtle, Fly River Turtle



John Cann / CBFTT / Daly R., Northern Territory, Australia



left: Fred Parker / Balimo, Western Prov., Papua New Guinea / hatchling
right: Philip M. Hall / Lake Murray, Western Prov., Papua New Guinea



** (orange dot = probably introduced or trade; stars = type localities)

Distribution: Australia (*Northern Territory*), Indonesia (*Papua*), Papua New Guinea (*Southern*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 149,082 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 983,664 sq. km

Size (Max SCL): male 41.3 cm, female 52.5 cm (Georges et al. 2006; Ceballos et al. 2013; P.M. Hall, unpubl. data); **(Max CCL):** male 53.0 cm [estimated], female 61.0 cm (Georges and Kennett 1989; Eisemberg et al. 2011)

CBFTT Account: Georges, Doody, Eisemberg, Alacs, and Rose (2008)

IUCN Red List: Endangered (EN A2bd+4bd) (Eisemberg, van Dijk, Georges, and Amepou 2018). Previously: Vulnerable (VU) (ATTWG 2000); Vulnerable (VU) (TFTSG *in* Baillie and Groombridge 1996)

CITES: Appendix II (2005)


Synonymy:

Carettochelys insculptus Ramsay 1886:158
Carettocchelys insculptus, Carettochelys insculpta, Carettochelys insculpta insculpta
Type locality: "Fly River, New Guinea." Restricted to "Strickland River...upper right hand branch of the Fly River" by Waite (1905:110); and to "Turtle Camp, Strickland River [Papua New Guinea] [6º25'30" S, 142º04'30" W]" [yellow star] by Mackay (2003:39).
Type specimen: AMS R3677, holotype, see Cogger (1979), Cogger et al. (1983), Shea and Sadlier (1999), and Cann and Sadlier (2017).
Etymology:* Latin, *insculptus*, etched or carved; referring to the sculpturing of the carapace.

Carettochelys insculpta canni Wells 2002a:1 (unavailable name) (07:78, 08:10, 10:43)
Carettochelys canni
Type locality: "near Ooloo Crossing, Daly River, Northern Territory [Australia]" [green star].
Type specimen: NTM 1213, holotype.
Etymology:* Eponym honoring John R. Cann (1938–), Australia; herpetologist who studied turtles extensively in Australia and New Guinea.

Family Trionychidae Gray 1825

Softshell and Flapshell Turtles
(includes 2 subfamilies, 13 genera, 35 species, 47 taxa)
Amydae Oppel 1811:9 (*partim*)
Trionyces Schmid 1819:18
Trionicidae Gray 1825:212
Trionychoidea Fitzinger 1826:5
Trionychidae Bell 1828c:515
Trionycidae Bonaparte 1831:63
Trionychides Pictet 1853:455
Trionychidi Portis 1890:22



Trionychidae Species Richness

Subfamily Cyclanorbinae Lydekker 1889

Flapshell Turtles
(includes 3 genera, 7 species, 9 taxa)
Cyclanosteina Gray 1864b:94
Cyclanorbinae Lydekker 1889:x
Cyclanorbidae Deraniyagala 1939:290

Cyclanorbis Gray 1854a (25)*

Sub-Saharan Flapshell Turtles
(includes 2 species, 2 taxa)
Cryptopus Duméril and Bibron 1835:499 (junior homonym, not = *Cryptopus* Latreille 1829 [= Crustacea])
Type species: *Cryptopus senegalensis* Duméril and Bibron 1835, by subsequent designation by Bour et al. (1995:82).
Etymology:* Greek, *krypto*, to hide, and *pus*, foot, leg; referring to the ability to hide the legs under the plastral cutaneous flaps.
Cyclanorbis Gray 1854a:135
Type species: *Cyclanorbis petersii* Gray 1854a [= subjective synonym of *Cryptopus senegalensis* Duméril and Bibron 1835], by original monotypy.
Etymology:* Greek, *kyklos*, circle, ring, and Latin, *orbis*, circle, orb; referring to the rounded oval carapace.
Comment:* The name *Cyclanorbis* credited to W.K.H. Peters by Gray.
Cyclanosteus Gray 1856a:201
Type species: *Cyclanosteus senegalensis* [= *Cryptopus senegalensis* Duméril and Bibron 1835], by subsequent designation by Günther (1865:108).
Etymology:* Greek, *kyklos*, circle, ring, and *an-*, without, and *osteon*, bone; referring to the lack of bones in the flexible carapace margin.
Cryptopodus Duméril 1856:374 (*nomen novum*)
Etymology:* Greek, *krypto*, to hide, and *pus*, foot, leg; referring to

the ability to hide the legs under the plastral cutaneous flaps.
Comment:* Justified replacement name for *Cryptopus*.

Tetrathyra Gray 1865a:205
Type species: *Tetrathyra baikii* Gray 1865a [= subjective synonym of *Cryptopus senegalensis* Duméril and Bibron 1835], by original monotypy.
Etymology:* Greek, *tetra*, four, and *thyreos*, shield, door stone; referring to the four plastral callosities (two pairs).


Baikiea Gray 1869a:215
Type species: *Baikiea elegans* Gray 1869a, by original monotypy.
Etymology:* Eponym honoring William B. Baikie (1824–1864), United Kingdom; explorer of Africa who obtained the types.

Cyclanorbis elegans (Gray 1869a)
Nubian Flapshell Turtle


Luca Luiselli / White Nile, South Sudan



Maurice Rodrigues / No data / captivity


** (stars = type localities)

Distribution:* Benin (?), Cameroon, Central African Republic, Chad, Equatorial Guinea; Ethiopia, Ghana, Nigeria, South Sudan, Sudan, Togo, Uganda

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 296,634 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 4,284,609 sq. km

Size (Max SCL):* male 67.6 cm, female 70.0 cm (Gramentz 2008; Baker et al. 2015 CBFTT); **(Max CCL):*** female 73.5 cm (Luiselli et al. 2022; pers. comm.).

CBFTT Account: Baker, Diagne, and Luiselli (2015)

IUCN Red List: Critically Endangered (CR A2bcd+4bcd) (Baker, Luiselli, and Diagne 2016). Previously: Near Threatened (NT) (TFTSG 1996)

CITES: Appendix II (2017)

Synonymy:

Baikiea elegans Gray 1869a:222
Cyclanorbis elegans, Cyclanosteus elegans
Type locality: "Africa." Restricted to "West Africa" by Gray (1873j:86); and to "Niger River drainage in west Africa" [yellow star, location estimated] by Webb (1975a:349).
Type specimen: NHMUK 1946.1.22.15 (formerly 1865.5.3.35), lectotype, designated by Webb (1975a:348).
Etymology:* Latin, *elegans*, elegant, fine, select; referring to the fine spotted markings on hatchlings.


Cyclanorbis oligotylus Siebenrock 1902c:810
Type locality: "Nubien...oberen Nil" [Nubia...upper Nile] [Sudan] [green star, location estimated].
Type specimens: NMW 1682, 1848, syntypes (2), see Tiedemann et al. (1994) and Gemel et al. (2019).
Etymology:* Greek, *oligos*, few, minor, little, and *tylos*, callus, bulge; referring to the fewer plastral callosities in this taxon.

Cyclanorbis senegalensis (Duméril and Bibron 1835) (17:75)
Sahelian Flapshell Turtle, Senegal Flapshell Turtle


William R. Branch / Northern Prov., Sierra Leone



Pearson McGovern / Senegal


** (orange dots = probable trade; stars = type localities)

Distribution: Benin, Burkina Faso, Cameroon, Central African Republic, Chad, Ethiopia, Gambia, Ghana, Guinea, Guinea-Bissau, Ivory Coast (Côte d'Ivoire), Liberia, Mali, Mauritania, Niger, Nigeria, Senegal, Sierra Leone, South Sudan, Sudan, Togo, Uganda (?)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 1,949,980 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 7,374,057 sq. km

Size (Max SCL): male 18.7 cm, female 35.5 cm (Pritchard 2001; Gramentz 2008); **(Max CCL):** female 37.9 cm (McGovern et al. 2021 CBFTT)

CBFTT Account: McGovern, Diagne, Diagne, Luiselli, and

Meylan (2021)

IUCN Red List: Vulnerable (VU A2bcd+4bcd) (Diagne, Luiselli, Trape, Rödel, Baker, Chirio, Petrozzi, and Segniagbeto 2016). Previously: Near Threatened (NT) (TFTSG 1996)

CITES: Appendix II (2017)

Synonymy:

Cryptopus senegalensis Duméril and Bibron 1835:504
Emyda senegalensis, Cyclanosteus senegalensis, Cyclanorbis senegalensis
Type locality: "Sénégal" [yellow star, location estimated].
Type specimen: MNHN 4151, holotype, discussed Bour et al. (1995).
Etymology:* Toponym for Senegal.

Cyclanorbis petersii Gray 1854a:135
Cyclanosteus petersii, Cycloderma petersii
Type locality: "West Africa, River Gambia" [The Gambia] [green star, location estimated].
Type specimen: NHMUK 1947.3.6.23, holotype.
Etymology:* Eponym honoring Wilhelm K.H. Peters (1815–1883), Germany; zoologist, herpetologist, and curator at the Museum für Naturkunde, Berlin.

Cycloderma senegalense Duméril 1861a:168 (*nomen novum*)
Etymology:* Toponym for Senegal.
Comment: Unjustified emendation for *senegalensis*.

Tetrathyra baikii Gray 1865a:205
Type locality: "West Africa, River Niger?" Restricted to "Niger River drainage in west Africa" [blue star, location estimated, possibly near Lokoja, Nigeria] by Webb (1975a:350).
Type specimen: NHMUK 1947.3.6.25 (formerly 1865.9.13.1), lectotype, designated by Webb (1975a:349).
Etymology:* Eponym honoring William B. Baikie (1824–1864), United Kingdom; explorer of Africa and Nigeria, who obtained the types.

Cyclanosteus senegalensis equilifera Gray 1865b:425
Type locality: "the Niger" [Niger River, West Africa] [blue star, location estimated, possibly near Lokoja, Nigeria].
Type specimen: NHMUK, holotype, not identified, figured (f.2) and in Gray (1870:f.38a), see Gray (1873i:87).
Etymology:* Latin, *aequus, equi-*, equal, and *ferre*, to bear; referring to the arrangement and number of the plastral callosities.

Cyclanosteus senegalensis normalis Gray 1865b:425
Type locality: "the Niger" [Niger River, West Africa] [blue star, location estimated, possibly near Lokoja, Nigeria].
Type specimen: NHMUK number?, holotype, not identified.
Etymology:* Latin, *normalis*, normal; referring to the arrangement and number of the plastral callosities.

Cyclanosteus senegalensis callosa Gray 1865b:425
Type locality: "the Niger" [Niger River, West Africa] [blue star, location estimated, possibly near Lokoja, Nigeria].
Type specimen: NHMUK 1865.5.3.75, holotype, figured (f.1) and in Gray (1870:f.38), see Gray (1873i:87).
Etymology:* Latin, *callum*, callus; referring to the arrangement and number of the plastral callosities.

Tetrathyra vaillantii Rochebrune 1884:36
Type locality: "Sénégambie...Podor, Dagana, Saldé, Leybar, Thiouk, Maringouins, Babagaye, Sorres, Diouk" [Senegal and The Gambia] [pink star = Podor, Senegal].
Type specimens: Not located, syntypes, two figured (pl.4,f.1-2).
Etymology:* Eponym honoring Léon Louis Vaillant (1834–1914), France; vertebrate zoologist at the Muséum National d'Histoire Naturelle, Paris.

Cycloderma Peters 1854

Central African Flapshell Turtles
(includes 2 species, 2 taxa)

Cycloderma Peters 1854:216
Type species: *Cycloderma frenatum* Peters 1854, by original monotypy.
Etymology:* Greek, *kyklos*, circle, ring, and *derma*, skin; referring to the rounded carapace covered with skin instead of scutes.

Heptathyra Cope 1860:294
Type species: *Heptathyra aubryi* [= *Cryptopodus aubryi* Duméril 1856], by original monotypy.
Etymology:* Greek, *hepta*, seven, and *thyreos*, shield, door stone; referring to the seven plastral callosities.

Aspidochelys Gray 1860a:6
Type species: *Aspidochelys livingstonii* Gray 1860a [= subjective synonym of *Cycloderma frenatum* Peters 1854], by original monotypy.
Etymology:* Greek, *aspidos*, a shield, and *chelys*, turtle; referring to the shield-like appearance of its shell.

Cycloderma aubryi (Duméril 1856) (10:31)


Aubry's Flapshell Turtle



Olivier S.G. Pauwels / Gamba, Ogooué-Maritime Prov., Gabon



left: Laurent Chirio / Lambarene, Gabon
right: James H. Harding / No data / captivity



** (orange dot = probable trade; star = type locality)

Distribution: Angola (*Cabinda*), Cameroon, Central African Republic, Congo (DRC), Congo (ROC), Gabon

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 571,218 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 1,566,358 sq. km


Size (Max SCL): male 40.0 cm, female 50.7 cm; **(Max CCL):** male 46.6 cm, female 61.0 cm (Gramentz 1998; Maran 2006a; Gramentz 2008)

IUCN Red List: Vulnerable (VU A2bcd+4bcd) (Chirio, Diagne, and Pauwels 2017). Previously: Least Concern (LC) [Not Listed] (Behler 1996)

CITES: Appendix II (2017)
Synonymy:
Cryptopodus aubryi Duméril 1856:374 (10:31)
Cryptopus aubryi, Cycloderma aubryi, Heptathyra aubryi
Type locality: "Gabon." Restricted to "mouth of the Gabon river at present-day Libreville" by Bour 2008d:41 [yellow star].
Type specimen: MNHN 8006, holotype, discussed by Bour et al. (1995) and Bour (2008d).
Etymology:* Eponym honoring Charles E. Aubry-Lecomte (1821–1879), France / Gabon; civil servant and naturalist who worked in Gabon and collected the types.

Cycloderma frenatum Peters 1854
Zambezi Flapshell Turtle


Martin Grimm / Rufiji R., Tanzania



Wulf Haacke / CBFTT / Lake Malawi, Malawi / adult, juvenile


**
(orange dot = possibly introduced; star = type localities)
Distribution: Malawi, Mozambique, Tanzania, Zambia, Zimbabwe
Introduced:* Tanzania (*Zanzibar*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 250,649 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 1,283,943 sq. km
Size (Max SCL): male 46.5 cm, female 49.0 cm [estimated]; **(Max CCL):** male 56.0 cm, female 58.9 cm (Gray 1860a; Peters 1882; Gramentz 2008; Broadley and Sachsse 2011 CBFTT)
CBFTT Account: Broadley and Sachsse (2011)
IUCN Red List: Endangered (EN A3d) (van Dijk 2016). Previously: Near Threatened (NT) (TFTSG 1996)

CITES: Appendix II (2017)
Synonymy:
Cycloderma frenatum Peters 1854:216
Cyclanosteus frenatus, Heptathyra frenata
Type locality: "In fluminibus Zambeze et Licuare..[&]..Tette et Sena..[&]..terra Boror" [Mozambique]. Restricted to "Sambesi bei Tete" [Zambezi River, Tete, Mozambique] [yellow star] by Fritz et al. (1994:169).
Type specimen: ZMB 4815, lectotype, designated by Fritz et al. (1994:169); Uetz et al. (2019) listed ZMB 47, 49–50, 4815, 8243, and 8432 as syntypes.
Etymology:* Latin, *frenum*, reins, bridle, *frenatum*, having reins; referring to the black line markings along the sides of the neck and between the eyes.
Aspidochelys livingstonii Gray 1860a:6
Heptathyra livingstonii
Type locality: "Mozambique in tributaries of River Zambesi?" [yellow star, location estimated].
Type specimen: NHMUK 1947.3.4.16, holotype.
Etymology:* Eponym honoring David Livingstone (1813–1873), United Kingdom (Scotland); physician, missionary, and explorer of Africa, who collected and presented the type.

Lissemys Smith 1931 (11:14)

Indian Flapshell Turtles

(includes 3 species, 5 taxa)

Emyda Gray 1830e:19 (junior homonym, not = *Emyda* Rafinesque 1815 [= *Emys*]) (10:7)

Type species: *Emyda punctata* [= *Testudo punctata* Lacepède 1788 (*nomen suppressum*) = *Testudo punctata* Bonnaterre 1789], by original monotypy.

Etymology:* Greek, *emydos*, turtle.

Lissemys Smith 1931:xxviii,154 (*nomen novum*)

Type species: *Lissemys punctata* [= *Testudo punctata* Lacepède 1788 (*nomen suppressum*) = *Testudo punctata* Bonnaterre 1789], by original designation.

Etymology:* Greek, *lissos*, smooth, and *emys*, turtle; referring to the shell bones covered with a smooth skin.


Comment: Justified replacement name for *Emyda*.

Lissemys ceylonensis (Gray 1856a) (11:14)


Sri Lankan Flapshell Turtle



Anslem de Silva / Udawalawe National Park, Sri Lanka



Anslem de Silva / Udawalawe National Park, Sri Lanka



(stars = type localities)

Distribution: Sri Lanka

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO): 62,568 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 74,759 sq. km

Size (Max SCL): female 37.0 cm (Deraniyagala 1939; Webb 1982; Itescu et al. 2014)

IUCN Red List: Vulnerable (VU A2d) (Praschag, de Silva, and de Silva 2021)

CITES: Appendix II (1995), as part of *Lissemys punctata*

Synonymy:

Emyda ceylonensis Gray 1856a:201

Emyda granosa ceylonensis, Lissemys ceylonensis, Lissemys punctata ceylonensis

Type locality: "Ceylon" [Sri Lanka] [yellow star, location estimated, possibly nr. Colombo].

Type specimens: NHMUK 1854.3.21.18, 1947.3.4.17, syntypes (2); 1947.3.4.17 listed as "holotype" by Webb (1980).

Etymology:* Toponym for Ceylon [Sri Lanka].

Lissemys punctata sinhaleyus † Deraniyagala 1953:5

Type locality: "near Ratnapura...Ceylon" [Sri Lanka] [green star].

Type specimen: RNM F283, holotype, fossil, hypoplastron.


Geological age: Late Pleistocene, Ratnapura Beds.

Etymology:* Toponym for Sinhala [Sanskrit name for Ceylon = Sri Lanka].

Lissemys punctata (Bonnaterre 1789) (09:44, 11:14, 21:101)

Indian Flapshell Turtle

(includes 3 subspecies)



**

(subspecies: *punctata* = red, *andersoni* = blue, *vittata* = green; overlap = intergrades; orange dots = probably introduced or translocated or trade)

Distribution:* Bangladesh, India (*Andhra Pradesh, Arunachal Pradesh, Assam, Bihar, Chhattisgarh, Goa, Gujarat, Haryana, Jammu, Jharkhand, Kerala, Madhya Pradesh, Meghalaya, Odisha, Punjab, Tamil Nadu, Sikkim [?], Telangana, Tripura, Uttarakhand, Uttar Pradesh, West Bengal*), Myanmar, Nepal, Pakistan

Introduced:* India (*Andaman Islands*), Singapore

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 2,239,325 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 5,352,100 sq. km

Size (Max SCL): male 23.0 cm, female 37.0 cm (see subspp.)

CBFTT Account: Bhupathy, Webb, and Praschag (2014)

IUCN Red List: Vulnerable (VU A2cd+4cd) (Rahman, Ahmed, Choudhury, Praschag, and Singh 2021). Previously: Least Concern (LC) (ATTWG 2000); Least Concern (LC) [Not Listed] (Behler 1996)

CITES: Appendix II (1995). Previously: Appendix I (1975)


Lissemys punctata punctata (Bonnaterre 1789) (11:14, 21:101)
Southern Indian Flapshell Turtle



Indraneil Das / CBFTT / Mamallapuram, Tamil Nadu, India



left: Indraneil Das / CBFTT / Mamallapuram, Tamil Nadu, India
right: Peter Praschag / CBFTT / Moyar R., Karnataka-Tamil Nadu border, India / juvenile



** (subspecies: *punctata* = red, *vittata* = green; orange dot = probably introduced, translocated, or trade, stars = type localities)

Distribution:* India (*Karnataka, Kerala, Tamil Nadu*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 153,098 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 258,762 sq. km

Size (Max SCL): No data located; Mean CL (SCL or CCL?): female 18.2 cm (Premkishore and Chandran 1996)

CITES: Appendix II (1995). Previously: Appendix I (1975)

Synonymy:

Testudo punctata Lacepède 1788:171, synopsis[table] (senior homonym and *nomen suppressum*, not = *Testudo punctata* Schoepff 1792 (*nomen suppressum*) [= *Clemmys guttata*]) (09:6)
Type locality: "les grandes Indes" [India]. Restricted to "Pondicherry, South Arcot (district), Tamil Nadu (state), India" [yellow star] by Webb (1980a:552).
Type specimen: MNHN 7978, holotype, type specimen figured (pl. opp.171), discussed by Webb (1980a) and Bour et al. (1995).
Etymology:* Latin, *punctata*, having small points; referring to the small punctate nodules on the surface of the carapacial bony disc.
Comment: Name suppressed by ICZN (2005a) as published in a rejected and invalid non-binomial work, see Savage (2003).

Testudo punctata Bonnaterre 1789:30 (senior homonym, not = *Testudo punctata* Schoepff 1792 (*nomen suppressum*) [= *Clemmys guttata*])
Trionyx (Emyda) punctatus, Trionyx punctatus, Emyda punctata, Trionyx punctata, Lissemys punctata, Lissemys punctata punctata, Trionyx punctatus punctatus
Type locality: "les grandes Indes" [India]. Restricted to "Pondicherry, South Arcot (district), Tamil Nadu (state), India" [yellow star] by Webb (1980a:552).
Type specimen: MNHN 7978, holotype, discussed by Webb (1980a) and Bour et al. (1995).
Etymology:* Latin, *punctata*, having small points; referring to the small punctate nodules on the surface of the carapacial bony disc.

Testudo sonnerati Meyer 1790:83 (*nomen novum et oblitum*) (09:8)
Etymology:* Eponym honoring Pierre Sonnerat (1748–1814), France; natural scientist and explorer who obtained the type.
Comment: Unjustified replacement name for *punctata*.

Testudo granulosa Suckow 1798:48 (*nomen novum*)
Etymology:* Latin, *granulum*, grain; referring to the granular and rough carapacial surface of juveniles.
Comment: Unjustified replacement name for *punctata*.

Testudo scabra Latreille *in* Sonnini and Latreille 1801:164 (junior homonym and *nomen novum*, not = *Testudo scabra* Linnaeus 1758 [= *Rhinoclemmys punctularia punctularia*])
Etymology:* Latin, *scabra*, rough, shabby, mangy, unclean; referring to the rough carapacial surface.
Comment: Unjustified replacement name for *punctata*.

Testudo granosa Schoepff 1801:127
Trionyx granosus, Cryptopus granosus, Emyda granosa, Emyda granosa granosa, Lissemys punctata granosa, Trionyx punctatus granosus
Type locality: "Coromandeliae" [India] [green star, location estimated].
Type specimens: ZMB, syntypes (2), apparently lost (Webb 1980a).
Etymology:* Latin, *granulum*, grain; referring to the granular and rough carapacial surface of juveniles.
Comment: Unjustified emendation of *granulosa*.

Testudo granulata Daudin 1801:81 (*nomen novum*)
Type locality: "les grandes Indes" [India]. Restricted to "Pondicherry, South Arcot (district), Tamil Nadu (state), India" [yellow star] by Webb (1980a:552).
Type specimen: MNHN 7978, holotype, by default, type specimen figured in Lacepède (1788:pl.opp.171) and Daudin (1801:pl.19,f.2), see Webb (1980a) and Bour et al. (1995).
Etymology:* Latin, *granulum*, grain; referring to the granular and rough carapacial surface of juveniles.
Comment: Unjustified emendation of *granulosa*.

Trionyx coromandelicus Geoffroy Saint-Hilaire 1809a:364 (*nomen novum*)
Type locality: "La Côte de Coromandel" [India] [green star, location estimated].
Type specimen: MNHN 7978, holotype, by default, discussed by Webb (1980a) and Bour et al. (1995).
Etymology:* Toponym for the Coromandel Coast, India.
Comment: Unjustified replacement name for *granosa* and *granulata*.

Emyda dura Buchanan-Hamilton *in* Anderson 1876:514 (*nomen nudum*)
Etymology:* Latin, *dura*, hard, strong; presumably referring to the hard bony carpace.


Lissemys punctata andersoni Webb 1980a (11:14, 21:101)
Spotted Northern Indian Flapshell Turtle



Shailendra Singh / CBFTT / Ghaghra R., Uttar Pradesh, India



left: Shailendra Singh / CBFTT / Yamuna R., Uttar Pradesh, India
right: Chittaranjan Baruah / CBFTT / Assam, India / juvenile



** (subspecies: *andersoni* = blue, *vittata* = green; overlap = intergrades; orange dots = probably introduced, translocated, or trade, star = type locality)

Distribution:* Bangladesh, India (*Arunachal Pradesh, Assam, Bihar, Chhattisgarh, Haryana, Jammu, Madhya Pradesh, Meghalaya, Punjab, Rajasthan, Sikkim [?], Tripura, Uttarakhand, Uttar Pradesh, West Bengal*), Myanmar, Nepal, Pakistan

Introduced: India (*Andaman Islands*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 1,343,471 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 3,218,279 sq. km

Size (Max SCL): male 23.0 cm, female 37.0 cm (Yadava and Prasad 1982; Agarwal 1987; Baruah et al. 2012; Ceballos et al. 2013; Bhupathy et al. 2014 CBFTT)

CITES: Appendix II (1995). Previously: Appendix I (1975)


Synonymy:

Lissemys punctata andersoni Webb 1980a:554
Lissemys andersoni
Type locality: "Belbari, Terai, southeastern Nepal, elevation 210 m" [yellow star] (see Mitchell and Rhodin 1996).
Type specimen: MNHN 1977.1486, holotype; Bour et al. (1995) erroneously listed MNHN 1977.1986.
Etymology:* Eponym honoring John Anderson (1833–1900), United Kingdom (Scotland) / India; physician, naturalist, and curator of the Indian Museum in Calcutta.


Lissemys punctata vittata (Peters 1854) (11:14, 21:101)
Central Indian Flapshell Turtle



Shailendra Singh / CBFTT / Mahanadi R., Odisha, India



Peter Praschag / CBFTT / Netravati R., Karnataka, India



** (subspecies: *punctata* = red, *andersoni* = blue, *vittata* = green; overlap = intergrades; orange stars = probably introduced, translocated, or trade, stars = type localities)

Distribution:* Bangladesh, India (*Andhra Pradesh, Chhattisgarh, Goa, Gujarat, Karnataka, Madhya Pradesh, Maharashtra, Odisha, Rajasthan, Telangana, West Bengal*), Pakistan

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 910,219 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 2,024,466 sq. km

Size (Max SCL): No data located

CITES: Appendix II (1995). Previously: Appendix I (1975)

Synonymy:


Emyda vittata Peters 1854:216
Emyda granosa vittata, Lissemys punctata vittata
Type locality: "India orientalis, Goa" [yellow star].
Type specimens: ZMB 46, 17772, syntypes (2), ZMB 46 erroneously listed as "holotype" by Webb (1980a), see Fritz et al. (1994).
Etymology:* Latin, *vitta*, band, ribbon, stripe, *vittata*, striped; referring to the head stripe.

Emyda granosa intermedia Annandale 1912a:171
Type locality: "Near Purulia, Manbhum Dist." [West Bengal, India] [green star].
Type specimens: ZSI 16764–65, syntypes (2), see Das et al. (1998), not located by Kundu et al. (2018a).
Etymology:* Latin, *intermedia*, in-between; referring to the relationship of this species to others.

Lissemys scutata (Peters 1868) (11:14)
Burmese Flapshell Turtle, Myanmar Flapshell Turtle


Gerald Kuchling / CRM 2 / Myanmar / trade: Ruili, Yunnan, China


Gerald Kuchling / Myanmar / trade: Ruili, Yunnan, China



(orange dot = probable trade; star = type localities)

Distribution: Myanmar
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 180,806 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 321,382 sq. km
Size (Max SCL): 25.0 cm (Platt et al. 2018)
IUCN Red List: Least Concern (LC) (Horne, Praschag, and Platt 2021). Previously: Data Deficient (DD) (ATTWG 2000); Data Deficient (DD) (TFTSG *in* Baillie and Groombridge 1996)
CITES: Appendix II (1995)
Synonymy:
Emyda scutata Peters 1868:449
Emyda granosa scutata, Lissemys punctata scutata, Trionyx punctatus scutatus, Lissemys scutata
Type locality: "Pegú" [Bago, Myanmar] [yellow star].
Type specimen: ZMB 6029, holotype, see Fritz et al. (1994).
Etymology:* Latin, *scutum*, shield, scute, *scutata*, having a shield; referring to the pronounced entoplastral callosity.
Emyda fuscomaculata Gray 1873c:308
Type locality: Not designated. Restricted to "Asia, Pegu?" [Myanmar] [yellow star] by Gray (1873j:89).
Type specimen: NHMUK 1946.1.22.4 (formerly 1868.4.3.138), holotype.
Etymology:* Latin, *fusca*, dark brown, and *maculata*, spotted, speckled, stippled; referring to the brownish carapace color in juveniles with dark brown spots.

Subfamily Trionychinae Gray 1825 (89)*
Softshell Turtles
(includes 10 genera, 28 species, 38 taxa)
Amydae Oppel 1811:9 (*partim*)
Trionyces Schmid 1819:18
Trionicidae Gray 1825:212
Trionychoidea Fitzinger 1826:5
Trionychinae Lydekker 1889:4

Amyda Schweigger *in* Geoffroy Saint-Hilaire 1809a (89)*
Asiatic Softshell Turtles
(includes 2 species, 5 taxa)
Amyda Schweigger *in* Geoffroy Saint-Hilaire 1809a:365
Type species: *Amyda javanica* Schweigger *in* Geoffroy Saint-Hilaire 1809a [= subjective synonym of *Testudo cartilaginea* Boddaert 1770], by original monotypy; invalid subsequent type species designation of *Amyda mutica* [= *Apalone mutica*] by Agassiz (1857a:399).
Etymology:* Greek, *emydos*, turtle.
Amida Duméril and Bibron 1834:416 (*nomen novum*)
Etymology:* Greek, *emydos*, turtle.
Comment:* Unjustified emendation of *Amyda*.
Potamochelys Fitzinger 1843:30
Type species: *Aspidonectes (Potamochelys) javanica* Wagler [= *Amyda javanica* Schweigger *in* Geoffroy Saint-Hilaire 1809a] [= subjective synonym of *Testudo cartilaginea* Boddaert 1770], by original monotypy.
Etymology:* Greek, *potamos*, river, and *chelys*, turtle.
Aspilus Gray 1864b:83 (junior homonym, not = *Aspilus* Schaum 1848 [= Coleoptera])
Type species: *Aspilus cariniferus* [= *Trionyx cariniferus* Gray 1856b] [= subjective synonym of *Testudo cartilaginea* Boddaert 1770], by original designation.
Etymology:* Greek, *a-*, without, and *spilos*, a spot or blemish.
Ida Gray 1873h:55 (junior homonym, not = *Ida* Gray 1838 [= Sauria] or *Ida* Gray 1840 [= Gastropoda] or *Ida* Robineau-Desvoidy 1863 [= Diptera])
Type species: *Ida ornata* [= *Trionyx ornatus* Gray 1861a] [= subjective synonym of *Testudo cartilaginea* Boddaert 1770], by original monotypy.
Etymology:* Eponym honoring Ida Pfeiffer (1797–1858), Austria; female explorer and first European woman to cross Borneo who collected specimens of *Amyda cartilaginea* on Ambon and Seram (now part of Indonesia), and donated them to the British Museum.

Amyda cartilaginea (Boddaert 1770) [17:76]
Asiatic Softshell Turtle
(includes 2 subspecies and several taxonomically unspecified populations of *A. cartilaginea* sensu lato)



(subspecies: *cartilaginea* = red, *maculosa* = blue, taxonomically unspecified populations of *A. cartilaginea* sensu lato = gray, orange dots = probable trade)

Distribution: Brunei, Indonesia (*Bali, Java, Kalimantan, Sumatra*), Malaysia (*East, West*), Singapore, Thailand

Introduced or Trade: Indonesia (*Lesser Sundas, Moluccas, Sulawesi*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 1,059,398 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 3,460,269 sq. km

Size (Max SCL): male 75.0 cm (see subspp.)

CBFTT Account: Auliya, van Dijk, Moll, and Meylan (2016) [includes *Amyda ornata*]

IUCN Red List: Vulnerable (VU A1cd+2cd [ver. 2.3]) (ATTWG 2000). Previously: Vulnerable (VU) (TFTSG *in* Baillie and Groombridge 1996)


TFTSG Provisional Red List: Vulnerable (VU) (2011)

CITES: Appendix II (2005)

Amyda cartilaginea cartilaginea (Boddaert 1770) [17:76]
South Sundas Softshell Turtle



Mark Auliya / Java, Indonesia / trade



(subspecies: *cartilaginea* = red, *maculosa* = blue, taxonomically unspecified populations of *A. cartilaginea* sensu lato = gray, orange dots = probable trade; stars = type localities)

Distribution: Indonesia (*Bali, Java, Kalimantan*), Malaysia (*East*)

Introduced or Trade: Indonesia (*Lesser Sundas [Lombok]), Moluccas [Ambon, Seram], Sulawesi*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 418,241 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 1,293,500 sq. km

Size (Max SCL): male 71.0 cm [estimated]; **(Max CCL):** male 80.0 cm (Kusrini et al. 2009; Auliya et al. 2016 CBFTT)

CITES: Appendix II (2005)

Synonymy:

Testudo cartilaginea Boddaert 1770:1
Gymnopus cartilaginea, Trionyx cartilagineus, Aspidonectes cartilagineus, Potamochelys cartilagineus, Amyda cartilaginea, Amyda cartilaginea cartilaginea
Type locality: Not known. Restricted to "Java" [Indonesia] [yellow star, location estimated] by Baur (1893a:220).
Type specimen: MNHN 4150, holotype, discussed by Bour et al. (1995) and Bour (2007c).
Etymology:* Latin, *cartilago*, cartilage; referring to the leather-like fibrous margins of the shell.

Testudo membranacea Blumenbach 1779:257 (*nomen dubium*)
Type locality: "Guiana" [Guyana, in error] [no star].
Type specimen: Originally ZMUG, now possibly ZFMK, apparently lost, type specimen figured in Schneider (1783:pl.1), see Wermuth (1956).
Etymology:* Latin, *membrana*, membrane; referring to the shell being covered with a leather-like skin.

Testudo boddaerti Schneider 1787:12 (*nomen novum*)
Trionyx boddaerti, Testudo boddarti, Testudo bodderti
Etymology:* Eponym honoring Pieter Boddaert (1730–1795), Netherlands; physician and naturalist who first described the species.
Comment: Unjustified replacement name for *cartilaginea*, see Wermuth (1956).

Testudo striata Suckow 1798:37 (*partim, nomen novum*)

Etymology:* Latin, *stria*, stripe, groove, *striata*, striped, grooved; referring to the striped or grooved shell pattern of this species.
Comment: Unjustified replacement name for *cartilaginea*. Name synonymized *in partim* under *Amyda cartilaginea cartilaginea*, *Pelodiscus sinensis*, and *Trionyx triunguis*.

Trionyx javanicus Schweigger *in* Geoffroy Saint-Hilaire 1809a:365 (senior homonym, not = *Trionyx javanicus* Gray 1830e [= *Nilssonia gangetica* or *Nilssonia leithii*])
Amyda javanica, Trionyx iavanicus, Tyrse javanica, Aspidonectes javanicus, Trionyx stellatus javanica, Aspilus javanicus, Potamochelys javanicus
Type locality: "Java et les îles voisines" [Indonesia] [yellow star, location estimated].
Type specimen: MNHN 9380, lectotype, designated by Bour et al. (1995:77), see Bour (2007c) and Ceríaco and Bour (2012).
Etymology:* Toponym for Java, Dutch Indies (now Indonesia).

Trionyx stellatus Geoffroy Saint-Hilaire 1809a:365 (*nomen novum*)
Potamochelys stellatus
Type specimen: MNHN 4150, holotype, discussed by Bour et al. (1995).
Etymology:* Latin, *stella*, star, *stellatus*, having stars; referring to the pattern on the shell.
Comment: Unjustified replacement name for *cartilaginea*.

Trionyx cariniferus Gray 1856b:67
Aspilus cariniferus, Trionyx carinifera
Type locality: "Moluccas" [Indonesia] [orange star, location estimated; probable trade specimen].
Type specimen: NHMUK 1946.1.22.95, holotype.
Etymology:* Latin, *carina*, keel, and *ferre*, to bear; referring to the distinct vertebral keel on the carapaceof juveniles.

Aspilus punctulatus Gray 1864b:84
Type locality: "Amboina or Ceram" [Indonesia] [orange star, location estimated; probable trade specimen].
Type specimens: NHMUK 1855.3.24.1 (3 specimens), syntypes (3).
Etymology:* Latin, *punctulum*, small spot, *punctulatus*, having small spots; referring to the distinct bright spots on the carapace, head, neck and extremities of juveniles.

Trionyx jeudi Gray 1869a:217
Type locality: "Java?" [Indonesia] [yellow star, location estimated].
Type specimen: NHMUK 1947.3.6.11 (formerly 1867.4.2.177), holotype.
Etymology:* Eponym honoring Theodoor Gerard van Lidth de Jeude (1788–1863), Netherlands; physician, veterinarian, and zoologist at Utrecht University who presented the type to the British Museum.

Testudo alba Boddaert *in* Gray 1873j:85 (*nomen nudum*)
Etymology:* Latin, *alba*, white; apparently referring to the color of an albino or preserved bleached specimen.

Trionyx trinilensis † Jaekel 1911:78
Type locality: "Pithecanthropus-schichten...Java...Trinil" [Indonesia] [gray star].
Type specimens: ZMB (MNB s/n or MB s/n), syntypes (2), fossil, plastral bones, figured (pl.15,f.12-13), see Karl (1987).
Etymology:* Toponym for Trinil, Java, Indonesia.
Geological age: Pleistocene, Trinil Beds.


Trionyx nakornsrithammarajensis Nutaphand 1979:209 (*nomen dubium*)
Amyda nakornsrithammarajensis
Type locality: "Southern Thailand (Nakorn Sri Thammaraj Province)" [green star, location estimated].
Type specimen: Not located, type specimen figured (f.139-140).
Etymology:* Toponym for Nakorn Sri Thammaraj Province, Thailand.
Comment: Unclear whether this described taxon is distinct or not, but it appears phenotypically closer to *A. c. cartilaginea* than to *A. c. maculosa* or *A. ornata*, and for now we consider it a taxonomically unspecified population of *A. cartilaginea* sensu lato.

Trionyx cartilageneus nakorn Nutaphand 1990:[8] (*nomen invalidum*)
Trionyx cartilagineus nakorn, Amyda cartilaginea nakorn
Etymology:* Toponym for Nakorn Sri Thammaraj Province, Thailand.
Comment: Unjustified emendation for *nakornsrithammarajensis*.

Amyda cartilaginea maculosa Fritz, Gemel, Kehlmaier, Vamberger, and Praschag 2014a (17:76)
North Sundas Softshell Turtle


Mark Auliya / CBFTT / Anjungan, West Kalimantan, Indonesia



left: Indraneil Das / Balai Ringin, nr. Serian, Sarawak, East Malaysia
right: Mark Auliya / CBFTT / Tanjung Lasa, West Kalimantan, Indonesia / juvenile


(subspecies: *cartilaginea* = red, *maculosa* = blue, taxonomically unspecified populations of *A. cartilaginea* sensu lato = gray; star = type locality)

Distribution: Brunei (?), Indonesia (*Kalimantan*, *Sumatra*), Malaysia (*East*, *West [?]*), Thailand (?)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 331,403 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 818,591 sq. km
Size (Max SCL): male 75.0 cm [estimated]; **(Max CCL):** male 85.0 cm (Mardiastuti 2008; Auliya et al. 2016 CBFTT)
CITES: Appendix II (2005)
Synonymy:

Amyda cartilaginea maculosa Fritz, Gemel, Kehlmaier, Vamberger, and Praschag 2014a:240
Type locality: "Nanga Badau, Kalimantan, Indonesia ("Nanga Bandang, Borneo")" [yellow star].
Type specimen: NMW 30204:3, holotype, see Gemel et al. (2019).
Etymology:* Latin, *macula*, spot, *maculosa*, rich in spots; referring to the carapacial markings.

Amyda ornata (Gray 1861a) (17:76) (90)*
Southeast Asian Softshell Turtle
(includes 3 subspecies)



**

(subspecies: *ornata* = red, *jongli* = blue, *phayrei* = green, orange dots = probable trade)

Distribution:* Bangladesh, Cambodia, India (*Assam*, *Mizoram*, *Tripura*), Laos, Myanmar, Thailand, Vietnam

Introduced: Indonesia (*Lesser Sundas*, *Sulawesi*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 711,630 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 1,886,167 sq. km

Size (Max SCL):* male 83.0 cm, female 54.0 cm (see subspp.)

CBFTT Account: Auliya, van Dijk, Moll, and Meylan (2016) [as part of *Amyda cartilaginea*]

IUCN Red List: Vulnerable (VU A1cd+2cd [ver. 2.3]) (ATTWG 2000), as part of *Amyda cartilaginea*. Previously: Vulnerable (VU) (TFTSG 1996), as part of *A. cartilaginea*


TFTSG Provisional Red List: Vulnerable (VU) (2015)

CITES: Appendix II (2005), originally as part of *A. cartilaginea*

Amyda ornata ornata (Gray 1861) (17:76)
Indochinese Softshell Turtle



David Emmett / CBFTT / Cardamom Mts. region, Cambodia



(subspecies: *ornata* = red, *phayrei* = green; orange dots = probable trade; star = type locality)

Distribution: Cambodia, Laos, Thailand, Vietnam

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 302,408 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 568,819 sq. km

Size (Max SCL): No data located

CITES: Appendix II (2005), originally as part of *A. cartilaginea*

Synonymy:

Trionyx ornatus Gray 1861a:41

Aspilus ornatus, *Ida ornata*, *Amyda ornata*, *Amyda ornata ornata*, *Amyda cartilaginea ornata*

Type locality: "Camboja" [Cambodia] [yellow star, location estimated, possibly nr. Angkor Wat].

Type specimen: NHMUK 1946.1.22.6–7, parts of one specimen (formerly 1860.11.14.1), holotype, collected by Henri Mouhot.

Etymology:* Latin, *ornatus*, decorated or adorned; referring to the ornate carapacial markings.


Amyda ornata jongli Praschag and Gemel 2022 (90)*
Northwest Asian Softshell Turtle



Peter Praschag / Sangu R. nr. Dohazari, Chittagong hill tracts, Bangladesh



Peter Praschag / Sangu R. nr. Dohazari, Chittagong hill tracts, Bangladesh



** (subspecies: *jongli* = blue, *phayrei* = green; orange dots = probable trade; star = type locality)

Distribution:* Bangladesh, India (Assam, Mizoram, Tripura), Myanmar

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 69,843 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 98,074 sq. km

Size (Max SCL):* No adult data located.

CITES: Appendix II (2005), originally as part of *A. cartilaginea*


Synonymy:*

Amyda ornata jongli Praschag and Gemel 2022:82
Type locality: "Sangu River near Dohazari, Chittagong hill tracts, south-eastern Bangladesh" [yellow star].
Type specimen: NMW 40287, skeletal specimen, holotype.
Etymology:* Vernacular (Bengali), *jongli*, derived from *jangala* (Sanskrit), local indigenous name for the species, meaning wild and ferocious; also meaning "coming from rivers of jungle woodlands."


Amyda ornata phayrei (Theobald 1868b) (17:76)
Myanmar Softshell Turtle, Burmese Softshell Turtle



Gerald Kuchling / CBFTT / Dokhtawady R., Myanmar



Gerald Kuchling / Dokhtawady R., Myanmar



** (subspecies: *ornata* = red, *jongli* = blue, *phayrei* = green, orange dots = probable trade; stars = type localities)

Distribution:* Myanmar, Thailand

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 339,379 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 811,970 sq. km

Size (Max SCL):* male 83.0 cm, female 54.0 cm [female estimated] (Kitana 1997; Cox et al. 1998)

CITES: Appendix II (2005), originally as part of *A. cartilaginea*

Synonymy:

Trionyx phayrei Theobald 1868b:18
Aspidonectes phayrei, *Amyda ornata phayrei*, *Amyda cartilaginea ornata*
Type locality: "montium Arakanensium, prope Bassein" [Arakan

Hills, nr. Bassein, Myanmar] [yellow star].
Type specimen: NHMUK 1868.5.18.2 (skull) and 1887.8.6.1 (shell), holotype.
Etymology:* Eponym honoring Sir Arthur Purves Phayre (1812–1885), United Kingdom / Myanmar; army officer and Commissioner in Burma (now Myanmar) who obtained the type.

Trionyx ephippium Theobald 1875:177
Type locality: "Tenasserim" [Myanmar] [Tanintharyi, Myeik District, Myanmar] [green star].
Type specimen: NHMUK 1946.1.22.8 (formerly 1881.7.8.8–9), holotype.
Etymology:* Greek, *ephippos*, meaning mounted, as on a horse, or a saddle; referring to the saddle-like carapacial marking in juveniles.

Trionyx phayrii Boulenger 1889:ix (*nomen novum*)
Amyda phayrii
Etymology:* Eponym honoring Sir Arthur Purves Phayre (1812–1885), United Kingdom / Myanmar; army officer and Commissioner in Burma (now Myanmar) who obtained the type.
Comment: Unjustified emendation or error for *phayrei*.

Apalone Rafinesque 1832 (89)*

North American Softshell Turtles
(includes 3 species, 9 taxa)

Aplaxia Rafinesque 1817:166 (*nomen oblitum*)
Type species: *Aplaxia nasica* Rafinesque 1817 (*nomen nudum*) [= *Trionyx nasica* Rafinesque 1822 (*nomen suppressum*] [= subjective synonym of *Trionyx spiniferus* LeSueur 1827], by original monotypy.
Etymology:* Greek, *a-*, without, and *plax*, plate, table, surface; referring to the lack of horny scutes on the shell.

Apalone Rafinesque 1832:64
Type species: *Apalone hudsonica* Rafinesque 1832 [= subjective synonym of *Trionyx spiniferus* LeSueur 1827], by original monotypy.
Etymology:* Greek, *apalos*, soft, and *chelone*, turtle; referring to the soft shell.

Mesodeca Rafinesque 1832:64
Type species: *Mesodeca bartrami* [= *Testudo bartrami* Daudin 1801] [= subjective synonym of *Testudo ferox* Schneider 1783], by original monotypy.
Etymology:* Greek, *mesos*, middle, between, and *deca*, ten; referring to ten small tubercles in the middle of the anterior carapacial margin.

Platypeltis Fitzinger 1835:109
Type species: *Platypeltis ferox* Fitzinger [= *Testudo ferox* Schneider 1783], by subsequent designation by Fitzinger (1843:30).
Etymology:* Greek, *platys* = flat, even, wide, and *pelte*, shield; referring to the relatively flat carapace.
Comment: Genus established as *Trionyx (Platypeltis)* without a type species.

Callinia Gray 1869a:221
Type species: *Callinia spinifera* [= *Trionyx spiniferus* LeSueur 1827], by subsequent designation by Stejneger (1907:514).
Etymology:* Latin, *callum*, callus or calloused; referring to the hard skin covering the bony carapace.

Euamyda Stejneger 1944:7
Type species: *Euamyda mutica* [= *Trionyx muticus* LeSueur 1827], by original monotypy.
Etymology:* Greek, *ef-*, *eu-*, real, correct, good, and *emydos*, turtle; referring to the conclusion that *Trionyx muticus* is the type species of this new genus rather than of *Amyda*, as proposed by Agassiz (1857a).

Apalone ferox (Schneider 1783)


Florida Softshell Turtle



Matthew Aresco / CRM 3 / Leon Co., Florida



John B. Iverson / Marion Co., Florida / adult, juvenile



** (orange dots = introduced or translocated; stars = type localities)

Distribution: USA (*Alabama, Florida, Georgia, South Carolina*)
Introduced:* USA (*Louisiana, Texas*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 247,090 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 532,015 sq. km
Size (Max SCL): male 32.4 cm, female 67.3 cm; **(Max CCL):** female 73.8 cm (Pritchard 2001; Ernst and Lovich 2009)
IUCN Red List:* Least Concern (LC) (Munscher, Morrison, Siders, Weber, and Walde 2025). Previously: Least Concern (LC) (van Dijk 2011); Least Concern (LC) [Not Listed] (Behler 1996)
CITES:* Appendix II, as *Apalone* spp. (2023). Previously: Appendix III (USA) (2016)
Synonymy:

Testudo ferox Schneider 1783:330
Emydes ferox, Trionyx ferox, Amyda ferox, Aspidonectes ferox, Platypeltis ferox, Amyda ferox ferox, Trionyx ferox ferox, Apalone ferox
Type locality: "Savannach und Alatama..&..Ost-Florida" [Savannah and Altamaha [Georgia] and East Florida] [USA]. Restricted to "Savannah river, Ga." [Georgia, USA] by Baur (1893a:220), to "East Florida" [USA] by Neill (1951:17), and to "Savannah River (at Savannah), Georgia" [yellow star] [USA] by Schwartz (1956b:8).
Type specimen: NHMUK 1947.3.6.17 (formerly 53a), holotype, figured in Pennant (1771:pl.10,f.1-3), see Boulenger (1889) and Stejneger (1944).
Etymology:* Latin, *ferox*, wild, ferocious; referring to the behavior of this taxon when captured or threatened.

Testudo mollis Lacepède 1788:137, synopsis[table] (*nomen suppressum*) (09:6)
Type locality: "sud de la Caroline, rivières de Savannah & d'Alatamaha..&..la Floride orientale" [USA].

Type specimen: Possibly MNHN, not located, type specimen figured (pl.opp.137).
Etymology:* Latin, *mollis*, soft; referring to the shell bones covered by a soft skin.
Comment: Name suppressed by ICZN (2005a) as published in a rejected and invalid non-binomial work, see Savage (2003).

Testudo mollis Bonnaterre 1789:25
Type locality: "rivières de Savannah, d'Alatamaha, de la Floride, & de la Caroline méridionale" [USA].
Type specimen: Possibly MNHN, not located, type specimen figured in Lacepède (1788:pl.opp.137).
Etymology:* Latin, *mollis*, soft; referring to the shell bones covered by a soft skin.

Testudo (ferox) verrucosa Schoepff 1795:90 (senior homonym, not = *Testudo verrucosa* Suckow 1798 [= *Rhinoclemmys punctularia punctularia*])
Testudo verrucosa, Trionyx verrucosus
Type locality: "Floridae Orientalis" [USA]. Restricted to "Halfway Pond...between...Palatka and Gainesville, Fla...somewhere in southwestern Putnam County" [Florida, USA] [green star, location estimated] by Harper (1940:717) by restriction of *nomen novum* replacement name *Testudo bartrami* Daudin 1801.
Type specimen: Not located, type specimen figured in Bartram (1791:pl.4-5).
Etymology:* Latin, *verruca*, wart, *verrucosa*, full of warts, warty; referring to the rough carapacial surface.

Testudo bartrami Daudin 1801:74 (*nomen novum*)
Chelys bartrami, Trionyx bartrami, Mesodeca bartrami
Type locality: "la Floride orientale" [Florida, USA]. Restricted to "Halfway Pond...between...Palatka and Gainesville, Fla...somewhere in southwestern Putnam County" [Florida, USA] [green star, location estimated] by Harper (1940:717).
Type specimen: Not located, type specimen figured in Bartram (1791:pl.4-5).
Etymology:* Eponym honoring William Bartram (1739–1823), USA; botanist, ornithologist, and explorer of the southern USA.
Comment: Unjustified replacement name for *verrucosa*.

Trionyx carinatus Geoffroy Saint-Hilaire 1809a:365
Aspidonectes carinatus
Type locality: Not known. Restricted to "Savannah River, Georgia" [USA] [yellow star] by Schmidt (1953:108).
Type specimen: MNHN, holotype, apparently lost, see Stejneger (1944) and Bour et al. (1995).
Etymology:* Latin, *carina*, keel, *carinata*, keeled; referring to the slight carapacial vertebral keel in juveniles.


Trionyx georgianus Geoffroy Saint-Hilaire 1809a:367 (*nomen novum*)
Type locality: "Les fleuves de la Géorgie et de la Caroline" [USA], but should be the same as for *ferox* by default, which was restricted to "Savannah River (at Savannah), Georgia" [USA] [yellow star] by Schwartz (1956b:8).
Etymology:* Toponym for Georgia, USA.
Comment: Unjustified replacement name for *ferox*.

Trionyx georgicus Geoffroy Saint-Hilaire 1809b:17 (*nomen novum*)
Type locality: "Les fleuves de la Géorgie et de la Caroline" [USA]. Restricted to "Savannah River, Georgia" [USA] [yellow star] by Schmidt (1953:109).
Etymology:* Toponym for Georgia, USA.
Comment: Unjustified emendation of *georgianus*.

Trionyx brongniarti Schweigger 1812:288 (*nomen novum*)
Testudo brongniarti, Platypeltis brongniarti
Type locality: Not known. Should be the same as for *carinatus* by default, which was restricted to "Savannah River, Georgia" [USA] [yellow star] by Schmidt (1953:108), but *brongniarti* instead incorrectly restricted to "St. John's River, Florida" [USA] [orange star, location estimated] by Schmidt (1953:109).
Type specimen: MNHN, holotype, same as *carinatus* by default, apparently lost, see Stejneger (1944) and Bour et al. (1995).
Etymology:* Eponym honoring Alexandre Brongniart (1770–1847), France; geologist, paleontologist, and zoologist
Comment: Unjustified replacement name for *carinatus*.

Trionyx harlani Bell *in* Harlan 1835:159
Type locality: "East Florida" [USA].
Type specimen: Not located, possibly originally in OUM, but also possibly a replacement name for *bartrami*.
Etymology:* Eponym honoring Richard Harlan (1796–1843), USA; physician, vertebrate paleontologist, and anatomist.

Apalone mutica (LeSueur 1827)
Smooth Softshell Turtle
(includes 2 subspecies)


**
(subspecies: *mutica* = red [light red = extirpated], *calvata* = blue; red dots = extirpated)

Distribution:* USA (*Alabama, Arkansas, Colorado, Florida, Illinois, Indiana, Iowa, Kansas, Kentucky, Louisiana, Minnesota, Mississippi, Missouri, Nebraska, New Mexico, North Dakota, Ohio, Oklahoma, Pennsylvania [extirpated], South Dakota, Tennessee, Texas, West Virginia, Wisconsin*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 652,923 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 2,972,374 sq. km
Size (Max SCL): male 26.6 cm, female 35.6 cm (see subspp.)
IUCN Red List:* Least Concern (LC) (Godwin 2025). Previously: Least Concern (LC) (van Dijk 2011); Least Concern (LC) [Not Listed] (Behler 1996)
CITES:* Appendix II, as *Apalone* spp. (2023). Previously: Appendix III (USA) (2016)

Apalone mutica mutica (LeSueur 1827)
Midland Smooth Softshell Turtle




Peter V. Lindeman / Scioto R., Pike Co., Ohio





John L. Carr / Ouachita Parish, Louisiana



(subspecies: *mutica* = red [light red = extirpated], *calvata* = blue; red dots = extirpated; star = type locality)

Distribution:* USA (*Alabama, Arkansas, Colorado, Illinois, Indiana, Iowa, Kansas, Kentucky, Louisiana, Minnesota, Mississippi, Missouri, Nebraska, New Mexico, North Dakota, Ohio, Oklahoma, Pennsylvania [extirpated], South Dakota, Tennessee, Texas, West Virginia, Wisconsin*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 618,656 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 2,814,354 sq. km

Size (Max SCL): male 26.6 cm, female 35.6 cm (Barko and Briggler 2006; Ernst and Lovich 2009)

IUCN Red List:* Least Concern (LC) (Godwin, in press)

CITES:* Appendix II, as *Apalone* spp. (2023). Previously: Appendix III (USA) (2016)

Synonymy:

Trionyx pusilla Rafinesque 1822:3 (*nomen suppressum*)
Type locality: "United States" [USA].
Type specimen: Not known or located.
Etymology:* Latin, *pusilla*, small, tiny, dwarfed; referring to the allegedly smaller size of this species compared to other softshells.
Comment: Name suppressed by ICZN (1984) as published in a rejected and invalid work, see Smith et al. (1980a).


Trionyx muticus LeSueur 1827:263
Aspidonectes muticus, Gymnopus muticus, Amyda mutica, Euamyda mutica, Trionyx muticus, Trionyx muticus muticus, Apalone mutica, Apalone muticus, Apalone mutica mutica
Type locality: "Newharmony, sur le Wabash, à peu de distance de son embouchure dans l'Ohio...l'Etat de l'Indiana" [USA] [yellow star].
Type specimen: MNHN 8813, lectotype, designated by Stejneger (1944:14) and Webb (1962:534), see also Bour et al. (1995) and Bour (2010).
Etymology:* Latin, *muticus*, smooth; referring to the smooth carapace without spicules.

Potamochelys microcephalus Gray 1864b:87
Callinia microcephala, Potamochelys microcephala
Type locality: "Sarawak" [Malaysia, in error]. Emended to "___?" [not known] by Boulenger (1889:262); restricted to "New Harmony, Indiana" [USA] [yellow star] by Schmidt (1953:110).
Type specimen: NHMUK 1947.3.6.22 (formerly 1856.9.19.22), holotype.
Etymology:* Greek, *mikro-*, small, and *kefale*, head; referring to the relatively small head compared to other softshells.

Apalone mutica calvata (Webb 1959) (17:77)
Gulf Coast Smooth Softshell Turtle



Barry Mansell / CRM 3 / Escambia Co., Florida



(subspecies: *mutica* = red, *calvata* = blue; star = type locality)

Distribution: USA (*Alabama, Florida, Louisiana, Mississippi*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 34,748 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 129,864 sq. km

Size (Max SCL): male 18.0 cm, female 28.7 cm (Mount 1975; Moler 2006a; Krysko et al. 2019)

IUCN Red List:* Data Deficient (DD) (Godwin, in press)

CITES:* Appendix II, as *Apalone* spp. (2023). Previously: Appendix III (USA) (2016)

Synonymy:

Trionyx muticus calvatus Webb 1959:519
Apalone mutica calvata, Apalone calvata
Type locality: "Pearl River, Roses Bluff, 14 miles northeast Jackson, Rankin County, Mississippi" [USA] [yellow star].

Type specimen: UIMNH 31071, holotype.
Etymology:* Latin, *calvatus*, bald, empty; referring to the lack of a striped pattern on the upper side of the snout.

Apalone spinifera (LeSueur 1827) (12:37)
Spiny Softshell Turtle
(includes 6 subspecies)




(subspecies: *spinifera* = red [light red = likely extirpated], *aspera* = blue, *atra* = tourmaline (Coahuila), *emoryi* = brown, *guadalupensis* = pink, *pallida* = green; overlap = intergrades; orange dots = introduced; red dots = extirpated)

Distribution: Canada (*Ontario, Québec*), Mexico (*Chihuahua, Coahuila, Nuevo León, Tamaulipas*), USA (*Alabama, Arkansas, Colorado, Florida, Georgia, Illinois, Indiana, Iowa, Kansas, Kentucky, Louisiana, Maryland, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma, Pennsylvania, South Carolina, South Dakota, Tennessee, Texas, Vermont, Virginia, West Virginia, Wisconsin, Wyoming*)
Introduced:* Mexico (*Baja California, Sonora, Chihuahua*); USA (*Arizona, California, Hawaii, Maryland, Massachusetts, Montana, Nevada, New Jersey, New York, North Carolina, Pennsylvania, Utah, Virginia*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 1,770,022 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 6,550,365 sq. km
Size (Max SCL): male 27.5 cm, female 54.0 cm (see subspp.)
IUCN Red List:* Least Concern (LC) (Doody, Gillingwater, Mali, Forstner, van Dijk, and Rhodin 2024). Previously: Least Concern (LC) (van Dijk 2011); Least Concern (LC) [Not Listed] (Behler 1996)
CITES:* Appendix II, as *Apalone* spp. (2023). Previously: Appendix III (USA) (2016)


Apalone spinifera spinifera (LeSueur 1827) (08:22)
Northern Spiny Softshell Turtle



John B. Iverson / Muscatine, Muscatine Co., Iowa / female



left: John B. Iverson / Buffalo Co., Nebraska
right: Ryan M. Bolton / Long Point National Wildlife Area, Ontario, Canada



(subspecies: *spinifera* = red [light red = likely extirpated], *aspera* = blue, *emoryi* = brown, *guadalupensis* = pink, *pallida* = green; overlap = intergrades; orange dots = introduced; red dots = extirpated; stars = type localities)

Distribution:* Canada (*Ontario, Québec*), USA (*Alabama, Arkansas, Colorado, Georgia, Illinois, Indiana, Iowa, Kansas, Kentucky, Louisiana, Maryland, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma,*

Pennsylvania, South Dakota, Tennessee, Vermont, Virginia, West Virginia, Wisconsin, Wyoming)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 1,086,381 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 4,377,847 sq. km

Size (Max SCL): male 27.5 cm, female 54.0 cm (Halk 1986; Barko and Briggler 2006)

IUCN Red List:* Least Concern (LC) (Doody, Gillingwater, Mali, Forstner, van Dijk, and Rhodin 2024)

CITES:* Appendix II, as *Apalone* spp. (2023). Previously: Appendix III (USA) (2016)

Synonymy:

Aplaxia nasica Rafinesque 1817:166 (*nomen nudum*)
Etymology:* Latin, *nasica*, having a pointed nose; referring to the long, tubular nose.

Trionyx ohiensis Rafinesque 1818:354 (*nomen oblitum*) (21:7)
Type locality: "the Ohio" [Ohio River, USA] [no star].
Type specimen: Not located, type specimen figured in manuscript drawing by Rafinesque in 1818, reproduced by Bell and Bauer (2020:f.7).
Etymology:* Toponym for the Ohio River.

Trionyx nasica Rafinesque 1822:3 (*nomen suppressum*)
Type locality: "western streams...United States" [USA].
Type specimen: Not known or located.
Etymology:* Latin, *nasica*, having a pointed nose; referring to the long, tubular nose.
Comment: Name suppressed by ICZN (1984) as published in a rejected and invalid work, see Smith et al. (1980a).

Trionyx spiniferus LeSueur 1827:258
Gymnopus spiniferus, Gymnopodus spiniferus, Aspidonectes spinifer, Gymnopus spinifer, Trionyx spinifer, Callinia spicifera, Callinia spinifera, Platypeltis spinifer, Tyrse spinifera, Amyda spinifera, Platypeltis spinifera, Amyda spinifer, Amyda spinifera spinifera, Trionyx spinifera, Trionyx spinifera spinifera, Amyda ferox spinifera, Trionyx ferox spinifera, Trionyx spinifer spinifer, Trionyx spiniferus spiniferus, Apalone spinifera, Apalone spiniferus, Apalone spinifera spinifera
Type locality: "Newharmony, sur le Wabash, à peu de distance de son embouchure dans l'Ohio...l'Etat de l'Indiana" [USA] [yellow star].
Type specimen: MNHN 8808, lectotype, designated by Webb (1962:491), discussed by Bour et al. (1995) and Bour (2010).
Etymology:* Latin, *spina*, spine, and *ferre*, to bear, to carry; referring to the spiny, cone-like skin projections on the leading edge of the carapace.

Trionyx ocellatus LeSueur 1827:261 (senior homonym, not = *Trionyx ocellatus* Gray 1832a [= *Nilssonia hurum*])
Type locality: "Newharmony, sur le Wabash, à peu de distance de son embouchure dans l'Ohio...l'Etat de l'Indiana" [USA] [yellow star].
Type specimen: MNHN 6957, lectotype, designated by Webb (1962:491), discussed by Bour et al. (1995) and Bour (2010).
Etymology:* Latin, *ocellus*, little eye, eyeball, *ocellatus*, having eye spots, ocellated; referring to the pattern of black speckles and eye spots on the carapace.

Apalone hudsonica Rafinesque 1832:64
Type locality: "River Hudson between the falls of Hadley, Glen and Baker, and further up to the source" [USA]. Restricted to "Hudson River, near Baker's Falls, Saratoga County, New York" [USA] [green star] by Webb (1962:491).
Type specimen: Not known or located, see Webb (1973).
Etymology:* Toponym for the Hudson River, New York, USA.

Trionyx annulifer Wied-Neuwied 1839:140 (*nomen novum*)
Etymology:* Latin, *annulus*, a ring, and *fero*, to bear; referring to the small carapacial rings.
Comment: Unjustified replacement name for *ocellatus*.

Tyrse argus Gray 1844:48
Trionyx argus
Type locality: "West Africa, Sierra Leone?" [in error]. Emended to "___?" [not known] by Boulenger (1889:260); restricted to "New Harmony, Indiana" [USA] [yellow star] by Schmidt (1953:110).
Type specimen: NHMUK 1947.3.6.18 (formerly 1843.9.6.12), holotype.
Etymology:* Mythonym for *Argus*, the Greek mythological giant with numerous eyes all over his body; referring to the pattern of black speckles and eye spots on the carapace.

Trionyx annulatus Gray 1856b:69 (*nomen novum*)
Etymology:* Latin, *annulus*, a ring; referring to the small carapacial rings.
Comment: Unjustified replacement name or error for *annulifer*.


Aspidonectes nuchalis Agassiz 1857a:402,406
Platypeltis nuchalis
Type locality: "Cumberland River..[&]..head waters of the Tennessee River" [USA]. Restricted to "Cumberland River near Nashville, Tennessee" [USA] [blue star] by Schmidt (1953:110).
Type specimen: MCZ 1623, lectotype, designated by Webb (1962:492), see also Barbour and Loveridge (1929) and Stejneger (1944).
Etymology:* Latin, *nucha*, neck, spinal cord, *nuchalis*, belonging to the neck; referring to the triangular dilation of the anterior vertebral keel.

Gymnopus olivaceus Wied-Neuwied 1865:55 (*nomen novum*)
Etymology:* Latin, *olivaceus*, olive-colored; referring to the carapacial coloration.
Comment: Unjustified replacement name for *ocellatus*.

Amyda spinifera hartwegi Conant and Goin 1948:1 (08:22)
Amyda ferox hartwegi, Trionyx ferox hartwegi, Trionyx spinifer hartwegi, Trionyx spiniferus hartwegi, Apalone spinifera hartwegi
Type locality: "Wichita, Sedgwick County, Kansas" [USA] [pink star].
Type specimen: UMMZ 95365, holotype.
Etymology:* Eponym honoring Norman E. Hartweg (1904–1964), USA; herpetologist and curator at the Museum of Zoology at the University of Michigan.

Apalone spinifera aspera (Agassiz 1857a)
Gulf Coast Spiny Softshell Turtle


Barry Mansell / CRM 3 / Santa Rosa Co., Florida


**

(subspecies: *spinifera* = red, *aspera* = blue, *pallida* = green; overlap [reddish olive, dark purple] = intergrades; orange dot = introduced; stars = type localities)

Distribution: USA (*Alabama, Florida, Georgia, Louisiana, Mississippi, North Carolina, South Carolina*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 263,860 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 696,348 sq. km

Size (Max SCL): male 20.3 cm, female 45.4 cm (Stejneger 1944; Moler 2006b)

IUCN Red List:* Least Concern (LC) (Doody, Gillingwater, Mali, Forstner, van Dijk, and Rhodin 2024)

CITES:* Appendix II, as *Apalone* spp. (2023). Previously: Appendix III (USA) (2016)

Synonymy:

Aspidonectes asper Agassiz 1857a:402,405

Platypeltis asper, Amyda spinifera aspera, Amyda ferox aspera, Trionyx ferox aspera, Trionyx spinifer asper, Trionyx spiniferus asper, Trionyx spiniferus asperus, Apalone spinifera asper, Apalone spinifera aspera

Type locality: "Mississippi" [USA]. Restricted to "Lake Concordia, La." [Louisiana, USA] [in error; orange star] by Baur (1893:220) and Schmidt (1953:109); and to "Pearl River at Columbus, Marion County, Mississippi" [USA] [yellow star] by Webb (1960:7).

Type specimen: MCZ 1597, lectotype, designated by Webb (1960:6), see also Reynolds et al. (2007); Uetz et al. (2019) erroneously listed MCZ 37172 as the holotype, but that specimen is a syntype of *Platypeltis agassizii* Baur 1888c.

Etymology:* Latin, *asper*, rough or thorny; referring to the anterior carapacial spines.

Platypeltis agassizii Baur 1888c:1122

Trionyx agassizii, Pelodiscus agassizii, Aspidonectes agassizii, Trionyx spiniferus agassizii, Amyda agassizii, Amyda ferox agassizii, Trionyx ferox agassizii

Type locality:* Not designated. Restricted to "Savannah River, Savannah, Georgia" [USA] [green star] by Neill (1951:16).


Type specimen: MCZ 37172, holotype, see Stejneger (1944), discussed also by Cochran (1961) and Reynolds et al. (2007).

Etymology:* Eponym honoring Louis Agassiz (1807–1873), Switzerland / USA; herpetologist at Harvard University who founded the Museum of Comparative Zoology.


Apalone spinifera atra (Webb and Legler 1960) (07:79, 08:23, 12:37)
Black Spiny Softshell Turtle, Cuatro Cienegas Softshell



Suzanne E. McGaugh / CBFTT / Tío Cándido, Cuatrociénegas Basin, Coahuila, Mexico



Suzanne E. McGaugh / CBFTT / Tío Cándido, Cuatrociénegas Basin, Coahuila, Mexico



** (subspecies: *atra* = tourmaline, *emoryi* = brown, *guadalupensis* = pink; star = type locality)

Distribution: Mexico (*Coahuila*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 535 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 2,116 sq. km

Size (Max SCL): male 24.7 cm, female 29.4 cm (Cerdá-Ardura et al. 2008 CBFTT; Legler and Vogt 2013)

CBFTT Account: Cerdá-Ardura, Soberón-Mobarak, McGaugh, and Vogt (2008)

IUCN Red List:* Critically Endangered (CR A4ce) (Doody, Gillingwater, Mali, Forstner, van Dijk, and Rhodin 2024). Previously: Critically Endangered (CR) (TFTSG 1996, 2016), listed originally as *Apalone ater*

CITES: Appendix I (1975)

USA ESA:* Endangered, as *Trionyx ater* (USFWS 1976)

Synonymy:

Trionyx ater Webb and Legler 1960:21

Trionyx spinifer ater, Trionyx spiniferus ater, Apalone spinifera ater, Apalone ater, Apalone spiniferus ater, Apalone spinifera atra, Apalone (Apalone) atra, Apalone atra

Type locality: "16 kilometers south of Cuatro Ciénegas, Coahuila... Mexico" [yellow star]. Emended to "Poza del Tío Cándido, 16 km southwest of Cuatrociénegas de Carranza, central Coahuila, México" by Cerdá-Ardura et al. (2008:2 CBFTT).

Type specimen: KU 46903, holotype, see Duellman and Berg (1962).

Etymology:* Latin, *atra*, black; referring to the color of the carapace.


Apalone spinifera emoryi (Agassiz 1857a) [(12:37)]
Texas Spiny Softshell Turtle



Vincenzo Ferri / Texas / captivity / juvenile



Viviana Ricardez / Kinney Co., Texas / adult male



** (subspecies: *spinifera* = red, *atra* = tourmaline, *emoryi* = brown, *guadalupensis* = pink, *pallida* = green; overlap [bright purple] = intergrades; orange dots = introduced; star = type locality)

Distribution: Mexico (*Chihuahua*, *Coahuila*, *Nuevo León*, *Tamaulipas*), USA (*Texas*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 159,367 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 645,870 sq. km

Size (Max SCL): male 27.3 cm, female 45.0 cm (Legler and Vogt 2013; Brown et al. 2020; C.J. Franklin, unpubl. data); male recorded to 36.1 cm (Degenhardt et al. 1996), but likely a misidentified female

IUCN Red List:* Data Deficient (DD) (Doody, Gillingwater, Mali, Forstner, van Dijk, and Rhodin 2024)

CITES:* Appendix II, as *Apalone* spp. (2023). Previously: Appendix III (USA) (2016)

Synonymy:

Aspidonectes emoryi Agassiz 1857a:392,407
Trionyx emoryi, *Platypeltis emoryi*, *Amyda emoryi*, *Amyda ferox emoryi*, *Trionyx ferox emoryi*, *Trionyx spinifer emoryi*, *Trionyx spinifera emoryi*, *Trionyx spiniferus emoryi*, *Apalone spinifera emoryi*
Type locality:* "lower Rio Grande of Texas, near Brownsville" [USA] and "Williamson County, Texas, in a stream emptying into the Rio Brazos" [USA]; restricted to "lower Rio Grande of Texas, near Brownsville" [USA] by lectotype designation by Webb (1962:514) [yellow star].
Type specimen: USNM 7855, lectotype, designated by Webb (1962:514), see also Cochran (1961), Reynolds et al. (2007), and Lieb (2023).
Etymology:* Eponym honoring William H. Emory (1811–1887), USA; army officer, civil engineer, and surveyor of the USA-Mexico border, who obtained the types.


Aspidonectes emyda Gray 1870c:95 (*nomen novum*)
Etymology:* Greek, *emydos*, turtle.
Comment: Unjustified replacement name or error for *emoryi*.

Aspidonectes georgii Gray 1870c:109 (*nomen novum*)
Etymology:* Eponym honoring George Stolley (1817–1911), Germany / USA; naturalist and geologist working with Agassiz who collected specimens of *emoryi*.
Comment: Unjustified replacement name for *emoryi* and *emyda*.

Apalone spinifera guadalupensis (Webb 1962)
Guadalupe Spiny Softshell Turtle



Peter V. Lindeman / Palmetto State Park, San Marcos R., Gonzales Co., Texas



** (subspecies: *atra* = tourmaline, *emoryi* = brown, *guadalupensis* = pink, *pallida* = green; overlap [bright purple] = intergrades; star = type locality)

Distribution: USA (*Texas*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 84,908 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 177,656 sq. km

Size (Max SCL): male 17.3 cm, female 35.2 cm (C.J. Franklin, unpubl. data)


IUCN Red List:* Data Deficient (DD) (Doody, Gillingwater,

Mali, Forstner, van Dijk, and Rhodin 2024)
CITES:* Appendix II, as *Apalone* spp. (2023). Previously: Appendix III (USA) (2016)
Synonymy:
Trionyx spinifer guadalupensis Webb 1962:517
Trionyx spiniferus guadalupensis, Apalone spinifera guadalupensis
Type locality: "15 miles northeast Tilden, McMullen County, Texas" [USA] [yellow star].
Type specimen: UMMZ 89926, holotype, see Kluge (1984) and Lieb (2023).
Etymology:* Toponym for the Guadalupe River, Texas, USA.

Apalone spinifera pallida (Webb 1962)
Pallid Spiny Softshell Turtle


Stanley E. Trauth / nr. Horatio, Sevier Co., Arkansas


Viviana Ricardez / Harris Co., Texas / female


**
(subspecies: *spinifera* = red, *aspera* = blue, *emoryi* = brown, *guadalupensis* = pink, *pallida* = green; overlap [bright purple, reddish olive, dark purple] = intergrades; star = type locality)
Distribution: USA (*Arkansas, Louisiana, Oklahoma, Texas*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 273,467 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 646,121 sq. km
Size (Max SCL): male 23.7 cm, female 47.6 cm (E.C. Munscher and A. Brinker et al., unpubl. data)
IUCN Red List:* Least Concern (LC) (Doody, Gillingwater, Mali, Forstner, van Dijk, and Rhodin 2024)
CITES:* Appendix II, as *Apalone* spp. (2023). Previously: Appendix III (USA) (2016)
Synonymy:
Trionyx spinifer pallidus Webb 1962:522
Trionyx pallidus, Trionyx spiniferus pallidus, Apalone spinifera pallida, Apalone spinifera pallidus
Type locality: "Lake Caddo, Caddo Parish, Louisiana" [USA] [yellow star].
Type specimen: LSUM 111652 (formerly TU 484), holotype.
Etymology:* Latin, *pallidus*, pale; referring to the light color of the carapace.

Chitra Gray 1844 (89)*
Narrow-headed Softshell Turtles
(includes 3 species, 4 taxa)
Chitra Gray 1844:49
Type species: *Chitra indica* [= *Trionyx indicus* Gray 1830e], by original monotypy.
Etymology:* Vernacular (Sanskrit / Bengali), *chitra*, beautiful, bright, picturesque; referring to the attractive markings on the carapace and extremities.

Chitra chitra Nutaphand 1986
Asian Narrow-headed Softshell Turtle
(includes 2 subspecies)


**
(subspecies: *chitra* = red, *javanensis* = blue)
Distribution: Indonesia (*Java, Sumatra*), Malaysia (*West*), Thailand
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 163,145 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 1,617,988 sq. km
Size (Max SCL): 140.0 cm (see subspp.)
IUCN Red List: Critically Endangered (CR A2cd) (Cota, Guntoro, As-singkily, Hamidy, and Kusrini 2019). Previously: Critically Endangered (CR) (ATTWG 2000); Critically Endangered (CR) (TFTSG *in* Baillie and Groombridge 1996)
CITES: Appendix I (2013). Previously: Appendix II, as *Chitra* spp. (2003)

Chitra chitra chitra Nutaphand 1986
Siamese Narrow-headed Softshell Turtle




Chris Tabaka / TCC / Thailand / captivity





left: Peter Paul van Dijk / Thailand / captivity / hatchling
right: Douglas B. Hendrie / No data / captivity / juvenile



(subspecies: *chitra* = red; star = type locality)

Distribution: Malaysia (*West*), Thailand

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 90,238 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 476,677 sq. km

Size (Max SCL): 140.0 cm (McCord and Pritchard 2003a; Ernst et al. 2006b; P.P. van Dijk, unpubl. data)

CITES: Appendix I (2013). Previously: Appendix II, as *Chitra* spp. (2003)

Synonymy:

Chitra chitra chitra Nutaphand 1986:65 (*nomen protectum*)
Chitra chitra
Type locality: "Khwae Noi and Khwae Yai rivers of Kanchanaburi Province and in the Mae Klong river of Ratburi Province" [Thailand]. Restricted to "[Mae Klong in Ratburi and Khwae River in Kanchanaburi]" by Nutaphand (1990:113); and to "Kanburi (= Kanchanaburi), where the Khwae Noi and the Khwae Yai Rivers join to form the Mae Klong River in Kanchanaburi Province, Thailand" [yellow star] by McCord and Pritchard (2003a:18).
Type specimen: Not located, specimen figured in Nutaphand (1990:104), lectotype, erroneously listed as "holotype" by McCord and Pritchard (2003a:18); specimen figured in Nutaphand (1986:65), paralectotype, erroneously designated "lectotype" by Webb and van Dijk (2004:96).
Etymology:* Vernacular (Sanskrit / Bengali), *chitra*, beautiful, bright, picturesque; referring to the attractive markings on the carapace and extremities.
Comment: Name conserved by ICZN (2005b), see McCord and Pritchard (2003b).

Chitra chitra javanensis McCord and Pritchard 2003a
Javanese Narrow-headed Softshell Turtle, *Labi-labi Bintang*



Dian Sartika / nr. Bojonegoro, Bengawan Solo, East Java, Indonesia



(subspecies: *chitra* = red, *javanensis* = blue; stars = type localities)

Distribution: Indonesia (*Java*, *Sumatra*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 72,906 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 369,631 sq. km

Size (Max SCL): 129.0 cm (Iskandar 2004)

CITES: Appendix I (2013). Previously: Appendix II, as *Chitra* spp. (2003)

Synonymy:

Chitra selenkae † Jaekel 1911:80 (*nomen suppressum*)
Type locality: "Pithecanthropus-schichten...Java...Trinil" [Indonesia] [gray star].
Type specimens: ZMB (MNB s/n or MB s/n), syntypes (4), fossils, nearly complete carapace plus plastral bones, figured (pl.15,f.1–2,11,14), see Karl (1987) and McCord and Pritchard (2003a).
Geological age: Pleistocene, Trinil Beds.
Etymology:* Eponym honoring Margarethe L. Selenka (1860–1922), Germany / Indonesia; female zoologist, paleontologist, and women's rights and peace activist, who worked at the Trinil site on Java, Netherlands Indies (now Indonesia).
Comment: Name suppressed by ICZN (2005b), see McCord and Pritchard (2003b).


Chitra chitra javanensis McCord and Pritchard 2003a:41
Chitra chitra javanica
Type locality: "Pasuruan River drainage, near Pasuruan, Probolinggo

District, East Java, Indonesia" (in error according to Iskandar 2004:129) [orange star]. Emended to "Bengawan (= river) Solo, between Kalitidu and Padangan, Bojonegoro District, East Java" [Indonesia] [yellow star] by Iskandar (2004:129).
Type specimen: MZB 199, holotype.
Etymology:* Toponym for Java, Indonesia.

Chitra indica (Gray 1830e) [10:7]
Indian Narrow-headed Softshell Turtle


Shekar Dattatri / CBFTT / Chambal R., Rajasthan, India


left: Chittaranjan Baruah / CBFTT / Assam, India
right: Ashutosh Tripathi / Chambal R., Uttar Pradesh, India



**
(stars = type localities)

Distribution:* Bangladesh, India (*Andhra Pradesh, Assam, Bihar, Gujarat, Haryana, Jammu, Jharkhand, Karnataka, Kerala, Madhya Pradesh, Maharashtra, Odisha, Punjab, Rajasthan, Tamil Nadu, Tripura, Uttar Pradesh, West Bengal*), Nepal, Pakistan
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 677,698 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 4,503,800 sq. km
Size (Max SCL): male 61.5 cm, female 99.0 cm [estimated] (Rashid and Swingland 1997); **(Max CCL):** female 110.0 cm (Das and Singh 2009 CBFTT)
CBFTT Account: Das and Singh (2009)
IUCN Red List: Endangered (EN A1cd+2cd [ver. 2.3]) (ATTWG 2000). Previously: Vulnerable (VU) (TFTSG *in* Baillie and Groombridge 1996)
TFTSG Provisional Red List: Endangered (EN) (2011, 2018)
CITES: Appendix II, as *Chitra* spp. (2003)
Synonymy:
Trionyx indicus Gray 1830e:18 [10:7]
Trionyx egyptiacus indicus, Trionyx aegyptiacus indica, Chitra indica, Gymnopus indicus, Trionyx aegyptianus indicus, Aspidonectes indicus
Type locality: "India." Restricted to "Fatehgahr, Ganges" [Uttar Pradesh, India] [yellow star] by Smith (1931:162).
Type specimen: RCSM 685, holotype, apparently lost, see Farkas (1994); specimen figured in Gray (1831f:pl.80), lectotype, listed as "type" by Webb (1980b), and erroneously as "neotype" by McCord and Pritchard (2003a).
Etymology:* Toponym for India.
Testudo chitra Buchanan-Hamilton *in* Gray 1831d:47 (*nomen nudum*)
Etymology:* Vernacular (Sanskrit), *chitra*, beautiful, bright; referring to the attractive markings on the carapace and extremities.
Gymnopus lineatus Duméril and Bibron 1835:491
Trionyx lineatus
Type locality: "le Gange" [India] [green star, location estimated].
Type specimen: MNHN 6968, holotype, discussed by Bour et al. (1995).
Etymology:* Latin, *linea*, line, guideline, border; referring to the striped pattern on the neck.

Chitra vandijki McCord and Pritchard 2003a
Burmese Narrow-headed Softshell Turtle, Myanmar Narrow-headed Softshell Turtle


Win Ko Ko / CBFTT / TCC / Chindwin R., Myanmar


Gerald Kuchling / CBFTT / Chindwin R., Myanmar


**
(orange dots = probable trade or temple ponds; star = type locality)

Distribution:* Myanmar, Thailand

Introduced:* China (*Yunnan*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 82,189 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 341,458 sq. km
Size (Max SCL):* ca. 100.0 cm (?) (Platt et al. 2014 CBFTT); max recorded: 40.8 cm (Platt et al. 2005); max CCL recorded: 86.0 cm (Hong et al. 2022)
CBFTT Account: Platt, Platt, Win Ko Ko, and Rainwater (2014)
IUCN Red List: Critically Endangered (CR A2cd+4cd) (Platt, Praschag, and Horne 2021)
CITES: Appendix I (2013). Previously: Appendix II, as *Chitra* spp. (2003)
Synonymy:
Chitra burmanica Jaruthanin 2002:40 (*nomen nudum*)
Etymology:* Toponym for Burma (= Myanmar).
Chitra vandijki McCord and Pritchard 2003a:39
Type locality: "Khayansat Kone Village (23°16.30'N; 95°58.99'E) on the Ayeyarwaddy River...Myanmar" [yellow star].
Type specimen: PCHP(CRI) 7059, holotype, figured (f.11); apparently lost (S. Thomson, pers. comm.); PCHP(CRI) 4896–97, 5050, paratypes (3).
Etymology:* Eponym honoring Peter Paul van Dijk (1967–), Netherlands / Malaysia / USA; herpetologist, conservationist, and a Co-Chair of the IUCN Tortoise and Freshwater Turtle Specialist Group who worked in Southeast Asia and the USA, who confirmed the occurrence of *Chitra* in Myanmar.


Dogania Gray 1844 (89)*

Malayan Softshell Turtles
(includes 1 species, 1 taxon)
Dogania Gray 1844:49
Type species: *Dogania subplana* [= *Trionyx subplanus* Geoffroy Saint-Hilaire 1809b], by original monotypy.
Etymology:* Anonym, *dogania*, possibly a fabricated name, possibly an unknown indigenous Vernacular name, or possibly an Eponym honoring someone named Dogan or Dugan.
Sarbieria Gray 1869a:211
Type species: *Sarbieria frenata* [= *Trionyx frenatus* Gray 1856b] [= subjective synonym of *Trionyx subplanus* Geoffroy Saint-Hilaire 1809b], by original monotypy.
Etymology:* Anonym, *sarbieria*, possibly a fabricated name, possibly an unknown indigenous Vernacular name, or possibly an Eponym honoring someone named Sarbier.

Dogania subplana (Geoffroy Saint-Hilaire 1809b)
Malayan Softshell Turtle


Edward O. Moll / Malaysia


left: Peter Paul van Dijk / CRM 2 / Khao Luang, Thailand / hatchling
right: Edward O. Moll / Malaysia


(orange dots = possible trade or questionable; stars = type localities)

Distribution: Brunei, Indonesia (*Java*, *Kalimantan*, *Sumatra*), Malaysia (*East*, *West*), Myanmar, Philippines (*Luzon*, *Mindanao*, *Mindoro*, *Palawan*), Singapore, Thailand
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 1,078,975 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 6,040,981 sq. km
Size (Max SCL): 31.0 cm (Grossmann and Grychta 1998; Pritchard 2001)
IUCN Red List: Least Concern (LC) (Cota Hamidy, Platt, Kusrini, Guntoro, Shepherd, and Schoppe 2021). Previously: Least Concern (LC) (ATTWG 2000); Least Concern (LC) [Not Listed] (Behler 1996)
CITES: Appendix II (2013)
Synonymy:
Trionyx subplanus Geoffroy Saint-Hilaire 1809b:11
Gymnopus subplanus, Amyda subplana, Dogania subplana, Dogania subplanus
Type locality: Not known. Possibly "Malaysia or Java, Indonesia" [yellow stars, locations estimated, possibly Penang or Jakarta] discussed by Bour et al. (1995:79), but not formally restricted.
Type specimen: MNHN A.5182, holotype, discussed by Bour et al. (1995).
Etymology:* Latin, *sub*, almost, under, below, and *plana*, flat, *subplanus*, almost flat; referring to the relatively flattened carapace.
Trionyx frenatus Gray 1856b:67
Potamochelys frenatus, Sarbieria frenata

Type locality: “Singapore” [green star].
Type specimens: NHMUK 1947.3.6.4–5 (formerly 1852.9.13.49, 1853.5.3.8), syntypes (2).
Etymology:* Latin, *frenum*, reins, bridle, *frenatus*, having reins; referring to the black line markings along the neck and head.

Gymnopus javanicus Bibron *in* Gray 1856b:67 (*nomen nudum*)
Etymology:* Toponym for Java, Indonesia.

Dogania guentheri Gray 1862c:265
Trionyx (Dogania) güntheri, Trionyx guentheri, Trionyx güntheri, Dogania güntheri
Type locality: “India” [in error, possible trade specimen, no star].
Type specimen: NHMUK 1947.3.6.6 (formerly 1860.3.19.1025), holotype.
Etymology:* Eponym honoring Albert C.L.G. Günther (1830–1914), Germany / United Kingdom; zoologist, herpetologist, ichthyologist, and curator at the British Museum of Natural History, London.

Trionyx dillwynii Gray 1873c:306
Type locality: “Borneo” [Malaysia or Indonesia] [blue star, location estimated, possibly Labuan].
Type specimen: NHMUK 1946.1.22.12 (formerly 1872.2.19.60), holotype, see Gray (1873j).
Etymology:* Eponym honoring Lewis L. Dillwyn (1814–1892), United Kingdom (Wales); industrialist, politician, and naturalist who studied the natural history of Labuan and Borneo.

Trionyx vertebralis Strauch 1890:113
Type locality: “aus dem Flusse Gabon im aequatorialen West-Afrika” [in error, no star], or “nicht aus Afrika, sondern aus irgend einem Theile Süd-Asiens stammt” [not from Africa, but from some part of South Asia]. Restricted to “Süd-Asien” [no star] by Wermuth and Mertens (1961:255).
Type specimen: ZIN 5391, holotype.
Etymology:* Latin, *vertebra*, vertebra, *vertebralis*, belonging to the vertebrae; referring to the dark longitudinal stripe down the center of the carapace in juveniles.

Trionyx pecki Bartlett 1895a:30
Type locality: “Borneo” [Sarawak, Malaysia] [pink star, location estimated, possibly Kuching].
Type specimen: Possibly SMK, holotype, originally live.
Etymology:* Eponym honoring Captain Herbert W. Peck (1860–1916), United Kingdom / Malaysia; administrative official in Sarawak (now part of Malaysia).

Nilssonia Gray 1872a (07:80, 11:15, 17:78) (89)*

Indian Softshell Turtles
(includes 5 species, 5 taxa)

Nilssonia Gray 1872a:332
Type species: *Nilssonia formosa* [= *Trionyx formosus* Gray 1869a], by original monotopy.
Etymology:* Eponym honoring Sven Nilsson (1787–1883), Sweden; zoologist, naturalist, and director of the Naturhistoriska Riksmuseet, Stockholm.

Isola Gray 1873h:51
Type species: *Isola peguensis* [= *Trionyx peguensis* Gray 1870c] [= subjective synonym of *Trionyx formosus* Gray 1869a], by original monotypy.
Etymology:* Vernacular (Old English), *isola*, isolated; possibly referring to the fact that only one species was described in this genus.


Aspideretes Hay 1903:274
Type species: *Aspideretes gangeticus* [= *Trionyx gangeticus* Cuvier 1825], by original designation.
Etymology:* Greek, *aspidos*, a shield, and *eretes*, oars; referring to the appearance of softshells as a shield with oars.
Comment:* Frétey (2024) noted that this name was published in December 1903, not 1904 as previously recorded by most authors.

Nilssonia formosa (Gray 1869a)

Burmese Peacock Softshell Turtle, Myanmar Peacock Softshell Turtle


Peter Paul van Dijk / TCC / Myanmar / captivity


Gerald Kuchling / Chindwin R., Myanmar


(orange dot = probable trade or introduced; stars = type localities)

Distribution: Myanmar

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 142,290 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 367,882 sq. km

Size (Max SCL): 57.0 cm (van Dijk *in* Pritchard 2001); **(Max CCL):** 65.0 cm (Ernst et al. 2006b; Itescu et al. 2014)

IUCN Red List: Critically Endangered (CR A2cd+4cd) (Horne, Platt, and Praschag 2021). Previously: Endangered (EN) (ATTWG 2000); Vulnerable (VU) (TFTSG *in* Baillie and Groombridge 1996)

CITES: Appendix II (2013)

Synonymy:

Trionyx formosus Gray 1869a:217
Nilssonia formosa, Aspidonectes formosus, Isola formosa, Trionyx formosa, Amyda formosus, Amyda formosa, Trionyx formosanus
Type locality: “Pegu” [Bago, Myanmar] [yellow star].
Type specimen: NHMUK 1946.1.22.11 (formerly 1868.4.3.143), holotype.
Etymology:* Latin, *formosus*, beautiful, splendid, pretty; referring to the fact that the attractive eye spots on the juvenile carapace.

Trionyx peguensis Gray 1870c:99
Isola peguensis
Type locality: “Pegu” [Bago, Myanmar] [yellow star].


Type specimen: NHMUK 1946.1.22.10 (formerly 1868.4.3.142), holotype.
Etymology:* Toponym for Pegu, capital of the Mon kingdom in ancient southern Myanmar.

Trionyx grayii Theobald 1875:176
Type locality: "neighbourhood of Thayet-myo...Irawadi valley" [Myanmar] [green star].
Type specimen: NHMUK 1947.3.6.9 (formerly 1881.7.8.3), holotype.
Etymology:* Eponym honoring John Edward Gray (1800–1875), United Kingdom; herpetologist, turtle taxonomist, and curator at the British Museum of Natural History, London.

Nilssonia gangetica (Cuvier 1825) (12:38)
Indian Softshell Turtle, Ganges Softshell Turtle


B.C. Choudhury / CRM 2 / India


Amtyaz Safi / Khyber Pakhtunkhwa Prov., Pakistan


(orange dots = probable trade or introduced; stars = type localities)

Distribution:* Afghanistan (?), Bangladesh, India (*Assam, Bihar, Chhattisgarh, Gujarat, Haryana, Jammu, Jharkhand, Madhya Pradesh, Meghalaya, Mizoram, Odisha, Punjab, Rajasthan, Tripura, Uttar Pradesh, Uttarakhand, West Bengal*), Nepal, Pakistan
Introduced:* India (*Andhra Pradesh, Odisha*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 729,897 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 3,349,942 sq. km
Size (Max SCL): male 77.0 cm, female 94.0 cm (Rashid and Swingland 1997; Pritchard 2001; Ernst et al. 2006b; Baruah et al. 2012)
IUCN Red List: Endangered (EN A2d+4d) (Ahmed, Choudhury, Das, and Singh 2021). Previously: Vulnerable (VU) (ATTWG 2000); Least Concern (LC) [Not Listed] (Behler 1996)
CITES: Appendix I (1975), originally as *Aspideretes gangeticus*
USA ESA:* Endangered, as *Trionyx gangeticus* (USFWS 1976)
Synonymy:

Trionyx gangeticus Cuvier 1825:186,203
Aspidonectes gangeticus, Tyrse gangetica, Isola gangetica, Aspideretes gangeticus, Trionyx gangeticus, Amyda gangetica, Nilssonia gangetica, Nilssonia gangetica gangetica
Type locality: "du Gange" [Ganges River, India] [yellow star, location estimated].
Type specimen: MNHN 9387, lectotype, designated by Bour et al. (1995:79).
Etymology:* Toponym for the Ganges River, India.

Trionyx javanicus Gray 1830e:19 (*partim*, junior homonym, not = *Trionyx javanicus* Schweigger *in* Geoffroy Saint-Hilaire 1809a [= *Amyda cartilaginea cartilaginea*]) (10:7)
Tyrse javanica
Type locality: "Java and India."
Type specimen: Not known or located.
Etymology:* Toponym for Java, Indonesia.
Comment:* Name synonymized *in partim* under *Nilssonia gangetica* and *Nilssonia leithii*.

Gymnopus duvaucelii Duméril and Bibron 1835:487 (*partim, nomen novum*)
Etymology:* Eponym honoring Alfred Duvaucel (1792–1824), France; naturalist, explorer, and collector for the Muséum National d'Histoire Naturelle, Paris, who collected the type of *Nilssonia gangetica*.
Comment:* Unjustified replacement name for *gangeticus* and *hurum*. Name synonymized *in partim* under *Nilssonia gangetica* and *Nilssonia hurum*.

Trionyx gangeticus mahanaddicus Annandale 1912b:262
Nilssonia gangetica mahanaddica
Type locality: "Sambalpur and Cuttack, Orissa" [India]. Restricted to "Cuttack, Orissa" [Odisha, India] [green star] by Smith (1931:167).
Type specimen: ZSI 17014, holotype, see Das et al. (1998) and Kundu et al. (2018a), listed as *Trionyx gangeticus mahanadicus*.
Etymology:* Toponym for the Mahanadi River, India.

Trionyx gangeticus mahanadicus Das, Dattagupta, and Gayen 1998:128 (*nomen invalidum*)
Etymology:* Toponym for the Mahanadi River, India.
Comment:* Unjustified emendation or error for *mahanaddicus*.


Nilssonia hurum (Gray 1830e) [10:7]
Indian Peacock Softshell Turtle



Chittaranjan Baruah / CBFTT / Brahmaputra R., India



Indraneil Das / CBFTT / Assam, India



** (orange dots = possible trade or introduced or questionable; stars = type localities)

Distribution:* Bangladesh, India (*Arunachal Pradesh, Assam, Bihar, Jammu, Jharkhand, Madhya Pradesh, Odisha, Manipur, Mizoram, Punjab, Rajasthan, Tripura, Uttar Pradesh, West Bengal*), Myanmar (?), Nepal, Pakistan

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 687,940 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 3,004,702 sq. km

Size (Max SCL): male 45.5 cm, female 60.0 cm (Rashid and Swingland 1997; Pritchard 2001; Das et al. 2010 CBFTT; Baruah et al. 2012)

CBFTT Account: Das, Basu, and Singh (2010)

IUCN Red List: Endangered (EN A2d+4d) (Das, Choudhury, Ahmed, Praschag, and Singh 2021). Previously: Vulnerable (VU) (ATTWG 2000); Least Concern (LC) [Not Listed] (Behler 1996)

CITES: Appendix I (1975), originally as *Aspideretes hurum*

USA ESA:* Endangered, as *Trionyx hurum* (USFWS 1976)

Synonymy:

Trionyx occellatus Gray 1830d:pl.78 (*nomen oblitum*)
Type locality: "India." Restricted to "Barrackpore, India" [West Bengal] [yellow star] by Webb (1980b:71).
Type specimen: NHMUK, holotype or syntypes, apparently lost, specimen figured (pl.78), lectotype, designated as "type" by Webb (1980b).
Etymology:* Latin, *ocellus*, small eye or eyeball; referring to the ocellated ringed eye spot pattern on the carapace of juveniles.

Trionyx hurum Gray 1830e:18 [10:7]
Isola hurum, Aspideretes hurum, Aspidonectes hurum, Tyrse hurum, Amyda hurum, Nilssonia hurum
Type locality: "India." Restricted to "Fatehgarh, Ganges" [Uttar Pradesh, India] [orange star] by Smith (1931:171), and to "Barrackpore (about 23 kilometers north Calcutta), West Bengal, India" [yellow star] by Webb (1980b:71) by lectotype designation.
Type specimen: NHMUK, holotype or syntypes, apparently lost, specimen figured in Gray (1835:pl.66), lectotype, designated as "type" by Webb (1980b).
Etymology:* Vernacular (Arabic), *haram*, forbidden; referring to the Muslim belief that turtle meat is impure and should not be consumed (see Theobald 1874).

Testudo chim Buchanan-Hamilton *in* Gray 1831d:47 (*nomen nudum*)
Etymology:* Vernacular (Bengali), *kachim*, turtle (see Das 1995).

Trionyx ocellatus Gray 1832a:directions (junior homonym and *nomen novum*, not = *Trionyx ocellatus* LeSueur 1827 [= *Apalone spinifera spinifera*])
Testudo ocellata, Gymnopus ocellatus
Etymology:* Latin, *ocellus*, small eye or eyeball; referring to the ocellated ringed eye spot pattern on the carapace of juveniles.
Comment: Emended name for *occellatus*.

Gymnopus duvaucelii Duméril and Bibron 1835:487 (*partim, nomen novum*)
Etymology:* Eponym honoring Alfred Duvaucel (1792–1824), France; naturalist, explorer, and collector for the Muséum National d'Histoire Naturelle, Paris.
Comment:* Unjustified replacement name for *gangeticus* and *hurum*. Name synonymized *in partim* under *Nilssonia gangetica* and *Nilssonia hurum*.

Trionyx sewaare Gray 1872a:336
Type locality: "Bengal" [India or Bangladesh]. Restricted to "Ganges" [India] [green star, location estimated] by Smith (1931:171).
Type specimen: Specimen figured in Hardwicke (1835:pl.36), lectotype, designated as "type" by Webb (1980b); NHMUK 1973.1051–2, possible paralectotypes (2).
Etymology:* Vernacular (Sanskrit), *sewa*, to serve, honor, or worship; native name for the species recorded by Hardwicke (see Gray 1872a).

Trionyx bellii Gray 1872a:337
Type locality: "Asia."
Type specimen: Possibly OUM, holotype, apparently lost.
Etymology:* Eponym honoring Thomas Bell (1792–1880), United Kingdom; herpetologist, dental surgeon, and Professor of Zoology at King's College, London.

Trionyx buchanani Theobald 1874:78
Type locality: "Bengal" [India or Bangladesh] [blue star, location estimated].
Type specimen: ZSI 1090, holotype, see Das et al. (1998) and Kundu et al. (2018a).
Etymology:* Eponym honoring Francis Buchanan-Hamilton (1762–1829), United Kingdom (Scotland) / India / Myanmar; physician and botanist who served with the East India Company in India and Burma (now Myanmar).

Trionyx hurum sivalensis † Lydekker 1889:9
Trionyx sivalensis
Type locality: "India...Siwalik Hills" [Punjab] [gray star, location estimated].
Type specimens: ZSI E.163, syntypes (2), fossil, carapace and plastral fragments, figured in Lydekker (1885:pl.27,f.3,3a); NHM(P) R.601, cast of plastral fragment.
Geological age: Late Pliocene to Early Pleistocene.
Etymology:* Toponym for the Siwalik Hills, India.


Nilssonia leithii (Gray 1872a)
Leith's Softshell Turtle, Deccan Softshell Turtle



Rahul Naik / CBFTT / Godavari R. nr. Nashik, Maharashtra, India



left: Shashwat Sirsi / CBFTT / Cauvery R., Karnataka, India / female
right: Veerappan Deepak / CBFTT / Moyar R., nr. Theni, Tamil Nadu, India / male



** (orange dots = questionable; stars = type localities)

Distribution:* India (*Andhra Pradesh, Chhattisgarh, Karnataka, Kerala, Madhya Pradesh, Maharashtra, Odisha, Tamil Nadu, Telangana*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 186,524 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 927,339 sq. km

Size (Max SCL): male 63.5 cm, female 54.8 cm (Moll and Vijaya 1986; Deepak and Vasudevan 2009); **(Max CCL):** male 72.0 cm, possibly to ca. 100 cm (Das et al. 2014 CBFTT)

CBFTT Account: Das, Sirsi, Vasudevan, and Murthy (2014)

IUCN Red List: Critically Endangered (CR A2cd+4cd) (Praschag, Das, Choudhury, and Singh 2021). Previously: Vulnerable (VU) (ATTWG 2000); Near Threatened (NT) (TFTSG *in* Baillie and Groombridge 1996)

CITES:* Appendix I (2023). Previously: Appendix II (2013)

Synonymy:

Trionyx javanicus Gray 1830e:19 (*partim*, junior homonym, not = *Trionyx javanicus* Schweigger *in* Geoffroy Saint-Hilaire 1809a [= *Amyda cartilaginea cartilaginea*]) [10:7]
Tyrse javanica
Type locality: "Java and India" [no star].
Type specimen: Not known or located.
Etymology:* Toponym for Java, Indonesia.
Comment:* Name synonymized *in partim* under *Nilssonia gangetica* and *Nilssonia leithii*.

Testudo gotaghol Buchanan-Hamilton *in* Gray 1831d:48 (*nomen nudum*)
Etymology:* Vernacular (Hindi?), *gotaghol*, probably a local indigenous name for the species.

Trionyx leithii Gray 1872a:334
Isola leithii, Aspideretes leithii, Amyda leithii, Nilssonia leithii
Type locality: "Poonah" [Poona (= Pune), Maharashtra, India] [yellow star].
Type specimens: NHMUK 1947.3.4.15 and 1947.3.6.7 (formerly 1869.8.28.10, 1870.7.11.1), syntypes (2).
Etymology:* Eponym honoring Andrew H. Leith (1827–1882), United Kingdom / India; physician, naturalist, and malacologist who worked in India on the Bombay Sanitary Commission and collected the types.

Aspilus gataghol Gray 1872a:339
Trionyx gataghol, Trionyx gatajhal
Type locality: "India." Restricted to "the vicinity of Barrackpore" [West Bengal, India] [in error, orange star] by Webb (1980b:72).
Type specimen: NHMUK, holotype or syntypes, apparently lost, specimen figured in Gray (1835:pl.65), lectotype, designated as "type" by Webb (1980b).
Etymology:* Vernacular (Hindi?), *gotaghol* or *gataghol*, probably a local indigenous name for the species.

Trionyx sulcifrons Annandale 1915b:341
Type locality: "Nagpur,...Central Provinces of India." [Nagpur, Maharashtra, India] [green star].
Type specimen: ZSI 17973, holotype, see Das et al. (1998) and Kundu et al. (2018a).
Etymology:* Latin, *sulci*, furrows, and *frons*, forehead, front; referring to the several small longitudinal furrows or lines between the eyes.

Nilssonia nigricans (Anderson 1875) (07:81, 17:78)
Black Softshell Turtle, Bostami Softshell




Chittaranjan Baruah / TCC / Madhab Temple, Hajo, Assam, India





left: Peter Praschag / Brahmaputra R., Assam, India (juvenile)
right: Craig B. Stanford / Chittagong Temple, Bangladesh (adult)



** (orange dots and star = temple pond populations; star = type locality)

Distribution:* Bangladesh, India (*Assam, Tripura, West Bengal*), Nepal

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 118,106 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 304,963 sq. km

Size (Max SCL): male 81.0 cm, female 74.0 cm (Smith 1931; Ahsan and Saeed 1989; Ahsan et al. 1991); **(Max CCL):** male 91.0 cm (Das 1995; Pritchard 2001; Ernst et al. 2006b)

IUCN Red List: Critically Endangered (CR A4cd) (Praschag, Ahmed, and Singh 2021). Previously: Extinct in the Wild (EW) (Asmat 2002); Critically Endangered (CR) (ATTWG 2000); Critically Endangered (CR) (TFTSG *in* Baillie and Groombridge 1996)

CITES: Appendix I (1975), originally as *Aspideretes nigricans*

USA ESA:* Endangered, as *Trionyx nigricans* (USFWS 1976)

Synonymy:

Trionyx nigricans Anderson 1875:284
Amyda nigricans, Aspideretes nigricans, Nilssonia nigricans
Type locality:* "Tanks at Chittagong, Bengal" [Bangladesh] [orange star].
Type specimens: ZSI 664, 1898, syntypes (2), see Das et al. (1998), Das (2009), and Kundu et al. (2018a).
Etymology:* Latin, *nigricans*, to become black, blackish; referring to the dark carapace color.

Palea Meylan 1987 (89)*


Wattle-necked Softshell Turtles
(includes 1 species, 1 taxon)
Palea Meylan 1987:77
Type species: *Palea steindachneri* [= *Trionyx steindachneri* Siebenrock 1906a], by original designation.
Etymology:* Latin, *palea*, wattle, chin lobe, wart; referring to the wart-like formations on the neck.

Palea steindachneri (Siebenrock 1906a)
Wattle-necked Softshell Turtle




Torsten Blanck / Hong Kong, China





left: Haitao Shi / Hainan, China
right: Torsten Blanck / Hong Kong, China / juvenile



(orange dot = probable trade or introduced; star = type locality)

Distribution: China (*Guangdong, Guangxi, Guizhou, Hainan, Yunnan*), Laos, Vietnam

Introduced: China (*Hong Kong*), Mauritius, Réunion, USA (*Hawaii [Kauai]*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 319,021 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 854,815 sq. km

Size (Max SCL):* male 36.0 cm, female 44.5 cm (Pritchard 2001; Ernst and Bogadek 2005; Ernst and Lovich 2009)

IUCN Red List: Critically Endangered (CR A2cd+4cd) (Fong, Hoang, Li, McCormack, Rao, Timmins, and Wang 2021). Previously: Endangered (EN) (ATTWG 2000); Near Threatened (NT) (TFTSG *in* Baillie and Groombridge 1996)

CITES: Appendix II (2013). Previously: Appendix III (China) (2005)

Synonymy:

Aspidonectes californiana Rivers 1889:233 (*nomen suppressum*)
Pelodiscus californianus
Type locality: "Sacramento River, near the city of Sacramento... California" [USA] [in error, trade specimen] [no star].

Type specimen: UCMZ, holotype, apparently lost, see Webb (1975b).
Etymology:* Toponym for California, USA.
Comment: Name suppressed by ICZN (1982), see Webb (1978).

Trionyx steindachneri Siebenrock 1906a:579 (*nomen protectum*)
Amyda steindachneri, Palea steindachneri, Pelodiscus steindachneri
Type locality: "Kau-Kongriver...Insel Hainan" [China]; emended to "Kau-Kong (= Gaugong He) River, Hainan Island, Hainan Prov., China" [yellow star] by Zhao and Adler (1993:174).
Type specimen: NMW 20373, holotype, see Tiedemann and Häupl (1980), Tiedemann et al. (1994), and Gemel et al. (2019).
Etymology:* Eponym honoring Franz Steindachner (1834–1919), Austria; ichthyologist and director at the Naturhistorisches Museum, Vienna.
Comment: Name conserved by ICZN (1982), see Webb (1978).

Pelochelys Gray 1864b (89)*


Asian Giant Softshell Turtles
(includes 3 species, 3 taxa)


Pelochelys Gray 1864b:89 (17:4, 17:79, 21:5)
Type species: *Pelochelys cantorii* Gray 1864b, by subsequent designation by Günther (1865:108).
Etymology:* Greek, *pelos*, mud, clay, swamp, and *chelys*, turtle.

Pelochelys bibroni (Owen 1853)

Southern New Guinea Giant Softshell Turtle, Striped New Guinea Softshell Turtle


Anders G.J. Rhodin / TCF / CCB / Lake Murray, Western Prov., Papua New Guinea


left: Anders G.J. Rhodin / Laloki R., Central Prov., Papua New Guinea / neotype [AMS, Sydney]
right: Anders G.J. Rhodin / Lake Murray, Western Prov., Papua New Guinea


**

(star = type locality)

Distribution: Australia (?) (*Queensland, possibly vagrant [?]*); Indonesia (*Papua*), Papua New Guinea (*Southern*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 254,251 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 465,106 sq. km

Size (Max SCL): 102.0 cm (Rhodin et al. 1993; Pritchard 2001; Itescu et al. 2014)

IUCN Red List: Vulnerable (VU A2d+4d) (Georges, Rhodin, Eisemberg, and Amepou 2020). Previously: Vulnerable (VU) (ATTWG 2000); Vulnerable (VU) (TFTSG *in* Baillie and Groombridge 1996)

CITES: Appendix II, as *Pelochelys* spp. (2003)

Synonymy:


Trionyx (Gymnopus) bibroni Owen 1853:185
Trionyx bibroni, Pelochelys bibroni
Type locality: "Australian." Restricted to "Laloki River, Astrolabe Range, 40 miles [ca. 65 km] from its entry into Redscar Bay (9°20'S 147°14'E), Central District, Papua New Guinea" [yellow star] by Webb (1995a:302) by neotype designation.
Type specimen: NHMUK 1946.1.22.13, 1947.3.4.5 (formerly RCSM 954–959, 1093–94; single specimen in parts), holotype, destroyed in World War II, see Webb (1995a); AMS 3425–26, 131315 (single specimen, in parts), neotype, designated by Webb (1995a:302).
Etymology:* Eponym honoring Gabriel Bibron (1806–1848), France; herpetologist at the Muséum National d'Histoire Naturelle, Paris.

Pelochelys cantorii Gray 1864b (17:4, 17:79, 21:5)

Asian Giant Softshell Turtle, Cantor's Giant Softshell Turtle


Annette Olsson / CBFTT / TCC / nr. Kratie, Mekong R., Cambodia


Indraneil Das / CBFTT / Johore, Malaysia / hatchling, adult


(orange dots = possible trade or uncertain; stars = type localities)

Distribution: Bangladesh, Brunei, Cambodia, China (*Fujian, Guangdong, Guangxi, Hainan, Zhejiang*), India (*Kerala,*

Odisha, Tamil Nadu, West Bengal), Indonesia (*Kalimantan, Sumatra*), Laos, Malaysia (*East, West*), Myanmar, Philippines (*Luzon, Mindanao, Palawan*), Singapore (*extirpated*), Thailand, Vietnam

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 1,514,742 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 14,176,406 sq. km

Size (Max SCL): 100.0 cm (Das 2008 CBFTT)

CBFTT Account: Das (2008)

IUCN Red List: Critically Endangered (CR A2cd+4cd) (Choudhury, Das, Horne, Li, McCormack, Praschag, Rao, and Wang 2021). Previously: Endangered (EN) (ATTWG 2000); Vulnerable (VU) (TFTSG *in* Baillie and Groombridge 1996)

CITES: Appendix II, as *Pelochelys* spp. (2003)

Synonymy:

Pelochelys cantorii Gray 1864b:90
Type locality: "Malacca" [Malaysia] [yellow star].
Type specimen: NHMUK 1947.3.6.21–22 (parts of one specimen) (formerly 1860.4.19.1444), holotype, see Webb (1995) and Das (2009).
Etymology:* Eponym honoring Theodore E. Cantor (1809–1860), Denmark / India; physician and naturalist for the East India Company in Calcutta, who also accompanied British troops in China.

Pelochelys cumingii Gray 1864b:90
Chitra indica cumingii
Type locality: "Philippines" [green star, location estimated, possibly nr. Manila, see Webb 2003].
Type specimens: NHMUK 1946.1.22.13 and 1947.3.4.5, syntypes (2).
Etymology:* Eponym honoring Hugh Cuming (1791–1865), United Kingdom; businessman, naturalist, and collector who obtained the types.


Pelochelys cantoris Boulenger 1889:ix (*nomen novum*)
Etymology:* Eponym honoring Theodore E. Cantor (1809–1860), Denmark / India; physician and naturalist for the East India Company in Calcutta, who also accompanied British troops in China.
Comment: Unjustified emendation for *cantorii*.

Pelochelys poljakowii Strauch 1890:118
Type locality: "Fu-tschau" [China]. Emended to "Fu-tschau (= Fuzhou Shi), Fujian Province, China" [blue star] by Zhao and Adler (1993:174).
Type specimens: Possibly ZIN 7896–7, syntypes (2), type specimens figured (pl.4,f.1–3).
Etymology:* Eponym honoring Ivan S. Polyakov (1845–1887), Russia; zoologist, anthropologist, and ethnographer at the Zoological Museum of the Russian Academy of Sciences, St. Petersburg, who collected the types.

Chitra minor † Jaekel 1911:80
Type locality: "Pithecanthropus-schichten...Java...Trinil" [Indonesia] [gray star].
Type specimens: ZMB (MNB s/n or MB s/n), syntypes (2), fossil, plastral fragments, figured (pl.15,f.3-4), see Karl (1987) and McCord and Pritchard (2003a).
Geological age: Pleistocene, Trinil Beds.
Etymology:* Latin, *minor*, small; referring to its smaller size.

Pelochelys signifera Webb 2003
Northern New Guinea Softshell Turtle


Anders G.J. Rhodin / CCB / Maprik, East Sepik Prov., Papua New Guinea


(star = type locality)

Distribution: Indonesia (*Papua*), Papua New Guinea (*Northern*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO): 111,493 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 260,834 sq. km

Size (Max SCL): ca. 100.0 cm (Rhodin et al. 2018 IUCN)

IUCN Red List: Vulnerable (VU A4cde) (Rhodin, Richards, Georges, Amepou, and Hamidy 2018)

CITES: Appendix II, as *Pelochelys* spp. (2003)

Synonymy:

Pelochelys signifera Webb 2003:100
Type locality: "Wanggar River, Weyland Range, Geelvinck Bay, N. New Guinea (Papua Province, Indonesia)" [yellow star].
Type specimen: NHMUK 1921.11.11.4, holotype.
Etymology:* Latin, *signum*, sign, feature, and *ferre*, to bear, to carry, *signifera*, sign bearing; referring to the clear reticulated and dot pattern on the top of the head, neck and carapace in juveniles.

Pelodiscus Fitzinger 1835 (07:82, 10:32, 11:16) (89)*

Chinese Softshell Turtles

(includes 7 species, 7 taxa)

Pelodiscus Fitzinger 1835:110

Type species: *Aspidonectes (Pelodiscus) sinensis* [= *Trionyx (Aspidonectes) sinensis* Wiegmann 1834], by subsequent designation by Fitzinger (1843:30).

Etymology:* Greek, *pelos*, mud, clay, swamp, and *diskos*, disk, discus, plate; referring to the habitat and carapace shape.

Comment: Genus established as *Trionyx (Pelodiscus)* without a type species.

Landemania Gray 1869a:211

Type species: *Landemania irrorata* Gray 1869a [= subjective synonym of *Trionyx (Aspidonectes) sinensis* Wiegmann 1834], by original designation.

Etymology:* Anonym, *landemania*, possibly a fabricated name, or possibly an Eponym honoring someone named Landeman.

Psilognathus Heude 1880:24

Type species: *Psilognathus laevis* Heude 1880 [= subjective synonym of *Trionyx (Aspidonectes) sinensis* Wiegmann 1834], by original monotypy.

Etymology:* Greek, *psilos*, thin, narrow, fine, weak, and *gnathos*, jaw; referring to the narrow upper jaw.

Temnognathus Heude 1880:25

Type species: *Temnognathus mordax* Heude 1880 [= subjective synonym of *Trionyx (Aspidonectes) sinensis* Wiegmann 1834], by original monotypy.

Etymology:* Greek, *temnein*, to cut, divide, split, and *gnathos*, jaw; referring to the sharp-edged jaws.

Gomphopelta Heude 1880:27

Type species: *Gomphopelta officinae* Heude 1880 [= subjective synonym of *Trionyx (Aspidonectes) sinensis* Wiegmann 1834], by original monotypy.

Etymology:* Greek, *gomfos*, wedge, stake, and *pelta*, small, crescent-shaped leather shield; referring to the humped carapace.

Coelognathus Heude 1880:29

Type species: *Coelognathus novemcostatus* Heude 1880 [= subjective synonym of *Trionyx (Aspidonectes) sinensis* Wiegmann 1834], by original monotypy.

Etymology:* Greek, *koilos*, hollow, and *gnathos*, jaw; referring to the cavities in the cutting edges of the jaws.

Tortisternum Heude 1880:31

Type species: *Tortisternum novemcostatum* Heude 1880 [= subjective synonym of *Trionyx (Aspidonectes) sinensis* Wiegmann 1834], by original monotypy.

Etymology:* Latin, *tortus*, twisted, and Greek, *sternon*, chest, sternum, thorax; referring to the irregular plastral callosities.

Ceramopelta Heude 1880:33

Type species: *Ceramopelta latirostris* Heude 1880 [= subjective synonym of *Trionyx (Aspidonectes) sinensis* Wiegmann 1834], by original monotypy.

Etymology:* Greek, *keramos*, brick, roof tile, and *pelta*, small, crescent-shaped leather shield; referring to the patterned carapace.

Coptopelta Heude 1880:34

Type species: *Coptopelta septemcostata* Heude 1880 [= subjective synonym of *Trionyx (Aspidonectes) sinensis* Wiegmann 1834], by original monotypy.

Etymology:* Greek, *kopto*, truncated, cut off, and *pelta*, small, crescent-shaped leather shield; referring to the anteriorly truncated carapace.

Cinctisternum Heude 1880:36

Type species: *Cinctisternum bicinctum* Heude 1880 [= subjective synonym of *Trionyx (Aspidonectes) sinensis* Wiegmann 1834], by original monotypy.

Etymology:* Latin, *cinctum*, belt, belted, and Greek, *sternon*, chest, sternum, thorax; referring to the transverse plastral depression.

Pelodiscus axenaria (Zhou, Zhang, and Fang 1991) (10:32, 11:16, 21:102) (91)*


Hunan Softshell Turtle, Sand Softshell Turtle, Central Chinese Softshell Turtle



Shiping Gong / Yudai R., Tongdao Co., Hunan, China / neotype



Shiping Gong / Yudai R., Tongdao Co., Hunan, China / neotype



(yellow dots = genotyped; gray dots = presumed; stars = type localities)

Distribution: China (*Guangdong*, *Guangxi*, *Hunan*, *Jiangxi*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 119,355 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 241,305 sq. km

Size (Max SCL):* male 12.9 cm (Gong et al. 2022)

IUCN Red List: Not Evaluated (NE)

TFTSG Provisional Red List: Data Deficient (DD) (2011, 2018)

CITES: Appendix II (2013). Previously: Appendix III (China) (2005)

Synonymy:

Trionyx axenaria Zhou, Zhang, and Fang 1991:382 (91)*

Pelodiscus axenaria, Pelodiscus axenarius

Type locality:* "Taoyuan, Pingjiang, Rucheng, Lingling, Shaoyang Counties, Hunan Province, China" [in Chinese]. Erroneously restricted to "Xiang River...Quanzhou County, Guangxi, China" [orange star] by Yang et al. (2011:22). Restricted to "Yudai River, Tongdao County, Hunan Province, China (26º09'N, 109º46'E)" [yellow star] by neotype designation by Gong et al. (2022:132). Restricted to "Taoyuan section of the Yuanjiang River, Changde City, Hunan Province, China" [brown star] by lectotype designation by Liu et al. (2024:343), setting aside the earlier neotype designation, but refuted by Gong et al. (2025:98) as an inaccurate lectotype designation.

Type specimen:* Not designated or located, a specimen figured (f.1-2); JNU 20210001, neotype, genotyped and designated by Gong et al. (2022:132), but set aside as neotype by Liu et al. (2024:339) in favor of HNNU 1991002ZGJ, genotyped and inaccurately designated as a lost lectotype, but refuted by Gong et al. (2025), restoring the neotype designation.

Etymology:* Latin, *arenaria*, of sand; referring to the local vernacular name of the species, Sand Softshell Turtle, but published with a typographic error as *axenaria*.

Pelodiscus huike Liu, Fu, Ma, Zhou, Blanck, Zhou, Xiao, and Mo 2024:346 (21)*

Type locality:* "Yudai River, Tongdao County, Hunan Province, China (26°09 N 109°46 E)" [yellow star].

Type specimen:* JNU 20210001, holotype, previously genotyped by Gong et al. (2022) as designated neotype of *P. axenaria*.

Etymology:* Vernacular (Chinese), *huike bie*, local indigenous name meaning gray softshell turtle.

Comment: Objective junior synonym of *P. axenaria* (see Gong et al. 2025).

Pelodiscus huangshanensis Gong, Peng, Huang, and Nie *in* Gong, Peng, Huang, Lin, Huang, Xu, Yang, and Nie 2021 (21:102)

Huangshan Softshell Turtle, Horse-hoof Softshell Turtle



Yan-An Gong / Huangshan, Anhui, China / female



Yan-An Gong / Huangshan, Anhui, China / male



** (star = type locality)

Distribution: China (*Anhui, Zhejiang [?]*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 6,037 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 9,442 sq. km

Size (Max SCL):* male 9.1 cm, female 11.6 cm (Gong et al. 2021)

IUCN Red List: Not Evaluated (NE)

TFTSG Provisional Red List: Data Deficient (DD) (2021)

CITES: Appendix II (2013), as part of *P. axenaria*

Synonymy:

Pelodiscus huangshanensis Gong, Peng, Huang, and Nie *in* Gong, Peng, Huang, Lin, Huang, Xu, Yang, and Nie 2021:140

Type locality: "Xikou Town, Xiuning County, Huangshan City, Anhui Province, China (118.02°N, 29.69°E, and 170 m A.S.L.)." Geographic coordinates corrected here to "29.69ºN, 118.02ºE."

Type specimen: ANU 20210001, holotype.

Etymology:* Toponym for Huangshan City, Anhui Province, China.

Pelodiscus maackii (Brandt 1857) (10:32, 11:16, 17:80)


Amur Softshell Turtle, Northern Chinese Softshell Turtle



Dayoung Lee / Sillim-dong, Gwanak-gu, Seoul, South Korea



left: Norbert Schneeweiss / Lake Khanka, Russia
right: Haitao Shi / nr. Nagoya, Japan / captivity



** (orange dots = probable historical introductions or trade; star = type locality)

Distribution: China (*Heilongjiang, Jilin, Liaoning, Nei Monggu*), Japan (*historical introduction [?]*), North Korea, Russia (*Amurskaya, Khabarovskiy, Primorskiy, Yevreyskaya*), South Korea

Introduced: USA (*Hawaii [Oahu]*)

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 244,908 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 1,352,651 sq. km

Size (Max SCL):* 50.0 cm (Pavlov 1932; Kuzmin 2002; Gong et al. 2018)

IUCN Red List: Not Evaluated (NE)

TFTSG Provisional Red List: Data Deficient (DD) (2011, 2018)

CITES: Appendix II (2013). Previously: Appendix III (China) (2005)

Synonymy:

Trionyx maackii Brandt 1857:110

Amyda maackii, Pelodiscus maackii

Type locality: "in fluminibus australioribus lateralibus Amuris, nominatim in fluviis Sungari et Ussuri, nec non in ipsius fluminis


Amuris parte inter fluvios modo dictos obvia" [in the more southern rivers next to the Amur, namely the Sungari and Ussuri rivers, and also in the part of the Amur river between these rivers] [Russia]. Restricted to "Southern Amur R. and Ussuri R." [Russia] [yellow star, location estimated] by Stejneger (1907:529).
Type specimens:* ZIN 3690, 4661, 5725, 5726, syntypes (4).
Etymology:* Eponym honoring Richard K. Maack (or Maak) (1825–1886), Estonia / Russia; educator and naturalist who explored the Amur River.

Trionyx maaki Chkhikvadze 1987:610 (*nomen novum*)
Etymology:* Eponym honoring Richard K. Maack (or Maak) (1825–1886), Estonia / Russia; educator and naturalist who explored the Amur River.
Comment:* Unjustified emendation of *maacki*.

Pelodiscus parviformis Tang 1997 (10:32, 11:16, 21:103)
Lesser Chinese Softshell Turtle


Mian Hou / Guilin, Guangxi, China


Shiping Gong / Longshen, Guangxi, China



**
(stars = type localities)

Distribution: China (*Guangxi*, *Hunan*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 22,224 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 34,760 sq. km
Size (Max SCL): 16.0 cm (Farkas et al. 2019)
IUCN Red List: Not Evaluated (NE)
TFTSG Provisional Red List: Critically Endangered (CR) (2018)
CITES: Appendix II (2013). Previously: Appendix III (China) (2005)
Synonymy:
Pelodiscus parviformis Tang 1997:13
Type localities: "Range: Quanzhou, Xing'an, Guanyang, Ziyuan, Lingchuan counties of Guangxi Auto. Region; Dong'an, Qiyang, Daoxian counties of Hunan Province. In Quanzhou and Xing'an [yellow stars, locations estimated] region it was common" [China].
Type specimens:* GXUA 95011, holotype; 95012, allotype.
Etymology:* Latin, *parva*, small, insignificant, and *formis*, shape, form, appearance; referring to the small body size.

Pelodiscus shipian Gong, Fritz, Vamberger, Gao, and Farkas 2022 (91)*
Chinese Stone Slab Softshell Turtle


Shiping Gong / Taoyuan, Hunan, China [= *P. axenaria*]


Shiping Gong / Taoyuan, Hunan, China [= *P. axenaria*]


**
(yellow dots = genotyped, gray dots = presumed; stars = type localities)

Distribution:* China (*Anhui*, *Hunan*, *Jiangxi*)
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 112,646 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 197,236 sq. km
Size (Max SCL):* male 10.3 cm, female 10.1 cm (Gong et al. 2022)
IUCN Red List:* Not Evaluated (NE)
TFTSG Provisional Red List:* Data Deficient (DD) (2022)
CITES:* Appendix II (2013), as part of *P. axenaria*
Synonymy:*
Pelodiscus shipian Gong, Fritz, Vamberger, Gao, and Farkas 2022:134 (91)*
Type locality:* "Liaohe River, Fengxin County, Jiangxi Province, China (28°43'N, 115°25'E)" [yellow star].
Type specimen:* JNU 20190011, holotype, genotyped by Gong et al. (2022).
Etymology:* Vernacular (Chinese), *shi pian*, local name for the species; referring to its appearance as a flat stone slab.
Pelodiscus jiangxiensis Hou, Lin, Li, Jiang, and Shi 2023:3 (*nomen nudum*) (91)*
Type locality:* "PingDing Town, YuJiang District, Yingtan City, Jiangxi Province, China" [green star].
Type specimen:* HNU JY20220001, holotype, genotyped by Hou et

al. (2023).
Etymology:* Toponym for Jiangxi Province, China.
Comment:* Name posted online as a preprint, but then withdrawn and not published.


Pelodiscus sinensis (Wiegmann 1834) (10:32, 11:16, 14:30, 17:80, 17:81, 21:104) (92)*
Chinese Softshell Turtle



Tien-Hsi Chen / Keelung R., nr. Taipei, Taiwan



Marc Dupuis-Desormeaux / No data / trade



** (orange dots = probable trade, introduced, or uncertain identification; stars = type localities)

Distribution: China (*Anhui, Beijing, Fujian, Guangdong, Guangxi, Hebei, Henan, Hong Kong, Hubei, Hunan, Jiangsu, Jiangxi, Macau, Nei Mongol, Ningxia, Shaanxi, Shandong, Shanxi, Sichuan, Tianjin, Zhejiang*), Taiwan
Introduced: Bosnia and Herzegovina, Brazil (*Pará*), Croatia, Guam, Indonesia (*Java, Kalimantan, Sumatra, Timor*), Iran, Japan (*Bonin Islands, Honshu, Kyushu, Ryukyu Archipelago, Shikoku*), Kyrgyzstan, Laos, Malaysia (*East, West*), Northern Mariana Islands, Philippines (*Bohol, Cebu, Latvia, Leyte, Luzon, Mindanao, Mindoro, Panay*), Serbia, Singapore, Slovenia, South Korea, Spain, Thailand, Timor-Leste, USA (*Hawaii [Kauai, Oahu], Virginia*), Vietnam
Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 867,951 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 3,303,687 sq. km
Size (Max SCL): 23.0 cm (Farkas et al. 2019)
IUCN Red List: Vulnerable (VU A1d+2d [ver. 2.3]) (ATTWG 2000). Previously: Least Concern (LC) [Not Listed] (Behler 1996)
TFTSG Provisional Red List: Vulnerable (VU) *or* Endangered (EN) (2011)

Synonymy:

Testudo rostrata Thunberg 1787:179 (*nomen suppressum*)
Emydes rostrata, Trionyx rostratus
Type locality: Not known.
Type specimen: UPSZTY 128 (formerly UUZM 128 and UUZM 2), holotype, see Thunberg (1828), Lönnberg (1896), Webb (1985, 1990) and Wallin (2001).
Etymology:* Latin, *rostrum*, snout, beak, *rostrata*, beaked; referring to the prominent beak.
Comment: Name suppressed by ICZN (1991), see Webb (1990).

Testudo striata Suckow 1798:37 (*partim, nomen novum*)
Etymology:* Latin, *stria*, stripe, groove, *striata*, striped, grooved; referring to the striped or grooved shell pattern of this species.
Comment: Unjustified replacement name for *rostrata*. Name synonymized *in partim* under *Amyda cartilaginea cartilaginea, Pelodiscus sinensis*, and *Trionyx triunguis*.

Testudo semimembranacea Hermann 1804:219 (*nomen suppressum*)
Type locality: "Japania" [Japan] [in error, trade specimen].
Type specimen: Possibly ZMUS, not located.
Etymology:* Latin, *semi*, half, and *membrana*, membrane; referring to the shell being covered with a leather-like skin.
Comment: Name suppressed by ICZN (1963), see Wermuth (1956) and Mertens and Wermuth (1961).

Trionyx stellatus var. *Japon* Temminck and Schlegel 1834:pls.5,7 (invalid vernacular name) (10:18, 14:30)

Trionyx (Aspidonectes) sinensis Wiegmann 1834:189 (*nomen protectum*) (17:81)
Trionyx sinensis, Pelodiscus sinensis, Tyrse sinensis, Amyda sinensis, Trionyx sinensis sinensis, Amyda sinensis sinensis, Pelodiscus sinensis sinensis
Type locality: "kleinen Insel im Tigerflusse, dicht bei Macao" [on a small island in the Tiger River, close to Macau] [China] [yellow star].
Type specimen: ZMB 38, lectotype, designated and genotyped by Stuckas and Fritz (2011:337).
Etymology:* Toponym for China.
Comment: Name conserved by ICZN (1963), see Wermuth (1956) and Mertens and Wermuth (1961), and by ICZN (1991), see Webb (1990).

Trionyx japonica Temminck and Schlegel 1838:139 (14:30)
Amyda japonica, Pelodiscus sinensis japonicus
Type locality: "Japon" [Japan] [in error, trade specimen].
Type specimens: RMNH 3259, 3264, syntypes (2), discussed by Hoogmoed et al. (2010).
Etymology:* Toponym for Japan.

Trionyx tuberculatus Cantor 1842a:16 (*nomen nudum*) (17:82)
Etymology:* Latin, *tuberculum*, small hump, nodule; referring to the nodules on the carapace of juveniles.

Trionyx tuberculatus Cantor 1842b:482 (17:82)
Potamochelys tuberculatus, Amyda tuberculata, Trionyx sinensis tuberculatus, Amyda sinensis tuberculata, Pelodiscus sinensis tuberculatus
Type locality: "Chusan...Island" [Zhoushan Island, Zhejiang, China] [green star].
Type specimens: NHMUK 1946.1.22.87–89, syntypes (3).
Etymology:* Latin, *tuberculum*, small hump, nodule; referring to the nodules on the carapace of juveniles.

Tyrse perocellata Gray 1844:48
Trionyx perocellatus, Potamochelys perocellatus, Landemania perocellata, Gymnopus perocellatus
Type locality: "Canton, China" [Guangzhou, Guandong, China] [pink star].
Type specimens: NHMUK 1946.1.22.87–89, syntypes (3), same as *Trionyx tuberculatus*.
Etymology:* Latin, *per*, very, high grade, and *ocellata*, having eye spots, ocellated; referring to the carapace of juveniles covered

with numerous small eye spots.

Trionyx schlegelii Brandt 1857:111

Amyda schlegelii

Type locality: "China borealis...et Japonia" [China and Japan].
Type specimen: Not known or located.
Etymology:* Eponym honoring Hermann Schlegel (1804–1884), Germany / Netherlands; ornithologist and herpetologist who worked in Austria and the Netherlands and was director of the Rijksmuseum van Natuurlijke Historie, Leiden.

Landemania irrorata Gray 1869a:216

Type locality: "Shanghai" [China] [brown star].
Type specimen: NHMUK 1946.1.22.9 (formerly 1862.11.1.299), holotype.
Etymology:* Latin, *irrorata*, covered with dew, sprinkled; referring to the head and carapace being covered with small white spots.

Trionyx peroculatus Günther *in* Gray 1869a:216 (*nomen nudum*)

Etymology:* Latin, *per*, very, high grade, and *ocellata*, having eye spots, ocellated; referring to the carapace of juveniles covered with numerous small eye spots.
Comment:* Unjustified emendation or error for *perocellata*.

Gymnopus simonii Duméril *in* David 1875:214 (*nomen nudum*)

Localities:* "China." Emended to "China and Ning-Pô" [Ningbo, Zhejiang, China] [orange star] by Bour et al. (1995:88).
Specimens:* MNHN 835, 1691, not types, discussed by Bour et al. (1995).
Etymology:* Eponym honoring G. Eugène Simon (1829–1896), France / China; agricultural engineer and French consul who collected specimens in China that were labeled as a new species by A.H.A. Duméril at the Muséum National d'Histoire Naturelle, Paris.

Psilognathus laevis Heude 1880:24

Type locality: "la région montagneuse au sud de la ville de Ning-kouo fou" [Ningguo County, Anhui, China] [light gray star].
Type specimens: Not located, type specimens figured (pl.2).
Etymology:* Latin, *laevis*, smooth, delicate; referring to the smooth plastral callosities.

Temnognathus mordax Heude 1880:26

Type locality: "environs de Chang-hai" [Shanghai, China] [brown star].
Type specimens: Not located, type specimens figured (pl.3).
Etymology:* Latin, *mordax*, biting; referring to the extremely defensive biting behavior.

Gomphopelta officinae Heude 1880:27

Type locality: "la Houai, vers l'endroit où cette rivière sort de la province du Ho-nan" [Huai River, Henan, China] [dark gray star].
Type specimens: Not located, type specimens figured (pl.4).
Etymology:* Latin, *officinae*, coming from a pharmacy; referring to the fact that a pharmacist had used this animal to attract customers.

Coelognathus novemcostatus Heude 1880:29

Type locality: "l'extrémité orientale du lac Tch'ao" [Lake Chao, Anhui, China] [white star].
Type specimens: Not located, type specimens figured (pl.5).
Etymology:* Latin, *novem*, nine, and *costatus*, having ribs; referring to the nine pairs of carapacial costal bones.

Tortisternum novemcostatum Heude 1880:31

Type locality: "lac Tch'ao, département de Lu-tcheou" [Lake Chao, Anhui, China] [white star].
Type specimens: Not located, type specimens figured (pl.6).
Etymology:* Latin, *novem*, nine, and *costatus*, having ribs; referring to the nine pairs of carapacial costal bones.

Ceramopelta latirostris Heude 1880:33

Type locality: "environs de Ngan-k'ing fou" [Anqing Shi, Anhui, China] [blue star].
Type specimens: Not located, type specimens figured (pl.7).
Etymology:* Latin, *latus*, broad, wide, flat, and *rostrum*, snout, beak; referring to the broad head and snout.

Coptopelta septemcostata Heude 1880:35

Type locality: "lacs de Tong-lieou (Ngan-houé)" [Dongliu, Anhui, China] [purple star].
Type specimens: Not located, type specimens figured (pl.8).
Etymology:* Latin, *septem*, seven, and *costatus*, having ribs; referring to the seven pairs of carapacial costal bones.

Cinctisternum bicinctum Heude 1880:37

Type locality: "marais de Ngan-K'ing" [Anqing Shi, Anhui, China] [blue star].
Type specimens: Not located, type specimens figured (pl.9).
Etymology:* Latin, *bi-*, two, and *cinctus*, belted, banded, girdled; referring to the depressed plastron.

Trionyx cartilagineus newtoni Ferreira 1897:114

Trionyx sinensis newtoni

Type locality: "Timor" [Timor-Leste] [in error, trade specimen]
Type specimen: Probably MUHNAC, holotype, presumed lost.
Etymology:* Eponym honoring Francesco Newton (1864–1909), Portugal; botanist who collected the type.

Amyda schlegelii haseri Pavlov 1932:27

Type locality: Not designated. Restricted to "Tzu ya ho, the river of Sien hien (central Chili)" [Ziya He, Xian County, Hebei, China] [light green star] by Licent *in* Pavlov (1932:34).
Type specimen: TNZ s/n, holotype.
Etymology:* Eponym honoring Brother Nicolas Haser (1883–1900s), France / China; priest who worked in China and collected the species.

Amyda schlegelii licenti Pavlov 1932:28

Type locality: "Tientsin" [Tianjin, China] [black star].
Type specimen: TNZ 14, holotype.
Etymology:* Eponym honoring Père Émile Licent (1876–1952), France / China; priest and naturalist who worked in China and collected the species.

Amyda schlegelii laoshanica Pavlov 1933:3

Type locality: "Chantong, Laoshan, near Tsingtao" [Qingdao Shi, Shandong, China] [red star].
Type specimen: TNZ 638, holotype.
Etymology:* Toponym for Laoshan Mountain region, Shandong.

Pelodiscus variegatus Farkas, Ziegler, Pham, Ong, and Fritz 2019 (21:103) (92)*


Vietnamese Softshell Turtle, Spotted Softshell Turtle



Peter Paul van Dijk / Vietnam / trade; Cuc Phuong Turtle Conservation Center



Thomas Ziegler / *left*: Ha Tinh Prov., *right*: Quang Binh Prov., Vietnam / juvenile



** (star = type locality)

Distribution:* China (*Hainan*, *Yunnan [?]*), Laos (?), Vietnam

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 137,017 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 511,105 sq. km

Size (Max SCL): male 11.6 cm, female 23.0 cm (Farkas et al. 2019)

IUCN Red List:* Near Threatened (NT A4cde) (Luiselli, Pham, and Rhodin, in press). Previously: TFTSG Provisional Red List: Critically Endangered (CR) (2019)

CITES: Appendix II (2013), as part of *P. parviformis*

Synonymy:

Pelodiscus variegatus Farkas, Ziegler, Pham, Ong, and Fritz 2019:72

Type locality: "Thai Thinh village, Kinh Mon District, Hai Duong Province, Vietnam" [yellow star].

Type specimen: IEBR 4480, holotype.

Etymology:* Latin, *varius*, diverse, *variegatus*, spotted; referring to the variegated carapace and large plastral blotches.

Rafetus Gray 1864b (89)*

Bicallosite Softshell Turtles

(includes 2 species, 2 taxa)

Rafetus Gray 1864b:81

Type species: *Rafetus euphraticus* [= *Testudo euphratica* Olivier *in* Daudin 1801], by original monotypy.

Etymology:* Vernacular (Arabic), *rafesh*, local indigenous name for the species.

Oscaria Gray 1873g:157

Type species: *Oscaria swinhoei* Gray 1873g, by original monotypy.

Etymology:* Eponym honoring Samuel Oscar Gray (1873–?), United Kingdom; grandnephew of the author (see Bour *in* Pritchard 2012:54).

Yuen Heude 1880:18

Type species: *Yuen leprosus* Heude 1880 [= subjective synonym of *Oscaria swinhoei* Gray 1873g], by subsequent designation by Stejneger (1907:514).


Etymology:* Vernacular (Chinese), *yuen*, local indigenous name for the species.

Rafetus euphraticus (Olivier *in* Daudin 1801) (17:83, 17:84)


Euphrates Softshell Turtle



Hanyeh Ghaffari / CBFTT / Balarood R., Khuzestan Prov., Iran



left: Hanyeh Ghaffari / CBFTT / Karkheh Lake Dam, Khuzestan Prov., Iran
right: Asghar Mobaraki / Looreh, Dez R., Khuzestan Prov., Iran



(star = type locality)

Distribution: Iran (*Ilam*, *Khuzestan*, *Lorestan*), Iraq, Syria, Turkey

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO): 263,030 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 508,066 sq. km

Size (Max SCL): male 53.7 cm, female 57.0 cm (H. Ghaffari, unpubl. data); **(Max CCL):** male 68.0 cm (Taskavak 1998; Pritchard 2001; Taskavak et al. 2016 CBFTT)

CBFTT Account: Taskavak, Atatür, Ghaffari, and Meylan (2016)

IUCN Red List: Endangered (EN A4c) (Ghaffari, Taskavak, Turkozan, and Mobaraki 2017). Previously: Endangered (EN) (ERASG 1996)

CITES: Appendix II (2017)

Synonymy:

Testudo euphratica Olivier *in* Daudin 1801:305 (17:83)

Trionyx euphraticus, Gymnopus euphraticus, Rafetus euphraticus, Pelodiscus euphraticus, Amyda euphratica, Tyrse euphratica

Type locality: "le Tigre et l'Euphrate" [Tigris and Euphrates rivers, Iraq]. Restricted to "Euphrates (Al Firat), vicinity of Anah, Al-Anbar, Iraq" [yellow star] by Bour et al. (1995:85).

Type specimen: Not located, specimen figured in Olivier (1807:pl.41), holotype, designated by Bour et al. (1995:85).

Etymology:* Toponym for the Euphrates River, Iraq.

Testudo rafcht Olivier 1807:328

Type locality: "le Tigre..[&]..l'Euphrate" [Tigris and Euphrates rivers, Iraq] [no star].

Type specimen: Not located, specimen figured in Olivier (1807:pl.41), holotype, see Bour et al. (1995).

Etymology:* Vernacular (Arabic), *rafcht, rafesh*, local indigenous name for the species.

Testudo rascht Gray 1830e:19 (*nomen novum*)

Etymology:* Vernacular (Arabic), *rafcht, rafesh*, local indigenous name for the species.

Comment: Unjustified emendation or error for *rafcht*.

Tyrse rafeht Gray 1844:49 (*nomen novum*)

Trionyx rafeht, Testudo rafeht

Etymology:* Vernacular (Arabic), *rafcht, rafesh*, local indigenous name for the species.

Comment: Unjustified emendation or error for *rafcht*.

Rafetus swinhoei (Gray 1873g) (07:83, 10:33, 11:17, 17:85)


Red River Giant Softshell Turtle, Yangtze Giant Softshell Turtle, Swinhoe's Softshell Turtle, *Con Giai*




Timothy E.M. McCormack / TCC / Dong Mo Lake, Vietnam



Joel Sartore / China / captivity / Suzhou Zoo / *left*: female, *right*: male



**



(light red and red dots = extirpated; stars = type localities)

Distribution: China (*Anhui [extirpated], Jiangsu [extirpated], Yunnan, Zhejiang [extirpated]*), Vietnam

Estimated Indigenous (Historical) Ranges (AOO and EOO):

Est. Indigenous Area of Occupancy (AOO):* 95,481 sq. km

Est. Indigenous Extent of Occurrence (EOO):* 1,014,780 sq. km

Size (Max SCL): male 86.0 cm, female 109.5 cm (Le and Pritchard 2009; Pritchard 2012)

IUCN Red List: Critically Endangered (CR A2acd; D) (Fong, Hoang, Kuchling, Li, McCormack, Rao, D, Timmins, and Wang 2021). Previously: Critically Endangered (CR) (ATTWG 2000); Least Concern (LC) [Not Listed] (Behler 1996) [in error, was provisionally assessed as either Critically Endangered (CR) or Endangered (EN), but final determination was erroneously recorded as Least Concern (LC)]

CITES: Appendix II (2013). Previously: Appendix III (China) (2005)

Synonymy:

Oscaria swinhoei Gray 1873g:157

Pelodiscus swinhoei, Trionyx swinhoei, Amyda swinhoei, Rafetus swinhoei

Type locality: "neighbourhood of Shanghai" [China] [yellow star].

Type specimen: NHMUK 1946.1.22.9 and 1947.3.6.13 (shell and skull separate) (formerly 1873.7.30.125), holotype, discussed by Farkas and Fritz (1998).

Etymology:* Eponym honoring Consul Robert Swinhoe (1836–1877), India / United Kingdom / China; naturalist and diplomat who worked in Formosa (Taiwan) and China and collected birds, mammals, and reptiles.

Yuen leprosus Heude 1880:20

Type locality: "le Houang-p'ou, à Chang-hai" [Huangpu Jiang, Shanghai, China] [green star, partly hidden].

Type specimens: Not located; possibly NZMC, see Pritchard (2012).

Etymology:* Latin, *leprosus*, leprous; referring to the appearance of the carapace.

Yuen maculatus Heude 1880:22

Pelochelys maculatus

Type locality: "le Houang-p'ou" [Huangpu Jiang, Shanghai, China] [green star, partly hidden].

Type specimens: Not located, possibly NZMC, see Pritchard (2012); type specimens figured (pl.1,1a).

Etymology:* Latin, *macula*, spot, *maculatus*, spotted, speckled, stippled; referring to the yellowish pattern of spots on the carapace, head, neck and throat.

Yuen elegans Heude 1880:23

Type locality: "Houang-p'ou" [Huangpu Jiang, Shanghai, China] [green star, partly hidden].

Type specimens: Not located, possibly NZMC, see Pritchard (2012).

Etymology:* Latin, *elegans*, elegant, fine, select; referring to the nice markings on the carapace.

Yuen viridis Heude 1880:23

Type locality: "Grand Lac, à Sou-tcheou" [Tai Hu, Suzhou Shi,

Jiangsu, China] [blue star].
Type specimens: Not located, possibly NZMC, see Pritchard (2012).
Etymology:* Latin, *viridis*, green; referring to the greenish carapace.

Yuen pallens Heude 1880:23
Type locality: "Grand Lac, près de Sou-tcheou" [Tai Hu, Suzhou Shi, Jiangsu, China] [blue star].
Type specimens: Not known or located, possibly NZMC, see Pritchard (2012).
Etymology:* Latin, *pallens*, pale, yellowish; referring to the light coloring of the carapace.

Trionyx swinhonis Boulenger 1889:ix (*nomen novum*)
Etymology:* Eponym honoring Consul Robert Swinhoe (1836–1877), India / United Kingdom / China; naturalist and diplomat who worked in Formosa (Taiwan) and China and collected birds, mammals, and reptiles.
Comment: Unjustified emendation or error for *swinhoei*.

Pelochelys taihuensis † Zhang 1984:71
Type locality: "Luojiajiao, Tongxiang County, Zhejiang" [China] [dark gray star].
Type specimen: Holotype, not located, subfossil, skull and costal bone.
Geological age: Holocene, Neolithic, Luojiajiao Relics.
Etymology:* Toponym for Taihu Lake, on the border of Jiangsi and Zhejiang, China.

Trionyx liupani † Tao 1986:23
Type locality: "Penghu Channel in the Taiwan Strait" [Taiwan, China] [light gray star].
Type specimen: Holotype, private collection of Chung-Wu Pan; NTUM FR 0002–0003, plaster model, fossil, skull and hypoplastron.
Geological age: Late Pleistocene.
Etymology:* Eponym honoring Mong-Song Liu (?–) and Chang-Wu Pan (or Chung-Wu Pan), (?–) Taiwan; fossil collectors who owned the types.

Rafetus hoankiemensis Ha 1995:4 (*nomen nudum*)
Etymology:* Toponym for Hoan Kiem Lake, Hanoi, Vietnam.

Rafetus hoguomensis Ha 1995:4 (*nomen nudum*)
Etymology:* Toponym for Ho Guom Lake (= Hoan Kiem Lake), Hanoi, Vietnam.

Rafetus leloii Ha 2000:104 (11:17)
Type locality: "Hoan Kiem Lake, Ha Noi" [Vietnam] [pink star].
Type specimen: RHK 01-1967, holotype, discussed by Farkas and Webb (2003).
Etymology:* Eponym honoring Le Loi (1385–1433), Vietnam; ancient king who, according to legend, returned a magic sword to the sacred turtle of Hoan Kiem Lake in Hanoi after his victory over Chinese occupiers.

Rafetus vietnamensis Le, Le, Tran, Phan, Phan, Tran, Pham, Nguyen, Nong, Phan, Dinh, Truong, and Ha 2010:950 (10:33, 11:17)
Type locality: Not known; holotype from "Hung Ky Pagoda, Hanoi" [Hoan Kiem Lake, Hanoi, Vietnam] [pink star].
Type specimen: RHK 1, holotype, same specimen as holotype of *Rafetus leloii*.
Etymology:* Toponym for Vietnam.
Comment:* Objective junior synonym of *Rafetus leloii*, based on the same holotype for both taxa.

Trionyx Geoffroy Saint-Hilaire 1809a (89, 93)*

African Softshell Turtles
(includes 1 species, 1 taxon)

Trionyx Geoffroy Saint-Hilaire 1809a:363 (93)*
Type species: *Trionyx* "tortue du Nil" [= *Trionyx egyptiacus* Geoffroy Saint-Hilaire 1809a] [= subjective synonym of *Testudo triunguis* Forskål 1775], by subsequent designation by Bory de Saint-Vincent (1828:77).
Etymology:* Greek, *tris*, three, and *onyx*, claw, nail; referring to the three claws on the feet.

Aspidonectes Wagler 1830b:134
Type species: *Aspidonectes aegyptiacus* [= *Trionyx aegyptiacus* Geoffroy Saint-Hilaire 1809b] [= subjective synonym of *Testudo triunguis* Forsskål 1775), by subsequent designation by Fitzinger (1843:30).
Etymology:* Greek, *aspidos*, a shield, and *nektos*, swimming.

Gymnopus Duméril and Bibron 1835:472 (*nomen novum*)
Etymology:* Greek, *gymnos*, bare, naked, and *pus*, foot; referring to the inability to hide the legs under any plastral flaps, as present in *Cryptopus* (= *Cyclanorbis*).
Comment:* Unjustified replacement name for *Trionyx* and *Aspidonectes*.

Tyrse Gray 1844:47
Type species: *Tyrse aegyptiacus* [= *Trionyx aegyptiacus* Geoffroy Saint-Hilaire 1809b] [= subjective synonym of *Testudo triunguis* Forskål 1775], by original designation.
Etymology:* Vernacular (Arabic), *tirse*, local indigenous name for the species.

Fordia Gray 1869a:219
Type species: *Fordia africana* Gray 1869a [= subjective synonym of *Testudo triunguis* Forskål 1775], by original monotypy.
Etymology:* Eponym honoring George H. Ford (1809–1876), South Africa; naturalist artist who illustrated for Sir Andrew Smith and the British Museum of Natural History, London.

Trionyx triunguis (Forskål 1775) [(10:29, 11:18)]
African Softshell Turtle, Nile Softshell Turtle



Oguz Türkozan / Dalaman, Turkey







left: Luca Luiselli / Nigeria
right: Pearson McGovern / Senegal




**



**



**



**

(orange dots = introduced, trade, or uncertain; stars = type localities)

Distribution:* Angola, Benin, Burkina Faso, Cameroon, Central African Republic, Chad, Congo (DRC), Congo (ROC), Egypt, Equatorial Guinea, Eritrea, Ethiopia, Gabon, Gambia, Ghana, Greece (*vagrant*), Guinea, Guinea-Bissau, Israel, Ivory Coast (Côte d'Ivoire), Kenya, Lebanon, Liberia, Mali, Mauritania, Namibia, Niger, Nigeria, Senegal, Sierra Leone, Somalia, South Sudan, Sudan, Syria, Togo, Turkey, Uganda

Introduced: Israel (*inland*)

Mediterranean subpopulation: Egypt (*extirpated*), Greece (*vagrant*), Israel, Lebanon, Syria, Turkey

Estimated Indigenous (Historical) Ranges (AOO and EOO):
Est. Indigenous Area of Occupancy (AOO):* 2,198,985 sq. km
Est. Indigenous Extent of Occurrence (EOO):* 24,355,804 sq. km

Size (Max SCL):* male 108.4 cm, female 69.5 (?) cm (Taskavak and Akcinar 2009; McGovern et al. 2025 CBFTT). The previously recorded Max SCL of 120.0 cm by Herz and Rudolphi (2006) was based on full animal length, including head and neck (McGovern et al. 2025 CBFTT)

IUCN Red List: Vulnerable (VU A4bcd) (van Dijk, Diagne, Luiselli, Baker, Turkozan, and Taskavak 2017). Previously: Least Concern (LC) [Not Listed] (Behler 1996). Regional (Europe): Not Applicable (NA) (Cox and Temple 2009). Previously: Not Applicable (NA) (van Dijk, Tok, Ugurtas, Sevinç, Werner, Hraoui-Bloquet, Sadek, and Baha El Din 2004. Subpopulations (Mediterranean): Critically Endangered (CR C2a [ver. 2.3]) (ERASG 1996)

CITES: Appendix II (2017). Previously: Appendix III (Ghana)

(1976–2007)

Synonymy:

Testudo triunguis Forskål 1775:ix [10:29]

Amyda triunguis, Trionyx triunguis, Pelodiscus triunguis, Aspidonectes triunguis, Tyrse triunguis, Amyda triunguis triunguis

Type locality: "Nilo" [Nile River, Egypt] [yellow star, location estimated].

Type specimen: Not known or located, apparently lost (Webb, pers. comm. *in* Iverson 1992).

Etymology:* Latin, *tria*, three, and *unguis*, nail, claw; referring to the three claws on each of the feet.

Testudo striata Suckow 1798:37 (*partim*, *nomen novum*)

Etymology:* Latin, *stria*, stripe, groove, *striata*, striped, grooved; referring to the striped or grooved shell pattern of this species.

Comment: Unjustified replacement name for *triunguis*. Name synonymized *in partim* under *Amyda cartilaginea cartilaginea*, *Pelodiscus sinensis*, and *Trionyx triunguis*.

Trionyx egyptiacus Geoffroy Saint-Hilaire 1809a:366

Type locality: "l'Egypte" [Egypt]. Restricted to "the lower Nile River drainage" [Egypt] [yellow star, location estimated] by Bour et al. (1995:77).

Type specimen: MNHN 4147, lectotype, designated by Bour et al. (1995:77).

Etymology:* Toponym for Egypt.

Trionyx aegyptiacus Geoffroy Saint-Hilaire 1809b:12 (*nomen novum*)

Trionyx (Aspidonectes) aegyptiacus, Gymnopodus aegyptiacus, Gymnopus aegyptiacus

Etymology:* Toponym for Egypt.

Comment: Unjustified emendation or error for *egyptiacus*.

Testudo nilotica Shaw and Nodder 1810:180 (*nomen novum*)

Trionyx niloticus, Tyrse nilotica, Aspidonectes niloticus

Etymology:* Toponym for the Nile River, Egypt.

Comment: Unjustified replacement name for *aegyptiacus*.

Trionyx labiatus Bell 1835:unnumbered[text p.55]

Trionyx (Pelodiscus) labiatus

Type locality: "Africâ occidentali...Sierra Leone" [green star, location estimated].

Type specimen: OUM s/n, holotype, apparently lost, not listed by Nowak-Kemp and Fritz (2010); type specimen figured (part 4,pl.2-4).

Etymology:* Latin, *labium*, lip, *labiatus*, having lips; referring to the noticeably broad, fleshy lips.

Trionyx mortoni Hallowell 1844:120

Type locality: "Africa."

Type specimen: Possibly ANSP, holotype, apparently lost, not listed by Malnate (1971).

Etymology:* Eponym honoring Samuel G. Morton (1799–1851), USA; physician, paleontologist, and naturalist at the Academy of Natural Sciences, Philadelphia, who obtained the type.

Aspidonectes aspilus Cope 1860:295

Gymnopus aspilus

Type locality: "Rembo and Ovenga rivers, tributaries of the Fernando Vas, Equatorial West Africa" [Gabon] [blue star, location estimated].

Type specimen: Possibly ANSP or USNM, holotype, apparently lost, not listed by Cochran (1961), Malnate (1971), or Reynolds et al. (2007).

Etymology:* Greek, *a-*, without, and *spilos*, a spot or blemish; referring to the unpatterned shell border, extremities, neck, and head.

Fordia africana Gray 1869a:219

Type locality: "Upper Nile, Chartoum" [Sudan] [pink star].

Type specimens: NHMUK 1947.3.5.42, 1947.3.5.70, 1947.3.6.12 (skull from one specimen catalogued separately) (formerly 1862.9.2.33–34), syntypes (2).

Etymology:* Toponym for Africa.

Trionyx triunguis rudolfianus Deraniyagala 1948:30

Amyda triunguis rudolfianus

Type locality: "Ferguson's Gulf, Lake Rudolf, Africa" [Lake Turkana, Kenya] [white star].

Type specimen: UPSZTY 2821 (formerly UUZM 2821), holotype (A.G.J. Rhodin, unpubl. data).

Etymology:* Toponym for former Lake Rudolf (now Lake Turkana) in northern Kenya and southern Ethiopia.

TESTUDINES, sp. indet.

Named modern turtle taxa that cannot currently be identified as to what family or genus they may belong, or under what species they should be synonymized.

Testudo porphyrea Daudin 1801:142 (*nomen dubium et oblitum*)
Emys porphyrea, Emys (Clemmys) porphyrea
Type locality: "nouvelle Hollande" [Australia, in error?]
Type specimen: Not known or located, possibly in MNHN.
Etymology:* Greek, *porphyrea*, porphyry-like; referring to the reddish-ocher carapace with scattered small tawny and dark green speckles and spots similar to igneous porphyry rocks.
Comment: Described as being very similar to *Testudo scripta* Thunberg *in* Schoepff 1792 (= *Trachemys scripta scripta*) (Emydidae, USA), but synonymized under *Chelodina longicollis* (Chelidae, Australia) by Gray (1844, 1856b) before disappearing from later synonomies. No mention was made in the description of a long neck nor how many plastral scutes there were, but the two anterior marginals projected a bit, the tail had a small dorsal ridge, and there were tubercules around the base of the tail. It may not even have come from Australia.

Testudo costata Thunberg 1808:4 (*nomen nudum*)
Type locality: Not designated.
Type specimen: UUZM s/n, apparently lost, not listed by Lönnberg (1896), Holm (1957), or Wallin (2001).
Etymology:* Latin, *costata*, having ribs, ribbed; possibly referring to visible carapacial ribs in a juvenile specimen.
Comment: A specimen with this name attributed to Thunberg was donated to UUZM (Uppsala) by Thunberg (1808) from his personal collection and was still recorded as being in the collection as of Thunberg (1828), but no published description appeared and it has not been listed since. No details of its appearance are recorded, and what kind of turtle it was remains unknown.

Emys gronovii Merrem 1820:23 (*nomen dubium et oblitum*)
Emys gronouii
Type locality: Not known.
Type specimen: Not located, originally Museum Gronovianum 75.
Etymology:* Eponym honoring Laurentius Theodorus Gronovius (1730–1777), Netherlands; naturalist and ichthyologist who had a large private museum.
Comment: Description cited as sourced from Gronovius (1763:17, n.75), but *E. gronovii* was never recognized or synonymized by any subsequent authors, and what kind of turtle it was remains unknown.

Emyda pugnax Rafinesque 1822:3 (*nomen nudum*)
Emyda pugnar
Type locality: "United States" [USA].
Type specimen: Not known or located.
Etymology:* Latin, *pugnax*, pugnacious, warlike; referring to the aggressive behavior of this species.
Comment: Common name given as Fighting Tarapen.

Emyda morsitans Rafinesque 1822:3 (*nomen nudum*)
Emyda mordans
Type locality: "United States" [USA].
Type specimen: Not known or located.
Etymology:* Latin, *morsitans*, biting; referring to the aggressive behavior of this species.
Comment: Common name given as Biting Tarapen.

Emyda megalonyx Rafinesque 1822:3 (*nomen nudum*)
Emyda meg lonyx
Type locality: "United States" [USA].
Type specimen: Not known or located.
Etymology:* Greek, *megalos*, long, large, and *onyx*, claw, nail; referring to the long claws of this species.
Comment: Original name with a printing typo ("a" missing). Common name given as Long claw Tarapen.

Emyda striata Rafinesque 1822:3 (*nomen nudum*)
Type locality: "United States" [USA].
Type specimen: Not known or located.
Etymology:* Latin, *stria*, stripe, *striata*, striped; referring to the striped pattern of this species.
Comment: Common name given as Striped Tarapen.

Emyda granularis Rafinesque 1822:3 (*nomen nudum*)
Type locality: "United States" [USA].
Type specimen: Not known or located.
Etymology:* Latin, *granulum*, grain; probably referring to the rough surface of the shell.
Comment: Common name given as Granulated Tarapen.

Lepidemys Rafinesque 1822:3
Lepidemy
Type species: *Lepidemys bifida* Rafinesque 1822, by original monotypy.
Etymology:* Greek, *lepidotos*, covered with scales, and *emys*, turtle.
Comment: Original name with a printing typo ("s" missing).

Lepidemys bifida Rafinesque 1822:3 (*nomen nudum*)
Lepidemy bifida
Type locality: "United States" [USA].
Type specimen: Not known or located.
Etymology:* Latin, *bifida*, split in two, divided, bifid; possibly referring to a divided plastron, as in *Terrapene*.
Comment: Original name with a printing typo ("s" missing). Common name given as Bifid Tarapen.

SUPPLEMENT

Extinct Holocene Turtle and Tortoise Taxa Extant from ca. 10,000 BCE to 1500 CE

We include a supplementary listing here of those taxa of turtles and tortoises that persisted into the Holocene and prehistorical times, but became extinct between ca. 10,000 BCE (ca. 12,000 ybp) and 1500 CE. This listing has been updated since our previous fossil checklist (TEWG 2015), which also included species that became extinct earlier during the Pleistocene, and since our previous Holocene supplement in TTWG (2021). Maps in this supplement show point localities for the type and other referred specimens; distributional ranges shown are hypothesized only for the Holocene, not for earlier potentially larger ranges in the Pleistocene.

We record a total of 20 extinct Holocene taxa, of which 18 are described species or subspecies and two are still unnamed taxa; this represents a total increase of two species since TTWG (2021). All are terrestrial chelonians, most of them with large to giant body sizes, and most being vulnerable island forms; 16 are in the family Testudinidae (tortoises) and 2 in the extinct family Meiolaniidae (horned terrestrial chelonians). Most appear to have become extinct due to direct human subsistence hunting and overexploitation, although some may have also been adversely affected by climate change (see discussion in TEWG 2015).

The species and unnamed taxa recorded here are not included in our various lists or conservation status analyses elsewhere in this checklist, which concern only modern turtles and tortoises that have existed since 1500 CE.



Meiolaniidae and Testudinidae that became Extinct during the Holocene between ca. 10,000 BCE (ca. 12,000 ybp) and 1500 CE (map shows point localities and hypothesized Holocene ranges only, not earlier potentially larger Pleistocene ranges)

Order Testudines Batsch 1788

Turtles, Tortoises, and Terrapins; Chelonians

Suborder Meiolaniformes Sterli and de la Fuente 2013

Meiolaniform Chelonians

Meiolaniformes Sterli and de la Fuente 2013:839

Family Meiolaniidae Boulenger 1887b

Giant Horned Chelonians

Miolaniidae Boulenger 1887b:554

Meiolaniidae Simpson 1937:2

Comment:* Phylogenetic placement of the family Meiolaniidae remains somewhat controversial. Several authors have placed it on the stem of crown turtles, branching off before the Cryptodira–Pleurodira split (e.g., Joyce 2007; Sterli 2010, 2015; Sterli and de la Fuente 2013), but others (e.g., Gaffney 1983, 1985, 1996; Gaffney et al. 2007; Gaffney and Jenkins 2010) placed it among the Cryptodira. We list it here under Meiolaniformes, as recommended by Sterli and de la Fuente (2013), but the hierarchical placement of these suprageneric categories in our checklist is simply intended to be functional. We note also that Meiolaniidae are uniformly considered to be terrestrial chelonians most similar in skeletal adaptations to large modern tortoises (Sterli 2015).

Meiolania † Owen 1886

Giant Horned Tortoises

Meiolania † Owen 1886:315

Etymology:* Greek, *meion*, smaller, lesser, and *elaino*, wandering; referring to the mistaken assumption when described that this genus was a smaller relative of the fossil terrestrial monitor lizard genus *Megalania*.

Ceratochelys † Huxley 1887:232 (*nomen novum*)

Etymology:* Greek, *keras*, horn, and *chelys*, turtle; referring to the horned skull, and representing a turtle, not a lizard.

Comment: Unjustified replacement name for *Meiolania*.

Miolania † Boulenger 1887b:554 (*nomen novum*)

Etymology:* Greek, *meion*, smaller, lesser, and *elaino*, wandering.

Comment: Unjustified emendation of *Meiolania*.




Meiolania platyceps, Lord Howe Island, Australia [Late Pleistocene] [from Burke et al. 1983]


Meiolania mackayi † Anderson 1925

Walpole Island Giant Horned Tortoise

(Extinct, Early Holocene?**)**



Cranial horn core (Anderson 1925:pl.XXXII,f.5-6) / Walpole Island, New Caledonia





Distribution: New Caledonia (*Walpole Island*)

Estimated Possible Holocene Range: 4 sq. km
Size (Max SCL): ca. 70 cm [estimated] (Gaffney 1996)
Synonymy:
Meiolania mackayi † Anderson 1925:239
Type locality: "Walpole Island...about one hundred miles south-east from New Caledonia...22°38'S, 168°27'E."
Type specimen: AMF 17720, holotype, fossil, cranial horn core, figured (pl.32,f.5-6) and in Gaffney (1996:f.77c).
Geological age: Late Pleistocene to Early Holocene (Gaffney 1996).
Etymology:* Eponym honoring A.C. Mackay (1800s–1900s), Australia; engineer who worked for the Australian Guano Company on Walpole Island and collected the types in 1922.
Comment: Specimens found in phosphatic guano deposits in crevices in coral rock, presumed to be Late Pleistocene or Early Holocene. There is no evidence of early human presence on the island, but in view of its isolated and remote occurrence, this species was possibly the last Meiolaniidae to become extinct.

Genus indet.

Meiolaniidae sp. † [New Caledonia]
New Caledonia Giant Horned Tortoise
(Extinct, ca. 530 CE**)**


Seventh cervical vertebra (Gaffney et al. 1984:f.2) / New Caledonia



Distribution: New Caledonia (*New Caledonia Island, Tiga*)
Estimated Possible Holocene Range: 19,649 sq. km
Size (Max SCL): ca. 85 cm [estimated] (Gaffney et al. 1984; Gaffney 1996)
Synonymy:
Meiolaniidae, gen. et sp. indet. † (Gaffney et al. 1984:2)
Localities: "Main Pindai Cave, Nepoui Peninsula, New Caledonia, 21°20'12"S, 164°57'24"E." and "Tiga Island, Loyalty Group, New Caledonia, 21°8'S, 167°49'E."
Specimens: MNHN NCP 05 and NCT 01, subfossil, cervical vertebrae, figured (f.2).
Geological age: Late Holocene; ^{14}C age: 1720 ± 70 ybp, 230 CE ± 70 (160–300 CE), see Gaffney et al. (1984); calibrated age: 1820–1419 ybp (130–531 CE), see TEWG (2015).
Comment: This appears to be one of the last known Meiolaniidae to have survived into the Holocene; it occurred in association with modern humans and was evidently extirpated by them.

Suborder Cryptodira Cope 1869
Hidden-necked Turtles

Infraorder Durocryptodira Danilov and Parham 2006
Hardshell Hidden-necked Turtles

Superfamily Testudinoidea Batsch 1788
Testudinoid Turtles


Family Testudinidae Batsch 1788
Tortoises

Subfamily Manouriinae Gray 1869a
Manourine Tortoises

Manouria Gray 1854a
Indochinese Tortoises
Manouria Gray 1854a:133
Etymology:* Anonym, *manouria*, possibly a fabricated name, or possibly an Eponym honoring someone named Manour; or possibly Greek, *manos*, thin, and *ouria*, tail, but unlikely, since the specimens described were shells without soft parts.

Manouria oyamai † Takahashi, Otsuka, and Hirayama 2003
Ryukyus Tortoise
(Extinct, ca. 7800 BCE**)**


Akio Takahashi / Okinawajima, Japan / humerus



Distribution: Japan (*Ryukyu Archipelago*)
Estimated Possible Holocene Range: 23,779 sq. km
Size (Max SCL): ca. 45 cm [estimated] (Takahashi 2012; TEWG 2015; Takahashi et al. 2018)
Synonymy:
Manouria oyamai † Takahashi, Otsuka, and Hirayama 2003:198
Type locality: "Kamikurukubaru, Chinen village, Shimajiri District, Okinawa Island, Ryukyu Islands, Japan."
Type specimen: ESK 6150, holotype, fossil, incomplete skull, carapace, and appendicular bones, figured (f.4-9a).

Geological age: Late Pleistocene to Early Holocene, from ca. 28,160–23,050 ybp, until ca. 9865 ± 35 ybp, = ca. 7950–7880 BCE), see Otsuka et al. (2008) and Takahashi et al (2003).

Etymology:* Eponym honoring Seiho Oyama (1912–1996), Japan; businessman and amateur archaeologist who first collected this species.

Comment: Recorded from Okinawajima, Tokunoshima, Iejima, Miyakojima, and Yonagunijima islands in the Ryukyus. Migration patterns into the Ryukyu Archipelago of this and other Pleistocene taxa has been documented by Otsuka and Takahashi (2000) and Takahashi et al. (2008). Redescribed by Takahashi et al. (2018).

Subfamily Xerobatinae Gray 1874

North American Tortoises

Hesperotestudo † Williams 1950b

Western Hemisphere Tortoises

Eupachemys † Leidy 1877:232 (*nomen oblitum*)

Etymology:* Greek, *eu-*, genuine, correct, and *pachys*, thick, coarse, and *emys*, turtle; referring to the mistaken assumption that this taxon resembled the genus *Emys*, albeit with thicker carapace bones.

Comment:* Declared a *nomen oblitum* by TTWG (2021), corroborated and validated by Frétey (2024).

Hesperotestudo † Williams 1950b:25 (*nomen protectum*)

Etymology:* Greek, *hesperos*, the west, and Latin, *testudo*, turtle; referring to the range in North America, the "western world."

Comment:* Assigned to the subfamily Xerobatinae by Meylan and Sterrer (2000), although not corroborated by Vlachos and Rabi (2018). Name declared and validated as a *nomen protectum* by Frétey (2024).

Caudochelys † Auffenberg 1963b:69

Etymology:* Latin, *cauda*, tail, and Greek, *chelys*, turtle; referring to the tail without flattened or fused vertebrae.

Hesperotestudo crassiscutata † (Leidy 1889b)

North American Giant Tortoise, Southeastern Giant Tortoise

(Extinct, ca. 9500 BCE**)**



Jason R. Bourque, Florida Museum of Natural History / Sumter Co., Florida [UF 13406]



(red dot = Holocene locality; orange dots = Pleistocene localities)

Distribution: Holocene: USA (*Florida*)

Estimated Possible Holocene Range: 15,160 sq. km

Size (Max SCL): ca. 152 cm [estimated] (Auffenberg 1963b); 94.0 cm [measured] (Holman and Clausen 1984)

Synonymy:

Eupachemys obtusus † Leidy 1877:232 (*nomen oblitum*)

Eupachemys obtusa, Testudo obtusa

Type locality: "South Carolina...Ashley River, in the vicinity of Charleston" [USA].

Type specimen: ANSP 10197, holotype, subfossil, a single peripheral, figured (pl.34,f.4-5) and in Hay (1908b:f.614-615), see also Gillette (1977).

Geological age: Phosphate beds, exact age unknown, Cenozoic (Hay 1908b).

Etymology:* Latin, *obtusus*, blunted, blunt; referring to the single peripheral bone that has a blunt outer edge.

Comment: Hay (1908b) noted that *Eupachemys obtusus* was most similar to *Testudo crassiscutata*, and possibly synonymous and that *E. rugosus* Leidy 1889b was an erroneous name for *E. obtusus*; the names were subsequently synonymized by Auffenberg (1963b, 1974). The specific name *obtusus* is a *nomen oblitum*, not having been used since 1908, and should not replace the name *crassiscutata* through priority. Identification doubted by Vlachos (2018), who assigned it to Testudinidae sp. indet.

Eupachemys rugosus † Leidy 1889b:29 (*nomen novum et oblitum*)

Type locality: "Peace Creek, Florida" [USA].

Type specimen: Unknown.

Geological age: Late Pleistocene–Early Holocene, Rancholabrean.

Etymology:* Latin, *ruga*, wrinkle, rough; referring to the single peripheral bone with a rough surface.

Comment: Error for *Eupachemys obtusus* Leidy 1877. Vlachos (2018) suggested this may have been a new combination for a Recent *Emys rugosa*. However, Leidy simply cited the shape of the peripheral bones of "Red-leg Terrapin, *Emys rugosa*" as a comparison for the very large peripheral fossil bones he was describing, and stated that they were similar to the peripheral bone figured in Leidy (1877:pl.34,f.4-5) which he erroneously then referred to as *Eupachemys rugosus* rather than *E. obtusus*. The term "Red-leg Terrapin" in Pennsylvania during Leidy's time referred to a small freshwater turtle common in the local trade as a culinary substitute for the Diamondback Terrapin (*Malaclemys terrapin*), possibly the Wood Turtle, *Glyptemys insculpta*.

Testudo crassiscutata † Leidy 1889b:31

Geochelone crassiscutata, Geochelone (Caudochelys) crassiscutata, Caudochelys crassiscutata, Hesperotestudo crassiscutata, Hesperotestudo (Caudochelys) crassiscutata

Type locality: "Peace Creek, Florida" [USA]. Emended to "Peace River, near Arcadia, Hardee County, Florida, USA" by Vlachos (2018:39).

Type specimen: USNM 983, holotype, subfossil, partial plastron, figured (pl.6,f.4-7) and in Hay (1908b:f.616-617) and Auffenberg (1963b:f.10).

Geological age: Middle Pleistocene to Early Holocene, Rancholabrean, until ca. 9515 BCE, 11,465 ybp; ^{14}C age: 12,030 ± 200 ybp, see Clausen et al. (1979), calibrated age: 12,896–11,465 ybp (10,946–9515 BCE), see TEWG (2015).

Etymology:* Latin, *crassum*, thick, fat, coarse, dense, firm, and *scutata*, having a shield or scutes; referring to the strikingly thick carapace bones.

Comment: The species *T. crassiscutata* was well redescribed by Loomis (1927) and Auffenberg (1963b), who synonymized *T. sellardsi*, *T. luciae*, *T. ocalana*, and *T. distans* with *T. crassiscutata*. Clausen et al. (1979) and Holman and Clausen (1984) reported the find of a large specimen of *H. crassiscutata* from a Paleo-Indian site in Florida aged ca. 12,030 ybp that had been killed by humans. The specimen was associated with a large wooden stake that was still inside the body cavity and was radiocarbon-dated by Clausen et al. (1979).

Testudo ocalana † Hay 1916a:45

Gopherus ocalana, Geochelone ocalana

Type locality: "Ocala, Florida" [USA]. Emended to "Ocala, Marion County, Florida, USA" by Vlachos (2018:69).

Type specimen: USNM 8822, holotype, subfossil, an epiplastron, figured (pl.3,f.1) and in Auffenberg (1963b:f.18).

Geological age: Late Pleistocene.

Etymology:* Toponym for Ocala, Florida.

Testudo distans † Hay 1916a:48

Geochelone distans

Type locality: "Ocala, Florida" [USA]. Emended to "Ocala, Marion County, Florida, USA" by Vlachos (2018:66).

Type specimen: USNM 8819 (formerly FGS 4289), holotype, subfossil, an entoplastron, figured (pl.3,f.9) and in Auffenberg (1963b:f.19).

Geological age: Late Pleistocene.

Etymology:* Latin, *distans*, distant; probably referring to the extension of the pectoral scutes into the single entoplastron.

Testudo sellardsi † Hay 1916a:49

Geochelone sellardsi

Type locality: "Vero, St. Lucie County, Florida" [USA].

Type specimen: USNM 8817, holotype, subfossil, a xiphiplastron, figured (pl.8, f.6-8) and in Auffenberg (1963b:f.17).

Geological age: Late Pleistocene.

Etymology:* Eponym honoring Elias H. Sellards (1875–1961), USA; paleontologist and geologist at the University of Florida, Gainesville.

Testudo luciae † Hay 1916a:52

Geochelone luciae

Type locality: "Vero, St. Lucie County, Florida" [USA].

Type specimen: USNM 8818, holotype, subfossil, a hypoplastron, figured (pl.9,f.5) and iin Auffenberg (1963b:f.17).

Geological age: Late Pleistocene.

Etymology:* Toponym for St. Lucie County, Florida.


Hesperotestudo incisa † (Hay 1916a)

North American Dwarf Tortoise

(Extinct, ca. 6000 BCE**)**



Jason R. Bourque, Florida Museum of Natural History / Alachua Co., Florida [UF 3141]



(red dot = Holocene locality;
orange dots = Pliocene and Pleistocene localities)

Distribution: Holocene: USA (*Florida*)

Estimated Possible Holocene Range: 15,687 sq. km

Size (Max SCL): ca. 26 cm [*H. incisa*, estimated] (Bourque et al. 2012); 23.1 cm [*H. incisa*, measured] (Auffenberg 1963); ca. 20 cm [*H. mlynarskii*, estimated] (Auffenberg 1988); ca. 18 cm [*H. alleni*, estimated] (Auffenberg 1966)

Synonymy:

Testudo incisa † Hay 1916a:46

Gopherus incisa, Geochelone incisa, Geochelone (Hesperotestudo) incisa, Hesperotestudo incisa

Type locality: "Ocala, Marion county, Florida" [USA].

Type specimen: USNM (formerly FGS 4287), holotype, subfossil, a xiphiplastron, figured (pl.3,f.5).

Geological age: Rancholabrean, Sangamonian, Late Pleistocene.

Etymology:* Latin, *incisa*, incised; referring to the deep, rounded indentation on the posterior internal rim of the plastron.

Comment: Holman (1978) recorded specimens of this species from a sinkhole in Florida dated to approximately 7000–8000 ybp (although possibly older), during the Late Pleistocene or Early Holo-

cene. The species was synonymized with *Hesperotestudo turgida* (Cope 1892a) by Vlachos (2018), but in view of its markedly disjunct distribution from that species, and that its synonymization has not been universally accepted in the paleontological community, we continue to recognize it and the other synonymized taxa from Florida under *H. incisa* as distinct from *H. turgida*. In a preliminary study, Bourque et al. (2012) recognized *H. incisa*, *H. alleni*, and *H. mlynarskii* as a separate clade of small tortoises from Florida with a unique bony tail-shield buckler. Additionally, the shell shape of *H. incisa* differs markedly from that of *H. wilsoni* of the *H. turgida* clade.

Geochelone alleni † Auffenberg 1966:877

Geochelone (Hesperotestudo) alleni, Hesperotestudo alleni

Type locality: "McGehee Site, McGehee farm, T. 9 S., R. 17 E., S½NW¼ sec. 22, approximately 120 yards east of Florida Highway 45 and almost exactly 3 miles due north of Newberry, Alachua County, Florida" [USA].

Type specimen: UF 9370, holotype, fossil, partial carapace and complete plastron, figured (pl.102).

Geological age: Hemphillian faunal age, Middle Pliocene.

Etymology:* Eponym honoring Robert Allen (c.1940–), USA; museum technician and amateur paleontologist who worked at the Florida Museum of Natural History, Gainesville.

Comment: This species was synonymized with *Hesperotestudo turgida* (Cope 1892a) by Vlachos (2018), but is more likely synonymous with *H. incisa*.

Geochelone mlynarskii † Auffenberg 1988:592

Geochelone (Hesperotestudo) mlynarskii, Hesperotestudo mlynarskii

Type locality: "Coleman IIA site (SE 1/4, NW 1/4, Sec 7, T 20 S, R 23 E) near Coleman, Sumter County, Florida, U.S.A."

Type specimen: UF 18960, holotype, fossil, plastron, figured (pl.29).

Geological age: Late Irvingtonian mammal age, Middle Pleistocene.

Etymology:* Eponym honoring Marian Młynarski (1926–), Poland; paleontologist and herpetologist at the Institute of Systematics and Evolution of Animals, Polish Academy of Sciences.

Comment: This species was synonymized with *Hesperotestudo turgida* (Cope 1892a) by Vlachos (2018), but is more likely synonymous with *H. incisa*.


Hesperotestudo turgida † (Cope 1892a)

North American Plains Tortoise

(Extinct, ca. 9000 BCE**)**



Chris Sagebiel, Texas Vertebrate Paleontology Collections / Friesenhahn Cave, Bexar Co., Texas / *Hesperotestudo wilsoni*, TxVP 933-3732 / paratype



(red dots = Holocene localities;
orange dots = Pliocene and Pleistocene localities)

Distribution: Holocene: USA (*New Mexico, Oklahoma, Texas*)

Estimated Possible Holocene Range: 191,132 sq. km

Size (Max SCL): ca. 28 cm [*H. oelrichi*, estimated] (Holman 1972); ca. 24 cm [*H. turgida*, estimated] (Oelrich 1957); 23.5 cm [*H. johnstoni*, measured] (Auffenberg 1962); 22.6 cm [*H. wilsoni*, measured] (Milstead 1956); 17.0 cm [*H. riggsi*, measured] (Holman 1972)

Synonymy:

Testudo turgida † Cope 1892a:127

Gopherus turgida, Geochelone turgida, Geochelone (Hesperotestudo) turgida, Hesperotestudo turgida

Type locality: "Espuella near Dockum, from the same horizon as that of Crosby county, or the Blanco Canyon bed" [Texas, USA]. Corrected to "Three mi. N Dockum, near Esquella, Dickens County, Texas" [USA] by Auffenberg (1962:630).

Type specimen: ANSP 14689, holotype, fossil, fragments of a carapace, plastron, and mandible, figured in Oelrich (1957:f.1).

Geological age: Blanco beds, Blancan mammal age, Piacenzian, Late Pliocene.

Etymology:* Latin, *turgida*, swollen; referring to the shell bones appearing somewhat swollen.

Comment: Both Kuhn (1964) and Auffenberg (1974) erroneously

cited *Emys turgidus* Cope 1870b as the original name for this tortoise, but that name described a Cretaceous dermatemyid turtle from New Jersey, now synonymized under *Agomphus* (Hutchison and Weems 1998; Knauss et al. 2011). *Hesperotestudo turgida* was evaluated by Vlachos (2018) who synonymized the other taxa listed below and the taxa that we synonymize under *H. incisa*, but his synonymizations have not been universally accepted in the paleontological community. Based on the markedly disjunct distribution it is likely that *H. turgida* (including the synonymized taxa *riggsi*, *wilsoni*, *johnstoni*, and *oelrichi*) is restricted to the west and the Great Plains, and that *H. incisa* (including the synonymized taxa *alleni* and *mlynarskii*) is restricted to the east (Florida and Georgia).

Testudo riggsi † Hibbard 1944:72

Gopherus riggsi, *Geochelone riggsi*, *Geochelone (Hesperotestudo) riggsi*, *Hesperotestudo riggsi*

Type locality: "Locality No. 6, Seward county, Kansas" [USA]. Emended to "Univ. Kansas locality 6, Seward Co., Kansas, Sec. 36, T. 34 S., R. 31 W., XI Ranch, S side of Cimarron River" [USA] by Oelrich (1957:230).

Type specimen: KU 6789, holotype, fossil, nearly complete carapace and plastron, figured (f.1-2) and in Oelrich (1957:f.2-3).

Geological age: Saw Rock Canyon fauna, Middle Pliocene.

Etymology:* Eponym honoring Elmer S. Riggs (1869–1963), USA; paleontologist and curator at the Field Museum of Natural History, Chicago.

Comment: This species was synonymized with *Hesperotestudo turgida* (Cope 1892a) by Vlachos (2018).

Testudo wilsoni † Milstead 1956:168

Geochelone wilsoni, *Geochelone (Hesperotestudo) wilsoni*, *Hesperotestudo wilsoni*

Type locality: "Friesenhahn Cave in northern Bexar County, Texas" [USA].

Type specimen: TMM 933-3585, holotype, subfossil, an almost complete shell, lost; TxVP 933-3732 (formerly TMM), paratype, subfossil, complete shell, figured in Moodie and Van Devender (1979:f.1).

Geological age: Rancholabrean, Late Pleistocene, Late Wisconsinan to Early Holocene; ^{14}C age: ca. 11,040 ybp, 9090 BCE, see Auffenberg (1974) and Moodie and Van Devender (1979).

Etymology:* Eponym honoring John A. Wilson (1914–2008), USA; paleontologist and geologist at the University of Texas, Austin.

Comment: Discussed by Moodie and Van Devender (1979) who noted radiocarbon dating of specimens up to ca. 11,000–10,000 ybp in the Early Holocene of Oklahoma, New Mexico, and Texas, between Clovis and Folsom cultural horizons (Early Archaic). Preston (1979) noted specimens found associated with early man from the Plano Culture of Oklahoma at ca. 11,000 ybp. This species was synonymized with *Hesperotestudo turgida* (Cope 1892a) by Vlachos (2018), but the synonymization has not been universally accepted in the paleontological community.

Testudo rugosa † Johnston *in* Auffenberg 1962:627 (*nomen nudum*)

Etymology:* Latin, *ruga*, wrinkle, rough; referring to the rough surface of the shell.

Geochelone johnstoni † Auffenberg 1962:627

Geochelone (Hesperotestudo) johnstoni, *Hesperotestudo johnstoni*

Type locality: "Cita Canyon, Tule County, Texas" [USA]. Corrected to "Randal County, Texas" [USA] by Holman (1972:66).

Type specimen: PPHM 1540, holotype, fossil, shell, figured (f.1).

Geological age: Cita Canyon series, Early Pleistocene, probably between Nebraskan and early Kansan in age (Aftonian?).

Etymology:* Eponym honoring C. Stuart Johnston (1900–1939), USA; paleontologist who collected the type and described the species in an unpublished dissertation.

Comment: This species was synonymized with *Hesperotestudo turgida* (Cope 1892a) by Vlachos (2018).

Geochelone (Hesperotestudo) oelrichi † Holman 1972:59

Geochelone oelrichi, *Hesperotestudo oelrichi*

Type locality: "northwest corner of the Wilbur Magill pasture on the north side of a short reentrant of the main draw in the NW. 1/4, SE. 1/4, sect. 12, T. 30 N., R. 21 W., Brown County, Nebraska, Long Pine Quadrangle." [USA].

Type specimen: UMMP V56298, holotype, fossil, nearly complete carapace and plastron and appendicular bones, figured (f.20-24).

Geological age: Long Pine Formation, Early Pleistocene.

Etymology:* Eponym honoring Thomas M. Oelrich (1924–2008), USA; anatomist, histologist, and paleoherpetologist at the University of Michigan Medical School.

Comment: This species was synonymized with *Hesperotestudo turgida* (Cope 1892a) by Vlachos (2018).

Subfamily Testudininae Batsch 1788

Testudine Tortoises

Aldabrachelys Loveridge and Williams 1957

Indian Ocean Giant Tortoises


Aldabrachelys Loveridge and Williams 1957:225

Etymology:* Toponym for Aldabra Atoll, Seychelles, and *chelys*, turtle.


Aldabrachelys abrupta † (Grandidier 1868) [75]*

Madagascan Giant Tortoise

(Extinct, ca. 830–1570 CE**)**



Roger Bour / Madagascar




(star = type locality)

Distribution: Madagascar
Estimated Possible Holocene Range: 292,101 sq. km
Size (Max SCL): ca. 125 cm [estimated], 115.0 cm [measured] (Bour 1994)
Synonymy:
Testudo abrupta † Grandidier 1868:377
Geochelone abrupta, Geochelone (Asterochelys) abrupta, Asterochelys abrupta, Dipsochelys abrupta, Aldabrachelys abrupta
Type locality: "terrain marécageux à Amboulitsate, sur la côte occidentale de Madagascar." Emended to "marais d'Amboulitsate (Ambohisatrana) [= Ambolisatra], canton de Maromiandra, Madagascar" [red star] by Bour (1994:136).
Type specimen:* MNHN(P) MAD3500, lectotype, subfossil, incomplete carapace, designated by Bour (1994:136), figured in Bour (1994:f.33-35), see also Arnold (1979) and Bour (1985), genotyped by Kehlmaier et al. (2023).
Geological age: Late Holocene; ^{14}C age: 750 ± 370 ybp, 1200 CE ± 370 (830–1570 CE), see Burleigh and Arnold (1986); calibrated age: 1358–55 ybp (592–1895 CE), see TEWG (2015).
Etymology:* Latin, *abrupta*, cut off, abrupt, steep; referring to the marginal scutes angled perpendicularly to the plastron.
Comment: This species went extinct after the first arrival of humans from continental Africa, who reached Madagascar approximately 2000 years ago (Pedrono 2008). Radiocarbon dating of bones from the species indicate a range of dates that suggest possible persistence of occurrence into the very earliest part of the modern era, just after 1500 CE. Three uncalibrated ^{14}C aged specimens are known: 1) the youngest, at 750 ± 370 ybp (ca. 1200 CE ± 370 yrs = 830–1570 CE; Burleigh and Arnold 1986); 2) an intermediate specimen, 1910 ± 120 ybp (ca. 40 CE ± 120 yrs = 80 BCE–160 CE; Mahé and Sourdat 1972); and 3) the oldest, at 2035 ± 35 ybp (ca. 85 BCE ± 35 yrs = 120–50 BCE; Burleigh et al. 1982). The calibrated ages for these three specimens, respectively, calculate as: 1) 592–1895 CE [less precise range due to younger age]; 2) 197 BCE–390 CE; and 3) 163 BCE–52 CE. Although it is conceivable that the youngest of these specimens was from a tortoise that actually lived in the 1500s or later, this is unlikely, and we presume that the species may have gone extinct sometime between 1200 and 1300 CE. The skull was redescribed by Gerlach (2008), who noted also that this species was apparently sympatric with *A. grandidieri*.

Aldabrachelys grandidieri † (Vaillant 1885b)
Grandidier's Giant Tortoise
(Extinct, ca. 900 CE**)**


Roger Bour / Madagascar


Uwe Fritz / Madagascar / *left*: paralectotype; *right*: lectotype


** (star = type locality)

Distribution: Madagascar
Estimated Possible Holocene Range:* 184,891 sq. km
Size (Max SCL): ca. 135 cm [estimated], 124.0 cm [measured] (Bour 1994); **(Max CCL):** 152.0 cm (Vaillant 1885b)
Synonymy:
Emys gigantea † Grandidier 1868:378 (junior homonym [as *Testudo gigantea*], not *Testudo gigantea* Schweigger 1812)
Testudo gigantea
Type locality: "Etséré...Madagascar" [red star].
Type specimen: MNHN(P) MAD3501, lectotype (see below), subfossil.
Geological age: Late Holocene.
Etymology:* Greek, *gigas*, huge, gigantic; referring to the considerable body size.
Comment: Name published almost simultaneously by Milne-Edwards (1868), see Bour (1994).
Testudo grandidieri † Vaillant 1885b:876 (*nomen novum*)
Geochelone grandidieri, Dipsochelys grandidieri, Aldab-

rachelys grandidieri, Megalochelys grandidieri

Type locality: "Etséré...Madagascar." Emended to "Etseré (= Itsere), Belavenoka, canton de Morombe, Madagascar" [red star] by Bour (1994:137).

Type specimen:* MNHN(P) MAD3501 (formerly 1884.30A), lectotype, subfossil, complete carapace, designated by Bour (1985:51), figured by Bour (1985:f.10) and Bour (1994:f.33-35), see also Boulenger (1894) and Arnold (1979); MNHN(P) MAD3502 (formerly 1884.30B), paralectotype, genotyped by Kehlmaier et al. (2023).

Geological age: Late Holocene; ^{14}C age: 1250 ± 50 ybp (650–750 CE), see Burleigh and Arnold (1986), calibrated age: 1282–1066 ybp (668–884 CE), see TEWG (2015).

Etymology:* Eponym honoring Alfred Grandidier (1836–1921), France / Madagascar; naturalist and explorer who worked in Madagascar and first described this species.

Comment: Justified replacement name for *Testudo gigantea* (Grandidier 1868). This species went extinct shortly after the first arrival of humans from continental Africa, who reached Madagascar approximately 2000 years ago (Pedrono 2008). The most recent dated specimen recorded was from ca. 1250 ybp (Burleigh and Arnold 1986).

Testudo madagascariensis † Vaillant *in* Rothschild 1915:pl.34 (*nomen novum*)

Etymology:* Toponym for Madagascar.

Comment: Unjustified replacement name for *Testudo grandidieri* † Vaillant 1885b.

Astrochelys Gray 1873j [75, 94]*

Madagascan Tortoises

Astrochelys rogerbouri † Kehlmaier, Graciá, Ali, Campbell, Chapman, Deepak, Ihlow, Jalil, Pierre-Huyet, Samonds, Vences, and Fritz 2023 [94]*

Bour's Malagasy Tortoise

(Extinct, ca. 900 CE**)**



Uwe Fritz / Ambolisatra, Madagascar / holotype / tibia



(star = type locality)

Distribution:* Madagascar

Estimated Possible Holocene Range:* 17,975 sq. km

Size (Max SCL):* ca. 50 cm [estimated] (Kehlmaier et al. 2023)

Synonymy:*

Astrochelys rogerbouri † Kehlmaier, Graciá, Ali, Campbell, Chapman, Deepak, Ihlow, Jalil, Pierre-Huyet, Samonds, Vences, and Fritz 2023:6

Type locality:* "Ambolisatra, Madagascar" [red star].

Type specimen:* MNHN.F.MAD480, holotype, subfossil, nearly complete tibia, figured (f.3,S4), genotyped by Kehlmaier et al. (2023).

Geological age:* Late Holocene; ^{14}C age: 1233 ± 28 ybp (689–745 CE), calibrated age: 1262–1069 ybp (688–881 CE) (Kehlmaier et al. 2023).

Etymology:* Eponym honoring Roger Bour (1947–2020), France; herpetologist, paleontologist, and turtle taxonomist, former member and co-author in the Turtle Taxonomy Working Group, and expert on western Indian Ocean giant tortoises.

Comment:* Described from a single tibia, previously interpreted as belonging to a subadult extinct *Aldabrachelys abrupta* †, but shown to be deeply genetically divergent from that species and instead an extinct taxon sister to extant *Astrochelys yniphora*.


Chelonoidis Fitzinger 1835

South American Tortoises, Neotropical Tortoises

Chelonoidis alburyorum † Franz and Franz 2009 (21:105)

Bahama Banks Tortoise, Albury's Tortoise

(includes 3 subspecies)



(subspecies: *alburyorum* = red; *keegani* = purple (Turks), *sementis* = blue)

Distribution: Bahamas (*Acklins, Andros, Crooked Island, Eleuthera, Flamingo Cay, Great Abaco, Great Inagua, Long Island, Mayaguana, Moore's Island, New Providence, San Salvador*); Turks and Caicos (*Grand Turk, Middle Caicos*)

Estimated Possible Holocene Range: 16,574 sq. km

Size (Max SCL): male 71.0 cm, female 44.7 cm (see subspp.)

Chelonoidis alburyorum alburyorum † Franz and Franz 2009 (21:105)

Abaco Tortoise

(Extinct, ca. 1200 CE**)**



Nancy Albury / Great Abaco Island, Bahamas [Albury et al. 2018:f.6,B,D,F]



(subspecies: *alburyorum* = red)

Distribution: Bahamas (*Acklins, Andros, Crooked Island, Eleuthera, Flamingo Cay, Great Abaco, Great Inagua, Long Island, Mayaguana, Moore's Island, New Providence, San Salvador*)

Estimated Possible Holocene Range: 15,814 sq. km

Size (Max SCL): male 46.6 cm, female 42.8 cm (Franz and Franz 2009; Albury et al. 2018); **(Max CCL):** male 61.0 cm, female 55.0 cm (Franz and Franz 2009; Albury et al. 2018)

Synonymy:

Chelonoidis alburyorum † Franz and Franz 2009:5

Chelonoidis alburyorum alburyorum

Type locality: "Sawmill Sink (BNM Site Number AB-50), north of Cornwall Point, 37 km (23 mi) S of Marsh Harbour, Great Abaco Island, Little Bahamas Bank, The Bahamas (26º17' N, 77º12' W)."

Type specimen: UF 225400, holotype, subfossil, skull, shell, and associated appendicular elements, figured (f.2-15); topotypic material genotyped by Kehlmaier et al. (2021a).

Geological age: Late Holocene; ^{14}C cal age: ca. 930 BCE–1110 CE (Steadman et al. 2020).

Etymology:* Eponym honoring Nancy and Michael Albury (1955– and 1958–), Bahamas / USA; paleontologists and naturalists who helped curate and preserve the types.

Comment: Described from remarkably intact shells from an underwater sinkhole, with ^{14}C age of 2720–2580 ± 50 ybp (820–580 BCE); calibrated age: 2770–2500 ybp (820–550 BCE). Recent finds of charred fragments of this species from Lucayan Taíno culture middens in peat deposits at Gilpin Point, Abaco Island (Steadman et al. 2014; Hastings et al. 2014), are calculated to ^{14}C age of 960–910 ybp and 920–780 cal. ybp (ca. 1170 CE), shortly after the arrival of humans on Abaco at ca. 950 ybp. Other tortoise fragments from the Pleistocene or Holocene of Bahamas (Banana Hole on New Providence and on Andros), described by Auffenberg (1967) as *Geochelone* sp., may represent the same species. Discussed also by Franz and Franz (2009), Vlachos (2018), Albury et al. (2018, 2020), Steadman et al. (2020), and Franz et al. (2020). Successful extraction of ancient mitochondrial DNA from bones of this species by Kehlmaier et al. (2017) demonstrated its close phylogenetic relationship with modern *C. chilensis* and the *C. niger* complex from the Galápagos, and more distant relationship to the other modern South American *Chelonoidis* species, *C. carbonarius* and *C. denticulatus*.


Chelonoidis alburyorum keegani † Franz, Albury, and Steadman 2020 (21:105)

Turks Tortoise, Keegan's Tortoise

(Extinct, ca. 1200 CE**)**



plastron, humerus / Great Turk, Turks and Caicos [Franz et al. 2020:f.8,16]



(subspecies: *alburyorum* = red; *keegani* = purple (Turks), *sementis* = blue)

Distribution: Turks and Caicos (*Grand Turk Island*)

Estimated Possible Holocene Range: 50 sq. km

Size (Max SCL): male 71.0 cm, female 44.7 cm [estimated] (Franz et al. 2020)

Synonymy:

Chelonoidis alburyorum keegani † Franz, Albury, and Steadman 2020:13

Type locality: "Coralie Archaeological Site (GT-3)...21°30'N, 71°08'W...west shore of North Creek...northern end of Grand Turk Island, Turks and Caicos Islands, British Overseas Territory."

Type specimen: UF 453000, holotype, subfossil, complete plastron, figured (f.8A,C); topotypic material genotyped by Kehlmaier et al. (2021a).

Geological age: Late Holocene, ^{14}C cal age: ca. 700–1200 CE (Carlson 1999; Steadman et al. 2020).


Etymology:* Eponym honoring William F. Keegan (1959–), USA; archaeologist and curator at the Florida Museum of Natural History, Gainesville, who helped collect the types.

Comment: Charred tortoise bones found in conjunction with an early Taíno habitation site at Coralie that was occupied from ca. 700 to 1200 CE (Carlson 1999; Franz et al. 2001, 2020). No tortoise bones have been found in later habitation sites on the island (Carlson 1999). The Turks Islands are separated from the Caicos Islands by a deep strait and were never connected, even during the last glacial maximum at the end of the Pleistocene.


Chelonoidis alburyorum sementis † Franz, Albury, and Steadman 2020 (21:105)

Caicos Tortoise

(Extinct, ca. 1100 CE**)**



xiphiplastron, epiplastron / Middle Caicos, Turks and Caicos [Franz et al. 2020:f.14]



(subspecies: *alburyorum* = red; *keegani* = purple (Turks), *sementis* = blue)

Distribution: Turks and Caicos (*Middle Caicos Island*)

Estimated Possible Holocene Range: 710 sq. km

Size (Max SCL): male 60.0 cm [estimated] (Franz et al. 2020)

Synonymy:

Chelonoidis alburyorum sementis † Franz, Albury, and Steadman 2020:19

Type locality: "Indian (Head) Cave (site MC-37), 3 km west of the airport and Conch Bar Settlement, Middle Caicos Island, Turks and Caicos Islands, British Overseas Territory...21°49'N, 71°48'W."

Type specimen: UF 432441, holotype, subfossil, xiphiplastron, figured (f.14A,B); topotypic material genotyped by Kehlmaier et al. (2021a).

Geological age: Late Holocene, from ca. 10–770 CE until ca. 890–1020 CE (Franz et al. 2020; Steadman et al. 2020).


Etymology:* Latin, *sementis*, seeding, sowing; referring to the presumed ability of this tortoise to disperse seeds of native plants.

Comment: The Caicos tortoise bones were associated with human habitation, including Meillac style pottery dated to ca. 1200–500 ybp. Many of the bones were charred, indicating they had been consumed. Franz et al. (2001) hypothesized that the species had been extirpated by human predation in the late pre-Columbian historical period. The Caicos Islands are separated from the Turks Islands by a deep strait and were never connected, even during the last glacial maximum at the end of the Pleistocene.

Chelonoidis cubensis † (Leidy 1868b)
Cuban Giant Tortoise
(Extinct, ca. 9000 BCE?**)**




Carapacial costal and plastral fragments / Cuba [from Williams 1950b:pl.1-2]; *left*: Ned Gilmore, ANSP 8923, holotype; *right*: AMNH 6202



Distribution: Cuba
Estimated Possible Holocene Range: 58,349 sq. km
Size (Max SCL):* ca. 113.9 cm [estimated] (Joos et al. 2022)
Synonymy:
Testudo cubensis † Leidy 1868b:179
Testudo (Chelonoidis) cubensis, Geochelone cubensis, Hesperotestudo cubensis, Chelonoidis cubensis
Type locality: "Ciego-Montero, Cienfuegos, Cuba." Emended to "Chapepote spring at Baños de Ciego Montero, Santa Clara Province, Cuba" by Williams (1950b:9) and to "Chapapote (= Chapepote) spring, Baños de Ciego Montero, Cienfuegos (formerly Santa Clara) Province, Cuba" by Vlachos (2018:33).
Type specimen: ANSP 8923, holotype, subfossil, partial costal, figured in Williams (1950b:pl.1), see Gillette (1977) and Vlachos (2018).
Geological age: Pleistocene to Early Holocene.
Etymology:* Toponym for Cuba.
Comment: Redescribed by Williams (1950b). Auffenberg (1967) discussed this species; Meylan and Sterrer (2000) treated it as *Chelonoidis*. Karl (1995) described fosssils of smaller (CL ca. 40 cm) individuals of what might be this species from Late Pleistocene to Early Holocene deposits in San José de la Lamas, La Habana; this population apparently went extinct through human exploitation.

Chelonoidis dominicensis † Albury, Franz, Rimoli, Lehman, and Rosenberger 2018 [95]*
Northern Hispaniola Tortoise
(Extinct, ca. 8000 BCE?**)**





Nancy Albury / Dominican Republic [Albury et al. 2018:f.6,A,C,E]



**

Distribution: Dominican Republic, Haiti (?)
Estimated Possible Holocene Range:* 34,090 sq. km
Size (Max SCL): male ca. 60.0 cm [estimated] (Franz and Woods 1983), male 42.1 cm [measured] (Albury et al. 2018); **(Max CCL):** male 54.6 cm [measured] (Albury et al. 2018)
Synonymy:
Chelonoidis dominicensis † Albury, Franz, Rimoli, Lehman, and Rosenberger 2018:7
Type locality: "Oleg's Bat Cave (= Oleg's Bat House), 7 km west and inland of Bavaro...La Altagracia Province...southeastern coast...Dominican Republic...N18°42', W68°32'...elev. 22 m."
Type specimen: MHD 1000, holotype, subfossil, complete carapace and plastron, skull, cervical vertebrae, and appendicular bones, figured (f.2-8).
Geological age: Late Pleistocene to Early Holocene.
Etymology:* Toponym for the Dominican Republic.
Comment: This species was listed in TEWG (2015) as "*Chelonoidis* sp. [Hispaniola]" based on Franz and Woods (1983). ^{14}C dating of a specimen by Turvey et al. (2017) was unsuccessful. Similar deposits on Hispaniola have been dated from ca. 20,000 to less than 10,000 ybp (Franz and Woods 1983). Genetic sequencing of a specimen by Kehlmaier et al. (2021) was unsuccessful.

Chelonoidis gersoni † Viñola-López and Almonte 2022 [95]*
Gerson's Hispaniola Tortoise
(Extinct, ca. 8000 BCE?**)**






Plastron, holotype / Dominican Republic [Viñola-López and Almonte 2022:f.5,C,E,F]




**

Distribution:* Dominican Republic
Estimated Possible Holocene Range:* 2,408 sq. km
Size (Max SCL):* Plastron length 28.1 cm (Viñola-López and Almonte 2022); SCL ca. 32.5 cm, estimated here, based on comparison to *C. dominicensis* (Albury et al. 2018)
Synonymy:*
Chelonoidis gersoni † Viñola-López and Almonte 2022:20
Type locality: "Cueva del Mono, Pedernales Province, Dominican Republic."
Type specimen: MNHNSD FOS 23.001, holotype, subfossil, partial shell, figured (f.2–15).
Geological age: Late Pleistocene to Early Holocene.
Etymology:* Eponym honoring Gerson Feliz (1982–), Dominican Republic; field biology assistant in the Grupo Jaragua working on conservation of natural resources in Pedernales.
Comment: Viñola-López and Almonte (2022) considered *C. gersoni* distinct from *C. marcanoi*, which they considered conspecific with *C. dominicensis*.

Chelonoidis marcanoi † Turvey, Almonte, Hansford, Scofield, Brocca, and Chapman 2017 [95]*
Southern Hispaniola Tortoise
(Extinct, ca. 8000 BCE?**)**




humerus and carapace fragment / Dominican Republic [Turvey et al. 2017:f.2a,b,4i]




Distribution: Dominican Republic, Haiti (?)
Estimated Possible Holocene Range:* 7,307 sq. km
Size (Max SCL):* ca. 77.8 cm [estimated] (Joos et al. 2022)
Synonymy:
Chelonoidis marcanoi † Turvey, Almonte, Hansford, Scofield, Brocca, and Chapman 2017:4 (*nomen dubium*)
Type locality: "Cueva del Papayo, Pedernales Province, Dominican Republic."
Type specimen:* NHMUK PV R 36954, holotype, subfossil, humerus, figured (f.2a,b).
Geological age: Late Pleistocene to Early Holocene.
Etymology:* Eponym honoring Eugenio de Jesús Marcano Fondeur (1923–2003), Dominican Republic; botanist, zoologist, and cave explorer, who first discovered giant tortoise fossils on Hispaniola.
Comment:* This species was listed in TEWG (2015) as "*Chelonoidis* sp. [Hispaniola]" based on Franz and Woods (1983). ^{14}C dating of a specimen by Turvey et al. (2017) was unsuccessful. Similar deposits on Hispaniola have been dated from ca. 20,000 to less than 10,000 ybp (Franz and Woods 1983). Vlachos (2018) considered this taxon to be undefinable and a *nomen dubium*. Viñola-López and Almonte (2022) considered *C. marcanoi* distinct based on new material, and conspecific with *C. dominicensis*. However, Albury, Steadman, and Franz (2022, in litt.) continue to consider *C. marcanoi* a *nomen dubium* and recognize *C. dominicensis* and *C. gersoni* as distinct.

Chelonoidis monensis † (Williams 1952)
Mona Island Tortoise
(Extinct, ca. 1000 BCE?**)**



humerus and partial skull / Mona Island, Puerto Rico [Williams 1952:pl.46,44]




Distribution: Puerto Rico (*Mona Island*)
Estimated Possible Holocene Range: 66 sq. km
Size (Max SCL): ca. 50 cm [estimated] (Hansen et al. 2010)
Synonymy:
Testudo (Monachelys) monensis † Williams 1952:547
Testudo monensis, Geochelone (Monachelys) monensis, Geochelone monensis, Monachelys monensis, Chelonoidis monensis, Chelonoidis (Monachelys) monensis
Type locality: "Lirio Cave, Mona Island, West Indies" [Puerto Rico, USA].
Type specimen: AMNH 1969, holotype, subfossil, a first dorsal vertebra, figured (f.1e,pl.44,f.4).
Geological age: Late Pleistocene to Late Holocene, possibly until ca. 1050 BCE, 3000 ybp.
Etymology:* Toponym for Mona Island, Puerto Rico.
Comment: Mona Island is small and isolated, halfway between the larger islands of Hispaniola and Puerto Rico. Stone tools found on Mona date to ca. 3000 ybp, and an ancient cave drawing representing what appears to be a tortoise suggests the possibility that the species may have survived until early humans arrived (see http://blog.britishmuseum.org/2013/07/09/new-discoveries-of-cave-art-in-the-caribbean/; J. Cooper, unpubl. data). See also Vlachos (2018).

Solitudo † Valenti, Vlachos, Kehlmaier, Fritz, Georgalis, Luján, Miccichè, Sineo, and Delfino 2022 [96]*
Sicilian Giant Tortoises
Solitudo † Valenti, Vlachos, Kehlmaier, Fritz, Georgalis, Luján, Miccichè, Sineo, and Delfino 2022:1711
Type species:* *Testudo robusta* † Leith-Adams 1877, by original designation.
Etymology:* Latin, *solitudo*, solitude, loneliness; in allusion to the insular isolation of these tortoises, and *-tudo*, alluding to *Testudo* and the testudinid affinities of this taxon.

Solitudo sicula † Valenti, Vlachos, Kehlmaier, Fritz, Georgalis, Luján, Miccichè, Sineo, and Delfino 2022 [96]*
Sicilian Giant Tortoise
(Extinct, ca. 10,000 BCE**)**



Femur holotype (Valenti et al. 2022:f.4) / Bagheria, Sicily, Italy



**


Distribution:* Italy (*Sicily*)
Estimated Possible Holocene Range:* 2,424 sq. km
Size (Max SCL):* ca. 50–60 cm [estimated] (Valenti et al. 2022)
Synonymy:*
Solitudo sicula † Valenti, Vlachos, Kehlmaier, Fritz, Georgalis, Luján, Miccichè, Sineo, and Delfino 2022:1711
Type locality:* "Zubbio di Cozzo San Pietro Cave, Municipality of Bagheria, near Palermo, north-western Sicily, Italy."
Type specimen:* ZCSP US.0-Q2, holotype, an almost complete right femur, figured (f.4).
Geological age:* Late Pleistocene, AMS-dated age: 12,500 ± 500 ybp, (ca. 10,500 ± 500 BCE) (Valenti et al. 2022).
Etymology:* Toponym for Sicilian in Latin.
Comment:* Described from an archaeological funerary assemblage in a cave associated with human bones and potsherds, this is the most recently known surviving large-sized testudinid from the Mediterranean region. Although described from the latest Late Pleistocene, the species likely persisted locally into the earliest Holocene.

Testudinidae, Subfamily and Genus indet.

"?*Meiolania" damelipi* † White, Worthy, Hawkins, Bedford, and Spriggs 2010
Vanuatu Giant Tortoise
(Extinct, ca. 800 BCE**)**



humerus / Teouma, Efate, Vanuatu [Hawkins et al. 2016:S1,f.1,A1-4]




Distribution: Vanuatu (*Efate*, *Malakula*, *Santo*)
Estimated Possible Holocene Range: 10,459 sq. km
Size (Max SCL): ca. >100 cm [estimated] (White et al. 2010)
Synonymy:
?*Meiolania damelipi* † White, Worthy, Hawkins, Bedford, and Spriggs 2010:15512
Meiolania damelipi
Type locality: "Teouma Lapita site, Efate, Vanuatu."
Type specimen: AMF 136641, holotype, subfossil, humerus, figured (f.3A).
Geological age: Late Holocene; ^{14}C age: 2733–2741 ± 30 ybp, calibrated age: 2890–2760 ybp (940–810 BCE), see White et al. (2010).
Etymology:* Eponym honoring Willy Damelip (1900s–2007), Vanuatu; archaeologist who worked at the Vanuatu Cultural Center and helped collect the species.
Comment: Originally described by White et al. (2010) as possibly belonging to the extinct genus *Meiolania* (Meiolaniidae) with a question mark, Sterli (2015) questioned its identification as a Meiolaniidae, suggesting it might instead be a Testudinidae; this was subsequently corroborated by Rabi et al. (2019). However, the species has not yet been reassigned to a non-meiolaniid genus. The species was extirpated by modern humans of the Melanesian Lapita culture; archaeological midden specimens have consisted primarily of limb elements with direct evidence of butchering by humans (White et al. 2010; Hawkins et al. 2016).

Testudinidae sp. † [Fiji]
Fiji Giant Tortoise
(Extinct, ca. 1000 BCE?**)**



humerus / Naigani, Fiji [Hawkins et al. 2016:S1,f.1,B1–4]



Distribution: Fiji (*Viti Levu*, *Naigani*)
Estimated Possible Holocene Range: 11,156 sq. km
Size (Max SCL): undetermined
Synonymy:
?Meiolaniidae sp. † (Worthy, Anderson, and Molnar 1999:239)
Localities: "Voli Voli, Viti Levu, Fiji."
Specimens: NMNZ s/n, subfossil, ungual bone and humerus, figured in White et al. (2016:S1,f.1,B1-4).
Geological age: Late Holocene, ca. >2000–3000 ybp, see Worthy et al. (1999).
Comment: A subfossil ungual bone (age ca. > 2000–3000 ybp) from Viti Levu in Fiji from prior to human occupation was described by Worthy et al. (1999) as possibly being a tortoise and questionably compared to *Meiolania*. It was treated as an indeterminate taxon of Meiolaniidae by White et al. (2010), TEWG (2015), and Hawkins et al. (2016), but considered to be a Testudinidae by Rabi et al. (2019) and possibly the same taxon as ?*Meiolania damelipi*. Tortoise bones of what is presumably the same taxon were also documented from Naigani Island, Fiji, by Hawkins et al. (2016).

ANNOTATIONS

Comments on taxonomic change or other annotations in this checklist and previous checklists are indicated by superscripts. New annotations in each new checklist are bold dark blue underlined numbers in separate parentheses, with an asterisk, e.g.,(35)*. Earlier annotations from any of the previous checklists are indicated in subsequent checklists by two-part non-bold superscripts in separate parentheses that indicate the year of publication and the annotation number from that year, e.g.,(07:15, 17:50, 21:9). All annotations from previous checklists are listed at the end of this section, and all checklists are available as open-access publications online at **www.iucn-tftsg.org/checklist/**.

CURRENT 2025 CHECKLIST

1. New Distributional Data: This checklist includes 450 new or updated taxon maps with additional locality dots with or without revised Estimated Indigenous (Historical) Ranges. New maps with updated localities and revised ranges (422) are marked with a double asterisk (**), and those with only new or revised locality dots without revised ranges (28) are marked with a bullet (•). We have now also added color-coded stars to all taxon-specific maps to indicate type localities (original, restricted, inaccurately restricted, or estimated) for all recognized and synonymized taxa, except for those taxa with only general, very vague, or no type locality data.

We have utilized multiple sources for these updates, including expert input based on field data, localities from taxonomic literature cited below, new CBFTT species accounts, and IUCN Red List and TFTSG Provisional Red List assessments. We have also added several geographic distribution and range extension records from short notes or accounts in *Herpetological Review, Herpetology Notes, Check List, Journal of Threatened Taxa*, and other similar publications. In this 10th edition of the checklist, we have now also added a large number of selected locality records downloaded from the Global Biodiversity Information Facility (*www.gbif.org*) and/or iNaturalist (*www.iNaturalist.org*), with many of these records personally evaluated or viewed for verification, but have not cited these sources individually.

In addition, we have continued to add new or previously unrecorded locality records from several articles and books, both older and more recent (Williams 1960; Mittermeier et al. 1978; Shammakov 1981; Carrillo de Espinoza and Lamas 1985; Dundee and Rossman 1989; D'Amato and Morato 1991; Hossain and Sarker 1995; Palmer and Braswell 1995; Thirakhupt and van Dijk 1995; Bozhanski et al. 1996; Fritz and Buskirk 1997; Bringsøe and Buskirk 1998; Fritz and Ziegler 1999; Parham and Li 1999; Sutton and Christiansen 1999; Pawar and Choudhury 2000; Gonzales and Reichle 2002; Iftime 2002; Peregontsev 2002; Baha El Din et al. 2003; Rai 2004; Stuart and Platt 2004; Kitimasak et al. 2005; Khan 2006; Perälä 2006; Gibbs et al. 2007; Blanck and Kremser 2008; Schneider and Schneider 2008; Ahmed and Das 2009; Linthoi and Sharma 2009; Nguyen et al. 2009; Aryal et al. 2010; Chen and Lue 2010; Sousa et al. 2010; Ferronato et al. 2011; Gramentz 2011; Brito et al. 2012; Das et al. 2012; Platt et al. 2013; Todd et al. 2013; Bondarenko et al. 2014; Costa et al. 2015; Guyer et al. 2015; Rahman et al. 2015; Brakels et al. 2016; Maffei et al. 2016; Moraru et al. 2016; Struijk et al. 2016; Uhrin et al. 2016; Wagner et al. 2016; Bauer et al. 2017; Cassano 2017; Forth 2017, 2018; Madella-Auricchio et al. 2017; Martin et al. 2017; Akani et al. 2018; Engel et al. 2018; Power and Gilhen 2018; Brito et al. 2019b; Camper 2019; Martínez del Mármol et al. 2019; Duysebaeva et al. 2019; Souza et al. 2019; Tomović et al. 2019; Hmar et al. 2020, 2024; Kundu et al. 2020; Auer and Herz 2021; Davis et al. 2021; Enge et al. 2021; Pradhan et al. 2021; Ríos-Solís et al. 2021; Rosales-Martínez et al. 2021; Arteaga 2022; Dawson et al. 2022; Praschag et al. 2022; Reynolds 2022; Sánchez-Vialas et al. 2022; Srinivasulu and Kumar 2022; Turner et al. 2022; Agostini et al. 2023; Boulaouad et al 2023; Carr et al. 2023; Dorado-Rodrigues et al. 2023; Guedes et al. 2023; Mimila-Manzur et al. 2023; Tietz et al. 2023; Assis et al. 2024; Contreras-Calvario et al. 2024; Glorioso et al. 2024; Koné et al. 2024; Selman 2024; Van Devender et al. 2024; Hou and Shi 2025).

2. New or Revised Entry Field Data: In order for readers to easily identify taxon entry fields with new or revised data, we indicate such updated entries with a single asterisk (*) after the heading for the pertinent entry field(s): Distribution, Introduced, Estimated Indigenous Range, Size (Max SCL), CBFTT Account, IUCN Red List, TFTSG Provisional Red List, IUCN Green List, CITES, ESA, Type Locality, Type Specimen, Etymology, or Comment. New taxa added or resurrected in this checklist are indicated by an asterisk after the Synonymy heading.

3. Listed Names: We follow the opinions and recommendations of others in not including in the present checklist certain recent turtle names that have been considered unavailable or rejected or otherwise critiqued by the global taxonomic community (see Kaiser et al. 2013; Kaiser 2014; Rhodin et al. 2015; Jones 2017; TTWG 2017; Thomson et al. 2018, 2021; Krell 2021; Thiele et al. 2021; Wüster et al. 2021; Conix et al. 2021; Lien et al. 2021; Pyle et al. 2021; TTWG 2021; Denzer and Kaiser 2023, 2025; Shepherd 2023; Dubois 2025; Naish 2025). In addition, annotations and commentaries from previous checklists regarding these names have also been removed from the current checklist.

4. Testudines vs. Testudinata: In TTWG 2021 (Annotation 21:4) we discussed the question of which name and authorship should be used for the Order of all turtles and tortoises. For many years the prevailing name used was Testudines Linnaeus 1758, but Linnaeus used the name Testudines in a strictly vernacular Latin sense, so prevailing usage of the attribution of the name Testudines shifted to Batsch (1788), as also recently agreed by Joyce et al. (2004, 2020, 2021). We noted in our annotation that the name Testudinata Klein 1751 was not available since it was published pre-1758, while also arguing that Testudinata Klein *in* Behn 1760 was also not available, since it was simply a translation by Behn of Klein's 1751 work, and that according to ICZN (1999; Articles 1, 23) suprafamilial names are not subject to the Principle of Priority.

Recently, however, Dubois and Frétey (2024) have interpreted the suprageneric names attributed to Klein (1760) in the translation by Behn (1760) as nomenclaturally available and have suggested that the ordinal name for all turtles and tortoises should now be Testudinata Klein 1760. We disagree, and consider this name and attribution to be a *nomen oblitum*, having never been used by any authors until Dubois and Frétey (2024). McCranie (2018) recommended use of the name Testudinata, but attributed it to Behn (1760), not Klein. In any case, even if available, the Principle of Priority does not apply to suprafamilial names and therefore Testudinata Klein *in* Behn 1760 does not replace Testudines Batsch 1788, and we continue to use the latter designation.

5. Cheloidea: Orthography of this family-level name has recently been rendered as Chelidoidea by Gable et al. (2022), but since the stem of the name of the type genus *Chelus* is *Chel-*, the correct orthography of the name for this superfamily is Cheloidea (see ICZN 1999; Article 29), as initially proposed in TTWG (2021).

6. *Acanthochelys radiolata*: During preparation of an updated reassessment of the conservation status of *A. radiolata* for the

IUCN Red List, the estimated distribution of the species has been significantly revised to include an apparently disjunct population in the middle of the Rio São Francisco basin near the border between Bahia and Minas Gerais (Balestra et al. 2008). Further investigation and genomic analysis of specimens from this region is highly recommended.

7. *Mesoclemmys raniceps* and *M. jurutiensis*: Cunha et al. (2021) described *Mesoclemmys jurutiensis*, a new small putative species of toad-headed turtle from the Brazilian Amazon. The authors distinguished the taxon from its broadly sympatric congeners by differences in body size, head size, carapace shape, coloration, and mitochondrial DNA (cytochrome *c* oxidase, CO1). Based on that single locus, it is most closely related to *M. raniceps* and has thus far only been recorded from a very small area of western Pará south of the mainstem Rio Amazonas in the lower-most Rio Tapajós drainage basin. Based on the limited genetic and morphologic data and analysis presented, most of us consider it a subjective synonym of *M. raniceps*, as Iverson (2022a) did in his listing of turtle types, and as further analyzed and concluded by Gallego-García et al. (2023a). Clearly, further genetic analysis is needed, and determination of its possible range in its drainage basin and resolution of whether it is locally sympatric or possibly parapatric with respect to *M. raniceps*. It may be only a locally restricted variant or microendemic lineage of *M. raniceps*. We await additional data before making a more definitive determination and therefore synonymize it under *M. raniceps*.

In addition, Cunha et al. (2025) recorded ontogenetic phenotypic change in hatchlings of *M. raniceps* that initially had yellow head bands suggestive of the synonymized taxon *M. heliostemma* that gradually changed over a year to the black head without yellow bands typical of *M. raniceps*, providing further evidence of the synonymy of these two taxa, as also demonstrated by Cunha et al. (2019; see TTWG Annotation 21:15). We continue to treat *M. heliostemma* as a synonym of *M. raniceps*, with *M. wermuthi* and *M. nasuta* representing the other recognized members of this species complex.

8. *Mesoclemmys vanderhaegei* and *M. sabiniparaensis*: Cunha et al. (2022) described *Mesoclemmys sabiniparaensis*, a new putative toad-headed turtle from the northern Rio Araguaia basin of the Brazilian Amazon. They used shell and skull morphology and genetic analysis using mitochondrial DNA (cytochrome *c* oxidase, subunits I [COI] and 16S) to build a phylogenetic tree, which placed the new species as sister to *M. vanderhaegei* from the southern Rio Araguaia basin and further south. Based on the limited genetic and morphologic data presented, most of us consider it a subjective synonym of *M. vanderhaegei*, as analyzed and concluded by Gallego-García et al. (2023a). Clearly, further genetic and morphologic analysis is needed, and determination of the purported taxon's possible range in its drainage basin and resolution of whether it may be only a latitudinal variant of *M. vanderhaegei*. We await additional data before making a more definitive determination and therefore synonymize it for now under *M. vanderhaegei*.

9. *Phrynops geoffroanus*: Carvalho et al. (2022), in a continuation of the work previously conducted by Carvalho et al. (2016) and discussed in TTWG 2017 (Annotation 17:90), provided a detailed genetic and morphologic analysis of the *Phrynops geoffroanus* complex (as defined by Rhodin and Mittermeier 1983). They analyzed mitochondrial DNA and genomic ddRAD nuclear sequences from 71 specimens of *P. geoffroanus* collected from 22 rangewide localities in Brazil and Argentina, as well as morphometric data from 198 specimens; genetic comparisons were also made with specimens of *P. tuberosus* and *P. williamsi* of the *P. geoffroanus* complex, but not with *P. hilarii*.

The authors demonstrated the existence of at least four or potentially five geographically defined monophyletic lineages of *P. geoffroanus*. Two of these lineages are from the Amazon basin, one from the Rio Paraná basin, one from the Rio São Francisco basin and northeast coastal Brazil, and one from the Atlantic Forest and the Rio Doce region of southeast coastal Brazil, corroborating earlier morphological work on the *P. geoffroanus* complex (Rhodin and Mittermeier, unpubl. data).

Despite reaching the conclusion that these lineages represent separate cryptic species, the authors deferred from making nomenclatural recommendations, pending further analysis of the several available junior synonyms of *P. geoffroanus*. Nonetheless, in order to facilitate and clarify ongoing work on this species complex (Carvalho, Rhodin, and Mittermeier, unpubl. data), we provide restrictions of the type localities for *P. geoffroanus* and the synonymized taxon *Platemys waglerii* in the checklist text.

10. *Chelodina*: Kehlmaier et al. (2025), using the nine most variable and phylogenetically most informative nuclear loci (6071 bp) from the study by Thomson et al. (2021) and complete mitochondrial genomes (15,893 bp), examined genetic differentiation in all currently recognized *Chelodina* species. In line with previous investigations (Alacs 2008; Hodges 2015; Kehlmaier et al. 2019), there was strong evidence for multiple and ancient mitochondrial introgression events in several species (*Chelodina expansa*, *C. longicollis*, *C. walloyarrina* – for the latter species, see also Annotation 13), rendering the subgenera *Chelodina* sensu stricto and *Chelydera* in phylogenetic analyses non-monophyletic. However, in the analyses of the nuclear dataset, the recognition and monophyly of the subgenera was supported. The morphologically most distinct taxa (*C. expansa*, *C. longicollis*, *C. oblonga*, *C. parkeri*, *C. steindachneri*) were found well differentiated genetically, while the remaining species of the subgenera *Chelodina* sensu stricto and *Chelydera* were not or only weakly distinct. Except for synonymizing *C. walloyarrina* under *C. kuchlingi* (see Annotation 13), the authors refrained from reaching taxonomic conclusions or making taxonomic changes. Further research, preferably with genome-wide nuclear markers, is needed for disentangling this complex situation.

11. *Chelodina (Chelodina) mccordi*: Devi et al. (2021) analyzed mitochondrial and nuclear DNA of specimens of *C. mccordi* from Roti Island and Timor-Leste and compared them to samples of several *Chelodina* species from Australia. They corroborated the previous findings of Kehlmaier et al. (2019a) (TTWG Annotation 21:19) and also concluded that *C. mccordi mccordi* and *C. mccordi timorensis* were distinct at the subspecies level and that *C. mccordi roteensis* was a synonym of *C. mccordi mccordi*.

12. *Chelodina (Chelodina) novaeguineae* and *C. (C.) ipudinapi*: Joseph-Ouni and McCord (2022a) described *Chelodina (Chelodina) ipudinapi* as a new species from the Aramia River region of southwestern Papua New Guinea, characterizing it morphologically as a dwarf form most closely related to *C. novaeguineae*. However, no quantitative analyses of differences between these taxa were performed, and no analysis was done of the extent of variation in rangewide populations of *C. novaeguineae*. Furthermore, genetic analysis was not performed, and the lower Aramia River is not significantly isolated from the southern coastal lowland populations of *C. novaeguineae*. Although it is possible that *C. ipudinapi* may represent a diagnosable population, perhaps as a locally microendemic lineage in the Aramia watershed basin, we refrain from recognizing it as a named species or subspecies at this time, and regard it as a junior synonym of *C. novaeguineae*, pending genetic analysis and morphological comparison.

13. *Chelodina (Chelydera) kuchlingi*: Kehlmaier et al. (2025) examined genetic variation in northern Australian *Chelodina* using mitochondrial genomes and nine nuclear loci. Their dataset included sequences from historical museum material, including the holotype of *C. (C.) kuchlingi*, which had previously also been genotyped by Kehlmaier et al. (2019a). Resequencing its mitogenome using improved protocols optimized for formalin-preserved specimens provided evidence that the formerly reported apparent mitochondrial distinction of *C. (C.) kuchlingi* (Kehlmaier et al. 2019a; see TTWG Annotation 21:23) was based on a sequencing artifact. Instead, two historical specimens of *C. (C.) kuchlingi* are in fact genetically indistinguishable from turtles morphologically resembling *C. (C.) kurrichalpongo* that now occur on the Ord River floodplain of northern Western Australia. In addition, specimens of *C. (C.) walloyarrina* from the surrounding low coastal plain and from the western Kimberley sandstone plateau are not differentiated from *C. kuchlingi* on the nuclear genomic level. The authors concluded that *Chelodina walloyarrina* (McCord and Joseph-Ouni 2007) is a junior synonym of *Chelodina kuchlingi* Cann 1997 and that the extant *Chelodina* turtles from the Ord River floodplain are conspecific, as *Chelodina (Chelydera) kuchlingi*.

However, the genotyped samples of *C. walloyarrina* did not include the holotype of *C. walloyarrina* from the southwestern Kimberley plateau, nor were any samples of the closely related Arnhem plateau species, *C. burrungandjii,* genotyped and compared. Additionally, *C. kuchlingi* and sampled *C. walloyarrina* had markedly different mitochondrial DNA, interpreted by the authors as representing ancient mitochondrial introgression in *C. walloyarrina*, but possibly arguing against synonymization. Furthermore, a specimen of *C. kurrichalpongo* from the Finniss River, Northern Territory (near the type locality of *C. kurrichalpongo*) was indistinguishable from *C. kuchlingi* in its mitogenome and had only very slightly different nuclear DNA.

Despite the remaining uncertainties about the identities and phylogenetic relationships of *C. kuchlingi*, *C. kurrichalpongo* (including referred populations in the Western Australian coastal lowlands of the Ord River basin), *C. walloyarrina*, *C. burrungandjii*, and *C. rugosa* (sensu stricto), most of us tentatively support the synonymization of *C. walloyarrina* under *C. kuchlingi* at this time, as proposed by Kehlmaier et al. (2025), while noting that *C. kurrichalpongo* may either alternatively or in addition also need to be synonymized under that name. Clearly, more genetic analysis of these species and lineages with more samples from a wider geographic range is warranted and we encourage ongoing work to better unravel these complex relationships. Further work also needs to be done to improve the morphological description and diagnosis of *C. kuchlingi* as now redefined.

14. *Chelodina (Chelydera) kurrichalpongo* and *C. (C.) rugosa*: The synonymized *nomen nudum*, *Macrochelodina billabong*, previously listed under *Chelodina (Chelydera) rugosa*, has now been moved to the synonymy of *C. (C.) kurrichalpongo* in view of the location of its type locality in western Northern Territory, Australia. It could conceivably also be a synonym of the redefined *C. (C.) kuchlingi*.

15. *Chelodina (Chelydera) rugosa*: Kehlmaier et al. (2024b) analyzed the near-complete mitogenomic sequence of the holotype of *Chelodina intergularis*. They demonstrated that it is most similar to specimens of *Chelodina rugosa* from southern Papua New Guinea, thereby confirming its synonymization under that species. Unfortunately, no specimens of *C. rugosa* from Cape York, Queensland, were genotyped for comparison. However, they examined extensive historical records and concluded that the provenance of the holotype specimens of both *C. intergularis* and *C. rugosa* was most likely Somerset, northern Cape York, Queensland, Australia, further corroborating their synonymization.

16. *Elseya (Elseya) dentata*: Joseph-Ouni et al. (2022b,c,d,e), in a series of four successive papers, described external morphological features of three apparently allopatric populations of Australian *Elseya dentata* (sensu lato) from the Northern Territory and Western Australia. In their first paper they defined the distributional range of *E. dentata* (sensu stricto) as restricted to the Daly and Victoria river regions of western Northern Territory west of Arnhem Land. In their second paper they described a new species, *E. (E.) kalumburu*, from the Kimberley region of Western Australia, and in their third they described two new species, *E. (E.) sahul* and *E. (Pelocomastes) gondwana*, from the Roper River region of eastern Northern Territory east of Arnhem Land, noting that the latter was most closely related to their *E. (P.) oneiros* [= *E. (P.) lavarackorum*]. None of the descriptions, nor the fourth paper summarizing morphological traits of these populations, included quantitative analyses of differences among these taxa, no analyses were done of the extent of variation in rangewide populations, and no genetics were investigated.

Although it is possible that these three described taxa may represent diagnosable populations, we refrain from recognizing them as distinct species or subspecies at this time. We regard *E. kalumburu* and *E. sahul* as junior synonyms of *E. dentata*, and *E. gondwana* as a junior synonym of *E. lavarackorum*, all pending rangewide genetic analyses of the various populations in *E. dentata* (sensu lato) as well as within the subgenus *Pelocomastes*.

17. *Elseya (Hanwarachelys) novaeguineae*: Joseph-Ouni and McCord (2023a) described *Elseya (Hanwarachelys) caelatus berau* as a new subspecies from the Berau Gulf of the Vogelkop Peninsula, based on specimens collected in the 1990s, but since damaged and lost. They based their description solely on photographs of the lost specimens showing alleged phenotypic differences of shell shape in adults. No quantitative analyses of differences between this alleged taxon and *E. novaeguineae* or other "*E. caelatus*" were performed, nor any analyses of the extent of variation in rangewide populations of *Elseya* from the Vogelkop Peninsula. The Berau Gulf population is not geographically separated from the overall range of *E. novaeguineae*, and although it is possible that it may represent a diagnosable population, we refrain from recognizing it as a named taxon at this time. We regard it and *E. caelatus* itself as junior synonyms of *E. novaeguineae*, pending rangewide genetic analysis.

18. *Elseya (Hanwarachelys) novaeguineae* and *E. (H.) schultzei*: Thomson et al. (2023), in their analysis of the conspecificity of *E. lavarackorum* and *E. oneiros*, listed *Elseya caelatus* and *Elseya orestiad* as apparently valid species. However, their phylogeny based on skeletal and other morphological characters did not differentiate *E. caelatus* from *E. novaeguineae*, *E. schultzei*, and *E. rhodini*, and *E. orestiad* was found to differ only slightly from *E. rhodini*, but not from the other two taxa. Based on our previous evaluations (TTWG 2021; Annotations 21:27 and 21:28), we continue to treat *E. caelatus* and *E. orestiad* as subjective junior synonyms of *E. novaeguineae* and *E. schultzei*, respectively, pending defensible morphological analyses and further genetic analysis using a combination of nuclear and mitochondrial data.

19. *Elseya (Hanwarachelys) rhodini*: Joseph-Ouni and McCord (2022c) described *Elseya (Hanwarachelys) papua* as a new species from the Merauke River region of southeastern coastal Papua Province in Indonesia. They characterized it as morphologically distinct from the much larger *E. branderhorsti* in various phenotypic features regarding maxillary color and head shield, and distinguished it from *E. rhodini* by a non-flecked iris and less prominent carapacial spots in females. No quantitative analyses of differences between

these taxa were performed, and no analysis was done of the extent of variation in rangewide populations of any of these species. In addition, genetic analysis was not performed. The Merauke River population is not geographically separated from the overall range of either *E. rhodin* or *E. branderhorsti*, and although it is possible that *E. papua* may represent a diagnosable population, we refrain from recognizing it as a named taxon at this time. We regard it as a junior synonym of *E. rhodini*, pending phylogeographic analysis of the entire range of what is currently considered *E. rhodini*.

In addition, Joseph-Ouni and McCord (2023b) described five other new purported taxa from within the overall distribution of *E. rhodini*: *Elseya (Hanwarachelys) timika timika*, *E. (H.) timika aru*, *E. (H.) timika wamena*, *E. (H.) timika paniai*, and *E. (H.) freweri*. Once again, no quantitative analyses of differences between any of these taxa and *E. rhodini* were performed, and no analysis was done of the extent of variation in rangewide populations of any of these taxa. In addition, genetic analysis was not performed, nor any measurements provided for any specimens. Although it is possible that the population on the Aru Islands (*E. timika aru*) might be minimally distinct, and the two possibly isolated populations in the central highlands, *E. timika wamena* and *E. timika paniai*, may also represent diagnosable populations (or quite possibly translocated trade specimens), we refrain from recognizing them or the other two lowland taxa, *E. timika timika* and *E. freweri*, as named species or subspecies at this time. We regard all five purported taxa as junior synonyms of *E. rhodini*, pending rangewide genetic analysis of that species.

20. ***Elseya (Hanwarachelys) schultzei*:** Joseph-Ouni and McCord (2022b) described *Elseya (Hanwarachelys) nabire* as a new species from the Wanggar River region of northern coastal Papua Province in Indonesia, characterizing it morphologically as distinct from both *E. novaeguineae* and *E. schultzei* in that its brown carapace lacked darker spots or radiating lines generally found in the other two taxa and that it was smaller. However, no measurements were provided for any specimens, no quantitative analyses of differences between these taxa were performed, and no analysis was done of the extent of variation in rangewide populations of any of these species. In addition, genetic analysis was not performed. The Wanggar River population appears to be geographically contiguous with the westernmost apparent extent of *E. schultzei*. It is possible that *E. nabire* may represent a diagnosable population, perhaps as a locally microendemic lineage in the Wanggar River watershed basin. However, we refrain from recognizing it as a named species or subspecies at this time, and we regard it as a junior synonym of *E. schultzei*, pending phylogeographic analysis of this wide-ranging species.

21. ***Elseya (Hanwarachelys) schultzei* and *E. (Solomonemys) eidolon* and *E. (S.) auramemoria*:** Joseph-Ouni et al. (2022a) described *Elseya (Solomonemys)* as a new subgenus most closely related to *Elseya (Hanwarachelys)* and allegedly endemic and restricted to the oceanic Solomon Islands east of Papua New Guinea. They characterized *Solomonemys* morphologically as distinct from *Hanwarachelys* in several phenotypic characteristics, but most prominently that the neck length in *Solomonemys* is allegedly 1.5 to 2 times longer than the skull length. However, they did not provide any measurements of neck length or the neck to skull length ratio in either subgenus. As such, we are not convinced that *Solomonemys* is a distinct subgenus and instead synonymize it under *Hanwarachelys*.

They also described two new proposed species in this subgenus: *E. (S.) eidolon* from Malaita and *E. (S.) auramemoria* from Guadalcanal (Solomon Islands). Their descriptions of these two allegedly new species were based on single specimens of each. A third chelid turtle specimen located in the Australian Museum and allegedly collected on Malaita in 1965 (see below) was identified by the authors as a translocated *Wollumbinia [= Myuchelys] latisternum*, yet they concluded that their other two specimens likely represented a nearly extinct native endemic chelid fauna in the Solomon Islands. No quantitative analyses of differences between these alleged taxa and *E. schultzei* or other *Elseya* species were performed, and no analysis was done of the extent of variation in rangewide populations. In addition, genetic analysis was not performed.

Given that no confirmed records of any native freshwater turtles have ever been recorded from any of the islands east of Papua New Guinea, including New Britain, New Ireland, Bougainville, or any of the Solomon Islands, these isolated specimens from Malaita and Guadalcanal appear most likely to represent translocated trade animals, as previously suggested by Rhodin and Genorupa (2000). It is unlikely that either of these proposed species represent native diagnosable taxa from the Solomon Islands, and we regard them both as junior synonyms of *E. schultzei*, introduced via native trade, most likely from northeastern mainland Papua New Guinea.

The specimen of *Myuchelys latisternum* allegedly collected on Malaita in 1965 is interesting in that there is significant disagreement whether it was in fact collected there or not. Joseph-Ouni et al. (2022a) noted that it was collected by William Dawbin, but they suggested the possibility that the specimen may have subsequently been mislabeled, while they also noted that Cann asserted that Dawdin, who died in 1998, had actually collected the specimen there. However, Dawbin himself, in a pers. comm. to John Legler (1990, W. Dawbin, unpubl. data), stated that he did not remember obtaining and preserving the specimen on Malaita and that it had been found unlabeled on top of a drum at the Australian Museum containing multiple specimens he collected and preserved on Malaita. It was his opinion that someone had placed the turtle in the drum with his Malaita collection and that the specimen had thus become erroneously labeled and had not been obtained on Malaita.

22. ***Elseya (Pelocomastes) irwini*:** Joseph-Ouni et al. (2023a) provided a morphological description of *Elseya* specimens from the Johnstone River in Queensland and assigned them to the previously unavailable name *stirlingi* (Wells 2007b) (unavailable as per ICZN criteria; see Fritz and Havaš 2007, Georges and Thomson 2010, and TTWG 2021, Annotation 21:17), differentiating them from *Elseya irwini* in the Burdekin River and redescribing them as a subspecies of that species. They attributed the name *stirlingi* specifically to Wells and copied his original descriptive language (Wells 2007b), thereby making the name nomenclaturally available as *Elseya (Pelocomastes) irwini stirlingi* Wells *in* Joseph-Ouni et al. 2023a.

The Johnstone River taxon appears morphologically somewhat different from the population in the Burdekin River, and previous genetic work by Todd et al. (2014b) demonstrated statistically significant but shallow mitochondrial DNA differentiation between Johnstone and Burdekin specimens of *E. irwini*. Whether the Johnstone population is subspecifically distinct from that in the Burdekin, or only a separate evolutionarily significant lineage (ESU) remains unclear, and we prefer to retain both populations under one species, *Elseya irwini*, without any subspecific distinction, pending further genomic analysis.

We note also that both Wells (2007b) and Joseph-Ouni et al. (2023a) recorded inaccurate geographic coordinates for the stated type locality of *stirlingi*, apparently based on an unrecognized typographic error. The given coordinates by both sets of authors (17 38'S, 145 05'E) actually places the supposed type locality ca. 80

km west of the Johnstone River drainage basin, in the highlands at Mount Garnet, near the source of the Herbert River in the adjacent drainage basin. Correcting the longitude to 145 50'E (by simply reversing 05' to 50') would place the type locality squarely on the South Johnstone River as Wells (2007b) originally noted, and we have therefore now emended the type locality as such. Unfortunately, Joseph-Ouni et al. (2023a, Fig. 1) additionally mapped the type locality for *stirlingi* as supposedly being near Malanda on the North Johnstone River (while also giving the inaccurate coordinates); this is clearly inaccurate as well.

23. *Elseya (Pelocomastes) lavarackorum*: Joseph-Ouni et al. (2022f), in a more detailed follow-up to their previous description of *E. oneiros* (Joseph-Ouni et al. 2020), re-examined the available late Pleistocene fossil specimens previously designated as *E. lavarackorum* and compared them to extant specimens of *Elseya* "*oneiros*" from the same drainage basin (the Nicholson and Gregory rivers of northern Queensland) and other northern Australian *Elseya* lineages. Their new work focused on characterizing several skeletal characters in the two species, which the earlier paper did not. They concluded that the two species were different and that *E. lavarackorum* was an extinct late Pleistocene taxon and that *E. oneiros* represented the extant populations in this area. The authors additionally concluded that *E. oneiros* appeared to have also occurred in the late Pleistocene, based on a single partial plastron fossil with some characteristics of *E. oneiros*. The authors also suggested, based on the single character of a purported medial constriction of its axillary buttress suture, that *E. lavarackorum* should be reallocated to the subgenus *Elseya* rather than the subgenus *Pelocomastes.*

Further detailed examination and analysis of all relevant samples of these and other related taxa by Thomson et al. (2023) demonstrated that the purported differences noted between *E. lavarackorum* and *E. oneiros* by Joseph-Ouni et al. (2022f) were not diagnostic. Thomson et al. (2023) noted that the axillary buttress suture of the fossil specimen of *E. lavarackorum* was associated with extensive and prominent embedded plastral elements, which had led Joseph-Ouni et al. (2022f) to misinterpret its supposed medial constriction (i.e., there was no constriction). Additionally, they found that marked inter-individual variation in scute patterns rendered any of the differences noted by Joseph-Ouni et al. (2020, 2022f) to be of little taxonomic value. In summary, Thomson et al. (2023) demonstrated that the late Pleistocene fossil *E. lavarackorum* (from ca. 23,900 years ago), and the extant and late Pleistocene *E. oneiros* are not diagnosable as distinct taxa, and that both are referable to the subgenus *Pelocomastes*.

In a subsequent lengthy response to Thomson et al. (2023), Joseph-Ouni et al. (2023b) revisited and re-analyzed specimens of the two purported taxa (*E. lavarackorum*, based on the single Pleistocene fossil holotype, and many living *E. "oneiros"*) and again reached the conclusion that they were distinct species. However, despite examining many specimens of "*E. oneiros*", they provided no quantitative statistical analyses of their findings and did not address the extent of variation within the living taxon and the extent to which it does or does not encompass the single fossil *E. lavarackorum*. In addition, genetic analysis was not performed, nor were any measurements provided for any specimens. Further, they also claimed to have discovered evidence of the persistent survival of the Pleistocene form *E. lavarackorum* (which they had previously considered to be extinct) based on photographs of a single living turtle obtained in 1982. Clearly, further analysis will be needed to resolve these thorny taxonomic questions and we encourage Joseph-Ouni et al. to add genetic analysis to clarify and hopefully resolve some of the questions their work has produced, particularly their conclusion regarding the supposed co-occurrence of two living lineages in the region.

Joseph-Ouni et al. (2023b) also addressed the question of the subgeneric allocation of the *E. lavarackorum* fossil holotype. Originally placed in the subgenus *Pelocomastes* by Thomson et al. (2015), Joseph-Ouni et al. (2020) reallocated it to the subgenus *Elseya*. However, Thomson et al. (2023) confirmed through re-examination of the fossil holotype that it belongs in the subgenus *Pelocomastes*, yet Joseph-Ouni et al. (2023b) again challenged this interpretation. Further, they also declared the genus-level name *Pelocomastes* to be a *nomen dubium* and called for the creation of a new subgeneric name.

The genus-level name *Pelocomastes* was first made available as *Pelocomastes ampla* † by De Vis (1897), a taxon that Thomson (2000b) regarded as a subjective junior synonym of *Chelymys uberrima* † De Vis (1897) [= *Elseya (Pelocomastes) uberrima*], which Thomson et al. (2015) then designated as the type species for *Pelocomastes*. However, Joseph-Ouni et al. (2023b:plate 31) compared lectotype specimens of *P. ampla* and *C. uberrima* and concluded that they belong to different taxa, based on slightly different scute sulcus patterns on the first peripheral bone. Although *C. uberrima* is considered by both Thomson and Joseph-Ouni et al. to be a diagnosable extinct taxon, the question remains as to whether *P. ampla* is also diagnosable as a separate species or simply a slightly different variation of *C. uberrima*.

According to ICZN Article 12.2.5, the availability of a genus-level name created prior to 1931 depends on whether it was created with "one or more available specific names in combination with it, [...] provided that the specific name [...] can be unambiguously assigned to a nominal species-group taxon." Therefore, if *P. ampla* and *C. uberrima* are indeed different taxa, and if *P. ampla* is undiagnosable at the species level, then the type species of *Pelocomastes* reverts to *P. ampla*, possibly a subjective *nomen dubium*, and *Pelocomastes* would then not be considered an available genus-level name under ICZN rules. We leave it to the two author teams to work out the best course of action, but creating a new name for the subgenus *Pelocomastes* based on a living species in that subgenus may be the best approach, as recommended by Joseph-Ouni et al (2023b).

However this genus-level nomenclatural quandry becomes resolved, and despite this last attempt by Joseph-Ouni et al. to justify the recognition of *E. oneiros* as distinct from *E. lavarackorum*, we reaffirm our prior evaluation (TTWG 2021; Annotation 21:30) and continue to refer to the extant *Elseya* species in the Nicholson–Gregory drainage basin and westward into the Roper River as *Elseya lavarackorum* rather than either *E. oneiros* and/or *E. gondwana*, as recommended by Joseph-Ouni et al. (2022d,f) (see Annotation 16 above). If a single taxon is present, our evaluation of the available evidence is that it is referable to *E. lavarackorum*; if two genetic lineages are present, then further taxonomic work is needed. We urge these authors or others to carry out genetic analyses to clarify the phylogeography of these confusing taxa and to follow the guidelines for best practice for taxonomic publication requirements as recommended by Kaiser et al. (2013).

24. *Emydura australis*: Kehlmaier et al. (2024a) analyzed the genetic differentiation of red-faced vs. yellow-faced individuals referred to as *Emydura victoriae* or *E. subglobosa worrelli* or *E. tanybaraga* from several rivers in northern Australia. They sequenced the mitochondrial genomes (mitogenomes) and 10 nuclear loci of 20 samples from the Carson, Mitchell, and Ord River drainages in the Kimberley region of northern Western Australia and compared them to sequences from the Victoria, Daly, and other river drainages in the Northern Territory, including mitogenomes

of the type specimens of *E. victoriae* and *E. australis*. The results provided evidence for the occurrence of a variably colored (red- or yellow-faced) single species in the Mitchell River, but with two deeply divergent mitochondrial lineages, one of which may have originated from ancient hybridization and mitochondrial introgression from the *E. subglobosa worrelli* / *E. tanybaraga* complex.

Turtles from the Ord River and a single individual from the Carson River were red-faced. Turtles from the Ord River represented a maximally supported but weakly divergent mitochondrial clade and turtles from the Carson and Mitchell rivers represented another maximally supported and weakly divergent clade. In this latter clade, the single sample from the Carson River was weakly divergent. The name-bearing types of *E. australis* and *E. victoriae* were the successive weakly divergent sister taxa of the Ord River turtles. The next successive sister taxon was a sample of *E. macquarii*; other samples of the latter species were paraphyletic to those mentioned above. Notably, another sample of *E. victoriae* from the Daly River clustered in a deeply divergent mitochondrial clade together with *E. subglobosa worrelli*, *E. tanybaraga*, and the introgressed Mitchell River turtles. These results suggested that *Emydura* populations in this region represent a speciation continuum and that the evolutionary history within the genus is characterized by multiple hybridization and introgression events.

Based on comparison with previously published mitogenomes of type material, Kehlmaier et al. (2024a) concluded that *E. victoriae* (Gray 1842) is conspecific with *E. australis* (Gray 1841) and the *Emydura* turtles from the Ord, Carson, and Mitchell rivers, and recommended that the older name (*E. australis*) be given nomenclatural priority, despite the long history of uncertainty as to the identification and provenance of its type specimen. Based on clarification of the type locality of *E. australis* (see below), we agree with this recommendation, and therefore accept the proposed resurrection of *Emydura australis* (Gray 1841) from its prior erroneous synonymy under *E. macquarii* (Cogger et al. 1983) and also place *Emydura victoriae* (Gray 1842) in synonymy under *E. australis*.

The genetic analysis also demonstrated that the type specimen of *E. australis* nested with and is weakly divergent from specimens from the Ord River drainage on the Kimberley plateau, whereas the type specimen of *E. victoriae* from the Victoria River is weakly divergent from that clade, with both bootstrap values and posterior probabilities showing maximum but shallow support for this phylogeny. As a result, this suggests that the type specimen of *E. australis* is not from the Victoria River, but instead from the Kimberley region. However, while there is evidence of some genetic divergence among the populations of *E. australis* across its range, there is no evidence that these represent different taxa at the species or subspecies level or that *E. australis* and *E. victoriae* are separate taxa.

When originally described, the provenance of the type specimen of *E. australis* was designated as from "Western Australia?" by Gray (1841). However, Gray (1872d) later erroneously changed it to "Australia, Macquarie River" and then further changed it to "N. Australia, Macquarie River" (Gray 1873j), synonymizing *E. australis* under his concept of *Chelymys macquaria* (= *Emydura macquarii*), where it has since been variably synonymized or treated as a *nomen dubium* by many (e.g., Cogger et al. 1983; Georges and Thomson 2010; TTWG 2017, 2021), but recognized by others (e.g., Goode 1967; Cann 1998).

More recently, Cann and Sadlier (2017), who also recognized *E. australis*, documented that the type specimen of *E. australis* was most likely collected in 1838 or 1839 by George Grey in the vicinity of the Prince Regent River in Western Australia. His travelogue (Grey 1841) indicates that he traversed the regions south of Hanover Bay from near the mouth of the Prince Regent River, discovered and named Mount Lyell and the Gairdner and Glenelg rivers, and mentioned that he found emus along the Glenelg River and that he also found an inland freshwater turtle. Furthermore, Gray's (1841) description of the single type specimen of *E. australis* was published as an Appendix in Grey's travelogue. Based on the above findings and clarifications, we hereby restrict the type locality of *Hydraspis australis* Gray 1841 (= *Emydura australis*) to the "vicinity of the Glenelg and Prince Regent rivers, Western Australia, Australia" in the western Kimberley region.

25. *Pelomedusa* and other Sub-Saharan genera: Fritz et al. (2022) produced a nicely informative and integrative review of Sub-Saharan African turtle taxonomy and phylogenetics in a major article honoring the taxonomic legacy of Margaretha Hofmeyr. The article reviewed detailed aspects of the emerging taxonomy of *Pelomedusa*, *Pelusios*, *Centrochelys*, *Chersina*, *Chersobius*, *Homopus*, *Kinixys*, *Psammobates*, *Stigmochelys*, *Cyclanorbis*, and *Trionyx*. It also addressed some of the difficulties for researchers to share genetic samples across international borders under the regulations of the Nagoya Protocol of the Convention on Biological Diversity.

26. *Pelusios castaneus*: Guedes et al. (2023) analyzed the mitochondrial DNA phylogeography of *P. castaneus*, including several specimens collected on São Tomé and Principe islands. They demonstrated that the genetic affinities of these island specimens grouped with separate and distinct genetic lineages from Ivory Coast, Sierra Leone, Nigeria, and coastal Republic of Congo, indicating that they most likely represent historically introduced animals, although some may have arrived via rafting from continental rivers. Their work further strengthens the previous conclusions regarding the origin of *P. seychellensis* (see TTWG Annotation 21:35), in that one of their specimens from São Tomé grouped with the type specimen of *P. seychellensis* and several specimens from coastal Republic of Congo.

27. Podocnemidinae: Authorship of this family-level name was given as Gray 1870f in TTWG (2021). However, in accordance with the ICZN (1999; Article 36, Principle of Coordination), this name should instead be attributed to Cope 1869.

28. *Podocnemis vogli*: Cárdenas-Barrantes et al. (2024) sequenced 19 microsatellite loci in five populations of *P. vogli* along its distribution range in Colombia. They found significant genetic structure among them and recommended that the five populations corresponded to five independent management units, but recommended no taxonomic changes.

29. Chelonioidea and Chelydroidea (Americhelydia): Gable et al. (2022) analyzed thousands of high-confidence single-copy orthologs from 19 genome assemblies representative of extant turtle diversity and estimated a phylogeny with multispecies coalescent and concatenated partitioned methods. They also collected next-generation sequences from 26 turtle species and assembled millions of biallelic markers to reconstruct phylogenies. They found strong and consistent support for all nodes in currently inferred turtle phylogenies, notably for the PhyloCode clade (or hyperfamily) Americhelydia (Joyce et al. 2013, 2021) uniting the superfamilies Chelonioidea and Chelydroidea (encompassing the families Cheloniidae, Dermochelyidae, Chelydridae, Dermatemydidae, and Kinosternidae). We agree that these families are most closely related, as previously shown also by Barley et al. (2010) and Thomson et al. (2021). However, we are not incorporating hyperfamily designations (e.g., Americhelydia) in our checklist at this time.

30. Cheloniidae and Dermochelyidae: Phillips et al. (2022) analyzed the global phylogeography of all seven sea turtle species

based on mitochondrial DNA control region haplotypes from 709 samples, including 35 newly sequenced juvenile *Chelonia mydas*, *Eretmochelys imbricata*, and *Caretta caretta* from the Gulf of Mexico. The generated gene trees differ somewhat from earlier published versions by others.

31. *Lepidochelys kempii* and *L. olivacea*: Vilaça et al. (2022) analyzed the global phylogeography of *Lepidochelys* species, based on mitochondrial DNA sequences from 943 *L. olivacea* and 287 *L. kempii*, and 15 nuclear microsatellite loci from 285 *L. olivacea*. They found that the two species diverged ca. 7.5 million years ago, before the Panama Isthmus closure. The most ancient mitochondrial *L. olivacea* lineage, located in the Indian Ocean, was dated to ca. 2.2 mya. They identified three divergent and well-supported clades based on mitochondrial and nuclear DNA: 1) *L. kempii*; 2) Indian Ocean *L. olivacea*; and 3) all other *L. olivacea*. This last group also contained one Atlantic, one East Pacific, and two Indian-West Pacific subclades. No taxonomic recommendations were made.

32. *Chelonia mydas*: Álvarez-Varas et al. (2021) analyzed genetic structure in the two *C. mydas* shape-based color morphotypes in the Pacific Ocean (yellow and black) using a reduced-representation genome approach (9473 SNPs) and identifying candidate outlier loci (67 SNPs) of biological relevance in 158 specimens from eight Pacific foraging grounds. Their results supported genetic divergence between the morphotypes and evolutionary distinction of the enigmatic black *C. mydas* of the Eastern Pacific, probably arising from strong natal homing behavior. Although these populations of black turtles had previously been recognized as the separate species *C. agassizii*, they did not recommend any taxonomic changes, but suggested that the black *C. mydas* should be managed as an independent evolutionary unit, and that specific conservation strategies for each shape-based morphotype should be established.

33. *Testudo chloronotos*: Zug (2022) was unable to locate the name *Testudo chloronotus* Bechstein 1800 (as previously synonymized under *Chelonia mydas*) and questioned its existence despite its having been recorded on multiple occasions over the years. In fact, Bechstein named the species *Testudo chloronotos*, not *chloronotus*, and our previous checklist had recorded the erroneous spelling, as did Brongersma (1961), who also discussed the species and restricted its type locality, which we had also overlooked. These errors are now corrected.

34. Dermochelyidae: Authorship of this family-level name was given as Fitzinger 1843 in our earlier checklists, including TTWG (2021). However, based on analysis by Frétey (2022), and in accordance with the ICZN (1999), this name should instead be attributed to Baur 1888a, as Fitzinger based his family name Dermatochelydae on the rejected genus name *Dermatochelys*, rendering his family designation nomenclaturally unavailable. Baur (1888a) designated the family as Dermochelydidae, which according to Frétey (2022) must be corrected to Dermochelyidae based on ICZN guidelines. We hereby follow Frétey's recommendation and now allocate authorship of the family-name Dermochelyidae to Baur (1888a).

35. *Dermochelys coriacea*: Castillo-Morales et al. (2023) analyzed mtDNA (D-loop) genetics of Leatherback Turtles nesting in Oaxaca, Mexico. They demonstrated five different haplotypes separated into two apparently divergent lineages, which also seemed to correspond to two different morphotypes of animals as described through Local Ecological Knowledge (LEK), with the divergent lineage much rarer than the more common one. They suggested that it might be possible that the divergent lineage represented a different species or subspecies, and that one of the previously synonymized names, *Sphargis coriacea schlegelii* Garman 1884 or *Sphargis angusta* Philippi 1899, might need to be resurrected for that lineage.

However, Colombo et al. (2024) re-analyzed the same set of samples with improved genomic techniques, and demonstrated that the apparently divergent lineage was an artifact caused by the misidentification of a nuclear mitochondrial insertion (NUMT) as mitochondrial DNA. They also noted that NUMTs had not previously been demonstrated as occurring in sea turtles. They further suggested that the supposed morphological differences between the lineages were within the parameters of phenotypic variation for *D. coriacea* and rejected the prior finding by Castillo-Morales et al. (2023) that these were divergent lineages.

In addition to these studies, we (the TTWG) have reassessed the maximum recorded sizes of *D. coriacea*, as there has been some confusion and disagreement on these alleged values, both for straight-line carapace length (SCL) and curved carapace length (CCL) measurements and estimates. The previously cited values in our previous checklist have turned out to be incorrect and we have now updated them.

36. *Macrochelys temminckii*: Apodaca et al. (2023) investigated the phylogeography of *M. temminckii* based on an analysis of SNPs from 104 Alligator Snapping Turtles from 27 river drainages from Texas to Florida, including samples from the Apalachicola River drainage basin, representing the previously synonymized taxon *M. apalachicolae*. They found that specimens from the Apalachicola River watershed were genetically distinct from western populations of *M. temminckii* and eastern populations of *M. suwanniensis*, with little gene flow among them. They suggested that *M. apalachicolae* represented a distinct lineage that could be considered as a separate species, but held off on resurrecting it, pending morphological studies.

37. *Kinosternon*: Hurtado-Gómez et al. (2024) analyzed phylogeographic structure and taxonomic limits within the genus *Kinosternon*, based on molecular (mitochondrial and nuclear DNA), geographic, and taxonomic sampling of 116 specimens, focusing on relationships within the polytypic species *K. leucostomum* and the *K. scorpioides* group, and presented a fossil-calibrated nuclear time tree for the genus.

Their results revealed contrasting differentiation patterns for the *K. scorpioides* group and *K. leucostomum*, despite similar wide distributions. *Kinosternon leucostomum* showed only shallow genetic divergence across its range, and its two currently recognized subspecies, *K. l. leucostomum* and *K. l. postinguinale*, were recommended for synonymization into a monotypic species. However, unpublished morphological data comparing the two subspecies previously demonstrated enough differentiation to warrant potential continued recognition of these two taxa (Berry 1978), so we await further data and publications before considering synonymizing them.

The *K. scorpioides* group, on the other hand, was found to be deeply polyphyletic, with up to 10 distinct taxa, with some of them still undescribed. The work by Hurtado-Gómez and colleagues strongly supported the elevation of *K. s. albogulare* and *K. s. cruentatum* to species, which they recommended and with which we agree. They also noted that the Atlantic and Pacific versant populations of *K. cruentatum* were divergent enough to warrant species recognition, but did not make any specific nomenclatural recommendations.

That study also found that all *K. scorpioides* from Panama southward across South America formed a monophyletic group, but with substantial genetic diversity within the clade. However, because geographic sampling across most of the distribution was meager, and the types of the many synonymized South American

taxa were not genotyped, Hurtado et al. declined to suggest any taxonomic changes pending further study.

Furthermore, in view of the deep divergence demonstrated within the genus *Kinosternon*, the authors also proposed the recognition of three subgenera: *Kinosternon*, *Cryptochelys*, and *Thyrosternum*, with which we also agree. The subgenus *Cryptochelys* was found to include *K. angustipons*, *K. dunni*, and *K. leucostomum*. The subgenus *Thyrosternum* was demonstrated to encompass *K. baurii, K. durangoense, K. flavescens, K. steindachneri, K. stejnegeri*, and *K. subrubrum*. The subgenus *Kinosternon* included all the other species except for *K. alamosae*, which was not formally assigned to a subgenus; however, in view of its morphological features it most likely belongs to this subgenus, which is where we provisionally assign it, pending further genetic data.

38. *Kinosternon cruentatum* and *K. mexicanum*: Iverson and Berry (2024), in a continuation and expansion of the work by Hurtado-Gómez et al. (2024) that had demonstrated that the Pacific and Atlantic versant populations of "*K. cruentatum*" were specifically distinct, examined the type material of the various taxa synonymized under *K. cruentatum*. They determined morphologically that the holotype of *K. mexicanum* represented the Pacific versant population, and that the holotype of *K. cruentatum* was an Atlantic versant specimen, thereby applying these names to those two allopatric populations. Unfortunately, the type locality for *K. cruentatum* (originally simply "Amér. septentr.") had previously been uncritically restricted as coastal Pacific Oaxaca, necessitating a revised restricted type locality to coastal Atlantic Quintana Roo. We agree with these taxonomic and nomenclatural clarifications.

39. *Kinosternon hirtipes*: Weaver et al. (2022) investigated the genomic phylogeography of the five extant subspecies of *Kinosternon hirtipes*. They analyzed SNPs as well as species distribution models to assess levels of ecological and genetic differentiation among these subspecies based on data from 141 specimens from across the range of the species. They found deep genetic divergence between a clade formed by northern populations of *K. h. murrayi* (from Texas to northern Durango, and likely including central and southern Durango and Zacatecas) and a second distinct clade formed by southern populations of *K. hirtipes* in the Trans-Mexican Volcanic Belt (TMVB), including distinct lineages for *K. h. chapalaense*, *K. h. magdalense*, and *K. h. hirtipes*, and a separate monophyletic lineage composed of *K. h. tarascense* and specimens previously identified as "*K. h. murrayi*" from the TMVB. No genetic material was successfully analyzed from the southern disjunct population of *K. h. murrayi* from central and southern Durango and Zacatecas; its presumed identification as *K. h. murrayi* is based on ecological habitat data.

Weaver et al. (2022) recommended that the southern TMVB populations of "*K. h. murrayi*" be considered as synonymous with *K. h. tarascense*, and we agree. The new distributional data they presented also indicated that the revised combined range of *K. h. tarascense* plus southern TMVB "*K. h. murrayi*" extends over much of the central plateau of the TMVB in a historically somewhat contiguous fashion. However, the populations in the endorheic basin of Lago Pátzcuaro (the originally described *K. h. tarascense*) and in southern Querétaro (not yet genotyped) both appear to be disjunct. As a result of all these changes, the newly revised range of *K. h. murrayi* has decreased by about 15% and the range of *K. h. tarascense* has increased hugely, by over 5000%.

40. *Kinosternon scorpioides*: Caballero et al. (2022) performed a genomic analysis on 24 specimens of *K. scorpioides* from Colombia, including 15 *K. s. scorpioides* from three separate continental watersheds (Caribbean, Orinoco, and Amazon), as well as 9 *K. s. albogulare* from San Andrés Island. They analyzed a dataset of 35,507 SNPs with up to 98% missing data across these 24 samples as well as sequences of an intron of the R35 gene for a subset of 383 SNPs from 7 of their samples with 10% missing data. They concluded that each of their four watershed populations represented a separate Evolutionarily Significant Unit and that each should be treated as a separate management unit and considered for taxonomic elevation.

Due to their limited specimen sampling, lack of comparative genetic analysis of other Central or South American *K. scorpioides* outside of Colombia, excessive and unbalanced missing data across genetic samples, and other discrepancies in their results, more comprehensive phylogeographic studies are needed to delineate management or taxonomic units in Colombia and across the widespread range of *K. scorpioides* in South America (see Annotation 37 above for a subsequent study).

41. *Kinosternon sonoriense*: Molina-Serrano et al. (2025) discovered and described a disjunct population of what they identified as *K. sonoriense* in the Río Culiacan basin of Sinaloa, extending the range of the species substantially southward. They noted that six males in the population were significantly larger than four females, the opposite condition from populations further north, but unfortunately measured only curved carapace length (CCL) rather than straight-line carapace length (SCL), making comparisons to northern populations more difficult. We provide estimated maximum SCL measurements for this southernmost population based on SCL vs. CCL regression equations based on 19 specimens from northern populations (J.B. Iverson, unpubl. data) and encourage genetic analysis of this disjunct population.

42. Testudinoidea: Authorship of this family-level name was given as Fitzinger 1826 in TTWG (2021). However, in accordance with the ICZN (1999; Article 36, Principle of Coordination), this name should instead be attributed to Batsch 1788.

43. *Chrysemys picta* and *C. p. picta*: In a paper we previously overlooked, Schmidtler et al. (2014) argued that the original description of *Testudo picta* by Schneider (1783) was invalid since it was published as a "footnote" and that Schneider had not personally examined the type specimen, but relied on notes provided by Hermann (see our previous TTWG Annotation 12:13 for further details regarding our opinion whether Schneider or Hermann *in* Schneider should be considered the original author). Schmidtler and colleagues recommended that the official authorship of *Testudo picta* should instead be attributed to Gmelin (1789:1045), who provided a second description of the species, but also appropriately attributed it to both Schneider and Hermann.

We disagree with the interpretation by Schmidtler et al. (2014) that the Schneider (1783) publication of *Testudo picta* was invalid. We note that the name was formally published and recognizably described and is therefore nomenclaturally available according to the criteria of ICZN Articles 11 and 12. The name was not published as a junior synonym and that its original publication was somehow considered a "footnote" by either Schneider or Schmidtler et al. is irrelevant. All authors since the original description have attributed the name *Testudo picta* to Schneider (1783) and so it remains.

44. *Chrysemys picta picta* and *C. p. marginata*: The estimated ranges and apparent zone of intergradation between these two subspecies has been reviewed and adjusted for this checklist based on an in-depth analysis of distributional and morphological data, expanding on the mitochondrial genetic work by Starkey et al. (2003). We reviewed the means of costal disalignment percentage measurements (methodology as per Hartman 1958) for nearly 6,000 specimens from over 100 populations of both taxa and presumed intergrades and adjusted our map accordingly. Data sources used were Hartman (1958), Waters (1964, 1969), Ernst (1967, 1970),

Pough and Pough (1968), Ernst and Ernst (1971), Groves (1983), Rhodin (1993), Rhodin and Butler (199), Ultsch et al. (2001), Wright and Andrews (2002), Mann (2007), Ernst et al. (2015), and A.G.J. Rhodin (unpubl. data). As a result, the current map now demonstrates an increased area of overlap between these two taxa and a more extensive intergradation zone.

45. *Chrysemys picta bellii*: We have expanded the depiction of the native range of this subspecies into northern and eastern Arizona and southeastern Utah based on previously overlooked work by Lovich et al. (2014) who examined archaeological data and traditional cultural use patterns and demonstrated the former disjunct presence of the subpecies in a few localities along the Little Colorado River watershed basin.

46. *Graptemys oculifera*: Glorioso et al. (2024) documented an allopatric native population of this species in the Bogue Falaya of eastern Louisiana, separate from its previously assumed endemic distribution in the Pearl River watershed. Although the population was noted to be distinct based on genetic analysis of microsatellite loci, the authors did not recommend taxonomic recognition due to its lack of unique mitochondrial control region haplotypes. Instead, they recommended that it could be considered as a distinct population segment under the U.S. Endangered Species Act. It should also be noted that this disjunct population is close to and in the same watershed as the original type locality of Mandeville, Louisiana, as noted by Baur (1890a), validating that as the type locality for *G. oculifera*, as opposed to the erroneously emended restriction of the type locality by Cagle (1953a) to the Pearl River.

47. *Malaclemys terrapin*: Converse et al. (2017) analyzed microsatellite data to show that patterns of genetic diversity along the coastal range of *M. terrapin* are consistent with historical accounts of translocation and cultivation activities. They identified possible instances of human-mediated dispersal by estimating gene flow over historical and contemporary timescales, and recovered six genotypic clusters along the Gulf and Atlantic coasts with varying degrees of admixture, including increased contemporary gene flow from Texas to South Carolina, from North Carolina to Maryland, and from North Carolina to New York. The genetic structure they found did not match the currently recognized subspecific taxonomy, but they made no recommendations for any taxonomic changes, as they lacked sampling from certain critical areas.

48. *Pseudemys alabamensis*: Moreno et al. (2022) used mitochondrial and nuclear microsatellite markers to assess genetic variation within and among populations of *P. alabamensis* throughout its limited range and to estimate admixture with co-occurring *Pseudemys* species. They found no variation in mitochondrial DNA and an excess of homozygosity in microsatellite data, and also documented genetic distinction between the essentially disjunct populations in Alabama and Mississippi, but made no taxonomic recommendations. In addition, they found evidence of introgression and admixture between *P. alabamensis* and *P. concinna* and *P. floridana*, causing conservation concern for this endangered limited-range species.

49. *Pseudemys floridana* and *P. peninsularis*: The type locality for *Pseudemys floridana* was originally given by Le Conte (1830) as the St. John's River in east Florida. It was postulated by Carr (1938b) to be from the lower (northern) reaches of that river, and subsequently restricted as such by Bour (2003). However, the taxon that is native to St. John's River basin, including its lower reaches, is *P. peninsularis*, not *P. floridana,* as noted by Seidel (1994) and Seidel and Ernst (2017). The single remaining syntype of *P. floridana,* supposedly collected in St. John's River, is a subadult shell that has markings suggestive of a possible intergrade between the two taxa, as noted by Ward (1984). The syntype should be genotyped to help resolve this uncertainty, but for now we consider the type locality for *P. floridana* to lie just outside its native range, and indicate it as such on our map. If the remaining syntype of *P. floridana* is identified as being a *P. peninsularis*, it would create a nomenclatural problem for the name *P. floridana*.

50. *Pseudemys gorzugi*: Vandewege et al. (2024) analyzed population genomics of the Rio Grande Cooter, *P. gorzugi*, using single nucleotide polymorphisms (SNPs). They documented that populations inhabiting the Pecos and Black Rivers of New Mexico are distinct from those in the Rio Grande and Devils River of Texas, with admixed individuals at the confluence of the Rio Grande and Pecos. They made no taxonomic recommendations, but advised that the two populations be considered Management Units for conservation purposes.

51. *Trachemys*: Fritz et al. (2024) examined all North American, South American, and West Indian *Trachemys* species, except for *T. hartwegi*, using mitochondrial and nuclear DNA sequences (3221 and 3396 bp, respectively) representing four mitochondrial genes and five nuclear loci. They provided evidence that *T. nebulosa* has captured a deeply divergent foreign mitochondrial genome that renders the mitochondrial phylogeny of *Trachemys* paraphyletic; however, using nuclear markers, *Trachemys* (including *T. nebulosa*) was found to be clearly monophyletic. In addition to the mitochondrial lineage of *T. nebulosa*, they found six additional mitochondrial *Trachemys* lineages: 1) *T. venusta*, 2) *T. ornata* + *T. yaquia*, 3) *T. grayi*, 4) *T. dorbigni* + *T. medemi*, 5) *T. gaigeae* + *T. scripta*, and 6) *T. decorata* + *T. decussata* + *T. stejnegeri* + *T. terrapen*. These six mitochondrial lineages were each found to constitute well-supported clades, and each was corroborated by the nuclear markers.

In addition, according to nuclear DNA, *T. nebulosa* undoubtedly belongs to the genus *Trachemys*, but its deeply divergent lineage of mitochondrial DNA is, with weak support, sister to *Malaclemys terrapin*. According to a molecular clock, the mitochondrial lineages of *M. terrapin* and *T. nebulosa* diverged ca. 7.0 mya, i.e., their split predates the divergence age of the remaining mitochondrial lineages of other extant *Trachemys* taxa, suggestive of a very ancient hybridization event. The two subspecies of *T. nebulosa* and many Mexican and Central American subspecies of *T. venusta* were not clearly distinct in the studied genetic markers and the putatively diagnostic coloration and pattern traits of *T. venusta* subspecies were found to be more variable than previously reported.

For *T. gaigeae,* another mitochondrial capture event is likely because its mitochondrial genome is sister to *T. scripta*, although it is deeply divergent in its nuclear markers and resembles Mexican, Central and South American *Trachemys* species in morphology, sexual dimorphism and courtship behavior. The analyses also provided additional evidence supporting possible subspecies status of *callirostris* and *chichiriviche* under *T. venusta* rather than under *T. callirostris* as well as of *adiutrix* as a subspecies of *T. dorbigni* rather than a distinct species. The analyses also failed to identify *T. taylori* as a lineage distinct from *T. venusta* and the authors designated it as a subspecies, *Trachemys venusta taylori*, but see Annotation 55.

Despite these studies, most of us continue to view *callirostris* and *chichiriviche* as subspecies of *T. callirostris*, and *T. adiutrix* and *T. taylori* as distinct and separate species, not subspecies of *T. venusta* or *T. dorbigni*. See further discussion in Annotation 55.

52. *Trachemys grayi* and *T. venusta*: Phylogenetic work by Fritz et al. (2023) and Fritz et al. (2024) on the relationships and geographic distribution of the species *Trachemys grayi* and *T. venusta* suggested that the distribution of the subspecies *T. grayi emolli* actually extends southwards from Lake Nicaragua (Lago

Cocibolca) to include the Caribbean coast of Costa Rica and into western Caribbean Panama, and that *T. venusta uhrigi* does not extend into Costa Rica and Panama, but reaches its southernmost extent in southern Nicaragua. Further, their work suggested that *T. grayi panamensis* occurs throughout most of Panama, including the Caribbean versant. We accept these range modifications in our current maps.

In adjusting the ranges of *T. grayi* and *T. venusta* it also becomes necessary to address the question of the nomenclatural status of *Testudo panama* Perry 1811 from Panama, previously discussed in TTWG 2012 (Annotation 12:18). That name was appropriately and officially declared to be a *nomen oblitum* by Bauer and Petit (2004) whenever used as a synonym of *Emys venusta* Gray 1856 (making it a subjective senior synonym also of *Trachemys venusta uhrigi* McCord, Joseph-Ouni, Hagen, and Blanck 2010). However, since *Trachemys grayi* (Bocourt 1868) is now considered to be the only *Trachemys* species apparently occurring in Panama (and *T. venusta* apparently does not), and since the description and original plate depicting *Testudo panama* is as phenotypically similar to *Trachemys grayi* as it is to *Trachemys venusta*, therefore *Testudo panama* should now be considered a subjective senior synonym of *Trachemys grayi panamensis*, rather than *Trachemys venusta uhrigi*.

As a result, in accord with ICZN Articles 23.9.1 and 23.9.2, we hereby further designate *Testudo panama* Perry 1811 to also be a *nomen oblitum* whenever used as a synonym of *Emys grayi* Bocourt 1868, and whenever used as a synonym of either of the two currently accepted subspecies of *Trachemys grayi*: *Pseudemys scripta emolli* Legler 1990 and *Trachemys venusta panamensis* McCord, Joseph-Ouni, Hagen, and Blanck 2010, and declare all three to be *nomen protectum* names whenever considered synonymous with *Testudo panama* Perry 1811.

The name *grayi* has been in widespread use throughout the last 50 years under the generic names *Trachemys* or *Pseudemys*, either as a full species or as a subspecies of *scripta* or *venusta*: Pritchard (1967, 1979), Moll and Legler (1971), King and Burke (1989), Ernst and Barbour (1989), Legler (1990), Iverson (1992), David (1994), Merchán Fornelino (2002), Artner (2003), Bonin et al. (2006), Fritz and Havaš (2007), TTWG (2007b, 2009, 2010, 2011, 2012, 2014, 2017, 2021), Rhodin et al. (2008, 2018), McCord et al. (2010), Legler and Vogt (2013), and Fritz et al. (2023, 2024). In addition, the subspecies names *uhrigi*, *emolli*, and *panamensis* have each been in widespread use under either the genera *Pseudemys* or *Trachemys* and/or the species *scripta*, *venusta*, or *grayi* ever since their descriptions, each more than 10 years ago.

53. *Trachemys medemi*: Cuadrado-Ríos et al. (2025) sequenced and annotated complete mitogenomes of *T. medemi*. When their data were combined with the broader genetic data from Fritz et al. (2024), they corroborated the conclusion that *T. medemi* and *T. dorbigni* are most closely related and form a clade that also includes *T. venusta*, *T. ornata*, and *T. yaquia*. No taxonomic recommendations were made.

54. *Trachemys scripta elegans*: Reshetnikov et al. (2023) analyzed the distribution and degree of invasive success of introduced *Trachemys scripta elegans* across Europe and Asia, based on thousands of records from nearly 1,500 separate water bodies, adding many new nations to our checklist of where this species has been formally recorded. They found that in many regions where the climate is warm and favorable, the species is reproducing successfully. In other slightly colder regions, it is overwintering successfully, but without evidence yet of successful reproduction. In less climatically favorable areas (colder, drier), it is only occasionally found as casually released pet trade individuals. As global climate change and warming progresses, areas of current successful overwintering may become conducive to successful reproduction and increased invasiveness.

55. *Trachemys taylori, T. venusta,* and *T. scripta elegans*: Espindola et al. (2022) examined variation in one mitochondrial marker and one nuclear marker, as well as SNPs and microsatellites, from the ranges of *Trachemys taylori*, *T. venusta* (including the subspecies *cataspila*, *venusta*, and *iversoni*, although the latter was lumped with *venusta* by the authors), and *T. scripta elegans* along the Caribbean versant of Mexico. Their analyses supported genetic differentiation between these taxa, characterized by reduced gene flow, including in contact zones of sympatric populations. They also found that non-native introduced *T. s. elegans* constituted a genetically distinct cluster different from native *T. s. elegans*, but made no taxonomic recommendations.

Subsequently, Fritz et al. (2024), as part of their phylogeographic study of nearly all *Trachemys* (see Annotation 51 above), examined the genetic relationships between *T. taylori* and *T. venusta* (including *T. v. cataspila*). Their analyses failed to identify *T. taylori* as a lineage distinct from the polytypic *T. venusta*, and they designated the former as an additional subspecies of the latter. However, based on the previous work by Espindola et al. (2022), which demonstrated clear genetic differentiation between *T. taylori* and *T. venusta* and no gene flow between them, with orders of magnitude more nuclear loci, most of us reserve judgment on which analysis may be more reflective of the phylogenetic relationships, and prefer to continue to recognize *T. taylori* as a distinct species rather than treating it as a subspecies of *T. venusta*. Further work on the relationships of these two taxa is indicated, especially in view of the very small and isolated range of the highly threatened and allopatric *T. taylori*, endemic to the Cuatro Ciénegas basin.

56. *Glyptemys muhlenbergii*: Frétey (2024:12,85) argued for replacing the long-standing prevailing usage of the species name *muhlenbergii* with the alternative orthography *muehlenbergii*, based on his opinion that the name was originally described by Schoepff (1801) as *mühlenbergii*. He cited ICZN (1999; article 32.5.2.1), which states that when the German letter "*ü*" (umlaut) is used in a name, it must be replaced by "*ue*" for names created prior to 1985, and therefore recommended that the orthography of the species name now be changed to *Glyptemys muehlenbergii*.

However, Schoepff (1801:132) described his new species as *Testudo Muhlenbergii* (not using the German *ü* umlaut), both in his own heading for the species on p. 132 as well as on the caption for his plate XXXI. However, the printer of Schoepff's work had labeled all pages with small headers, and the printer's header on p. 132 reads *Testudo Mühlenbergii*. In his text, Schoepff does use the term *T. Mühlenbergii* once in describing his new species' differences from *T. punctata*, but since two of his three name usages are *muhlenbergii* instead of *mühlenbergii*, it is our opinion that *muhlenbergii* should be interpreted as the original orthography, and as such there would be no need to change the current orthography to *muehlenbergii*.

Most other turtle taxonomists since then have agreed, and the use of the *muhlenbergii* orthography has been in widespread prevailing use since the description, with only a few early exceptions. Prominent taxonomists that interpreted and used *muhlenbergii* as the correct orthography include Schweigger (1812), Gray (1830e, 1831d, 1844, 1856b), Duméril and Bibron (1835), Fitzinger (1835), Holbrook (1836, 1842a), Strauch (1865, 1890), Cope (1875b), Boulenger (1889), Siebenrock (1909a), Stejneger and Barbour (1917), Babcock (1919), Pope (1939), Carr (1952), Mertens and Wermuth (1955), Pritchard (1967), and Wermuth and Mertens (1961, 1977). Authors who used the *mühlenbergii* orthography

include Merrem (1820), Temminck and Schlegel (1834), Agassiz (1857a) and Gray (1870c, 1873j). Agassiz established his new genus *Calemys* based on *Testudo mühlenbergii* as the type species, and Gray (1870c, 1873j) copied his orthography, but erroneously recorded the name as *mühlenburgii*. Mertens and Wermuth (1955) pointed out that the usage of *mühlenburgii* by Gray (1870c) was "ex errore," and Wermuth and Mertens (1977) further noted that the use of *mühlenburgii* by Gray (1870c) (but altered by them to "*muehlenburgii*" in their synonymy) was "ex errore pro *T. muhlenbergii* Schoepff." No authors since Agassiz (1857a) have used the *mühlenbergii* orthography.

Based on this history, and on concern for the potential confusion caused for conservation legislation (e.g., CITES and ESA regulations) if the name of this Critically Endangered species were to be changed, we hereby follow ICZN (1999:xxviii) recommendation measure number 11 to declare the name *Testudo muhlenbergii* Schoepff 1801 a *nomen conservandum* and the names *mühlenbergii* and its alternative orthography *muehlenbergii* to be unavailable names. The following >10 authors of >25 taxonomic works in the last 50 years have all used *muhlenbergii*, and none since 1857 have used *mühlenbergii* or *muehlenbergii* as a valid name: Wermuth and Mertens (1977), Ernst and Bury (1977), Pritchard (1979), Berry and Shine (1980), King and Burke (1989), Ernst and Barbour (1989), Iverson (1992), Klemens (1993), Bickham et al. (1996), USFWS (1997, 2022c), Artner (2003), Fritz and Havaš (2007), TTWG (2007b, 2009, 2010, 2011, 2012, 2014, 2017, 2021); Rhodin et al. (2008, 2018b), Buhlmann et al. (2009), Ernst and Lovich (2009), van Dijk (2011, IUCN Red List), TCC (2011, 2018), Ceballos et al. (2013), Shoemaker and Gibbs (2013), Seidel and Ernst (2017), and Zappalorti (2023).

57. *Terrapene coahuila*: Cortés-Rodríguez et al. (2021) assessed genetic diversity and structure of the isolated and endemic *T. coahuila* in Cuatro Ciénegas, using two mitochondrial DNA genes (cytochrome b and D-loop), one nuclear gene (GAPD), and 12 nuclear microsatellite loci. Their results dated the time to the most recent common ancestor (*T. carolina*) for *T. coahuila* at ca. 10.1 mya, with highest diversification during the Holocene. They identified two genetic clusters, western and eastern, one on either side of the main historical barrier, the Sierra San Marcos. Demographic results supported a divergence between western and eastern populations from 47,500 to 110,950 years ago, and colonization of the southernmost western distribution (Churince) at 28,500–66,500 years ago, which experienced a recent bottleneck.

58. *Platysternon megacephalum*: Gong et al. (2023) genotyped 143 samples from across the geographic range of the Big-headed Turtle, using mitochondrial DNA fragments and genome-wide single nucleotide polymorphisms (SNPs) to investigate phylogenetic relationships, genetic divergences, and demographic history of the species and its three generally recognized subspecies: *P. m. megacephalum*, *P. m. peguense*, and *P. m. shiui*.

Their results suggested the presence of five distinct clades with separate demographic histories. They recommended that three of those clades be considered taxonomically distinct (*P. m. megacephalum*, *P. m. peguense*, and a separate clade from the Baise region of western Guangxi). They found weaker support for a separate clade in northern Vietnam and Hainan, previously recognized as *P. m. shiui*, but noted that specimens from those regions clustered more closely with *P. m. megacephalum* than *P. m. peguense*.

Based on this genetic differentiation, the authors suggested elevating the subspecies *P. m. peguense* to species level, and suggested synonymizing *P. m. shiui* under a monotypic species-level *P. megacephalum*. They left their newly discovered clade from the Baise region of western Guangxi as a temporarily unnamed cryptic taxon. However, whether that lineage is distinct enough to warrant taxonomic distinction will require more comprehensive sampling within the Guangxi basin to establish its level of distinctiveness.

Overall, we are not convinced that the authors have presented compelling genetic evidence that the two taxa *megacephalum* and *peguense* each warrant recognition at species rank, and we therefore continue to treat each of them as subspecies of *P. megacephalum* and also await further work on the lineage from the Baise region. Specifically, their results showed major discrepancies between their dated phylogeny using mitochondrial genes and the patterns of divergent trajectories in their genomic demographic history analysis. They also presented unrealistically high genetic differentiation fixation index (F_{ST}) values between lineages which challenged the reliability of their population differentiation estimates.

In addition, we are not convinced that the genetic relationships and minimal morphological details provided in this study offer enough evidence to synonymize *P. m. shiui*. The distribution of this subspecies in northern Vietnam and Hainan is congruent with a distinct clade in their mitochondrial tree (Clade II) which corresponds with a clade in their maximum likelihood tree based on genome-wide SNPs. The individuals from northern Vietnam and Hainan also formed a genetic cluster in their structure analysis and PCA. We therefore leave this clade as a valid subspecies of *P. megacephalum* pending further analysis. We do note, however, that the previous morphology-based suggestion by Ernst and Laemmerzahl (2002) that Hainan specimens of *Platysternon* are intergrades of *P. m. shiui* and *P. m. peguense* was not supported by the present study, but instead cluster genetically with the *P. m. shiui* lineage. We have adjusted our map accordingly.

Additional genomic work on these taxa by Ngo et al. (2023) largely corroborated these previous results. They sequenced the mitochondrial ND4 gene of 450 specimens of *P. megacephalum*, but only 15 were wild-collected and the rest were from trade confiscations without locality data. They noted significant differences between wild-collected *P. m. peguense* and confiscated *P. m. megacephalum*, and suggested that these differences supported possible elevation of the two taxa as distinct species rather than subspecies. They also noted that wild-collected *P. m. shiui* were most closely related to *P. m. megacephalum*. However, they made no specific taxonomic recommendations in either situation.

59. Geoemydidae fossils: Garbin et al. (2020) examined numerous fossil taxa from the Siwalik deposits (Miocene through Early Pleistocene age) of India and Pakistan that TTWG (2017, 2021) and others (e.g., Fritz and Havaš 2007) had previously listed as synonyms or *nomen dubium* names under the various extant species *Batagur dhongoka*, *Batagur kachuga*, *Geoclemys hamiltonii*, *Hardella thurjii*, *Melanochelys tricarinata*, and *Melanochelys trijuga*. They confirmed or agreed with many of the prior synonymizations, except: 1) they removed *Batagur cautleyi* † Lydekker 1885 from the synonymy of *Hardella thurjii*, but did not assign it to a new species, instead only identifying it as *Batagur* sp. indet. †, 2) they removed *Nicoria tricarinata sivalensis* † Lydekker 1889 from the synonymy of *Melanochelys tricarinata* and suggested reallocating it to *Melanochelys sivalensis* †, an action that created a junior homonym, requiring them to establish the replacement name *Melanochelys tapani* † (*nomen novum*) for this taxon, and 3) they recognized *Bellia sivalensis* † Theobald 1877 as distinct from *Melanochelys trijuga indopeninsularis* and allocated it to a resurrected *Melanochelys sivalensis* † (senior homonym), also synonymizing *Clemmys hydaspica* † Lydekker 1885, *Clemmys theobaldi* † Lydekker 1885, and *Clemmys punjabiensis* † Lydekker 1885 under this species. We acknowledge these revisions and reallocations and

have removed all these taxa from this checklist and will instead include them in the next TEWG fossil checklist.

60. *Batagur affinis*: Salleh et al. (2023), in a phylogeographic study of 651 base pairs of the mitochondrial D-loop in 120 *B. affinis* from two populations (30 samples each) from the west coast of peninsular Malaysia (*B. a. affinis*) and two (30 samples each) from the east coast (*B. a. edwardmolli*), recorded the presence of 46 unique mutations that defined six mitochondrial haplotypes. They recovered clear genetic differentiation between the two subspecies, but also a few *B. a. edwardmolli* haplotypes (3 of 30 animals) in their northern west coast (Kedah) sample of *B. a. affinis*. They interpreted this as a novel and natural range extension of *B. a. edwardmolli* from the east coast across the peninsula to the west coast, resulting in the presence of both subspecies in northwestern Malaysia. However, in view of the decades-long conservation practices of hatching, headstarting, and releasing *B. affinis* from various breeding facilities in peninsular Malaysia (Moll et al. 2015 CBFTT), it is at least as likely that specimens with east coast haplotypes were at some point in the past released into west coast waters, especially prior to the recognition of the two coastal forms as separate subspecies (Praschag et al. 2009a). Whether the co-occurrence of both haplogroups in the Kedah region is the result of human release or natural migration is an open issue that requires nuclear data on the extent of interbreeding and admixture. Until such data are available, we feel that the most likely interpretation is that the presence of *B. a. edwardmolli* haplotypes on the west coast of Malaysia is due to human translocation activities.

61. *Malayemys khoratensis* and *M. subtriguga*: Chaianunporn et al. (2024a) analyzed morphological traits of 109 *Malayemys* specimens along with mitochondrial DNA from 60 specimens from across northeastern and eastern Thailand. They found that both *M. khoratensis* and *M. subtrijuga* occur in northeastern Thailand. They clarified that the distribution of *M. khoratensis* includes the entire Chi River basin, the Mun River basin north of the Mun River, and the upper Mekong River, while the distribution of *M. subtrijuga* includes the Mun River basin south of the Mun River to the lower Mekong River and also a small part of the lower Chin River basin. We have updated our distribution maps accordingly.

Further genetic work by Chaianunporn et al. (2024b) on *M. khoratensis* identified 13 unique haplotypes from 64 samples. The total haplotype diversity was 0.82, while nucleotide diversity was 0.003. The haplotype network revealed no distinct population structure among the river basins (Chin, Mun, and Mekong), suggesting gene flow across these regions, due either to natural movement between basins or potential human-mediated translocations linked to the turtle trade.

62. *Morenia petersi*: The type locality for this species has always been listed as three sites from which Anderson (1879) examined four syntype specimens: a female from Huzurapur, Jessore District, two males from Furreedpore [= Faridpur], and one male from Dacca [= Dhaka], all in present-day Bangladesh. The female syntype from Jessore District has not been located (Iverson 2022a). Murthy and Das (2009) and Das (2009) listed two specimens from Faridpur in the ZSI collection (ZSI 155–156) as syntypes (actually paralectotypes; now apparently lost, according to Kundu et al. 2018). Fritz et al. (1994:164) identified and designated a single male in the ZMB collection (ZMB 8865, collected by Anderson with generalized locality data of "Lower Bengal") as the "holotype" (actually lectotype), figured by Anderson (1879:pl.59; reproduced by Bour 2009 and Das 2009). Hence, it is clear that the male in the ZMB collection was actually obtained by Anderson from Dhaka, Bangladesh, and we hereby restrict the type locality for *M. petersi* as such.

63. *Cistoclemmys*: Frétey (2024) suggested that authorship of the generic name *Cistoclemmys* should be attributed to Gray *in* Swinhoe 1863 since he also noted that Swinhoe (1863) was supposedly published earlier than Gray (1863e). However, Swinhoe credited Gray as the author and provided the same exact description within quotation marks as Gray had provided and also provided the exact page number from Gray's description, which therefore was clearly available earlier. Therefore, in our opinion, the attribution of the name *Cistoclemmys* remains Gray 1863e rather than Gray *in* Swinhoe 1863.

64. *Cuora amboinensis* (sensu lato) species complex: Blanck et al. (2023) analyzed morphology of 288 range-wide specimens of the *Cuora amboinensis* (sensu lato) species complex and mitochondrial and nuclear DNA phylogeography in 104 of these specimens. They concluded that populations from Assam, India, and the Philippines were each distinct at the species level, and described these populations as *C. praschagi* and *C. philippinensis*, respectively. They also concluded that the apparently endemic taxon from Myanmar previously identified as *C. amboinensis lineata* is a species distinct from the other members of the complex, and elevated it as *C. lineata*. They also noted that a disjunct population of what they interpreted as this taxon occurs in an isolated higher-altitude valley in Manipur, India. In addition, they concluded that the previously designated subspecies *C. amboinensis couro* and *C. amboinensis kamaroma* are sister taxa and separate from *C. amboinensis* (sensu stricto) and therefore designated each of them as subspecies of *C. couro* instead of as subspecies of *C. amboinensis*. At the same time, they restricted *C. amboinensis* (sensu stricto) to the eastern Indonesian islands of Sulawesi and the Moluccas, including Ambon, the original type locality for the species. They also noted that populations of *C. amboinensis* inhabiting the higher altitude central plateau of Sulawesi are genetically and morphologically distinct from the low-altitude Sulawesi populations referable to *C. amboinensis amboinensis*, and as a result described the upland form as the new subspecies *C. amboinensis aurantiae*.

Although we are somewhat uncertain as to whether the revised and new descriptions of these several taxa have indeed been placed at the most appropriate taxonomic levels, we currently accept all of them as valid. We note further that Blanck et al. (2023) also identified some genetically distinct animals from unknown localities (trade specimens) and that they were uncertain about taxonomic allocations for some insular populations, notably those from the Nicobar Islands, Palawan, and the Sulu Archipelago. Clearly, further phylogeographic work will be warranted on this large and widespread species complex.

65. *Cuora galbinifrons, C. bourreti,* and *C. picturata*: Ihlow et al. (2023) re-investigated the genetic relationships among the species of the *C. galbinifrons* complex based on 2 mitochondrial genes and 12 microsatellite loci obtained from 394 individuals from assurance colonies, zoos, private collections, and museum specimens (mostly without locality data). Their well-supported mitochondrial phylogeny confirmed *C. bourreti* as the sister clade to *C. galbinifrons* which together represented the sister clade to *C. picturata*. Their microsatellite analysis confirmed three distinct clusters that were in agreement with these species. Previously, Liu et al. (2019) argued for the relegation of *bourreti* to a subspecies of *galbinifrons* based on inconclusive genetic results. Based on the results presented by Ihlow et al. (2023), we continue to recognize the three taxa as full species.

66. *Cuora lineata*: Zug (2022) synonymized this taxon under *C. amboinensis kamaroma*, citing work by Ernst et al. (2016), which was based only on color morphology. However, Zug did not refer to Protiva et al. (2016) nor TTWG 2017 (Annotation 17:41) who

both disagreed with Ernst et al. and called for genetic analysis. Blanck et al. (2023) subsequently analyzed the phylogeography of this taxon (see Annotation 64 above), and confirmed its status as a distinct species.

67. *Cuora pani*: Li et al. (2023) analyzed mitochondrial and nuclear DNA in 14 specimens of *C. pani*. They confirmed the reciprocal monophyletic distinction between this species and *C. aurocapitata* as previously described by Spinks et al. (2012a) (see TTWG Annotation 12:22) and described three intraspecific haplotypes with low diversity and shallow distinction, but made no taxonomic recommendations.

68. *Mauremys annamensis* and *M. mutica*: Ngo et al. (2025) analyzed mitochondrial (ND4) and nuclear (R35 intron) DNA from 38 specimens of the Critically Endangered and nearly extirpated *M. annamensis*, of which 18 were obtained from local trade of known provenance in central Vietnam. They also compared these new samples to GenBank sequences of *M. mutica* from mainland China and from Hainan. They confirmed and corroborated earlier findings by Somerová et al. (2015; see TTWG Annotation 17:49) and Fong et al. (2019; see Annotation 21:69) that there are two mitochondrial lineages within *M. annamensis* that are geographically isolated and evolutionarily significant, and important to consider for conservation repatriation efforts, but they made no taxonomic recommendations.

The results of their analysis of *M. mutica* suggested that the population of that species from Hainan appeared likely to have evolved through introgression between a lineage distantly related to true *M. mutica* from mainland China and *M. annamensis* and a lineage closely related to the two mitochondrial haplotypes of *M. annamensis* from central Vietnam. Once again, no taxonomic recommendations were made.

69. *Mauremys leprosa saharica*: In our previous checklists, we listed the name *Emys fraseri* Gray 1869c as a "*nomen dubium, partim*" in the synonymies of both *Mauremys leprosa leprosa* (Schoepff *in* Schweigger 1812) and *M. l. saharica* Schleich 1996a, and we listed *M. l. erhardi* Schleich 1996a in the synonymy of *M. l. leprosa*, following Fritz et al. (2006a) and Fritz and Havaš (2007). However, Veríssimo et al. (2016) showed that *Mauremys leprosa* haplotype lineage B (*M. l. saharica*, with synonyms *M. l. erhardi*, *M. l. zizi* Schleich 1996a, and *M. l. vanmeerhaegei* Bour and Maran 1999), is distinct and separate from haplotype lineage A (*M. l. leprosa*, with synonyms *M. l. atlantica* Schleich 1996a, *M. l. marokkensis* Schleich 1996a, and *M. l. wernerkaestlei* Schleich 1996a). Haplotype lineage B (*M. l. saharica*) occurs in northern, southern, and eastern Morocco and Algeria, both north and south of the Atlas Mountains, and characterizes specimens collected close to the type localities for *E. fraseri*, *M. l. saharica*, *M. l. erhardi*, *M. l. zizi*, and *M. l. vanmeerhaegei*. As such, *Emys fraseri* is not a *nomen dubium*, but instead, a senior subjective synonym of these other more recently named taxa. However, no authors since Gray (1873a) have recognized *Emys fraseri* as a valid taxon, and it has been universally synonymized in different generic combinations under *Emys leprosa* Schoepff *in* Schweigger 1812 by all authors since, starting with Boulenger (1889).

Therefore, *E. fraseri* can be designated a *nomen oblitum* under the criteria of ICZN Art. 23.9, as it meets the requirements of both paragraphs 23.9.1.1 and 23.9.1.2. Specifically, 1) *Emys fraseri* has not been recognized as valid after 1899, and 2) the younger names *M. l. saharica*, *M. l. erhardi*, *M. l. zizi*, and *M. l. vanmeerhaegei* have been used as valid, or as subjective synonyms of *M. l. saharica*, in the following 25 works by more than 10 authors over more than 10 years: Schleich (1996a), Schleich et al. (1996), Bour and Maran (1999), Keller and Busack (2001), Artner (2003), Fritz et al. (2005c, 2006a), Bonin et al. (2006), Fritz and Havaš (2007), Rhodin et al. (2008), TTWG (2009, 2010, 2011, 2012, 2014, 2017, 2021), Lovich et al. (2010), Palacios et al. (2015), Velo-Antón and Pinya (2015), Veríssimo et al. (2016), Bertolero and Busack (2017; CBFTT), Loulida et al. (2019, 2021), Martínez del Mármol et al. (2019), and Toledo et al. (2021).

As such, *Emys fraseri* Gray 1869c is hereby declared a *nomen oblitum* whenever considered synonymous with either *Mauremys leprosa saharica* Schleich 1996a, *Mauremys leprosa erhardi* Schleich 1996a, *Mauremys leprosa zizi* Schleich 1996a, and/or *Mauremys leprosa vanmeerhaegei* Bour and Maran 1999. The name *Mauremys leprosa saharica* Schleich 1996a is declared a *nomen protectum* and remains in usage for the concerned taxon.

70. *Mauremys nigricans* and *Damonia crassiceps*: We noted previously (TTWG 2017; Annotation 17:48) that the taxon *Damonia crassiceps* Gray 1870c, which had traditionally been synonymized under *Malayemys subtrijuga*, was interpreted by Sumontha et al. (2016) as not representing a species of *Malayemys*, but they did not suggest an alternative taxonomy. Dawson et al. (2025) have now demonstrated through morphometric analysis of the iconotype figures of *D. crassiceps* with comparisons to the three species of *Malayemys* and *Mauremys nigricans* (Gray 1834a), that *D. crassiceps* is most similar to *M. nigricans*. They have therefore removed it from the synonymy of *M. subtrijuga* and reallocated it to the synonymy of *M. nigricans*. We agree with this taxonomic reallocation.

71. *Melanochelys trijuga trijuga*: The synonymized junior taxon *Emys belangeri* Lesson 1831b was described as having been collected by Charles Bélanger and known from two regions in India: "Bengale" (= West Bengal, likely nr. Calcutta [Kolkata]) and "Carnate" (= the Carnatic region of the southeastern Indian coast). Smith (1931) implicitly restricted the type locality to "Bengal" without comment, but this locality is within the range of *M. t. indopeninsularis* (Annandale 1913) and should have resulted in *belangeri* replacing *indopeninsularis* as the older valid name for that northern Gangetic basin taxon.

However, *M. t. indopeninsularis* has been globally accepted since its description in 1913, and *E. belangeri* has been universally considered a junior synonym of *M. t. trijuga* since Gray (1844), based on the morphology depicted in the drawing of its only known type specimen (Lesson 1831b:pl.1, reproduced by Bour and Schmidtler 2014:f.9E). That type specimen has not been found at the MNHN in Paris where it was originally held and figured (Smith 1931; Bour and Schmidtler 2014). As such, the figure of the lost type specimen becomes the designated lectotype and we thereby also restrict the type locality to the Carnatic region where Charles Bélanger was stationed for a long time (Pondicherry [Puducherry], the French port in Tamil Nadu), where he collected many animals (Bélanger 1834), including *E. belangeri*, as noted by Lesson (1831b). We hereby reject the previously restricted type locality of "Bengal" for *E. belangeri* as suggested by Smith (1931) and instead restrict and emend the type locality to "Carnate (Carnatic region), nr. Pondicherry, Tamil Nadu, India," which places it correctly within the distribution of *M. t. trijuga* (as currently understood) and avoids the need to consider having the name *belangeri* take precedence over the name *indopeninsularis* for the northern taxon.

72. *Sacalia bealei* and *S. quadriocellata*: In a continuation of their previous work (see TTWG Annotation 21:73), Lin et al. (2022) analyzed the phylogeography of *Sacalia*, using more specimens from a wider geographic area. Mitochondrial cytochrome b and 16 nuclear genes from 87 specimens of *S. bealei* and *S. quadriocellata* from many localities were sequenced and analyzed. Three major mitochondrial clades were identified: one for *S. bealei* and two for *S.*

quadriocellata, with five mitochondrial subclades and three nuclear clusters for various populations of *S. quadriocellata*. In addition, they inferred that a few specimens from a contact zone of the two species are of hybrid origin (*S. bealei* x *S. quadriocellata*) and could represent an incipient hybrid speciation process. However, in the face of the presented evidence, this conclusion requires additional investigation. One of the mitochondrial subclades of *S. quadriocellata*, endemic to Hainan, had previously been referred to as a distinct species by the authors, *S. insulensis*, and its distinctiveness was corroborated. However, they noted that more genomic and morphological work was needed, because recognizing the Hainan population as a species would render the other mitochondrial clades paraphyletic. The authors considered that a future solution could be to recognize multiple additional *Sacalia* species, if additional evidence supports this. We continue to consider *S. insulensis* to be a junior synonym of *S. quadriocellata*.

73. **Rhinoclemmydinae:** Frétey (2024:67) suggested that the orthography of this subfamily name, attributed to Gray (1873j), that we had designated as Rhinoclemmydinae in our previous four checklists (TTWG 2012, 2014, 2017, 2021), was rendered incorrectly according to Article 29 of the Code (ICZN 1999) and should be changed to Rhinoclemmyinae. However, Frétey (2024) overlooked that Article 29.5 of the Code rules that the prevailing usage has to be maintained even if the derivation of the name is not in accordance with the grammatical procedures explained in the same Article. We therefore maintain our usage of Rhinoclemmydinae.

74. ***Testudo scabra*:** Frétey (2024:74) noted that the designation by Brown (1908) of *Testudo scabra* as the type species of *Trachemys* may technically invalidate the declaration by Rhodin and Carr (2009) that the name is a *nomen oblitum*, since it does not meet the requirement of ICZN Article 23.9.1.1 that it has not been used as a valid name since 1899. As such, Frétey (2024) suggested that a petition to the ICZN would need to be submitted to request the use of their plenary powers to suppress the name; otherwise it might technically have precedence over the widely used and established name *Rhinoclemmys punctularia*. We would be in support of such a petition if it was felt to actually be needed.

75. ***Aldabrachelys*, *Astrochelys*, and *Cylindraspis*:** Kehlmaier et al. (2023), in a wide-ranging in-depth study, combined analysis of ancient DNA with phylogenetic, ancestral range, and molecular clock analyses of radiocarbon dates and paleogeographic evidence to delineate the diversity and biogeography of extinct and extant tortoises of Madagascar and the western Indian Ocean.

They genotyped several fossils of extinct Madagascar giant tortoises, including *Aldabrachelys abrupta* and *A. grandidieri*, and discovered that a single tibia previously thought to represent a smaller subadult specimen of the former species was actually deeply genetically divergent in both mitochondrial and nuclear DNA, and described it as an extinct large taxon, *Astrochelys rogerbouri*, sister to extant *A. yniphora*.

They also genotyped several specimens of the extinct Mascarene *Cylindraspis* tortoises, including some of the types, and presented evidence regarding the evolution and dispersal of these and other giant and large tortoises out of Africa to Madagascar, the granitic Seychelles, the Mascarenes, and Aldabra. Genotyping for the first time one of the only two alcohol-preserved specimens of *Cylindraspis* hatchlings (MNHN 1991, 1991A), originally described as the syntypes of *Testudo rotunda* Latreille *in* Sonnini and Latreille 1801, a name previously interpreted as a *nomen dubium* and possibly representing either *C. peltastes* or *C. vosmaeri*, was confirmed as representing the latter species.

Mitogenomic analysis of extinct and extant tortoises from the granitic Seychelles and Aldabra, including many type specimens, failed to detect any differences between the currently designated subspecies (*Aldabrachelys gigantea gigantea*, *A. g. arnoldi*, *A. g. daudinii*, and *A. g. hololissa*), although a subfossil specimen from the granitic Seychelles differed slightly from the other more modern specimens. As a result, Kehlmaier et al. suggested that the various subspecies of *A. gigantea* not be recognized as distinct.

However, nuclear genomics have still not been fully analyzed for the *A. gigantea* complex and morphological differences among the subspecific taxa are reasonably prominent. In addition, the one specimen of "*A. g. hololissa*" analyzed genetically by Kehlmaier et al. (a paralectotype, NHMUK 1888.3.20.1) was previously identified by Gerlach and Canning (1998) as representing an Aldabran *A. g. gigantea* and removed by them from the type series of *A. g. hololissa*. In view of these inconsistencies, most of us continue to treat the different morphotypes from the granitic Seychelles as recognized subspecies, separate from the Aldabran *A. g. gigantea*.

76. ***Aldabrachelys gigantea*:** Çilingir et al. (2022a,b) analyzed genome-wide SNPs from 33 *A. gigantea* from each of the three main islands in the Aldabra Atoll, Seychelles, and used a chromosome-level *de novo* assembled genome plus low-coverage resequencing of 30 individuals from wild populations and two zoo individuals. They detected patterns of within-island population structure, notably on Grande Terre, but no differentiation between the islands.

77. ***Chelonoidis carbonarius*:** Gallego-García et al. (2023b), in a detailed study of genomic phylogeography of *Chelonoidis carbonarius* in Colombia, and building on previous single-gene mitochondrial studies (Vargas-Ramírez et al. 2010a) that had suggested two separate genetic lineages in Colombia, demonstrated that populations in western and eastern Colombia, isolated from each other by the Andes, were markedly different. Using a combination of restriction-site-associated DNA sequencing and environmental niche modelling, they provided three independent lines of evidence that strongly support the recognition of species-level distinctions between the East and West lineages: allopatric reproductive isolation, local adaptation, and ecological divergence. However, they recommended that the two Colombian populations be provisionally treated as separate evolutionarily significant units (ESUs) for conservation purposes, pending further rangewide genomic analysis.

78. ***Chelonoidis niger*:** Gaughran et al. (2025) analyzed whole-genome relationships among all 13 extant taxa of Galapagos Giant Tortoises to assess potential support for delimiting each of them as species rather than subspecies. They suggested that their findings of model-based and divergence-based results indicated that these taxa form a species complex consisting of a minimum of 9 species, with most of their analyses supporting 13 species. They found mixed support for the species status of taxa living on the same island (e.g., on Isabela), with some models suggesting multiple populations of a single species per island. They suggested that their results made clear that Galapagos Giant Tortoise taxa represent different stages in the process of speciation, with some taxa further along in that evolutionary process than others.

We agree that their results indicate that all extant taxa of Galapagos tortoises are distinct evolutionary lineages, and that some are more distinct and others less distinct from each other, but whether these lineages represent separate species or subspecies remains an open question. Indeed, the authors noted several areas of ambiguity in how best to interpret their results and noted that although the preponderance of evidence supported several genetically distinct taxa, whether these distinct taxa should be considered species or subspecies was a more nuanced question.

In view of the shallow genomic divergences found among the lineages of these insular taxa of *Chelonoidis niger* as compared to the deeper divergences found among various regional populations

of continental *C. carbonarius* (Gallego-García et al. 2023b, see Annotation 77 above) and other tortoise species (Kehlmaier et al. 2021a), most of us continue to maintain our more conservative view of considering all the Galapagos tortoise taxa to be distinct at the subspecies level.

79. *Chelonoidis niger chathamensis*: Jensen et al. (2022a) sequenced mitochondrial DNA from five subfossil giant tortoise shells collected in 1906 from a cave on central San Cristóbal, Galapagos, as well as from extant tortoises from eastern San Cristóbal. One of the subfossil shells was radiocarbon dated to ca. 1540 CE, and another is the holotype of *C. n. chathamensis* (CAS 8127; Van Denburgh 1907, 1914). They discovered that the haplotype of these five subfossil specimens is markedly different from that of the extant tortoises on eastern San Cristóbal, currently referred to as *C. n. chathamensis*, and all other known Galapagos giant tortoises, and is most closely related to the saddlebacked taxa from Española (*C. n. hoodensis*) and Pinta (*C. n. abingdonii*). They also noted that the mitochondrial haplotypes of an extant specimen of *C. n. chathamensis* from eastern San Cristóbal, as well as one of the original paratypes of *C. n. chathamensis* (CAS 8133, collected alive on central San Cristóbal in 1906), are most closely related to the dome-shelled taxon from eastern Santa Cruz, *C. n. donfaustoi*.

The authors did not determine whether the extant lineage on eastern San Cristóbal is distinct enough to warrant naming as a separate taxon, preferring to analyze nuclear genomic markers first, but its degree of mitochondrial haplotype differentiation is equal to or greater than that which separates the other currently recognized *C. niger* taxa (Jensen et al. 2022a). Further, as we previously noted (TTWG 2007b; Annotation 07:59), the taxon *C. n. chathamensis*, described by Van Denburgh (1907, 1914) from central San Cristóbal, was regarded by Pritchard (1996) to have become extinct in its central and western range by 1933 (see Banning 1933), and he considered the extant population of tortoises in eastern San Cristóbal to be an undescribed subspecies. Tapia et al. (2020) also questioned whether the extinct population from the central and western parts of the island was a separate taxon from the extant population in the east.

Based on these considerations, we wonder whether the extant and extinct populations of tortoises on San Cristóbal may be separate taxa of *C. niger*, or simply different mitochondrial lineages. If their divergence is well supported by nuclear data and if they become regarded as separate taxa, one would then technically need to consider *C. n. chathamensis* to be extinct, as recorded by Fritz and Havaš (2007), and the extant population to be an unnamed taxon in need of formal description, as suggested by Pritchard (1996).

However, the name *chathamensis* has come to refer primarily to the extant population in the east, despite its holotype now identified as representing the extinct central (and possibly western) populations. Creating a new name for the extant taxon would be counter to prevailing usage of the name *chathamensis* and could cause nomenclatural instability with potential consequences for conservation efforts for this IUCN Red List Endangered taxon. Alternatively, since one of the original type series paratypes of *C. n. chathamensis* (CAS 8133; alive when collected) has essentially the same mitochondrial haplotype as the extant eastern population, as shown by Jensen et al. (2022a), one could also consider invoking ICZN Article 75.6 to petition the ICZN to set aside the subfossil holotype and designate CAS 8133 as a neotype, thus maintaining prevailing usage of the name *chathamensis*. The extinct central San Cristóbal taxon would then require a new name and type specimen.

80. *Chelonoidis niger phantasticus*: As previously noted (TTWG 2021; Annotation 21:85), a second specimen of this giant tortoise lineage from Fernandina in the Galápagos Islands was discovered in 2019 and genetically analyzed, with preliminary unpublished results at that time apparently confirming its identification. Verification of its conspecificity with the single specimen of *C. n. phantasticus* previously known (the holotype collected in 1906) has now been documented by Jensen et al. (2022b), who sequenced both mitochondrial and nuclear genomes of both specimens and compared them to all other extant Galápagos tortoise lineages and the recently extinct *C. n. abingdonii*. This verifies that *C. n. phantasticus* is not extinct, though only a single living female is currently known.

Jensen et al. (2022b) also elucidated nuclear DNA genomic monophyly of all the saddlebacked taxa (*C. n. phantasticus, abingdonii, duncanensis,* and *hoodensis*) and one semi-saddlebacked taxon (three specimens from the extant population of *C. n. chathamensis* from eastern San Cristóbal), a pattern not seen in previously generated mitochondrial DNA phylogenies. Notably, the mitochondrial haplotype of two other specimens from the extant population from eastern San Cristóbal is more closely related to dome-shelled *C. n. donfaustoi* (Jensen et al. 2022a).

81. *Geochelone elegans*: The synonymized taxa *Peltastes stellatus maura* and *Peltastes stellatus sebae* were cited in previous editions of this checklist as described by Gray (1870c; published September 1870). However, they actually appeared first in Gray (1870b; published June 1870). The only difference between the two descriptions was that the latter name was spelled as *sebae* in the first publication, and *seba* in the second.

82. *Geochelone elegans*: Sahoo et al. (2024) genotyped 82 wild *G. elegans* covering the entire range of the species using two mtDNA regions and six polymorphic microsatellites, and added morphometric analysis. Their results revealed two major clades on the Indian subcontinent: a relatively invariable northwestern group from Kutch and Saurashtra in Gujarat and Malwa Plateau and Aravali in Rajasthan, and a highly diverse and disjunct southern group from the Eastern Ghats, Deccan Plateau, Nilgiri Hills, and Coromandel Coast, with a closely related subgroup from Sri Lanka. Their results clarified some of the earlier less robust results from similar studies by Gaur et al. (2006) and Vamberger et al. (2020b) (see TTWG Annotation 21:92), as well as Kundu et al. (2023). The two deeply divergent groups apparently split around 2 mya during the Pleistocene and the authors recommended they be considered separate subspecies, with the southern clade referred to *Geochelone elegans elegans* (Schoepff 1795) and the northwestern clade to *Geochelone elegans stellata* (Schweigger 1812).

We agree that the results of the genetic and morphological analyses demonstrate taxonomic structure that warrants recognition of the two clades, but the proposed name *G. e. stellata* is unfortunately nomenclaturally unavailable for the northwestern group, as it is clearly an objective junior synonym of *G. e. elegans* (see detailed explanation in TTWG 2017; Annotation 17:65). Instead, we note that the taxon name *Testudo actinodes* Bell 1828a is apparently available as the potential name for the northwestern group.

Gray (1870b, 1870c) examined variation within his concept of the species *Peltastes stellatus* (= *G. elegans*), and noted that his "var. *actinoides*" (= *actinodes* Bell 1828a) was distinguished from his other varieties by having 15 or more yellow rays on the carapacial scutes, whereas his "var. *elegans*" had fewer yellow rays (8–9) on the costal scutes. Five syntypes of *T. actinodes* were present in the Oxford University Museum when examined by Nowak-Kemp and Fritz (2010). We have now examined detailed curatorial photos of these five syntype specimens and note that one of them (OUM 8556) has 13 radiating yellow lines on each of the carapacial scutes, with OUM 8555 and 8558 also having 11 and 12 lines, respectively, whereas OUM 8559 and 8560 have

only 7 costal lines each and appear to be similar to animals from the southern group. We also note that Sahoo et al. (2024) depict their northwestern clade animals as having multiple radiating costal lines and their southern group as having significantly fewer lines, and photos we have seen of animals from Gujarat in northwestern India have many more radiating yellow carapacial scute lines than those from southeastern India. On the other hand, specimens of *G. elegans* from Sri Lanka also have multiple radiating lines (de Silva 2003) similar to the ones from the northwestern group.

Although we cannot be certain whether the name *T. actinodes* Bell 1828a actually refers to the northwestern subspecies, or whether that clade may require a new or different name, it is clear that *G. elegans stellata* (Schweigger 1812) is not the correct name for the taxon and should not be used.

To clarify this situation, the syntypes of *T. actinodes* Bell 1828a need to be genotyped to ascertain their provenance, and if found, also the holotypes of *Peltastes stellatus maura* Gray 1870b and *P. stellatus sebae* Gray 1870b, two other nominal taxa synonymized under *Geochelone elegans*. However, in view of the uncertainty of the provenance of these three taxa, and until such time that the uncertainty is hopefully resolved, we make the strong recommendation that the northwestern clade of *Geochelone elegans* needs to be referred to as *G. elegans actinodes* rather than the unavailable objective junior synonym *G. elegans stellata*, and we initiate use of this revised nomenclature here in our checklist, pending further genetic and morphological work.

83. *Indotestudo elongata*: Ihlow et al. (2024) analyzed the phylogeography of *I. elongata* based on three mitochondrial gene fragments (ND4, cytochrome b, and COI) for 52 individuals. They found genetic homogeneity across its wide range, supported by a shallow mean uncorrected p distance of 1.2% in cytochrome b, indicating a surprising lack of biogeographic structuring. In addition, multivariate statistical analyses of morphometric and phenotypic characters of 166 tortoises from across the range showed no statistically significant differences among biogeographic units. Archaeological evidence suggests that *I. elongata* has been used as a food item for millennia. The authors hypothesized that its current wide distribution and the absence of genetic differentiation could be the result of significant range expansion caused by extensive human-mediated trade dispersal over prehistorical and historical time frames.

84. *Pyxis*: Bell (1827) described the genus *Pyxis* with the type species *Pyxis arachnoides* Bell 1827. This generic name continued to be used consistently as valid in the herpetological literature until the mid-20th century. However, Williams (1950) noted that *Pyxis* Bell 1827 (erroneously cited as 1825) was a junior homonym preoccupied by *Pyxis* Chemnitz 1784 (describing a Brachiopoda and a Bivalvia) and he therefore established the new replacement name *Bellemys* Williams 1950a (*nomen novum*). Williams (1952) also used *Bellemys* instead of *Pyxis*, but Loveridge and Williams (1957) reverted to using *Pyxis*, stating that "*Pyxis* Bell 1825 [sic] is used rather than *Bellemys* Williams 1950a, 1952, in as much as the supposedly preoccupying name *Pyxis* Chemnitz 1784 is not binomial, and the next use of the name (Humphreys in the "Museum Calonnianum" 1797) has now been outlawed by ruling of the International Commission on Nomenclature (Decision 51)."

Subsequently, Wermuth and Mertens (1977), apparently following the conclusion by Loveridge and Williams (1957), stated that ICZN (1956) in their Direction 32, which included Opinion 51 (= "Decision" 51), had labeled the earlier Chemnitz (1784) name a "*nomen illegitimum*." This statement was uncritically repeated by Bour (1981) and Fritz and Havaš (2007). As a result, the name *Pyxis* Bell 1827 has since 1957 continued to be consistently used as valid and has not been considered a junior homonym.

However, the assertions regarding the unavailability of *Pyxis* Chemnitz 1784 by Loveridge and Williams (1957) and Wermuth and Mertens (1977) were erroneous, and neither ICZN Opinion 51 nor Direction 32 (ICZN 1956) addressed the Chemnitz (1784) publication, or rejected the publication itself or Chemnitz' name *Pyxis*. Instead, ICZN (1956) rejected for nomenclatural purposes the "Museum Calonnianum" by Humphreys (1797) in its Opinion 51. Unfortunately, despite the assertion as such by Loveridge and Williams (1957), Humphreys (1797) did not use or refer to the generic name *Pyxis*, although three species-level Chemnitz names were used.

In addition, ICZN (1944) in their Opinion 184 had previously ruled that all species-level names in the several volumes of the Martini and Chemnitz "Conchylien-Cabinet" publications (including Chemnitz 1784) were unavailable, since they did not use binomial nomenclature consistently; however, they also noted that the included generic-level names were potentially available. Subsequently, ICZN (1954), in their Direction 1, affirmed their earlier Opinion 184, rejecting all species-level names from Chemnitz (1784) as unavailable, but pointedly did not make a decision regarding the availability of generic-level names from that publication, referring to ICZN case File Z.N.(S.) 800 that left the question open. In doing so, it would appear that *Pyxis* Chemnitz 1784 could possibly still be available as a senior homonym, and that *Bellemys* Williams 1950a could potentially still be a valid replacement name for *Pyxis* Bell 1827.

However, ICZN case File Z.N.(S.) 800 was closed in 1985 (F. Welter-Schultes, ICZN, pers. comm.) and two years later ICZN (1987) officially listed the Martini and Chemnitz volumes as totally rejected works due to their failure to use binomial nomenclature, this time also affecting generic names since the work did not meet the requirements of ICZN Article 11.4, instituted in 1961. As a result, *Pyxis* Chemnitz 1784 clearly became an unavailable name and not valid for the purposes of priority or homonymy, and *Bellemys* Williams 1950a was no longer needed as a replacement name for *Pyxis* Bell 1827.

85. *Testudo (Testudo) graeca buxtoni* and *T. (T.) g. zarudnyi*: Ranjbar et al. (2022) combined phylogenetic (mitochondrial DNA) and morphometric analyses to explore the contact zone between the two subspecies of *Testudo graeca* (*T. g. buxtoni* and *T. g. zarudnyi*) in central Iran. Each subspecies was identified as a monophyletic clade, but with some geographic intergradation. In addition, they identified four subclades within *T. g. buxtoni*, and even suggested that those might be worthy of subspecific recognition. However, their morphometric analysis showed significant geographic variation that did not match the genetic variation, and they made no formal taxonomic changes. The analysis suggested that the two subspecies diverged from one another after the uplifting of the Zagros Mountains during the early Pliocene.

86. *Testudo (Testudo) graeca tripolitania*: Poterala et al. (2024) described a new subspecies of the Mediterranean Spur-thighed Tortoise, *Testudo graeca tripolitania*, from northwestern Libya, separating it from *T. graeca nabeulensis*, redefined as occurring only in Tunisia and eastern Algeria. They used mitochondrial DNA (mtDNA) sequencing, and found that several specimens imported to the U.S. from Libya and specimens sampled in the wild in northwestern Libya shared the same haplotype as the distinctive "lineage E" haplotype previously documented by Graciá et al. (2017a), but at that time of unknown provenance. They also noted that the earlier described taxon, *Testudo flavominimaralis* Highfield and Martin 1989c, although possibly a senior synonym, had been

declared a *nomen dubium* by Pieh and Perälä (2002) and that its low carapace height ratio was inconsistent with their new higher-domed subspecies. In addition, the holotype of *T. flavominimaralis* was the only described and curated specimen of that taxon, and is now apparently lost and unavailable for genotyping. Unless the original type material of *T. flavominimaralis* is located and genotyped, we agree that it represents a *nomen dubium* and that *T. g. tripolitania* appears to be a valid and distinct taxon.

87. *Testudo (Chersine) hermanni*: Chiocchio et al. (2022) analyzed the phylogeography of *T. hermanni* in Italy, genotyping 174 individuals at 7 microsatellite loci. They identified three main genetic clusters, including a fairly small hotspot of intraspecific diversity in the Calabrian region of southern Italy. Another relatively small distinct cluster was also found on Sicily, with a more extensive bicoastal cluster along the east and west coastal regions of the more northern portions of the peninsula. This larger cluster also demonstrated areas of admixture with the southern cluster. They made no taxonomic recommendations, but urged the need for conservation actions to secure the long-term persistence of viable tortoise populations, notably in the southern cluster.

88. *Carettochelys insculpta*: Young et al. (2025) analyzed and reconstructed phylogenetic relationships and patterns of genetic diversity of *Carettochelys insculpta* from Australia and New Guinea using a genome-wide dataset of 15,081 single nucleotide polymorphisms (SNPs) and two mitochondrial loci from 251 rangewide specimens. Their analysis revealed that genetic drift coupled with low population sizes in Australia compared with New Guinea has resulted in reduced genetic diversity of Australian populations. They also observed unexpectedly high levels of genetic structuring both among the Australian populations and between Australian and New Guinean populations, with three evolutionarily significant units (ESUs) delineated as having diverged during the Pleistocene: northern Australia, southern Indonesian Papua, and southern Papua New Guinea. However, without an accompanying morphological analysis, they were unwilling to recommend any changes to the current taxonomy. They regarded the distinct lineages that they identified as lineages within a single widespread species subject to a complex history of isolation and interconnection in the Australo-Papuan region reflective of sea level changes and an early vicariance event that isolated the New Guinea and Australian populations. Dispersal along paleodrainages of the submerged continental shelf since that event has been limited by paleogeographic barriers to dispersal despite the dramatic sea level changes during the glacial cycles.

89. Trionychinae: Chen et al. (2025) analyzed the complete mitochondrial genomes of *Palea steindachneri* and *Pelodiscus axenaria*, and compared them to all other genera of the subfamily Trionychinae. The phylogeny they generated matched those previously found for mitochondrial and nuclear DNA by Engstrom et al. (2004), Kundu et al. (2018c), and Thomson et al. (2021) for the Trionychinae, and further corroborated their findings of three monophyletic clades (essentially infrafamilies) consisting of: 1) the Asian genera *Amyda*, *Dogania*, *Nilssonia*, *Palea*, and *Pelodiscus*, 2) the genera *Chitra*, *Pelochelys*, and *Trionyx*, and 3) *Apalone* and *Rafetus*. Engstrom et al. (2004) had previously coined the PhyloCode names Amydona, Gigantaestuarochelys, and Apalonia, respectively, for these three well-defined clades.

90. *Amyda ornata jongli*: Praschag and Gemel (2022) described this new subspecies of Asian softshell turtle from Bangladesh, India, and western Myanmar based on morphological differences from *A. o. phayrei* from central and eastern Myanmar and western and central Thailand, and *A. o. ornata* from the greater Mekong region of Cambodia, Laos, and Vietnam. The new subspecies was previously identified as an unnamed taxon of *A. ornata* sensu lato based on deep genetic differences from other lineages of *A. ornata* as elucidated by Fritz et al. (2014). We accept this new subspecies.

91. *Pelodiscus axenaria*, *P. shipian*, and *P. huike*: Gong et al. (2022) described *Pelodiscus shipian* as a new species of small Chinese softshell turtle of the *P. axenaria* complex. They found that it was most closely related to *P. huangshanensis*, but differentiated by its distinct mitochondrial haplotype, small body size, and keeled carapace. It was also distinct from their definition of *P. axenaria*, a poorly known species for which they also designated a neotype in order to define that taxon better genetically, since the original syntype specimens could not be located. Specimens of what they described as the new species *P. shipian* had previously been noted to be genetically distinct by Gong et al. (2021), but without a morphological or taxonomic description at that time.

The same taxon (*P. shipian*) was subsequently also genetically analyzed and provisionally described as the new species *Pelodiscus jiangxiensis* by Hou et al. (2023) in a publicly distributed online preprint that was never actually published. The authors commendably withdrew their paper before publication when they became aware of the prior description of the same taxon by Gong et al. (2022). We agreed with these findings and planned to accept *P. shipian* as a new and distinct species of the *P. axenaria* complex, and also to place *P. jiangxiensis* as a *nomen nudum* in its synonymy.

However, Liu et al. (2024) then subsequently claimed to have rediscovered the lost type material of *P. axenaria*. They analyzed sequence variation in the mitochondrial cytochrome b gene of three specimens of the purported original syntypes of *P. axenaria* together with previous data, and these sequences clustered in the clade of *P. shipian* as defined by Gong et al. (2022). Therefore, Liu et al. (2024) stated that the work by Gong et al. (2022) had apparently confused the species involved, such that specimens of their new taxon, *P. shipian*, were genetically identical to the newly located purported syntypes of *P. axenaria*, and the specimens of what Gong et al. had identified as *P. axenaria* were actually a new and unnamed taxon.

As a result, Liu et al. (2024) then described this allegedly new taxon as *Pelodiscus huike*, and placed both *P. shipian* and *P. jiangxiensis* in the synonymy of *P. axenaria*. They also designated a lectotype from the purported original syntypes of *P. axenaria*, in order to set aside and invalidate the neotype designated by Gong et al. (2022). Furthermore, they suggested that the neotype of *P. axenaria* that Gong et al. (2022) had designated was actually a specimen of their new species, *P. huike*, and instead designated it as the holotype of that new species while setting it aside as the neotype of *P. axenaria*.

However, Gong et al. (2025) then examined and analyzed photographs of the specimens of *P. axenaria* that Liu et al. (2024) had suggested represented original syntypes, and the one they had designated as lectotype. Gong et al. found that it was not the same animal as the figured specimen from the original description by Zhou et al. (1991), nor was it the same animal that had previously been labeled as the supposed holotype of *P. axenaria* by Zhou and Li (2013), which also did not correspond to the original figure. The characteristic plastral markings were different and there were notable inconsistencies in the labeling of the specimens. In addition, in view of the challenges of sequencing DNA from formalin-preserved specimens, Gong et al. (2025) also questioned whether the DNA sequences allegedly obtained from the long-term (30-year) formalin-preserved specimens identified by Liu et al. (2024) as types of *P. axenaria* were accurate, because the sequences were evidently obtained using standard Sanger sequencing approaches.

Gong et al. (2025) concluded that neither the specimens nor the DNA sequences examined by Liu et al. (2024) could be equated with any actual type material of *P. axenaria*, and since their holotype of *P. huike* was the same specimen as the previously designated neotype of *P. axenaria*, Gong et al. therefore noted the name *P. huike* to be an objective junior synonym of *P. axenaria*. We agree with this interpretation, and therefore recognize *P. shipian* as an accepted species, and *P. huike* as a synonym of *P. axenaria*, while also noting that the taxonomic relationships and geographic distributions of these and other morphologically challenging *Pelodiscus* species have been most difficult to elucidate and unfortunately still remain unclear.

92. ***Pelodiscus sinensis* and *P. variegatus*:** Liu et al. (2024) analyzed variation in sequences of the mitochondrial cytochrome b gene of all *Pelodiscus* species, focusing on the *P. axenaria* complex (see Annotation 91 above). One of their ancillary findings was that specimens of *P. sinensis* and *P. variegatus* had very low uncorrected p distances between them, indicating very close relatedness. They suggested that their divergence might only be at the subspecific level, but did not make any specific taxonomic or nomenclatural recommendations. However, Gong et al. (2018) had previously reported this mitochondrial DNA similarity between these taxa, but found what was later named *P. variegatus* distinct in the studied nuclear DNA sequences. The subsequent description of *P. variegatus* concluded that this species has introgressed mitochondria (Farkas et al. 2019), explaining the apparent similarity.

93. ***Trionyx*:** Both Bour et al. (1995) and Frétey (2024) noted that in addition to the accepted designation by Bory de Saint-Vincent (1828:77) of *Trionyx egyptiacus* (= *Trionyx triunguis*) as the type species for *Trionyx*, prior to that, Bell (1828c:516) had designated *Trionyx coromandelicus* (= *Lissemys punctata punctata*) as the type species of *Trionyx*, and that this overlooked earlier designation might threaten the continued use of the name *Trionyx* for *T. triunguis*. Both suggested that the ICZN should be petitioned to use their plenary powers to suppress the type designation by Bell (1828c). We would be in support of such a petition if it was felt to actually be needed.

94. ***Astrochelys rogerbouri*:** Kehlmaier et al. (2023) genotyped several fossils of extinct Western Indian Ocean giant tortoises, including *Aldabrachelys abrupta* and *A. grandidieri* from Madagascar, and discovered that a single tibia previously thought to represent a smaller subadult specimen of *A. abrupta* was actually deeply genetically divergent in both mitochondrial and nuclear DNA, and described it as an extinct large tortoise, *Astrochelys rogerbouri*, sister to extant *A. yniphora*.

95. ***Chelonoidis dominicensis, C. gersoni,* and *C. marcanoi*:** Based on new subfossil material, Viñola-López and Almonte (2022) considered *C. dominicensis* to be conspecific with and a junior subjective synonym of *C. marcanoi* and present on both the northern and southern Hispaniola paleo-islands. They described *C. gersoni* as a new species from the southern paleo-island. However, Albury et al. (pers. comm., *in litt.*), consider *C. dominicensis* and *C. gersoni* to both be valid species, and the holotype of *C. marcanoi* to be unidentifiable and a *nomen dubium*. Pending the analysis of more material, we tentatively agree, but continue to list the species as potentially distinct.

96. ***Solitudo* and *S. sicula*:** Valenti et al. (2022) described this new genus and species of large extinct Mediterranean tortoise from Sicily from the Late Pleistocene and possibly the earliest Holocene. They suggested that it belongs to a previously unrecognized clade that includes other large-sized tortoises from Mediterranean islands, such as Malta and Menorca. A review of the pertinent taxa indicated that the remains described represented the geologically youngest large-sized tortoise of the Mediterranean region.

PREVIOUS CHECKLIST ANNOTATIONS

2007 Checklist Annotations

TTWG 2007b (CRM 4:173–199) [07:1-105]

07:1. Both IUCN (The World Conservation Union, http://www.iucnredlist.org) and CREO (Committee on Recently Extinct Organisms, http://creo.amnh.org) have designated 1500 AD as their official cutoff date for determining what constitutes a recently extinct species, and we follow their criteria in our checklist.

07:2. ***Chelydra***: Phillips et al. (1996) elevated *acutirostris* and *rossignoni* to full species status and retained the subspecies *osceola*. See Shaffer et al. (in press) for a complete review.

07:3. ***Macrochelys* [formerly *Macroclemys*]**: Although *Macroclemys* has been the most commonly used name, Webb (1995b) showed that *Macrochelys* is the oldest available name.

07:4. ***Chelonia mydas***: Bowen et al. (1992) showed that recognition of the taxon *agassizii* Bocourt 1868 renders *mydas* paraphyletic, and *agassizii* is no longer generally recognized as either a distinct species or subspecies. See Parham and Zug (1996) and Karl and Bowen (1999) for a complete review.

07:5. ***Eretmochelys imbricata***: Fritz and Havas (2006, 2007) did not list *bissa* as a valid taxon, but no argumentation for this opinion was given. Genetic data (Okayama et al. 1999) have suggested significant separation of Atlantic from Pacific stocks.

07:6. ***Kinosternon* species**: Serb et al. (2001) elevated two former subspecies of *flavescens* (*arizonense* and *durangoense*) to full species status.

07:7. ***Kinosternon chimalhuaca***: This new species name appeared prematurely and erroneously first in the hobbyist literature, with the full original description published a few months later (Berry et al. 1996, 1997).

07:8. ***Kinosternon scorpioides scorpioides***: Includes the previously recognized subspecies *seriei* Freiberg 1936 and *carajasensis* Cunha 1970 in synonymy (Cabrera and Colantonio 1997).

07:9. ***Sternotherus***: This genus was included as a junior synonym of *Kinosternon* by Iverson (1992) and David (1994) based on work by Seidel et al. (1986) and Iverson (1991). However, this view was never widely accepted, and Iverson (1998) showed that the species referred to either *Sternotherus* or *Kinosternon* formed reciprocally monophyletic clades and recommended that both genera be used.

07:10. ***Sternotherus depressus***: Whereas some earlier authors had placed this taxon as a subspecies of *minor*, Walker et al. (1998) showed that *depressus* was genetically distinct from *minor*.

07:11. ***Chrysemys picta dorsalis***: This subspecies of *Chrysemys picta* was elevated to full species status by Starkey et al. (2003), who recognized two distinct genetic lineages: *C. dorsalis* and *C. picta*. They did not find genetic support for the other subspecies of *C. picta* (*belli, marginata*) but did not recommend that they be abandoned. Fritz and Havas (2006, 2007) argued that full specific status of *dorsalis* was not fully demonstrated and retained it and the other two taxa as subspecies of *C. picta*, agreeing also with Ernst et al. (2006a).

07:12. ***Graptemys ouachitensis sabinensis***: Based on molecular and morphologic data, Stephens and Wiens (2003) suggested that *sabinensis* may not be closely related to *ouachitensis*. However, statistical support for this was weak, and they did not discuss or recommend a taxonomic change. Further study of this complex may warrant the elevation of the sympatric taxon *sabinensis* to full species status.

07:13. ***Pseudemys concinna concinna***: Includes the previously recognized subspecies *hieroglyphica* Holbrook 1836, *mobilensis* Holbrook 1838, and *metteri* Ward 1984 in synonymy (Seidel 1994).

07:14. ***Pseudemys concinna floridana***: This taxon was previously considered a separate species, but was designated a subspecies of *concinna* by Seidel (1994). Jackson (1995) argued for the retention of *floridana* as a full species, but Seidel (1995) rejected this argument.

07:15. ***Pseudemys concinna suwanniensis***: Previously considered a subspecies of *concinna*, Seidel (1994) argued for the elevation of this taxon to full species status, but Jackson (1995) argued for its subspecific status.

07:16. ***Pseudemys gorzugi***: This taxon was previously considered a subspecies of *concinna*, but was elevated to species status by Ernst (1990) without argumentation, but then supported through analysis by Seidel (1994).

07:17. ***Pseudemys peninsularis***: This taxon was previously considered a subspecies of *floridana*, but was elevated to species status by Seidel (1994). Jackson (1995) argued for the retention of *peninsularis* as a subspecies of *floridana*, but Seidel (1995) reaffirmed his recognition.

07:18. ***Trachemys* species**: Seidel (2002) recommended elevating nine Mesoamerican taxa, previously recognized as subspecies of *Trachemys scripta*, to species rank.

07:19. ***Trachemys* subspecies**: Seidel (2002) also recommended reassigning five taxa, previously subspecies of *scripta*, to subspecies of his various elevated *Trachemys* species.

07:20. ***Trachemys dorbigni***: Includes the previously recognized subspecies *brasiliensis* Freiberg 1969 in synonymy, based on morphologic work (del Barco and Larriera 1993).

07:21. ***Emydoidea* and the turtles formerly known as *Clemmys***: The four traditional species of *Clemmys* (*guttata* [type], *insculpta*, *muhlenbergii*, and *marmorata*) do not form a monophyletic group with respect to the two monotypic genera *Emys orbicularis* and *Emydoidea blandingii* in phylogenies based on DNA data (Bickham et al. 1996; Burke et al. 1996; Lenk et al. 1999; Feldman and Parham 2002). While there is a general agreement that *insculpta* and *muhlenbergii* are sister-species and should be placed in the genus *Glyptemys* (Holman and Fritz 2001; Parham and Feldman 2002), there are two schemes presented for *marmorata* and *blandingii*. Holman and Fritz (2001) recommended that *marmorata* be placed in the monotypic genus *Actinemys*, retaining both *Emys orbicularis* and *Emydoidea blandingii* as additional monotypic genera. Other authors (Bickham et al. 1996; Feldman and Parham 2002; Parham and Feldman 2002) recommended that *marmorata* and *blandingii* be placed into an expanded *Emys*, a scheme favored in the most recent analysis of variation in *marmorata* (Spinks and Shaffer 2005).

07:22. ***Emys* or *Actinemys marmorata***: Previously, two subspecies were distinguished, including *pallida* Seeliger 1945, but genetic analysis by Spinks and Shaffer (2005) demonstrated that the typical and previously recognized subspecies *pallida* were within the same phylogeographic clade and so *pallida* should not be considered valid.

07:23. ***Emys orbicularis iberica***: Includes the recently described subspecies *kurae* Fritz 1994 in synonymy (Fritz 1998).

07:24. ***Emys orbicularis persica***: Includes the recently described subspecies *orientalis* Fritz 1994 in synonymy (Fritz 1998).

07:25. **Mexican *Terrapene carolina***: Stephens and Wiens (2003) suggested that Mexican subspecies of *T. carolina* may warrant full species status. While this convention has also been adopted previously (Smith et al., 1996), almost all other workers recognize these as subspecies.

07:26. **Platysternidae**: Krenz et al. (2005) confirmed that nuDNA placed *Platysternon* solidly within the Testudinoidea, and Parham et al. (2006a) supported this finding with mtDNA.

07:27. ***Platysternon megacephalum***: Ernst and Laemmerzahl (2002) synonymized two subspecies of *megacephalum* (*vogeli* Wermuth 1969 and *tristernalis* Schleich and Gruber 1984) with the nominate subspecies.

07:28. **Testudinoidae or Testuguria**: Shaffer et al. (1997) coined the name 'Testudinoidae' for the clade that united Testudinidae with Bataguridae/Geoemydidae. Joyce et al. (2004) listed Testudinoidae as an undesirable derivative of *Testudo* being to similar to both 'Testudinidae' and 'Testudinoidea.' In that same paper, the authors coined the new clade name 'Testuguria' for that same clade (while neglecting to list Testudinoidae as an objective senior synonym). Parham et al. (2006a) explicitly argued for the use of Testuguria over Testudinoidae.

07:29. **Bataguridae or Geoemydidae**: Both names are being used to refer to this group of predominantly Asian testudinoids. McDowell (1964) used the name Batagurinae for this group (as a subfamily) which was changed to Bataguridae (as a family) by Gaffney and Meylan (1988). Bour and Dubois (1986) showed that Geoemydidae has priority, and David (1994), Spinks et al. (2004) and others have embraced this view. However, this approach was questioned by Joyce et al. (2004) who, working in a rank-free phylogenetic taxonomy framework, recommended the continued use of Bataguridae. In the interest of reconciling phylogenetic nomenclature with traditional Linnaean rules of priority, Parham et al. (2006a) endorsed a phylogenetic codification of Geoemydidae.

07:30. ***Batagur***: Praschag et al. (2007b) and Le et al. (2007) demonstrated that species of *Kachuga* were genetically paraphyletic with respect to those referred to *Batagur* and *Callagur* and recommended that only one genus be recognized, and the name *Batagur* has priority.

07:31. ***Batagur baska***: The subspecies *ranongensis* Nutaphand 1979 is not well differentiated and has been synonymized under *baska* by Fritz and Havas (2006, 2007), but no specific morphologic or genetic analysis has yet been performed to formally evaluate the status of this taxon.

07:32. ***Cuora***: Phylogenies based on DNA data (Honda et al. 2002a; Stuart and Parham 2004; Parham et al. 2004; Spinks et al. 2004) have shown that continued recognition of the genus *Pyxidea* for *mouhotii* would render *Cuora* paraphyletic. All of these studies recommended expanding *Cuora* to include *mouhotii*. Other schemes for *Cuora* have not been published in the recent scientific literature, though there has been some use of *Cistoclemmys* for *flavomarginata* and *galbinifrons* (e.g., Zhao et al. 1997; Zhao 1997; Yasukawa and Ota 1999).

07:33. **Hybrid species**: The validity of six taxa of *Cuora*, *Mauremys* [including *Ocadia*], and *Sacalia* recently described from pet trade specimens has been refuted by genetic studies that have shown them to be based on hybrids (Parham et al. 2001; Wink et al. 2001; Spinks et al. 2004; Stuart and Parham 2004, 2007). The taxa shown to be hybrids are: *Cuora galbinifrons serrata* Iverson and McCord 1992b, *Mauremys iversoni* Pritchard and McCord 1991, *Mauremys pritchardi* McCord 1998, *Ocadia glyphistoma* McCord and Iverson 1994, *Ocadia philippeni* McCord and Iverson 1992, and *Sacalia pseudocellata* Iverson and McCord 1992a.

07:34. ***Cuora flavomarginata sinensis***: Some authors recognize this taxon as a valid subspecies (McCord and Iverson 1991; Fong et al. 2002) while others synonymize it with *flavomarginata* (Yasukawa and Ota 1999; Fritz and Havas 2006, 2007).

07:35. ***Cuora galbinifrons***: The taxa *bourreti* and *picturata*, originally described as subspecies of *Cuora galbinifrons*, were elevated to species rank by Stuart and Parham (2004) based on concordance of morphological with molecular differentiation. Fritz et al. (2006c) returned *bourreti* to subspecies rank based on osteological characters shown by market specimens, and suggested that *picturata* warrants the same ranking; Fritz and Havas (2006, 2007) subsequently listed *picturata* at subspecies rank based on morphologically intermediate pet trade specimens. Includes the previously recognized *hainanensis* Li 1958 in synonymy (Zong and Pan 1989; Iverson and McCord 1992b).

07:36. ***Cuora trifasciata***: Blanck et al. (2006a) recommended that *Cuora trifasciata* be split into two species (including their newly named species *cyclornata* and its new subspecies *meieri*) based on paraphyletic mtDNA haplotypes and morphological differences. Spinks and Shaffer (2007) showed that *trifasciata* as traditionally recognized is monophyletic based on nuDNA and therefore recommended that *cyclornata* should not be recognized, pending additional study.

07:37. ***Cuora yunnanensis***: This species has been listed as extinct by the IUCN since 2000 (www.iucnredlist.org), based on several decades of not finding any surviving animals despite intensive searches. Recently, a pair of animals representing this species were found in markets (Zhou and Zhao 2004; Zhou 2005), with subsequent confirmation through genetic analysis (He et al. 2007).

07:38. ***Cyclemys***: Iverson (1992) recognized two taxa of *Cyclemys* (*dentata* and *tcheponensis*). Later, *atripons* and *pulchristriata* were described and *oldhamii* was resurrected (Iverson and McCord 1997; Fritz et al. 1997). Genetic analysis by Guicking et al. (2002) also supported the validity of *shanensis*.

07: 39. ***Geoemyda***: Yasukawa et al. (1992) elevated *japonica* to species status (previously considered a subspecies of *spengleri*).

07:40. ***Hardella thurjii***: Praschag et al. (2007b) found no genetic or morphologic evidence for continued recognition of the subspecies *indi* Gray 1870c, and synonymized it under *thurjii*.

07:41. ***Heosemys annandalii* [formerly in *Hieremys*]**: Spinks et al. (2004) showed that *annandalii* was nested among species of *Heosemys*. Diesmos et al. (2005) formally moved *annandalii* into *Heosemys*.

07:42. ***Leucocephalon yuwonoi* [formerly in *Geoemyda* or *Heosemys*]**: Originally described as a species of *Geoemyda* (McCord et al. 1995), Fritz and Obst (1996) placed *yuwonoi* in *Heosemys*. McCord et al. (2000) showed that *yuwonoi* was not closely related to the type species of *Geoemyda* or *Heosemys*, but instead sister to *Notochelys platynota*, and erected a new genus, *Leucocephalon*, for *yuwonoi*.

07:43. ***Malayemys macrocephala***: Brophy (2004) proposed the recognition of this species as distinct from *subtrijuga* based on morphological grounds.

07:44. ***Mauremys* [including species formerly in *Annamemys*, *Chinemys*, or *Ocadia*]**: Iverson and McCord (1994) included *annamensis* under an expanded *Mauremys*. Subsequent phylogenies based on DNA data (Honda et al. 2002b; Barth et al. 2004; Feldman and Parham 2004; Spinks et al. 2004) showed that the genera *Ocadia* and *Chinemys* rendered *Mauremys* paraphyletic. Based on these results, some authors (Feldman and Parham 2004; Spinks et al. 2004) recommended synonymizing *Ocadia* and *Chinemys* under *Mauremys*. Barth et al. (2004) presented this same scheme as well as one that would retain

Chinemys and *Ocadia* and further divide *Mauremys* into the genera *Cathaiemys* and *Emmenia*. Barth et al. (2004) did not favor one scheme over the other and a competing scheme for *Mauremys* has not been formally proposed in the scientific literature.

07:45. ***Mauremys leprosa***: Fritz et al. (2006a) explicitly synonymized several subspecies of *leprosa* recently described by Schleich (1996a) (*atlantica, erhardi, marokkensis, wernerkaestlei*, and *zizi*) plus *vanmeerhaeghei* Bour and Maran 1998 [1999], and only recognized *leprosa* and *saharica*.

07:46. ***Mauremys reevesii***: Iverson et al. (1989) and Barth et al. (2003, 2004) refuted the validity of the terminal taxon *megalocephala* Fang 1934, but it has continued to be recognized by Chinese researchers (Guo et al. 1997; Zhao 1997; Zhang et al. 1998), and Fritz and Havas (2006, 2007) listed it as a separate taxon with speculation about its relationships.

07:47. ***Melanochelys trijuga edeniana***: The subspecies *wiroti* Reimann 1979 was recognized by Iverson (1992), but David (1994) suggested that it was synonymous with *edeniana*, and Fritz and Havas (2006, 2007) followed this arrangement.

07:48. ***Pangshura* [formerly in *Kachuga*]**: Das (2001) and Schleich and Kästle (2002) used the name *Pangshura* to refer to small-bodied *Kachuga*. A phylogeny based on DNA data (Spinks et al. 2004) showed that *Kachuga* was paraphyletic and so removed *flaviventer, smithii, sylhetensis, tecta*, and *tentoria* into the genus *Pangshura*. Praschag et al. (2007b) using mtDNA confirmed the well-supported monophyly of *Pangshura*.

07:49. ***Pangshura tentoria flaviventer***: Schleich and Kästle (2002) elevated *flaviventer* to full species status based on sympatry with *circumdata*, but Praschag et al. (2007b) performed a phylogeographic analysis and retained *flaviventer* as a subspecies of *tentoria*.

07:50. ***Siebenrockiella leytensis* [formerly in *Heosemys*]**: Diesmos et al. (2005) placed *leytensis* into the genus *Siebenrockiella* based on strong genetic evidence for its sister relationship to *S. crassicollis*.

07:51. ***Vijayachelys silvatica* [formerly in *Geoemyda*]**: This species was originally named as a species of *Geoemyda*. However, a molecular study by Praschag et al. (2006) suggested a distant relationship with that genus and they recommended that it be placed in the new monotypic genus *Vijayachelys*.

07:52. **The *Geochelone* complex**: This generic complex includes the genera *Geochelone, Aldabrachelys, Astrochelys, Angonoka, Centrochelys, Chelonoidis, Dipsochelys*, and *Stigmochelys*. Lapparent de Broin (2000b), Gerlach (2001, 2004), Le et al. (2006), and Fritz and Bininda-Emonds (2007) recommended dividing the *Geochelone* complex into several genera, although their schemes differ somewhat. A general consensus on a generic-level revision for some members of the group is lacking while in other areas (e.g., *Astrochelys radiata*, *Chelonoidis*) there is agreement.

07:53. ***Aldabrachelys* or *Dipsochelys***: Bour (1982a) originally recommended that Aldabran tortoises (*dussumieri* or *gigantea*) be placed in the genus *Dipsochelys* instead of *Aldabrachelys*. However, *Aldabrachelys* is still widely used, including sometimes by Bour (Austin et al. 2003), though *Dipsochelys* is favored by others (Palkovacs et al. 2002, 2003; Gerlach 2004). There is recent disagreement regarding the type specimen of *Testudo gigantea*, the type species of *Aldabrachelys*, that was presumed lost. Frazier (2006) designated a neotype for *T. gigantea*, an act that would seemingly validate the use of both *Aldabrachelys* and the terminal taxon *gigantea*. Around the same time, Bour (2006c) rediscovered the original lost type specimen, which is actually an individual of the South American tortoise *Chelonoidis denticulata*. If this claim is correct, then the names *Aldabrachelys* or *gigantea* might not be applicable to Aldabran tortoises. Whether Frazier's neotype designation or Bour's specimen rediscovery prevails nomenclaturally remains a matter of ongoing debate, but since Bour (2006c) was the most recently published authority we use the name *dussumieri* rather than *gigantea* in our list.

07:54. ***Aldabrachelys* or *Dipsochelys* species**: Gerlach and Canning (1998) recognized six species of tortoises in Aldabra, Madagascar, and the Seychelles (three of which were extinct: *abrupta, daudinii*, and *grandidieri*). The two species from Madagascar became extinct prior to modern times (*abrupta* Grandidier 1868 in ca. 1250 AD and *grandidieri* Vaillant 1885b in ca. 950 AD) so we do not include them in our list of modern taxa. Palkovacs et al. (2002, 2003) questioned the validity of multiple extant species based on their analysis of genetic data, recognizing only a single living taxon (*Dipsochelys dussumieri*). Gerlach and Bour (2003) reemphasized the validity of the extant species based on the observation that the hatchlings are diagnostic. Fritz and Havas (2006, 2007) recognized only one extant species of Indian Ocean giant tortoise which they referred to *Aldabrachelys gigantea*, but did not address the findings of Gerlach and Bour (2003) or Bour (2006c). As we consider the issues surrounding the validity of these species as remaining unresolved, we list all these species as potentially valid.

07:55. ***Aldabrachelys* or *Dipsochelys dussumieri***: Iverson (1992) listed this species as *Geochelone gigantea* Schweigger 1812. Many authors now use *dussumieri* for the Aldabra tortoise (see above), but others persist in using the older name *gigantea* (e.g., Fritz and Havas 2006, 2007), and others have used the name *elephantina* Duméril and Bibron 1835 (David 1994; Devaux 2007).

07:56. ***Astrochelys* or *Angonoka yniphora***: Le et al. (2006) named *Angonoka* for *yniphora* because of its uncertain phylogenetic position. Fritz and Bininda-Emonds (2007) recovered a weak sister relationship between *yniphora* and *Astrochelys radiata* under some algorithms and recommended that *yniphora* be placed in *Astrochelys*.

07:57. ***Chelonoidis petersi***: According to Cabrera (1998), citing morphologic and osteologic work by Fernández (1988), *Chelonoidis chilensis* should be divided into two species, *chilensis* and *petersi* Freiberg 1973, but he considered the taxon *donosobarrosi* Freiberg 1973 to be synonymous with *chilensis*. Fritz and Havas (2006, 2007) speculated that *petersi* may not be valid and synonymized it under *chilensis*, citing phenotypic plasticity in other tortoise species as a reason for not accepting the reported differences between *petersi* and *chilensis*.

07:58. ***Chelonoidis nigra***: Most recent authors have considered the various taxa of Galapagos tortoises as subspecies of *nigra* (e.g., Pritchard 1996; Caccone et al. 1999; Fritz and Havas 2006, 2007), but Caccone et al. (2002) and Russello et al. (2005, 2007) treated them as distinct species. The nomenclatural and survival status of these taxa were discussed in detail by Pritchard (1996).

07:59. ***Chelonoidis nigra chathamensis***: This taxon described from western Chatham Island (San Cristóbal) appears to have been extirpated from its original range, but a population of tortoises persists on eastern Chatham Island that was considered a possible separate subspecies by Pritchard (1996). Pending genetic analysis and resolution of this issue we continue to list *chathamensis* as the extant taxon from Chatham, whereas Fritz and Havas (2006, 2007) listed it as extinct, but made no mention of the extant population.

07:60. ***Chelonoidis nigra duncanensis***: This taxon from Duncan Island (Pinzón) was historically usually referred to *ephippium* Günther 1875a, but Pritchard (1996) demonstrated that *ephippium* was a synonym of *abingdonii* and therefore resurrected the old *nomen nudum* name *duncanensis* Garman 1917.

07:61. ***Chelonoidis nigra nigra***: The nominotypical subspecies *nigra* from Charles Island (Santa Maria or Floreana) is considered to be extinct and is therefore included separately on this list.

07:62. ***Chelonoidis nigra phantastica***: This taxon was listed by Fritz and Havas (2006, 2007) as extant, but Pritchard (1996) considered it probably extinct.

07:63. ***Chelonoidis nigra porteri***: This taxon from Indefatigable Island (Santa Cruz) has often been referred to *nigrita* Duméril and Bibron 1835, but most recent authors, including Pritchard (1996) and Fritz and Havas (2006, 2007) have used *porteri*.

07:64. ***Chelonoidis nigra vicina***: This widespread taxon from Albemarle Island (Isabela) was previously recognized as one of several valid taxa on that island, including *becki* Rothschild 1901, *microphyes* Günther 1875a, *guentheri* Baur 1889, and *vandenburghi* De Sola 1930. Pritchard (1996) synonymized *microphyes, guentheri*, and *vandenburghi* under *vicina*, and recognized only *vicina* and *becki* from Albemarle.

07:65. ***Cylindraspis indica***: Includes the recently described *borbonica* Bour 1978 in synonymy, based on genetic work by Austin and Arnold (2001).

07:66. ***Cylindraspis vosmaeri***: Fritz and Havas (2006) credited Fitzinger 1826 with authorship of this name, but corrected it to Suckow 1798 in their 2007 checklist.

07:67. ***Homopus***: A separate taxon of *Homopus* was referred to *H. bergeri* Lindholm 1906 by Branch (1989). However, that name was a junior synonym of *Psammobates tentorius verroxii* Smith 1839b (Branch 1992; Boycott and Bourquin 2000), and the new taxon was recently described as *H. solus* by Branch (2007).

07:68. ***Indotestudo travancorica***: This taxon was previously considered a subspecies of *forstenii* (Hoogmoed and Crumly 1984; Iverson 1992), but was resurrected to species status by Pritchard (2000) based on morphology, a conclusion supported by mtDNA analysis by Iverson et al. (2001c).

07:69. ***Kinixys belliana***: Fritz and Havas (2006, 2007) recognized only *belliana* and *nogueyi*, following Broadley (1993) uncritically, but others (Iverson 1992; David 1994; Iverson et al. 2001a) also recognized *domerguei* and *zombensis*. As the phylogeography of this broadly distributed species complex

has not been analyzed, we list the four most widely recognized subspecies.

07:70. ***Pyxis arachnoides***: The three recognized subspecies have recently been confirmed as genetically distinct lineages (Chiari et al. 2005).

07:71. ***Stigmochelys* or *Psammobates pardalis***: Based on genetic analysis, Le et al. (2006) recommended that this taxon be included in an expanded genus *Psammobates*. Fritz and Bininda-Emonds (2007) argued for the retention of a monophyletic *Psammobates* exclusive of *pardalis*. Le at al. (2006) also found a high level of mitochondrial divergence between two specimens assigned to the two subspecies *pardalis* and *babcocki*. In conjunction with morphological distinctions between these two taxa (Loveridge and Williams 1957; Broadley 1989), the preliminary genetic data suggest that they may be different at the species level.

07:72. ***Testudo* or *Agrionemys***: The species *horsfieldii* and *hermanni* have been alternatively placed in the genera *Testudo* or *Agrionemys* (Khosatzky and Mlynarski 1966; Gmira 1993, 1995) and *hermanni* also recently in *Eurotestudo*. Lapparent de Broin (2000a,b) and Parham et al. (2006b) supported the placement of *horsfieldii* in the genus *Agrionemys*, but suggested that a new genus name was needed for *hermanni*. Later Lapparent de Broin et al. (2006a) created the name *Eurotestudo* for *hermanni*, but Fritz and Bininda-Emonds (2007) demonstrated that older genus names (*Chersine* and *Medaestia*) are available for that species. Fritz and Bininda-Emonds (2007) recovered a weakly monophyletic clade that included *horsfieldii, hermanni*, and the three core species of *Testudo* (*graeca, kleinmanni*, and *marginata*). Based on this phylogeny they recommended that all of these species be placed in the genus *Testudo*. The genetic support for some nodes within this clade is not strong and the decision to lump or split is subjective (e.g., whether *Agrionemys* should be used for *horsfieldii* is open to debate), therefore the taxonomy of this group may remain in flux for some time.

07:73. ***Testudo graeca***: This species complex has been the subject of massive taxonomic revisions at the species and subspecies level. These revisions have resulted in the naming and elevation of numerous taxa (e.g., Perälä 2002a,b,c). Several studies (van der Kuyl et al. 2002, 2005; Harris et al. 2003; Carretero et al. 2005; Parham et al. 2006b,c; Fritz et al. 2007a) have explicitly refuted the validity of many of these taxonomic acts. Fritz et al. (2007a) proposed a taxonomic scheme that recognized five mitochondrial clades in the eastern part of the range of *T. graeca* as subspecies, but did not address the status of several North African subspecies. Since this is the most recent taxonomic suggestion, it is listed here. However, in their recent checklist, Fritz and Havas (2006, 2007) included not only the eleven taxa we list, but also *anamurensis* Weissinger 1987, *antakyensis* Perälä 1996, *floweri* Bodenheimer 1935, *nikolskii* Chkhikvadze and Tuniyev 1986, *pallasi* Chkhikvadze and Bakradze 2002, and *perses* Perälä 2002c. The relationships within this species complex remain uncertain and we expect its taxonomy to continue fluctuating.

07:74. ***Testudo kleinmanni***: Baha el Din (2006), Široký and Fritz (2007), and Attum et al. (2007) explicitly refuted the validity of *werneri* Perälä 2001 as a species distinct from *kleinmanni*.

07:75. ***Testudo marginata***: Fritz et al. (2005b) explicitly refuted the validity of *weissingeri* Bour 1996 as a subspecies of *marginata*.

07:76. ***Testudo hermanni***: Fritz et al. (2006b) explicitly refuted the validity of *hercegovinensis* Werner 1899 (previously resurrected by Perälä 2002b) and recommended that *boettgeri* be considered a subspecies of *hermanni*.

07:77. ***Testudo horsfieldii***: In a conference proceedings, Perälä (2002a) elevated two subspecies of *horsfieldii* (*kazachstanica* and *rustamovi*) to full species status. This was accepted by Lapparent de Broin et al. (2006b), but warrants reconsideration, especially considering the evidence for unjustified taxonomic inflation in related tortoises in the same work (van der Kuyl et al. 2002, 2005; Fritz et al. 2005b, 2006b; Parham et al. 2006b,c).

07:78. ***Carettochelys insculpta canni***: This subspecies from northern Australia described by Wells (2002a) was only weakly defined as different from the nominotypical subspecies from New Guinea. We list it tentatively pending further analysis, as did Fritz and Havas (2006), although they excluded it from their 2007 checklist.

07:79. ***Apalone spinifera atra***: This taxon has usually been designated a subspecies of *spinifera* (usually with the original spelling *ater*), but others (e.g., Flores-Villela 1993; David 1994) have listed it as a full species, though usually without specific argumentation.

07:80. ***Aspideretes* or *Nilssonia***: Engstrom et al. (2004) found *Aspideretes* to be paraphyletic with respect to *Nilssonia formosa* based on morphologic and genetic criteria. Praschag et al. (2007a) formally synonymized *Aspideretes* into an expanded concept of *Nilssonia* based on their analysis of mtDNA of all five included taxa.

07:81. ***Aspideretes* or *Nilssonia nigricans***: Recent morphologic and genetic work on this species previously known only from a single captive population has demonstrated that it also occurs in the wild (Praschag and Gemel 2002; Praschag et al. 2007a).

07:82. ***Pelodiscus***: The genus has recently been recognized as including up to four separate species by some authorities (David 1994; Zhao 1997; Chen et al. 2005, 2006; Fritz and Havas 2006, 2007). Relationships within the genus are far from resolved and also complicated by translocation and mixing of huge numbers of farm-raised individuals from many parts of the range.

07:83. ***Rafetus swinhoei***: Includes the recently described *Pelochelys taihuensis* Zhang 1984 (Farkas 1992) and *Rafetus leloii* Duc 2000 in synonymy (Farkas and Webb 2003).

07:84. ***Acanthochelys macrocephala***: Includes the recently described *Phrynops chacoensis* Fritz and Pauler 1992 in synonymy (Fritz and Pauler 1999).

07:85. ***Chelodina***: This genus was split into three genera by Wells and Wellington (1985), using *Chelodina* for the narrower-headed shorter-necked species (*longicollis, novaeguineae*), and establishing *Macrochelodina* for the broader-headed longer-necked species (*oblonga, expansa, rugosa, siebenrocki*), and *Hesperochelodina* for *steindachneri*. Iverson et al. (2001b) refuted the availability of the name *Hesperochelodina*, but validated *Macrochelodina*. Georges et al. (2002) retained *Chelodina* for the entire genus, but identified three phylogenetic clades within the genus and recommended recognition of three subgenera (but did not name them). Fritz and Havas (2006, 2007) accepted two of these clades (*Chelodina* and *Macrochelodina*) as separate genera.

07:86. ***Chelodina canni***: This taxon is the same as the previously described *rankini* Wells and Wellington 1985, but that name was declared invalid as a *nomen nudum* by Iverson et al. (2001b). Wells (2007a) recently disputed this interpretation and redescribed *rankini*, but *canni* McCord and Thomson 2002 retains nomenclatural precedence and *rankini* Wells 2007a is therefore a junior synonym of *canni*.

07:87. ***Chelodina mccordi roteensis***: This recently named subspecies described in the hobbyist literature needs genetic confirmation of its distinctiveness, but we recognize it pending further analysis.

07:88. ***Chelodina oblonga***: Thomson (2000a) showed that the holotype of *oblonga* Gray 1841 is a specimen of what is currently regarded as *Chelodina rugosa* Ogilby 1890. An application is before the International Commission for Zoological Nomenclature (ICZN) to conserve current usage of the name *C. rugosa* Ogilby 1890 for the northern snake-necked turtle and to apply the earlier available name *Chelodina collei* Gray 1856a to the long-necked species of southwestern Australia, while retaining the nomenclatural availability of the name *oblonga* for potential future designation of distinct populations of *rugosa* (Thomson 2006). Though no decision has yet been rendered by the ICZN, Fritz and Havas (2006, 2007) used the name *colliei* for this southwestern population. Georges et al. (2002) found support that this taxon represents a third subgenus under *Chelodina*, but did not formally establish it under a generic-level name.

07:89. ***Chelodina timorensis***: This species recently described in the hobbyist literature by McCord et al. (2007b) was also described a few months later as a new subspecies of *mccordi* (‘*timorlestensis*’) by Kuchling et al. (2007), but the McCord et al. description has chronologic precedence. Concerns surrounding the history and methodology of the description of *timorensis* by McCord et al. are discussed by Kuchling et al. (2007) and serve to emphasize our recommendations (made in our other chapter in this volume) to follow certain procedural guidelines for descriptions of new taxa (Turtle Taxonomy Working Group 2007a).

07:90. ***Chelodina kuchlingi***: This species was described from a single specimen, leading to doubts about its validity (Georges and Thomson 2006; Fritz and Havas 2006, 2007), but it remains listed pending further exploration of its remote area of provenance.

07:91. ***Chelodina rugosa***: The species *siebenrocki* Werner 1901 was considered valid by Rhodin and Mittermeier (1976) and Rhodin and Genorupa (2000), but synonymized under *rugosa* by Georges et al. (2002) based on weakly differentiated allozymes within the broader *rugosa* complex.

07:92. ***Elseya***: This genus has been recognized as consisting of two separate lineages (Georges and Rose 1996; Georges and Thomson 2006). It was subsequently split into two genera, *Elseya* and *Wollumbinia*, by Wells (2007c), with *latisternum* designated genotype of *Wollumbinia*. Papers by Wells (2002a,b; 2007a,b,c) and Wells and Wellington (1985) have been self-published without any peer review and also highlight our recommendations to follow certain procedural guidelines for descriptions of new taxa (Turtle

Taxonomy Working Group 2007a).

07:93. ***Elseya branderhorsti***: This species was considered valid by Rhodin and Genorupa (2000), Thomson et al. (2006), and Georges and Thomson (2006).

07:94. ***Elseya jukesi***: The name *jukesi* Wells 2002b was a *nomen nudum* since no type specimen was designated, but the species was recently redescribed by Wells (2007b).

07:95. ***Elseya schultzei***: This species was listed by Thomson et al. (2006) and Georges and Thomson (2006), but neither morphologic nor genetic data have been analyzed from the type population and its status remains unclear.

07:96. ***Elseya stirlingi***: The previously named taxon *stirlingi* Wells and Wellington 1985 was declared invalid as a *nomen nudum* by Iverson et al. (2001b) (though spelled erroneously as *sterlingi*), but was recently redescribed as a valid species by Wells (2007b).

07:97. ***Elseya* or *Wollumbinia bellii***: The taxon *dorriani* Wells 2002b is a *nomen nudum* without a type designation, but was recently considered a valid subspecies of *bellii* by Wells (2007c).

07:98. ***Emydura macquarii***: The taxonomy of *E. macquarii* was previously reviewed by Georges and Adams (1996). Later, Cann et al. (2003) and McCord et al. (2003) described two new subspecies, but taxa previously described by Cann in 1998 (*binjing, dharra, dharuk,* and *gunabarra*), plus *signata* Ahl 1932 were not specifically evaluated by those authors. However, these taxa were all recognized as subspecies of *macquarii* by Fritz and Havas (2006, 2007), and since phylogeographic variation in the *macquarii* species complex has not yet been fully resolved with adequate genetic work, we tentatively list all these subspecies as valid, pending further analysis.

07:99. ***Emydura subglobosa worrelli***: Originally described as *Tropicochelymys worrelli*, this taxon was synonymized under *Emydura victoriae* Gray 1842 by Iverson (1992) and the nomenclatural validity of the species name confirmed by Iverson et al. (2001b). Cann (1998) considered it a distinct species, but Georges and Thomson (2006), partially based on electrophoretic work by Georges and Adams (1996), concluded that it was best referred to as a subspecies of *subglobosa* Krefft 1876. Fritz and Havas (2006, 2007) also listed it as a subspecies of *subglobosa*, but Georges et al. (2006) referred to it as a species, though without providing data or argument.

07:100. ***Phrynops***: Wermuth and Mertens (1977) divided this genus into three subgenera: *Phrynops*, *Batrachemys*, and *Mesoclemmys*. Cabrera (1998) and Georges et al. (1998) elevated these subgenera to generic level. McCord et al. (2001) further divided the remaining monophyletic *Phrynops* into a total of four genera (*Bufocephala*, *Phrynops*, *Ranacephala*, and *Rhinemys*). Joyce et al. (2004) did not accept the taxonomic acts of McCord et al. (2001). Bour and Zaher (2005) synonymized *Bufocephala* and *Ranacephala* with *Mesoclemmys*, but recognized *Rhinemys* as distinct.

07:101. ***Mesoclemmys heliostemma***: Rueda-Almonacid et al. (2007) questioned the validity of this taxon which is completely sympatric with *raniceps*, suggesting that it may simply represent a juvenile color morph of that taxon, and recommended genetic analysis.

07:102. ***Pelomedusa subrufa***: Gasperetti et al. (1993) recommended that the two previously recognized subspecies (*nigra* Gray 1863b and *olivacea* Schweigger 1812) be abandoned.

07:103. ***Pelusios seychellensis***: The taxonomic status of this species is unclear. Gerlach and Canning (2001) concluded that it is extinct.

07:104. **Podocnemididae or Podocnemidae**: Cope (1868) used the name Podocnemididae to refer to this clade. Baur (1893b) later referred to this group as Podocnemidae. Joyce et al. (2004) phylogenetically defined Baur's name (Podocnemidae) to refer to this clade.

07:105. ***Podocnemis unifilis***: This long-recognized species was briefly referred to as *P. cayennensis* Schweigger 1812 by David (1994), but that name was previously often used for what is now recognized as *P. erythrocephala* (Mittermeier and Wilson 1974), and most authors have continued to use *unifilis*.

2008 Checklist Annotations

Rhodin et al. 2008 (000.1-38.checklist.v.1) [(08:2-25)]

08:2. ***Chelodina* or *Macrodiremys***: The southwestern long-necked turtle of Australia (*Chelodina oblonga* or *colliei*, see annotation below) represents one of three lineages that were considered unnamed subgenera of *Chelodina* by Georges et al. (2002). McCord and Joseph-Ouni (2007b) created the name *Macrodiremys* for *oblonga*/*colliei*, designating *Chelodina oblonga* Gray 1841, as type species by original designation and monotypy,and elevated this to a full monotypic genus. Whether *Chelodina* sensu stricto will be a subgenus of *Chelodina* sensu lato along with *Macrodiremys* and *Macrochelodina* or if all three will be used as full genera is subjective and not yet stable.

08:3. ***Chelodina* or *Macrodiremys oblonga***: Within *Chelodina*, the specific epithet *oblonga* has long been applied to a long-necked species in southwestern Australia. Thomson (2000a) showed that the holotype of *oblonga* Gray 1841 is a specimen of what is currently regarded as *Chelodina rugosa* Ogilby 1890 from northern Australia. An application (Thomson 2006, 2007) is before the International Commission of Zoological Nomenclature (ICZN) to conserve current usage of the name *C. rugosa* Ogilby 1890 for the northern snake-necked turtle and to apply the next available name, *Chelodina colliei* Gray 1856a, to the long-necked species of southwestern Australia. Separately, McCord and Joseph-Ouni (2007b) designated the holotype of *colliei* as the neotype of *oblonga* which would render *colliei* a junior synonym of *oblonga* which would be incompatible with an identification of the holotype and name *oblonga* as pertaining to the northern Australian taxon. We list the southwestern long-necked species as *oblonga* because McCord and Joseph-Ouni (2007b) is the latest published action but note that, given the differing taxonomic acts and opinions, this name may remain unstable in the coming years.

08:4. ***Macrochelodina* or *Chelodina walloyarrina***: McCord and Joesph-Ouni (2007b) described the new species *Macrochelodina walloyarrina* based on morphological criteria.

08:5. ***Chelydra serpentina***: Shaffer et al. (2008) recommended synonymization of *Chelydra serpentina osceola* Stejneger 1918 into *Chelydra serpentina* (Linnaeus 1766) based on range-wide patterns in variability of mtDNA.

08:6. ***Cyclemys***: Fritz et al. (2008b) performed a revision of the genus based on molecular and morphological data that included the description of three new species (*enigmatica*, *fusca*, and *gemeli*).

08:7. ***Cyclemys dentata***: Stuart and Fritz (2008) analyzed mtDNA from type specimens of *Cyclemys belli* Gray 1863e, *Cyclemys orbiculata* Bell 1834 and *Cyclemys ovata* Gray 1863e, and confirmed their previous morphology-based synonymizations with *Cyclemys dentata* (Gray 1831d) as accurate.

08:8. ***Cyclemys oldhamii***: Stuart and Fritz (2008) placed the names *shanensis* Annandale 1918, and *tcheponensis* Bourret 1939a, into the synonymy of *oldhamii*, based on the absence of significant genetic variation between the type specimen of *oldhamii*, topotypes of *shanensis*, and samples of *tcheponensis* from near the type locality.

08:9. ***Batagur***: Praschag et al. (2008) examined mtDNA variation within *Batagur baska* sensu lato and recommended that the southern populations should be elevated to full species status and for which the available name *affinis* Cantor 1847 should be used.

08:10. ***Carettochelys insculpta***: Fritz and Havas (2007) and Georges et al. (2008) indicated that *Carettochelys insculpta canni* Wells 2002a is not an available name because it had not been published in accordance with criteria established by the International Code of Zoological Nomenclature.

08:11. ***Kinixys nogueyi***: This taxon was treated as a full species by McCord et al. (2005) with minimal argumentation for the change from traditional recognition as a subspecies of *K. belliana*.

08:12. ***Rhinoclemmys punctularia***: *Testudo scabra* Linnaeus 1758 has previously been referred (as a *nomen dubium*) to the synonymies of both *Melanochelys trijuga* (Schweigger 1812) and *Rhinoclemmys punctularia* (Daudin 1801) (see Fritz and Havas 2007). Examination of the holotype specimen still extant in the Uppsala Linnaean collection indicates that it appears to be a *Rhinoclemmys punctularia* (Rhodin and Carr, in press) [2009]. However, since the name *Testudo scabra* has not been used as the name for a recognized taxon since the early 1800s, it remains a *nomen oblitum* and does not replace the name *punctularia* Daudin 1801, recognized and used as valid since its description, and also protected by the ICZN (1963) as a *nomen protectum*.

08:13. ***Aldabrachelys* or *Dipsochelys***: The generic and specific names of the Aldabra tortoise are still being debated (reviewed in TTWG 2007). The original type specimen of *Testudo gigantea* Schweigger 1812 is a *Chelonoidis denticulata* from Brazil (Bour 2006c), but since the name *gigantea* had been associated with tortoises from Aldabra for a long time, Frazier (2006) designated a neotype from Aldabra, leading to some confusion. The matter is currently being petitioned to the International Commission on Zoological Nomenclature (Frazier 2008).

08:14. ***Chelonoidis nigra vicina***: The type locality for *Testudo microphyes* Günther 1875a was given as follows: "I suppose that the specimen… has come from Hood's Island," and Pritchard (1996) concluded that the name

microphyes was a *nomen dubium* since its type specimen was not identifiable as either a Hood Island tortoise or any other recognizable taxon. Fritz and Havas (2007) indicated that Günther (1877) had subsequently designated the type locality for *microphyes* as "Tagus Cove, northern Albemarle Island" and they placed *microphyes* in the synonymy of *Testudo vicina* Günther 1875a, but this is not necessarily correct. Günther (1877) simply referred later-collected specimens from Tagus Cove to his type-based concept of *microphyes*—the name *microphyes* therefore remains a *nomen dubium* until such time as the original type specimen can perhaps be identified as to its exact provenance using genetic analysis.

08:15. ***Agrionemys* or *Testudo horsfieldii***: Vasilyev et al. (2008) demonstrated minimal mitochondrial variation between populations of *horsfieldii* Gray 1844 and *kazachstanica* Chkhikvadze, Amiranashvili, and Ataev 1990 and so recommended that these taxa be considered subspecies of *Agrionemys horsfieldii*. Elsewhere in the paper they referred to *A. h. rustamovi* as a third subspecies, but made no definitive comment on taxonomic status or validity.

08:16. ***Podocnemis unifilis***: The terminal taxon *lata* Bell *in* Gray 1830e has previously been included under the synonymy of *Peltocephalus dumerilianus* (Schweigger 1812) by many previous authors, including Fritz and Havas (2007), but Bell (*in* Gray 1830e) described *Chelys (Hydraspis) lata* as having a depressed black shell and orange-spotted head more typical of *Podocnemis unifilis* or *P. expansa*. Later, Gray (1870f) placed *Hydraspis lata* in the synonymy of his concept of *Chelonemys dumeriliana* (= *Podocnemis unifilis*) and added *P. unifilis, P. cayennensis*, and *P. erythrocephala* to the same synonymy, while differentiating *Peltocephalus tracaxa* (= *Peltocephalus dumerilianus*) as a distinct taxon. Though *Chelys (Hydraspis) lata* Bell *in* Gray 1830e is an older name than *Podocnemis unifilis* Troschel 1848, it is a *nomen oblitum* not used for a valid taxon since its description, and therefore does not replace *unifilis* as the valid name for the species.

08:17. ***Trachemys decussata***: Authorship of this taxon was actually first by Bell (*in* Griffith and Pidgeon 1830) with a plate. Seidel (1988a) listed the author as Gray 1831:28 (= Gray 1831d), and Fritz and Havas (2007) listed the author as Gray 1831:11 (= Gray 1830e). Griffith and Pidgeon 1830 was published in September 1830, whereas Gray 1830e was published in December 1830, and Gray 1831d was published in May 1831.

08:18. ***Graptemys geographica***: The name *lesueuri* Gray 1830d (= *lesueurii* Gray 1830e) was shown by Bour and Dubois (1983) to be a junior synonym of *geographica* LeSueur 1827 rather than a senior synonym of *pseudogeographica* Gray 1831d as recorded by Fritz and Havas (2007).

08:19. ***Graptemys pseudogeographica***: The name *pseudogeographica* Gray 1831d was originally published only as a *nomen nudum* of a LeSueur manuscript name in junior synonymy under *geographica* LeSueur 1827, but gradually achieved wide usage by many authors. Holbrook (1842) was the first to actually describe the taxon under the name *pseudogeographica*, and arguably his name should perhaps be associated with it, but Stejneger and Barbour (1917) established the name as *pseudogeographica* Gray 1831d as pointed out by Bour and Dubois (1983), who agreed that Gray should be listed as the author.

08:20. **Cryptodira and Pleurodira**: These subordinal names were based on the French vernacular names, Cryptodères and Pleurodères, originally used by Duméril and Bibron 1834. Cope (1864, 1865, 1868b) has generally been credited with authorship of these names, and he was the first to use the exact names Cryptodira in 1868 and Pleurodira in 1865, but previously used the name Pleurodera in 1864. Cope was preceded by Lichtenstein (1856) who used Cryptodera and Pleurodera as subordinal names, but in a printed catalogue distributed to zoological colleagues and museums, and not apparently sold in bookstores, so therefore perhaps not nomenclaturally available.

08:21. ***Cuora evelynae***: In the previous checklist (TTWG 2007), *Cuora flavomarginata* (Gray 1863e) had three subspecies. Ernst et al. (2008) recommended returning *Cuora evelynae* Ernst and Lovich 1990 to full species status, based on new morphological and previously published molecular data. They also argued that the mainland populations assigned to taxon *sinensis* Hsü 1930 are synonymous with nominotypical *flavomarginata*.

08:22. ***Apalone spinifera***: In the previous checklist (TTWG 2007), seven subspecies of *spinifera* LeSueur 1827 were listed. McGaugh et al. (2008) performed a rangewide phylogeographic study that uncovered patterns of discordant molecular and morphological variation. These authors conservatively refrained from making sweeping nomenclatural changes, but noted that there was "little utility" in recognizing the taxon *hartwegi* Conant and Goin 1948 and synonymized it under *spinifera*.

08:23. ***Apalone spinifera atra***: In the previous checklist (TTWG 2007), *atra* Webb and Legler 1960 was referred to as a subspecies of *spinifera* LeSueur 1827 as it is here. This taxon is sometimes considered a full species (e.g., argumentation cited in Flores-Villela 1993). Several recent studies on *atra* (McGaugh 2008, McGaugh and Janzen 2008, McGaugh et al. 2008) have argued in support of subspecies status based on low levels of genetic distinctiveness and habitat-driven color variation. The subspecific status of *atra* was followed by Cerdá-Ardura et al. (2008).

08:24. ***Sacalia quadriocellata***: Shi et al. (2008) performed a mitochondrial survey of *S. quadriocellata* based on known-locality and trade specimens. They found that populations on Hainan are genetically distinct and can also be diagnosed by morphological characters. They did not elevate these populations to species status, but noted that eventual study might validate this conclusion, in which case the name *Sacalia insulensis* (Adler 1962) would be available. They also noted that samples of *S. quadriocellata* from northern Vietnam are genetically distinct from those from Laos and the type region of central Vietnam, but there appear to be no obvious morphological differences between these populations.

08:25. ***Rhinoclemmys flammigera***: Barrio-Amorós and Narbaiza (2008) elevated *Rhinoclemmys punctularia flammigera* Paolillo 1985 to species status based on distinct head coloration pattern and allopatric isolated distribution.

2009 Checklist Annotations

TTWG 2009 (000.39-84.checklist.v.2) (09:3-49)

09:3. **Chelydridae**: Chandler and Janzen (2009) analyzed the phylogenetic position of the Chelydridae based on nucleotide sequence data, and found weak support for a sister group relationship with either the Kinosternoidea (Kinosternidae + Dermatemydidae) or Chelonioidea (Cheloniidae + Dermochelyidae). In a more extensive analysis, Barley et al. (in press) show that Chelydridae is sister to Kinosternoidea.

09:4. ***Macrochelys temminckii***: Roman et al. (1999) showed that *M. temminckii* could be divided into three major mitochondrial clades which they treated as Evolutionarily Significant Units (ESUs). They noted that samples from the Suwannee drainage in Florida showed high divergence from the rest of the range. Echelle et al. (2009) performed a microsatellite study and further subdivided *M. temminckii* into six ESUs. They also noted that the Suwannee population was the most distinct and concluded that it might eventually be recognized as a distinct taxonomic unit.

09:5. ***Cheloniidae***: Bowen and Karl (2007) reviewed population genetics and phylogeography of marine turtles and while they noted mtDNA divergence between Indo-Pacific and Atlantic *Chelonia mydas* and *Eretmochelys imbricata*, they recognized no taxa below the species level.

09:6. **Lacepède 1788 and Bonnaterre 1789**: The International Commission on Zoological Nomenclature (ICZN) previously rejected the names created by Lacepède in his 1788 *Histoire Naturelle des Serpens* and subsequent editions since they were published in non-binomial works (ICZN 1987a). However, all names published in his earlier 1788 volume, *Histoire Naturelle de Quadrupèdes Ovipares* (which contained all his turtle descriptions), continued to be nomenclaturally available until recently, when they were also rejected as being published in a non-binomial work (ICZN 2005). A few of these turtle names from the 1788 *Histoire Naturelle de Quadrupèdes Ovipares* volume had already been individually suppressed by the ICZN (1963). Therefore, Bonnaterre (1789), who republished Lacepède's descriptions with proper binomials, becomes the authorship source for these rejected Lacepède turtle names.

09:7. ***Testudo nasicornis***: *Testudo nasicornis* Lacepède 1788 was described as distinct from *Testudo caouana* Lacepède 1788 (= *Caretta caretta*) based on the possession of a soft nasal projection and on being fit for human consumption like *Testudo marina* (= *Chelonia mydas*). The species was included in the synonymy of various other marine turtle taxa until Loveridge and Williams (1957) placed it alongside *T. caouana* in the synonymy of *Caretta caretta* (Linnaeus, 1758). Bonnaterre (1789) provided an amplified description and drawing of *Testudo nasicornis*, but his concept of the taxon differed from that of Lacepède, illustrating a specimen (pl. 3, fig. 3; hereby designated as the holotype) with 13 large carapacial scutes, costal tubercles forming an interrupted lateral keel, a strongly serrated carapacial margin, and apparently two pairs of prefrontal scales. The 13 carapacial scutes are inconsistent with *Caretta* but correspond to *Chelonia* or *Eretmochelys*, while the serrated carapacial margin and apparently split prefrontals indicate that the figured specimen is an *Eretmo-*

chelys imbricata (Linnaeus, 1766). As long as the name *Testudo nasicornis* was nomenclaturally occupied by Lacepède's 1788 description, Bonnaterre's 1789 description could simply be dismissed as an incorrect subsequent attribution. However, now that ICZN Opinion 2104 (ICZN 2005) has rejected availability of all of Lacepède's turtle names, Bonnaterre's becomes the first available usage of the name *Testudo nasicornis*. Bonnaterre's description is clearly attributable to *Eretmochelys imbricata*, and thus we consider *Testudo nasicornis* Bonnaterre 1789 to be a subjective junior synonym of that taxon, while retaining *Testudo nasicornis* Lacepède 1788 as a *nomen rejectum* attributable to *Caretta caretta*.

09:8. **Meyer 1790 species names**: Meyer (1790), in a long-overlooked review article, provided short diagnoses and replacement names (*nomina nova*) for four species described by Lacepède (1788). These names have never appeared in any subsequent synonymies and are clearly *nomina oblita*. The Meyer names are *Testudo bomarii* for *Testudo viridisquamosa* (= *Lepidochelys kempii* or *Chelonia mydas*), *Testudo lauanna* for *Testudo caouana* (= *Caretta caretta*), *Testudo sonnerati* for *Testudo punctata* (= *Lissemys punctata punctata*), and *Testudo rubra* for *Testudo subrufa* (= *Pelomedusa subrufa*).

09:9. ***Eretmochelys imbricata***: This checklist (TTWG 2007b; Rhodin et al. 2008) has previously treated *bissa* as a valid subspecies of *imbricata* in the absence of a definitive, data-based synonymization. Whereas genetic results have demonstrated distinct genetic lineages in the Atlantic and Indo-Pacific Oceans, no genetic studies or reviews (Okayama et al. 1999; Bowen and Karl 2007) have argued for continued recognition of the subspecies *bissa*. The phylogenetic structure within *Eretmochelys* is comparable to that within *Chelonia*, for which only a single monotypic species is currently recognized, and thus we now treat *bissa* as a synonym of *imbricata*.

09:10. ***Kinosternon arizonense***: The authorship of this name was given as *arizonense* Gilmore 1922 in our previous checklist as well as TTWG (2007b) and Fritz and Havas (2007). However, the article appeared in February 1923 and the year is therefore corrected.

09:11. ***Kinosternon hirtipes***: The authorship of this name was given as Wagler 1833 in our previous checklist as well as in TTWG (2007b) and Fritz and Havas (2007), whereas previous authors (e.g., Iverson 1992) have recognized *hirtipes* Wagler 1830, a name sometimes interpreted as a *nomen nudum*. However, the ICZN (1999) rules for availability of names published prior to 1931 (Article 12) state that species names must be accompanied by a "description or definition" or by an "indication". The 1830 citation for *hirtipes* is not accompanied by a description or definition, but is accompanied by an indication—the associated illustration of the holotype of the taxon being named (see Article 12.2.7). Wagler 1830 is therefore the correct authorship designation.

09:12. **Deirochelyinae**: Spinks et al. (2009b) performed a phylogenetic analysis of the Emydidae based on mitochondrial and nuclear DNA. Their mitochondrial phylogeny did not recover a monophyletic Deirochelyinae, instead it placed *Deirochelys* as the sister to the rest of Emydidae. In contrast, their nuclear data recovered a monophyletic Deirochelyinae.

09:13. ***Pseudemys***: Spinks et al. (2009b) performed a phylogenetic analysis of Emydidae based on mitochondrial and nuclear DNA. Although their sampling within *Pseudemys* was limited and uneven, their samples of *concinna* and *floridana* did not yield a monophyletic *P. concinna* with respect to *peninsularis* (mtDNA and nuclear DNA) or *nelsoni* (nuclear DNA only). Further genetic and morphological study of known locality samples will be necessary in order to resolve the taxonomic status of the terminal taxa within *Pseudemys*.

09:14. ***Trachemys***: Seidel (2002) made several recommendations about the specific and subspecific taxonomy of *Trachemys*, as reflected in our earlier checklist. Jackson et al. (2008) performed a mitochondrial survey of the genus and supported those taxonomic revisions.

09:15. ***Trachemys scripta***: Rhodin and Carr (2009) demonstrated that the original authorship of the taxon name *scripta* should be attributed to Thunberg *in* Schoepff (1792), rather than just Schoepff.

09:16. ***Emys* or *Actinemys* and *Emys* or *Emydoidea***: Using nuclear markers, Spinks and Shaffer (2009) re-emphasized a close phylogenetic relationship among *marmorata*, *orbicularis/trinacris*, and *blandingii* as was previously shown from mitochondrial DNA (see Feldman and Parham 2002 and case summary in annotation 07:21 of TTWG 2007b). Spinks and Shaffer (2009) also showed that those species share a complex evolutionary history including prehistoric hybridization, and that *blandingii* and *orbicularis/trinacris* are sister taxa. In light of this evidence they strongly recommended that all these species be included in the genus *Emys* rather than continued recognition of the genera *Actinemys* and *Emydoidea*. Other authors (Iverson et al. 2008) argue for the continued recognition of all three genera in this clade.

09:17. ***Emys orbicularis orbicularis***: Fritz et al. (2009b) demonstrated that the mitochondrial DNA differentiation of the two previously recognized subspecies *colchica* and *luteofusca* were insufficient to continue to recognize them as distinct and therefore synonymized both under *orbicularis*.

09:18. ***Emys orbicularis fritzjuergenobsti***: Velo-Antón et al. (2008) performed a genetic analysis of multiple populations of *Emys orbicularis* on the Iberian peninsula and found no significant genetic divergence between the two previously-defined subspecies *hispanica* and *fritzjuergenobsti*, and therefore synonymized the former under the latter.

09:19. ***Emys orbicularis persica***: Fritz et al. (2009b) demonstrated that the mitochondrial DNA differentiation of the previously recognized subspecies *iberica* was insufficient to continue to recognize it as distinct and therefore synonymized it under *persica*, thereby also bringing *kurae* under the synonymy of *persica*.

09:20. **Geoemydidae and *Rhinoclemmys***: Le and McCord (2008) evaluated the molecular phylogeny of *Rhinoclemmys* and other geoemydid genera and affirmed the monophyly of the Geoemydidae, but recommended that *Rhinoclemmys* be afforded subfamilial recognition as the Rhinoclemminae, a grouping concept first proposed by Gray (1873j) as the Tribe Rhinoclemmyina.

09:21. ***Batagur affinis edwardmolli***: Praschag et al (2009) assessed the taxonomic status of *B. affinis* using mitochondrial and nuclear genetic analysis, and described the populations of Cambodia and the eastern coast of Peninsular Malaysia as the new subspecies *edwardmolli*, with the populations of western Peninsular Malaysia and Sumatra (Indonesia) retained as the nominotypical subspecies *affinis*.

09:22. ***Cuora galbinifrons* complex**: In our previous checklists we listed the three taxa *galbinifrons*, *bourreti*, and *picturata* as subspecies of *galbinifrons*. However, the preponderance of well-documented evidence now supports the elevation of these three taxa to recognition as three closely related full species, based on both morphology and genetics (Stuart and Parham 2004, Spinks et al. 2009a).

09:23. ***Cuora trifasciata***: Spinks et al. (2009a) assessed the validity of the recently described *Cuora cyclornata* Blanck, McCord, and Le 2006a using a combination of mitochondrial and nuclear genetic markers. Their genetic evidence elucidates a complex history of introgression involving *Cuora trifasciata* and the *Cuora pani* complex. They also provide a critique of the morphological analysis of Blanck et al. (2006), concluding that the available evidence is not sufficient to diagnose *C. cyclornata*. Following their previous analysis of this group (Spinks et al. 2006), they continue to recommend that *C. cyclornata* be considered a junior synonym of *C. trifasciata*.

09:24. ***Cyclemys* species**: Praschag et al. (2009b) analyzed mitochondrial and nuclear genes in these species, and found that *gemeli* and *fusca* were distinct but closely related, and that *atripons*, *dentata*, and *pulchristriata* were also well-differentiated and formed a well-supported clade. The taxonomy of the genus *Cyclemys* has been subject to intense debate over the past several years and will likely continue for some time before it is stabilized.

09:25. ***Geoemyda spengleri***: Gong et al. (2009a) demonstrated phylogeographic structure in mitochondrial DNA within this taxon.

09:26. ***Mauremys***: Hirayama et al. (2007) recommended splitting the genus *Mauremys* (sensu Feldman and Parham 2004) into five genera (*Mauremys*, *Cathaiemys*, *Chinemys*, *Ocadia*, and an unnamed new genus) based on the morphology of the palate. The relative utility of single character typological taxonomies and monotypic genera versus restricting familiar names to well-defined evolutionary clades has been discussed elsewhere for *Mauremys* and turtles in general (Parham and Feldman 2002; Feldman and Parham 2004; Spinks et al. 2004; Turtle Taxonomy Working Group 2007a; Spinks et al. 2009). Following the philosophy outlined in these papers we retain the larger aggregate *Mauremys* and recommend consideration of subgenera for phenetically distinctive subclades (e.g., Parham and Feldman 2002; Smith and Chizsar 2006).

09:27. ***Mauremys caspica***: Fritz et al. (2008a) performed a rangewide genetic survey of *Mauremys caspica*. Their study revealed discordant patterns of morphological and genetic differentiation in this species. They did not recommend abandoning the current subspecies, but highlighted the need for future taxonomic revision.

09:28. ***Rhinoclemmys punctularia flammigera***: Barrio-Amorós and Narbaiza (2008) elevated this taxon to species status based on a brief statement about head coloration and allopatric distribution, a change we reflected in our previous checklist; however, based on the relative lack of supportive data, we treat it again as a subspecies pending further analysis.

09:29. ***Aldabrachelys* or *Dipsochelys***: The nomenclatural validity of the

generic and specific names of the Aldabra tortoise (*Aldabrachelys gigantea* or *Dipsochelys dussumieri*) is currently being debated (reviewed in TTWG 2007b and Rhodin et al. 2008). There is recent disagreement regarding the type specimen of *Testudo gigantea*, the type species of *Aldabrachelys*, that was presumed lost. Frazier (2006) designated an Aldabran neotype for *T. gigantea*, an act that would validate the use of both *Aldabrachelys* and the terminal taxon name *gigantea*. Around the same time, Bour (2006c) reported to have rediscovered the original lost type specimen, which is actually an individual of the South American tortoise *Chelonoidis denticulata*, making *gigantea* and *Aldabrachelys* junior synonyms of *denticulata* and *Chelonoidis*, respectively, which would thereby result in the use of *Dipsochelys dussumieri* as the valid name for the Aldabra tortoise. Whether Frazier's neotype designation or Bour's specimen rediscovery prevails nomenclaturally remains a matter of major ongoing debate. The matter has been petitioned to the International Commission on Zoological Nomenclature (Frazier 2009), with multiple commentaries on both sides of the issue (Zug et al. 2009; Bour et al. 2009; Takahashi et al. 2009), but no decision has yet been made by the ICZN.

09:30. ***Aldabrachelys* or *Dipsochelys* species**: Gerlach and Canning (1998) recognized six species of tortoises in Aldabra, Madagascar, and the Seychelles (three extant: *gigantea* or *dussumieri*, *arnoldi*, and *hololissa*; and three extinct: *abrupta*, *daudinii*, and *grandidieri*). The two species from Madagascar became extinct prior to modern times (*Testudo abrupta* Grandidier 1868 in ca. 1250 AD and *Testudo grandidieri* Vaillant 1885b in ca. 950 AD) so we do not include them in our list of modern taxa. Palkovacs et al. (2002, 2003) rejected the validity of multiple extant species inhabiting the Indian Ocean Islands based on their analysis of genetic data, recognizing only a single living taxon (*gigantea* or *dussumieri*). Gerlach and Bour (2003) re-emphasized the validity of their recognized species based on the observation that the hatchlings are diagnostic. Further morphologic and genetic research is clearly needed to determine whether more than one taxon of giant tortoise persists on the Indian Ocean Islands. Based on their lack of demonstrable genetic differences, we suggest that these morphologically-defined taxa be listed as subspecies of *gigantea*/*dussumieri* pending further analysis.

09:31. ***Astrochelys yniphora***: Le et al. (2006) proposed the genus name *Angonoka* for the single taxon *yniphora*, but Fritz and Bininda-Emonds (2007) showed that this species is closely related to *radiata* and so placed both species in the genus *Astrochelys* Gray 1873j. The genus name *Angonoka* has not been adopted by other authors and we recommend the use of the genus name *Astrochelys* for both *radiata* and *yniphora*.

09:32. ***Chelonoidis nigra* species complex**: Many recent authors and our previous two checklists have considered the various taxa of Galápagos tortoises as subspecies (e.g., Pritchard 1996; Caccone et al. 1999; Beheregaray et al. 2003; Fritz and Havas 2007; TTWG 2007b; Rhodin et al. 2008). However, previous authors have considered them as full species based on morphology (Bour 1980a; Fritts 1983; Ernst and Barbour 1989) and recently several researchers (Caccone et al. 2002; Russello et al. 2005, 2007; Poulakakis et al. 2008; Chiari et al. 2009) have re-elevated them to species based on congruent patterns of mitochondrial and nuclear variation. Given the allopatric distribution of Gálapagos taxa, combined with the concordant patterns of mitochondrial, nuclear, and morphological variation, we support their recognition as distinct species. In raising these taxa to species, we prefer to highlight their close monophyletic relationship (as distinct from mainland South American *Chelonoidis*) by listing them as a species complex.

09:33. ***Chelonoidis nigra***: The correct epithet for the extinct Floreana tortoise is *nigra* Quoy and Gaimard 1824. Poulakakis et al. (2008) used the epithet *elephantopus* Harlan 1827, but this is in error because that name is younger and the now-lost holotype of *elephantopus* cannot be assigned to any island based on descriptions (Pritchard 1996). Extinct on Floreana since the 1850s, hybrid descendants of this species were recently discovered on Isla Isabela (Poulakakis et al. 2008; Parham 2008) indicating that the lineage persists but has interbred with *becki* Rothschild 1901. Captive cross-breeding of these *becki* x *nigra* hybrids could be used to partially reconstitute the *nigra* lineage.

09:34. ***Chelonoidis abingdonii***: Hybrid descendants of *C. abingdonii* have recently been found on Volcan Wolf on Isla Isabela (Russello et al. 2007), and since only a single male (Lonesome George) of this species survives, the lineage could be partially reconstituted by captive cross-breeding.

09:35. ***Chelonoidis porteri***: Chiari et al. (2009) performed an extensive analysis of morphological, mitochondrial, and nuclear genetic variation in the two separate populations of tortoises on Santa Cruz presently referred to the taxon *porteri*. They demonstrated that the Cerro Fatal population is genetically and morphologically distinct from the La Caseta population and warrants a formal new taxon description, currently in preparation.

09:36. ***Chelonoidis vicina***: Pritchard (1996) previously synonymized *guentheri* Baur 1898 under this taxon (see TTWG 2007b) based on lack of morphological distinctiveness. Recent genetic work by Ciofi et al. (2006) has confirmed a lack of significant genetic distinctiveness between these previously recognized taxa on southern Isabela Island.

09:37. ***Chersina angulata***: Daniels et al. (2007) have demonstrated that this taxon includes two parapatric mitochondrial lineages. These lineages are morphometrically distinct and also show ecological and behavioral differences. Taken together, these data suggest the existence of more than one taxon within *C. angulata* and the matter is under further study (Daniels et al. 2007; Hofmeyr 2009).

09:38. ***Cylindraspis indica***: In our previous checklist (Rhodin et al. 2008) we followed Fritz and Havas (2007) in synonymizing *Testudo tabulata africana* Schweigger 1812 under *Chersina angulata* (Schweigger 1812). However, Bour (1985) previously identified the type specimen of *africana* as being a *Cylindraspis graii*, and Bour (2008a) reaffirmed it as a synonym of *Cylindraspis indica*.

09:39. ***Kinixys belliana nogueyi***: This taxon was treated as a full species by McCord et al. (2005) with minimal argumentation for the change from traditional recognition as a subspecies of *K. belliana*, and we followed that usage in our previous checklist. However, we now agree with Branch (2008) and traditional usage, and therefore restore *nogueyi* to a subspecies of *belliana*.

09:40. ***Testudo* or *Chersine* or *Agrionemys***: The type species and synonymizations of the genera *Chersine* Merrem 1820 and *Medaestia* Wussow 1916 have recently come under discussion. Bour and Ohler (2008) argued that *Testudo graeca* Linnaeus 1758 is the type of *Medaestia* and that *Testudo hermanni* Gmelin 1789 is the type of *Chersine*, whereas Fritz and Kraus (2008) concluded that *hermanni* is the type species for both. In either case, the oldest available generic name for *hermanni* or the clade (*hermanni* + *horsfieldii*) is *Chersine* Merrem, 1820, of which *Eurotestudo* and perhaps *Medaestia* are objective junior synonyms. *Agrionemys* is a subjective junior synonym if *horsfieldii* is considered congeneric with *hermanni* outside the genus *Testudo*, but remains available for a monotypic genus containing *horsfieldii*.

09:41. ***Testudo graeca graeca***: Fritz et al. (2009c) demonstrated that the mitochondrial haplotype of topotypic *T. g. whitei* is identical to samples of *T. g. graeca*, a taxon with overall low genetic variation. They therefore reconfirmed placement of *whitei* Bennett *in* White 1836 in the synonymy of *graeca* Linnaeus 1758.

09:42. ***Testudo graeca marokkensis***: Fritz et al. (2009c) demonstrated that *T. g. lamberti* and *T. g. marokkensis* share the same mitochondrial haplotype. They also questioned the morphometric analyses and proposed geographical separation of *lamberti* and *marokkensis* that were used to justify these taxa, and recommended combining them into a single subspecies. Since both *lamberti* and *marokkensis* were proposed in the same publication (Pieh and Perälä 2004), they invoked the principle of first reviser and chose *marokkensis* as the valid name.

09:43. ***Testudo* or *Agrionemys horsfieldii***: A recent study by Hitschfeld et al. (2008) showed that carpal osteological characters used previously to elevate the subspecies *kazachstanica* and *rustamovi* to species level are ontogenetically variable. In addition, Fritz et al. (2009a) have demonstrated the presence of three major mitochondrial haplotype clades that do not correspond well with the presently understood geographic distribution of the three currently recognized morphologically-defined subspecies. Whether or not to continue to recognize *kazachstanica* and *rustamovi* as distinct subspecies remains uncertain, but we retain them on the list pending further analysis and resolution.

09:44. ***Lissemys punctata***: Rohilla et al. (2009) demonstrated some geographic differentiation in allozymes in this wide-ranging taxon.

09:45. ***Wollumbinia* or *Myuchelys***: Thomson and Georges (2009) described the new genus *Myuchelys* for these taxa (but not including *dorsii* Wells 2009), choosing not to recognize the previous description of *Wollumbinia* Wells 2007c as nomenclaturally available. Whether Wells' work, distributed online without adequate hardcopy dissemination, is nomenclaturally available needs to be decided by the International Commission on Zoological Nomenclature, and we therefore list both names.

09:46. ***Wollumbinia* or *Myuchelys dorsii***: Wells (2009) described the new species *Wollumbinia dorsii*, but whether the name is nomenclaturally available is open to question, as the description was distributed online without adequate hardcopy dissemination. As Australian chelid taxonomy is in a state of flux and the validity of Wells' multiple papers in his Australian Biodiversity Record is

under question, we list the name here. However, we make no determination as to its validity either nomenclaturally or taxonomically.

09:47. ***Wollumbinia* or *Myuchelys latisternum***: Wells (2009) resurrected the taxon *Wollumbinia spinosa* (originally *Euchelymys spinosa* Gray 1871a) as a separate valid species, based on the supposed distinctiveness of the single holotype without known locality data collected in 1866 (as noted by Cann 1998). Whether this resurrection will be accepted as valid is open to question, as no further specimens of the taxon have been identified and its distribution is unknown. We therefore retain *spinosa* as a junior synonym of *latisternum* pending further data.

09:48. **Podocnemididae**: Vargas-Ramírez et al. (2008) performed mitochondrial and nuclear genetic analysis of all eight extant species and demonstrated strong support for the Madagascan genus *Erymnochelys* being sister to a strongly monophyletic South American *Podocnemis*, and the South American *Peltocephalus* being sister to *Erymnochelys* + *Podocnemis*. This phylogenetic analysis renders the occasionally used subfamilial clade name Podocnemidinae Broin 1988 (for *Podocnemis* and *Peltocephalus*) paraphyletic.

09:49. ***Podocnemis unifilis***: As we noted in our previous two checklists (TTWG 2007b; Rhodin et al. 2008), most authors since Troschel 1848 have used the name *unifilis* for this species (the yellow-spotted river turtle), though some early authors erroneously used the epithet *dumeriliana* Schweigger 1812. Recently, the name *unifilis* was referred to the synonymy of *cayennensis* Schweigger 1812 by David (1994), but the latter name has historically been used for what is now recognized as *P. erythrocephala* (the red-headed Amazon River turtle) (Mittermeier and Wilson 1974). Bour (2006a) then redescribed what he concluded to be one of the original three specimens used by Schweigger in his concept of *cayennensis* and designated it as lectotype, but noted that this specimen was actually a representative of the taxon currently known as *unifilis*, and he recommended that the name *cayennensis* therefore be used instead of *unifilis*. However, since the measurements of the lectotype provided by Bour do not correspond exactly with those originally provided by Schweigger, and since Schweigger evidently had examined three specimens for his description, it remains unclear whether the lectotype has been correctly identified. In view of the long history of stable usage of the epithet *unifilis* for the yellow-spotted river turtle, we recommend its continued usage; suppression of *cayennensis* by petition to the ICZN may be needed for nomenclatural stabilization.

2010 Checklist Annotations

TTWG 2010 (000.85-164.checklist.v.3) (10:4-49)

10:4. **Testudines**: In a paper published too close to our manuscript deadline to fully analyze its implications, Dubois and Bour (2010b) discuss the distinction between nomenclature at family-series and class-series rank, and its application to the widely used Order group name Testudines Batsch 1788. Arguing that group names established at a family-series level cannot be applied at a class-series level, and that the family-group name based on the genus *Testudo* is already validly applied at the Family level (as "Testudinidae Batsch 1788"), they conclude that the name Testudines Batsch 1788 cannot also be applied to the Order of turtles, but do not suggest an available name for the Order. However, the International Code of Zoological Nomenclature does not regulate use of names above the superfamily level, and there remain a variety of uncertainties and possible alternative interpretations on the validity, format, use, and authorship attribution of these names. Additionally, some modern authors continue to use and defend the use of the original name Testudines Linnaeus 1758 to designate all modern turtles, even though the name was used primarily in a vernacular fashion in the original publication. Therefore, we do not make any changes at this time, but continue to refer to all turtles as the Order Testudines Batsch 1788, and expect to revisit this issue in more detail in a future checklist.

Additionally, in recent years, the rank level of turtles has been recommended by some to be elevated from its traditional rank of Order within the Class Reptilia to full Class rank on its own (e.g., Collins et al. 2010). Under this scheme, Reptilia would be the Class containing only squamates and tuataras (traditionally known as the Lepidosauria), Class Eusuchia would contain the crocodiles, Mammalia the mammals, and Aves the birds. This possible class-level rank for turtles is to some extent supported by studies indicating the paraphyly of the traditional Class Reptilia with regard to birds. However, other studies indicate a sister-group relationship between turtles and diapsid reptiles, or placement of turtles within diapsids, and hence, uncertainty about the phylogenetic relationship of turtles to other groups abounds (e.g., Laurin and Reisz 1995, deBraga and Rieppel 1997, Kirsch and Mayer 1998, Modesto and Anderson 2004, Bhullar and Bever 2009). Considering that the monophyly of turtles has never been challenged, and that ranking of turtles at class-level provides no improved resolution of the group's phylogenetic position, but simply shifts its distinctiveness to a different rank, we continue to treat turtles as an Order, with no implied judgment of its placement among other living and fossil tetrapod groups.

10:5. ***Caretta caretta***: Dubois and Bour (2010a) noted that Garsault (1764) depicted and named a marine turtle as *Testudo marina*, which they considered a junior synonym of *Caretta caretta* (Linnaeus 1758) based on morphology and geography. *Testudo marina* Garsault 1764 is also a senior homonym of *Testudo marina* Wilhelm 1794, a junior synonym of *Dermochelys coriacea* (Vandellius 1761).

10:6. ***Chrysemys dorsalis* or *picta dorsalis***: Phylogeography of the entire *Chrysemys picta* complex was studied by Starkey et al. (2003), who demonstrated two distinct mtDNA genetic lineages: *dorsalis* and *picta*. They recommended elevating *dorsalis* to species status, but did not find genetic support for the other traditional subspecies (*bellii*, *marginata*), although they recommended that they not be abandoned. Ernst et al. (2006a) documented morphologic intergradation between *dorsalis* and *marginata* in Missouri, but did not express an opinion as to the validity of the proposed elevation of *dorsalis* by Starkey et al., although they referred to their work. Fritz and Havas (2007) suggested that full specific status of *dorsalis* was not fully demonstrated by Starkey et al.'s data and retained it and the other two taxa as subspecies of *picta*. Iverson et al. (2008) agreed with Starkey et al.'s analysis and listed *dorsalis* as a full species, as have other recent authors and database managers (e.g., McAllister et al. 2007, NatureServe), while others retain *dorsalis* at subspecies rank (e.g., Ernst and Lovich 2009, ITIS). The sequencing of the entire *Chrysemys picta* genome is currently in draft form and should help resolve this problem. We now choose to list this taxon provisionally at species rank, recognizing the validity of arguments on either side of the issue, which remains unresolved.

10:7. **Gray 1830e and Gray 1831d**: The date of publication of Gray 1830e (A Synopsis of the Species of the Class Reptilia) is cited by most sources as 1831, since the title page of Griffith and Pidgeon, Volume 9 (Reptilia), in which the Synopsis appears as a Supplement, is dated 1831. However, Gray's Synopsis is dated on its first page as having been written in October 1830, and Volume 9 of Griffith and Pidgeon was actually published in three separate sections from 1830 to 1831 (see Cowan 1969). The first section, Part 25 (pp. 1–192), which included Griffith and Pidgeon's own text on Chelonia (plus part of Sauria), appeared in September 1830 (this part also includes three new Bell and Gray names that we cite as Bell 1830a and Gray 1830c). The second section, Part 26 (part of Sauria plus Ophidia and probably including Gray's Supplement) was apparenty published in December 1830; this date was interpolated by Cowan (1969) as the planned three-month time interval between the publication dates of Parts 25 and 27. The last section, Part 27 (Batrachia plus 18 plates) was published in March 1831. Cowan (1969) did not indicate when Gray's Supplement was published, nor with which Part it appeared. It was certainly not published in Part 25 in September 1830, when only pp. 1–192 of the main text were published, and prior to the October date recorded by Gray on his Synopsis, nor was it published with Part 27 in March 1831 with the Batrachia and plates. The page header for the first few pages of Gray's supplement has "Order Ophidia" printed at the top, and was therefore printed at the same time as the Ophidia section published in Part 26.

In his later publication, Gray 1831d (Synopsis Reptilium; or Short Descriptions of the Species of Reptiles), dated on p. viii as having been written in January 1831, on p. 77 Gray referred to the exact pagination for the citation for *Hydraspis lata* in the earlier published Griffith version (p. 17 in Gray 1830e). Therefore, Gray 1830e was available for page-citation in January 1831, and was therefore published with Part 26, probably in December 1830. Also, in Gray's own listing of his publications (Gray 1873k) he recorded 1830 as the date for publication of his Synopsis in Griffith's Vol. 9. In addition, Cogger et al. (1983) recognized 1830 as the date of publication for this work, as do we.

The date of publication of Gray 1831d has always been considered to be 1831, but the exact date has not been recorded. As noted, it is dated on p. viii as having been written in January 1831. The back cover of the publication lists other books by Gray already available for sale at the same time. Listed as already published are Gray's Illustrations of Indian Zoology, Parts 1 through 6, with a statement that a total of 20 parts were to be completed, one published every three months. Sawyer (1953) recorded that Part 1 of Indian Zoology was published on 6 January 1830, Part 2 on 30 March 1830, Part 3 on 15 July

1830, Part 4 on 6 October 1830, Part 5 on 25 January 1831, Part 6 on 7 April 1831, and Part 7 on 27 July 1831. Gray 1831d was therefore published between Parts 6 and 7, i.e., between April and July 1831, and we have chosen May as the probable month of publication.

10:8. ***Graptemys gibbonsi* and *G. pearlensis***: Ennen et al. (2010) analyzed morphological and genetic variation in *Graptemys gibbonsi* throughout its range and concluded that the western population inhabiting the Pearl River system of eastern Louisiana and western Mississippi is sufficiently distinct to warrant description as a full species, *Graptemys pearlensis*, with *Graptemys gibbonsi* restricted to the Pascagoula River system of eastern Mississippi only.

10:9. ***Pseudemys concinna, floridana*, and *peninsularis***: These three taxa remain difficult to resolve morphologically and genetically, and their taxonomic relationships have vacillated among various views held by Seidel (1994, 1995), Jackson (1995, 2006), and Thomas and Jansen (2006). Our checklist has historically listed *floridana* as a subspecies of *concinna*, and *peninsularis* as a separate species, based primarily on Seidel's work, and we continue to do so. However, recent ongoing fieldwork (e.g., Jensen et al. 2008; M. Aresco and D. Jackson, in litt.) potentially supports recognition of *floridana* and *concinna* as separate species, based on marked differences in their habitat preferences across wide areas of sympatry, with *peninsularis* apparently a subspecies of *floridana*. In view of the long history of taxonomic uncertainty surrounding these taxa and the unresolved nature of the data, we now note these conflicting views by providing alternative listings of *floridana* as either a subspecies of *concinna* or a possible separate species, and *peninsularis* as either a species or possible subspecies of *floridana*. However, we make no actual change in the taxonomic status of these turtles at this time, but await further field data and genetic analyses of this difficult species complex, both of which are ongoing.

10:10. ***Trachemys venusta***: McCord et al. (2010) described three new subspecies of *Trachemys venusta* based on patterns of head and neck stripes, carapace and plastron patterns and coloration, plastral scute formulae, maximum sizes, and distribution. We provisionally list these subspecies pending genetic analysis.

10:11. **Duméril 1805**: This reference has historically been given as 1806, as that date is printed on its frontispiece, but recent work by Gregory (2010), brought to our attention by R. Bour, has shown that it was actually published in November 1805.

10:12. ***Emys*, *Actinemys*, and *Emydoidea***: Wiens et al. (2010) analyzed multiple mitochondrial and nuclear loci for many emydid terminal taxa. They considered that their results did not provide phylogenetic support for the placement of *Actinemys* and *Emydoidea* in *Emys* (as recommended most recently by Spinks and Shaffer 2009); instead, Wiens et al. recommended recognizing *Actinemys* and *Emydoidea* as monotypic genera, with *Actinemys* apparently more closely related to *Clemmys*. We note the discordance among various published data sets regarding the relationships and analyses of the *Emys* + *Emydoidea* + *Actinemys* group, and hence, we retain our previous listings pending greater resolution.

10:13. ***Emys orbicularis orbicularis***: Dubois and Bour (2010a) noted that Garsault (1764) depicted and named a freshwater turtle as *Testudo terrestris*, which they identified as an *Emys orbicularis* (Linnaeus 1758), and which they considered a subjective junior synonym of *E. o. orbicularis* based on geography. *Testudo terrestris* Garsault 1764 is also a senior homonym of *Testudo terrestris* Forskål 1775; however, the latter name has extensive usage over the past half century, and is a *nomen protectum* (ICZN 1963), thus safeguarding its continuing usage.

10:14. ***Emys orbicularis galloitalica***: Pedall et al. (2011) investigated genetic differentiation of populations of Italian, Corsican, Sardinian, and Sicilian *Emys orbicularis* populations, based on mtDNA and polymorphic microsatellite loci. They found no significant divergence of Corsican and Sardinian populations from populations of the southwestern Italian mainland, supporting the view that the subspecies *capolongoi* (Sardinia) and *lanzai* (Corsica) described from these islands are invalid. Their results also suggested that Sardinian and Corsican populations represented reintroduced populations following earlier extirpations of native taxa. While Pedall et al. (2011) did not explicitly synonymize *capolongoi* and *lanzai* into *galloitalica*, they indicated synonymization to be warranted and we consider their results adequate justification to do so. This leaves the status of the subspecies *ingauna*, restricted to a small isolated area in Liguria in the middle of the range of *galloitalica*, unresolved. Fritz and Havas (2007:184) noted that *ingauna* could be synonymous with *galloitalica*, but no data-supported analyses of the status of *ingauna* have apparently been published since its original description; in the absence of evidence to the contrary, we continue to recognize *ingauna* as a valid subspecies.

10:15. ***Emys* or *Actinemys marmorata***: An extensive mitochondrial and nuclear gene study by Spinks et al. (2010) indicated that southern and northern lineages, with a zone of contact somewhere in the central Coast Range of California, exist and may well be diagnosable. However, given the lack of concordance between the traditionally defined ranges of *pallida* in the south and *marmorata* in the north (as previously recognized subspecies), and with two nuclear and four mitochondrial lineages identified, we follow Spinks et al. in waiting for any formal recognition of this variation pending publication of a much larger, ongoing genetic analysis, and we do not re-elevate *pallida* from synonymy at this time.

10:16. ***Glyptemys muhlenbergii***: The name *Emys biguttata* Say was previously recorded as being published in 1824; however, though the paper was read in 1824, it was not published until 1825, in Volume 4, Part 2 of the *Journal of the Academy of Natural Sciences*.

10:17. ***Terrapene***: The genus name *Cistudo* was previously attributed to Say 1825 as a *nomen novum*, as also recorded by Fritz and Havas (2007); however, an Errata sheet published with Say's article in Volume 4, Part 2 of the *Journal of the Academy of Natural Sciences* noted that the name should have been *Cistuda* (as originally published by Fleming 1822). However, many subsequent authors used *Cistudo* as a valid generic name, often citing Say as the original author. The first authors to do so appear to have been Duméril and Bibron (1835), but many others, including Gray (1856b), Agassiz (1857a), and Boulenger (1889) followed suit.

10:18. ***Mauremys japonica***: Hoogmoed et al. (2010) indicated that this species name was actually published in 1834 rather than 1835 as previously recorded by most other authors.

10:19. ***Chelonoidis carbonaria* and *C. denticulata***: Vargas-Ramírez et al. (2010a) investigated phylogeographic differentiation in *carbonaria* and *denticulata*, and found distinct mitochondrial clades in *carbonaria* but only weak differentiation in *denticulata*. They recommended further investigation, but proposed no taxonomic changes at this time.

10:20. ***Chelonoidis carbonaria***: Nowak-Kemp and Fritz (2010) examined the holotype of *Testudo hercules truncata* Gray 1830e in the Bell collection of the Oxford University Museum and determined that it is a *C. carbonaria*, rather than a *C. denticulata*, where it was previously synonymized as a *nomen dubium*.

10:21. ***Chelonoidis chilensis* and *C. petersi***: The validity of the taxa *petersi* and *donosobarrosi* remains subject to debate; the latest contribution to the case was made by Vinke et al. (2008), who considered *petersi* synonymous with *chilensis* based on syntopic occurrence and lack of consistent morphological differentiation of populations attributed to *petersi* vs. *chilensis*. We consider the situation unresolved and tentatively retain our recognition of distinct southern and northern taxa (*chilensis* and *petersi*) within the *chilensis* group, pending further analysis of range-wide patterns of morphological and molecular variation.

10:22. ***Gopherus agassizii***: Cooper's description appeared in print in a section of the *Proceedings of the California Academy of Sciences* that was printed and distributed in 1861, not 1863 when the completed volume (including wrappers dated 1863) was issued (R.B. Murphy, pers. comm.). The two bird species described by Cooper in the same paper (Whitney's Owl, *Athene whitneyi*, now *Micrathene whitneyi*, and Lucy's Warbler, *Helminthophaga luciæ*, now *Vermivora luciae*) are consistently attributed to Cooper 1861 in the ornithological literature.

10:23. ***Homopus***: The date of authorship of the genus *Homopus* has traditionally been given as 1835, referring to Tome 2 of Duméril and Bibron (1835:145). However, Roger Bour (pers. comm.) has drawn our attention to the fact that the name was actually created and diagnosed in Tome 1 of Duméril and Bibron (1834:357).

10:24. ***Homopus signatus***: Daniels et al. (2010) investigated systematics and phylogeography of *Homopus signatus* using mitochondrial and nuclear DNA, neither of which supported the recognition of the two traditional subspecies, *signatus* and *cafer*. As a result they recommended abandoning subspecies designations for *H. signatus*, and we concur.

10:25. ***Homopus signatus***: Our previous checklists and earlier checklists by other authors have consistently listed *Testudo juvencella* Gray 1831d:14 as a junior synonym of this taxon. However, Gray attributed the name to Daudin, and in fact, it was formally described by Daudin in 1802; hence we correct this oversight.

10:26. ***Stigmochelys***: Le et al. (2006) proposed placing *pardalis* in the genus *Psammobates*, while Fritz and Bininda-Emonds (2007) argued for placement in the monotypic genus *Stigmochelys*. Since then, at least three peer-

reviewed publications have consciously (i.e. citing Le et al. 2006, among their references) chosen to use *Stigmochelys*, as did two ecological papers (which did not refer to Le et al. 2006) and the CITES Nomenclature Specialist (CITES 2010); in contrast, only a single peer-reviewed paper (Spinks et al. 2009) was published after 2006 using the combination *Psammobates pardalis*, and this was to name the species as an outgroup. Noting the emerging consensus, we no longer accept *Psammobates* as an alternative name for *Stigmochelys*.

10:27. ***Stigmochelys pardalis***: Fritz et al. (2010a) analyzed phylogeographic patterns in mitochondrial DNA and found that seven, largely parapatric, mtDNA lineages could be identified; these clades did not correspond to the traditional subspecies (*pardalis* vs. *babcocki*) nor to the pronounced geographic size variation. They concluded that there is no rationale for recognizing subspecies within *S. pardalis*.

10:28. ***Testudo graeca buxtoni***: The taxon *Testudo ecaudata* Pallas 1814 has historically been listed in the synonymy of *Testudo graeca*, and Fritz and Havas (2007) and our earlier checklists synonymized it under *T. g. buxtoni* as a *nomen dubium*. Wermuth and Mertens (1961) synonymized it under *T. g. ibera*, and in their later checklist (Wermuth and Mertens 1977) again under *T. g. ibera*, but with a question mark. However, in an overlooked reference, Darevsky and Mertens (1973) examined the unpublished plate from Pallas (1814) depicting the type specimen of *ecaudata*, and determined it to be a variety of the South African *Psammobates tentorius verroxii* (Smith 1839b), despite the fact that the specimen was allegedly obtained in the forests of northern Persia along the Caspian Sea. Despite being a co-author, Mertens was evidently not completely convinced of the synonymization, and only added a question mark to the name in his 1977 checklist, while retaining it under the synonymy of *T. g. ibera*. Because of the uncertainty of the identification of *ecaudata* by Darevsky and Mertens, we solicited input from several specialists regarding the identity of the figured specimen. Examination by Roger Bour, Ernst Baard, Brian Henen, Oguz Türkozan, and Jim Buskirk confirmed that it was not a *Psammobates*, but a juvenile specimen of a *Testudo*, most likely referable to *T. graeca*.

10:29. **Forskål or Forsskål**: The spelling of the family name of Pehr Forsskål has varied through the years, with widespread usage of both single and double 's' spelling (as in our previous checklists). Dubois and Bour (2010a) declared the "Forsskål" spelling to be a mistake, but whether to spell his name with a single or double 's' depends on its usage. In the original paper describing *Testudo terrestris* and *T. triunguis* his name is given on the title page in Latin as Petrus Forskål with one 's', but in his own Swedish vernacular he always spelled his name as either Petter or Pehr Forsskål, with a double 's'. Current references to his name are reasonably evenly split between the two spellings (as determined by a search on Google), with the scientific literature preferring Forskål and the sociohistorical literature preferring Forsskål. We therefore now spell his name as rendered on the original publication, Forskål, just as we render the name Carolus Linnaeus in the original published Latin form rather than the name he was known by in his own Swedish vernacular, Carl von Linné.

10:30. ***Testudo* or *Agrionemys horsfieldii***: Three new taxa from this species complex have recently been described by Chkhikvadze and colleagues: *Agrionemys bogdanovi, A. kazachstanica kuznetzovi*, and *A. kazachstanica terbishi*. In addition, they have elevated two other taxa, *A. horsfieldii rustamovi* and *Testudo baluchiorum*, to species status. The taxonomy of Central Asian steppe tortoises remains in a state of flux; in contrast to the deeply-dissected arrangement of several species and subspecies in *Agrionemys* as proposed by Chkhikvadze and colleagues based on morphological characters, Fritz et al. (2009a) found phylogeographic structure as evidenced by mtDNA to be in weak agreement with morphologically-defined taxa, suggested synonymy of *rustamovi* with *horsfieldii*, and recognized only a single species, placed in *Testudo*, with implicit recognition of subspecies *kazachstanica* and an unnamed ESU in the Fergana valley. To minimize nomenclatural changes in a highly dynamic situation, and to attempt a middle road between the views of Chkhikvadze et al. and Fritz et al., we retain the recent arrangement of a single species with several subspecies, with the newly-described taxa *bogdanovi*, *kuznetzovi*, and *terbishi* provisionally listed as subspecies of *horsfieldii*, keeping *rustamovi* as a subspecies, and not elevating *baluchiorum* from synonymy. We anticipate that further changes will occur in the near future. Additionally, we drop *Chersine* as an alternative generic name for the *horsfieldii* species complex, as all authors use either *Testudo* or *Agrionemys* for this group. This issue was previously discussed in TTWG 2009.

10:31. ***Cycloderma aubryi***: Duméril (1856) described this species under the name *Cryptopodus aubryi*, sp. nov., in his text, but labeled the plate *Cryptopus aubryi*. *Cryptopodus* is considered an *ex errore* name for *Cryptopus* Duméril and Bibron 1835.

10:32. ***Pelodiscus***: Fritz et al. (2010b) carried out preliminary genetic analyses of softshells of the *Pelodiscus sinensis* group, and demonstrated the taxonomic validity and species status of *P. maackii*; further taxonomic research was called for to elucidate the status and proper name of the lineages currently recognized by some as *P. axenaria* and *P. parviformis*, as we continue to do in this checklist.

10:33. ***Rafetus swinhoei***: Le et al. (2010) described a purported new species of giant softshell, *Rafetus vietnamensis*, with the type specimen a complete mounted skeleton located in the Hung Ky pagoda in Hanoi, but without locality data, while at the same time indicating that *R. leloii* Ha 2000 was probably not a taxonomically valid description. The study analyzed mostly the same materials as Le and Pritchard (2009), who reached the opposite conclusion, that Vietnamese records of *Rafetus* all pertain to *R. swinhoei*. A critical re-assessment of the description of *R. vietnamensis* is in progress (M. Le, B. Farkas, pers. comm.). Based on the conclusions of Le and Pritchard (2009), we provisionally list *R. vietnamensis* in the synonymy of *R. swinhoei*.

10:34. ***Chelodina* subgenera**: Georges and Thomson (2010) summarized the history and rationale of grouping species of Australasian snake-necked turtles and naming these groups, and concluded that the preferred nomenclatural arrangement would be to place all species in the genus *Chelodina*, while recognizing three subgenera within *Chelodina*: subgenus *Chelodina* for the narrow-headed species traditionally assigned to 'Group A' related to *C. longicollis*, subgenus *Macrodiremys* for the single species *C. colliei* (= *C. oblonga* of many authors), and subgenus *Macrochelodina* for the broad-headed species of 'Group B' related to *C. expansa*. However, as explained in annotation 10:39, the name *Macrodiremys* may be invalid.

10:35. ***Chelodina (Chelodina) gunaleni***: Georges and Thomson (2010) considered *gunaleni* McCord and Joseph-Ouni 2007a as synonymous with *novaeguineae* Boulenger 1888b, but provided no data supporting their conclusion; we retain *gunaleni* as distinct pending further analysis.

10:36. ***Chelodina (Chelodina) mccordi***: Georges and Thomson (2010) synonymized *Chelodina timorensis* (McCord et al. 2007a) under *C. mccordi* (Rhodin 1994b), without recognizing any subspecies of *mccordi*, instead referring to the taxa *roteensis* and *timorensis* as Diagnosable Terminal Taxa or ESU's (Evolutionarily Significant Units). Kuchling et al. (2007) interpreted these taxa as subspecies of *mccordi*, and we provisionally agree until further work clarifies their taxonomic status. The CITES Nomenclature Specialist (CITES 2010) also synonymized *timorensis* under *mccordi*.

10:37. ***Chelodina (Macrochelodina) rugosa***: Georges and Thomson (2010) reiterated their earlier (Georges and Thomson 2006) synonymy of *C. kuchlingi* Cann 1997d into *C. rugosa* Ogilby 1890, and we now synonymize it as well.

10:38. ***Chelodina (Macrochelodina) walloyarrina***: Georges and Thomson (2010) treated *walloyarrina* McCord and Joseph-Ouni 2007b as synonymous under *burrungandjii* Thomson et al. 2000, referring to morphological and molecular information available in the literature, and their own research indicating hybridization and introgression of *burrungandjii* with *rugosa*. However, we provisionally retain *walloyarrina* as distinct until published molecular data resolves the issue.

10:39. ***Chelodina* (subgenus name undetermined) *oblonga* or *colliei***: Thomson (2006), Georges and Thomson (2010) and Kuchling (2010) summarized the convoluted nomenclatural history of the name *Chelodina oblonga* Gray 1841, and attributed it to the northern populations (currently named *C. rugosa* Ogilby 1890, a name provisionally retained by them) as a senior synonym, pending the outcome of ICZN Case 3351 (Thomson 2006). With the name *oblonga* thus potentially unavailable for the snake-necked turtle of southwestern Western Australia, they referred to this turtle either by its next available name, *Chelodina colliei* Gray 1856a [Georges and Thomson], or *C. oblonga* (= *C. colliei*) [Kuchling] to maintain prevailing usage. In an attempt to retain traditional usage of the name *oblonga* for the southwestern snakeneck, McCord and Joseph-Ouni (2007a) had designated a neotype for *oblonga* Gray 1841, being the lectotype of *C. colliei*. However, as a type specimen already exists for *oblonga*, their neotype designation is invalid (Kuchling 2010).

There is general agreement that the southwestern snakeneck represents a distinct lineage within the *Chelodina* group (Burbidge 1967, Goode 1967, Kuchling 1988, Georges and Adams 1992, Georges et al. 1998, 2002), warranting recognition at subgenus level alongside *Macrochelodina* and *Chelodina* sensu stricto. The genus name *Macrodiremys* was created by McCord and Joseph-Ouni (2007a) to recognize this lineage. They designated *oblonga* as

its type species, in the mistaken belief that their neotype designation had fixed the name *oblonga* to the southwestern taxon. However, with *oblonga* being applicable to the northern taxon (*rugosa*) by virtue of its original valid holotype, *Macrodiremys* becomes a junior synonym of *Macrochelodina*, in contrast to the intent of McCord and Joseph-Ouni. Kuchling (2010), as first reviewer, listed possibilities under which articles of the Code the name *Macrodiremys* could potentially be fixed to the southwestern longneck, but stated his conviction that it was reasonable to await an opinion on case 3351 of the ICZN plenum to be published before such nomenclatural acts should be attempted, and expressed his hope that in their ruling on case 3351, the ICZN plenum would take suitable action to solve these problems. Independently, Georges and Thomson (2010) made an explicit attempt to correct the error and follow the intent of McCord and Joseph-Ouni 2007a under Article A 67.13.1 of the Code by correcting the type species of *Macrodiremys* to *Chelodina oblonga* McCord and Joseph-Ouni 2007a = *C. colliei* Gray 1856a, making it potentially available as a genus-group name for the southwestern snakeneck. However, Kuchling (2010) had already pointed out the possibility that a nomenclatural act such as the attempted correction by Georges and Thomson (2010) could be invalid under the ICZN Code.

Further nomenclatural acts in this group of animals must await the outcome of the anticipated ICZN plenary decision to avoid further complicating this issue. For that reason, we list *Macrodiremys* as a synonym of *Chelodina* sensu lato and we do not employ a subgenus designation for the southwestern species in this year's checklist, recognizing well that the southwestern snakeneck warrants its own subgenus and that a name is potentially available for it.

10:40. ***Elseya novaeguineae***: Georges and Thomson (2010) returned *E. schultzei* Vogt 1911 to the synonymy of *E. novaeguineae* Meyer 1874, and also tentatively placed *E. novaeguineae* in the genus *Myuchelys*. However, we retain the species in *Elseya* pending further genetic analysis.

10:41. ***Emydura m. macquarii***: Georges and Thomson (2010) reviewed the complicated history of the name *Hydraspis australis* Gray 1841, and supported the conclusion of Cogger et al. (1983) that the name is a junior synonym of *E. m. macquarii*. The name *australis* had previously been inconsistently used for a variety of *Emydura* populations in northern Australia (e.g., Cann 1998); Georges and Thomson (2010) instead used the name *victoriae* for some of these populations. We interpret the name *australis* as a *nomen dubium* and agree with its synonymization under *macquarii*.

10:42. ***Emydura m. macquarii***: Georges and Thomson (2010) synonymized the previously recognized subspecies *E. m. binjing*, *E. m. dharra*, *E. m. dharuk*, *E. m. gunabarra*, and *E. signata*, into typical *E. m. macquarii*, based on lack of distinction based on allozyme electrophoresis, and they attributed variation in shell shape and body size to phenotypic plasticity. They considered that the Bellinger River population of *Emydura macquarii* is genetically unremarkable and that it was almost certainly established by introduction of animals from adjacent drainages.

10:43. ***Myuchelys* and Wells taxa**: Thomson and Georges (2009) described the new genus *Myuchelys* and noted that the name *Wollumbinia* Wells 2007c, used for the same group of species, but proposed online in an unpublished web-based document without adequate hardcopy dissemination, is nomenclaturally unavailable. They also considered all web-based names proposed by Wells in his *Australian Biodiversity Record* (Wells 2002, 2007a,b,c, 2009), as unpublished and nomenclaturally unavailable. In previous editions of this checklist we included *Wollumbinia* as an alternative genus name and listed some of the species described in the other documents; however, based on arguments against availability of all these names proposed by Wells (Fritz and Havas 2007, Thomson and Georges 2009, Georges and Thomson 2010), we now concur in considering these names unavailable under the ICZN Code and treat each of them as an unavailable name (*nomen illegitimum*) placed in synonymy.

10:44. ***Phrynops geoffroanus* and *tuberosus***: The taxonomic status and distribution of these two taxa remains problematic, without clear consensus of taxon limits or range delimitations. Pritchard and Trebbau (1984) documented that *tuberosus* is isolated from the broad range of *geoffroanus* and restricted to a small upland area in eastern Venezuela, extreme western Guyana, and northern Roraima in Brazil. Métrailler and Le Gratiet (1996) documented that neither *tuberosus* nor *geoffroanus* occurs in French Guiana, and neither taxon has ever been recorded in Suriname or lowland Guyana. However, McCord et al. (2001), based on an examination of three specimens of what they identified as *tuberosus* from eastern Venezuela, the Brazil–Guyana border, and Piauí, northeastern Brazil, depicted large apparently well-defined allopatric ranges for both taxa, with *tuberosus* supposedly distributed throughout the Guyanan lowlands, the lower Brazilian Amazon, and northeastern Brazil. Rueda-Almonacid et al. (2007), while acknowledging that the systematics of the *Phrynops geoffroanus* complex (Rhodin and Mittermeier 1983) remained controversial and unresolved, nevertheless reproduced the vastly different ranges of *tuberosus* and *geoffroanus* depicted by McCord et al. We choose at this time to instead recognize the earlier documented restricted view of the distribution of *tuberosus*, since it is based on more extensive fieldwork, while also noting that the *P. geoffroanus* complex remains in need of a thorough range-wide phylogeographic study.

10:45. ***Pelomedusa subrufa***: Phylogeographic patterns of mitochondrial and nuclear DNA variation in *Pelomedusa subrufa* were analyzed by Vargas-Ramírez et al. (2010b) and Wong et al. (2010). Both studies reached the same conclusion that the species is structured into three major clades: northwestern, eastern, and southern lineages, with evidence that Madagascan populations may have been introduced there in prehistoric times. Vargas-Ramírez et al. (2010b) suggested that valid taxonomic units may currently be included in the synonymy of *Pelomedusa subrufa*, and recommended further study to clarify the taxonomy.

10:46. ***Podocnemis erythrocephala***: We previously listed the name *Podocnemis agassizi* Coutinho 1868 as a synonym, as did other earlier checklists. However, although Coutinho described a new species of turtle in 1868 that he proposed to name in honor of "M. Agassiz", no formal name was created at that time. Later, Göldi (1886) translated and reprinted Coutinho's earlier writings and created the binomen *Podocnemis agassizii*, attributing it to Coutinho, but recommending that it be changed to *Podocnemis coutinhii* since the name *agassizii* had already been used for another turtle, *Chelonia agassizii*.

10:47. ***Podocnemis sextuberculata***: Coutinho (1868) provided an excellent description of this species that he named *Podocnemis pitiu*, a name overlooked in our previous checklists as well as other earlier checklists.

10:48. ***Podocnemis unifilis***: We continue to use this name for the Yellow-spotted River Turtle, as it has a long and continuous usage, although the name *Emys cayennensis* Schweigger 1812 has recently been shown to have priority based on the available type material (Bour 2006a). However, the name *cayennensis* was for a long time erroneously applied to *Podocnemis erythrocephala* and it would introduce unacceptable confusion to use that name for what has nearly always been referred to as *unifilis*. We therefore maintain prevailing usage of *unifilis* and retain the name *cayennensis* in synonymy as a provisional *nomen rejectum* pending petition to the ICZN (R.C. Vogt et al., in prep.).

10:49. ***Podocnemis unifilis***: Coutinho (1868) provided an excellent description of this species that he named *Podocnemis tracaya*, a name overlooked in our previous checklists as well as other earlier checklists.

2011 Checklist Annotations

TTWG 2011 (000.165-242.checklist.v.4) [(11:5-20)]

11:5. ***Malaclemys terrapin***: Parham et al. (2008) demonstrated, based on radiometric dating of a fossil found in a cave on Bermuda, that the small population of diamondback terrapins present there was the result of a natural dispersal event dating from before human presence on the island. As a result, we now consider the Bermuda population as native rather than introduced. Genetic comparison of Bermuda samples with U.S. populations also demonstrated closest similarity to samples from the Carolinas, so we provisionally list the Bermuda population under the subspecies *M. t. centrata* pending further studies.

11:6. **Genus *Trachemys***: Fritz et al. (2011c) assessed the phylogeny and taxon boundaries of *Trachemys* from Central and South America based on mitochondrial and nuclear DNA. Their nuclear data were largely uninformative, but based on their mitochondrial and combined analyses, they concluded that the Pacific Coast taxa *grayi*, *emolli*, and *panamensis* form a distinct clade that warrants recognition at the species level, as *Trachemys grayi*, with three subspecies. Fritz and co-authors also recommended combining the taxa *cataspila*, *ornata*, *venusta*, *callirostris*, and *chichiriviche* as subspecies under a single species, *Trachemys ornata*. They further proposed placing *adiutrix* at subspecies level in *dorbigni*, and considered the taxon *uhrigi* to be synonymous with typical *venusta*. Their analysis did not include samples of the taxon *iversoni*, whose range on the Yucatan Peninsula is embedded in the range of the redefined *ornata* group, nor did they sample most *Trachemys* taxa from the arid western region of Mexico.

While Fritz et al. (2011c) provide a useful set of hypotheses for future testing, we recognize that taxonomy of *Trachemys* will remain dynamic and expect further changes when additional nuclear data, and deeper geographic

sampling of field-verified specimens, are brought to bear on the problem. We therefore provide the alternative taxonomies of Seidel (2002) and Fritz et al. (2011c) in this year's checklist.

11:7. ***Emys/Emydoidea/Actinemys***: Fritz et al. (2011b) reviewed competing generic concepts for Blanding's, Western Pond, and European Pond turtles, and reiterated their support for recognition of *Emydoidea* and *Actinemys* as separate genera distinct from *Emys*.

The TTWG members continue to be deeply divided in their perspectives on whether to recognize three genera, or a single genus, for the four species concerned (*blandingii, marmorata, orbicularis,* and *trinacris*). The challenge is that available primary research findings result in different relationship trees among the four, based on mitochondrial and nuclear DNA and on morphology (Spinks and Shaffer 2009; Wiens et al. 2010; Fritz et al. 2011b). Overall, the majority of data indicate that *blandingii, marmorata, orbicularis,* and *trinacris* collectively are each others' closest relatives. Whether to recognize this by combining the four species in an expanded genus *Emys*, or to emphasize the morphological distinctiveness of *blandingii* and *marmorata* from (*orbicularis* + *trinacris*) by recognizing *Actinemys* and *Emydoidea* as monotypic genera, is largely a subjective matter. Correspondingly, the herpetological taxonomic community and recent scientific literature have not come to a clear consensus or prevailing usage. With further research in progress, we agree that the situation remains undecided, and continue to present alternative taxonomies in this year's checklist, with no implication that this represents the preferred or supported arrangement of individual TTWG members.

11:8. ***Terrapene carolina***: Butler et al. (2011) conducted morphological and molecular analyses to address the status of lineages within *Terrapene carolina*. They found that box turtles phenotypically corresponding to *T. c. carolina, T. c. bauri,* and *T. c. triunguis* all occur within the range of *T. c. major,* and that the latter does not demonstrate a diagnosable morphology. They also found that *carolina, bauri,* and *triunguis* possess divergent mtDNA haplotypes, which are present alongside a fourth, distinct, haplotype in the range of *T. c. major*. Butler et al. interpreted these findings as the introgressed genetic signal of the extinct Pleistocene *T. c. putnami* perpetuating in a morphologically mixed population, and advocated equating the taxon *major* with *putnami*; they argued that *major,* which has precedence over *putnami,* should only be used to refer to the large extinct form.

Their genetic analysis also placed *bauri* as sister to *ornata,* and *triunguis* as sister taxon to the [*ornata* + *bauri*] clade. Butler et al. thus suggested that *bauri* should be elevated to species status, although this would leave the remaining taxa in *T. carolina* paraphyletic by continued inclusion of *triunguis*. Butler et al. did not address the status of *mexicana* and *yucatana,* which have variously been considered subspecies of *carolina* or full species each. Furthermore, their analysis did not support the previously recognized species groups (e.g., Milstead 1969; Spinks et al. 2009).

We are reluctant to change the widely-recognized taxonomic arrangement for a species complex of notable conservation and legislative significance based on a single study that relied heavily on a short mtDNA segment. Until the alternative taxonomy presented by Butler et al. is independently corroborated by further research, and potentially finds widespread acceptance in the herpetological and taxonomic communities, we prefer to retain the traditional arrangement and defer possible adoption of their hypothesis until a later version of this checklist.

11:9. ***Cuora flavomarginata evelynae***: In previous checklists we had overlooked that Ota et al. (2009) had transferred *evelynae* back to subspecies status under *Cuora flavomarginata*, but we now follow them here.

11:10. ***Aldabrachelys gigantea*** or ***Dipsochelys dussumieri***: Gerlach (2011a) studied morphological development of juveniles of the three forms of Indian ocean giant tortoises reared under identical captive conditions, and concluded that animals consistently develop into the morphotypes characterized by their parents. Gerlach thus concluded that the morphotypes cannot be explained by environment alone, and may have a genetic basis, supporting the recognition of *arnoldi* and *hololissa* as taxonomically distinct from *gigantea/dussumieri*.

11:11. ***Gopherus agassizii* and *G. morafkai***: Murphy et al. (2011) investigated taxonomic problems affecting the Desert Tortoise. They designated the sole remaining of three syntypes of *Gopherus agassizii* as lectotype and genetically confirmed that it originated from California. They also determined that the holotype of *G. lepidocephalus* originated from the Mojave Desert population, at least based on mitochondrial DNA, reconfirming *lepidocephalus* as a junior synonym of *agassizii*. A suite of morphological, molecular and ecological differences between the Mojave and Sonoran Desert populations led Murphy and co-authors to describe the Sonoran form, long recognized as an Evolutionarily Significant Unit, as a new and distinct species, *Gopherus morafkai*. Further research will be necessary to determine if the southernmost populations of *G. morafkai* in the Sinaloan thornscrub ecosystem also deserve taxonomic recognition (Lamb et al. 1989; Murphy et al. 2011).

11:12. ***Testudo graeca* and *T. marginata***: Chkhikvadze et al. (2011) described *Testudo dagestanica* from Lake Papas, Dagestan, and considered the taxa *anamurensis, pallasi, terrestris, weissingeri,* and *zarudnyi* (variously considered subspecies or synonyms of *T. graeca* or *T. marginata* in earlier versions of this checklist) as valid subspecies of *T. marginata*. However, Parham et al. (2006) and Fritz et al. (2007a) placed the Dagestan population of *Testudo* firmly in *T. graeca* based on a combination of morphological and genetic characters, although those studies differed in their assignment of this population to *T. g. ibera* and *T. g. armeniaca*, respectively. Until further data become available supporting the radical changes proposed by Chkhikvadze et al., we take a conservative approach and make no changes to the taxonomy of *T. graeca* or *T. marginata*.

11:13. ***Testudo marginata***: Perez et al. (2012) studied the effects of landscape features and demographic history on the genetic structure of *Testudo marginata* using microsatellites. They found that their samples from Sardinia clustered with samples from northern Greece, suggesting that the Sardinian population may have originated from a small founder population approximately 200 generations ago, while the source population from which those founders originated was estimated to be very large. In contrast, the samples from the 'dwarfed' *marginata* population of the southwestern Peloponnesus demonstrated a low but significant differentiation from all other *marginata* populations. While Perez et al. did not recommend taxonomic recognition of the dwarf population, they did emphasize the conservation significance of what in effect is an Evolutionarily Significant Unit (ESU).

11:14. **Genus *Lissemys***: Praschag et al. (2011) analyzed variability across the range of the genus *Lissemys*, based on 2286 bp of mitochondrial DNA sequences, with additional morphological and biogeographical considerations. They concluded that *scutata* is a distinct divergent lineage and reaffirmed Webb's (1982) conclusion to recognize *scutata* as a full species. They found that the Sri Lankan population shows similar divergence, and recognized this population as a full species, *Lissemys ceylonensis* (Gray 1856a). The remaining populations of *Lissemys* fell into three lineages, one broadly corresponding to the spotted northern taxon *andersoni* (though with some intergradation with unspotted animals in Orissa) inhabiting the Indus-Ganges-Brahmaputra systems, and the other two lineages comprised of unspotted populations in peninsular India. Praschag et al. recommended recognition of the two peninsular lineages by the names *punctata* for the southernmost Indian lineage, and *vittata* for the central Indian lineage. Because the three mainland Indian lineages are more closely related to each other than to either *scutata* or *ceylonensis*, and intergradation between at least *andersoni* and *vittata* is known, Praschag et al. proposed recognition of *vittata* and *andersoni* at subspecies rank under *punctata*. They also concluded that *Emyda granosa intermedia* Annandale 1912, traditionally placed in synonymy of the southern, unspotted form *punctata* (see Webb 1980a:554), is based on intergrades between *vittata* and *andersoni* and cannot be used as the valid name for any *Lissemys* taxon. As *intermedia* is clearly not a valid synonym of *andersoni*, and its type locality (Purulia, western West Bengal) is far outside the redefined range of *L. p. punctata*, we take this opportunity to place *intermedia* into the synonymy of the central unspotted taxon *L. p. vittata*.

11:15. **Genus *Nilssonia***: In earlier versions of this checklist, we used the provisional designation '*Nilssonia* or *Aspideretes*' as a transitional phase between the widespread usage of *Aspideretes* since Meylan's (1987) morphological analysis, and the more recent findings that *Aspideretes* is paraphyletic with regard to *Nilssonia* and consequent recommendation to synonymize *Aspideretes* into *Nilssonia* (Engstrom et al. 2004; Praschag et al. 2007; Fritz and Havas 2007). We note that the species *gangetica, hurum, leithii,* and *nigricans* are now widely accepted as belonging in the genus *Nilssonia*, making *Aspideretes* a synonym.

11:16. **Genus *Pelodiscus***: Yang et al. (2011) evaluated the validity of *P. parviformis* and inferred it to be a distinct species based on the results of their morphological and molecular analyses. Stuckas and Fritz (2011) sequenced DNA from the lectotype of *P. sinensis* and analyzed its placement in relation to other *Pelodiscus*. They found it distinct from the lineages identified as *P. axenaria* and *P. parviformis*, and concluded that *P. sinensis* is not a senior synonym of either of these two names. Consequently, Stuckas and Fritz proposed recognition of *axenaria, maackii, parviformis,* and *sinensis* as valid

species; they recognized that older names may be available for some of these, but recommended use of these four names for the time being. The respective distribution ranges of the different forms, and possible areas of co-occurrence, remain unclear; the reported occurrence of both *axenaria* and *parviformis* in Guangxi and Hunan warrants further research.

11:17. ***Rafetus swinhoei***: Farkas et al. (2011) reviewed the description of *Rafetus vietnamensis* Le et al. 2010. They declared *vietnamensis* an objective synonym of *R. leloii*, and reasserted their view that *R. leloii* (and thus *R. vietnamensis*) is a subjective synonym of *R. swinhoei*.

11:18. ***Trionyx triunguis***: Gidis et al. (2011) sequenced up to seven genes of 20 known-locality samples and reported shallow divergence among Mediterranean Coast, Nile, and Cameroon samples, in contrast to the results of Guçlu et al. (2009), who found four different haplotypes for the four unknown-locality Sub-Saharan specimens that they compared to their Mediterranean samples.

11:19. **Pelomedusidae**: Fritz et al. (2011a) examined the phylogeny of *Pelusios* and *Pelomedusa* species based on three mitochondrial and three nuclear DNA fragments. They reported divergent lineages within *Pelusios rhodesianus* and *P. sinuatus*, found no clear differentiation of *P. chapini* from *P. castaneus*, and attributed the nesting of the sole *P. williamsi* sequence within *P. castaneus* to misidentification of the *williamsi* sample in GenBank. They also demonstrated very shallow divergences within *P. castanoides*, suggesting that populations in Madagascar and the Seychelles were only recently colonized; these findings agreed with those of Silva et al. (2010), who found limited mtDNA differentiation of Seychelles from Madagascar specimens of *P. castanoides*, but cautioned that their sampling of Madagascar material was limited. Fritz et al. (2011a) did not propose explicit changes to taxonomy, but indicated that *chapini* could be re-instated to subspecies rank under *P. castaneus*, and that the recognition of subspecies of *P. castanoides* could be unwarranted; they also suggested the existence of cryptic taxa within *P. rhodesianus* and *P. sinuatus*, and reaffirmed the view of Vargas-Ramírez et al. (2010) that *Pelomedusa subrufa* represents a species complex.

11:20. **Podocnemididae**: Gaffney et al. (2011) analyzed the phylogeny of fossil and living species of Podocnemididae in the context of their earlier analysis of extinct related groups (Gaffney et al. 2006). They reconfirmed the monophyly of the family Podocnemididae, and largely agreed with França and Langer (2006) in not recognizing Erymnochelyinae as a subfamily.

2012 Checklist Annotations

TTWG 2012 (000.243-328.checklist.v.5) (12:6-44)

12:6. **Testudines**: Crawford et al. (2012) and Lourenço et al. (2012) analyzed the placement of turtles in a wider context based on molecular phylogeny, and each team concluded that their data provided strong support for turtles being the sister group to Archosauria (i.e., Crocodilians + Birds, and extinct related groups), rejecting hypothesized relationships of turtles as sister group to Lepidosauria (lizards, snakes, and tuataras), or as the most basal branch of the reptilian (including birds) lineage. Lourenço et al. (2012) estimated the divergence of turtles and archosaurs as dating back to the late Permian around 255 million years ago (MYA), and dated the divergence between Cryptodira and Pleurodira at about 157 MYA in the late Jurassic. Guillon et al. (2012) analyzed mitochondrial and nuclear DNA sequences available in GenBank, including whole genomes for a few species, and constructed a phylogeny for the group involving 230 turtle species representing all families and nearly all genera. Their results reconfirmed the monophyly of Testudines and of Pleurodira and Cryptodira, added support for the placement of *Platysternon* among Testudinoids, placed *Dermatemys* in the Kinosternoids, placed Trionychoids (Trionychidae + Carettochelyidae) as sister to all other Cryptodires (a finding shared with Lourenço et al. [2012] and Wang et al. [2012]), and indicated the need for additional research to better resolve various groups of chelids and testudinids at the genus level.

12:7. **Cheloniidae**: The recognition of higher taxa within the Family Cheloniidae has been inconsistent, at least since Gray's (1825) recognition of an imprecisely defined Carettidae. The inconsistency has been due primarily to the uncertainty regarding the phylogenetic relationships of *Natator* and *Eretmochelys*, and the relationships of fossil to living cheloniids. However, in the midst of the uncertainty of most of the relationships within this family, the close (sister) relationship of *Caretta* and *Lepidochelys* has not been disputed at least since Deraniyagala (1934), who initially argued for the recognition of the latter as the Carettidae, with the remaining taxa in the Cheloniidae. By 1952, Deraniyagala had changed his position to subfamilial recognition of these two groups. Some authors have followed this latter arrangement of recognizing these groups at subfamily rank (e.g., Mlynarski 1976; Pritchard and Trebbau 1984; Smith and Smith 1979), but most have followed Zangerl and Turnbull (1955) and Zangerl (1958) in recognizing these groups as two tribes within the Cheloniidae: Carettini (including *Caretta* and *Lepidochelys*) and Chelonini (including the remaining taxa: *Chelonia*, *Natator*, and *Eretmochelys*).

Impetus for continuing to recognize the Carettini as a tribe (rather than a subfamily) emerges from 1) the strong support for *Caretta* as sister to *Lepidochelys* (virtually every study since Deraniyagala); 2) the controversy about the phylogenetic positions of *Natator* and *Eretmochelys* (i.e., the possible paraphyly of the living Cheloniinae or Chelonini; review in Bowen et al., 1993); and 3) the uncertain phylogenetic relationships of numerous fossil cheloniid taxa, rendering any hierarchy likely to introduce paraphyly (e.g., see Parham and Fastovsky 1998).

Fortunately, four recent publications, by Naro-Maciel et al. (2008; based on two mitochondrial and five nuclear genes), Parham and Pyenson (2010; based on osteology), Duchene et al. (2012; based on the entire mitogenome), and Guillon et al. (2012; based on all available GenBank sequences), seem to reveal the relationships among the living genera quite definitively. These studies clearly resolved *Natator* as sister to *Chelonia*, and *Eretmochelys* as sister to (*Caretta* + *Lepidochelys*), with each of these two clades being reciprocally monophyletic. However, Duchene et al. (2012) explicitly recognized these clades as subfamilies, whereas Naro-Maciel et al. (2008) explicity referred to them as tribes; Parham and Pyenson (2010) defined the tribe Carettini for (*Caretta* + *Lepidochelys*) only, but did not use or define group names for other groups below family level, while Guillon et al. (2012) took no position. In an effort to recognize recent research consensus, to promote stability, and until the relationships among fossil cheloniid taxa are better resolved, we here recognize the two living clades as subfamilies (Carettinae and Cheloniinae), and hope that this will stimulate further research and discussion of the phylogeny of living and extinct cheloniid sea turtles.

12:8. **Cheloniidae**: Vilaça et al. (2012) reviewed the occurrence of natural interspecific hybrids among marine turtle species, and using nuclear markers demonstrated that hybridization among marine turtle species is very common along the Brazilian coast. Most of the hybridization involves male *Eretmochelys* and female *Caretta*, but problematic introgression is occurring among all four genera *Caretta*, *Eretmochelys*, *Lepidochelys*, and *Chelonia*. Vilaça et al. hypothesized that the incidence of this hybridization may have escalated only about 40 years ago, and may be the result of overhunting and local warming of the beaches due to coastal deforestation.

12:9. **Cheloniidae and Dermochelyidae**: Duchene et al. (2012) studied variation across the entire mitochondrial genome of all seven living marine turtle species, and demonstrated divergent intraspecific haplotype clades in the Pacific versus Atlantic and Indian Ocean basins for *Eretmochelys imbricata*, *Chelonia mydas*, and *Dermochelys coriacea*. However, they made no recommendations regarding intraspecific taxonomy.

12:10. ***Chelonia mydas***: Shamblin et al. (2012) demonstrated distinctive mitogenomic haplotype frequencies among the nesting populations of Green Turtles at Buck Island (US Virgin Islands), Aves Island (Venezuela), Suriname, and Tortuguero (Costa Rica), and recommended that these populations receive separate management unit status. However, although these populations are genetically distinct, the authors made no recommendations for taxonomic changes.

12:11. ***Kinosternon subrubrum steindachneri***: Bourque (2012a, 2012b) analyzed the phylogenetic placement of two new fossil *Kinosternon* taxa, and in the course of his morphological analysis found that *Kinosternon subrubrum steindachneri* was placed as sister taxon to the (*K. subrubrum* + *K. baurii*) group; consequently he suggested (2012a) and then elevated (2012b) *steindachneri* to full species rank. Regrettably, he did not specify whether his data for *K. subrubrum* were based on characters of *K. s. subrubrum*, *K. s. hippocrepis*, or a combination of these. The extreme morphological similarity of *baurii* to *hippocrepis* was already noted by Iverson (1992), and we provisionally retain *steindachneri* at subspecies rank under *subrubrum* pending further analysis.

12:12. ***Chrysemys***: Gemel and Grillitsch (2008) reported that Wagler (1821) had nomenclatorally occupied the genus name *Hydrochelys* for the species *Testudo picta* (now *Chrysemys picta*), and explicitly qualified *Hydrochelys* Wagler 1821 as a *nomen oblitum* and invalid, and qualified *Chrysemys* Gray 1844 as *nomen protectum* and valid.

12:13. ***Chrysemys picta***: The description of *Testudo picta* has generally been attributed to Schneider (1783). However, careful reading of his work indicates that the description of *picta* was based on descriptive information contained in letters from Johann Hermann of Strasbourg, and shows no indica-

tion that Schneider had access to the actual specimen, instead adding information from Hermann's letters for the sake of completeness of his (Schneider's) monograph of turtles.

Article 50.1.1 of the Code (ICZN 1999) states "However, if it is clear from the contents that some person other than an author of the work is alone responsible both for the name or act and for satisfying the criteria of availability other than actual publication, then that other person is the author of the name or act." In the case of *picta*, it is not evident how much Schneider was directly quoting from Hermann's writings, so we continue to attribute authorship of the name to Schneider, rather than 'Hermann *in* Schneider 1783'.

As an aside, we note that Hermann used different spellings for his surname over time, involving single or double 'r' and single or double 'n', as well as using French (Jean) and German (Johann or Johannes) versions of his given name. We elect to use the spelling 'Hermann' as that matches the spelling on his death certificate (indicating his preferred spelling later in life), and the spelling employed by Gmelin six years later for the tortoise named for him, *Testudo hermanni*.

12:14. ***Graptemys***: Freedman and Myers (2012) identified and sequenced a species-specific mitochondrial control region marker and two nuclear markers in a population of *Graptemys pseudogeographica* that had hybridized historically with *G. geographica*, after which geological events prevented further inbreeding for several generations. Both mitochondrial and nuclear introgression was documented. They provided additional evidence for hybridization events among *G. pseudogeographica*, *G. ouachitensis*, and *G. geographica* in other parts of the range, and combined with literature records of *Graptemys* hybridization events in captivity, concluded that reproductive isolation in *Graptemys* has evolved incompletely. In the context of conflicting evidence for monophyly vs. paraphyly of *G. pseudogeographica* and *G. ouachitensis*, they suggested that the two groups may maintain reproductive isolation in parts of their range while interbreeding in other areas.

12:15. ***Graptemys ouachitensis***: Brown et al. (2012) examined mitochondrial DNA control region sequence variation throughout the range of the subspecies *ouachitensis*, and found 18 haplotypes forming two minimally divergent groups. They also found modest divergence between *o. ouachitensis* and *o. sabinensis*, and slightly greater divergence between *ouachitensis* and *pseudogeographica*. They acknowledged earlier studies suggesting that *sabinensis* might warrant species status, and noted the need for additional studies to clarify the relationship among these taxa.

12:16. ***Pseudemys***: Using mitochondrial DNA, Jackson et al. (2012) inferred phylogenetic relationships of the taxa within *Pseudemys*. While recovering the genus as a strongly supported monophyletic group, they found no support for monophyly of the traditionally recognized redbelly (*rubiventris*) and cooter (*concinna*) species groups. They also did not find great support for monophyly of individual taxa other than *gorzugi* and *texana*, suggesting mitochondrial introgression since the glacial retreat in the Pleistocene, or recent speciation, as possible explanations. They made no specific recommendations for taxonomic changes based on their results, and indeed cautioned against making taxonomic changes within this highly complex group without comprehensive data.

12:17. ***Trachemys taylori***: McGaugh (2012) analyzed population genetic structure of *Trachemys taylori*, including a detailed genetic assessment of potential hybridization, and found significant genetic differentiation among populations, consistent with population bottlenecks, but detected no genetic evidence for hybridization of *T. taylori* with invasive *T. scripta elegans*.

12:18. ***Trachemys venusta***: Perry (1810) published a description of *Testudo panama*, a name overlooked and/or ignored ever since, until Bauer and Petit (2004) discussed it. They attributed it to the genus *Trachemys* based on the illustration and text description, and based on provenance and color pattern, regarded *Testudo panama* as referring to *Trachemys venusta* (Gray 1856b). Invoking ICZN Article 23.9, Bauer and Petit (2004) asserted that *Testudo panama* Perry 1810 was to be regarded as a *nomen oblitum* and invalid, and *Emys venusta* Gray 1856b as a *nomen protectum* whenever the two are considered as synonyms.

12:19. ***Emys orbicularis hellenica***: Schreiber 1875, in his section on *Cistudo lutaria*, listed and defined eight varieties, including var. h), to which he attributed "*Emys Hoffmanni* Fitzinger Mus. Vindob." Our previous inclusion in earlier checklists of *Emys orbicularis hoffmani* [sic] Schreiber 1875 as a synonym of *E. o. hellenica* referred in fact to a subsequent use of Fitzinger's (1835) name *Emys (Emys) hofmanni* (with a minor spelling difference) rather than to a new taxon description, and as such we now delete it from our listing of synonymized primary taxa.

12:20. ***Terrapene ornata***: Joyce et al. (2012) described a new fossil box turtle and carried out a phylogenetic analysis of extant and fossil box turtle species based on morphological and osteological characters. The fossil taxon *T. longinsulae* Hay 1908, from an undefined location in the general region of Long Island, Kansas (possible age ranging from Miocene to Pleistocene; see Joyce et al. 2012:185), scored identical for all available characters to the character states shown by both living *T. ornata ornata* and *T. o. luteola*, including two unique shared characters. They therefore placed *T. longinsulae* in the synonymy of *T. ornata*.

12:21. **Geoemydidae**: The deep divergence of living New World from Old World geoemydids has been well-established based on mitochondrial and nuclear DNA data (Spinks et al. 2004; Le and McCord 2008; Guillon et al. 2012), and has recently been dated at 57 mya (Lourenço et al. 2012). Le and McCord (2008) first proposed the recognition of living New World taxa (genus *Rhinoclemmys*) as the subfamily Rhinoclemminae. Given that the divergence of the latter is as old as, or older than, that of the widely accepted emydid subfamilies (dated at 37 mya by Dornburg et al. 2011; 57 mya by Lourenço et al. 2012), we support the subfamilial status of the genus *Rhinoclemmys*. However, we amend Le and McCord's name to Rhinoclemmydinae to correspond to the proper case for its ending *-emys* (genitive *emydis*), and we note that the group may also include North American fossil species of the genera *Echmatemys* (Hervet 2004) and *Bridgeremys* (Hutchison 2006). Correspondingly, we apply the subfamily name Geoemydinae to all Eurasian geoemydid taxa.

12:22. ***Cuora***: Phylogeny and species boundaries were studied for the genus *Cuora* by Spinks et al. (2012a) using a combination of mitochondrial and nuclear genes. While noting discrepancies between the results of the mitochondrial and nuclear data sets, concordance with traditional, morphology-based phylogenies for *Cuora* was greatest with the nuclear dataset, which they considered to represent the most reliable estimates of phylogeny and species boundaries. The species *amboinensis*, *flavomarginata*, *mouhotii*, and *yunnanensis* emerged clearly resolved, whereas the *trifasciata* and *galbinifrons* taxon clusters were less well resolved. *Cuora aurocapitata* and *C. pani* showed signs of introgression of *trifasciata* mtDNA, but were reciprocally monophyletic based on nuclear markers. The position of animals attributed to '*cyclornata*' was highly variable depending on what markers were analyzed, a result that was interpreted as being most consistent with introgression, and leading to the recommendation to consider animals with '*cyclornata*' phenotype as part of *C. trifasciata* rather than a distinct separate taxon. *Cuora zhoui* emerged as a strongly divergent species based on nuclear DNA, but its mtDNA association with *trifasciata* may indicate mitochondrial introgression. The monophyly of *C. mccordi* was well supported, though its phylogenetic position was different when nuclear and mtDNA datasets were analyzed. While some doubts were expressed about the species status of *mccordi*, the authors recommended continued treatment as a valid species for the time being. The members of the *galbinifrons* group (*bourreti*, *galbinifrons*, and *picturata*) were strongly supported as a clade of three genetically and morphologically diagnosable taxa, and while some potential gene flow was detected between these taxa, the authors recommended that they continue to be recognized as valid separate species.

12:23. ***Cuora amboinensis***: Ernst et al. (2011) evaluated the potential to use shell morphology and postorbital stripe to separate the subspecies *amboinensis* and *kamaroma* in the Philippines, as alternative approaches to the usual plastron pattern character used to differentiate these subspecies. They found that shell shape and postorbital stripe data do not necessarily correspond to plastron pattern, suggesting that animals previously identified as *kamaroma*, and as such representing a potential co-occurrence of two different subspecies in some areas of the Philippines (i.e., Palawan and Sulu), may in effect be *amboinensis* animals with a plastral pattern resembling that of *kamaroma* through introgression or phenotypic plasticity.

12:24. ***Cuora galbinifrons***: Bourret described this species in a work dated 1939 on the title page, but it was not actually published until 1940 (R. Bour, pers. comm.). We have therefore changed the date of publication from 1939 to 1940.

12:25. ***Heosemys spinosa***: Spinks et al. (2012b) examined variation in mitochondrial (cytochrome b) and nuclear (11 loci) DNA in a large sample of *Heosemys spinosa* that lacked locality data. Two clades were clearly identified by the mtDNA analysis, and were supported by preliminary morphological analysis, but not recovered in the nuDNA analysis. Future identification of the geographic provenance of these two clades is critical for conservation management as well as establishing the taxonomic implications of this work.

12:26. ***Mauremys caspica***: Vamberger et al. (2013) investigated population structure and history of *Mauremys caspica* based on an analysis of 14

microsatellite loci and cytochrome b mtDNA sequences from nearly range-wide samples (but lacking Iraq). Their results found two clusters of mitochondrial haplotypes, and four microsatellite clusters, with each mtDNA haplotype cluster comprising two of the microsatellite clusters. Specimens from Bahrain and Saudi Arabia were found to constitute a distinct microsatellite cluster, and were thus viewed as representing native (and endangered) populations. The authors proposed that each of the four identified phylogenetic clusters (i.e., Central Anatolia; eastern Turkey and Syria; Dagestan, Azerbaijan, and Iran; and Bahrain and Saudi Arabia) be treated as distinct management units. Their results did not support the validity of any of the three previously morphologically-defined subspecies, and they therefore proposed that the subspecies no longer be recognized, a recommendation that we follow here.

12:27. ***Orlitia borneensis***: Palupcikova et al. (2012) analyzed mitochondrial (cytochrome b) and nuclear (R35) sequences, and shell and scute morphometrics among *Orlitia borneensis* specimens in European collections. Most specimens lacked precise locality data because they originated from a single confiscation in 2001, but three known-locality specimens from Borneo and Sumatra were added to the sample series (no reference specimens from West Malaysia were available). Haplotype diversity in cytochrome b was found to be relatively high, with three main haplotype groups identified; nucleotide diversity was low, and phylogenetic structure was poorly supported. The three known-origin animals clustered within one of the main haplotype groups, suggesting that the confiscated animals covered much of the species' genetic diversity. Only minimal variation was found in R35 sequences. Geometric morphometrics demonstrated morphological similarity of all examined specimens. These results led the authors to conclude that all examined animals represented a single conservation unit.

12:28. ***Aldabrachelys gigantea* or *Dipsochelys dussumieri***: The latest contribution to the ongoing debate in the ICZN (case #3463) on the scientific name for the Aldabra Tortoise came from Ceríaco and Bour (2012), who traced the history of the specimen claimed to be the type of *Testudo gigantea* Schweigger from the MNHN Paris via the defunct Royal Cabinet of Natural History of Ajuda in Lisbon to the collections made in Brazil and other parts of South America by Alexandre Rodrigues Ferreira during his travels there during 1783–1792. The authors thus provided further evidence that Schweigger's type specimen of *Testudo gigantea* originated in Brazil and represents a *Chelonoidis denticulata*, providing support for possibly precluding the use of the name *gigantea* for the Aldabra Tortoise. At the time of writing of this annotation, the ICZN is voting regarding the use of either *gigantea* or *dussumieri*, and a result is expected to be published in the next issue of the Bulletin of Zoological Nomenclature.

12:29. ***Geochelone* or *Centrochelys sulcata***: Bour and Henkel (2012) described a captive colony of tortoises of different species maintained in extensive enclosures in southern France and documented a female *Centrochelys sulcata* producing viable, morphologically aberrant, hatchlings that were interpreted as hybrids between this sulcata female and a male *Dipsochelys dussumieri* [or *Aldabrachelys gigantea*], with which copulation had been observed. From each of three clutches of 20 eggs each, normal *sulcata* hatched from 15 eggs after 75–90 days, one hatchling emerged after 140–170 days and was a suspected *sulcata-dussumieri* hybrid, and 4 eggs failed to develop. The authors also reported the repeated production of hybrid hatchlings between *C. sulcata* and *Astrochelys radiata* at the A Cupulatta facility in Corsica; these either died just before emergence from the egg, or within days after hatching. These records add to the growing catalog of documented intergeneric turtle hybrids (see also annotation 12:8 above).

12:30. ***Chelonoidis chilensis***: The number of species in the *Chelonoidis chilensis* complex recognized by various authors has ranged from one to three (see TTWG 2010, annotation 10:21). However, based on an examination of a mitochondrial gene (cytochrome b) and 10 microsatellite loci, Fritz et al. (2012a) found negligible genetic variation among populations and concluded that this complex represents only a single species, *C. chilensis*. Previously described morphological variation among populations is apparently explained by Bergmann's Rule, with body size increasing with latitude. Hence, *C. petersi* and *C. donosobarrosi* were placed in the synonymy of *C. chilensis*.

12:31. ***Chelonoidis nigra* species complex**: Only two papers published in the past year bear on the taxonomy of the Galapagos tortoises, and both continue to recognize the various taxa at the species level (see TTWG 2009, annotation 09:32). Garrick et al. (2012) reported the discovery of hybrid tortoises on Volcan Wolf that were F1 hybrids between a purebred Floreana tortoise (*C. nigra*; erroneously referred to in their paper as *C. elephantopus*) and the local Volcan Wolf tortoise (*C. becki*). This led the authors to speculate that translocated Floreana tortoises might still exist on northern Isabela. Actual rediscovery of genetically pure individuals of Floreana tortoises would reverse their current IUCN status as Extinct, and be cause for great celebration.

Based on DNA sequences from three mitochondrial genes from extant and museum specimens of Galapagos tortoises, Poulakakis et al. (2012) attempted to resolve the phylogenetic relationships and reconstruct the biogeographic history of the living and extinct taxa. Prior to their study, they recognized eleven extant and four extinct species of Galapagos tortoises, though one of the latter has not yet been described. It should be noted that the taxa they recognized a priori did not completely agree with those recognized by the TTWG. In any case, the results of Poulakakis et al. (2012) suggested that the following populations represent "independent evolutionary units" and they applied the term "species" to them: *abingdonii* (extinct in 2012, Pinta), *becki* (Volcan Wolf, northern Isabela), *chathamensis* (San Cristobal), *darwini* (Santiago), *elephantopus* (extinct, Floreana; recognized as *nigra* by the TTWG), *ephippium* (Pinzón; recognized as *duncanensis* by the TTWG), *hoodensis* (Española), *porteri* (La Reserva, Santa Cruz), and *vicina* (central and southern Isabela, and including the names *microphyes*, *guentheri*, and *vandenburghi*; matching previous TTWG checklists), an unnamed extant species (Cerro Fatal, Santa Cruz), and an unnamed extinct species (Santa Fe). They explicitly noted that two taxa were not likely to be independent evolutionary units: *wallacei* (Rabida) being subsumed under *vicina* (as already reflected in previous TTWG checklists), and *phantastica* (extinct, Fernandina; possibly introduced by humans) under *porteri*. However, in the face of pronounced morphological differences between the extreme saddleback *phantastica* and the greatly domed *porteri*, the TTWG believes that additional support must be presented before synonymization of *phantastica* is warranted. Finally, while Poulakakis et al. (2012) laudably extracted DNA from bones of museum specimens from extinct lineages, it is critical that the genotyping of actual type specimens of all possible named Galapagos tortoise taxa (including those of synonyms) be completed before final allocations of names to existing populations can be done with full confidence.

12:32. ***Gopherus flavomarginatus***: Ureña-Aranda and Espinosa de los Monteros (2012) examined variation in a mitochondrial gene in the Bolson Tortoise and not only found no geographic structure in that variation, but also noted that the existing variation was the lowest ever reported for a tortoise. They attributed this low variability to the post-Pliocene collapse of this once more widely distributed (New Mexico to central Mexico) species.

12:33. ***Gopherus polyphemus***: Based on a single mitochondrial gene fragment (ND4), Ennen et al. (2012) examined geographic variation in Gopher Tortoises and found two major haplotype assemblages that overlapped in distribution in the Appalachicola-Chattahoochee River basin. They also found some (albeit weak) support for the distinction of the Federally Threatened portion of the western assemblage to the west of the Mobile River Basin (USFWS 1987c). In a more comprehensive study of both mitochondrial (cytochrome b) and nuclear microsatellite markers, Clostio et al. (2012) confirmed the Appalachicola basin as the transition region between the distinctive western and eastern lineages. In addition, based on both mtDNA and nuDNA, they noted the distinction of the populations west of the Mobile River and those in western Georgia. They concluded that the tortoises in each of these four regions should be managed independently, but they made no specific taxonomic recommendations.

12:34. ***Gopherus* species indeterminate:** *Testudo australis* Girard 1858 was described based on a specimen reputedly originating from New Zealand. Clearly this specimen must have been transported by humans, as no testudinids or other non-marine turtles are known from New Zealand, living or fossil. The name has generally been overlooked or ignored, except by Boulenger (1889) who attributed it with doubt to *Gopherus polyphemus*. Based on communication with Robert Murphy and Steve Gotte, the specimen cannot be found in the USNM collection, if indeed it ever was there. Girard's description of a uniform near-black tortoise agrees nearly perfectly with *Gopherus*, except for the small nuchal (cervical) scute, which is usually wide in *Gopherus*. The scutellation on the top of the head and on the forearms, the absence of thigh spurs, as well as the shape of the caudal (supracaudal) scute are all characters that eliminate *Manouria*. *Chelonoidis* species are eliminated from consideration by the presence of a nuchal in *australis*. Therefore, we tentatively attribute *Testudo australis* Girard 1858 to the genus *Gopherus*. Attribution to any particular species is challenging: based on size and coloration, *G. berlandieri* and *G. flavomarginatus* are easily excluded, and while the Pacific location of New Zealand suggests possible seafaring or trade links with California and therefore *G. agassizii* and/or *G.*

morafkai, morphological details such as the large head with rounded snout, and upward curving of the gulars, are more reminiscent of *G. polyphemus*. We therefore include *Testudo australis* Girard 1858 as a *nomen dubium* and *nomen oblitum* under *Gopherus* species indeterminate.

12:35. ***Kinixys***: Kindler et al. (2012) examined the phylogeography, phylogeny, and taxonomy of all currently recognized *Kinixys* taxa, based on examination of sequence data for three mtDNA fragments (12S, ND4, cytochrome b) and three nuclear loci (C-mos, ODC, R35). Their findings indicated that the savannah taxa, traditionally recognized as subspecies or affiliated species of *K. belliana*, represent three deeply divergent clades, which are paraphyletic with respect to the rainforest species *K. erosa* and *K. homeana*.

To reflect this phylogeny, the authors recognized their East African cluster as a valid species, for which they followed the first reviser (Bour 1979) to apply the name *zombensis* in preference over *zuluensis*. They also elevated *nogueyi* to full species status, and reconfirmed *lobatsiana*, *natalensis*, and *spekii* as evolutionarily distinct and ranked as full species.

They did not, however, present a clear arrangement of recognized taxa; their new delineation of species is presented in their Fig. 2, implying that no subspecies are recognized. They noted that their samples of *domerguei* from Madagascar clustered with *zombensis*, and it appears that they intended to transfer *domerguei* from the synonymy of *K. b. belliana* (following Broadley 1992, 1993; Fritz and Havas 2007; Branch 2008) to that of *K. zombensis*. In their text (Kindler et al. 2012:198), the five-clawed tortoises of the Central African Republic are specifically combined with the four-clawed West African animals to form the distinct species *K. nogueyi*; however, these Central African Republic records are mapped (Fig. S1) and listed (Table S1) as '*belliana belliana*' in the online supporting material.

While the TTWG generally believes that the results presented by Kindler et al. (2012) represent major advances in our understanding of *Kinixys* phylogeny, we prefer to retain a slightly more traditional arrangement for *domerguei*, pending further data. While *domerguei* may not be genetically recognizable based on the examined genes, it is morphologically well established (e.g., Bour 2006e), and we continue to recognize it as a valid taxon at subspecies rank under *zombensis* (to which it appears most closely related). Hence, we recognize the new combinations *Kinixys zombensis zombensis* and *K. z. domerguei*.

12:36. ***Testudo graeca***: Parham et al. (2012) extracted mtDNA sequence data from Iranian tortoises of the *T. graeca* complex, specifically the holotype of *T. g. zarudnyi* and topotypes of *T. g. buxtoni* and *T. g. perses*. Their results confirmed the previous work of Fritz et al. (2007a, 2009) regarding the existence of two distinctive mitochondrial haplotypes in Iran, one in the northwest, and one in eastern and central regions.

Using AFLPs (amplified fragment length polymorphisms) from samples across the range of the *T. graeca* species complex, Mikulíček et al. (2013) identified four geographically defined genetic groups: 1) western Mediterranean (Morocco and Spain to Libya); 2) Balkans-Middle East (Bulgaria and Romania to southwestern Russia, Azerbaijan, eastern Turkey, and southern Israel); 3) Caucasian (extreme eastern Turkey, Russia, and southeastern Azerbaijan to northwestern Iran); and 4) central and eastern Iran. These groups generally correspond to the mtDNA haplotype lineages identified by Fritz et al. (2007a, 2009), but with one to four mitochondrial lineages per AFLP group.

Considered together, these two data sets suggest that the western Mediterranean and central-eastern Iran groups are both divergent and allopatric, and thus could be considered as distinct species. The oldest available names for these groups are *T. graeca* (including *cyrenaica*, *lamberti*, *marokkensis*, *nabuelensis*, *soussensis*, and *whitei*) and *T. zarudnyi*, respectively. The remaining two groups come into contact in the eastern Caucasus, with some mtDNA evidence of gene flow between them. Hence, there remains uncertainty as to whether they represent distinct evolutionary units worthy of species status. The oldest available names for those two groups are *T. terrestris* (Balkans-Middle East) and *T. buxtoni* (Caucasus; including *armeniaca*, *perses*, and *pallasi*). Unfortunately, variation in neither set of markers corresponds well with described morphological variation in this complex, on which the traditional taxonomy has been based.

Pending future studies of variation in nuclear markers and a re-examination of morphological variation to determine minimally plastic characters that correspond to the currently defined genetic groups (i.e., dismissing color and general body size and shape), we prefer to retain *T. graeca* as a single species, with no changes in the subspecies recognized from our previous checklist.

12:37. ***Apalone spinifera***: McGaugh (2012) examined variation at ten microsatellite loci for populations of *Apalone spinifera* within and outside of the Cuatro Ciénegas basin in Mexico. She found considerable divergence among all sampled populations (particularly eastern versus western basin localities), but found no genetic variation associated with carapacial color variation within the basin (the basis for the original description of *A. s. atra*). She reported negligible differentiation between softshells sampled inside the basin compared to those outside, but provided no indication whether additional *A. s. emoryi* were sampled since the range-wide analysis of *A. spinifera* reported by McGaugh et al. (2008) and did not provide a detailed comparison between *Apalone* within the Cuatro Cienegas basin versus outside the basin. As McGaugh did not make an explicit taxonomic recommendation to synonymize *atra* with *emoryi*, we conservatively continue to recognize *atra* as before, at subspecies level.

12:38. ***Nilssonia gangetica***: An analysis of mitochondrial and nuclear DNA sequence data by Liebing et al. (2012) confirmed the monophyly of the genus, resolved the placement of *N. formosa* as sister to other *Nilssonia*, and identified significant intraspecific genetic variation within *N. gangetica* corresponding to river basin of origin. Populations from the Brahmaputra, Mahanadi, and combined Indus and Ganges basins were each genetically distinct, and worthy of separate management. The authors did not support species recognition of the three identified units, but noted that if these units were given subspecies status, the name *mahanaddica* (Annandale 1912b) is available for the Mahanadi basin population, and the nominate trinomial would apply to the Indus-Ganges population. No name has been applied to the Brahmaputra population. Until the three intra-specific populations are adequately characterized and named, we continue to recognize *N. gangetica* as monotypic.

12:39. **Chelidae**: It was assumed by Gaffney (1977) that the three long-necked taxa in the Chelidae (*Chelodina*, *Hydromedusa* and *Chelus*) formed a monophyletic lineage. However, Pritchard (1984) proposed that these three taxa were not necessarily closely related, based on the major structural differences in how they arrived at their long-necked condition, a position generally consistent with the phylogenetic arrangement previously proposed by Baur (1893a). Sequencing of 12S mtDNA was used to demonstrate that the Australian radiation of the Chelidae formed a monophyletic group (Seddon et al. 1997; Georges et al. 1998) and that the shorter-necked South American taxa (including *Chelus*) were also a monophyletic group, with *Hydromedusa* a third lineage (Georges et al. 1998). At that time Georges et al. (1998) proposed names for the three lineages; Chelinae [as Chelidinae] for *Chelus* and its South American relatives; Chelodininae for the Australasian species; and Hydromedusinae for the genus *Hydromedusa*, a clade that also includes the fossil genus *Yaminuechelys* (de la Fuente et al. 2001) and hence is not monotypic. Further morphological work agreed with Pritchard (1984) and supported the molecular results with differences in skull, shell, and cervical structures among the three monophyletic lineages (Thomson 2000a), indicating that what gross similarity there is between the long-necked forms was a result of convergence and not of shared ancestry. In the ten or so years since these publications, numerous studies have added further support to this phylogenetic arrangement, which was summarized and further supported by Guillon et al. (2012). Therefore we now include these three subfamily assignments in this edition of the checklist.

We are aware of the family-group name Hydraspidina Bonaparte 1836, based on the genus *Hydraspis* Bell 1828, whose type species is *Testudo* [now *Chelodina*] *longicollis*, as a potential source for the subfamily name for chelids of the Australia-New Guinea region. However, as Bonaparte's concept of *Hydraspis* differed fundamentally from that of Bell, to the point of excluding *longicollis* from *Hydraspis* by properly placing it in *Chelodina* Fitzinger 1826, revival of the family-group name Hydraspidina (as Hydraspidinae for this subfamily) would lead to unnecessary confusion. In the absence of strict ICZN priority requirements for family-group names, we therefore use the name Chelodininae Baur 1893b for the chelids of the Australia-New Guinea region.

12:40. ***Mesoclemmys dahli* and *M. zuliae***: Based on both mitochondrial and nuclear gene sequence variation, Vargas-Ramírez et al. (2012a) confirmed the species distinction of *Mesoclemmys dahli* and *M. zuliae*, and identified two geographically isolated, weakly divergent populations within *M. dahli* (Córdoba and Cesar Departments, Colombia). Further sampling will be necessary to determine if this divergence has taxonomic implications.

12:41. ***Mesoclemmys heliostemma* and *M. raniceps***: Because some authors (e.g., Rueda-Almonacid et al. 2007) have questioned whether *M. heliostemma* is distinct from *M. raniceps*, Molina et al. (2012) undertook a multivariate morphometric study of those two taxa (and *M. gibba*). Their results supported the validity of *M. heliostemma*, and identified eight new localities that significantly expand the known range of the species across the Amazon basin.

12:42. ***Pelusios castanoides***: Analysis by Fritz et al. (2012c) of mito-

chondrial genes of *P. castanoides* revealed that samples from Madagascar and the Seychelles were weakly differentiated from each other, and significantly different from the sampled mainland populations. However, sparse sampling from the mainland precluded any final conclusions about the origins of the Malagasy and Seychelles populations, and therefore the validity of the currently named, endemic subspecies of the Seychelles. As the population of Madagascar is related more closely to that of the Seychelles than to the continental African populations, a subspecies designation for these combined populations might be warranted. The oldest available name is *kapika* Bour 1979; however, we defer any changes to our checklist at this time.

12:43. ***Pelusios subniger***: Parallel analysis by Fritz et al. (2012c) of mtDNA samples of *Pelusios subniger* from mainland Africa, Madagascar, and the Seychelles demonstrated the existence of a cryptic, unnamed taxon from the Democratic Republic of the Congo, but no significant variation among all other sampled populations. Based on these results, the authors suggested that *subniger* was introduced by humans to both Madagascar and the Seychelles, and recommended that the supposed endemic Seychelles subspecies *P. s. parietalis* be placed into the synonymy of a monotypic *P. subniger*. However, reanalysis of the morphological differentiation that originally distinguished these taxa has not been carried out, and we are reluctant to make a definitive change at this time, particularly for a taxon with such significant regional conservation concerns. We await further analysis before amending this taxonomy.

12:44. ***Podocnemis lewyana***: Based on 10 microsatellite loci and a mitochondrial DNA fragment, Vargas-Ramírez et al. (2012b) found low genetic variability across the range of *Podocnemis lewyana*, consistent with a significant historic population bottleneck. They did identify three weakly differentiated genetic management units: 1) the Upper Magdalena River Basin; 2) the Lower Magdalena, Lower Cauca, and San Jorge Basins; and 3) the Sinú River Basin, but made no taxonomic recommendations.

2014 Checklist Annotations

TTWG 2014 (000.329-479.checklist.v.7) (14:1-50)

14:1. [*Annotation text previously included here now deleted from this checklist*.]

14:2. ***Caretta caretta***: Considerable confusion has surrounded the names and authors and dates of publication of the turtle descriptions published in various outputs of the *Expédition Scientifique de Morée*, currently cited in our checklist as Valenciennes (1833), Bibron and Bory de Saint-Vincent (1833), and Bory de Saint-Vincent (1835). See the detailed clarification below in annotation 14:25 for *Emys orbicularis hellenica*. The name *Chelonia pelasgorum* was first published on plate 6 by Valenciennes (1833), but rendered as *C. pelasgica* in the subsequent text by Bibron and Bory de Saint-Vincent (1833), where they synonymized it with *Chelonia caouanna* (= *Caretta caretta*).

14:3. ***Eretmochelys***: In last year's checklist, in annotation 13:7, we outlined our reasoning for recognizing the subfamilies Cheloniinae and Carettinae. Unfortunately, we made an editorial error in listing *Eretmochelys* under the Cheloniinae when, in fact, studies have shown that it is more closely related to *Caretta* and *Lepidochelys*, and belongs in the Carettinae, as we pointed out in our annotation. We correct the error in this year's checklist.

14:4. ***Eretmochelys imbricata***: The junior synonym *Chelonia grisea* Eschscholtz 1829b has for many years been incorrectly cited as *Chelonia griseam* in our previous checklists and in Fritz and Havaš (2007) and extensively on the web. Having finally successfully accessed the obscure original publication, we now note this long-standing error and correct it.

14:5. ***Chelonia mydas***: We note also that in the same obscure publication cited above, Eschscholtz (1829b) also described *Chelonia castanea* from Surinam as a new species. The name has been overlooked since its description and is a *nomen oblitum* and junior synonym of *C. mydas*.

14:6. ***Dermochelys coriacea***: The original citation for the junior synonym *Dermatochelys porcata* is actually Wagler (1830b), not Wagler (1833) as listed in our previous checklists, but which contained no new turtle descriptions.

14:7. ***Dermochelys coriacea***: In our previous checklists we had included the name *Testudo marina* Wilhelm 1794 in the synonymy of *D. coriacea*, based on its inclusion in older checklists (e.g., Fritz and Havaš 2007). However, examination of Wilhelm's (1794) work indicates that his use of the name *Testudo marina* was as an incorrect collective group name for "marine" species (sea turtles and softshells), as he also grouped most "terrestrial" turtles (testudinids and kinosternids) under the incorrect group name *Testudo terrestris*, and all other freshwater turtles under the group name *Testudo fluviatilis*. In discussing separate species under these group names, he used names previously described by other authors (including a description of the Leatherback, using the name *Testudo coriacea*), but not in a consistently binomial manner. His work therefore has no standing nomenclaturally, and we have removed the name *Testudo marina* Wilhelm 1794 from the synonymy of *D. coriacea*.

On the other hand, Ranzani (1832) published a description in Latin of the Leatherback Turtle in which he described it as *Testudine coriacea marina*. This description is valid (as the trinomen *Testudo coriacea marina*), as per ICZN Code Article 11(h)(ii) allowing for the use of adjectival Latin descriptions, as previously noted by Smith and Rhodin (1986) in regard to the validity of the original authorship of *Testudo coriacea* Vandellius 1761.

14:8. ***Dermatemys mawii***: González-Porter et al. (2013) presented microsatellite data that supported their previous mitochondrial DNA studies (González-Porter et al. 2011) in recognizing populations of *Dermatemys mawii* in the Papaloapan River drainage as genetically distinctive. However, they made no taxonomic recommendations based on their results. In addition, they also identified a small sample of genetically divergent individuals in the Sarstun and Salinas River basins along the southeastern distribution of the species that they speculated might represent a cryptic taxon.

14:9. **Kinosternidae**: Iverson et al. (2013) sequenced three mtDNA and three nuclear markers for every recognized species and most subspecies of kinosternids. Their analyses revealed three well-resolved clades within the Kinosterninae, corresponding to *Sternotherus*, a previously unnamed clade that they described as the new genus *Cryptochelys*, and *Kinosternon* sensu stricto. Their molecular data support for *Cryptochelys* was strong, but data support for non-monophyly of *Kinosternon* with respect to *Sternotherus* was weak. The identified groups are broadly consistent with morphological and biogeographical features. Their new genus *Cryptochelys* was diagnosed based on an extensive set of morphological and molecular characters, and contains the designated type species *leucostoma*, as well as *acuta*, *angustipons*, *creaseri*, *dunni*, and *herrerai*.

As we are aware of a parallel study of kinosternid phylogenetics, currently in review, that reaches different taxonomic conclusions, we present the recommended taxonomy of Iverson et al. (2013) as an additional alternative to the traditional arrangement, in the knowledge that we will revisit kinosternid taxonomy again in our next edition, and hopefully come to a consensus position then.

14:10. ***Kinosternon abaxillare***: A multivariate analysis of morphometric data by Berry (1978) demonstrated the distinctiveness of the endemic, allopatric taxon *K. scorpioides abaxillare* from the parapatric *K. s. cruentatum*. In addition, preliminary molecular sampling of the *K. scorpioides* complex by Iverson et al. (2013) suggested that *K. s. abaxillare* was more closely related to *K. oaxacae* than to *K. s. cruentatum* (or any other *K. scorpioides*). Given both the morphometric and molecular evidence, the latter authors followed Alvarez del Toro (1972, among many others) and suggested that *K. abaxillare* be recognized as a full species. Until more thorough geographic and molecular sampling is completed, we acknowledge both options in this checklist, but treat the taxon as more likely a species.

14:11. ***Kinosternon chimalhuaca***: In 1996, while the original, full description by Berry et al. (1997) of *K. chimalhuaca* was in press, it was shared with Manfred Rogner for inclusion in his forthcoming book. However, though unintended, Rogner's abbreviated version, clearly attributed to Berry et al., was published first (in 1996). Hence, although many authors have cited Berry et al. (1997) as the original description for this taxon, the proper attribution should be Berry, Seidel, and Iverson *in* Rogner (1996). The ICZN has now been petitioned (Rogner et al. 2013) to officially confirm this proper authorship and date, which has already been used in all previous TTWG checklists.

14:12. ***Kinosternon subrubrum steindachneri***: Based on osteology, Bourque (2012) recommended that *K. s. steindachneri* be elevated to full species status, as it was originally described. Preliminary molecular data provided by Iverson et al. (2013) supported this conclusion. However, until a more complete, range-wide study of molecular and morphological variation of the *K. subrubrum-baurii* complex is available, we here retain *steindachneri* as a subspecies of *K. subrubrum*.

14:13. **Staurotypinae or Staurotypidae**: Highlighting the extensive divergence of the staurotypines from the kinosternines based on morphology (Hutchison 1991), genetics (Iverson et al. 2103), karyotype (Bickham and Carr 1983), and sex determination mechanisms (Ewert et al. 2004), Iverson et al. (2013) followed Bickham and Carr (1983) in recognizing the Staurotypidae as a separate family. Within the TTWG we have differing opinions on the

appropriate ranking of this taxonomic node, and recognize that the views and actions of the wider turtle taxonomic community will determine its eventual accepted ranking; until consensus emerges, we provide alternative rankings in the checklist.

14:14. ***Graptemys***: In a historical review of the taxonomic history of the genus *Graptemys*, Lindeman (2013:20) mentioned two genus names from an unpublished manuscript by Georg Baur: *Neoclemmys* (intended to include *pseudogeographica* and *oculifera*) and *Megaloclemmys* (for *pulchra*), while *Graptemys* would have been retained for *geographica* and *kohnii*. However, Lindeman only used these names in a conditional manner, without formal status as valid taxa, and hence the names *Neoclemmys* and *Megaloclemmys* were not made available according to Article 15.1 of the International Code of Zoological Nomenclature (see also annotation 14:19 for *Graptemys intermedia*).

14:15. ***Graptemys flavimaculata***: Using microsatellite loci, Selman et al. (2013) demonstrated a significant degree of genetic structure across the range of the species in the Pascagoula River basin, with the greatest divergence between the main Pascagoula basin and the lowland Escatawpa River tributary, historically separate drainages. Although they urged that at least these two units be managed separately, they made no taxonomic recommendations.

14:16. ***Graptemys ouachitensis* or *G. o. ouachitensis***: See annotation 14:19 below regarding the taxon *sabinensis*, previously listed as a subspecies of *ouachitensis*, but now conditionally elevated to full species status, therefore also necessitating listing the Ouachita Map Turtle as a full species, rather than the nominate subspecies.

14:17. ***Graptemys pulchra***: In his historical review of the taxonomic history of *Graptemys*, Lindeman (2013:20) also made reference to Baur's manuscript names for the species he eventually described as *Graptemys pulchra*; we hereby designate these names, *G. alabamensis* and *G. grandis*, as *nomina nuda*, and associate them with the synonymy of *G. pulchra*, as they were considered for application to that taxon.

14:18. ***Graptemys sabinensis* or *G. o. sabinensis***: Originally described as a subspecies of *Graptemys pseudogeographica* by Cagle (1953a), the Sabine Map Turtle (*G. sabinensis*) was later classified by Vogt (1980) as a subspecies of *G. ouachitensis*, and most subsequent authors have followed that arrangement. However, based on a small sample of skulls, Ward (1980) believed that *sabinensis* was so distinctive that it warranted species status. Recent analyses of morphology, mitochondrial DNA, and nuclear DNA have generally failed to resolve the relationships of *sabinensis* with confidence, and the interrelationships of the "narrow-headed *Graptemys*" remain largely unresolved (Stephens 1998; Stephens and Wiens 2003; Myers 2008; Wiens et al. 2010; Brown et al. 2012). Based on these previous studies and his own extensive examinations of Gulf Coast *Graptemys* specimens, Lindeman (2013) noted that *sabinensis* is allopatric, non-intergrading, and diagnosable morphologically, and concluded that it should be recognized as a full species. While subspecific as well as specific recognition can each be supported, we conclude that enough uncertainty remains regarding this lineage to list it as either a species or subspecies. Further sampling of the nuclear genome and more strongly supported phylogenetic trees will be necessary to settle this issue as well as the relationships across the entire genus *Graptemys*.

14:19. ***Graptemys sabinensis* or *G. o. sabinensis*:** In a checklist of turtles of Louisiana, Beyer (1900) listed "*Malacoclemmys intermedia* Baur" from the "southern and southwestern parts", referring to a manuscript name by Georg Baur for the taxon subsequently described as *Graptemys pseudogeographica sabinensis* by Cagle (1953a). The name was based on specimens from the Mermentau River basin, now in the Tulane University Museum collection, sent to Baur by Joseph Gustave Kohn (Lindeman 2013). The original publication of the name *Malacoclemmys intermedia* is therefore attributable to Beyer and pre-dates the name *sabinensis* Cagle by 53 years, but is clearly a *nomen nudum*. Furthermore, it has not been used in over 113 years and would have the status of *nomen oblitum*. In his discussion of the history of the name *Graptemys intermedia* used by Baur in his unpublished manuscript, Lindeman (2013) published Baur's original manuscript drawings (Fig. 2.4) and a photograph of the Kohn specimens on which Baur had intended his diagnosis to rest (Fig. 8.18). Lindeman also clearly identified *intermedia* as a synonym of the taxon he recognized by the name of *G. sabinensis*. However, Lindeman only, and consistently, used the name *G. intermedia* in a conditional manner, without formal status of valid taxon, and the name *intermedia* as used by Lindeman, published after 1960, has not been made available according to Article 15.1 of the International Code of Zoological Nomenclature.

14:20. ***Pseudemys***: Based on three mitochondrial and ten nuclear gene loci, Spinks et al. (2013) examined variation across all recognized taxa of the genus *Pseudemys*. Their analyses revealed essentially no support for currently recognized species groups, species, or subspecies. Only *P. gorzugi* was consistently recovered as monophyletic across all their analyses, while their molecular evidence identified three geographically cohesive groups that do not correspond to current species boundaries. They concluded that the genus *Pseudemys* has probably been oversplit taxonomically. However, they made no explicit recommendations for change until a much larger and more definitive, multi-character data set is brought to bear on this complex. Thus we retain *Pseudemys* essentially unchanged from the previous checklist.

14:21. ***Trachemys***: In their book on Mexican turtles, Legler and Vogt (2013) continued to follow the taxonomy for *Trachemys* as used by Legler (1990) (i.e., all Mexican taxa as subspecies of *T. scripta*), and did not provide data or rationale to refute the phylogenetic data and taxonomic opinions published since 1990. Because implementing that taxonomy here would reverse 23 years of increased understanding and progress toward a stable classification of this complex genus, we have not incorporated the *Trachemys* taxonomy presented by Legler and Vogt (2013) in our current checklist.

14:22. ***Trachemys* (Caribbean)**: Parham et al. (2013) examined variation in mitochondrial and nuclear DNA markers for *Trachemys* populations across the Greater Antilles. They identified the morphologically distinct population reported by Tuberville et al. (2005) in northwestern Jamaica as *T. d. decussata*, representing a significant range extension from eastern Cuba. Parham et al. (2013) demonstrated the monophyly of West Indian taxa, as well as evidence of hybridization between *T. decorata* and *T. stejnegeri* in the southern Dominican Republic, and between *T. terrapen* and *T. d. decussata* in northwestern Jamaica. The authors were unable to determine whether the presence of *decussata* on Jamaica and localized hybridization with *terrapen* was the result of natural or human-mediated dispersal.

Their data also supported the continued recognition of the subspecies *T. s. stejnegeri* (Puerto Rico) and *T. s. vicina* (Hispaniola), with occasional gene flow (natural or human-mediated) between them. They also acknowledged that morphological and genetic data suggest the recognition of *T. d. decussata* (eastern Cuba) and *T. d. angusta* (western Cuba) as full species, but declined to make that recommendation pending further sampling in Cuba.

Finally, they speculated that the occurrence of *T. d. angusta* on the Cayman Islands was "non-native". Given that Grand Cayman was periodically inundated even in the latest Pleistocene (20–25 thousand years ago; Iturralde-Vinent 2006), and that the prevailing winds and currents would make a natural colonization from the northwest difficult, we concur that the Grand Cayman populations of *T. d. angusta* are likely the result of human introduction. This is further supported by Echternacht et al. (2011) who, in their review of the herpetofauna of the Cayman Islands, explicitly stated that since no *Trachemys* fossils have been found in peat deposits on the island (which contained many other vertebrates), they presumed *T. decussata* to be introduced.

In addition, Parham et al. (2013) analyzed a small sample of *Trachemys* from Central America that yielded results indicating genetic similarity of *T. venusta* and *T. emolli*, demonstrating the need for further sampling and analysis to evaluate the sweeping taxonomic changes proposed by Fritz et al. (2012) and the subspecies described by McCord et al. (2010). Until such additional information becomes available, Parham et al. (2013) recommended taxonomic conservatism and cautious interpretation of preliminary results, and proposed no taxonomic changes.

14:23. ***Trachemys* (Central America)**: Using mitochondrial DNA sequence data for *Trachemys* downloaded from the European Nucleotide Archive and new data from two Honduran specimens, McCranie et al. (2013) confirmed that the range of the taxon *emolli* extends from northwestern Costa Rica to southeastern El Salvador (see Ibarra Portillo et al. 2009). Because of the extensive overlap in their analysis with the data used by Fritz et al. (2012), McCranie et al. (2013) supported their earlier taxonomic recommendations. McCranie et al. (2013) also commented on the status of the taxon *T. v. uhrigi*, originally described from Honduras, but subsequently reported from Colombia, Costa Rica, Nicaragua, and Panama, and argued that the diagnostic coloration of *uhrigi* is not exhibited consistently by Caribbean Honduran specimens (and presumably occurs in individuals of *T. venusta* as far away as Colombia), and suggested that *T. v. uhrigi* has no taxonomic validity. In contrast, Páez et al. (2012) listed *T. v. uhrigi* as being the subspecies occurring in the Colombian Departments of Antioquia and Chocó. However, until additional molecular data are forthcoming from the Caribbean versant of Central America, particularly from the Yucatan peninsula, Honduras, Nicaragua, Costa Rica, Panama, and

Colombia, we adhere mainly to the alternative taxonomies of Seidel (2002) and Fritz et al. (2012).

14:24. ***Emys* (sensu lato)**: In a molecular analysis of emydine turtles, Angielczyk and Feldman (2013) found strong support for a monophyletic *Emys* (including *orbicularis*, *blandingii*, and *marmorata*) using mtDNA sequence data, but strong support for a paraphyletic *Emys* using 14 nuclear genes. The combined data set resolved a monophyletic *Emys*, but the results were apparently driven by the much more variable mitochondrial genome. Despite some uncertainty about the monophyly of *Emys* sensu lato (see also Wiens et al. 2010), we retain both options of a narrow and a broad definition of the genus *Emys*, pending even more genetic data.

14:25. ***Emys orbicularis hellenica* and *Mauremys rivulata***: Considerable confusion has surrounded the names and authors and dates of publication of the turtle descriptions published in various outputs of the *Expédition Scientifique de Morée*, currently cited in our checklist as Valenciennes (1833), Bibron and Bory de Saint-Vincent (1833), and Bory de Saint-Vincent (1835). Sherborn and Woodward (1901) documented that the zoology sections dealing with vertebrates, in which turtle descriptions (*Chelonia pelasgorum*, *Emys hellenica*, *Emys iberica*, and *Emys rivulata*) appear, were all first published sequentially in 1833 in looseleaf "*livraisons*" with sets of plates ("*planches*") and text, and later all the plates were re-published as a bound volume in 1835. Confusion about the sequence of publication of the turtle plates and the text has arisen due to the imprint of "1832" on the frontispiece of the text, but it actually appeared in 1833, after the plates. That the unbound turtle plates by Valenciennes (1833:pls.6–9) were published first was clearly documented in the subsequent text by Bibron and Bory de Saint-Vincent (1833:61, lines 5–9, 21–23, footnote 2), who referred to the specifically numbered plates as coming from the "troisième séries".

The name *Emys hellenica* was first published in the third series of *planches* (plate 8, figures 2–2a), where it was attributed to Valenciennes in the legend, and then subsequently described (as *Cistuda hellenica*) on pages 61–62 of the text, where it was attributed to Bibron and Bory de Saint-Vincent. Proper original attribution of the name is therefore *Emys hellenica* Valenciennes *in* Bory de Saint-Vincent 1833:pl.8. That name was subsequently synonymized, as "Cistude hellénique", with *Cistudo europaea*, another synonym of *Emys orbicularis*, by Duméril and Bibron (1835:227), but is today recognized as a valid subspecies of that taxon (Fritz et al. 2005; Fritz and Havaš 2007).

According to Bibron and Bory de Saint-Vincent (1833), in their text on page 61, lines 5–9, under the synonymy of *Cistuda europaea* (= *Emys orbicularis*), they noted that Valenciennes (1833) had "for unknown reasons" ("*on ne sait pourquoi*") named the juvenile specimen of *Emys* on plate 9 as *iberica* [not *Emys iberica* Eichwald 1831]. The original typeface in the legend of plate 9 reads "*Emyde ibérienne. Emys iberica. Val.*"; however, in at least some contemporary copies of the subsequently bound atlas (Bory de Saint-Vincent 1835), a small printed label in similar text reading "*des anciens*" has been pasted over "*ibérienne*", and a second label reading "*antiquorum*" pasted over "*iberica*" and the first part of "*Val.*" We do not know if these labels were originally added as an "erratum" to all looseleaf copies at the time of their original publication in 1833, or more likely only to some of them when bound into the atlas in 1835, because some copies today lack the labels (e.g., that in the Paris Museum, but apparently not those in the British Museum [Gray 1844:31] or the Museum of Comparative Zoology at Harvard [Loveridge and Williams 1957:213]). It must also be noted that Bibron and Bory de Saint-Vincent (1833) commented on the name *iberica* printed on the original plate, but made no mention of the name *antiquorum*. This uncertainty greatly complicates these names and their authorship. If the "*antiquorum*" labels were added to all copies of this work by the publisher, then *Emys iberica* Valenciennes was technically never described, and *Emys antiquorum* would presumably be attributable to Bory de Saint-Vincent, the editor of the 1835 atlas, although the remaining partial exposure of the name "*Val.*" led Gray (1844:31) to attribute the name *antiquorum* to Valenciennes when he (Gray) synonymized that name with *Cistudo europaea* [= *Emys orbicularis*]. If the labels were inconsistently added to only some copies of the original work, then the name *antiquorum* would have no nomenclatural status. Pending the availability of additional historic information about the consistency of this labeling and the reasons behind it, we here attribute the name *Emys iberica* to Valenciennes *in* Bory de Saint-Vincent (1833), and declare *Emys antiquorum* Bory de Saint-Vincent (1835) a *nomen novum* and *nomen nudum*; and interpret both names as junior synonyms of *Emys orbicularis hellenica* (Valenciennes *in* Bory de Saint-Vincent 1833).

14:26. ***Emys orbicularis*, ssp. indet.**: Rook et al. (2013) synonymized the two Upper Pliocene fossil taxa, *E. major* and *E. latens*, described by Portis (1890), with *Emys orbicularis*, citing an unpublished thesis by Chesi (2009). The fossils were from Valdarno (d'Arno valley) along the northwest Italian Ligurian coast, within the range of the present-day subspecies *E. o. galloitalica*. However, given the subsequent Pleistocene and Holocene climate-associated range shifts of *E. orbicularis*, we do not associate these fossil names with any current subspecies at this time.

14:27. ***Terrapene***: Martin et al. (2103) examined variation in two mitochondrial genes and one nuclear gene across all previously recognized taxa of *Terrapene* (except *T. nelsoni klauberi*). Both mtDNA and a single nuclear gene supported the monophyly of *T. ornata*, *T. carolina* (including *T. coahuila*), and *T. nelsoni*. All analyses confirmed the distinctiveness of *T. nelsoni* and *T. ornata*, but found no support for distinction between *T. o. ornata* and *T. o. luteola*, and hence, they recommended the synonymy of the latter. For mtDNA only, they found significant divergence within the previously recognized *T. carolina* group taxa (including *T. coahuila*), and identified a western clade (including *triunguis*, *mexicana*, and *yucatana*) and an eastern clade (all others, including *coahuila*); the western clade was strongly supported based on mtDNA, but the eastern clade had only very weak support. The authors recommended the recognition of the western clade as a full species, *T. mexicana*, with three subspecies (*mexicana*, *triunguis*, and *yucatana*). They were not able to resolve the relationships among *bauri*, *major*, and *carolina*, and retained them as subspecies of *T. carolina*. Similarly, the relationship of *T. coahuila* to Gulf Coast *T. carolina* was suggested but not resolved, and they recommended continued recognition of *T. coahuila* at the species level. Meanwhile, Legler and Vogt (2013) treated *T. mexicana* and *T. yucatana* as full, monotypic species, and continued to recognize *T. o. luteola* as the subspecies of *T. ornata* inhabiting Mexico.

As the suggested taxonomic rearrangements of Martin et al. (2013) and Legler and Vogt (2013) are not based on a comprehensive analysis of morphology, mitochondrial DNA, and nuclear genes, and show disagreement with both the traditional taxonomy of *Terrapene* and the molecular phylogeny presented by Butler et al. (2011), we consider that the phylogenetic relationships within this genus remain insufficiently resolved for us to adopt such significant taxonomic changes, especially in light of the desire for minimal fluctuations in taxonomy of this genus, given its extensive inclusion in State, Federal, and CITES legislation.

14:28. ***Terrapene putnami***: This species was described by Hay (1906) based on a single fossil hypoplastron from the Alafia River basin in Florida with imprecise stratigraphic data, but presumably Late Pleistocene. The taxon has been widely assumed to represent virtually all eastern North American fossil *Terrapene* material from the Miocene to the late Pleistocene, which is clearly an exaggerated concept of the taxon (Ehret et al. 2011). This has become increasingly problematic because of recent molecular analyses of extant taxa that suggested that *T. putnami* should be synonymized with *T. carolina major* (Butler et al. 2011; followed in TTWG 2012) or that argued that *putnami* be retained as an extinct subspecies of *T. carolina* (Martin et al. 2013). To facilitate future attempts to resolve the relationships among living and fossil turtles of the *T. carolina* complex, Ehret et al. (2013) proposed to the ICZN the designation of a neotype for *T. putnami* with precise locality and stratigraphic data, and consisting of a nearly complete carapace and plastron with numerous associated postcranial elements.

We here maintain *T. putnami* as a synonym of *T. carolina major* as recommended by Butler et al. (2011), until the ICZN makes a ruling and additional research clarifies the relationships of extant and fossil members of the *T. carolina* group.

14:29. ***Mauremys annamensis***: *Clemmys guangxiensis* was described by Qin (1992) based on two market specimens supposedly originating from Guangxi, China. Iverson and McCord (1994) speculated that the type series might be a composite of *Mauremys mutica* and *M. iversoni* (the latter now known to be of hybrid origin between *Cuora trifasciata* and *M. mutica*; Parham et al. 2001). As a result, we have previously included *C. guangxiensis* in the synonymies of both *C. trifasciata* and *M. mutica*. However, Hu et al. (2013) provided sequence data from a single mitochondrial gene for four specimens of *Mauremys* from Guangxi purported to be *M. guangxiensis*, along with two *M. mutica* from the same province. When included in a phylogenetic analysis with sequences of *Mauremys* downloaded from GenBank, they determined that their four specimens of *M. guangxiensis* were nearly identical to *M. annamensis* (which is endemic to Vietnam) and not *M. mutica*. Assuming their four specimens represented the same taxon as described by Qin (1992), Hu et

al. (2013) interpreted their results as indicating that *M. guangxiensis* was either synonymous with *M. annamensis* or a subspecies of the latter. However, their analysis did not address the possibility of a hybrid origin for *guangxiensis*. Pending further study of specimens being referred to *M. guangxiensis*, we add *guangxiensis* to the synonymy of *annamensis* (as partim, hybrid), while retaining its inclusion in the synonymy of *M. mutica* and *C. trifasciata*.

14:30. ***Mauremys japonica* and *Pelodiscus sinensis***: Temminck and Schlegel's publications in Fauna Japonica are usually recorded as having been published in 1835. However, their chapter on "Les Cheloniens" (pp. 1–80, plates 1–9) was actually published in 1834 (see Hoogmoed et al. 2010) and only contained invalid vernacular names. Their valid names *Trionyx japonica* = *Pelodiscus sinensis* and *Emys vulgaris japonica* = *Mauremys japonica* were not published until 1838 when Schlegel wrote and published (on p. 139) his dated explanation of the previously published plates and for the first time provided Latin names for the two new species described earlier in French (*Trionyx stellatus* Var. Japon [pls. 5 and 7] = *Trionyx japonica* and *Emys palustris* Var. Japon [pls. 8 and 9] = *Emys vulgaris japonica*). Although the species name "*palustris*" was used on the plate, in the text it was corrected to "*vulgaris*", but never with a specific "var. Japon" modifier attached to it.

14:31. ***Rhinoclemmys***: Based on both mitochondrial and nuclear DNA data, Vargas-Ramírez et al. (2103) identified significant phylogeographic structuring within *R. melanosterna*, but found conflicting phylogenetic relationships among the allopatric/parapatric members of the *R. punctularia* group (including *R. funerea, R. diademata*, and *R. melanosterna*). They recommended no taxonomic changes without further geographic and genome sampling.

14:32. ***Aldabrachelys gigantea***: After several years of vigorous debate, the ICZN (2013b) published their decision (Opinion 2316) regarding the appropriate scientific name for the Aldabra Tortoise (Case 3463; Frazier 2008, 2009). The Commission ruled to conserve the long-term use of the specific name *Testudo gigantea* Schweigger (1812) for this tortoise, to affirm the neotype designation of Frazier (2006), and to suppress the more recently used name *Testudo dussumieri* Gray (1831d). One effect of this action was also to validate the genus name *Aldabrachelys* Loveridge and Williams (1957) over *Dipsochelys* Bour (1982a). Comments were published in BZN 66:80–87, 169–186, 274–290, 352–357; 67: 71–90, 170–178, 246–254, 319–331; 68: 72–77, 140–143, 294–300. With 83 published comments, this represented the most extensive correspondence received by the Commission on a Case to date.

14:33. ***Chelonoidis carbonaria***: The original citation for the junior synonym *Testudo boiei* is actually Wagler (1830a), not Wagler (1833) as listed in our previous checklists; the latter contained no new turtle descriptions.

14:34. ***Gopherus berlandieri***: In 1850, Berlandier described two terrestrial turtles from the "llanos" of Tamaulipas, Mexico: *Testudo bicolor* (not to be confused with *Terrapene bicolor* Bell 1826, synonym for *Testudo* [= *Cuora*] *amboinensis* Daudin 1801) and *Testudo tuberculatu* [sic] (not to be confused with *Testudo tuberculata* Pennant *in* Schoepff 1801 [= *Dermochelys coriacea*]). His ample description leaves little doubt that the names referred to a juvenile and adult male, respectively, of *Xerobates* [= *Gopherus*] *berlandieri* Agassiz 1857a, and hence should be considered senior subjective synonyms of the latter. However, since 1850, *T. tuberculatu* has only been mentioned by True (1882), as *T. tuberculata*. In 1980 Berlandier's manuscript was translated and republished, with both species recorded again, as *T. bicolor* and *T. tuberculata*, with distinct diagnostic characters, "and they are common on both banks of the Rio Bravo." No other publications seem to include these names, both considered here as being *nomina oblita*.

14:35. ***Testudo graeca***: Based strictly on morphology Chkhikvadze et al. (2013) continued to argue for the recognition of six taxa of tortoises in the Caucasus (*Testudo graeca ibera, T. g. nikolskii, T. g. armeniaca, T. marginata pallasi, T. m. buxtoni*, and *T. dagestanica*). However, genetic sampling by Fritz et al. (2007a), Mashkaryan et al. (2013), and Mikulíček et al. (2013), including specimens from within the ranges of each of those six purported taxa, supports only the recognition of three taxa in the area (*T. g. ibera, T. g. buxtoni*, and *T. g. armeniaca*). Because the unreliability of morphology in establishing species boundaries within the genus *Testudo* has been well documented (Parham et al. 2006; Fritz et al. 2007a, 2009; Mikulíček et al. 2013; Danilov et al. 2013; among others), we have not followed Chkhikvadze et al. (2013), pending further genetic sampling.

14:36. ***Testudo* or *Chersine hermanni***: Perez et al. (2014) examined variation in mitochondrial DNA and nuclear microsatellites across the range of *T. hermanni*, and found substantial geographic differentiation based on distance between sites. They documented the greatest divergence between the recognized subspecies, with the eastern subspecies (*T. h. boettgeri*) ranging westward to and including the Po River valley in northeastern Italy. Their data also demonstrated the effects of thousands of years of human-mediated dispersal of these tortoises. Although they noted that the isolated population in France could have been established via natural or human dispersal, their data supported the hypothesis that the Spanish, Corsican, and Sardinian populations were likely the result of prehistoric human introductions of animals with Sicilian genotypes.

14:37. ***Testudo* or *Chersine h. hermanni***: Lapparent de Broin et al. (2006b) analyzed all known fossil specimens of *T. globosa*, *T. oriens*, and *T. seminota*, and concluded that *oriens* and *seminota* were synonymous with *globosa* and that *globosa* was apparently synonymous with western *T. h. hermanni*.

14:38. ***Testudo* or *Agrionemys h. kazachstanica***: *Agrionemys kazachstanica terbishi* was described by Chkhikvadze (2009) based on a mummified specimen, supposedly from Mongolia, in the Kohovd University collection (Kohovd City, Mongolia). Ansorge et al. (2012) reported that the type specimen has been lost, that the herpetologist who collected the specimen believed that it was a pet brought from Kazachstan, and that there is no confirmed record of an extant tortoise indigenous to Mongolia. They recommended that *Testudo horsfieldii terbishi* (Chkhikvadze 2009) should be regarded as a *nomen dubium* and allocated to the synonymy of *Testudo horsfieldii*; they also recommended that Mongolia should be excluded from its distribution range. We adopt these recommendations and associate the taxon with *T.* or *A. h. kazachstanica* by virtue of it having originally been described as a subspecies of that taxon.

14:39. ***Acanthochelys* and *Platemys***: Sequence data from two mitochondrial genes analyzed by Huebinger et al. (2013) supported the sister group relationship between *Platemys* and *Acanthochelys*, the monophyly of the latter, and the possibility that *A. radiolata* as currently defined morphologically may be polyphyletic. We continue to affirm the recognition of both genera, even though *Platemys* is monotypic.

14:40. ***Phrynops geoffroanus***: The name *Emys tritentaculata* was listed by Cuvier (1829) as attributed to Auguste de Saint-Hilaire, a botanist who traveled in Brazil and subsequently deposited several chelid turtles in the Paris Museum (R. Bour, unpubl. data). Based on the name, suggestive of several barbels, we assign it tentatively to the synonymy of *P. geoffroanus* pending further study of Saint-Hilaire's original specimens. The name *E. tritentaculata* does not refer to the American Box Turtle, *Terrapene carolina*, as originally synonymized by Wermuth and Mertens (1961) and followed by several others since then.

14:41. ***Platemys platycephala***: The name *Emys carunculata* Cuvier 1829 was listed by Wermuth and Mertens (1961, 1977) and Fritz and Havaš (2007) as an *ex errore* name for *E. canaliculata* Spix 1824, itself a synonym of *P. platycephala*. However, the name *E. carunculata* was attributed by Cuvier (1829) to Auguste de Saint-Hilaire, while in the same paragraph also listing *E. canaliculata* as attributed to Spix. The two names clearly represent different *nomina nuda*. Pending further studies of the chelid turtles that Saint-Hilaire collected in Brazil (R. Bour, unpubl. data), we leave *E. carunculata* in the synonymy of *P. platycephala* for the present.

14:42. ***Chelodina (Macrochelodina) kuchlingi***: This species, originally described by Cann (1997d), was synonymized under "*Chelodina rugosa*" (now *Chelodina oblonga*) by Georges and Thomson (2010). Their original basis for the synonymization was "that names that are available under the Code, but that apply to supposed taxa, unsupported by scientific evidence either in the original account or subsequently, are placed in synonymy." They also indicated that *C. kuchlingi* was described from a single specimen of uncertain origin with a long history of captivity and so was treated as a junior synonym of "*C. rugosa*" (now *C. oblonga*), citing Georges and Thomson (2006) who had questioned the distinction between *C. kuchlingi* and *C. rugosa*, but did not synonymize them. The synonymization by Georges and Thomson (2010) was subsequently followed by us (TTWG 2010) and Kennett et al. (2014) in their recent CBFTT species account for *C. oblonga* (see link under that species). However, the synonymization has recently been challenged by Kuchling (CCB, in review, and in litt.), who has provided data that *C. kuchlingi* is an apparently demonstrably valid and distinct range-restricted species, with a more extensive distribution in northeastern Western Australia (including the Ord River basin) than noted in the original description. Kuchling also raised serious concerns about the conservation status and potential regional development threats to *C. kuchlingi*. Georges (in litt.) has acknowledged the

difference in opinion, but stands by his opinion that the original description was deficient, and that there has been insufficient evidence presented to date that *C. kuchlingi* is a valid taxon.

We take note of this on-going controversy here and, based on our own principles of making only data-driven taxonomic changes in the checklist, acknowledge that our original decision to follow the hypotheses of Georges and Thomson (2006, 2010) to synonymize *C. kuchlingi* was likely premature, and also inconsistent with our continued recognition at that time of other species also synonymized by Georges and Thomson at the same time (e.g., *C. gunaleni* and *C. walloyarrina*). Given the potential conservation threats to this range-restricted species and the lack of data supporting the prior synonymization, we therefore now reverse our earlier decision and resurrect *C. kuchlingi* from its synonymy with "*C. rugosa*" (now *C. oblonga*) and await further data-driven analyses from Kuchling, Georges, and others.

14:43. ***Chelodina oblonga* (formerly *C. rugosa*)**: Thomson (2000a) demonstrated that the holotype of *Chelodina oblonga* Gray 1841 is actually a specimen of what had over the last ca. 40 years been referred to as *Chelodina rugosa* Ogilby 1890 from northern Australia. The ICZN was petitioned (Thomson 2006, 2007) to conserve current usage of the name *C. rugosa* for the Northern Snake-necked Turtle and to apply the next available name, *Chelodina colliei* Gray 1856a, to the Southwestern Snake-necked Turtle, instead of the commonly and erroneously used name *C. oblonga*. We previously discussed this ICZN case in our second checklist (Rhodin et al. 2008). Recently, in their Opinion 2315, the ICZN (2013a) declined to support the petition to give precedence to the younger, recently used name *C. rugosa* over the older name *C. oblonga* for the Northern Snake-necked Turtle. Although the latter species has been known as *C. rugosa* since 1974 and was listed as such in previous editions of this checklist, we now follow the ruling of the ICZN and use the name *Chelodina (Macrochelodina) oblonga* Gray 1841 for the Northern Snake-necked Turtle, although the name *Chelodina (Macrochelodina) rugosa* Ogilby 1890 remains an available name in the synonymy of *C. oblonga*. The decision by the ICZN has also been followed by Kennett et al. (2014) in their recently published account on the Northern Snake-necked Turtle in this CBFTT monograph series.

14:44. ***Macrodiremys***: In an attempt to conserve usage of the name *Chelodina oblonga* for the Southwestern Snake-necked Turtle, McCord and Joseph-Ouni (2007b) designated the lectotype of *Chelodina colliei* (set by Thomson 2000a) as the neotype of *Chelodina oblonga*. At the time this was done, there was already an open case before the ICZN (Thomson 2006) concerning whether to use the name *C. oblonga* or *C. rugosa* for the Northern Snake-necked Turtle (see annotation 14:43). It should also be noted that the setting of a neotype where an extant holotype (or lectotype) already exists can only be done by the ICZN. In their subsequent Opinion (ICZN 2013a), it was ruled that, considering the confusion over these names and the potential for further confusion, that the Principle of Priority should be followed, and that *C. oblonga* should maintain priority over *C. rugosa* for the northern taxon. By associating the new name *Macrodiremys oblonga* to the lectotype of *C. colliei*, McCord and Joseph-Ouni (2007b) had effectively erected a new nominal species as a junior objective synonym of *C. colliei*. Thus, since *M. oblonga* was the type species for the new genus *Macrodiremys*, then in effect so was also its senior objective synonym, *C. colliei*. Fortunately, the latter was not already a type species for another genus. Georges and Thomson (2010) reduced the various genera of snake-necked turtles to subgeneric status, all under the oldest genus name *Chelodina*; this has been recognized in previous editions of the checklist (TTWG 2012), however, the subgenus name for *C. colliei* was left undetermined because of the uncertainty surrounding the case. In this checklist edition, now that the ICZN Opinion has been published, this matter can be rectified by restoring the subgeneric name *Macrodiremys*, which follows the intent of McCord and Joseph-Ouni (2007b).

14:45. ***Elseya* and *E. schultzei***: Based primarily on mtDNA data, Georges et al. (2014) identified three reciprocally monophyletic, deeply divergent clades within the taxon formerly recognized as *Elseya novaeguineae*: 1) the Birds Head (Kepala Burung, Vogelkop, or Doberai Peninsula) population of western Indonesian New Guinea, 2) the population on the New Guinea mainland north of the Central Range, and 3) the mainland population south of the Central Range. They also demonstrated some phylogeographic structure within each of those three clades, and confirmed the genetic distinction of *E. branderhorsti* of the southern lowlands / Fly River floodplain as separate from the *E. novaeguineae* clades. They suggested that these three clades each deserved species rank, and they followed Rhodin and Genorupa (2000) in noting that the southern form is distinct and undescribed and that the name *E. schultzei* (Vogt 1911) is available for the northern population. They also implied that the name *E. novaeguineae* should be applied to the Birds Head population (the source of the type). We follow these recommendations and now recognize *E. schultzei* as a full species (again), and await additional work in progress to determine the appropriate name for the southern form, whose populations we retain under *E. novaeguineae* pending further work.

14:46. ***Flaviemys* and *F. purvisi***: Using molecular data only, Le et al. (2013) concluded that the species known as *Myuchelys purvisi* is the sister taxon to all other taxa that were included in *Emydura*, *Elseya* and/or *Myuchelys*. To correct this paraphyly, they erected a new monotypic genus, *Flaviemys*, with type species *Elseya purvisi* Wells and Wellington 1985, by original designation and monotypy. There is also support for this in previous studies, where *Flaviemys purvisi* and *Myuchelys georgesi* were perceived as a cryptic species pair, very similar by appearance, but on analysis were found not to be sister taxa (Georges and Adams 1992; Georges et al. 1998; Thomson and Georges 2009; Georges and Thomson 2010; Fielder et al. 2012; Fielder 2013). We follow this new taxonomy here.

14:47. ***Pelusios castaneus seychellensis***: Based on mitochondrial gene sequence data from all known lineages of *Pelusios*, Stuckas et al. (2013) found that the lectotype of *P. seychellensis* was nested among specimens of the West African *P. castaneus*. They concluded that *P. seychellensis* was most likely based on specimens of *P. castaneus* not native to the Seychelles Islands, and recommended the synonymy of *P. seychellensis* with *P. castaneus*. However, Bour (1983) identified significant morphological differences between these two taxa, and recently (Bour 2013a) argued that *seychellensis* might represent an ancient prehistoric introduction of *castaneus* to the islands by humans that has subsequently diverged morphologically from the ancestral population. He recommended the use of the subspecific designation *P. castaneus seychellensis* until additional comparisons (especially morphological) can be made between *castaneus* and *seychellensis*, a recommendation we have adopted. See also the pertinent discussion of the geographic occurrence of this species [*P. castaneus*] in the distributional data appendix below.

14:48. ***Podocnemis erythrocephala***: The forgotten names *Emys bitentaculata* and *Hydraspis bitentaculata* were not listed or synonymized by Wermuth and Mertens (1961, 1977) or Fritz and Havaš (2007). Gray (1830e) first placed the Cuvier manuscript name *Emys bitentaculata* under his concept of *Chelys* (*Hydraspis*) and subsequently (Gray 1831d) described it himself as *Hydraspis bitentaculata*: "*Testa rufa, subtus pallide lutea nigro maculate, scutello nuchal nullo.*" Fitzinger (1835) synonymized both names under his concept of *Hydraspis (Podocnemis) tracaxa* (which also included *Podocnemis expansa* in part and some chelid turtles). Gray's description did not identify the species very well, but the combination of a red shell (*testa rufa*), two barbels (*bitentaculata*), and lack of a nuchal scute (*scutello nuchal nullo*) suggests that it is indeed a *Podocnemis*, and we tentatively place it as most likely synonymous with *P. erythrocephala*, which shares those features, including a red shell margin in juveniles.

14:49. **Humboldt *Podocnemis* names**: Alexandre de Humboldt first published the names *Testudo arrau* (= *P. expansa*) and *Testudo terekay* (= *P. unifilis*) in the French version of his original work (Humboldt 1819a:243). This work was subsequently translated into English (Humboldt 1819b:482), and later (Humboldt 1820:415) into German. These various translations have caused some confusion in the literature, and some authors (including our previous TTWG checklists) have even attributed the names to Humboldt *in* Gray (1831d:77). However, the French version remains the original source for these names. Both names are considered *nomina oblita* (see also annotation 14:50 for *P. unifilis*).

14:50. ***Podocnemis unifilis***: *Emys cayennensis* was described from French Guiana by Schweigger (1812), but was incorrectly applied to *Podocnemis erythrocephala* for most of its history (reviewed in Pritchard and Trebbau 1984; but see David 1994 and Bour 2006a). In 1819 Humboldt (see annotation 14:49) described *Testudo terekay* from Venezuela; however, this obscure work was ignored by most subsequent authors (but see Schinz 1833). In 1830 Bell (*in* Gray 1830e) described *Chelys (Hydraspis) lata* from Guyana and this name was also ignored by most subsequent authors until Rhodin et al. (2008) declared it a *nomen oblitum* (see also Schneider et al. 2012). All three of these names apply to the taxon *Podocnemis unifilis* that was finally described from Guyana by Troschel (1848), and the latter name has been applied to the Yellow-spotted Amazon Turtle by most (but not all) authors over the last 165 years. In light of this complicated nomenclatural history, and in an

effort to ensure the stability of Troschel's name, Vogt et al. (2013) petitioned the ICZN to conserve the name *Podocnemis unifilis* Troschel 1848 for the Yellow-spotted Amazon Turtle, giving it precedence over *Emys cayennensis* whenever the two are considered synonymous. Furthermore, they declared the names *Testudo terekay* Humboldt 1819a and *Chelys (Hydraspis) lata* Bell *in* Gray 1830e as *nomina oblita*. Our checklist reflects this arrangement, pending an ICZN ruling.

2017 Checklist Annotations

TTWG 2017 (CRM 7, checklist.v.8) (17:1-105)

17:1. **Phylocode Classification**: Crawford et al. (2015) performed a genome-scale analysis of turtle phylogeny, sequencing 2381 ultraconserved element loci representing a total of 1,718,154 bp of aligned DNA sequences in 32 turtle taxa representing 14 turtle families. Their recovered phylogeny corresponded to well-supported clades that they concluded were consistent with the temporal appearance of clades and paleobiogeography. They recommended the alternative hierarchical Phylocode classification of turtles presented at the beginning of our Checklist. We continue to use our ICZN-compliant Linnaean classification as outlined on the same page.

In addition, Pereira et al. (2017) recently used publicly available databases for nucleotide sequences and composed a dataset comprising 13 loci for 294 living species of Testudines, accounting for all living genera and 83% of extant species diversity (as recognized in our checklist); they constructed a Phylocode-based classification scheme somewhat at variance with both Crawford et al. (2015) and our checklist.

17:2. ***Macrochelys***: Previous molecular studies have demonstrated significant diversity within the long-recognized widespread species *Macrochelys temminckii*: mtDNA data supported the distinction of three geographic clades, and microsatellite data suggested that six genetic clusters were recognizable (Roman et al. 1999; Echelle et al. 2010). Based on additional mtDNA data, a morphometric (osteological) analysis, and an examination of all available fossil material for the genus, Thomas et al. (2014) demonstrated the existence of three distinct, Recent, geographically separated populations: *M. suwanniensis* (Suwannee River drainage), *M. apalachicolae* (Apalachicola River and nearby lesser drainages), and a restricted *M. temminckii* (Yellow-Conecuh to Mobile Bay to Mississippi to Neches River basins). Independently, Murray et al. (2014) examined morphometric variation in the skull of *Macrochelys* across most of its range. Their results demonstrated distinct drainage-specific differences in skull morphology, supported the distinction of the Suwannee River turtles, and suggested that "further splitting may eventually be warranted" among the basins west of the Suwannee. Subsequently, Folt and Guyer (2015) reconsidered the published data, critiqued the methods of Thomas et al. (2014), and concluded that the evidence to date supported the distinction of *M. suwanniensis*, but that *M. apalachicolae* was not adequately distinguished from *M. temminckii* (sensu stricto), and thus recommended their synonymization, with which we concur.

17:3. [*Annotation text previously included here now deleted from this checklist.*]

17:4. [*Annotation text previously included here now deleted from this checklist.*]

17:5. ***Caretta caretta***: Continued examination of intraspecific genetic structure within this species by Shamblin et al. (2014) using mitochondrial gene sequences from samples from 42 nesting rookeries identified 59 different haplotypes. However, the authors made no taxonomic recommendations, and we agree.

17:6. ***Eretmochelys imbricata***: Gaos et al. (2016) examined mtDNA haplotype diversity among nesting populations of Hawksbills along the eastern Pacific Ocean. Despite the low genetic diversity across their samples, their phylogenetic analysis suggested that Eastern Pacific Hawksbills are more closely related to those from the Indo-Pacific rather than to those in the Atlantic, despite the recent closure of the Panamanian Portal. In addition, despite the low haplotype diversity overall, differences among the four major eastern Pacific rookeries (El Salvador, Nicaragua, Costa Rica, and Ecuador) led the authors to recommend that these four rookeries should be considered separate management units for conservation purposes.

17:7. ***Lepidochelys olivacea***: This species was described by Eschscholtz in 1829 in two separate publications (Eschscholtz 1829a, 1829b), the earlier of them in general overlooked until recently (Flores-Villela et al. 2016). Historically, and in our previous checklists, the *Zoologischer Atlas* (Eschscholtz 1829b) has always been credited as the source for the name *Chelonia olivacea*, and although we previously listed his other publication from the journal *Die Quatember* (Eschscholtz 1829a) as the source for two other new names (*Chelonia castanea* and *Chelonia grisea*) (TTWG 2014), we assumed that his *Zoologischer Atlas* had been published first and had nomenclatural priority. However, Flores-Villela et al. (2016) have now shown that the *Quatember* article was published earlier (January 1829) than the *Atlas* (after May 1829), and is therefore the original source for the name *olivacea*. In addition, the relatively overlooked name *Chelonia grisea*, previously synonymized by others and us under *Eretmochelys imbricata*, has been shown by Flores-Villela et al. (2016) to more likely represent an anomalous *Chelonia mydas*, and we follow their suggestion to reassign it as such.

17:8. ***Chelonia mydas***: Okamoto and Kamezaki (2014) examined plastral coloration and head and shell morphometrics for *Chelonia* captured in the Western Pacific off Japan. Their data demonstrated the presence of two distinct phenotypes ("yellow" and "black") and they argued that the black form represented *C. agassizii*, and should be recognized as a full species. However, given previous studies (see annotation 07:4), we continue to consider *C. agassizii* to be a synonym of *C. mydas*, unless integrated range-wide analyses of morphology, coloration, and genetics conclusively demonstrate significant lineage divergence.

Naro-Maciel et al. (2014) examined 15 microsatellite markers across 19 green turtle rookeries across the Atlantic Ocean. These data revealed a strong barrier to dispersal between the northern and southern Atlantic, but not a degree meriting taxonomic changes.

17:9. **Wagler 1830b and 1830c**: Gutsche and McCranie (2016) stated that Wagler's 1830 publication *Natürliches System der Amphibien* was issued as two separate parts: first the text (Wagler 1830b), and then the plates (Wagler 1830c), although dates for the publication of each were not provided. We and others had always treated these two parts as a single publication, but evidently they appeared as separate issues. As a result, the name *Dermatochelys* was established in Wagler 1830b, but the species *Dermatochelys porcata* first appeared in Wagler 1830c. Similarly, the name *Cinosternon hirtipes* was only a *nomen nudum* in Wagler 1830b, but described by association with the figures in Wagler 1830c, as we previously noted in our annotation 09:11. These changes do not affect any taxonomic or nomenclatural considerations. We also note in passing that the description of *Testudo boiei* Wagler 1830a (= *Chelonoidis carbonarius*) appears to have preceded these others.

17:10. ***Dermochelys coriacea***: Using mtDNA sequences and nuclear microsatellite markers, Molfetti et al. (2013) and Dutton et al. (2013) demonstrated considerable genetic structure within *Dermochelys coriacea* in the Atlantic Ocean basin. Although they advised the recognition of their units for conservation purposes, they made no taxonomic recommendations.

17:11. ***Kinosternon***: Iverson et al. (2013) sequenced three mitochondrial and three nuclear markers for all recognized species of kinosternid turtles, and their analysis revealed strong support for a monophyletic clade of *Sternotherus*, a second monophyletic clade of primarily Meso-American taxa that they named *Cryptochelys*, and a third restricted monophyletic clade of the remaining species formerly included in *Kinosternon*. However, their support for the non-monophyly of the traditional inclusive genus *Kinosternon* was weak. Subsequently, Spinks et al. (2014b) sequenced 14 additional nuclear loci for most (but not all) recognized species, and their analysis of the nuclear data alone supported the reciprocal monophyly of *Sternotherus* and the traditional *Kinosternon* (sensu lato), but did not support a monophyletic *Cryptochelys*. Because two independent data sets have produced different conclusions, we retain the alternative generic arrangement from our last edition until new data emerge or new analyses are performed to settle this complicated issue.

17:12. ***Kinosternon steindachneri***: Bourque (2016) treated this taxon as a distinct species based on morphology, and argued that no morphological or genetic data have been presented to support its continued recognition as a subspecies of *K. subrubrum* (Iverson 1998; Bourque 2012a,b; Iverson et al. 2013; Spinks et al. 2014; Bourque and Schubert 2015). We concur, and therefore now list *K. steindachneri* as a distinct species.

17:13. ***Kinosternon stejnegeri***: Iverson (1979a) synonymized extant *K. flavescens stejnegeri* Hartweg 1938 with the Pliocene–Pleistocene fossil taxon, *K. arizonense* Gilmore 1923, based on similar morphology. Considered a subspecies, *K. f. arizonense*, by Iverson (1979b), it was subsequently recognized as a distinct species based on genetic analysis by Serb et al. (2001). Recently, McCord (2016) examined all available Pliocene material of *K. arizonense* and compared it with extant specimens, and concluded that the fossil material differs significantly from the extant material and is actually more similar

morphologically to *K. flavescens* (sensu stricto). As a consequence, he restricted the name *arizonense* to the Pliocene fossils and resurrected the old name *stejnegeri* for the extant species. This conclusion was also tentatively accepted by Joyce and Bourque (2016), and we follow these recommendations here, accepting the resurrected name *K. stejnegeri* for the extant species previously known as *K. arizonense*.

17:14. ***Kinosternon subrubrum hippocrepis***: Bourque (2016) elevated this taxon to a full species based on morphological data from living and fossil forms; however, without a range-wide analysis of morphometric and/or molecular data supporting that change, we continue to recognize *hippocrepis* as a subspecies of *K. subrubrum*.

17:15. ***Sternotherus depressus***: Scott and Rissler (2015) reported a 32–56% decline in the historical range of *S. depressus*, as well as significant unidirectional mtDNA introgression from *S. minor peltifer*. This hybridization is changing the morphology of *S. depressus* and severely threatens its continued distinction.

17:16. ***Sternotherus minor peltifer***: Bourque (2016) argued that this taxon should be recognized as a full species based on previous published phylogenetic analyses using molecular (Iverson et al. 2013) and morphological data (Bourque and Schubert 2015). In addition, Guyer et al. (2016), based on color differences and the mitochondrial DNA study by Walker et al. (1995), also recommended elevating *S. m. peltifer* to species status. However, Walker et al. (1995) sampled part of only one mitochondrial gene and included no samples from the previously hypothesized area of intergradation along the Gulf Coast (Iverson 1977). Because animals that appear to be morphological intergrades have been described (Iverson 1977), we consider these recommendations premature and continue to recognize *peltifer* as a subspecies of *S. minor*.

17:17. **Emydidae**: Seidel and Ernst (2017) provided an extensive review of the history and taxonomic status of the phylogeographic relationships of the family and its two subfamilies, but recommended no significant taxonomic changes.

17:18. ***Chrysemys picta***: Jensen et al. (2014a) examined genetic variation among populations of *C. picta* at the northwestern range limit in British Columbia (*C. p. bellii*). Although they documented very little variation in regional mitochondrial sequences, they found two unique mitochondrial haplotypes when compared to previously published range-wide data. In contrast, based on nine microsatellite loci, they identified six distinct local population clusters. They urged that these six geographic units be managed separately, but made no taxonomic recommendations; however, their data did support the continued recognition of *C. dorsalis* and *C. picta* as separate species. In addition, Jensen et al. (2015a) examined variation in one mitochondrial and one nuclear gene across the range of the genus. The nuclear gene provided no resolution, but the mitochondrial data demonstrated the reciprocal monophyly of *C. dorsalis* versus *C. picta* (*sensu stricto*), but no clear pattern among the subspecies of the latter. Because the available evidence for the recognition of *C. dorsalis* as a species vs. subspecies is primarily mitochondrial (see TTWG annotations 07:11 and 10:6), we continue to list two alternatives for its classification.

17:19. ***Graptemys***: Praschag et al. (2017) examined over 3200 bp of mtDNA sequence data and 7800 bp of nuclear DNA (across 12 loci) for 89 specimens of all recognized taxa of the genus *Graptemys*. The mtDNA data supported the distinctiveness of *G. geographica* (sister to all other *Graptemys*), and the divergence of the broad-headed clade from the narrow-headed clade. Most species in the broad-headed clade were resolved as reciprocally monophyletic, but in the narrow-headed clade only *G. caglei* and *G. versa* were well supported as monophyletic. Principal component analyses using coded phased nuclear DNA sequences revealed eight clusters: 1) *G. geographica*; 2) *G. barbouri*; 3) *G. caglei*; 4) *G. versa*; 5) *G. sabinensis*; 6) *G. ernsti*, *G. gibbonsi*, *G. pearlensis*, and *G. pulchra*; 7) *G. flavimaculata*, *G. nigrinoda*, and *G. oculifera*; and 8) *G. ouachitensis* and *G. pseudogeographica*. The authors concluded that species recognitions of *G. geographica*, *G. barbouri*, *G. caglei*, *G. versa*, and *G. sabinensis* were all well supported, but that the remaining taxa were oversplit. They suggested 1) that *G. flavimaculata* and *G. nigrinoda* should be relegated to subspecies of *G. oculifera* (see also Mertens and Wermuth 1955); 2) that *G. ernsti*, *G. gibbonsi*, *G. pearlensis*, and *G. pulchra* are conspecific; and 3) that *G. ouachitensis* should be relegated to a subspecies of *G. pseudogeographica*. Their results supported the earlier view by Walker and Avise (1998) that *Graptemys* is taxonomically oversplit, and they made some suggestions for possible changes, but they presented no explicit taxonomic revisions, and we await further clarification before changing the checklist.

Selman (2017), in an analysis of the incidence of presence of carapacial rings vs. blotches in *G. flavimaculata*, documented that blotches are the typical pattern in all populations examined, both for recently observed as well as historically preserved specimens. This was interpreted as representing the pervasive presence of a derived character, and distinctly different from *G. oculifera*, which consistently shows the ancestral pattern of carapacial rings, but he made no taxonomic recommendations.

17:20. ***Graptemys caglei***: Ward et al. (2013) examined microsatellite variation across the range of *Graptemys caglei* in the Guadalupe and San Marcos Rivers of Texas. They found weak but identifiable divergence between populations in the Upper Guadalupe Rive versus those in the Middle Guadalupe and San Marcos Rivers. They made no taxonomic recommendations, but noted that turtles in these two regions also differed in life history, morphology, and coloration.

17:21. ***Graptemys ernsti* and *G. barbouri***: Godwin et al. (2014) confirmed the previously unrecognized presence of both *Graptemys ernsti* and *G. barbouri* in the Choctawhatchee River basin, and demonstrated hybridization between those two species in that basin. Based on the available evidence they concluded that both species were likely present in that basin prior to human intervention. In a follow-up study, Ennen et al. (2016) examined morphometric, colorimetric, microsatellite and mitochondrial DNA variation in *G. ernsti*, across the three inhabited basins. They demonstrated no morphometric differences between turtles in two of the basins (no data for the Pea River, a tributary of the Choctawhatchee), weak colorimetric differences between turtles from the Yellow vs. Conecuh/Pea basins, and both microsatellite and mtDNA evidence for the distinction of turtles from the Yellow River vs. those from the other basins. They declined to recognize those populations taxonomically, and hesitated to even recommend their recognition as Evolutionarily Significant Units, and we concur with these conclusions. Although it is now clear that both *G. ernsti* and *G. barbouri* are established in the Choctawhatchee River basin, whether either of them is native to that system is still an open question (Godwin et al. 2014; Ennen et al. 2016).

17:22. ***Graptemys nigrinoda***: Ennen et al. (2014) examined morphometric and colorimetric characters, as well as sequence variation in a single mitochondrial gene, across the range of *Graptemys nigrinoda*. Morphological variation was primarily clinal, and mitochondrial haplotypes differed by less than 0.3% and were not related to geography. They recommended the synonymy of *G. n. delticola* with the nominate form, and we have followed their recommendation.

17:23. ***Graptemys pseudogeographica***: Lindeman et al. (2015) demonstrated that False Map Turtles from the Calcasieu River basin in southwestern Louisiana differ from those in all other basins in having a unique eye color and a variable chin pattern. They made no taxonomic recommendations, but encouraged further range-wide morphological and genetic study of this species in light of these findings.

17:24. ***Malaclemys terrapin***: Hart et al. (2014) examined 12 nuclear microsatellite loci from 21 populations of *Malaclemys terrapin* from across the species' range, and identified four genetic clusters that did not correspond to currently accepted, morphology-based subspecies descriptions. Despite gaps in their coastal sampling along the east coast of Florida and Georgia, and the Gulf Coast of Florida, their recommended management units were Massachusetts (part of the range of *M. t. terrapin*), New York to South Carolina (most of the range of *M. t. terrapin*, and part of the range of *M. t. centrata*), the Florida Keys to Tampa Bay (the range of *M. t. rhizophorarum* and part of that of *M. t. macrospilota*), and Louisiana and Texas (the combined ranges of *M. t. pileata* and *M. t. littoralis*). In addition, based on 16 microsatellite loci (12 shared by Hart et al. 2014), Drabeck et al. (2014) also found low diversity and an absence of structure among populations of *M. terrapin* along the Gulf Coast of Louisiana and Texas. Both studies questioned the current subspecies designations, but made no explicit taxonomic recommendations, and we agree. More studies with better geographic sampling are clearly needed.

17:25. ***Pseudemys concinna***: In a generally overlooked publication, Guérin (1829) provided an illustration as indication of a species that he named *Emys concinna*, preceding the description of *Testudo concinna* by Le Conte (1830). However, the species figured appears to represent what is now considered *Trachemys scripta elegans*, described by Wied-Neuwied (1839). Whereas *Emys concinna* Guérin 1829 is technically a senior homonym of *Testudo concinna* Le Conte 1830 (later placed in the genus *Emys*) and precedes the description of *Emys elegans* Wied-Neuwied 1839, it is also a name and attribution that has never been used since its appearance and we therefore declare it a *nomen oblitum*, not requiring any change in status or usage of the two subsequently published widely-used and accepted names.

17:26. **Central American *Trachemys***: The taxonomy of Meso-American *Trachemys* sliders has been a quagmire for decades. Because of the level of disagreement among recent authors concerning this group, with no clear resolution in sight, our last checklist listed as many as three taxonomic options for some taxa. However, in an attempt to settle the confusion, Parham et al. (2015) examined variation in one mitochondrial and one nuclear gene across every named *Trachemys* taxon from Mexico. Their analysis revealed that the samples of "*ornata*" from Acapulco used by Fritz et al. (2012) clustered with turtles from the Caribbean versant, and not with confirmed *ornata* from near its type locality. Hence, the species name *ornata* should be restricted to the western versant of Mexico, and the name *venusta* should be restricted to the eastern drainages of Mexico and Central America. Furthermore, they confirmed the finding of Fritz et al. (2012) that *grayi* is not closely related to *venusta*, but the sister taxon of *emolli*. We agree with these insights (although we remain equivocal regarding whether *emolli* is a distinct species or a subspecies of *grayi*) and have modified our checklist accordingly. Additionally, Fritz et al. (2012) showed that *panamensis* is very closely related to *grayi*, and should be considered a subspecies of that taxon, rather than of *venusta*, further corroborating the split between western and Caribbean versant taxa; we have now made that change in our checklist. Also, as a consequence of the recognition that *ornata* is a western versant taxon, *callirostris* is now considered either a distinct species or a subspecies of *venusta*, rather than of *ornata*, as it was designated in our previous checklist.

17:27. ***Trachemys adiutrix* or *T. dorbigni adiutrix***: Using a niche modelling approach, Rodrigues et al. (2016) supported the subspecies status of the Maranhao slider. The invasive potential of the nominotypical subspecies was better explained when the different climatic niches of both taxa were combined, reflecting the common pattern of lack of niche conservatism between subspecies.

17:28. ***Trachemys grayi***: The species *Callichelys concinna* Gray 1873a, described from San Mateo, Tehuantepec, Oaxaca, Mexico, has unfortunately been overlooked in all previous checklists, including our own (Siebenrock 1909a; Wermuth and Mertens 1961, 1977; Fritz and Havaš 2007; TTWG 2014). However, Gray (1873f) and Boulenger (1889) synonymized it with *T. grayi*, and Harfush-Meléndez and Buskirk (2008) noted it in their analysis of the distribution of *T. grayi*, and we now include it in our checklist.

17:29. ***Clemmys guttata***: Davy and Murphy (2014) examined variation in 11 nuclear microsatellite loci across Canadian populations of *Clemmys guttata*. They identified significant genetic structure, with a minimum of six distinct subpopulations, with the most distinctive population being in Hastings County, Ontario. They made no taxonomic recommendations, but strongly urged management and protection of the latter population, which numbers less than 50. An expansion of this study across the entire range of *C. guttata* would be invaluable.

17:30. ***Emys orbicularis orbicularis***: The identity of the Pleistocene fossil names *Clemmys schlotheimii* † Fitzinger 1835 and *Trionyx schlotheimii* † Fitzinger 1835 has previously been uncertain, and we did not even include *T. schlotheimii* in our previous checklists. However, Karl and Paust (2014) examined the original fossil material, designated lectotypes of both taxa, and confirmed that both are synonyms of *Emys orbicularis*.

17:31. ***Emys orbicularis orbicularis***: The Pleistocene fossil taxon *Testudo (Emys) canstadiensis* † Plieninger 1847 was included under the synonymy of *Testudo hermanni* in our previous checklist (TTWG 2014), based on presumed synonymy suggested by Auffenberg (1974); however, the specimen is actually an *Emys orbicularis*, based on synonymization by Karl and Tichy (2002), and we corrected it in our fossil checklist (TEWG 2015), and herein.

17:32. ***Emys orbicularis occidentalis***: Based on both mitochondrial DNA sequences and 15 nuclear microsatellite markers, Stuckas et al. (2014) identified two distinct genetic units within *Emys orbicularis* in North Africa, one from Morocco (identified as *E. o. occidentalis*) and the other from eastern Algeria and Tunisia (undescribed). The former was found to be very similar to Iberian specimens (identified as *E. o. fritzjuergenobsti*). The authors declined to name the new taxon for want of morphological data, but synonymized *E. o. fritzjuergenobsti* under *E. o. occidentalis*, and we reflect that change here.

17:33. ***Emys orbicularis persica***: We spell the name of this taxon as *Emys orbicularis persica*, following Fritz (1998). However, Eichwald (1831:196) recognized three varieties ("var.") of *Emys europaea* Schneider 1783: the nominotypical, not named, α by inference, and *β* and *γ*, with names given as "*ibericae* var. *β*" and "minoris var. *γ persicae*." One might therefore assume that the valid names should be *Emys europaea persicae* and *Emys europaea ibericae*. However, the names *ibericae* and *persicae* are adjectives, referring to the countries of Iberia and Persia, respectively, and as a result, ICZN Article 11.9.2 applies: "An adjectival species-group name proposed in Latin text but written otherwise than in the nominative singular because of the requirements of Latin grammar is available provided that it meets the other requirements of availability, but it is to be corrected to the nominative singular if necessary." Therefore, *persica* and *iberica* are the valid names, and these were subsequently used by Eichwald (1840:47).

17:34. ***Emys trinacris***: Vamberger et al. (2015) examined mitochondrial and nuclear microsatellite variation (15 markers) in *Emys* from southern Italy. Their results revealed negligent gene flow between *E. orbicularis* and *E. trinacris*, with intergradation evident between *E. o. galloitalica* and *E. o. hellenica*. Their data support the continued recognition of *E. trinacris* as a species, and *galloitalica* and *hellenica* as subspecies of *E. orbicularis*.

17:35. ***Emys* or *Actinemys pallida***: Spinks et al. (2014a) examined 89 nuclear single nucleotide polymorphisms (SNPs) and a mitochondrial gene sequence from range-wide samples of *Emys* or *Actinemys marmorata*, and compared those results with earlier work (Spinks et al. 2010) using nuclear gene sequence data. The mitochondrial data resolved four poorly supported clades, whereas the previous nuclear sequence data revealed only two main groups with considerable admixture between them. However, the SNP analysis demonstrated strong support for two geographic clusters: a northern group from the southern San Joaquin valley to Washington, and a southern group from the Central Coastal Range of California and southern California to Baja California. These results are remarkably (though not perfectly) consistent with Seeliger's (1945) morphological work, who described the subspecies *pallida* for the southern populations. Furthermore, secondary analyses by Spinks et al. (2014a) of the molecular data within each of these two groups recognized two subgroups in the southern cluster, one from the coastal range and southern California, and the other from Baja California. Although populations from Baja were also recognized as morphologically distinct (though undescribed) by Seeliger (1945), Spinks et al. only noted that species recognition may emerge from future studies. Hence, they recommended that the two California lineages be recognized as separate species (rather than subspecies as defined by Seeliger 1945), and we follow that recommendation here.

17:36. ***Emys* or *Emydoidea blandingii***: Based on variation across 12 microsatellite loci, Davy et al. (2014) identified four distinct genetic units among populations of *Emydoidea blandingii* in Ontario, Canada, and suggested that they should be managed separately. Similarly, based on eight microsatellites, Sethuraman et al. (2014) identified moderate but significant differentiation among Midwestern USA populations, with four or five unique genetic clusters, but were unable to explain the close genetic relationship between a population in western Nebraska to those in eastern Illinois, rather than with intervening populations in Iowa. McCluskey et al. (2016) examined variation in seven microsatellites in this species in New York, and identified two or possibly three genetic units there. None of these studies made taxonomic recommendations. Range-wide genetic studies of *E. blandingii* are clearly needed, as is careful population management of this genetically diverse species.

17:37. ***Terrapene carolina* complex**: These North American box turtles remain the center of considerable taxonomic controversy (see TTWG 2014, annotation 14:27). Based primarily on mtDNA data, Martin et al. (2013) recommended the recognition of the western forms (*triunguis*, *mexicana*, and *yucatana*) as a polytypic species (*T. mexicana* being the oldest name) separate from the eastern forms (under *T. carolina*). However, in a brief summary of the turtle taxonomy issues, Fritz and Havas (2013) declined to accept this major change for the genus *Terrapene*, primarily because of evidence for introgression between the proposed species. Martin et al. (2014) responded with evidence of interspecific hybridization between other well-accepted species pairs, and reaffirmed their conclusion that *T. mexicana* was a distinct species by the phylogenetic species concept. Fritz and Havas (2014) replied with three lines of evidence as to why they considered it to be premature to recognize *T. mexicana* as distinct from *T. carolina*. First, intergradation is common where they are sympatric, and indeed, Butler et al. (2011) demonstrated panmixia (rather than occasional hybridization) in those areas. Second, *mexicana* was resolved as monophyletic by Martin et al. (2013) with relatively weak support based on mtDNA, and no support when based on a single nuclear gene. Third, previous study of mitochondrial and nuclear DNA sequence data (Spinks et al. 2009; seven nuclear loci) did not resolve *triunguis* as reciprocally monophyletic relative to eastern forms of *T. carolina*. A recent analysis of 30 nuclear and one mitochondrial loci (Spinks et al. 2016) indicated that while *triunguis* appears to be monophyletic and distinct, it is nested within a paraphyletic *carolina*;

given the phylogenetic uncertainty surrounding much of this complex and the demonstrated hybridization among taxa, these authors also recommended that the traditional *carolina* taxonomy (all taxa as subspecies of *carolina*) be retained pending a comprehensive, genomic-level analysis of all contained taxa. Unfortunately, some authors (e.g., Guyer et al. 2015; Powell et al. 2016) have chosen to recognize the western populations as *T. triunguis* (rather than *T. mexicana*), without justification. Since the taxonomic issues here appear relatively unresolved, we continue to recognize the competing alternative taxonomies, pending additional sampling of the nuclear genome.

17:38. ***Terrapene carolina carolina***: In a nearly range-wide study of variation across eleven microsatellite loci for the subspecies *T. c. carolina*, Kimble et al. (2014) found surprisingly little genetic structure, except that the Appalachian Mountains represented a modest barrier to gene flow among populations.

17:39. ***Platysternon megacephalum***: In the present checklist we reallocate the placement of *Platysternon megacephalum tristernalis* Schleich and Gruber 1984 by moving it from the synonymy of *P. m. megacephalum* into the synonymy of *P. m. peguense*. Ernst and Laemmerzahl (2002) originally placed *P. m. tristernalis* in the synonymy of the nominotypical subspecies, which was followed by Fritz and Havaš (2007) and all previous editions of our checklist. However, Vetter and van Dijk (2006) identified the holotype of *tristernalis* from southwestern Yunnan as representing *peguense*, the taxon occurring in adjacent northwestern Laos, and Zheng et al. (2013) identified animals apparently from the *tristernalis*-topotypic area of southwestern Yunnan as *peguense*, both morphologically and genetically, and we agree with these assessments. In addition, Ernst and Laemmerzahl (2002) noted that the populations on Hainan appear to be intergrades between *P. m. peguense* and *P. m. shiui*, as also reflected on our current map.

Using the duplicate control region sequences of the mitochondrial genome of 20 *P. megacephalum* representing all three subspecies, Zheng et al. (2013) found these morphologically defined taxa to also be genetically distinct, with *P. m. megacephalum* and *P. m. shiui* identified as sister taxa. More comprehensive geographic surveys and the application of additional genetic markers to samples from across the species' entire range should hopefully refine current taxonomy and provide a better understanding of the geographic ranges of the individual taxa.

17:40. ***Cuora***: Tiedemann et al. (2014) sampled 16 nuclear microsatellite markers across all members of the *Cuora trifasciata* complex (*C. aurocapitata, C. pani, C. zhoui, C. trifasciata*, and the controversial Vietnamese *C. cyclornata*) (see previous annotations on the status of *cyclornata* Blanck et al. 2006a: TTWG 2007b, 07:36; TTWG 2009, 09:23; TTWG 2012, 12:22). The analyses by Tiedemann et al. (2014) confirmed the genetic distinction of each of these five taxa, and they recommended their recognition at the species level. In addition, their data corroborated the genetic distinction of the two morphotypes within *cyclornata* that were previously described as *C. c. cyclornata* and *C. c. meieri* (Blanck et al. 2006a). Finally, they also corroborated the genetic distinction of two morphotypes within the restricted *C. trifasciata*, one from the Chinese mainland (*C. t. trifasciata* according to the authors) and the other an undescribed subspecies from Hainan Island.

Independently, Li et al. (2015) sequenced the entire mitochondrial genome of nine species of *Cuora*; their phylogenetic analysis revealed that *C. trifasciata* (*sensu lato*) is polyphyletic, and that Chinese (*trifasciata*) and Vietnamese (*cyclornata*) populations are distinctly different. They also recommended the species recognition of *C. cyclornata*.

Given that morphology and mitochondrial and nuclear DNA data each support the recognition of *C. cyclornata* as a species distinct from *C. trifasciata*, with the former comprising two genetic lineages, we now follow Blanck et al. (2006a) in recognizing *C. cyclornata* with two subspecies, *C. c. cyclornata* and *C. c. meieri*.

17:41. ***Cuora amboinensis***: In a morphological and colorimetric study of populations of *Cuora amboinensis*, Ernst et al. (2016) recommended the synonymy of *C. a. lineata* with *C. a. kamaroma*, and of *C. a. couro* with *C. a. amboinensis* (suggesting that *C. a. couro* was an intergrade between *C. a. kamaroma* and *C. a. amboinensis*). However, their samples of *lineata* and *couro* were small, their analyses and conclusions were based primarily on color patterns, and they did not discuss the size of the inguinal scute (apparently diagnostic for at least *lineata*), nor did they present statistical or graphical results of their morphometric analyses (i.e., separately from color data).

Subsequently, Protiva et al. (2016) analyzed shell shape and mitochondrial DNA from *C. amboinensis* from primarily Borneo, Sumatra, and Seram. They disagreed with the findings of Ernst et al. (2016), and documented morphologic and genetic differentiation between Bornean vs. Sumatran populations, affirming the distinctiveness of *C. a. couro* from Sumatra and *C. a. kamaroma* from Borneo. Additionally, they identified a deeply divergent lineage from Seram that they indicated might require taxonomic recognition based on further work and improved sampling. Based on this, we consider the synonymizations by Ernst et al. (2016) to be premature, and await comprehensive genetic studies of *C. amboinensis* before altering the taxonomy.

17:42. ***Cuora aurocapitata, C. cyclornata*, and *C. trifasciata***: Blanck et al. (2017) analyzed variation in these three *Cuora* species from across their range, using morphometric principal components analysis (PCA) and microsatellite data, comparing their results with previous genetic work by Tiedemann et al. (2014) and Li et al. (2015). They concluded that all three species are polytypic and described a new subspecies within each: *C. aurocapitata dabieshani*, *C. cyclornata annamitica*, and *C. trifasciata luteocephala*. We tentatively accept these named taxa as new subspecies pending further analysis, but note that the described lineages are not completely distinctive and that the mixing of individuals across the clades suggests that this may be too fine an application of the lineage approach to defining taxa.

17:43. ***Cuora flavomarginata***: The map for this species has been updated and revised extensively by re-evaluating most previously recorded localities. We have also added many localities provided by T. Blanck based on compilations of Chinese survey literature and field data.

17:44. ***Cyclemys atripons* and *C. oldhamii***: Vamberger et al. (2017b) examined a recently discovered *Cyclemys* population from Phnom Kulen National Park in northwestern Cambodia using external morphology, 17 unlinked microsatellite loci, and the mitochondrial cytochrome b gene. Morphologically, the turtles resemble *C. oldhamii*, but have mitochondrial haplotypes of *C. atripons*, while having microsatellite loci distinct from *C. atripons*. The authors concluded that this population represents either a natural hybrid swarm of *C. atripons* and *C. oldhamii* or a distinct undescribed species with introgressed mitochondria of *C. atripons*, without drawing taxonomic consequences. This underscores that genetic differentiation in *Cyclemys* is complex and still incompletely understood.

17:45. ***Cyclemys pulchristriata***: Using a turtle kept in Shanghai Zoo, Li et al. (2017) published a complete mitochondrial genome (mt genome) assigned to *Cyclemys pulchristriata* and calculated a phylogenetic tree using complete mt genomes of other geoemydid species from GenBank. The topology of their tree conflicted with the trees published by Fritz et al. (2008) and Stuart and Fritz (2008) using the mitochondrial cytochrome b gene. When the cytochrome b sequences of the mt genomes used by Li et al. (2017) are compared to the data set of Fritz et al. (2008), it turns out that the sequences of '*C. atripons*' and '*C. pulchristriata*' used by Li et al. match the *C. pulchristriata* from the data set of Fritz et al., while the sequence of '*C. dentata*' used by Li et al. match the *C. atripons* from the data set of Fritz et al. (Fritz, pers. comm.). The data of Fritz et al. (2008) are considered taxonomically reliable. This underlines the pitfalls of sequencing animals with unclear identification and using uncritical taxonomic identifications of GenBank data.

17:46. ***Malayemys* species**: Brophy (2004) examined morphometric variation in *Malayemys subtrijuga* Schlegel and Müller 1845 (sensu lato) across its known range and concluded that turtles from the Mekong River basin (Laos, Cambodia, and Vietnam) differed (primarily in color pattern) from those in the Chao Phrya River basin (Thailand), and resurrected the name *macrocephala* (Gray 1859) for *Malayemys* from the latter basin.

Despite the facts 1) that Schlegel and Müller (1845) reported the type locality of *subtrijuga* as "Java", 2) that at least three syntypes are known (RMNH 6082, 6084–6085: Hubrecht 1881; although two others [NHMUK 1947.3.4.53 (specimen "m" listed in Boulenger 1889) and MNHN 7964 may also be among the original type series]; King and Burke 1989; Iverson 1992), 3) that field data associated with the RMNH syntypes indicate collection "most probably in the most western province of Bantam [=Bantan]" (Hubrecht 1881), 4) that numerous specimens of this taxon from "Java" were available in the 1800s (in at least eight European Museums; see Specimens Examined in Brophy 2004), 5) that this species has been recorded from at least six localities on Java (Brophy 2002); and 6) that Brophy's own morphometric analysis (2002: Figs. 30–31; 2004: Figs. 6–7) demonstrated that Javan specimens were distinct from specimens from both the Chao Phrya and Mekong basins (with the latter two samples actually overlapping in morphometric hyperspace), Brophy (2004) argued a) that the type locality of *subtrijuga* ("Java") was in error, b) that *Malayemys* does not occur nor has it occurred on Java, and c) that the

holotype must have originated from the Mekong basin. Hence, Brophy (2004) assigned the name *subtrijuga* to the population of *Malayemys* inhabiting the Mekong basin (but refrained from restricting the type locality of *subtrijuga*, citing uncertainty about whether the Javan specimens might be native), and the name *macrocephala* to the population in the Chao Phrya basin. Furthermore, Brophy (2004) distinguished these two populations primarily on color patterns of the head.

More recently, Sumontha et al. (2016) and Ihlow et al (2016) observed that the head pattern in *Malayemys* from the Mekong tributaries of the Khorat Plateau (northeastern Thailand) and adjacent Laos, differed from that of specimens from elsewhere in the Mekong basin. Based solely on color differences, Sumontha et al. (2016) described a population from the northern Khorat Plateau as *Malayemys isan*. Nearly simultaneously (but see below), Ihlow et al. (2016) examined color pattern and some morphometrics, as well as mtDNA and nuclear microsatellite variation across three Southeast Asian mainland regions of *Malayemys* distribution (Chao Phraya basin, Mekong basin, and Khorat Plateau), and concluded that each of these regions has its own distinctive species, based on each of their data sets, and therefore described the form that they recorded from three locations on the Khorat Plateau as *M. khoratensis*. Two of those locations of *M. khoratensis* are effectively sympatric with the *M. isan* population, and color patterns in the two taxa are very similar, suggesting that the two taxa are subjectively synonymous. Although the description of *M. isan* by Sumontha et al. appeared online first, their paper did not comply with ICZN standards for digital publication, whereas that by Ihlow et al. did (see below); therefore, we conclude that *M. isan* is a junior subjective synonym of *M. khoratensis*.

Unfortunately, neither of these papers mentioned the possibility of taxonomically recognizing the Javan population as a fourth, distinctive taxon, whose continued occurrence is confirmed by ongoing exports for commercial trade (UNEP-WCMC 2017). If further phylogenetic research demonstrates the Javan population to be taxonomically distinct, the name *Emys subtrijuga* Schlegel and Müller 1845 would apply to that population, as would its later synonyms *Cistuda gibbosa* Bleeker 1857b, *Emys nuchalis* Blyth 1863, and *Damonia oblonga* Gray 1871. Furthermore, Gray 1870c also described *Damonia crassiceps* from "China", the locality probably erroneous but possibly representing the Mekong River basin population. Finally, none of these papers have addressed Schweigger's 1812 name *Emys herrmanni*, which is considered by most authors a *nomen dubium*, synonymous with *M. subtrijuga*.

It is problematic that none of these type specimens have been included in any morphometric or genetic analysis to date. For example, if a Javan population does or did exist, and it was found to be distinctive (as Brophy's morphometric analyses suggested), then the name *subtrijuga* should be applied there, and the Mekong population might be identifiable as *crassiceps*.

In conclusion, until a more comprehensive analysis of coloration and pattern, morphometrics, and both mitochondrial and nuclear DNA variation among all populations (including Java) is undertaken, with the inclusion of all of the type specimens mentioned above, we cannot be certain which name applies precisely to which population of *Malayemys*. Finally, the TTWG is not unanimous in its support for the recognition of *macrocephala* or *khoratensis* as full species, and whether or not *isan* is synonymous with or distinct from *khoratensis*. However, to minimize taxonomic changes as we await further data, we here recognize three species in the genus *Malayemys*: *subtrijuga*, *macrocephala*, and *khoratensis*.

17:47. ***Malayemys khoratensis***: In early 2016 a distinctive clade of Snail-eating turtles (*Malayemys*) from the Khorat Plateau of eastern Thailand was described as a new species with two different names in two separate articles (Ihlow et al. 2016; Sumontha et al. 2016). The name *M. khoratensis* Ihlow et al. 2016 was published on 6 April 2016, the date when the electronic version of the article, published in the journal *PLoS One*, met the ICZN Code for nomenclatural availability of electronic articles under article 8.5.3 (ICZN 2012). The name *M. isan* Sumontha et al. 2016 was published electronically on 26 March 2016 in the journal *Taprobanica*; however, the Zoobank registration of this article failed to meet the requirements of ICZN article 8.5.3.1, rendering the electronic version unavailable for the purposes of nomenclature. It must instead be considered as published when it first met the requirements of ICZN article 8.4.1, i.e., when numerous identical hard copies printed on paper became available. There is no evidence of the journal *Taprobanica 8(1)* having been printed prior to 13 April 2016, when copies were requested and were simultaneously sent to the archiving libraries as listed on the journal's website (Thomson and Lambertz, in press). Therefore, the date of publication of the journal *Taprobanica 8(1)* and the contained article by Sumontha et al. (2016) is to be corrected to 13 April under ICZN article 21.4, the first date for which there is evidence of its physical existence. As a result, the name *Malayemys khoratensis* has nomenclatural priority, and *Malayemys isan* becomes a junior subjective synonym (see above annotation 17:46).

17:48. ***Damonia crassiceps* Gray 1870c**: This taxon has been listed under the synonymy of *Malayemys subtrijuga* by most authors since Smith (1931). However, Sumontha et al. (2016) questioned its identification as representing either this species or genus, based on the description of the drawing of the species as lacking facial stripes, but did not suggest an alternative taxonomy.

17:49. ***Mauremys annamensis* and *M. mutica***: Zhao et al. (2016a) sequenced the entire mitochondrial genomes of three specimens of *Mauremys mutica* without published locality data, but purported to originate from Taiwan, China, and Vietnam or Hainan. Phylogenetic analysis revealed that the latter genome was more similar to that of *M. annamensis* than to the other more northerly mutica genomes, rendering *M. mutica* paraphyletic. Zhao et al. (2016b) examined variation in the mitochondrial barcode gene COI among larger sample sizes of *M. annamensis* and *M. mutica*. They identified reciprocally monophyletic northern (including Taiwan) and southern (Vietnam, Hainan and *M. annamensis*) clades, but the latter showed no monophyly among the constitutent populations. They speculated that *M. annamensis* might be of hybrid origin. Furthermore, Zhou et al. (2015) compared the entire mitochondrial genome of samples of eight species of *Mauremys*, and their results also nested *annamensis* within *mutica*, sister to *mutica* from southern China (near Vietnam). Unfortunately, they included no material from Vietnam, and most of their samples were purchased from the pet trade or food markets. More extensive geographic sampling (including *M. m. kami* from the Ryukyus) and nuclear analysis will be necessary before any taxonomic changes are warranted.

Independently, Somerová et al. (2015) examined variation in a mitochondrial gene and a nuclear intron in European zoo specimens of *M. annamensis*, and also resolved *M. mutica* as paraphyletic with respect to *annamensis*. In addition, their analysis also revealed two distinct, reciprocally monophyletic mitochondrial clades within *annamensis*, which they recommended be maintained separately in captive breeding operations. Unfortunately, the natural geographic provenance of these two clades was unknown.

17:50. ***Mauremys caspica* and *Mauremys rivulata***: Using microsatellite loci and nuclear and mitochondrial DNA sequences, Vamberger et al. (2017a) showed that the two species hybridize only rarely along their contact zone in Turkey. However, there is evidence for introgression between both species. In addition, they found hybrid evidence on Cyprus and, unexpectedly, terrapin populations in Israel and Jordan turned out to be a hybrid swarm of the two species, morphologically resembling *M. rivulata*. Ecological paleomodeling suggested that the two species formerly had temporary contact across what is now the Syrian Desert during more humid climatic episodes.

17:51. ***Mauremys leprosa***: An analysis of variation in two mitochondrial markers and one nuclear gene across the range of *Mauremys leprosa* by Veríssimo et al. (2016) confirmed two distinct genetic lineages that generally correspond to the two currently recognized subspecies. One is distributed from southern France through Iberia to Morocco, north of the Atlas Mountains (*M. leprosa leprosa*); the other occurs from northern Libya and Tunisia westward to Morocco, both north and south of the Atlas mountains (*M. leprosa saharica*); in northern Morocco there is secondary contact between these subspecies. In northeastern Iberia and southern France, some native populations have been genetically impacted by introduced *M. l. saharica* (Palacios et al. 2015).

17:52. ***Mauremys reevesii***: The historic native distribution of this species has been difficult to determine accurately, as the species has been traded extensively in China for several thousand years, and has apparently been introduced to Taiwan and Japan in historic times (see the genetic analysis by Suzuki et al. 2011). The populations in Korea may also have been introduced prehistorically, but could represent natural distribution during an interglacial warming period. Our previous distribution map in the last checklist was based on Iverson (1992) and the EmySystem database plus input from the CBFTT account by Lovich et al. (2011). We have now updated and revised this map extensively by re-evaluating most localities and restricting the distribution to areas below approximately 600 m elevation. We have also added more localities provided by T. Blanck (based on compilations of Chinese survey literature and field data) and D. Gaillard. While generally a lowland species, some populations (e.g., in Anhui) appear to occur up to elevations of ca. 500 m in hill regions, but in more southern regions (e.g., Hunan and Jiangxi) the species appears to be limited to lowland areas below 300 m (T. Blanck, unpubl. data).

Most northern Chinese localities for *M. reevesii* appear to be trade specimens from ports and market centers (as also concluded by Pope 1935:46). Many southern and eastern coastal specimens are also trade specimens from coastal ports and markets or possibly locally invasive from markets. However, ancient Chinese writings seem to indicate that this species apparently occurred along the south coast, at least in Guangdong (T. Blanck, unpubl. data). Native wild populations do occur in the Chengdu basin of the upper Yangtze in Sichuan and all along the central Yangtze lowland basin, notably in Anhui, Guizhao, Hubei, Hunan, and Jiangxi. There also appears to be a native population in the Wei Valley of Shaanxi. Other scattered records north of the Yangtze may or may not represent native populations. Further genetic studies of Korean and southern and eastern coastal specimens with comparisons to specimens from the central Yangtze and Sichuan portions of the range are clearly needed.

Oh et al. (2017) examined geographic variation in mitochondrial cytochrome b sequences across the range of *M. reevesii*. Their results were complicated by the translocation of this species by humans since prehistoric times. They suggested that the two main natural populations in China and Korea were weakly but distinctly divergent genetically; however, introgression as a result of translocations is diluting that difference, precluding any taxonomic recognition.

17:53. ***Mauremys rivulata***: Vamberger et al. (2014) examined variation in a mitochondrial gene fragment and across 13 microsatellite loci for samples from throughout the range of *Mauremys rivulata*. The mitochondrial sequence data showed no evident structuring. However, the microsatellite data revealed a distinct genetic break in southern Turkey, separating eastern and western populations. This break is likely to be caused by introgressed alleles from *M. caspica* in the eastern portion of the distribution range of *M. rivulata* (Vamberger et al 2017a). Vamberger et al. (2014) attributed the lack of clear substructure among the western populations of *M. rivulata* to trans-Mediterranean dispersal.

17:54. ***Melanochelys trijuga parkeri***: This taxon was described as a large-bodied, low-shelled subspecies endemic to Sri Lanka, but recent surveys at all historically recorded localities have encountered mainly the more common and widespread smaller Sri Lankan subspecies *M. t. thermalis* (A. de Silva, unpubl. data). These two taxa need genetic evaluation to determine if and how they may be distinct, and if distinct, whether they are undergoing introgression and intergradation.

17:55. ***Aldabrachelys* and *Cylindraspis***: Based on contemporary Indian Ocean currents, and historic fluctuations in sea levels, Wilmé et al. (2017) proposed that insular Indian Ocean populations of tortoises were the result of introductions by humans approximately 4000 ybp. However, given that previous work dates the colonization of these islands by tortoises at 9.5–22 mybp, and fossils of *Aldabrachelys* have been dated to at least 138,000 ybp (Cheke and Hume 2008; Gerlach and Paquette 2014; Cheke et al. 2017; Hansen et al. 2017; Hawlitschek et al. 2017), this hypothesis is untenable.

17:56. ***Aldabrachelys gigantea* and *Testudo***: Besnard et al. (2016) used shotgun sequencing to elucidate the entire mitochondrial genome of *Aldabrachelys gigantea*. Phylogenetic analysis of the aligned sequence of this tortoise with the available mitochondrial genomes of twelve other tortoise species generally supported the results of Le et al. (2006), except for the placement of *Malacochersus* and *Testudo horsfieldii*. However, the analysis by Besnard et al. suggested a paraphyletic genus *Testudo*, although with reduced bootstrap support (77–78%). Synthesis of these data with nuclear markers is needed before any taxonomic changes are warranted.

17:57. ***Aldabrachelys gigantea***: Turnbull et al. (2015) examined variation in body size and sexual size dimorphism among four subpopulations of Aldabra tortoises. They speculated that these differences might have a genetic basis, and recommended population genetic studies.

17:58. ***Chelonoidis***: The gender of the tortoise genus name *Chelonoidis* Fitzinger (1835) has long been assumed to be feminine (e.g., Agassiz 1857) because its root is the feminine Greek noun *Chelone* (not *Chelonos* as assumed by Olson and David [2014]), and the Latinized suffix *-oides* (from the Greek *-eides*, meaning form or shape [or "like" in English]). However, according to the ICZN (Article 30.1.4.4) genus names ending in *-oides* must be considered masculine unless the original author indicated gender directly or indirectly (i.e., by usage). Because Fitzinger gave no direct indication or orthography indicating that he considered the name *Chelonoidis* to be feminine (or neuter), Olson and David (2014) argued that it should be considered masculine. However, Fitzinger did provide indirect evidence that he considered the genus name masculine, by his clear rendition of other reptile genera he described as masculine (e.g., *Dracontoidis* and *Elapoidis*). In either case, the conclusion is the same—that a strict application of the Code would render *Chelonoidis* masculine, and require the emendation of the suffix of many of the species currently recognized within that genus. The name *vicina* is treated as a noun in apposition (Art. 31.2.2) and thus is unaffected by the gender of the genus name. While the conversions to *carbonarius, denticulatus, niger*, and *phantasticus* imply disruption to a few well-established and widely-used names, they should not lead to any significant confusion, and in the interest of Code compliance, we adopt these changed endings.

17:59. ***Chelonoidis chilensis***: Sánchez et al. (2015) examined variation in the karyotype of *Chelonoidis chilensis* across its range. They identified two karyomorphs, one from tortoises in the Dry Chaco Ecoregion, and one from the Monte of Steppes and Plains Ecoregion. However, these karyomorphs were independent of the external morphotypes of *donosobarrosi, petersi*, or *chilensis*, and therefore they followed Fritz et al. (2012a) in recognizing only a single species (*C. chilensis*) in this complex.

17:60. ***Chelonoidis niger* complex**: Molecular studies of Galápagos tortoises have surged over the last several years, although most of the work has been more directed at population genetics than establishing species boundaries and their taxonomic implications. The general working assumption is that separate island populations previously recognized as subspecies of *Chelonoidis niger* [= *C. nigra*] (or synonyms thereof) are now accepted as species (e.g., Caccone et al. 2002; Russello et al. 2005, 2007; Poulakakis et al. 2008, 2012, 2015; Chiari et al. 2009; TTWG 2009 [annotation 09:32], 2014; Edwards et al. 2014; Garrick et al. 2014), and we continue to recognize them as such, now also resurrecting the three previously synonymized southern Isabela taxa (*guntheri*, *microphyes*, and *vandenburghi*) from their synonymy under *C. vicina*. However, Loire et al. (2013) examined the population genomics of five Galápagos tortoise taxa (purportedly representing three named island populations) based on 248 nuclear genes. Their results suggested panmixis across their samples, with little genetic differentiation, and they questioned the species-level recognition of the various island taxa. Expansion of this study to include Galápagos-wide sampling would clearly be helpful in settling some of the taxonomic issues facing these tortoises.

Edwards et al. (2014) examined 14 microsatellite loci for tortoises from southern Isabela from the ranges of *C. vicina* and *C. guntheri* (the latter was synonymized with the former based on mtDNA data in Poulakakis et al. 2012). Not only did they find support for the distinction of those two taxa, but they also identified a third, unnamed, geographically intermediate genetic cluster they referred to as the "*aplastados*" [flattened] type. Although they made no taxonomic recommendations, and an expansion of this study to include other Isabela populations is needed, these data warrant our removal of *C. guntheri* from the synonymy of *G. vicina*, with full species recognition. Simultaneously, based on 12 microsatellite loci and mtDNA sequence data, Garrick et al. (2014; see also Emerson and Faria 2014) identified two distinct but coalescing genetic lineages within *C. becki* on northern Isabela that appear to represent two different colonizations from Santiago (*C. darwini*). They also found some evidence of introgression of *C. vandenburghi* into the genome of *C. becki*. Hence, they concluded that "species boundaries in the group may be somewhat porous," and questioned the species-level status of *C. darwini* relative to *C. becki*.

Clearly, the evolutionary history of colonizations and divergences by tortoises in the Galápagos has been very complex, and much more reticulate than previously realized. Sorting out this history is an on-going challenge. However, as we noted in a previous checklist (TTWG annotation 12:31), genotyping of the many type specimens of named Galápagos tortoises is desperately needed, so that the correct names may be applied to all genetic lineages. Until those data are available, confusion and nomenclatural uncertainty will no doubt continue, such as some authors' use of *elephantopus* instead of *niger* or *nigra* (see TTWG annotation 09:33), and *ephippium* instead of *duncanensis* (e.g., Poulakakis et al. 2012; Jensen et al. 2015b; Hennessy 2015).

17:61. ***Chelonoidis donfaustoi***: Chiari et al. (2009) and Poulakakis et al. (2015) analyzed mitochondrial DNA and nuclear microsatellite characters in the two giant tortoise populations on Santa Cruz (Reserva and Cerro Fatal). Both studies found the populations to be genetically different and Poulakakis et al. (2015) demonstrated that the Reserva population represented the previously described *C. porteri*, and that the Cerro Fatal population was distinct and most similar to *C. chathamensis* from San Cristóbal. Despite minimal morphological differences from *C. porteri*, and that the holotype of *C. porteri* is a hybrid with Reserva nuclear DNA and Cerro Fatal mtDNA, they described the Cerro Fatal population as a new species, and we tentatively agree.

17:62. ***Chelonoidis niger***: Olson (2015) investigated the history of some

early names applied to Galapagos tortoises. He concluded that the provenance of the single type specimen (MNHNP 9550) of both *Testudo californiana* Quoy and Gaimard 1824a and *Testudo niger* Quoy and Gaimard 1824b (= *Chelonoidis niger*) was "extremely unlikely" to be determined based on historical information, and hence the names should be considered *nomina dubia* and their use abandoned. Olson also concluded that one of the two syntypes of *Testudo nigrita* Duméril and Bibron 1835, designated the lectotype by Günther (1875a:268), was apparently lost, and since the paralectotype (MNHNP 9313) is a juvenile, he also recommended treating that name as a *nomen dubium*.

In addition, Olson and Humphrey (2017) investigated the origin of the type specimen of *Testudo elephantopus* Harlan 1827, and concluded that it may have come from Charles Island (Floreana), and suggested that the name *elephantopus* therefore replace the name *Testudo niger* Quoy and Gaimard 1824b, currently used for that island's species.

However, genetic analysis of the types currently in progress will hopefully demonstrate their geographic origins satisfactorily, and the allegedly lost lectotype of *nigrita* (NHMUK 1949.1.4.37) actually remains extant at the British Museum (P. Campbell, in litt., 2017). We therefore consider all these recommendations by Olson (2015) and Olson and Humphrey (2017) as premature and unnecessary at this time.

17:63. ***Chelonoidis phantasticus***: In our previous checklists we indicated that this species was Extinct, based on that supposition by Pritchard (1996), but its formal conservation status has not previously been assessed using IUCN Red List criteria (IUCN 2001). This has now been done by the TFTSG using updated Red Listing guidelines in regard to determining whether a species is actually Extinct or not (IUCN 2016). Although only a single individual of this species has ever been collected (in 1906), sightings and signs from 1964, 2009, and 2013 suggest that a few individual tortoises may remain extant in the exceedingly fragmented and hard-to-access landscape of Fernandina, most of which is covered by uninhabitable lava flows (A. Rhodin, L. Cayot, J. Gibbs, R. Kiester, and W. Tapia, in press). As such, the TFTSG now determines this species to be Critically Endangered (Possibly Extinct) and has submitted that assessment to the IUCN Red List for publication.

17:64. ***Chersobius, Homopus,* and *Psammobates***: Based on three mitochondrial and two nuclear genes, Hofmeyr et al. (2017) resolved a paraphyletic *Homopus* with respect to *Chersina*, and resurrected the genus *Chersobius* for the five-toed species (*signatus*, *boulengeri*, and *solus*) formerly in the genus *Homopus*, and restricted the genus *Homopus* to the four-toed species (*areolatus* and *femoralis*). We have tentatively accepted those changes here. The authors also noted strong support (Hofmeyr and Daniels, in prep.) for deep genetic divergence among the currently recognized subspecies of *Psammobates tentorius*, suggesting that they deserved species status. They also indicated the presence of phylogeographic structure within *Chersina angulata* and *Homopus signatus*, suggesting possible future taxonomic changes in those taxa. We await these forthcoming publications before making any further changes to the checklist.

17:65. ***Geochelone elegans***: Schweigger (1812:325) described *Testudo stellata* based on the same specimens and plate previously described and figured by Schoepff (1795:111) as *Testudo elegans*. However, Schweigger coined the new name *stellata* as a *nomen novum* because Schoepff had based his description of *elegans* partly on a Seba (1734) specimen (pl. 79, fig. 3) that Schweigger (1812:325) concluded had been incorrectly identified, and he instead synonymized that figure under *Testudo rotunda* Latreille. Schweigger therefore coined a new name for Schoepff's species, renaming it *stellata* and stating: "Habitat in India orientali. (Schoepf. sub falso nomine: test. elegans Seb.)". However, the description of *T. elegans* by Schoepff was nomenclaturally available, and therefore *T. stellata* Schweigger has the same type specimen as *T. elegans*, according to Article 72(e) of the ICZN Code, and thus becomes its objective junior synonym.

17:66. **Date of Publication of Blyth 1854 ["1853"]**: The date of publication of this article in issue number 7 of volume 22 of the Journal of the Asiatic Society of Bengal has historically been listed as 1853, the printed date on its page-headers. However, a note in the Proceedings published in the same issue 7 on p. 684 is dated 4 January 1854, indicating that the issue was actually not published until 1854, consistent also with the imprint date of 1854 on the bound volume of all 7 issues of volume 22. This change of date affects the following taxa: *Testudo phayrei* [= *Manouria emys phayrei*], *Testudo elongata* [= *Indotestudo elongata*], *Testudo megalopus* [= *Geochelone elegans*], *Testudo iberia* [= *Testudo graeca ibera*], and *Homopus burnesii* [= *Testudo horsfieldii horsfieldii*].

17:67. ***Gopherus***: In the recent book on the biology of North American tortoises (edited by Rostal et al. 2014), Bramble and Hutchison (2014) reviewed and augmented the previous morphological data for the five known living species at that time, as well as the fossil forms, and concluded that the available evidence (including molecular data) supported the reciprocal monophyly of *Xerobates*, including *agassizii*, *berlandieri*, and *morafkai*, and *Gopherus* (*sensu stricto*), including *flavomarginatus* and *polyphemus*. They recommended recognition of *Xerobates* and *Gopherus* as sister genera. In the same volume, Murphy (2014) provided a critical review of the systematics of the broad genus *Gopherus* and its members, and also concluded that *flavomarginatus* and *polyphemus* are sister taxa, and that *berlandieri* is sister to the *agassizii* group (including *morafkai*; see also Reid et al. 2011, not cited in Rostal et al. 2014). However, he made no recommendation about the use of *Xerobates*, and included all species in the broader genus *Gopherus*, as most authors since Crumly (1994) have done. Although most of the other authors of chapters in this book used the genus *Gopherus* (*sensu lato*), in his review of fossils, Franz (2014) recognized *Xerobates* and *Gopherus*. As noted by Murphy (2014), broad nuclear sampling is needed to resolve definitively the relationships within this species group. Until such time, and considering that the name *Gopherus* could still apply to all currently included species even if the two identified clades are reciprocally monophyletic, we retain *Gopherus* as the sole genus name.

17:68. ***Gopherus evgoodei***: The genetic distinctiveness of the southern Sonoran / northern Sinaloan population of Desert Tortoises has long been recognized (reviews in Edwards et al. 2015a,b, 2016), and has now been formally described as a new species by Edwards et al. (2016).

17:69. ***Indotestudo forstenii***: The type specimen of *I. forstenii* was recorded by Schlegel and Müller (1845) as having been collected by Forsten on "Gilolo, Indischen Archipel" (now Halmahera, Indonesia). However, no further specimens of *I. forstenii* have ever been recorded or found anywhere on that island since that time (D. Iskandar, A. Riyanto, pers. comm.), and the species does not appear to occur on Halmahera or its adjacent islands. The collector, Eltio Alegondas Forsten, spent a few months in 1841 collecting botanical and zoological specimens on Ternate island just off the west coast of Halmahera, where his collectors also obtained some material from mainland Halmahera. However, he also made extensive collections later in 1841–1842 on northern Sulawesi, while headquartered at Gorontalo (http://www.nationaalherbarium.nl/FMCollectors/F/ForstenEA.htm), where *I. forstenii* does occur. We hypothesize that the type specimen was therefore probably obtained by Forsten near Gorontalo, Sulawesi, Indonesia, and that it was then mislabeled as to its exact origin, or possibly that it was acquired from Ternate or Halmahera as a regionally traded specimen originating from Sulawesi. We no longer consider Halmahera to be part of the natural range of *I. forstenii*, and hereby formally restrict its type locality to "near Gorontalo, Sulawesi, Indonesia".

17:70. ***Testudo***: Based on a phylogenetic analysis of morphological traits for twelve fossil and the five extant species in the genus *Testudo* (sensu lato), Luján et al. (2016) recognized three monophyletic subgenera: *Chersine* for *hermanni*, *Agrionemys* for *horsfieldii*, and *Testudo* for *graeca*, *marginata*, and *kleinmanni*. Additionally, Vasilyev et al. (2014) studied variation in the 12S rRNA mitochondrial gene and three RAPD markers among individuals of a number of species of *Testudo* (sensu lato). Although *graeca*, *marginata*, *kleinmanni* (including synonymous *werneri*), and *horsfieldii* were resolved as monophyletic in the mtDNA analysis, *hermanni* was not; however, the RAPD analysis resolved each of those taxa as monophyletic. Support for the recognition of three monophyletic clades (corresponding to *Chersine*, *Agrionemys*, and *Testudo* [sensu stricto]) was ambiguous, with no support from the mtDNA, and only weak support from the RAPD data. Based on the combination of morphology of extant and fossil species and acknowledging the results of molecular genetics studies of *Testudo* by others, Luján et al. (2016) recommended that *Chersine* and *Agrionemys* (as well as the fossil *Paleotestudo*) be recognized as valid subgenera of *Testudo* (sensu lato). We now adopt that recommendation here, discontinuing our previous listing of alternative generic designations.

17:71. ***Testudo (Testudo) graeca***: In a paper previously not recorded in our checklist, Türkozan et al. (2010) analyzed morphometric variation in a large sampling of *Testudo graeca* from throughout Turkey. They determined that the putative subspecies *anamurensis* and *antakyensis* are not distinct from each other or from *terrestris* (and therefore synonymized under *terrestris*), and that *armeniaca* and *perses* are distinct. Further, they noted that the populations from northern and southern Turkey, possibly corresponding to the *ibera* and *terrestris* mtDNA clades, respectively, also appeared to be morphometrically distinct. They made no taxonomic recommendations other than maintaining a conservative approach pending additional morphologic and genetic assessment

of more material from both within and outside Turkey. Our checklist has reflected these conclusions for many years, although we have listed *perses* as a synonym under the distinct taxon *buxtoni* since 2007, a placement reconfirmed by Parham et al. (2012) (see TTWG annotations 07:73 and 12:36).

Our current updated map with its subspecies delineations has benefitted from review and input by Oguz Türkozan and Peter Mikulíček. Notably, unpublished mtDNA genetic work in progress by Türkozan and colleagues now appears to confirm the earlier impression that *ibera* and *terrestris* represent well-separated northern and southern populations of *T. graeca*. Their distribution patterns reflect clear separation by the so-called Anatolian Diagonal, with only a few areas of intergradation in southwestern Turkey. Additionally, *buxtoni* appears to intergrade slightly with *terrestris* in the southeastern corner of Turkey.

Using the cytochrome b gene, Javanbakht et al. (2017) examined the phylogeography of the subspecies of *T. graeca* in Iran and Transcaucasia and refined the knowledge of their distribution. Based on species distribution models, they showed that the distribution ranges changed little since the Last Glacial Maximum.

Using three mitochondrial DNA fragments (cytochrome b, 12S, ND4 plus adjacent DNA coding for tRNAs), Graciá et al. (2017a) studied differentiation of the *Testudo graeca* complex. According to fossil-calibrated molecular clock calculations, they inferred a dual diversification burst. The eastern subspecies, including the last common ancestor of the North African taxa, radiated in the Mio-Pliocene, whereas a second radiation in North Africa took place during the Pleistocene. Based on a Libyan tortoise of unknown exact provenance, a new North African lineage was discovered. The recent introduction of most Western European populations was confirmed, with the exception of populations in southeastern Spain, which are older.

17:72. ***Testudo (Testudo) graeca* and *T. (T.) g. terrestris***: Meiri et al. (2011) examined body size variation in *Testudo graeca* across the Levant, and demonstrated that these tortoises follow Bergmann's Rule, with the smallest tortoises found at the southern end of the range in the Negev Desert. The small Negev tortoises had earlier been decribed as *T. floweri* by Bodenheimer (1935), and these authors agreed that this taxon is synonymous with *T. graeca*. Werner et al. (2016) then expanded that analysis to include all *T. graeca* (sensu lato; see TTWG 2014) from Morocco to Iran. They also found a general correlation of body size with latitude. However, their analyses of sexual size dimorphism demonstrated that it seems to scale differently among tortoises in Anatolia versus the Levant, and concluded that this might suggest that those two populations might deserve consideration as "separate entities" (presumably with taxonomic consequences).

17:73. ***Testudo (Testudo) kleinmanni***: Based on morphological differences between *Testudo kleinmanni* on opposites sides of the Nile River, Perälä (2001) described the eastern population as *T. werneri*. Subsequent studies (see TTWG 2007, annotation 07:74) did not support this distinction and *werneri* was synonymized with *kleinmanni*. However, Werner (2016) has re-emphasized the apparent morphological differences between the populations and has claimed that some tortoises from west of the Nile have been translocated to the east to northern Sinai, potentially confounding the recent studies. Therefore, he suggested that *werneri* should be recognized as a distinct subspecies. Until further genetic work tests this scenario, we continue to retain *werneri* in synonymy.

17:74. ***Testudo (Chersine) hermanni***: Based on an analysis of 17 nuclear microsatellite loci and broad geographic sampling, Zenboudji et al. (2016) identified three major genetic clusters within *Testudo hermanni hermanni*: one continental (Spain, France and Italy), one insular (Corsica, Sardinia, and Sicily), and one an island supporting two clusters (Menorca). There is still debate as to which populations are autochthonous and which are the result of human introductions (or reintroductions). Perez et al. (2014; see also TTWG 2014, annotation 14:36), based on mtDNA and microsatellites, concluded that the Spanish, Corsican, and Sardinian populations were likely introduced from Sicily by humans. However, although Zenboudji et al. (2016) agreed that the contemporary population on Sardinia is the result of introductions from Sicily following an earlier extirpation on Sardinia, they argued that the Corsican population is natural and genetically distinct. They also concluded that the population in northeastern Spain (Albera) is a relict, natural population. Menorca supports two distinct genetic populations. The one in the west is closely related to mainland populations, and hence is likely an introduction from a now extirpated mainland source. The eastern population is more distinct, but more closely related to the other insular populations, but whether this represents an ancient autochthonous divergence or an ancient human introduction of a lineage of unknown provenance could not be determined. Zenboudji et al. (2016) also noted a number of examples of mismatched genotypes in several populations, representing introduced individuals (and their offspring) from other populations. Finally, they recommended that six populations be considered genetic management units for conservation purposes: 1) Albera (Spain) + western Menorca; 2) France (Var); 3) Italy; 4) Sardinia + Sicily; 5) Corsica; and 6) eastern Menorca.

17:75. ***Cyclanorbis senegalensis***: Based on mitochondrial DNA sequence data, Mazuch et al. (2016) reported the presence of *C. senegalensis* in western Ethiopia for the first time, and also found significant genetic divergence between western (Togo and Benin) and eastern (Ethiopia) populations. They called for further phylogeographic study, but made no taxonomic recommendations.

17:76. ***Amyda cartilaginea* or *A. ornata***: Fritz et al. (2014a) examined variation in mitochondrial and nuclear DNA and color patterns across the range of *Amyda cartilaginea* (sensu lato). Their mtDNA tree was not highly resolved; however, because of the concordance of the identified clades with color pattern and geography (paleodrainages), the authors recognized two named species with four named subspecies within what was previously recognized as *A. cartilaginea*. They restricted *A. cartilaginea* (sensu stricto) to the southern portion of the Greater Sundas, with the nominate subspecies in southern Borneo and Java, and a new subspecies, *maculosa*, in the northern portion of the Greater Sundas (northwestern Borneo and southern Sumatra). Populations of *A. cartilaginea* (sensu lato) in northern Sumatra and Peninsular Malaysia were not assessed or assigned to subspecies. They also identified, but did not describe, a potential candidate species from northeastern Borneo. They resurrected the older name *ornata* (Gray 1861) for the Southeast Asian mainland species, with the nominate subspecies confined to the Mekong basin, and resurrected the name *phayrei* (Theobald 1868) for the subspecies in Thailand and Myanmar. They also identified, but did not describe, a third possible subspecies from Bangladesh. Regrettably, no specimens from southern Myanmar, southern peninsular Thailand, or Peninsular Malaysia were included. In view of the relatively weak resolution of their named mtDNA clades, but reasonable concordance of those clades with paleogeographic drainage basins and color patterns, we list their proposed taxonomic changes provisionally, pending additional geographic sampling and further genetic and morphologic work.

17:77. ***Apalone mutica calvata***: In a guide to Alabama turtles, Guyer et al. (2015) elevated *Apalone mutica calvata* to a full species based on its distinctive color pattern, its geographic isolation, and the mtDNA data (part of only one gene) presented by Weisrock and Janzen (2000). Powell et al. (2016) followed their recommendation. However, because of the small sample size and small DNA fragment available to Weisrock and Janzen (2000) and incomplete sampling in the presumed area of overlap in southeastern Louisiana, we consider this change premature, and continue to recognize *calvata* as a subspecies of *A. mutica*.

17:78. ***Nilssonia* and *N. nigricans***: Based on mitochondrial Cytochrome C Oxidase Subunit I (mtCOI) barcode sequences, Kundu et al. (2016) provided a phylogenetic analysis of the Asian softshell turtles allied with the genus *Nilssonia*. Their results confirmed the presence of *N. nigricans* in the wild in Assam, India, as well as provided additional support for the synonymy of *Aspideretes* with *Nilssonia* (see also TTWG 2011, annotation 11:15).

17:79. [*Annotation text previously included here now deleted from this checklist*.]

17:80. ***Pelodiscus sinensis* and *P. maackii***: Suzuki and Hikida (2014) examined mitochondrial cytochrome b sequence data from specimens of *Pelodiscus* across Japan and compared them with previously published data from Fritz et al. (2010b) from across the wider species range. They identified two lineages from Japan, corresponding to the previously recognized *P. sinensis* and *P. maackii*. The latter was widely distributed in Japan, whereas the former had only a sporadic distribution, leading the authors to conclude that *maackii* was native to Japan and *sinensis* was introduced. They also noted that if future nuclear DNA data supported this scenario, the older name *japonicus* (Temminck and Schlegel 1838) could be available for the lineage currently recognized as *maackii* (Brandt 1857).

17:81. ***Pelodiscus sinensis***: The publication date of F.J.F. Meyen's *Reise um die Erde*, in which *Trionyx (Aspidonectes) sinensis* (= *Pelodiscus sinensis*) was first described by A.F.A. Wiegmann, has been a source of controversy. Although frequently cited as appearing in 1835, Bauer and Adler (2001) in a previously overlooked publication determined that the description of *sinensis* first appeared in print in 1834, and we have now corrected that date here.

17:82. ***Pelodiscus sinensis***: As clarified by Adler (2016), the name *Trionyx tuberculatus* was first used by Cantor (1842a), but without a description or indication, and thus a *nomen nudum*; then subsequently formally and validly described a few months later (Cantor 1842b).

17:83. ***Rafetus euphraticus***: We note that in the original description of *Testudo euphratica* published by Daudin (1801) that he specifically credited Olivier with providing him with the description of the new species. We therefore amend the authorship of the original name to *Testudo euphratica* Olivier *in* Daudin 1801.

17:84. ***Rafetus euphraticus***: Ihlow et al. (2014) examined variation in two mitochondrial genome fragments across nearly the entire range of the species, and found no significant genetic variation.

17:85. ***Rafetus swinhoei***: Le et al. (2014) analyzed sequence data from two mitochondrial loci and one nuclear gene from all known populations of *Rafetus swinhoei* in China and Vietnam. Their results demonstrated minimal divergence among populations, and warranted no taxonomic changes. Despite the long history in China of human use and transport of turtles, the authors cautioned against speculation that the Chinese distribution is unnatural.

17:86. **Chelidae**: A few recent authors have followed Storr (1978) in using the family name Cheluidae (Michael and Lindermeyer 2010; Wilson and Swann 2013), a grammatically correct derivation from the Greek type genus *Chelus*, genitive singular *Cheluos*. However, the spelling Chelidae has been in prevailing usage globally since Lindholm (1929) and most authors continue to use it. Under ICZN Article 29.5, the original spelling of a family-level name is to be maintained when it is in prevailing usage, whether or not its derivation from the name of the type genus is in accordance with grammatical procedures. We therefore regard the name Cheluidae as an unjustified emendation, given the long-standing prevalence of Chelidae in the literature, and we strongly recommend and concur with the continued usage of Chelidae.

17:87. ***Acanthochelys radiolata***: Garbin et al. (2016) examined morphological variation in *Acanthochelys radiolata* across its range. Although they described significant variation in this species (especially in color and shape), they could identify no geographic pattern to this variation and concluded that *A. radiolata* is a single, highly variable species. In addition, they considered records from Mato Grosso and Sao Paulo States to be in error.

17:88. ***Chelus fimbriata***: Zug (1977) clarified that the genus name for the matamata is *Chelus*, not *Chelys*, as originally proposed by Dumeril (1805:76). Most authors since then have treated the generic name as feminine, rendering the species name as *Chelus fimbriata*. However, Ferreira et al. (2016), following the recommendation of Vlachos (cited as a pers. comm., but also discussed in Vlachos 2015), argued that the genus name *Chelus* is actually masculine (and the species name therefore *Chelus fimbriatus*) because it was supposedly based on Latinization of the classical Greek feminine word *χέλυς*, and according to ICZN (1999) Art. 30.1.3, a Latinized suffix should take the gender of the Latin suffix. However, according to Appendix B of the 1961 and 1985 ICZN Codes, as well as Art. 30.1.2 and 30.1.3 of the 1999 Code, there is a distinction between transcription or transliteration vs. Latinization of a Greek word. Specifically, for the Greek letter *upsilon* (*υ*), its Latin equivalent (i.e., transliteration) is '*u*', whereas the Latinized version is '*y*' (ICZN 1985, App. B). Hence, the name *Chelus* is considered a transliteration, not a Latinization, of *χέλυς*, whereas *Chelys* is considered a Latinization. Unfortunately, Vlachos (2015) transposed the terms "transliteration" and "Latinization" in his Table 1—he rightly used "Latinized form" to name his column L (where he gives "*y*" as the Latinized form of *upsilon*), but used "transliteration" for it in the caption, causing considerable confusion. *Chelus* and *Chelys* should not be considered homonyms (1999: Art. 56.2; 1985: Art. 56.b), and in accordance with ICZN 1999 Art. 30.1.2 (1985: Art. 30a), the gender of both *Chelus* and *Chelys* is feminine. The correct species name is therefore *Chelus fimbriata*, as it has been rendered for several decades, and we continue to use it as such.

17:89. ***Mesoclemmys raniceps***: Rivas et al. (2015) discussed the alleged distribution of this species in Venezuela, concluding that of the three specimens recorded from there, two were misidentified, and one (the type specimen of *Hydraspis maculata*), was misidentified and incorrectly restricted to Venezuela. Hence, we delete Venezuela from the range of *M. raniceps*, and synonymize *H. maculata* under *M. raniceps* instead of under *M. nasutus* where it was previously placed.

17:90. ***Phrynops geoffroanus* and *P. tuberosus***: In a previous checklist (TTWG 2010, annotation 10:44), we noted that the distribution and taxonomy of these two taxa remained highly problematic and subject to differing opinions as to their extent and inter-relationships, and in our last checklist (TTWG 2014) we provided coarse and uncritical distribution maps for them. In this checklist we now present new revised distributional maps that document most known localities of these taxa, sourced from a combination of the EmySystem database (with several corrections), recently published literature, and previous research on many collected specimens by Rhodin and Mittermeier during their work on the *P. geoffroanus* complex (Rhodin and Mittermeier 1983; A.G.J. Rhodin and R.A. Mittermeier, unpubl. data). Morphological analysis at that time indicated the presence of several apparently allopatric and differentiated populations of *P. geoffroanus* corresponding to various level 3 and 4 hydroshed basins, with several areas of close parapatry and possible intergradation. We now depict these distributional patterns in our revised maps, showing the distributional limits of the two taxa to delineate the hydroshed-restricted extent of their mostly separate populations, and showing the various subpopulations of *P. geoffroanus*. Many of these differentiated populations have now also been preliminarily substantiated as genetically distinct lineages by Carvalho et al. (2016). We anticipate that the *P. geoffroanus* complex will eventually be recognized as a polytypic species complex with several distinct lineages, notably the southeastern Brazilian coastal, the Rio São Francisco basin, the Rio Paraná basin, the lower Amazon basin, the Colombian, and the Peruvian-Bolivian populations.

17:91. ***Platemys platycephala melanonota***: Mendes-Pinto et al. (2011) reported the collection of a *Platemys platycephala* in southwestern Pará, Brazil, that was diagnosable as the subspecies *P. p. melanonota*. This record lies nearly 2000 km from the known range of *melanonota*, well within the known range of the nominate subspecies, and calls into question the validity of the subspecies *melanonota* which was described based primarily on color patterns. A re-evaluation of geographic variation and genetics in this species is needed.

17:92. ***Chelodina (Chelodina) longicollis***: Hodges et al. (2015) studied mitochondrial phylogeography of *Chelodina longicollis*, a highly terrestrially mobile freshwater species, to determine if its population genetic structure would correspond to hydrological boundaries or not. They found two ancient haplogroups broadly with east-west partitioning across the Great Dividing Range, but made no taxonomic recommendations. Each haplogroup was characterized by complex genetic structure, demographically stable subpopulations, and signals of isolation by distance; but the patterns were also overlaid with signatures of introgression and recent gene flow, likely facilitated by late Pleistocene and ongoing anthropogenic landscape changes.

17:93. ***Chelodina (Macrochelodina) expansa***: Hodges et al. (2014) carried out a phylogeographic study of *Chelodina expansa* based on mitochondrial gene variation to identify two major clades of mitochondrial haplotypes. The first comprised populations from the inland Murray-Darling Basin and the Mary River in southeast Queensland; the second comprised populations from coastal catchments north of the Mary River. They did not regard it as appropriate to provide taxonomic recognition for these populations east and west of the Great Dividing Range (as previously proposed for other similarly distributed taxa by Cann 1998), because the mitochondrial analysis demonstrated that the morphological variation observed was not concordant with the spatial population structure defined by the molecular data (views subsequently discussed further by Spinks et al. 2015). [*Annotation text previously included here now deleted from this checklist.*]

17:94. ***Chelodina (Macrochelodina) kuchlingi* and *C. walloyarrina***: Both of these purported taxa remain enigmatic and poorly known, with their apparent distinctiveness incompletely evaluated, and we remain uncertain whether they are indeed valid and recognizable taxa (as either species or subspecies), or poorly differentiated lineages of *C. oblonga* (formerly *C. rugosa*) and *C. burrungandjii*, respectively, or possibly their hybrid intergrades. Ellis and Georges (2015), in their catalogue of turtle type specimens held at the Western Australian Museum, synonymized *C. kuchlingi* under *C. oblonga* and *C. walloyarrina* under *C. burrungandjii*, following the earlier recommendation by Georges and Thomson (2010), but their action was not based on new data or analysis. However, we note that prior to the description of *Chelodina walloyarrina* from the Kimberleys by McCord and Joseph-Ouni (2007b), and not addressed in our earlier comment (TTWG 2010, annotation 10:38), is that Thomson et al. (2000) noted morphological differences between Kimberley specimens (= *C. walloyarrina*) and *C. burrungandjii* from Arnhem Land, while Georges et al. (2002) compared the still undescribed Kimberley form and *C. burrungandjii* from Arnhem Land species using allozyme data (45 independent nuclear loci), which showed no fixed differences between these populations. As we noted in our prior checklist (TTWG 2014, annotation 14:42), further analysis of the phylogeographic relationships among all these lineages is still in progress by Kuchling, Georges, and others, and based on our precautionary

principles in regard to data-driven analysis in the recognition and protection of biodiversity, we remain reluctant to formally synonymize these taxa until more conclusive genetic and morphologic data emerge.

17:95. ***Elseya***: Thomson et al. (2015) divided the broad genus *Elseya* into three subgenera based on genetics and skeletal morphology: *Elseya (Elseya)*, *Elseya (Hanwarachelys)*, and *Elseya (Pelocomastes)*.

17:96. ***Elseya flaviventralis***: Allopatric populations of the *Elseya dentata* group from the Arnhem Land region of Northern Australia were first suggested to be unique by Legler (1981) based on morphology. Subsequent allozyme electrophoretic analyses by Georges and Adams (1992, 1996) and morphological work by several authors (reviewed in Thomson and Georges 2016) supported the distinction of these populations at the species level, and Thomson and Georges (2016) formally described this species as *Elseya (Elseya) flaviventralis*, which we accept.

17:97. ***Elseya rhodini***: As indicated in our last checklist (annotation 14:45), the population of *Elseya* (formerly included under *E. novaeguineae*) from the southern versant of New Guinea, previously noted by Rhodin and Genorupa (2000) and by Georges et al. (2014) to be genetically and morphologically differentiated from other New Guinean *Elseya*, has since been formally described as *Elseya (Hanwarachelys) rhodini* by Thomson et al. (2015). We accept this new species as distinct.

17:98. ***Emydura macquarii krefftii***: Todd et al. (2014a) examined variation in mitochondrial and nuclear microsatellite markers across the widespread range of *Emydura macquarii krefftii*. The mitochondrial data revealed 1) a distinct divergence between northern (Burdekin River northward) and southern (Fitzroy River and southward) populations of *E. m. krefftii*, 2) that *E. m. emmotti* was most closely related to but highly divergent from the northern *krefftii* clade, 3) that *E. m. nigra* and *E. m. macquarii* were nested within the southern *krefftii* clade, and 4) that populations in the upper Burnett River were highly divergent from other southern clade populations. Their examination of twelve microsatellite loci within only *krefftii* also demonstrated support for the same north-south divergence as for the mtDNA data, but did not support the distinction of the upper Burnett population (possibly a human translocation, according to the authors), or the taxonomic distinction among the subspecies *nigra*, *macquarii*, and southern *krefftii*. They made no explicit taxonomic recommendations, but did note that the north-south divergence was within the range exhibited for other accepted chelonian species. Further analysis is clearly indicated.

17:99. ***Myuchelys purvisi***: Legler (1981) foreshadowed splitting the genus *Elseya* into two major clades, one containing *Elseya dentata* and related species, the other containing *Elseya latisternum* and its close relatives, many of which were undescribed at that time. A subsequent study based on 54 allozymes loci (Georges and Adams 1994) established that Legler's "*latisternum* group" was indeed a clade (monophyletic), and this was the foundation for the description of the new genus *Myuchelys* (Thomson and Georges 2009) with four contained species: *M. purvisi, M. georgesi, M. bellii*, and *M. latisternum* as type. Unfortunately, morphological characters that diagnose that genus are symplesiomorphies.

Two subsequent analyses based on a limited set of nuclear and mitochondrial sequence data were equivocal on the monophyly of *Myuchelys*. Georges et al. (1998) recovered *Myuchelys* as paraphyletic with respect to *Elseya* based on analyses of two mitochondrial genes (12S rRNA and 16S rRNA), but without statistical support. More recently, analyses of a single nuclear locus (c-mos) provided moderate support (83% bootstrap support values) for grouping *Emydura macquarii*, *Myuchelys latisternum*, and *M. georgesi* as a clade to the exclusion of *M. purvisi*, a result confirmed by analysis of mtDNA (Fielder et al. 2012). However, Georges and Adams (1994), Georges et al. (1998), and Fielder et al. (2012) all maintained that the uncertainty surrounding incongruence among these analyses should preclude taxonomic revisions and therefore did not propose revisions to correct the potential paraphyly of *Myuchelys* with respect to *Emydura*.

Subsequently, Le et al. (2013) generated phylogenies for the chelid genera *Elseya, Emydura, Myuchelys, Elusor,* and *Rheodytes* using two mtDNA markers and a single nuDNA marker. Their phylogeny also recovered *Myuchelys* as paraphyletic, again owing to the position of *M. purvisi*, and they assigned *purvisi* to a new genus, *Flaviemys*, to maintain monophyly of *Myuchelys*. A more comprehensive analysis (Spinks et al. 2015) using 13 independent nuclear DNA markers recovered *Myuchelys*, including *M. purvisi*, as a well-supported clade, in agreement with the previous allozyme data. Thus, based on the weight of evidence, the taxonomic revision of Le et al. (2013), using their more limited sequence information, was considered by Spinks et al. (2015) to be premature.

On the basis of conflicting evidence from the mitochondrial and nuclear DNA evidence, and because *M. purvisi* and *M. georgesi* are morphologically similar and have been regarded as a cryptic species pair (but see Thomson and Georges 1996), we have retained *Myuchelys* based on the weight of evidence suggesting that it is a well-supported monophyletic group, and placed *Flaviemys* into its synonymy, following the recommendation of Spinks et al. (2015).

17:100. **Pseudemydurinae, *Pseudemydura, P. umbrina***: The affinities of *Pseudemydura* among the Chelidae are not well established. Early work using serological comparisons revealed that *Pseudemydura, Emydura-Elseya*, and *Chelodina* formed an unresolved trichotomy (Burbidge et al. 1974). Many of the defining morphological characters of *Pseudemydura* have been regarded as autapomorphies, and so not useful for phylogenetic analysis. Nevertheless, Gaffney (1977) placed *Pseudemydura* as sister to all the remaining extant chelids of both Australasia and South America (as the new subfamily Pseudemydurinae), while admitting that his case, based on a single retained primitive character, was weak. Subsequent DNA sequence studies were unable to resolve the conundrum. 12S mitochondrial rRNA data were unable to conclusively establish the relationships of *Pseudemydura*, but tentatively resolved it as sister to the other Australasian shortnecked genera—*Emydura, Elseya, Myuchelys, Rheodytes* and *Elusor* (Seddon et al. 1997). Additional sequence analysis from mt 16S rRNA and nuclear c-mos supported this arrangement, but bootstrap support remained poor (Georges et al. 1998). A more recent analysis, using the same data drawn from GenBank, resolved *Pseudemydura* as sister to *Chelodina* (Guillon et al. 2012). Subsequently, in a broad analysis of previously published sequence data representing 13 mitochondrial and nuclear DNA loci and including 83% of all extant turtle species (as per our checklist), Pereira et al. (2017) resolved *Pseudemydura* as sister to all short-necked Australasian chelids, with those two clades sister to *Chelodina*. Most recently, Zhang et al. (2017) analyzed the whole mitogenome of *Pseudemydura* and demonstrated it to be sister to all Australasian chelid turtles (subfamily Chelodininae), with strong bootstrap support; consequently, they proposed resurrection of the subfamily Pseudemydurinae Gaffney 1977, which we adopt here.

17:101. ***Pelomedusa***: Vargas-Ramírez et al. (2010) and Wong et al. (2010) reported deeply divergent mitochondrial clades within what was then recognized as the monotypic species *Pelomedusa subrufa*, with less, but concordant, variation in nuclear DNA markers. The extent of mitochondrial divergences of up to 20% of the cytochrome b gene exceeds pairwise divergences between any other congeneric chelonian species. Based on mitochondrial DNA variation, Petzold et al. (2014) and Nagy et al. (2015) identified at least thirteen terminal clades (see also Fritz et al. 2014b). Petzold et al. (2014) recommended full species status for some of these clades. They restricted *P. subrufa* (sensu stricto) to southern Angola, Botswana, southeastern Democratic Republic of the Congo, Madagascar (presumably introduced), Malawi, Namibia, South Africa, and Tanzania. Older names were resurrected from the synonymy of *P. subrufa* for three of the clades: *P. galeata* (Schoepff 1792; South Africa), *P. gehafie* (Rüppell 1835; Eritrea and possibly Sudan), and *P. olivacea* (Schweigger 1812; Benin, Burkina Faso, Niger, Nigeria, and Senegal). Six other species were newly described: *P. barbata* (Saudi Arabia and Yemen), *P. kobe* (Tanzania), *P. neumanni* (Kenya and Tanzania), *P. schweinfurthi* (Central African Republic and South Sudan), *P. somalica* (Somalia and Ethiopia; see also Fritz et al. 2015b), and *P. variabilis* (Ghana and Ivory Coast). The authors also identified two clades that they considered candidate species but did not describe: one from Cameroon, and the other from Sudan. Further candidate species could correspond to distinct clades within *P. galeata* and *P. somalica* (Fritz et al. 2015b), and Nagy et al. (2015) added another possible candidate species from the southeastern Democratic Republic of the Congo. All these studies, however, relied primarily on mitochondrial data and preliminary geographic sampling, but also included some morphologic work and sequencing and allocation of type specimens (see also Fritz et al. 2015a). We remain uncertain whether all taxa described and/or identified are valid and at what systematic level (species vs. subspecies vs. ESUs), pending corroborating data from nuclear genes. However, based on the data and extensive analyses presented, we tentatively accept the proposed taxonomic arrangement while urging and awaiting further work, notably analysis of nuclear loci, more detailed morphologic work, and further geographic sampling, especially from areas between assigned species. We document these unsampled and unassigned areas as *Pelomedusa subrufa* (sensu lato) species complex.

17:102. ***Pelusios***: Kindler et al. (2016) examined sequence variation in three mitochondrial and three nuclear genes in *Pelusios*, and found no

phylogenetic structure in *P. chapini*, *P. gabonensis*, or *P. nanus*; however, they identified significant structure within *P. rhodesianus* (with two deeply divergent clades), *P. carinatus*, and *P. castaneus*. Both their mtDNA and nDNA data also suggested that *rhodesianus* was paraphyletic with respect to *carinatus*. But because their geographic sampling was incomplete and there was discordance between their mtDNA and nDNA data, the authors made no new taxonomic decisions regarding these six taxa.

17:103. ***Pelusios castaneus seychellensis***: Kindler et al. (2016), in their wider phylogeographic study of *Pelusios*, found that their mtDNA data nested the lectotype of *P. seychellensis* deep within a variable *P. castaneus*, and sister to specimens from the Republic of Congo, agreeing with Stuckas et al. (2013), leading them to treat *seychellensis* as a junior synonym of *castaneus*. We addressed the status of *seychellensis* previously in annotation 14:47 and interpreted it at that time as a subspecies of *P. castaneus*. We maintain that interpretation at this time; however, we acknowledge the possibility that the types of *seychellensis* could conceivably have been mislabeled or based on transported specimens, in which case *seychellensis* should indeed be synonymized with *castaneus*.

17:104. ***Podocnemis erythrocephala***: Santos et al. (2016) examined variation in a single mitochondrial gene across the range of *Podocnemis erythrocephala*. They identified considerable genetic structure among populations, with four distinct genotypes that they deemed "management units," but made no taxonomic recommendations.

17:105. ***Podocnemis sextuberculata***: *Podocnemis sextuberculata* and *Pentonix americana* (*nomen dubium*) (= *Pelomedusa subrufa*) were first described on p. 13 of Cornalia (1849), but the descriptions (with minor changes) were reprinted on p. 312 in the Appendix of Osculati (1850), who collected the holotype of *sextuberculata* described by Cornalia. The two publications have created some confusion about dates and pagination for these two names, and we correct our previously cited paginations here.

2021 Checklist Annotations

TTWG 2021 (CRM 8, checklist.v.9) (21:1-105)

21:1. **New Distributional Data**: We have updated almost all our distributional maps, including adding separate maps for all subspecies; this checklist includes 492 new or revised maps with additional locality dots and/or adjusted distributional ranges. New or significantly revised maps (468) are marked with an asterisk; those with very minimal changes (24) are marked with a bullet.

We have utilized multiple sources for these updates, including expert input based on field data, localities from taxonomic literature cited in the new annotations below, new CBFTT species accounts, and IUCN Red List and TFTSG Provisional Red List assessments. We have also added several geographic distribution and range extension records from short notes or accounts in *Herpetological Review, Herpetology Notes, Check List, Mesoamerican Herpetology, Herpetozoa, Journal of Threatened Taxa, Catalogue of American Amphibians and Reptiles*, and other similar publications, including selected verified records from *iNaturalist.org*, but have not cited most of these individually.

In addition, we have added new or previously unrecorded locality records from several articles and books, both older and more recent (Said-Aliev 1979; Iverson 1992; Das 1995; Devaux 2000; Mitrus and Zemanek 2000; Bour et al. 2001; Bringsøe et al. 2001; Lee and Carey 2001; Lee and Ross 2001; Savage 2002; Spawls et al. 2002; Stebbins 2003; Zavialov et al. 2003; Khabibullin 2004; Široký et al. 2004; Chirio and Ineich 2006; Gong et al. 2006, 2009b, 2012; Rosen et al. 2006, Diesmos et al. 2008; Gramentz 2008; Schneider and Schneider 2008; Valakos et al. 2008; Avila-Pires et al. 2010; Fong and Qiao 2010; Largen and Spawls 2010; Auer 2011; Li et al. 2011; Bondarenko and Dujsebayeva 2012; Buckner et al. 2012; Trape et al. 2012; Ferri and Diagne 2013; Rodríguez Schettino et al. 2013; Safi and Khan 2014; Khan et al. 2015; Lemos-Espinal et al. 2015, 2017, 2019; Ramírez-González and Canseco-Márquez 2015; Sethy et al. 2015; Htun and Platt 2016; Keller et al. 2016; Powell et al. 2016; Trebbau and Pritchard 2016; Valdez-Villavicencio et al. 2016a,b; Alqahtani 2017; Bondarenko and Peregontsev 2017; Bury 2017; Cann and Sadlier 2017; Dujsebayeva et al. 2017; Ferrara et al. 2017; Ge et al. 2018; Kunz et al. 2018; Macip-Ríos et al. 2018; McCranie 2018; Jadhav et al. 2018; Murphy 2018; Platt et al. 2018; Rodrigues and Lima-Ribeiro 2018; Yang et al. 2018; Brito et al. 2019a; Forero-Medina et al. 2019; Ihlow et al. 2019; Krysko et al. 2019; Nekrasova et al. 2019; Petrozzi et al. 2019; Sánchez et al. 2019; Vamberger et al. 2019b; Xiaoyou et al. 2019; Yudha et al. 2019; Berry et al. 2020; Brown et al. 2020; Ihlow et al. 2020; Kagayama et al. 2020; Le Duc et al. 2020; Mital et al. 2020; Nguyen et al. 2020; Smith et al. 2020; Ayaz et al. 2021; Brown et al. 2021; Cunha et al. 2021; Eustace et al. 2021; Jain et al. 2021; Thapa et al. 2021; Vásquez-Cruz et al. 2021; Weitzman et al. 2021; Xiao et al. 2021).

21:2. **Testudines**: Joyce et al. (2021) provided a detailed phylogenetic nomenclature (PhyloCode) for all fossil and living turtles, in which they largely followed our TTWG (2017) nomenclature for the higher categories of all living turtles. However, they did not recognize the clades of Cheloniinae, Carettinae, Geoemydinae, or Rhinoclemmydinae that we accept as subfamilies, but recognized Testudininae, Podocnemidinae, and Erymnochelyinae, and named the new clades Peltocephalinae and Emydurinae. We continue to use traditional ICZN-compliant Linnaean ranking categories and nomenclature, but recognize the hierarchical value and utility of the PhyloCode system, and have now recognized both Peltocephalinae and Emydurinae as valid subfamily names in our Linnaean hierarchy (see next Annotation).

21:3. **Testudines**: Thomson et al. (2021) published a comprehensive global phylogeny with estimated divergence times for nearly all turtle lineages, based on new nuclear DNA sequence data from 15 nuclear loci for 591 individual turtles, covering all 14 extant families, nearly all extant genera (lacking only *Pseudemydura*, *Rhinemys*, and *Natator*), and 279 of the 348 (80%) extant species of turtles recognized by TTWG (2017). Thomson and colleagues demonstrated that both suborders Cryptodira and Pleurodira are monophyletic and strongly supported; they additionally noted deeply divergent monophyletic separation of all hardshelled Cryptodira, for which they used the PhyloCode name Durocryptodira, from the softshelled Trionychoidea, for which they used the PhyloCode name Trionychia. Since neither PhyloCode names nor Linnaean names above the family-group level are regulated by the ICZN, we now use these terms to represent infraorders of Cryptodira to reflect this deep phylogenetic divergence. We have also changed the order in which we place these suborders within our checklist, now placing Pleurodira first and Cryptodira second.

In the pleurodiran Podocnemididae, Thomson et al. (2021) recovered three recognized subfamilies—Erymnochelyinae, Peltocephalinae, and Podocnemidinae—we have therefore now also incorporated these into our checklist. They also recovered a monophyletic clade for Podocnemididae and Pelomedusidae as sister to the Chelidae and referred to it as Pelomedusoides, the hyperfamily name proposed by Broin (1988); we instead utilize the superfamily name Pelomedusoidea for that clade. In addition, we use the adjusted superfamily name Cheloidea for its sister clade containing the Chelidae, instead of the hyperfamily name Cheloides proposed by Gaffney et al. (2006).

Thomson et al. (2021) demonstrated the monophyly of the clade containing Chelydridae, Dermatemydidae, and Kinosternidae, supportive of an expanded Chelydroidea, as previously recommended by Crawford et al. (2015), which we now adopt, removing the previous concept of a more restricted Kinosternoidea, as recommended by Joyce et al. (2004) and others.

Thomson et al. (2021) also demonstrated that all currently defined extant families are monophyletic and strongly supported, including well-supported subfamilial diversification within several families. In the Geoemydidae they recovered three strongly supported separate clades, including the previously recognized subfamily Rhinoclemmydinae, but their results separated the old concept of Geoemydinae into two reciprocally monophyletic clades corresponding to the subfamilies Batagurinae and a restricted Geoemydinae. We have therefore incorporated these subfamilies into the checklist and rearranged the respective genus and species placements accordingly.

In the Testudinidae, Thomson et al. (2021) recovered three reciprocally monophyletic clades corresponding to the core Testudininae plus a separate clade for *Gopherus*, which we allocate to the previously recognized subfamily clade Xerobatinae, and a clade for *Manouria*, which we designate as the subfamily Manouriinae (type genus *Manouria* Gray 1854a), based on the previous family-group level concepts of Manourina Gray 1869a and Manouriina Baur 1888b.

At the genus level, Thomson et al. (2021) found several cases of non-monophyly that warrant further discussion. They confirmed that *Chersina* rendered the older concept of a broad *Homopus* (with five included species)

paraphyletic, corroborating the results of Hofmeyr et al. (2017), but recommended combining these two genera into a broader *Chersina*. However, we note the strong evidence of divergence within *Homopus* presented by Hofmeyr et al. (2017) and Hofmeyr and Branch (2018), and instead accept their recommended split of *Chersobius* from *Homopus* and the recognition of a monotypic *Chersina* (see also TTWG Annotations 88 and 17:64).

Similarly, Thomson and colleagues confirmed that the monotypic *Stigmochelys* is sister to a monophyletic *Psammobates*. They recommended combining these two genera into a broader *Psammobates* to convey their close evolutionary relatedness. However, in view of the marked morphological differences between *Stigmochelys* and *Psammobates*, we continue to recognize them as separate genera, as per our previous TTWG Annotations (07:71 and 10:26).

In addition, the analysis by Thomson and colleagues rendered the South American chelid genus *Mesoclemmys* paraphyletic with respect to *Phrynops* by the recovery of *M. hogei* as sister to the remaining species of *Mesoclemmys* and *Phrynops*. The divergence of *M. hogei* had previously been suggested by McCord et al. (2001) who established *Ranacephala* as a monotypic genus for the species, separating it from *Phrynops*; however, that action was not accepted by Joyce et al. (2004) or Bour and Zaher (2005), who instead assigned *hogei* to *Mesoclemmys*, which we accepted (TTWG 2007b, TTWG Annotation 07:100). However, in view of the strong nuclear genomic data recovered by Thomson et al. demonstrating the deep divergence of *hogei* from both *Mesoclemmys* and *Phrynops*, as well as the morphological details recorded by McCord et al. (2001), the combined morphological and genetic results of Holley et al. (2020), and the osteological distinctiveness preliminarily noted in Reed et al. (1991), we have now re-assessed our position and restore the allocation of *hogei* to *Ranacephala*.

We note also that Thomson et al. recognized broader concepts of the genera *Emys* and *Geochelone* than we do, and that they did not recognize the genera *Actinemys*, *Emydoidea*, and *Centrochelys*, which we recognize either fully or provisionally, but they included species from all three in their analysis.

Thomson et al. (2021) favored the use of an inclusive generic concept of *Emys* including *marmorata+pallida*, *blandingii*, and *orbicularis+trinacris*. However, they recovered an apparent non-monophyly of *Emys* with respect to *Terrapene*, with *marmorata+pallida* (= *Actinemys*) as a monophyletic sister group to a clade containing two reciprocally monophyletic subgroups: *blandingii* (= *Emydoidea*) plus *orbicularis+trinacris* (= *Emys*), and *Terrapene*. In contrast, earlier work by Spinks and Shaffer (2009) and Spinks et al. (2009b, 2016) recovered *marmorata+pallida*, *blandingii*, and *orbicularis+trinacris* as a clade sister to *Terrapene*. Based on sequence data from 30 nuclear genes, 4 mitochondrial genes, and the combined data set, Spinks et al. (2016) provided the largest molecular data set concerning these taxa, and found strong statistical support for the monophyly of the inclusive *Emys* clade, as well as the three contained lineages (*marmorata+pallida*, *blandingii*, and *orbicularis+trinacris*). Wiens et al. (2010) and Fritz et al. (2011b) found evidence for separate generic status for these three lineages and treated them as such (see also TTWG Annotations 07:21, 09:16, 10:12, and 11:7). In view of these mixed results, and the continuing use in the wider herpetological community of the separation of these lineages as genera, our checklist now treats *Actinemys* as a separate genus without equivocation as to its possible congeneric status with *Emys*. Most of us also favor the use of *Emydoidea* as separate from *Emys*, but note that some prefer to use *Emys* for the Blanding's Turtle.

21:4. **Testudines**: McCranie (2018) recommended that the group name Testudinata Behn 1760 be used for the order of all turtles, as opposed to Testudines Batsch 1788, based on Behn's use of the name Testudinata in his Tabula Generalis as an "Ordo". However, Behn's (1760) work was a direct translation of Klein (1751), a pre-1758 publication with no nomenclatural standing, and Behn's Table is simply a copy with German translation of the Table presented by Klein (1751). Additionally, the name that both Klein and Behn used for their "Ordo III" of Quadrupedia was actually Depilata, which included the "Familias" of Testudinata (for the sole genus *Testudo*), Cataphracta (for the genera *Crocodilus*, *Caimanus*, and *Lacerta*), and Nuda (for various amphibian and reptilian genera, including *Salamandra*, *Gecko*, *Cordylus*, *Scincus*, *Chamaeleo*, and *Batrachus*). We have for a long time listed Testudinata Klein *in* Behn 1760 in our checklist as a synonym of Testudines Batsch 1788, but since the Principle of Priority of the ICZN (Art. 23) does not pertain to suprafamily names (see ICZN Art. 1.2.2 and TTWG Annotation 10:4), and as no one else has ever used this name and authorship for the order of all turtles, we consider it a virtual *nomen oblitum*, and continue to use Testudines Batsch 1788 as the preferred ordinal name for all turtles and tortoises, as do most recent authors.

Furthermore, Savage (2020) argued that because the name Testudines Batsch 1788 was originally proposed as a family name, it should be excluded from applying to an order. He also argued that the name Chelonia Macartney and Ross 1802 was applied to a genus, and is similarly unavailable as an order name. His resolution was to use Testudinata Oppel 1811 for the order of turtles. However, we disagree and continue to use the older name Testudines Batsch 1788, as do most recent authors.

21:5. [*Annotation text previously included here now deleted from this checklist.*]

21:6. **Publications by Cope**: Two dates of publication for papers by Edward D. Cope have been corrected: Cope 1868b was actually published in 1869 (on Feb. 6), and Cope 1876 was actually published as a separate in 1875 (on Nov. 26); these dates were confirmed by Osborn (1929), but had been overlooked.

21:7. **Publications by Rafinesque**: Bell and Bauer (2020) evaluated a previously overlooked publication by Rafinesque (1818), in which he published three new turtle names based on animals captured along the Ohio River and in Kentucky: *Testudo bigibbosa*, *Testudo chlorops*, and *Trionyx ohiensis*. Bell and Bauer (2020) also accessed and published part of Rafinesque's archived field notes from that expedition, including his original drawings of the three new turtle species.

Testudo bigibbosa from the Ohio River, also referred to alternatively as *Testudo nodosa* in Rafinesque's notes, and then later also published by Rafinesque (1822) as *Emyda nodosa* (as a *nomen nudum*) is certainly a *Graptemys*, most likely *G. ouachitensis*, although possibly *G. pseudogeographica*, with the drawing of a male specimen with a narrow and markedly serrated shell with clear dark vertebral knobs and dark costal spots.

Testudo chlorops, also referred to alternatively as *Testudo radiata* in Rafinesque's notes, and then later published by Rafinesque (1822) as *Emyda semiradiata* (as a *nomen nudum*), and noted to come from Kentucky and incorrectly referred to as being terrestrial, is clearly a *Trachemys scripta elegans* based on the detailed drawings of the carapace and plastron with their characteristic markings.

Trionyx ohiensis was questionably allocated by Bell and Bauer (2020) to *A. spinifera* (as opposed to the smaller *A. mutica*), but the fact that Rafinesque described it as "*the* [emphasis ours] large softshell of the Ohio," and the clear depiction on his drawing of many small spiny tubercles on the anterior margin of the carapace, clearly refers it to *A. spinifera spinifera*. Thus, Rafinesque's *ohiensis* preceded the description of *spinifera* by LeSueur (1827), but it was declared a *nomen oblitum* by Bell and Bauer (2020), with which we agree. The other overlooked Rafinesque names are clearly *nomina nuda*, as also determined by Bell and Bauer (2020).

21:8. **Pleurodera and Cryptodera**: McCranie (2018), referencing Gutsche (2016), stated that the authorship of Pleurodera and Cryptodera was not solely by Lichtenstein (1856), as we and other authors had previously recorded it, but by Lichtenstein and von Martens (1856). A review of the work in question indicates that, in fact, Lichtenstein acknowledged both Weinland and von Martens for their editorial help and work, but not that he considered either a co-author. However, Lichtenstein did note that he worked closely together with von Martens to coin some of the new names used, so we have now updated the citation accordingly.

In addition, we had previously considered this work to be an unpublished catalogue (as stated by Lichtenstein himself in his introduction) and therefore "perhaps not nomenclaturally available" (Rhodin et al. 2008, TTWG Annotation 08:20). McCranie (2018) criticized our position and selectively quoted us (eliminating our modifying word "perhaps") and instead considered the work and the names available and implied their nomenclatural priority over Cryptodira Cope 1869 and Pleurodira Cope 1864. However, once again, since the Principle of Priority of the ICZN (Art. 23) does not pertain to suprafamily names (see ICZN Art. 1.2.2 and TTWG Annotation 10:4), the names Pleurodera and Cryptodera (Lichtenstein and von Martens 1856) do not have priority and we continue the use of the names authored by Cope, as do most authors.

21:9. **Pleurodira, Chelidae, and Podocnemididae**: Ferreira et al.

(2018) performed an extensive morphologic analysis of the Pleurodira, using 245 characters in 77 fossil taxa and 24 living species. Their interpretation of relationships among the Chelidae suggested that the long-necked South American and Australian taxa (*Chelodina, Hydromedusa,* and *Chelus*) were monophyletic, and placed them in the clade Chelina, whereas all the shorter-necked South American and Australian taxa (except for *Pseudemydura*) were placed in the clade Chelini. They acknowledged that their results did not corroborate most phylogenetic analyses that regard South American and Australian chelids as divergent and in separate subfamilies, but they did not recommend choosing one alternative interpretation over the other. For the Podocnemididae, they placed *Erymnochelys* and *Peltocephalus* in the subfamily Erymnochelyinae and *Podocnemis* in the Podocnemidinae.

21:10. ***Chelus fimbriata***: Ceríaco and Bauer (2017) documented and published the entire previously unpublished manuscript description by Alexandre Rodrigues Ferreira from 1784 of the Matamata Turtle as *Testudo torticollis*. Since the name was published as a junior synonym of *Chelus fimbriatus* (= *Chelus fimbriata*), it has no nomenclatural validity under ICZN Article 11.6, and is therefore considered a *nomen nudum*. The same unpublished name by Ferreira was previously inaccurately referred to by Carvalho (1972) as *Testudo corticolis*, also a *nomen nudum*.

21:11. ***Chelus fimbriata*** **and** ***Chelus orinocensis***: Vargas-Ramírez et al. (2020) examined phylogeographic differentiation of the Matamata across its range using mitochondrial DNA and nuclear intron sequences and 1905 ddRAD loci corresponding to 685,247 bp. Their analysis revealed the existence of two morphologically distinct and genetically deeply divergent evolutionary lineages that apparently separated in the late Miocene, corresponding to the approximate time when the Orinoco Basin was established. They described the Matamata from the Orinoco, Río Negro, Rio Branco, and Rio Essequibo basins as the new species *Chelus orinocensis*, restricting *C. fimbriata* to the Amazon and the eastern Guianas. Two specimens, one each from the Rio Branco of northern Brazil and the Rio Essequibo of Guyana, showed conflicting mitochondrial and nuclear results suggestive of past gene flow and old mitochondrial introgression from the Amazon lineage into the Orinoco lineage, and the authors assigned these specimens only tentatively to *C. orinocensis*, pending further study. We agree with the specific separation of *C. orinocensis* from *C. fimbriata*, noting that the Rio Branco and Essequibo specimens are morphologically most similar to Orinoco specimens, as previously pointed out by Pritchard and Trebbau (1984), Sánchez-Villagra et al. (1995), and Pritchard (2008 CBFTT). Cunha et al. (2021) documented many new localities for both species.

21:12. ***Mesoclemmys***: In a paper aimed primarily at dating the diversification of chelid turtles, Holley et al. (2020) combined molecular and morphological analyses (see TTWG Annotation 33 for *Myuchelys purvisi*) and found that the genus *Mesoclemmys*, as currently recognized, emerged as paraphyletic. However, as this was likely a by-product of their analytical process, and without deeper taxonomic and genetic sampling in their dataset, the authors themselves did not advocate changing the taxonomy of *Mesoclemmys*.

Nevertheless, based on subsequent work by Thomson et al. (2021) we acknowledge that *Mesoclemmys* is indeed paraphyletic and in need of revision, including recognition that the included species *M. hogei* is best separated and placed into the previously described monotypic genus *Ranacephala* (see TTWG Annotation 3 for Testudines).

21:13. ***Mesoclemmys dahli***: Gallego-García et al. (2018) investigated genetic structure of the critically endangered endemic *M. dahli* across its complete geographic range in Colombia, using 3211 neutral and adaptive-loci associated SNPs. They documented significant genetic structuring, with at least four subpopulations with surprisingly moderate to high levels of genetic diversity. However, extant subpopulations are severely fragmented, isolated, and very small (effective population sizes < 50), with significant inbreeding and little to no contemporary gene flow among them. They made no taxonomic recommendations (see also Vargas-Ramírez et al. 2012a and TTWG Annotation 12:40), but noted the increased conservation risk associated with these population parameters and recommended the implementation of a genetic rescue strategy in order to reduce inbreeding.

In a subsequent paper, Gallego-García et al. (2019) employed Restriction-site Associated DNA sequencing (RADseq) to examine fragmentation and gene flow among populations of *M. dahli*. They identified six genetically isolated populations divided among three higher level genetic clusters. Given the results of both papers, they hypothesized that these divergences might be the result of geographic (and hence genetic) fragmentation and strong natural selection for adaptation to anthropogenic landscape changes, particularly forest clearing. They made no taxonomic recommendations and did not suggest managing those clusters independently, but rather supported restoring gene flow via habitat corridors or translocation among ecologically similar forest and grassland habitats to increase genetic diversity.

21:14. ***Mesoclemmys gibba***: Ettmar (2019:152), citing previous observations by Bour and Pauler (1987) and Bour and Zaher (2005), noted that western Amazonian populations of this species, corresponding to the *Emys stenops* morphotype, appear to be differentiated from the eastern morphotypes, corresponding to *Emys gibba*. Whether these differences warrant recognition of *stenops* as a separate taxon will require further evaluation.

21:15. ***Mesoclemmys heliostemma, M. raniceps,*** **and** ***M. wermuthi***: Cunha et al. (2019) incubated eggs of *M. raniceps* and produced clutches of hatchlings matching the phenotypic appearance of both *M. raniceps* and *M. heliostemma* in the same clutch, demonstrating that these two "species" are color morphs of the same taxon, and synonymized *heliostemma* under *raniceps*. This was further confirmed by Vogt et al. (2020). We agree with this synonymization.

Cunha et al. further demonstrated, based on the presence of prominent dark head stripes and lack of white pretibial skin flaps, that the previously described species *Phrynops wermuthi* Mertens 1969 (= *Mesoclemmys wermuthi*) was distinct from *M. raniceps* / *M. heliostemma* and a separate species. They also noted that the holotype of *Hydraspis maculata* Gray 1873c, previously considered a synonym of *M. raniceps*, had head stripes and appeared to be the same taxon as *wermuthi*, but that it was of unknown provenance, as previously noted by Rivas et al. (2015). They suggested that the name *H. maculata* was available (as *Mesoclemmys maculata*) and that *P. wermuthi* was its synonym. However, Ettmar (2019) concurrently synonymized *H. maculata* with *M. zuliae* (Pritchard and Trebbau 1984), which also has dark head stripes, lending further uncertainty to its allocation.

Since the identity and provenance of the single type specimen of *H. maculata* are both problematic, and the species has not been recognized as valid since initially being synonymized with *Mesoclemmys nasuta* by Boulenger (1889), and *P. wermuthi* was considered to be distinct as either a valid species or subspecies by many authors in multiple publications for about 20 years after its description in 1969, in accordance with the criteria of ICZN article 23.9.1, we interpret *Hydraspis maculata* Gray 1873c to be a *nomen oblitum*. We therefore recognize *Mesoclemmys wermuthi* (Mertens 1969) as valid and distinct from *M. raniceps* / *M. heliostemma*.

Cunha et al. (2019) also noted that Molina et al. (2012), who had demonstrated morphological differences between *M. heliostemma* and *M. raniceps*, had in fact interpreted specimens with head stripes and black pretibial flaps (actually *M. wermuthi*) as instead being *M. raniceps*, thereby further confirming the difference between *M. wermuthi* and their *M. heliostemma*, which was actually the true *M. raniceps*. Range-wide genetic analysis of these two confusing Amazonian species is badly needed.

21:16. ***Phrynops geoffroanus***: The junior synonym *Emys depressa*, historically attributed to Merrem (1820), should actually be attributed to Wied-Neuwied *in* Merrem (1820), as previously noted by Bour (2008c), since Merrem specifically credited Wied-Neuwied as having sent him the description of *Emys depressa*.

21:17. **Chelodininae**: Cann and Sadlier (2017; hereafter C&S), in their beautifully illustrated and richly detailed volume on Australian and New Guinean turtles, used a taxonomy not currently accepted by most authorities or TTWG, based only on qualitative morphological distinctions. Their taxonomic arrangements included the following disagreements with our previous TTWG checklists (TTWG 2007b, 2009, 2010, 2011, 2012, 2014, 2017) and other recently published scientific literature (e.g., Iverson et al. 2001; Georges 2002; Fritz and Havas 2007; Rhodin et al. 2008; Thomson and Georges 2009; Georges and Thomson 2010; Todd et al. 2014a; Georges and Thomson 2016; Georges et al. 2018a; Rhodin et al. 2018b; and many others, but see Cogger 2018 for a supportive view).

C&S recognized two subspecies under *Chelodina (Chelodina) canni*: *C. (C.) canni canni* (western populations) and *C. (C.) canni rankini* (eastern populations). Although future research may support the recognition of these two populations as separate taxa, the name *rankini* is unavailable (Iverson et al. 2001; TTWG Annotation 07:86), and hence not recognized by us.

C&S elevated *C. (C.) mccordi timorensis* from a subspecies to a full

species, without any justification, but based on Kuchling et al. (2007; TTWG Annotations 07:89 and 10:36), we retain it as a subspecies. Subspecies status is further supported by the results of Kehlmaier et al. (2019a), that revealed shallow mitogenomic divergence between *C. m. mccordi* and *C. m. timorensis* (see Annotation 19 below).

C&S recognized four subspecies under *Chelodina (Macrochelodina) oblonga* (=*rugosa*): *C. (M.) oblonga oblonga*, *C. (M.) oblonga rugosa*, *C. (M.) oblonga burrungandjii*, and *C. (M.) oblonga walloyarrina*. Following the work of previous authors and TTWG, we recognize *C. (M.) oblonga* (=*rugosa*), *burrungandjii*, and *walloyarrina* as full species (TTWG Annotations 08:94-95, 10:38, and 14:42-43), and *rugosa* as a synonym of *oblonga* (Annotations 07:88, 08:29, 10:39, and 14:42-43).

Without any analysis to support their usage, C&S elevated *C. (M.) siebenrocki* from the synonymy of *C. (M.) oblonga* (=*rugosa*) and recognized it as a distinct species, although they acknowledged that it was indistinguishable from their *C. (M.) oblonga rugosa*. Based on previous work by Georges (2002) and Georges and Thomson (2010; see also TTWG Annotation 07:91), we retain *siebenrocki* as a synonym of *rugosa*, but see Annotations 22 and 24 below regarding genotyping of these and other taxa of *Chelodina* by Kehlmaier et al. (2019a).

C&S recognized and used the name *Elseya jukesi*, although they acknowledged that the validity of the name is in question. The name *jukesi* is unavailable (Iverson et al. 2001; see also TTWG Annotation 07:94) and we use the available name *flaviventralis* (Georges and Thomson 2016) for this taxon.

C&S recognized *Elseya stirlingi* (from the Johnstone River in Queensland) as distinct from the nearby disjunct population of *Elseya irwini* (restricted by them to the Burdekin River), although they acknowledged that the validity of the name *stirlingi* is in question. Because Thomson and Georges (2009) rendered *stirlingi* an unavailable name, we continue to include both populations under *Elseya irwini* pending future analyses evaluating the potential divergence of these disjunct river populations.

C&S used the generic name *Wollumbinia* Wells and Wellington for the Saw-Shelled Turtles, instead of *Myuchelys* Thomson and Georges, although they acknowledged that its validity was in question. Georges and Thomson (2010) indicated that *Wollumbinia* was an unavailable name (see also Iverson et al. 2001, Fritz and Havas 2007, and TTWG Annotations 07:92, 09:45, and 10:43 for discussions of taxonomic names proposed by Wells and Wellington), and we recognize this genus with the available name *Myuchelys*.

C&S recognized several subspecies of *Emydura macquarii* that were synonymized under *E. m. macquarii* by Georges and Thomson (2010; see TTWG Annotations 07:98 and 10:42): *E. m. signata*, *E. m. binjing*, *E. m. dharra*, *E. m. gunabarra*, and *E. m. dharuk*. We follow Georges and Thomson (2010) in synonymizing these taxa under *E. m. macquarii*. In addition, C&S transferred the currently recognized subspecific taxa *E. m. emmotti*, *E. m. krefftii*, and *E. m. nigra* (see Todd et al. 2014a) from *E. macquarii* to *E. krefftii*, justifying the establishment of *E. krefftii* as a distinct species based solely on head color and its exceptionally deep shell. There is no support for the species *E. krefftii* as defined by C&S in the recent range-wide phylogeographic analysis of the genus *Emydura* elucidated by Georges et al. (2018a). We follow Todd et al. (2014a) in retaining these three taxa as subspecies of *E. macquarii*. However, using dense genomic sampling of single-nucleotide polymorphisms (SNPs), Georges et al. (2018a) identified at least five distinct evolutionary units within this complex, but declined from making any taxonomic recommendations, pending further studies, including morphology: 1) the Murray-Darling Basin (to which the name *macquarii* could be restricted), 2) the Lake Eyre Basin (to which the name *emmotti* applies), 3) the Hunter River Basin (to which the name *gunabarra* applies), 4) and a single coastal form (to which the name *krefftii* applies) that might be divisible into four geographic variants that correspond to the three available names *krefftii*, *signata*, *nigra*, and one unnamed variant.

C&S recognized both *Emydura australis* and *E. victoriae* as distinct species; however, *E. australis* was synonymized with *E. m. macquarii* by Cogger et al. (1983) and Georges and Thomson (2010) (see TTWG Annotation 10:41). C&S argued that *E. australis* is most similar to *E. victoriae*, and that they might even be synonymous, in which case *E. australis* Gray 1841 would have nomenclatural priority over *E. victoriae* Gray 1842. Pending further study, we follow Georges and Thomson (2010) and retain *australis* as a *nomen dubium* under the synonymy of *E. m. macquarii*.

C&S elevated *Emydura worrelli* to a distinct species separate from its previous recognition as a subspecies of *E. subglobosa*, based on the degree of reddish color differences between these two taxa. However, Georges and Adams (1996) previously demonstrated a lack of genetic differentiation between them and we continue to retain *worrelli* as a subspecies of *subglobosa* (e.g., Georges and Thomson 2010; TTWG Annotation 07:99).

C&S also recognized and described, but did not name, an alleged new species of *Elseya* from the Daintree River in Queensland, and an alleged new subspecies of *Emydura macquarii* from the Tweed and Richmond Rivers in New South Wales.

Reviews of this book by Georges (2018) and Spencer (2019) provided additional critique of the taxonomy used by C&S, and Sadlier and Cann (2018) provided a defense of their positions.

21:18. ***Chelodina gunaleni***: Kehlmaier et al. (2019a) demonstrated that the mitogenome of the holotype of *C. gunaleni* differed from that of its closest sister lineage, *C. novaeguineae/reimanni*, supporting its distinctness, but at a low divergence level consistent more with subspecific rather than specific separation. However, they made no recommendation to change the current taxonomy.

21:19. ***Chelodina mccordi***: Kehlmaier et al. (2019a) demonstrated that the mitogenome of the holotype of *C. mccordi roteensis* (McCord et al. 2007b) did not differ significantly from the paratype of *C. mccordi* (Rhodin 1994a), and therefore synonymized *roteensis* under *mccordi*, which we accept. They also found that the holotypes of *C. timorlestensis* (Kuchling et al. 2007) and *C. timorensis* (McCord et al. 2007a) were identical to each other and sister to *C. mccordi*, and concluded that the differences between *mccordi* and *timorensis* were consistent with subspecific separation.

21:20. ***Chelodina reimanni* and *C. novaeguineae***: Kehlmaier et al. (2019a) demonstrated that the mitogenome of the holotype and paratype of *C. reimanni* (Philippen and Grossmann 1990) showed negligible divergence from that of the lectotype of *C. novaeguineae* (Boulenger 1888), and at a low level consistent more with subspecific rather than specific separation, but retained both as distinct species and made no taxonomic changes.

21:21. ***Chelodina (Chelydera)* and *Chelodina (Macrochelodina)***: Shea et al. (2020) noted that the previous designation by McCord and Joseph-Ouni (2007a) of a neotype for the Southwestern Snake-necked Turtle, *Chelodina oblonga* Gray 1841, and their creation of the new subgenus *Macrodiremys* for *C. colliei* Gray 1856a (= *C. oblonga*) was invalid. Instead, Shea et al. (2020) restored the subgeneric name *Macrochelodina* Wells and Wellington 1985 for *C. oblonga* (as it had been originally designated), and synonymized *C. colliei* under *C. oblonga*. Further, since *Macrodiremys* was designated a junior synonym of *Macrochelodina*, this left the previous concept of *Macrochelodina* without a valid name, therefore Thomson and Georges (in Shea et al. 2020) proposed *Chelydera* as the necessary new subgeneric name for the Northern Snake-necked Turtle, *C. rugosa* Ogilby 1890, and the other broad-shelled long-necked species (such as *C. expansa*), with *C. parkeri* Rhodin and Mittermeier 1976 designated as its type species.

21:22. ***Chelodina rugosa* (previously *C. oblonga*) and *Chelodina oblonga* (previously *C. colliei*)**: Kehlmaier et al. (2019a) genotyped the holotype of *Chelodina oblonga*, which had previously been interpreted by Thomson (2000a), based on morphology, as representing a Northern Snake-necked Turtle, *C. rugosa* (see also Kennett et al. 2014 CBFTT and TTWG Annotation 14:43). Instead, Kehlmaier et al. (2019a) demonstrated genetically that the specimen was referable to the Southwestern Snake-necked Turtle, as originally allocated, and therefore returned the name *C. oblonga* to that species, but considered it a *nomen dubium* (with which we do not agree), suggesting that *C. colliei* Gray 1856a should be used as the name for that species. They also recommended that *C. rugosa* be resurrected as the correct name for the Northern Snake-necked Turtle. Subsequently, as noted in the previous annotation, Shea et al. (2020) invalidated the subgeneric name *Macrodiremys* for *C. oblonga*, returning it to its originally intended subgenus *Macrochelodina*, synonymized *C. colliei* under *C. oblonga*, and Thomson and Georges (in Shea et al.) created the new subgenus *Chelydera* for a resurrected *C. rugosa* and the other broad-shelled long-necked species.

21:23. ***Chelodina kuchlingi***: Kehlmaier et al. (2019a) demonstrated that the mitogenome of the holotype of *C. kuchlingi* differs significantly from Northern Territory *C. rugosa* [= *C. kurrichalpongo*] and other *Chelodina* species, including the holotype of *C. siebenrocki* [= *C. rugosa*] from Papua New Guinea, validating its status as a separate lineage.

Kuchling (2020) subsequently documented that the originally-designated type locality of "Kalumburu" for the holotype of this species was incorrectly recorded when it was accessioned into the Western Australian Museum, and that instead, it was likely collected in Parry Creek in the Ord River floodplain. As such, he restricted the known range of the species to the limited Ord River basin and the type locality to "Parry Creek." He also noted that the species had not been collected since 1974 and that *C. rugosa* [= *C. kurrichalpongo*] had recently invaded the Ord River floodplain as a result of anthropogenic habitat disturbance, possibly leading to hybridization or introgression with *C. kuchlingi* and its possible gradual replacement, although focused surveys would be needed to ascertain its conservation status and possible persistence.

21:24. ***Chelodina kurrichalpongo***: Joseph-Ouni et al. (2019a) described this species from the lowlands of the Northern Territory as *Macrochelodina kurrichalpongo*, recognizing the western populations of the Northern Snake-necked Turtle (*C. rugosa*, previously *C. oblonga*) as a separate species from the eastern populations, based on morphological differences of shell shape, coloration of the body and iris, and referring to previously described mtDNA differences between these populations elucidated by Alacs (2008) in an unpublished thesis and Kehlmaier et al. (2019a). They also recognized *C. siebenrocki* as a valid subspecies of *C. rugosa*. Following their previous works, they recognized *Macrochelodina* as a genus as opposed to a subgenus of *Chelodina*. They recognized the eastern Gulf of Carpentaria and Cape York populations as *M. rugosa*, reverting to the name previously used before the origin and identity of the holotype of *M. oblonga* was questioned by Thomson (2000a), but clarified by Kehlmaier et al. (2019a) and since by Shea et al. (2020).

Georges et al. (2002) examined variation in 24 allozyme loci for 7 populations (60 individuals) of *Chelodina rugosa* and related taxa across their range. Despite documenting some fixed allelic differences between populations of *C. rugosa* from the Northern Territory vs. *C. rugosa* from Queensland and *C. siebenrocki* from New Guinea, they concluded that these populations comprised a single species based on their exceptionally low level of genetic differentiation, less than would be expected by chance given the sample sizes and less than would be expected from isolation by distance within a single species. However, Kehlmaier et al. (2019a) demonstrated deep genetic divergence between *C. rugosa* populations from the Darwin area of Northern Territory vs. those from New Guinea (the holotype of *C. siebenrocki*) and *C. rugosa* of unknown origin and suggested that the group comprised more than one species. They also suggested that the name *C. intergularis* might be applicable to one of these species.

Pending further genetic analysis of these populations and the holotypes of *C. rugosa*, *C. intergularis*, and *C. kurrichalpongo*, and further assessment of the diagnostic characters to demonstrate their consistency or variation among individuals from across the range of *C. kurrichalpongo* and *C. rugosa*, we tentatively recognize *C. kurrichalpongo* from the Northern Territory as a valid taxon, although some of us consider it a synonym of *C. rugosa*. We also retain *C. siebenrocki* in the synonymy of *C. rugosa* until additional data are available to resolve the incompatibility of the nuclear and mtDNA evidence.

21:25. ***Chelodina rugosa* and *C. siebenrocki***: In addition to their description of *Macrochelodina* [*Chelodina*] *kurrichalpongo*, Joseph-Ouni et al. (2019a) recognized two subspecies of *M.* [*C.*] *rugosa*: the nominate form, *M.* [*C.*] *r. rugosa* from Australia, and *M.* [*C.*] *r. siebenrocki* from New Guinea, based on their morphological assessment, although they provided no quantitative data to support their assertion. They also synonymized *Chelodina intergularis* Fry 1915 with *M.* [*C.*] *r. rugosa* from Cape York, although it is possible that the holotype, lacking a specific type locality, may conceivably have come from the Northern Territory, and as such might have nomenclatural priority over *M. kurrichalpongo*.

Kehlmaier et al. (2019a) demonstrated that the mitogenome of the holotype of *C. siebenrocki* differed only slightly from a *C. rugosa* of unknown origin (but possibly Queensland), agreeing with the prior characterization of the minor differences between these taxa as noted by Rhodin and Mittermeier (1976) and Georges et al. (2002), and chose to retain *siebenrocki* in the synonymy of *rugosa*. They also found that this *C. rugosa*/*siebenrocki* lineage was markedly different from *C. rugosa* from the Northern Territory, a taxon subsequently described by Joseph-Ouni et al. (2019a) as *M. kurrichalpongo*.

21:26. **Emydurinae**: Joyce et al. (2021) provided a detailed phylogenetic nomenclature (PhyloCode) for all fossil and living turtles, in which they largely followed our TTWG (2017) nomenclature for the higher categories of all living turtles. However, they established and defined the new suprageneric clade Emydurinae for the short-necked Chelidae from the Australia-New Guinea region as sister to the Pseudemydurinae. Gaffney (1977) had previously erected and named this clade as the infrafamily Emydurodd. Based on subsequent work on the phylogenetic relationships of the Chelidae by Seddon et al. (1997), Georges et al. (1998), Guillon et al. (2012), Pereira et al. (2017), and Zhang et al. (2017), we had previously recognized the subfamilies Chelodininae (see TTWG Annotation 12:39) and Pseudemydurinae (see TTWG Annotation 17:100) for the Australasian Chelidae. Thomson et al. (2021) recently corroborated this clade of *Emydura*, *Elseya*, *Elusor*, *Myuchelys*, and *Rheodytes* as a monophyletic group sister to the Chelodininae. We therefore now also recognize the subfamily Emydurinae.

21:27. ***Elseya novaeguineae* (*Elseya caelatus*)**: Based on phenotypic differences of shell shape, color, size, and carapacial serrations, Joseph-Ouni and McCord (2019a) described a new species with two subspecies (*E. caelatus caelatus* and *E. c. ayamaru*) from the western Vogelkop peninsula and the Raja Ampat Islands of West Papua, Indonesia, separating these taxa from *E. novaeguineae* from the eastern Vogelkop and southern Bomberai peninsulas. They did not consider or discuss previously published molecular evidence of mitochondrial haplotype diversity in these populations as delineated by Georges et al. (2014). Both the molecular and morphological data suggest that there is substantial variation within *E. novaeguineae* distributed across the Vogelkop and Bomberai peninsulas (Georges et al. 2014; Thomson et al. 2015). However, neither data set is sufficiently complete or internally consistent enough to resolve the question of whether this is lineage diversity within a single species or an indication of a species complex with multiple taxa. The molecular data (Georges et al. 2014) indicate two substantive mitochondrial clades within Vogelkop and Bomberai *E. novaeguineae*, but the distribution of haplotypes conflicts with a clear split between the populations assigned by Joseph-Ouni and McCord (2019a) to their western species, *E. caelatus*, vs. those they assigned to eastern and southern *E. novaeguineae*. The holotype and western type locality site they selected for *E. caelatus*, Salawati Island, actually has an eastern *E. novaeguineae* haplotype (Georges et al. 2014). Given our preference for only changing taxonomy when the data are compelling and corroborative, most of us recognize neither *Elseya caelatus* nor its subspecies *E. c. ayamaru* at this time and place both names in the synonymy of *Elseya (Hanwarachelys) novaeguineae*.

We note additionally that the genus name *Elseya* is feminine, and therefore the specific epithet *caelatus* (= past participle, from the verb *caelare*: I carve, to carve) needs to be rendered in the Latin feminine, *caelata*, rather than in the masculine *caelatus*, as originally written. Gray (1867) proposed the generic name *Elseya* to honor Dr. Elsey, the collector, by adding an "*a*", which most often indicates feminine gender, though not always. He included two species in his new genus: *dentata* (a feminine adjective) and *latisternum* (*sternum* = a neutral noun), thereby choosing the feminine gender for *Elseya*. As such, the epithet needs to be *caelata* instead of *caelatus*. The epithet *ayamaru* was established as a noun in apposition and therefore does not need to agree with the gender of the genus.

21:28. ***Elseya schultzei* (*Elseya orestiad*)**: Based on phenotypic differences of shell shape, color, small size, and other morphological features, Joseph-Ouni and McCord (2019b) described *Elseya orestiad* as a new species based on eight specimens from a very limited distribution around Lake Sentani in the Cyclops Mountains of northeastern Papua, Indonesia, differentiating it from the larger, widespread, and sympatric *E. schultzei* which also occurs in the Tami River catchment. However, no quantitative analyses of morphological differences were performed, nor any indication of the extent of variation in *E. schultzei*, nor was any genetic analysis done. Although it is possible that *E. orestiad* may represent a distinct taxon, we refrain from recognizing it at this time, and most of us regard it as a junior synonym of *E. schultzei*, pending genetic analysis.

21:29. ***Elseya albagula* and *E. irwini***: Todd et al. (2014b) analyzed the phylogenetic relationships of nearly all major populations and taxa of Australian and New Guinean *Elseya* based on molecular data comprising both mitochondrial and nuclear loci. They identified the Daintree River population in northern Queensland as a putative undescribed species, *Elseya* sp. [Daintree], and sister to *E. irwini*. They also documented shallow genetic

differentiation of *E. albagula* from the Fitzroy River as compared to the rest of its range, but did not suggest the need for nomenclatural distinction. [*Annotation text previously included here now deleted from this checklist.*]

21:30. ***Elseya lavarackorum* (*Elseya oneiros*)**: Joseph-Ouni et al. (2020) described apparent shell scutellation differences between two Late Pleistocene fossils (one partial carapace and one plastron) of *E. lavarackorum* from the Riversleigh site and several living specimens of *E. lavarackorum* from the Nicholson and Gregory River drainages, and concluded that the living specimens were not the same species as the fossils, in contradiction to the earlier conclusion by Thomson et al. (1997) that the fossil and living specimens belonged to the same species. Joseph-Ouni et al. (2020) re-named the living form as a new species, *Elseya (Pelocomastes) oneiros*, but also assigned a third Late Pleistocene fossil (plastron only) from Riversleigh to their new species.

Thomson et al. (1997) had based their conclusions of conspecificity on several skeletal characters and a single scutellation character that demonstrated a close relationship between the fossil and living specimens, yet Joseph-Ouni et al. (2020) did not address or describe any of the skeletal characters, focusing instead on a variety of other apparently variable scutellation differences. However, basing these variable differences on only single carapacial and plastral fossil specimens without regard to the extent of variation in living specimens does not, in our opinion, constitute adequate demonstration of significant enough specific differences to warrant recognition of separate taxa, and as such, most of us consider *E. oneiros* to be a junior synonym of *E. lavarackorum*. In addition, Joseph-Ouni et al. (2021) erroneously placed *E. lavarackorum* in the subgenus *Elseya*, despite the clear allocation of that species to the subgenus *Pelocomastes*, as demonstrated by its skeletal morphology (Thomson et al. 1997, 2015).

The known distribution of *E. lavarackorum* has recently been extended markedly westward through the work of Georges (2021), who genotyped specimens of what appeared phenotypically to be *E. dentata* from the Roper River in the Northern Territory, but identified them as E. *lavarackorum*.

21:31. ***Emydura gunaleni***: Smales et al. (2019a) described this isolated new species of *Emydura* from the western Vogelkop peninsula in West Papua, Indonesia, differentiating it morphologically from the widespread *E. subglobosa* occurring more than 500 km to the southeast and other more distant members of the genus. *Emydura gunaleni* is characterized by unambiguous discrete diagnostic characters (e.g., intergular completely separates the pectorals, distinctive pencil thin light stripe behind the eye) in addition to qualitative diagnostic characters of color and shell shape, and we therefore recognize this taxon as distinct.

21:32. ***Emydura subglobosa angkibaanya***: Joseph-Ouni et al. (2019b) described this taxon from a very limited distribution in the Jardine River at the northern tip of Cape York, Australia, distinguished from *E. s. subglobosa* from New Guinea by some morphological and color differences. However, only limited quantitative analyses of differences between these taxa were performed, and no analysis was done of the extent of variation in rangewide populations of *E. subglobosa*, including *E. s. worrelli* from Australia. Furthermore, genetic analysis was not performed, and the Jardine River population has only been separated from mainland New Guinea populations for about 12,000 years, a time frame relatively unlikely to have led to subspecific separation. Although it is possible that *E. s. angkibaanya* may represent a diagnosable population, perhaps at an ESU level, we refrain from recognizing it as a named subspecies at this time, and most of us regard it as a junior synonym of *E. s. subglobosa*, pending genetic analysis.

21:33. ***Myuchelys purvisi***: Kehlmaier et al. (2019a) demonstrated that the mitochondrial genome of *M. purvisi* differed markedly from other *Myuchelys* species, and clustered instead with those of *Elusor* and *Rheodytes*, which partially corroborated the previous findings of Le at al. (2013), who had noted the paraphyly of *Myuchelys* and erected the genus *Flaviemys* for *purvisi* to resolve this situation. However, Kehlmaier et al. agreed with Spinks et al. (2015) that the nuclear genomic analysis supported *purvisi* as a member of *Myuchelys*, corroborated earlier by nuclear allozyme data (Georges and Adams 1992), and suggested that the discordance between the nuclear and mitochondrial data was evidence for ancient mitochondrial introgression.

In a paper aimed primarily at dating the diversification of chelid turtles, Holley et al. (2020) generated phylogenetic trees based on: 1) mitochondrial and nuclear DNA data downloaded from GenBank, 2) previously published (but slightly modified) morphological data, and 3) the combined morphological and molecular data (largely driven by the more inclusive molecular data set). The morphological analysis was hampered by incomplete sampling of extant taxa (and did not include *M. purvisi*). However, the molecular and combined analyses produced trees with potential taxonomic implications. The latter trees both supported the recognition of *Flaviemys purvisi* as sister to *Elusor* rather than *Myuchelys*, contrary to Spinks et al. (2015) and Kehlmaier et al. (2019a), most likely caused by the mitochondrial signal in the data set. In view of the vagaries of mitochondrial genomic relationships and the more robust nuclear genomic evidence, we follow Georges and Adams (1992), Spinks et al. (2015), and Kehlmaier et al. (2019a) in continuing to recognize *purvisi* as a member of *Myuchelys* rather than a monotypic *Flaviemys*.

21:34. ***Pelomedusa galeata***: Vamberger et al. (2018) analyzed phylogeographic patterns of mitochondrial and nuclear DNA in *Pelomedusa galeata* and *P. subrufa* (sensu stricto) and inferred shifts of climatically suitable refugia since the Last Glacial Maximum using a modeling approach. Their data demonstrated distinct intraspecific genetic structuring in South African *P. galeata*, indicative of speciation between the southwestern and southeastern populations (for which existing names are available), but no evidence of genetic structuring in the more widespread *P. subrufa*. They made no specific taxonomic or nomenclatural recommendations.

21:35. ***Pelusios castaneus seychellensis***: After further review of papers by Bour (1983, 2013a), Stuckas et al. (2013), Kindler et al. (2016), and Bour et al. (2016 CBFTT) (see also TTWG Annotations 14:47 and 17:103), we acknowledge that: 1) differences in shell coloration (black plastron) and head morphology apparently differentiate *seychellensis* from typical western African *P. castaneus*, 2) but genetic findings support its close sister relationship to *P. castaneus* from coastal Republic of Congo, and 3) there is a distinct possibility that the only three known museum specimens of *seychellensis* may well have been mislabeled when collected or accessioned, or even brought to the Seychelles as part of historic trade routes, similar to specimens of *P. castaneus* recorded from Guadeloupe, Cape Verde, and São Tomé.

Combined, these considerations now lead us to conclude that it is unlikely that *seychellensis* was introduced prehistorically to the Seychelles and then became extinct. It is more likely that the population of *P. castaneus* from which the three type specimens of *seychellensis* originated has not yet been located, and it is highly unlikely to have been the Seychelles. No specimens of *P. castaneus* from the western Central African coast either south or north of the Republic of Congo were genetically sampled by Kindler et al. (2016), so it is entirely possible that the *seychellensis* genotype may have originated from an unsampled area along that coast. In addition, the geographic distribution of *P. castaneus* is discontinuous, with three apparently allopatric populations (Bour et al. 2016 CBFTT): 1) coastal West Africa, 2) inland Cameroon and Central African Republic, and 3) coastal western Central Africa. Genomic analysis by Kindler et al. (2016) also suggests that these three regions may represent separate lineages. As a result, we now accept the earlier recommendations by Stuckas et al. (2013) and Kindler et al. (2016) and synonymize *P. c. seychellensis* under *P. castaneus*. In doing so, we also remove it from the list of turtle taxa that have gone extinct since 1500 CE.

If the actual population from which the three specimens of *seychellensis* originated is positively identified genetically, and if the morphological features present in them are also present in that population, then the epithet *seychellensis* would still be available for it, if taxonomically warranted. Specimens of *P. castaneus* from forested regions of Nigeria often have dark plastra similar to *seychellensis* (Bour et al. 2016 CBFTT:f.4), and other populations with dark plastra may occur elsewhere.

21:36. ***Pelusios sinuatus***: Vamberger et al. (2019a) performed a phylogeographic analysis of this widespread species using mtDNA and nuclear DNA and noted deep division between northern clades and southern clades with a boundary along roughly the northern border of South Africa. They acknowledged the southern clade as the nominotypical subspecies *Pelusios sinuatus sinuatus* and recognized the northern taxon as a distinct subspecies, *Pelusios sinuatus bottegi* (Boulenger 1895a), which we accept.

21:37. ***Pelusios williamsi***: In a previous paper not included for annotation in our previous checklists, Fritz et al. (2012b) combined analyses of mitochondrial and nuclear DNA of all species of *Pelusios* and demonstrated that *P. williamsi* is sister to *P. castanoides*. Further, they noted that some

DNA sequences stored in GenBank, specifically earlier submitted samples of "*P. williamsi*," were misidentified as to what species they represented. They did not suggest any novel taxonomic rearrangements, but offered an important critique of how identification procedures at GenBank could be improved, which in view of how critical genetic analyses have become in taxonomy, are hopefully being addressed and resolved.

21:38. [*Annotation text previously included here now deleted from this checklist.*]

21:39. ***Podocnemis erythrocephala***: Based on mitochondrial and microsatellite markers, Michels and Vargas-Ramírez (2018) demonstrated significant differences between Amazon and Orinoco river basin populations of the species, that may have diverged during the Last Glacial Maximum, ca. 25,000 to 38,000 yrs ago. They recommended no taxonomic changes, but considered these divergent populations as distinct Management Units.

21:40. ***Podocnemis sextuberculata***: Viana et al. (2017) examined variation in the mitochondrial control region, which revealed that populations of this species in the Xingu River were divergent from those across the rest of the range in the main Amazon basin. The authors recommended them as two separate Management Units, with no taxonomic implications.

21:41. ***Podocnemis unifilis***: Confusion over the identity of the type specimen of *Emys cayennensis* (see TTWG Annotation 14:50) led Vogt et al. (2013) to petition the ICZN to conserve the name *Podocnemis unifilis* Troschel 1848 for the Yellow-spotted Amazon Turtle, giving it precedence over *Emys cayennensis* (an older name previously used erroneously as a senior synonym of *Podocnemis erythrocephala*). This petition was accepted and the ICZN (2019) has used its plenary power to give the name *unifilis* Troschel 1848, as published in the binomen *Podocnemis unifilis*, precedence over the name *cayennensis* Schweigger 1812, as published in the binomen *Emys cayennensis*, whenever the two are considered to be synonyms. This means that *Podocnemis unifilis* is a protected name and that the name *E. cayennensis* is still an available name and will appear in the synonymy of *P. unifilis* whenever they are considered to be the same species.

21:42. ***Caretta caretta***: The question of the availability of the name *Testudo caretta* Linnaeus 1758 for the Loggerhead Turtle, as originally discussed by Wallin (1985), was reopened by McCranie (2018). Wallin had suggested that since Linnaeus did not personally examine any specimens of what he called *T. caretta*, basing his description of that species only on literature citations, and since his concept of *T. mydas* (also described in 1758), was based on a composite of specimens of *T. mydas* and actual *T. caretta*, that *Testudo caretta* Linnaeus 1758 was therefore an invalid name. Instead, Wallin (1985) had suggested that the name *T. caretta* was available as *Testudo caretta* Walbaum 1782. This recommendation was again noted by McCranie (2018), who also remarked that Wallin's suggestion had not been followed in the recent literature.

However, neither author addressed the fact that Walbaum (1782) is a non-binomial work, as originally noted by Wermuth (1956), and that all new turtle names in that work (*T. verrucosa*, *T. macropus*, *T. tabulata*, *T. tabulata campanulata*, and *T. signata*) are considered unavailable *nomen illegitimum* names (e.g., Wermuth and Mertens 1961, 1977; Fritz and Havaš 2007; Rhodin et al. 2008; TTWG 2009, 2017, etc.). A review of Walbaum (1782) confirms that he did not employ binomial Linnaean nomenclature for any of his species descriptions, e.g., referring to the Loggerhead Turtle by the full original Linnaean descriptor: "*Testudo (Caretta) pedibus pinniformibus, unguibus, palmarum plantarumque binis, testa ovata, acute serrata.* L." Furthermore, we note that there is no requirement by ICZN that for a name to be available it needs to be based on actual examined specimens (see also Annotation 65). We therefore reject the suggestion that "*Testudo caretta* Walbaum 1782" replace *Testudo caretta* Linnaeus 1758, agreeing with all other recent authors.

21:43. ***Chelonia mydas***: Shamblin et al. (2018) examined variation in the mitochondrial control region for nesting *C. mydas* along the east coast of the United States, and compared their results with previously published data from the southeastern USA and the northern Caribbean. They found that haplotype frequencies from populations in South Carolina and North Carolina were significantly different from those from the other populations, and considered them as a distinct subpopulation of *C. mydas*. They made no taxonomic recommendations but suggested Management Unit status for these northern populations.

Jensen et al. (2019) examined variation in the mitochondrial control region across 127 rookeries of *C. mydas* distributed globally. They found strong phylogenetic structure and recognized 11 lineages, along with the geographically isolated Mediterranean population, as 12 Management Units, but made no taxonomic recommendations. They concluded that the greatest lineage diversity was in the western Pacific and hence was likely the center of origin for the species, and that the species likely colonized the Atlantic basin in a single event.

21:44. **Chelydroidea**: As per Crawford et al. (2015) and Thomson et al. (2021), we now also recognize the expanded superfamily Chelydroidea that includes the three families Chelydridae, Dermatemydidae, and Kinosternidae, removing the previous concept of a more restricted Kinosternoidea (which excluded Chelydridae), as per Joyce et al. (2004) and others.

21:45. ***Kinosternon***: In our previous two checklists we listed the alternative header designations of "*Kinosternon* or *Cryptochelys*" for those species of *Kinosternon* that were included in *Cryptochelys* by Iverson et al. (2013). We deferred a decision at that time as to which generic name best reflected the underlying phylogeny, as outlined in our TTWG Annotations 14:9 and 17:11, based on work by Spinks et al. (2014b). However, since then, community usage has not generally adopted *Cryptochelys*, and we have therefore now synonymized it with *Kinosternon*.

21:46. ***Kinosternon chimalhuaca***: In previous TTWG Annotations (07:7 and 14:11) we discussed the inadvertent early publication of this name in a book by Rogner (1996), and our first request to the ICZN to validate that error (Rogner et al. 2013), but since then the ICZN has recommended that we instead request suppression of the earlier name, and we have therefore done so (Rogner et al. 2016). A final determination by the ICZN is still pending, but we now revert to using the Berry et al. (1997) date of publication for this species.

21:47. ***Kinosternon cora***: Loc-Barragán et al. (2020) described this range-restricted mud turtle from the coastal plain of extreme southern Sinaloa and Nayarit, Mexico, based on morphologic and phylogenetic analysis (two mitochondrial and four nuclear loci) comparing and differentiating it from *K. sonoriense* and *hirtipes* and its sister species, *K. vogti*, as well as all other Kinosternidae. The species is distinctive and we accept it as valid.

21:48. ***Kinosternon vogti***: López-Luna et al. (2018) described this range-restricted mud turtle from the coastal Bahía de Banderas region of Jalisco, Mexico, based on a morphologic analysis comparing and differentiating it from *K. integrum* and *K. chimalhuaca*. Males have an unusual distinctive and bright yellow snout not known in other *Kinosternon* species. Genetic analysis was not done in the original description, but Loc-Barragán et al. (2020) performed a phylogenetic analysis of their new species, *K. cora*, comparing its mitochondrial DNA to *K. vogti* and most other Kinosternidae, and demonstrated that *K. vogti* and *K. cora* are sister species most closely related to *K. sonoriense*, *hirtipes*, and *chimalhuaca*, and significantly differentiated from other *Kinosternon*. Subsequently, López-Luna et al. (2021) analyzed a broader sampling of mitochondrial DNA of *K. vogti* and demonstrated its close sister relationship to *K. hirtipes* and *K. chimalhuaca*. The species is distinctive and we accept it as valid.

21:49. ***Sternotherus intermedius, Sternotherus minor,* and *Sternotherus peltifer***: Scott et al. (2018) analyzed range-wide phylogeography of *Sternotherus minor* and *S. depressus* based on 4430 RAD loci (48,207 polymorphic SNPs) scored for 193 ingroup and 12 outgroup specimens. Based on phylogenetic analysis, Bayesian clustering, and the near-absence of hybrid or significantly admixed genotypes, they concluded that what had previously been interpreted as a zone of intergradation between the subspecies *S. m. minor* and *S. m. peltifer* actually represents a new species, *S. intermedius*, and that *S. peltifer* and *S. minor* are shallowly diverged, but specifically distinct. We are uncertain as to whether *S. intermedius* is valid or not, and whether *S. peltifer* is distinct enough to warrant species rank, but accept both for now, pending further analysis.

21:50. ***Sternotherus odoratus***: Legler and Vogt (2013) discussed two museum specimens of this species collected in El Sauz, Chihuahua, Mexico, in 1903, about 300 km southwest of the nearest confirmed record in Texas, USA. They maintained that the provenance of the specimens was unequivocal (as opposed to being mislabeled), and that further collecting efforts in the same area since that time have failed to find any more; they concluded that the species once occurred in Mexico, but has since been extirpated.

21:51. **Staurotypinae**: In our previous two checklists we listed the alternative designation of "Staurotypinae or Staurotypidae" for the

monophyletic grouping of *Claudius* and *Staurotypus*. There is broad agreement concerning the reciprocal monophyly of these two genera and the remainder of the Kinosternidae (*Kinosternon* and *Sternotherus*), but we deferred a decision as to which suprageneric category (subfamily or family) best reflected the level of differentiation between the two lineages, as outlined in our previous TTWG Annotation (14:13), based on recommendations for the use of Staurotypidae by Bickham and Carr (1983) and Iverson et al. (2013). Additional data and discussions since then (Joyce et al. 2013; Spinks et al. 2014b; Joyce and Bourque 2016) have supported usage favoring Staurotypinae, and since we prefer to maintain stability in suprageneric categories (see TTWG Guidelines for Taxonomic Changes in the introduction), we have decided to continue use of the subfamily designation Staurotypinae.

21:52. **Fossil North American Testudinoidea**: Vlachos (2018) reviewed and analyzed published morphological features of all described North American fossil Testudinoidea (Testudinidae, Emydidae, and Geoemydidae [Rhinoclemmydinae]) and made many taxonomic recommendations. Most of the Pleistocene fossil taxa that were included as synonyms of Recent taxa in our previous TTWG checklists, or as extinct species in our TEWG (2015) checklist, have as a result been reassessed and taxonomically reassigned. Vlachos unfortunately labeled his synonymized or undiagnosable species as *nomen invalidum* taxa; however, the names are not nomenclaturally invalid or unavailable, but simply taxonomically synonymized.

Chrysemys: Based on what Vlachos considered to be diagnostic features of vertebral-marginal scute contacts in *Chrysemys timida* † Hay 1908b from Nebraska, he recognized it as separate and distinct from Recent *Chrysemys picta bellii*, disagreeing with the previous synonymization by Preston (1979). We acknowledge this new assessment and have therefore removed *C. timida* from the synonymy of *C. p. bellii* and from this checklist and will incorporate it in our next updated edition of TEWG (2015).

Pseudemys: Based on review of the type specimen of *Deirochelys floridana* † Hay 1908b (listed as a synonym of Recent *Pseudemys nelsoni* in TTWG 2017), with comparison to extant *Pseudemys* and *Deirochelys*, Vlachos concluded that *D. floridana* was a *nomen dubium* undiagnosable beyond *Deirochelys* sp. indet., disagreeing with Jackson (1964, 1974), who had identified it as a *Pseudemys* with closest affinities to Recent *P. nelsoni*. In view of the taxonomic disagreement and lack of a clear diagnosis, we have removed it from the synonymy of *P. nelsoni*. Vlachos also evaluated the type of *Trachemys jarmani* † Hay 1908b and felt that it was undiagnosable; therefore, since Jackson (1974) was also uncertain about its identity and had referred it to *Chrysemys* (= *Pseudemys*) sp. indet., we have also removed this taxon from the synonymy of *P. nelsoni*. Additionally, the single nuchal of *Pseudemys extincta* † Hay 1908b was determined by Vlachos to be undiagnosable, disagreeing with Preston (1979), and he removed *P. extincta* from the synonymy of Recent *Pseudemys rubriventris*, with which we also agree.

Trachemys: Vlachos examined the type series and other collected material of *Trachemys sculpta* † Hay 1908b from Florida and concluded that the species is distinct and separate from Recent *Trachemys scripta scripta*, based on wedge-shaped costals and hexagonal neural plates. We have therefore removed *T. sculpta* from the synonymy of *T. s. scripta* and from this checklist. He also examined type figures of *Emys euglypha* † Leidy 1889a (type specimens apparently lost) and *Trachemys delicata* † Hay 1916a (a single costal bone), and interpreted both as being undiagnosable. We have therefore also removed *E. euglypha* and *T. delicata* from the synonymy of *T. s. scripta*. He found *Emys petrolei* † Leidy 1868a, *Pseudemys bisornatus* † Cope 1878, and *Trachemys trulla* † Hay 1908b to be undiagnosable as well, and we have therefore removed all three from the synonymy of Recent *Trachemys scripta elegans*.

Actinemys: Based on a review of *Clemmys hesperia* † Hay 1903, which had recently been determined to be Late Miocene instead of Pliocene, Vlachos was unable to identify the plastral fragment as anything but a Testudinoidea sp. indet., and therefore removed it from the synonymy of *Actinemys marmorata*, as we have done as well.

Terrapene: Vlachos also provided extensive taxonomic re-evaluations of all fossil Pleistocene taxa of Box Turtles. We acknowledge these and have generally followed his recommendations; removing most of them from the synonymy of *T. carolina* and from this checklist and incorporating them in our next updated edition of TEWG (2015). A few are retained as junior synonyms of Recent taxa based on his recommendations.

Cistudo eurypygia † Cope 1870b from Maryland and *Toxaspis anguillulatus* † Cope 1899 from Pennsylvania were synonymized with each other, and *Terrapene eurypygia* recognized as a separate and distinct species from Recent *Terrapene carolina carolina*. *Terrapene antipex* † Hay 1916 from Florida and *Terrapene bulverda* † Hay 1921 from Texas were placed in the synonymy of *Terrapene canaliculata* Hay 1907 from Georgia and *T. canaliculata* in the synonymy of *Terrapene putnami* † Hay 1906 from Florida, which was itself separated out as a diagnosable extinct species and removed from the synonymy of either Recent *Terrapene carolina bauri* or *Terrapene carolina major.* This is in agreement with Ehret et al. (2013), but disagrees with Butler et al. (2011), who synonymized *T. putnami* under Recent *T. c. major,* and Martin et al. (2013), who recognized it as a separate extinct subspecies, *T. carolina putnami* (see TTWG Annotations 11:8 and 14:28).

Terrapene llanensis † Oelrich 1953 from Kansas, previously synonymized under Recent *Terrapene carolina triunguis*, was considered indistinguishable from *T. canaliculata* and was therefore synonymized under the extinct species *T. putnami*. *Cistudo marnochii* † Cope 1878 from Texas, *Testudo munda* † Hay 1921 from Tennessee, and *Terrapene impressa* † Hay 1924 from Texas were determined to be *nomina dubia*, only identifiable as *Terrapene* sp. indet., and have been removed from this checklist.

Terrapene formosa † Hay 1916a and *Terrapene innoxia* † Hay 1916a, both from the Pleistocene of Florida, were synonymized with Recent *T. carolina*. In view of the provenance of *T. formosa* from mid-peninsular Florida, we have tentatively synonymized it under *T. c. bauri* rather than *T. c. major,* where it was previously placed; *T. innoxia* stays under *T. c. bauri*.

Terrapene singletoni † Gilmore 1927 from Florida was determined to be a separate and distinct species, following Auffenberg (1958), and has been removed from its synonymy with *T. c. bauri*. *Trachemys nuchocarinata* † Hay 1916a, based on only a partial nuchal, was considered an undiagnosable Testudinoidea and removed from the synonymy of *Terrapene c. bauri*, where Auffenberg (1958) had placed it. *Terrapene whitneyi* † Hay 1916b from Texas was considered synonymous with Recent *Terrapene ornata ornata* instead of *T. c. triunguis*, and we have therefore reallocated it.

Gopherus: Based on what he considered to be diagnostic features of pleural-peripheral bone contacts and gular-entoplastron relationships in the Mexican Pleistocene species *Gopherus auffenbergi* † Mooser 1972 and *Gopherus pargensis* † Mooser 1980, Vlachos synonymized *G. pargensis* (see TTWG 2015) with *G. auffenbergi*, but recognized *G. auffenbergi* as a separate and distinct species from Recent *Gopherus berlandieri*. We agree with this assessment and have therefore now removed *G. auffenbergi* from the synonymy of *G. berlandieri* and from this checklist and will incorporate it in our next updated edition of TEWG (2015). He also removed *Gopherus huecoensis* † Strain 1966 from the Pleistocene of Texas from the synonymy of Recent *Gopherus flavomarginatus*, based on a more elongated medial carpal centrale and possible separation between the distal radius and the distal carpals in the drawings of the species. Instead, he synonymized it with the larger-bodied Pleistocene species *Testudo hexagonata* † Cope 1893, a synonym of *Testudo laticaudata* † Cope 1893, as discussed in TEWG (2015). We do not agree that these findings are necessarily differentially diagnostic of separate species status for the single specimen of *G. huecoensis*, and therefore tentatively retain it in the synonymy of *G. flavomarginatus,* as previously concluded by Bramble (1982) and Reynoso and Montellano-Ballesteros (2004). Additionally, Vlachos considered the Pleistocene taxa *Testudo atascosae* † Hay 1902 from Texas and *Gopherus praecedens* † Hay 1916a from Florida as undiagnosable *nomina dubia* of Testudinidae, sp. indet., and did not synonymize them under any other species, and we agree, removing them from the synonymy of *G. polyphemus*.

21:53. ***Graptemys***: Thomson et al. (2018) performed a range-wide phylogeographic time-calibrated divergence analysis of all *Graptemys* taxa using 18 nuclear and 2 mitochondrial genes from 55 individuals representing all Recent taxa. Their phylogeny and divergence time estimates suggested that diversification within the genus has been both more recent and more rapid than has so far been suspected. Their results confirmed the existence of three reciprocally monophyletic clades of map turtles: 1) a deeply divergent clade corresponding to *geographica*, 2) a distinct megacephalic clade for *barbouri*, *ernsti*, *pulchra*, *gibbonsi*, and *pearlensis*, with each species distinctive at a shallow level, and 3) a distinct western clade comprised of a monophyletic sawback clade (*flavimaculata, oculifera*, and *nigrinoda*) and a monophyletic Texas clade (*versa* and *caglei*), but with a paraphyletic and

unresolved narrow-headed false map turtle complex (*pseudogeographica*, *kohnii*, *ouachitensis*, and *sabinensis*) within the western clade that may require taxonomic revision. They made no specific recommendations for taxonomic changes, but did not agree with Praschag et al. (2017), who had suggested that all of the megacephalic taxa except *barbouri* were conspecific, and that among the sawbacks *flavimaculata* and *nigrinoda* should be treated as subspecies of *oculifera* (see TTWG Annotation 17:19). However, Thomson et al. supported the view previously affirmed by Lindeman (2013) that the narrow-headed *sabinensis* is a distinct species (see TTWG Annotation 14:18), as also supported by Praschag et al. (2017).

21:54. ***Malaclemys terrapin***: Lovich and Hart (2018) reviewed the previously published taxonomic literature on this species and concluded that it is oversplit, and that there are only four discrete genetic lineages, or Management Units, that do not match the current taxonomy of seven subspecies (following Hart et al. 2014; see TTWG Annotation 17:24). They made no taxonomic recommendations other than recommending further phylogeographic studies based on more complete distributional sampling and more robust genetic analyses, which are currently underway.

21:55. ***Trachemys grayi emolli***: The taxonomic status of the old unused name *Emys valida* Le Conte 1860 has been uncertain for a long time. In our previous checklists we listed it as a synonym of *Trachemys venusta venusta*, in accordance with the checklist by Fritz and Havas (2007). It was originally synonymized with question marks under *Chrysemys ornata* by Boulenger (1889) and under *Pseudemys ornata ornata* by Wermuth and Mertens (1961), and then without a question mark under the synonymy of *Pseudemys scripta venusta* by Smith and Smith (1980). The taxon was based on a single specimen collected in "Honduras" (no precise locality given) by Le Conte's son, and was described as having a shell length "nearly two feet long" (= nearly 60 cm, but possibly representing curved carapace length). The shell has since been lost, but the head and soft parts of the holotype remain (Malnate 1971; McCranie 2018).

McCord et al. (2010), based on the stated shell size and an examination of the head markings and tomial morphology of the type specimen, synonymized *E. valida* with what they defined as the Pacific versant taxon *Trachemys venusta grayi* (= *T. grayi grayi*), which they stated reaches a maximum shell length of 60 cm. They also restricted the type locality of *E. valida* to the "Pacific coastal drainages of Honduras" and declared the name *Emys valida* Le Conte 1860 a *nomen oblitum*. We agree with them and Smith and Smith (1980) that the name has not been recognized as a valid species since its description, but in our opinion, it should not be synonymized with *T. grayi grayi* (as currently defined), which occurs along the Pacific versant of southern Mexico, Guatemala, and El Salvador, but not in Honduras, and is smaller, with a maximum shell length to about 40 cm (McCranie et al. 2013; Legler and Vogt 2013). Nor should it be synonymized with *T. venusta venusta*, which is a Caribbean versant taxon that does not occur in Honduras and only reaches a maximum shell length of about 42 cm, nor with *T. venusta uhrigi* from the Caribbean versant of Honduras, which only reaches a CL of about 44 cm (McCord et al. 2010; Legler and Vogt 2013). McCranie (2018) also disagreed with McCord et al. (2010) and stated that *E. valida* should not be synonymized with *T. grayi* (= *T. grayi grayi*), and left it in the synonymy of Caribbean versant *T. venusta* (= *T. venusta venusta*).

Where in Honduras the type specimen was collected is therefore of relevance. Le Conte's son, John Lawrence LeConte [1825–1883], who collected the specimen, was a physician and well-known entomologist who explored Honduras in 1857 as the surgeon for the survey team working on the Honduras Interoceanic Railway that transversed the country from Port Caballos (now Puerto Cortes) on the Caribbean coast to the Gulf of Fonseca on the Pacific coast (Squire 1870). It is therefore reasonable that he may have collected the type specimen in the Pacific versant of Honduras, and we agree with the restricted type locality by McCord et al. (2010). However, instead of agreeing with their synonymization, we instead provisionally synonymize *E. valida* Le Conte 1860 with *T. grayi emolli*, which does occur along Pacific coastal Honduras (McCranie et al. 2013), and reaches a carapace length of at least about 54 cm (Ernst 2008). Since *E. valida* is a *nomen oblitum*, it should not take nomenclatural precedence over *T. grayi emolli* (Legler 1990). This provisional synonymization may be resolved through genetic analysis of the remaining soft tissue parts of the holotype of *E. valida*, as suggested by McCranie (2018).

21:56. ***Trachemys hartwegi***: Forstner et al. (2014) and Parham et al. (2015) elevated the subspecies *Trachemys gaigeae hartwegi* to species status based on mitochondrial and nuclear DNA differences between it and *T. g. gaigeae*. Parham et al. (2015) also demonstrated its close relatedness to *T. taylori* and *T. venusta*. We overlooked these findings regarding *hartwegi* in our last checklist (although we noted other taxonomic updates by Parham et al. in TTWG Annotation 17:26), and accept *T. hartwegi* as a full species at this time.

21:57. ***Trachemys medemi***: Vargas-Ramírez et al. (2017) analyzed mtDNA and nuclear DNA sequences of *Trachemys* from the isolated Río Atrato basin of northwestern Colombia and compared them with data from many other *Trachemys* (including taxa variously classified by them or others as species or subspecies of *T. grayi*, *T. venusta*, *T. callirostris*, and *T. dorbigni*). They concluded that the Río Atrato lineage was distinctive, and described it as a new species, *T. medemi*, more closely related to *T. dorbigni*, the more distant and older radiation of *Trachemys* into South America, than to any of the more geographically proximate taxa. We accept their interpretation and also recognize this new species. In our previous checklist we had included these populations under *Trachemys venusta uhrigi*.

21:58. ***Trachemys scripta* and *T. gaigeae***: Two recent papers (Vamberger et al. 2020a; Parham et al. 2020) have investigated the genetics and phylogeography of the three recognized subspecies of the widespread species *Trachemys scripta*, with Parham et al. (2020) also examining the relationships of western *T. scripta elegans* with *T. gaigeae*.

Vamberger et al. (2020a) sampled mitochondrial and nuclear DNA sequences and microsatellite loci of several populations of *T. s. scripta*, western and eastern *T. s. elegans*, and *T. s. troostii*. They found that only the quickly evolving microsatellite loci discriminated *T. s. troostii* and western *T. s. elegans* from eastern *T. s. elegans* and *T. s. scripta*, while eastern *T. s. elegans* and *T. s. scripta* were not distinct from each other in any sampled marker system. They also found that western *T. s. elegans* is morphologically distinct and differs from eastern *T. s. elegans* and *T. s. scripta* with respect to microsatellites as much as *T. s. troostii* does, and concluded that the currently recognized subspecies are weakly differentiated. However, if subspecies within *T. scripta* are continued to be recognized, they recommended that distinct subspecies status should be considered for western *T. s. elegans*. They did not examine specimens of *T. gaigeae* from the Rio Grande drainage to determine how closely related that taxon is to western *T. s. elegans*. Their findings challenged the current intraspecific systematics of *T. scripta*, but they refrained from synonymizing any taxa, since abandoning the well-established and morphologically distinct subspecies of *T. scripta* was not deemed desirable.

Parham et al. (2020) sampled mitochondrial DNA sequences and SNPs from ddRAD libraries of several populations of *T. s. scripta*, western and eastern *T. s. elegans*, *T. s. troostii*, and *T. gaigeae* from the Rio Grande drainage. They found that some samples of *T. s. troostii* were not genetically distinct from *T. s. elegans* and some shared morphological affinities with *T. s. scripta*. They also found genetic admixture and morphological intergrades suggesting introgression between *T. gaigeae* and *T. s. elegans* in the Pecos and lower Rio Grande rivers of Texas and New Mexico, but recommended that these populations continue to be considered *T. s. elegans*.

The taxonomic status of *T. s. troostii* is particularly complicated: Vamberger et al. (2020a) recovered it as distinct in their microsatellite data, whereas Parham et al. (2020) concluded that it represents an intergrade between *T. s. elegans* and *T. s. scripta* and recommended that it should be synonymized. Unfortunately, neither study sampled the lower Tennessee River basin and the sampling and analysis of *T. s. troostii* in the Cumberland River basin was not very robust; hence, it appears premature to make any conclusions about its taxonomic status.

In our opinion, more data based on thorough geographic sampling across the range of *T. s. elegans*, and especially in the areas of potential intergradation among *T. s. scripta*, *T. s. elegans*, and *T. s. troostii* in eastern Tennessee, are needed before we are comfortable making any definitive taxonomic decisions or changes affecting any of these taxa.

Although it appears possible based on Parham et al. (2020) that *T. gaigeae* and *T. s. elegans* could potentially be considered conspecific, with the Pecos and lower Rio Grande rivers the areas of intergradation and co-occurrence, further sampling and analysis is strongly recommended to evaluate their relationships with each other and other closely related *Trachemys* taxa.

21:59. ***Trachemys terrapen***: The currently prevailing hypothesis that this taxon is native to Jamaica and possibly prehistorically introduced to the Bahamas remains unresolved and problematic. Seidel and Ernst (2017) postulated that the presence of *T. terrapen* in the Bahamas might have been the result of introduction by British trade from Jamaica, but also noted that fossil evidence suggested a natural origin of the species in the Bahamas. Indeed, Olson et al. (1990) documented the presence of *Trachemys* fossils from a possibly Late Pleistocene site at the southern tip of San Salvador island just northeast of Great Bahama Bank, and Berman (1994) excavated subfossils of *Trachemys* from the same island from a Lucayan archeological site dated to the Late Holocene (660–860 CE). A few Pleistocene fossils of *Trachemys*, identified as *T. terrapen,* have also been found in breccia cave deposits on Jamaica, dated as being up to about 100,000 yrs old (Morgan 1993).

Recent specimens of apparently native *T. terrapen* have only been found on the islands of Great Bahama Bank (Cat, Eleuthera, Exuma, Long, and South Andros), all of which would have been emergent as a single large palaeoisland during the last glacial maximum ca. 17,000 yrs ago. Populations of *Trachemys* sp. found on Little Bahama Bank and the developed island of New Providence (Nassau) on Great Bahama Bank all appear to be hybrids of various *Trachemys* species: *terrapen*, *decussata*, *stejnegeri*, and *scripta*, notably *T. s. elegans*. Comparative genetic analysis of Jamaican and native Bahamian populations of *T. terrapen* is needed to ascertain whether they possibly represent separate evolutionary lineages or Management Units, but based on fossil evidence, and the previous widespread presence of the extinct Holocene tortoise *Chelonoidis alburyorum* in the Bahamian Archipelago (Franz et al. 2020; Steadman et al. 2020; Kehlmaier et al. 2021a), there is a possibility that the Bahamian population of *T. terrapen* is native, as previously suggested by Seidel (1996) and Lee and Ross (2001). Based on the age of *Trachemys* fossils from the Bahamas and Jamaica, it appears possible that natural dispersal of the species may have occurred from Jamaica to the Bahamas during the Late Pleistocene, although prehistoric human-mediated introduction by the indigenous Lucayan people who arrived in the Bahamas Archipelago from the south at about 600 to 750 CE (Fall et al. 2021) is also a distinct possibility. In either case, *T. terrapen* has been present in the Bahamas since about that time or even earlier.

21:60. ***Emydoidea blandingii***: Jordan et al. (2019) examined variation in two mitochondrial and three nuclear genes across the range of *E. blandingii*. Although they found relatively low levels of genetic variation in either data set, both supported two divergent lineages: one in eastern Ontario, eastern New York, and Nova Scotia, and another across the rest of the species' range (from central Ontario and western New York to Nebraska). They theorized that this divergence was likely reflective of previous refugial isolation of populations on either side of the Appalachian Mountains. In addition, the mitochondrial data set alone suggested a divergence between populations in Nebraska from those across the rest of the western genetic cluster. However, all of these divergences were shallow and they did not suggest any taxonomic recognition, but highlighted the need for appropriate conservation management.

21:61. ***Emys orbicularis***: Based on dense sampling across the Iberian Peninsula and adjacent regions, Pöschel et al. (2018) examined genetic differentiation and gene flow in *E. orbicularis* using 15 microsatellite loci and 1031 bp of the mitochondrial cytochrome b gene. Their results confirmed the presence of three distinct taxa (*E. o. orbicularis*, *E. o. galloitalica*, *E. o. occidentalis*). According to simulations using Approximate Bayesian Computing, *E. o. orbicularis* circumvented the Pyrenees in the Holocene and invaded the northern Iberian Peninsula, leading to local hybridization with *E. o. galloitalica* along the northern Mediterranean coast of the peninsula, and with *E. o. occidentalis* south of the Pyrenees. Pure populations of *E. o. occidentalis* occur in most of the Iberian Peninsula; however, the population along the inland course of the Ebro River has introgressed mitochondrial haplotypes of *E. o. orbicularis*. Hybridization between the three subspecies was found to be surprisingly low, with steep genetic clines typically seen in distinct species. Yet, the authors recommended no taxonomic changes, pending more comprehensive analyses involving unstudied populations from outside the study region.

21:62. ***Emys trinacris***: Speybroeck et al. (2020), representing the Taxonomic Committee of the Societas Europaea Herpetologica, reviewed several studies concerning the relationships between *Emys trinacris* and *E. orbicularis* (e.g., Fritz et al. 2005a, 2007b; Spinks and Shaffer 2009; Vamberger et al. 2015; Vamberger and Fritz 2018). Although without full consensus, they took a conservative stance and decided to treat *trinacris* as a subspecies of *E. orbicularis*, pending further studies resolving the complicated relationships of this complex. However, we continue to treat *E. trinacris* as a separate species because of its allopatry and morphologic and genetic distinction.

Vecchioni et al. (2020) studied the phylogeography of *E. trinacris* across its range, using 16 microsatellite loci. Analysis of 245 specimens demonstrated geographically based structuring into five well-characterized clusters; the authors recommended that they be considered as independent Management Units.

21:63. ***Glyptemys insculpta***: Based on variation in nine microsatellite loci across the range of *G. insculpta* in Canada, Bouchard et al. (2019) observed low-level divergence between watershed populations on the northern versus southern sides of the St. Lawrence River, and a tendency for divergence among individual sub-drainage systems. Their results confirmed isolation-by-watershed for wood turtle genetic structure, with implications for population management, but they did not recommend that individual watershed-based populations warranted recognition as Evolutionarily Significant Units or even as Management Units. However, they did recommend that separate watershed populations be considered as potential Designatable Units under Canadian conservation laws.

21:64. ***Terrapene***: The North American box turtles have continued to receive taxonomic attention, with some improved resolution; see also previous TTWG Annotations 11:8, 14:27, and 17:37. However, the genus remains one of the most troubling among chelonian taxa.

Vitek (2018) quantified shell morphology of fossil and extant USA *Terrapene carolina*-group populations based on geometric morphometrics of 156 carapacial landmarks. In her PCA analysis, *T. c. carolina, T. c. bauri, T. c. major*, and *T. c. triunguis* all overlapped extensively in carapacial morphospace and jackknife assignment of individuals to subspecies returned relatively low classification accuracy (*carolina*, 42–60%; *major* 40–58%; *bauri*, 59–65%; *triunguis*, 63–74%). Based on this morphometric overlap, Vitek could not identify diagnosable differences among the four taxa, and concluded that there was clinal variation in size-corrected shell shape, rather than distinct groups corresponding to recognized subspecies.

Based on sequencing data from > 11,000 ddRADseq loci sampled for 320 individuals, Martin et al. (2020) investigated levels of introgression among eastern USA populations of *Terrapene* (*carolina, triunguis, major, ornata*), but excluding *bauri*. They found support for five groups: *carolina* across much of the eastern USA, two groups of *major*, one in Alabama and Mississippi and another in the Florida Panhandle, *triunguis* across central USA, and *ornata* in the western prairie states, with weaker support for an additional split between northeastern (Illinois, Wisconsin, Iowa) and southwestern (Kansas, Texas, Colorado, Nebraska) populations of *ornata*. They also recovered variable genetic admixture between *carolina* and *major* across Alabama and the Florida Panhandle, *carolina* and *triunguis* in Georgia and South Carolina, and *major* and *triunguis* in Mississippi and southern Alabama, but a lack of admixture between the two populations of *major*. The authors made no taxonomic recommendations, but did recommend the recognition of the two populations of *major* as Evolutionarily Significant Units (ESUs) or Management Units (MUs). In addition, despite the admixture between *carolina–major* and *triunguis*, they argued that the genomic clines between these taxa were steep enough to support their recognition as separate species.

In a follow-up study employing ddRADseq analysis of nearly 15,000 SNPs, Martin et al. (2021) generated phylogenetic trees by three methods for all taxa of *Terrapene* except *klauberi*. They also employed species delimitation and machine learning protocols to further examine possible species limits in the genus. Sampling levels varied, but notably were either a single or at most a few individuals for *nelsoni, bauri, yucatana, mexicana*, and *coahuila*, often with large sampling gaps. A time-calibrated tree (IQ-TREE) recovered the strongest statistical support for clades containing *T. ornata* (with variable support for reciprocally monophyletic subspecies *T. o. ornata* and *T. o. luteola*), *T. nelsoni*, *T. bauri*, and a large clade with all remaining taxa. Their SVDquartets species tree resolved *T. ornata* sister to *nelsoni*, and that clade was sister to the remaining taxa, which comprised a monophyletic group with limited statistical support for most contained subclades. Their PoMo tree was more highly resolved, and returned *T. ornata* as sister to *nelsoni*, and that clade as sister to the remaining taxa.

Within that larger group, *mexicana* + *yucatana* were recovered as a clade sister to the rest, followed by *triunguis* as sister to the remaining taxa; in addition, *bauri* was resolved as sister to *carolina*. In all of these analyses, *coahuila* was resolved as sister to or nested within the clade of the western (Mississippi and southern Alabama) population of *major*.

The species delimitation analyses of Martin et al. (2021) returned a variety of results, depending on the method and data sets used. The consistently recovered species were *ornata* (but not the subspecies within it), *bauri, triunguis, carolina* and *major*. Based on these results, the authors concluded that recognition of *T. ornata* as a full species was strongly supported, although the evidence for the recognition of *luteola* as a subspecies was weak (see also TTWG Annotation 14:27). They also supported the recognition of *T. nelsoni* as a full species. The most consistent divergence among the remaining samples was between the *mexicana–yucatana–triunguis* and *carolina–bauri–major–coahuila* clades.

Given these results and the morphological distinctions previously noted by Minx (1996), the authors argued for species recognition for *T. mexicana* (including *triunguis* and *yucatana* as subspecies), as they did in 2013 (see TTWG Annotation 17:37). The authors also noted the enigmatic position of *T. coahuila* as a full species deep within the *carolina* clade, most closely related to the western population of *major*, but did not suggest taxonomic revision. Finally, because the boundaries of, and relationships among *carolina, bauri*, and *major* were inconsistently resolved, the authors recommended they be retained as subspecies of *T. carolina*.

Although the work of Martin et al. is a major step forward, there is still very considerable uncertainty about the lineages within *Terrapene*, their interrelationships, and the resolution of a consistent, lasting taxonomy. *Terrapene ornata* and *T. nelsoni* are almost universally well supported, and we agree that they should be considered full species. In addition, barring any new evidence, we retain within *nelsoni* its two contained subspecies, *nelsoni* and *klauberi*, but place *luteola* in the synonymy of *ornata* rather than retain it as a separate subspecies.

In view of the marked discrepancies among the various trees and delimitation models, we consider four possibilities for the remaining taxa *carolina, bauri, major, triunguis, mexicana*, and *yucatana*: 1) that all should continue to be recognized as subspecies of *carolina*, 2) that all should be recognized as separate species, 3) that they should be recognized as two species (*carolina* and *mexicana*) with three subspecies each, or 4) that *triunguis, mexicana*, and *yucatana* should be considered as separate species (especially in view of their widely allopatric distributions), and that *carolina* should include either both subspecies *major* and *bauri*, or possibly only *major*, with *bauri* possibly recognized as a separate species. We anticipate more genomic work, and especially more detailed field sampling, to be necessary to further clarify these complex relationships.

Despite these uncertainties and multiple possibilities, and in light of the results of these most recent genetic analyses by Martin et al. (2020, 2021) and the morphologic data provided by Minx (1996), most of us are reasonably confident in 1) recognizing *T. nelsoni, T. ornata, T. carolina, T. mexicana, T. yucatana*, and *T. coahuila* as separate species, 2) sinking *luteola* as a subspecies of *T. ornata*, with the recognition that more complete genetic sampling may revise this decision, and 3) retaining *major* and (with less certainty) *bauri* as subspecies of *T. carolina*. As for *triunguis*, which Martin and colleagues view as a subspecies of *T. mexicana*, we agree that it resolves as a separate lineage in most analyses, but do not agree that there is strong enough support for its phylogenetic position to place it unambiguously within, or sister to, *mexicana*. Combined with its unique color pattern and restricted, primarily first and second generation-restricted hybridization (Martin et al. 2020), and in view of its genetic distinctiveness, we tentatively recognize the Three-toed Box Turtle as a separate species, *T. triunguis*, rather than as a subspecies of either *carolina* or *mexicana*.

21:65. ***Terrapene carolina carolina***: Linnaeus (1758) cited his description of *Testudo carolina* as being sourced only from Edwards (1751:pl.205). He did not cite any of his own previous publications on turtle specimens that he had examined (Linnaeus 1749a, 1749b, 1754), as he did for three other species he described (*Testudo mydas, Testudo lutaria*, and *Testudo geometrica*), nor any museum specimens, as he did later (Linnaeus 1766) for *Testudo denticulata* (Museum De Geer), when he also cited his own previous work (Linnaeus 1764) as the source for *Testudo serpentina*. It therefore appears that he did not examine any actual specimens of *T. carolina* in his or others' collections, nor have any been identified or listed since (Linné and Thunberg 1780; Thunberg 1828; Lönnberg 1896; Andersson 1900; Holm 1957; Ernst and McBreen 1991; Dodd 2001; Wallin 2001). As such, the specimen figured in Edwards (1751:pl.205) should be recognized as the holotype of *Testudo carolina* Linnaeus 1758, and we formally recognize it as such. This is also consistent with and similarly reasoned as for the other plate by Edwards (1751:pl.204), where the specimen figured there was recognized by Bour (1987a) as the holotype for *Testudo graeca* Linnaeus 1758, a species also cited as being sourced only from Edwards (1751:pl.204).

21:66. ***Testudo carinata***: There has been considerable long-standing confusion surrounding the identity of *Testudo carinata* Linnaeus 1758, which was not described as being based on any known specimens or previously cited references. Its brief original diagnosis (*Testudo pedibus digitatis, testa gibbosa, scutellis dorsalibus quatuor anterioribus carinatis, sterna integro* [Tortoise with digitated feet, gibbose shell, four anterior dorsal scutes carinated, plastron entire]) did not change in subsequent publications by Linnaeus (1766) or Gmelin (1789), and Linné and Thunberg (1780) did not list the species as being present in the Linnaean collection in Uppsala (UUZM). However, Lacepède (1788:164) described a ca. 15 cm specimen in the collection of Lamarck in Paris that he suggested was possibly the same species as *T. carinata* Linnaeus 1758, calling it La Bombée, a reference to its high-domed shell (shell depth measured ca. 43% of its length). Subsequently, Shaw (1802:35) described a smaller, ca. 7.6 cm specimen, in the Leverian Museum in London as also questionably representing *T. carinata*, based on its moderately convex shell and prominent yellow vertebral carina. Both of these specimens were probably *Terrapene* spp. (in which shell depth typically measures ca. 39–50% of shell length) and Merrem (1820:28) synonymized *T. carinata* Linnaeus 1758 and the Lacepède and Shaw descriptions under *Terrapene clausa* (= *Terrapene carolina*). Subsequently, Gray (1831d:18), who may have seen the actual Lacepède and Shaw specimens, synonymized *T. carinata* Linnaeus 1758 under *Cistudo carolina* = *Terrapene carolina carolina* (Linnaeus 1758), where it has stayed ever since, although without clarity, in essence a *nomen dubium*.

However, Holm (1957) and Wallin (2001) noted that a Linnaean-era specimen in the UUZM collection (from Museum Gustavi Adolphi, MGA) labeled *T. carinata* by Thunberg (1810, 1828) and possibly representing the type specimen of *Testudo carinata* Linnaeus 1758, was not a *Terrapene carolina*, but a *Nicoria spengleri* (= *Geoemyda spengleri*), potentially threatening the nomenclatural priority of *Testudo spengleri* Gmelin 1789. This specimen (UUZM 277, formerly MGA 46), labeled by Thunberg, is a typical *G. spengleri* with a relatively flat (depth ca. 35% of length) markedly tricarinate shell with a prominent plastral anal notch, clearly not conforming to the initial diagnosis of *T. carinata* Linnaeus 1758. The specimen was accessioned into the Uppsala collection by Thunberg (1810) at the same time as the type of *Testudo tricarinata* Retzius *in* Schoepff 1792 (= *Kinosternon scorpioides*), and possibly identified initially by Thunberg as a *Testudo tricarinata* Bory de Saint-Vincent 1804 (a junior synonym of *Testudo spengleri*), but then replacing the specific epithet *tricarinata* with *carinata* in order to avoid confusion with the Retzius specimen with the same name. In conclusion, *Testudo carinata* Thunberg 1810 is a *nomen nudum* and = *Geoemyda spengleri*, *Testudo carinata* Linnaeus 1758 is a *nomen dubium* and most likely = *Terrapene carolina carolina*. The specimen UUZM 277 (MGA 46) has no status as a type specimen of *T. carinata* Linnaeus, and the nomenclatural priority of *Geoemyda spengleri* is not affected.

21:67. ***Batagur baska*** **and** ***B. kachuga***: Vyas (2017) reported on a specimen of "*Batagur baska*" recently collected from Rudramata Dam, Kutch, Gujarat, western India, but the photo of the large female specimen is that of a *Batagur kachuga* (P. Praschag, S. Singh, and P.P. van Dijk, pers. comm.), and most likely represents a released trade specimen. However, previous records of *B. baska* from the nearby Indus River drainage of Pakistan are known and are reviewed by Vyas (2017). Murray (1884a) reported *B. baska* to occur in the Indus River, Sindh, Pakistan, and a specimen in the Vienna museum (NMW 1841) labeled as having been collected in the 1850s in the Indus River Delta of Sindh was genotyped by Praschag et al. (2008) and shown to be *B. baska*. In addition, archeological specimens of *B. baska* have been described from Mohenjo-daro, Indus River, Pakistan (Seymour Sewell and Guha 1931), as well as from Sarnath, Ganges River, Uttar Pradesh, India (Nath 1959), both sites far inland from the typical

coastal habitat of the species and are either representative of previous trade in the species or misidentified specimens. It remains possible, however, that a disjunct population of *B. baska* once existed in the Indus River Delta and the coastal mangrove wetlands extending east to the Gulf of Kutch in westernmost Gujarat, India.

21:68. ***Cuora galbinifrons* complex**: Liu et al. (2019) examined the concordance of morphometric and molecular data for specimens in the *C. galbinifrons / bourreti / picturata* complex from a turtle farm (i.e., without locality data). Although their morphometric data showed some differences among the three taxa, they failed to standardize their data for body size in their multivariate analysis. Hence, those results likely reflect differences in body size rather than relative morphometrics. Their phylogenetic analysis of sequence data from a single nuclear gene (Rag1) was poorly resolved, and employed a distantly related outgroup rather than a congener. Their phylogenetic analysis of the entire mitogenome for ten *Cuora* species employed only a distantly related trionychid outgroup, but nevertheless fully resolved all ten taxa, with *picturata* sister to the *bourreti-galbinfron*s clade. Finally, they repeated the latter analysis focusing only on partial COI mitochondrial DNA gene sequences. That analysis also employed a distant outgroup, and the resulting tree was poorly resolved, with no evidence of repicrocal monophyly for any of the three taxa. However, despite these inconclusive results, the authors argued that *picturata* should be recognized as a full species but that *bourreti* should be relegated to a subspecies of *galbinifrons*. Pending more complete and appropriate morphometric and molecular analyses, we continue to recognize the three taxa as full species.

21:69. ***Mauremys annamensis* and *M. mutica***: In a follow-up to their earlier preliminary study (Fong et al. 2007) that had suggested that *M. mutica* was paraphyletic to *M. annamensis*, Fong et al. (2019) examined variation in the ND4 mitochondrial gene and the R35 nuclear gene from a wider sampling of wild-caught, captive, and museum specimens of *M. annamensis* and *M. mutica* across Vietnam, China, and Taiwan (see also Zhou et al. 2015 and Zhao et al. 2016a,b; TTWG Annotation 17:49).

The mitochondrial phylogeny identified two well-supported clades: a "true" *mutica* clade including topotypic specimens from eastern China and samples from Taiwan; and a clade including *mutica* from Hainan Island and Vietnam and *annamensis* from Vietnam. The latter clade included three subclades: one for *mutica* specimens from Hainan, a second including *annamensis* from turtle farms and seizures in Vietnam, and a third including known-locality *annamensis*, GenBank sequences from controversial "*M. guangxiensis*" specimens, and Vietnamese *mutica* specimens.

The nuclear phylogeny also identified two major clades, one including Hainan *mutica* specimens and the other including a monophyletic *annamensis* clade nested within the Chinese and Taiwanese *mutica* specimens. Both analyses were hampered by the inclusion of samples of questionable provenance and original identification, but it is clear that there are at least two distinct clades within *mutica*: eastern China and Taiwan (nominotypic material) and Hainan and Vietnam (for which the name *schmackeri* Boettger 1894 may be applicable). However, the relationship between *mutica* and *annamensis* remains unclear based on the discordance between the mitochondrial and nuclear DNA trees. Broader nuclear sampling, including material from southern China and the Ryukyu Islands (*M. mutica kami*) will be necessary to resolve the taxonomic relationships among these various taxa; in the meantime we continue to recognize *mutica* and *annamensis* as full species.

21:70. ***Mauremys japonica***: Suzuki and Hikida (2011) analyzed the phylogeography of this endemic Japanese species using mtDNA sequences (cytochrome b gene and control region). They analyzed 238 wild-caught specimens from 43 localities and determined that there were two major haplotype groups: a western one from Kyushu and western Honshu, and an eastern one from Shikoku and eastern Honshu, representing the result of previous isolation of the two lineages in separate glacial refugia, with a secondary postglacial contact zone in Chugoku District of western Honshu. They made no taxonomic recommendations.

21:71. ***Mauremys reevesii***: Bu et al. (2019) examined variation in 12 microsatellite markers for five captive populations of *M. reevesii* in China, and found two genetic clusters: one from Anhui Province in the north and the other from southern China and Hainan Island. Since the samples represented long-term captive-breeding populations, and were not necessarily representative of local, natural populations, the authors made no taxonomic judgments. However, given the evidence of significant geographic variation in *M. reevesii* (see Iverson et al. 1989 and TTWG Annotation 17:52), range-wide molecular studies of known-locality specimens are sorely needed.

21:72. ***Melanochelys trijuga***: Schweigger (1812) described *Emys trijuga* based on a specimen collected by Lechenault [Leschenault] on the island of Java, Indonesia, far from the native range of *Melanochelys trijuga* in India. The holotype specimen was deposited in MNHN, but has been lost (Bour and Schmidtler 2014). Fortunately, Nikolaus Michael Oppel [1782–1820], who also worked for a while at MNHN (Oppel 1811) and was friendly with Schweigger, painted many MNHN turtle specimens between 1807 and 1810, including the type of *Emys trijuga* Schweigger, and these paintings have been made accessible by Bour and Schmidtler (2014). The holotype of *Emys trijuga* (shown in their Fig. 9 and Oppel's Plates 15–16), labeled by Oppel as "*Testudo trijuga* Schweiger," appears to be a Southeast Asian *Malayemys subtrijuga* (or *M. macrocephala*), instead of *trijuga*, which has been recognized as an Indian species since Duméril and Bibron (1835) and as *Melanochelys trijuga* since Gray (1869a). Bour and Schmidtler (2014) also pointed out that original illustrations of the type specimens of both *Emys herrmanni* Schweigger 1812 and *Emys belangeri* Lesson 1831b represent specimens of what is now considered to be the polytypic species *Melanochelys trijuga*, but that recognizing either of these names, or recognizing *Emys trijuga* Schweigger 1812 as having nomenclatural priority over *Emys subtrijuga* Temminck and Schlegel 1845, would create widespread nomenclatural chaos, affecting the names of several taxa and genera. Instead, they recommended maintenance of nomenclatural stability through the designation of a neotype for *Emys trijuga* Schweigger 1812. Reflecting these findings, we have now moved *Emys herrmanni* Schweigger 1812 from the synonymy of *Malayemys subtrijuga* to the synonymy of *Melanochely trijuga*, but we leave *Emys trijuga* as the type species for *Melanochelys*, pending a neotype designation.

21:73. ***Sacalia quadriocellata***: Lin et al. (2018a) recognized the population of this species from Hainan as specifically distinct from mainland Guangdong populations, based on differences in coloration pattern and previously-reported mtDNA variation, and resurrected the name *Clemmys quadriocellata insulensis* Adler 1962 for it, elevating it to species status as *Sacalia insulensis*. This had previously been preliminarily recommended by Shi et al. (2008) based on genetics (see TTWG Annotation 08:24). Subsequently, Lin et al. (2020) compared the nearly complete mitogenome of a single specimen each of *S. quadriocellata*, *S. bealei*, and *S. insulensis* from Hainan with three *Sacalia* sequences from GenBank. They found that *S. bealei* was 7.8% divergent from the other samples, and that their single Hainan sample differed from mainland *S. quadriocellata* by 2.8%. However, in a range-wide phylogeographic study of mtDNA variation in *S. quadriocellata*, Le et al. (2020) identified three distinct geographic clades, one from southern Vietnam and central Laos, a second from northern Vietnam and northeastern Laos, and a third from southern Guangdong and Hainan, but did not consider any of them distinctive enough to warrant taxonomic recognition and specifically did not recognize *S. insulensis* as separate. As such, we continue to retain *S. insulensis* in the synonymy of *S. quadriocellata*.

The type locality of *Sacalia quadriocellata* Siebenrock 1903, originally designated only as "Annam," was subsequently restricted by Siebenrock (1909) to "Annam, Phuc Son" [= Phuoc Son, Quang Nam, Vietnam], which would suggest that the name *quadriocellata* should, if warranted, be applied to the southern Vietnam clade. However, the type (NMW 23393) should be examined genetically to confirm or reject the type locality restriction. That determination is essential if future genetic sampling confirms the existence of three reciprocally monophyletic clades, and argues that those three clades merit taxonomic recognition.

21:74. ***Rhinoclemmys punctularia punctularia***: Siciliano et al. (2014) described finding two specimens of *R. punctularia* on the coastal plain of Rio de Janeiro and suggested that they might represent a native disjunct population or possibly introduced animals from the Amazon basin, the closest known population at that time occurring in Amazonian Maranhão (Pereira et al. 2013). Since then, a few more specimens of *R. punctularia* were discovered in coastal Bahia (do Valle et al. 2016), and more recently, a significant population with several animals was found on the coastal plain of Espirito Santo (Oliveira et al. 2019), suggesting that the taxon may indeed be native to the Atlantic Forest coastal plain, although in scattered disjunct locations. However, we continue to consider the two records from

Rio de Janeiro to probably represent introduced specimens. Oliveira et al. (2019) postulated that their population in Espirito Santo represented an undescribed species of *Rhinoclemmys*, but a published description has not yet appeared.

21:75. ***Manouria emys***: Based on two mitochondrial markers (COI and cytochrome b), Kundu et al. (2018b) found shallow support for the genetic distinction between the two recognized subspecies of *M. emys*. They also examined pectoral scute morphology in a small sample of tortoises from northeastern India, Myanmar, and Bangladesh, as well as captive trade specimens in that region. They found widely separated pectoral scutes in *M. e. emys* and both broad and very narrow contact of the pectoral scutes with the plastral midline in *M. e. phayrei*, and claimed to have found all three morphotypes in northeastern India within the range of *M. e. phayrei*. Further sampling of known-locality specimens will be necessary to clarify this discordance.

21:76. ***Gopherus***: Román-Palacios and Wiens (2018) generated a time-calibrated phylogeny for tortoises (Testudinidae) based on four mitochondrial and one nuclear gene in a study of evolutionary diversification rates. Their chronogram found that 10 of 11 tortoise clades that diverged in the Oligocene (older than ca. 23 mya) are currently recognized as including two or three separate genera. The one clade with only one genus, *Gopherus*, contains two clades that are apparently older, having diverged at ca. 26 mya by their calculation, but are not currently recognized as separate genera. Their analysis supports the previous arguments by Bramble (1982), Bramble and Hutchison (2014), and Franz (2014) (see TTWG Annotation 17:67) that these two morphologically and genetically distinct clades should perhaps be recognized as the genera *Gopherus* (for *polyphemus* and *flavomarginatus*) and *Xerobates* (for *agassizii, morafkai, evgoodei*, and *berlandieri*). However, the broader time-calibrated phylogenetic study by Thomson et al. (2021) found that the divergence between the two clades of *Gopherus* was younger than the divergence between the two congeneric species of *Manouria*, in contrast to Román-Palacios and Wiens (2018) who dated the divergence of the *Manouria* species at about 20 mya. Based on the morphological evidence alone, there is a good argument for the recognition of these two deep and old clades of Gopherine tortoises, and in conjunction with their deep genetic divergence, we suggest that they should perhaps be considered as subgenera, as previously concluded and recommended by Gerlach (2001).

21:77. ***Gopherus berlandieri***: Savage (2021) provided formal justification of the *nomen oblitum* status of the two names *Testudo tuberculata* Berlandier 1850 and *Testudo bicolor* Berlandier 1850, with full documentation of the *nomen protectum* status for *Xerobates berlandieri* Agassiz 1857, as per ICZN Article 23.9. The same conclusions had previously been reached by us (TTWG Annotation 14:34) and Bour (2017), with *nomen oblitum* status suggested for both names, but without full and formal ICZN-recommended justification of *nomen protectum* for *X. berlandieri*.

21:78. ***Gopherus polyphemus***: Gaillard et al. (2017) analyzed range-wide phylogeography of this species based on 20 microsatellite loci using 933 individuals from 47 sampling sites. They defined five separate well-delineated Management Units (Western, Central, West Georgia, East Georgia, and Florida), with major river systems and physiographic provinces forming the boundaries between the genetically defined regions, but did not recommend any taxonomic changes.

21:79. ***Astrochelys yniphora***: Mandimbihasina et al. (2020) examined variation in mitochondrial sequence data and nuclear microsatellite data across living and extirpated populations of *A. yniphora*, arguably the most critically endangered tortoise species in the world, now nearly extinct in the wild due to illegal poaching. The mitochondrial data were not well resolved geographically, whereas the microsatellite data identified four geographically discrete genetic clusters. No taxonomic changes were recommended, but three of the genetic clusters were identified as separate Management Units. The authors advised that these genetic clusters be treated as conservation units for both wild and captive management to maintain the breadth of genetic diversity present in the wild.

21:80. ***Chelonoidis carbonarius***: The alleged fossil taxon *Geochelone atlantica* † was described by López-Jurado et al. (1998) based on three bone specimens embedded in a chalky tufa-like matrix thought to represent an extinct Quaternary tortoise endemic to Sal Island, Cape Verde (see also TEWG 2015). However, Kehlmaier et al. (2021b) studied the type material using ancient DNA approaches and AMS radiocarbon dating and unexpectedly found that the type material actually matches extant specimens from Paraguay representing the disjunct southern population of *Chelonoidis carbonarius*, and that the single tortoise to which the type specimens belong was still alive from at least 1962 to 1974. They could not disentangle the history of the specimens that led to this confusion, but options are that the type material may have been misidentified as to its origin, or that the tufa-like matrix may have been fabricated.

21:81. ***Chelonoidis chilensis***: Sánchez et al. (2017) presented a phylogenetic and divergence time analysis of *C. chilensis* using mitochondrial cytochrome b sequence data from 111 individuals sampled across the known geographic range of the species. The resulting trees showed similar topologies and support values, with *C. chilensis* always recovered as a monophyletic group sister to *C. niger* and composed of two major reciprocally monophyletic and largely allopatric clades corresponding closely to the Dry Chaco and Monte ecoregions, respectively. Although these clades were well defined, they were less deeply divergent than intraspecific clades in other *Chelonoidis* species, and no recommendations were made for nomenclatural novelties nor a return to the older taxonomy of two (or three) purported Chaco taxa as species or subspecies: *C. chilensis* and *C. petersi* (and *C. donosobarrosi*), agreeing with the previous recommendations by Fritz et al. (2012a) and Sánchez et al. (2015) (see TTWG Annotations 12:30 and 17:59).

21:82. ***Chelonoidis niger***: Miller et al. (2018) analyzed single nucleotide polymorphisms (SNPs) for 117 individuals representing all 11 extant taxa of Galapagos giant tortoises (*Chelonoidis niger*). They detected genetic lineages concordant with these extant taxa, but noted significant additional structure within the polymorphic taxon *becki* from Wolf Volcano on Isabela Island, suggestive of successive independent colonizations from Santiago Island. Their SNP-based estimates of diversity and differentiation were significantly correlated with those derived from previously published nuclear microsatellite loci and mitochondrial DNA sequences, but provided higher resolution to detect individuals with admixed ancestry than previously used genetic markers.

Based on four mitochondrial and one nuclear gene, Román-Palacios and Wiens (2018) generated a time-calibrated phylogeny of all tortoises (Testudinidae), including the extinct genus *Cylindraspis*. Their analysis indicated that no tortoise species diverged less than 2 mya except for 10 of the 11 sampled taxa from the Galapagos Islands. They discussed this recent rapid diversification, but made no comment on the taxonomic status of these populations.

Poulakakis et al. (2020) analyzed mitochondrial genomes of all 11 extant and 2 recently extinct taxa of Galápagos tortoises to infer that dispersal of the ancestral form of *Chelonoidis* occurred from the two oldest Galápagos islands (San Cristóbal and Española) to Santa Cruz, Santiago, and Pinta, and was followed by colonization events from several sources within the archipelago. Using a molecular clock, they inferred that divergence within the Galápagos archipelago commenced ca. 1.5 mya.

Kehlmaier et al. (2021a) conducted an independent fossil-calibrated molecular clock analysis comparing mitochondrial genomes of all species groups of extant tortoises, including the same 13 Galápagos taxa and the recently extinct five *Cylindraspis* species from the Mascarenes, and 11 subfossil Holocene specimens of extinct *Chelonoidis* (from 700–2700 ybp) from the Bahamas archipelago. They observed that the inter-island differentiation within the Bahamas and within the Galápagos were much younger (ca. 1.5 mya and 2.0 mya, respectively) than the genetic differentiation between any other congeneric pair of tortoise species (ca. 4.0–26.8 mya). Even the divergences between two samples of *Chelonoidis carbonarius* (ca. 3.7 mya) exceeded those between the respective tortoises from the Bahamas and the Galápagos. They concluded that each archipelago harbored only one species of tortoise, and that the taxa from the Galápagos, currently regarded as distinct species, should be returned to subspecies status, as the extinct Bahamas taxa were already considered (Franz et al. 2020).

Thomson et al. (2021), in their comprehensive global turtle phylogeny using 15 nuclear loci (and no mitochondrial genomes, so completely independent from Kehlmaier et al.), also showed very shallow differentiation for five *Chelonoidis* taxa from the Galápagos, with less divergence than between any other congeneric tortoise species, except for *Kinixys belliana* and *K. spekii* (however, the latter case conflicted with earlier reported deeper genetic differentiation patterns for *Kinixys* [Kindler et al. 2012] and may have resulted from sample misidentification [H.B. Shaffer, pers. comm.]).

In view of the strong corroborating evidence from these recent comprehensive studies, as well as similar findings and recommendations from earlier work by Loire et al. (2013), Fontaine (2017), and Loire and Galtier (2017), we now agree that the *Chelonoidis* taxa from the Galápagos are probably more conservatively and appropriately considered subspecies of a single species, *C. niger,* rather than separate species as treated by Miller et al. (2018), Poulakakis et al. (2020), Caccone (2020), Gibbs et al. (2020), and others, including ourselves in most of our previous checklists (see TTWG Annotations 07:58, 09:32, 12:31, and 17:60). As a result, we have returned all these taxa to subspecific status pending further analysis. We note that full genomic analysis of all Galápagos taxa, including type specimens, may help further clarify their relationships—Kehlmaier et al. (2021a) highlighted that the mitochondrial genomes of 13 taxa from the Galápagos (data from Kehlmaier et al. 2017 and Poulakakis et al. 2020) corresponded to only 6–7 clades, whereas 15 distinct taxa (one unnamed) are currently recognized (Frazier 2020; Caccone 2020).

21:83. ***Chelonoidis niger***: Sizes provided for these taxa are maximum straightline carapace length (SCL) and maximum curved carapace length (CCL) of the largest known specimens in the Galápagos, as opposed to specimens in captivity elsewhere. These records were sourced from published field studies and the extensive unpublished database of over 35,000 measurements of tortoises by the Galapagos Conservancy, the Galapagos National Park Directorate, and the Giant Tortoise Restoration Initiative, generously shared by James P. Gibbs. In addition, summarized data from this database found in Chiari (2020), although stating that "maximum CCL" measurements were documented in Table 8.1, were actually 95th percentile sizes, while Fig. 8.6(A) provided a graphic indication of maximum recorded sizes. We additionally provide these 95th percentile sizes here.

21:84. ***Chelonoidis niger niger***: In our previous checklist (see TTWG Annotation 17:62) we commented on the just-published article by Olson and Humphrey (2017) that discussed the probable island origin of the lost type specimen of *Testudo elephantopus* Harlan 1826 as likely being Floreana (Charles Island). We do not disagree with this assessment, and have therefore now tentatively moved this taxon from its previous placement under *Chelonoidis*, sp. indet., into the synonymy of *Chelonoidis niger niger* (Quoy and Gaimard 1824b). We also note that we overlooked the clarification by Olson and Humphrey that Harlan's work was actually published in 1826 (November), not 1827; we therefore correct the date accordingly. Although we place *Testudo elephantopus* in the synonymy of *C. n. niger*, we retain it as a *nomen dubium*, as its origin is not totally certain, and the holotype is lost and not available for genetic analysis. In addition, as we noted previously, and pending genetic analysis of the holotype of *Testudo nigra* Quoy and Gaimard 1824b, we do not agree with Olson and Humphrey that the name *elephantopus* should replace *niger* as the currently accepted name for the extinct Floreana Giant Tortoise.

21:85. ***Chelonoidis niger phantasticus***: This spectacular giant tortoise, known from only a single old male specimen collected on Fernandina in 1906, had previously generally been assumed to be extinct, but was recently assessed by the TFTSG and IUCN as Critically Endangered (Possibly Extinct) (see TTWG Annotation 17:63) based on scattered signs of its possible persistent survival (Hendrickson 1965; Rhodin et al. 2017 Red List Assessment; Kiester 2019). An expedition in early 2019 to Fernandina by the Galapagos Giant Tortoise Restoration Initiative (Tapia 2019; Tapia et al. 2020) and Animal Planet host Forrest Galante (Liu 2019) led to the exciting discovery and collection of a live female tortoise from a relatively inaccessible area on the western side of Fernandina. Genomic analysis by Adalgisa Caccone at Yale University confirmed its identity in May 2021 as *C. n. phantasticus* (A. Caccone, pers. comm., unpubl. data).

21:86. ***Chelonoidis niger porteri***: We previously listed *Testudo planiceps* Gray 1854b under *Chelonoidis*, sp. indet., based on Pritchard (1996), but based on examination of its skull by Günther (1875a) it would appear to be consistent with his concept of *Testudo nigrita* Duméril and Bibron 1835, a *nomen dubium* subjective senior synonym of *Testudo porteri* Rothschild 1903. We therefore tentatively place *T. planiceps* in the synonymy of *Chelonoidis niger porteri*, pending genetic analysis of the types of *T. planiceps* and *T. nigrita*.

21:87. ***Chelonoidis niger* subsp. [?]**: Poulakakis et al. (2012) analyzed mitochondrial DNA from four specimens of *Chelonoidis* tortoise bones collected on Santa Fé (Barrington Island) by Rollo Beck in 1906, who found only appendicular bones and some subfossil eggs there, but no shells or skulls (Van Denburgh 1914). Poulakakis et al. (2012) showed that the mtDNA of these Santa Fé specimens was distinct from and sister to the clade of *abingdonii* (Pinta), *hoodensis* (Española), *chathamensis* (San Cristóbal), and *donfaustoi* (eastern Santa Cruz) (the latter still unnamed at that time). Several authors since then have referred to these bones from Santa Fé as an undescribed extinct species (Poulakakis et al. 2015; Frazier 2020; Caccone 2020).

However, the original provenance of the bones found on Santa Fé had previously been questioned by Pritchard (1996:65), who suggested that since only leg and pelvic bones were found, the specimens may have been brought in as provisions from a ship anchored in the island's easily accessible harbor and that the animals had perhaps been obtained elsewhere and butchered before being brought onto the island. On the other hand, Townsend (1925) recorded that 22 tortoises were removed from Santa Fé in 1839 and one in 1853, and Van Denburgh (1914) noted anecdotal reports of tortoises still present there in about 1876 and 1890, but none thereafter.

Two specimens of *chathamensis* were used by Poulakakis et al. (2012) as comparisons to the four Santa Fé specimens. One was the female paratype of *chathamensis* (CAS 8133) collected by Beck in 1906 from the now-extinct population from central San Cristóbal (see Pritchard 1996:59, map 5), and the second was a recent specimen of *chathamensis* from the extant population at Media Luna in northeastern San Cristóbal. There also existed a now-extinct population of tortoises in southwestern San Cristóbal (Pritchard 1996; Tapia et al. 2020). No specimens from this extinct southwestern population have been genotyped, but they could hypothetically have been distinct from central and northeastern *chathamensis* and possibly the same haplotype as the Santa Fé animals and could therefore conceivably have been the source of the limb bones brought to Santa Fé and consumed there. We list this Santa Fé taxon here as distinct, but consider it an unnamed extinct subspecies of *C. niger*, rather than a species.

21:88. ***Chersina angulata***: Spitzweg et al. (2020) sampled the mitochondrial cytochrome b gene and 14 microsatellite loci across the range of *C. angulata* in southern Africa. Both data sets suggested the existence of two lineages (a western clade, including southern Namibia and western South Africa; and an eastern clade, in southern South Africa) occupying different climatic niches. They estimated that these lineages diverged in the Pliocene (about 3.8 mya), but hybridize in areas of sympatry, suggesting conspecificity. The authors concluded that the two clades each warrant subspecific recognition; however, they cannot yet be named until type material of the synonymized taxa can be genotyped and their origins clarified. Their microsatellite analysis also identified two genetic sub-clusters (western and southwestern) within the western subspecies, which they recognized as important Management Units.

21:89. ***Chersobius* and *Homopus***: As previously noted (TTWG Annotation 17:64), Hofmeyr et al. (2017) resolved a paraphyletic *Homopus*, and resurrected the genus *Chersobius* for the five-toed species (*signatus*, *boulengeri*, and *solus*) formerly in the genus *Homopus*, and restricted the genus *Homopus* to the four-toed species (*areolatus* and *femoralis*). We tentatively accepted those changes in our previous checklist (TTWG 2017). Hofmeyr and Branch (2018) have now corroborated those genetic findings with a morphologic analysis that lends further support for this split, and we therefore now fully accept this taxonomy.

21:90. ***Cylindraspis***: Kehlmaier et al. (2019b) analyzed mitogenomics of all five extinct giant tortoise species of *Cylindraspis* from the Mascarenes in the western Indian Ocean. They found that *Cylindraspis* was monophyletic and sister to a diverse and deeply structured clade containing genera from sub-Saharan Africa (*Centrochelys, Chersina, Chersobius, Homopus, Kinixys, Psammobates,* and *Stigmochelys*), India (*Geochelone*), Madagascar (*Astrochelys* and *Pyxis*), Aldabra (*Aldabrachelys*), and South America plus the Bahamas and Galápagos (*Chelonoidis*), and these two clades were sister to the clade containing *Malacochersus, Indotestudo,* and *Testudo*. Within *Cylindraspis, C. triserrata* from Mauritius was deeply divergent from the remaining four species, resolved as two clades. *Cylindraspis inepta*, also from Mauritius, was sister to *C. indica* from Réunion, and the two species from Rodrigues, *C. vosmaeri* and *C. peltastes*, were sister taxa with a genetic divergence similar to that between *C. inepta* and *C. indica*. Their data documented that the evolutionary divergence within *Cylindraspis* is much older and deeper than the radiations within other extant and extinct giant tortoise lineages from either Aldabra and Madagascar (*Aldabrachelys*), the Bahamas (*Chelonoidis*), or the Galápagos archipelago (also *Chelonoidis*).

The extinction of *Cylindraspis* not only represents the loss of one of the five major radiations of extant Testudininae but also that of a truly ancient vertebrate lineage.

21:91. ***Cylindraspis vosmaeri***: The trinomen *Testudo indica vosmaeri* Schoepff 1795 (or as "1792" or "1793") is occasionally cited as the first use of the specific epithet *vosmaeri* (e.g., Hubrecht 1881, Vaillant 1893). However, Schoepff (1795:103) used "Testudo indica. *Vosmaeri"* as the heading for a specimen of *Testudo indica* described by Vosmaer, just as he also used the heading "Testudo indica. *Perrault*" for a specimen of *Testudo indica* described by Perrault. As such, Schoepff's supposed trinomen has no ICZN Code-compliant standing as an available name, as pointed out by Hoogmoed et al. (2010). Suckow (1798) was the first to use the trinomen in a Code-compliant manner.

21:92. ***Geochelone elegans***: Vamberger et al. (2020b) sequenced three mitochondrial genes for *G. elegans*, representing each of the three primary disjunct distribution areas from Pakistan across India to Sri Lanka. They found little variation across their samples, with three weakly to moderately supported clades that did not correspond to the geographic distribution patterns. One clade contained samples from Pakistan, southeast mainland India, and western Sri Lanka. A second clade was identified from Gujarat in northwestern India, geographically close to the Pakistan samples. The third clade included only individuals from eastern Sri Lanka, not far from samples of the second clade. The authors speculated that these unclear geographic patterns, as compared to the rangewide patterns previously reported by Gaur et al. (2006), may reflect the long-term illegal commercial trade and resultant indiscriminate release of confiscated tortoises into non-native areas.

21:93. ***Homopus areolatus***: Hofmeyr et al. (2020) examined the mitochondrial ND4 gene and the nuclear prolactin gene from samples across the range of *H. areolatus*. Although the nuclear locus exhibited no evidence of geographic structure, the mitochondrial gene revealed two well-supported genetic clades strongly associated with ecological niche divergence, one in southwestern South Africa and the other across the more central and eastern parts of southern South Africa. They estimated their divergence as the late Pliocene (>2.5 mya). Because their study was based on only a single variable marker, they recommended no taxonomic changes, but suggested the recognition of the two clades as Management Units.

21:94. ***Psammobates tentorius***: Zhao et al. (2020a, 2020b, 2020c, 2020d, 2021) investigated the rangewide phylogeography of mitochondrial and nuclear DNA and microsatellites in the three currently recognized subspecies of the *P. tentorius* species complex: *tentorius*, *trimeni*, and *verroxii*. Zhao et al. (2020a) initially identified seven mtDNA clades and inferred five to seven putative species or operational taxonomic units (OTUs), rather than just the three recognized subspecies, though the single conservative nuclear gene generated incongruent results. Their ongoing analyses demonstrated *verroxii* as currently defined to be paraphyletic, with a distinctive northwest-southeast geographic divide separated by the lower Orange River, and that most mitochondrial clades appeared to be geographically and historically climatically delimited by distributional barriers (rivers and mountain ranges). However, microsatellite analyses revealed broad admixture between some of the geographically adjacent clades, not necessarily supportive of distinct species. They made no specific nomenclatural recommendations, but in their two most recent papers (Zhao et al. 2020d, 2021) they suggested that four taxa should be considered for recognition pending further morphological studies: *tentorius*, *trimeni*, *verroxii* (southeastern clade), and *verroxii* (northwestern clade).

The *P. tentorius* species complex has a confusing and prolific nomenclatural history, with no less than 26 different nominal taxa synonymized under the three currently recognized subspecies. Genetic analysis of DNA of the pertinent type specimens of many of these taxa will need to be accomplished to help elucidate what names may need to be applied to the various clades being identified.

Within the current concept of *P. t. verroxii*, with its 14 synonymized nominal taxa, we note that the clade southeast of the lower Orange River would apparently keep the name *verroxii*. However, its restricted type locality (north of Aliwal North, north of the origin of the upper Orange River and south of the Caledon River) is beyond the southeastern distributional limit of the clade as sampled and depicted by Zhao et al. (2020d).

For the paraphyletic *verroxii* clade north of the lower Orange River, the primary candidate taxon name would appear to be *Testudo oscarboettgeri* Lindholm 1929, a justified *nomen novum* replacement name for *Testudo boettgeri* Siebenrock 1904a (a junior primary homonym of *Testudo graeca boettgeri* Mojsisovics 1889) with its type specimen from Great Namaqualand, Namibia. In addition, both *Homopus bergeri* Lindholm 1906 and *Psammobates depressa* FitzSimons 1938 appear to belong to the northern *verroxii* clade.

21:95. ***Stigmochelys pardalis***: Spitzweg et al. (2019) and Dajčman et al. (2021) analyzed population genetic and phylogeographic variation in the southern portion of the range of Leopard Tortoises using microsatellite loci and mitochondrial DNA. They found two major genetic clusters, reflecting a north-south divergence: a southern cluster distributed in southern coastal South Africa, and a northern cluster in Namibia to Mozambique. While the southern cluster showed little variation, substructuring occurred in the north. The results of both studies were concordant, but Dajčman et al. (2021) had denser sampling in the north, resulting in the identification of one additional northern subcluster as compared to Spitzweg et al. (2019). According to Dajčman et al. (2021), the northern cluster consisted of four sub-clusters: 1) northwestern South Africa to Namibia; 2) north-central South Africa; 3) northeastern South Africa; and 4) far eastern South Africa and Mozambique, with considerable admixture among the sub-clusters. The two studies did not recommend any taxonomic changes, and Spitzweg et al. (2019) concluded that Leopard Tortoises occurring in the southern part of the Western Cape province of South Africa could either be native, or had expanded their range recently, or represented short-distance transplants (possible introductions). Dajčman et al. (2021) urged the recognition of the five genetic clusters as Management Units.

21:96. ***Testudo graeca***: Schweiger and Gemel (2020) documented and corrected the confusing history regarding the type locality of the subspecies *Testudo graeca graeca* Linnaeus 1758. The original description by Linnaeus was based on an illustration, Plate 204 in Edwards (1751), and the type locality given as "Africa." Edwards had indicated that he received the illustrated specimen from Thomas Rawlings of "Santa-Cruz in West Barbary," but Houttuyn (1764), erroneously cited as Leven (1764) by Bour (1987) and TTWG (2017), recorded the origin of Edwards' depicted animal as simply "Santa Cruz in Barbarie," dropping the "West" qualifier. Strauch (1862) then erroneously interpreted "Santa Cruz" as referring to the "alte spanische Festung bei Oran in der Algérie" (on the Mediterranean coast of Algeria in the region that was previously known as Barbary), and the type locality for *T. g. graeca* has been interpreted as such ever since. However, Schweiger and Gemel (2020) documented that Edwards (1747) had previously specifically noted that Thomas Rawlings lived in "Santa Cruz, in that Part of Barbary which lies without the Streights of Gibraltar, on the Atlantic Ocean," i.e., in coastal Morocco, not Algeria, and that the town of Santa Cruz had undergone several name changes over the years and is currently known as Agadir (30.4278° N, 9.5981° W). Since the vicinity of Agadir is also the type locality for the subspecies *T. g. soussensis* Pieh 2001, Schweiger and Gemel (2020) declared *soussensis* to be a junior subjective synonym of *graeca*.

The clarification of the actual type locality of *T. g. graeca* also meant that the subspecies in northeastern Morocco to central Algeria needed a new name. Schweiger and Gemel attributed the name *T. mauritanica* Duméril and Bibron 1835 to this population formerly regarded universally as *T. g. graeca*. Duméril and Bibron had originally designated the distribution (*patrie*) of *T. mauritanica* as extremely widespread: "Mauritanie..[&]..les côtes occidentales de la mer Caspienne..[&]..les environs d'Alger." Later, Mertens and Wermuth (1955) restricted the type locality to "Umgebung von Algier" [Algeria], possibly based on the statement by Duméril and Bibron that among the specimens from which they had described their new species were many obtained from Parisian markets that had come from Algiers. However, as Mertens and Wermuth did not designate a lectotype among the many available syntypes (specimens and descriptions cited by Duméril and Bibron), their type locality designation was invalid (ICZN Art. 76.1). According to Schweiger and Gemel, no remaining syntypes of *T. mauritanica* are currently identifiable at MNHN, and they therefore designated MNHN 0.1937, labeled as coming from "Alger," as the neotype of *mauritanica*, agreeing with the previously restricted type locality of Algiers.

However, Schweiger and Gemel discounted the fact that Bour (1987) had previously validly designated Plate 204 in Edwards (1751), already the holotype of *T. graeca*, as also being the lectotype of *T. mauritanica*, as Duméril and Bibron (1835) had listed that plate in the synonymy of their

new species, qualifying it as a syntype (ICZN Art. 72.4.1). Schweiger and Gemel also overlooked the fact that both Cochran (1961) and Reynolds et al. (2017) documented that a "cotype" (= syntype) of *T. mauritanica* was obtained by the USNM in 1881 from the Jardin des Plantes in Paris (an affiliate of MNHN). As such, USNM 10980 (formerly MNHN 22, an alcoholic adult male, CL 110 mm, from Algeria) is the only known remaining physical paralectotype of *T. mauritanica.* Since both a lectotype and paralectotype of *T. mauritanica* exist, the neotype designation by Schweiger and Gemel (2020) is superfluous and invalid, and *mauritanica* remains a junior objective synonym of *graeca*. Further, the type locality of *mauritanica* must agree with the origin of its lectotype (ICZN Art. 76.2), so it is hereby restricted to Agadir, Morocco.

Other possible names to consider for the subspecies in northeastern Morocco to central Algeria are *T. pusilla* Linnaeus 1758 and *T. whitei* Bennett *in* White 1836. *Testudo pusilla* has a confusing history, in that Linnaeus (1758) first listed its description as sourced only from Worm (1655), Grew (1681), and Ray (1693), but then later (Linnaeus 1766) also added Edwards (1751:pl.204) as a source for its description, removing that source from his prior description of *T. graeca* (Linnaeus 1758). Whether Linnaeus examined any specimens of *T. pusilla* is unknown, but Linné and Thunberg (1780) listed one specimen as present in the Linnaean collections, but it was never recorded again and is presumably lost. *Testudo pusilla* is appropriately listed as a junior synonym of *T. graeca*, and its type locality, previously restricted by Strauch (1862) to "Santa-Cruz (alte spanische Festung bei Oran in der Algérie)" is hereby corrected to Agadir, Morocco, as for *T. graeca*.

Testudo whitei also has an uncertain history, with an unknown provenance of the holotype obtained from the pet trade and kept for a long time in England (Bennett 1836; White 1836). Gray (1873j) designated its origin as "S. Europe," but Highfield and Martin (1989b) concluded that the holotype resembled animals from Algeria and restricted the type locality to "Algier and its environs, Algeria." As such, *T. g. whitei* becomes the new apparent name for the *T. graeca* subspecies occurring from northeast Morocco through central Algeria, pending genetic evaluation of the holotype to confirm or reject whether it matches the mitochondrial haplotype for that population as identified by Fritz et al. (2009c).

Despite these irregularities and prior confusion concerning the type specimens and type localities for *T. graeca*, *T. mauritanica*, and *T. whitei*, what Schweiger and Gemel (2020) have demonstrated is that the subspecies of *T. graeca* occurring from northeastern Morocco to central Algeria should no longer be called *T. g. graeca*. It is now instead referred to *T. g. whitei* (instead of *T. g. mauritanica*), and *T. g. graeca* replaces *T. g. soussensis* as the valid name for the subspecies in southwestern Morocco.

21:97. ***Testudo graeca***: Turkozan et al. (2018) examined variation in two mitochondrial genes (cytochrome b and ND4) for populations of the four subspecies of *T. graeca* in Turkey (*armeniaca, buxtoni, ibera, terrestris*), and their phylogenetic analysis generally supported the monophyly of each of them, although the support values were not high. In addition, they reported that *ibera* and *terrestris*, *armeniaca* and *ibera*, and *armeniaca* and *buxtoni* are found syntopically in some locations, where they hybridize. They also examined morphometric variation among the four subspecies and found that only *armeniaca* was morphometrically distinct from the others. However, they concluded that the available molecular and morphological data were insufficient to warrant the recognition of subspecies in Turkey. Until broader sampling of the genome is done across the entire *T. graeca* complex, we continue to recognize these taxa as subspecies of *T. graeca*.

21:98. ***Testudo graeca armeniaca, T. g. buxtoni,* and *T. g. ibera***: In a paper describing the ecomorphology of *T. gracea* subspecies in Armenia, northwestern Iran, and far eastern Turkey, where the three taxa *armeniaca, buxtoni,* and *ibera* occur and hybridize, Arakelyan et al. (2018) found that ecological habitat conditions of tortoises were correlated with significant external morphological plasticity. Their results were also concordant with previous findings of admixed mitochondrial and microsatellite loci and the existence of a contact zone with extensive gene flow between the three subspecies occurring in this area (Mashkaryan et al. 2013).

21:99. ***Testudo hermanni***: Biello et al. (2021) examined variation across seven microsatellite loci in *T. hermanni* from 28 regions across its range. Their best supported model confirmed the divergence between the two recognized subspecies *hermanni* and *boettgeri*. However, a secondary well-supported model also emerged, subdividing *T. h. hermanni* into four genetic subclusters, and *T. h. boettgeri* into two. Five of these subclusters had previously been identified by Perez et al. (2014) and the sixth was from a previously unsampled population in Calabria at the southwestern tip of continental Italy (see also TTWG Annotations 14:36 and 17:74). Since the purpose of their paper was for conservation and forensics (e.g., determining the provenance of trade-confiscated animals), they did not discuss taxonomy. Although the subclusters qualify as Management Units, we concur that they do not warrant taxonomic designation.

Djurakic and Milankov (2019) examined variation in the partial mitochondrial cytochrome b gene (518 bp) across the range of *T. hermanni boettgeri*. They identified three primary genetic clusters, one corresponding to *T. h. hermanni* (see also TTWG Annotation 17:74) and two within *T. h. boettgeri*: 1) Croatia, Bosnia and Herzegovina, and northern Greece, and 2) Romania, Serbia, Bulgaria, North Macedonia, and most of Greece). They also compared carapace morphometrics between a population in each of the two *boettgeri* genetic clusters and found subtle but statistically significant differences, but made no taxonomic recommendations.

In a follow-up study, Djurakic and Milankov (2020) undertook another combined morphological and molecular analysis of variation across Greek populations of *T. h. boettgeri*. Their analysis of a small (266 bp) portion of the mitochondrial 12s rRNA locus revealed three haplotypes differing in only one to three bases from three regions: 1) northern mainland Greece, 2) southern mainland Greece, and 3) the Peloponnese peninsula. Their analysis of plastral morphometry showed no difference between the two mainland groups, but both of them differed significantly from tortoises in the Peloponnese. Again, the authors made no taxonomic or management recommendations.

21:100. ***Testudo hermanni hermanni***: The fossil species *Testudo globosa* † Portis 1890, with its junior synonyms, *Testudo oriens* † Portis 1890 and *Testudo seminota* † Portis 1890, were included in the synonymy of *T. h. hermanni* in our previous checklists based on Lapparent de Broin et al. (2006b, 2006c) (see TTWG Annotation 14:37), but Corsini et al. (2014) considered *T. globosa* distinct and separate, so we have now removed it and its two synonyms, and will include them in our planned revised edition of TEWG (2015).

21:101. ***Lissemys punctata***: As a follow-up to the study by Praschag et al. (2011; see TTWG Annotation 11:14) of mtDNA sequence variation in *L. punctata*, Bhaskar and Mohindra (2019) undertook additional geographic sampling, but examined only one (cytochrome b) of the three genes studied by Praschag et al. The results of both studies were concordant, supporting the recognition of three subspecies (*punctata*, *andersoni*, and *vittata*). However, both studies identified two well-supported subclades within *L. p. vittata*: one apparently restricted to the states of Goa and Karnataka in western central India, and the other ranging widely across the rest of central India. Although Praschag et al. (2011) did not discuss this divergence, presumably because they only had two samples from the restricted area, Bhaskar and Mohindra, with 13 samples, suggested that *vittata* may represent two subspecies, pending further study.

21:102. ***Pelodiscus axenaria* and *Pelodiscus huangshanensis***: Yu et al. (2019) analyzed the full mitochondrial genome of a single softshell turtle from Huangshan, Anhui, that they identified as *P. axenaria*, although occurring far from its previously defined range. Subsequently, Gong et al. (2021) described a new small-sized *Pelodiscus* species from Huangshan, southern Anhui province, China, based on molecular and morphological evidence, and identified the specimen that Yu and colleagues had examined as belonging to their new species. Based on phylogenetic analyses of the mitochondrial cytochrome b and ND4 genes, *P. huangshanensis* is most closely related to *P. axenaria* and another unspecified lineage previously characterized by Gong et al. (2018). *Pelodiscus huangshanensis* was described as having a very limited distribution and a narrow ecological niche confined to clear rivers with fine sand in the slightly upland source area of the Xian'anjiang River. No genetic comparisons were made with *P. sinensis* specimens from near the type locality of *P. huangshanensis*, and the authors' statements about the maximum carapace length size of the new species varied from 84.0 to 101.6 to 115.6 mm, suggesting some confusion as to its size, although it is apparently the smallest known softshell turtle. Its habitat in higher gradient upland sandy-bottomed clear rivers apparently differentiates it from the larger *P. sinensis* which tends to inhabit a variety of nearby habitats, including more lowland rivers

with muddy bottoms. According to the authors, the known population of *P. huangshanensis* is highly threatened due to habitat destruction and hunting pressure. We tentatively accept this new species pending more genetic analysis and improved definition of its distribution and whether or not it occurs sympatrically with *P. sinensis*.

21:103. ***Pelodiscus parviformis* and *Pelodiscus variegatus***: Gong et al. (2018) and Farkas et al. (2019) analyzed genetic and morphological characteristics of all *Pelodiscus* species and concluded that *P. parviformis*, previously considered to have two disjunct populations, one in central China and one in Vietnam, was a composite of two species. They documented distinct mitochondrial and nuclear DNA differences as well as diagnostic morphological and color-pattern differences between these populations, and Farkas et al. (2019) described the Vietnamese one as a new species, *Pelodiscus variegatus*, noting that it also occurred on Hainan Island in China. We agree with this assessment.

21:104. ***Pelodiscus sinensis***: Xiong et al. (2019) analyzed mtDNA variation in this wide-ranging species using the partial cytochrome b gene (922 bp). They identified four separate genetic lineages (Yangtze River, Xijiang River, Yellow River, and Taiwan) and suggested that the species probably originated in the Yangtze River basin, with specimens from Japan apparently having been introduced from the Yellow River lineage. Further, they noted evidence of lineage mixing between some specimens of *P. sinensis*, *P. parviformis*, and *P. maackii*, and suggested that the latter two taxa should be considered subspecies of *P. sinensis*. We await further genetic analysis before following this recommendation, and we also note that it is unclear whether the analyzed turtles were native or farmed animals and we abstain from any conclusions on the distribution of these species based on these data.

21:105. ***Chelonoidis alburyorum***: In recent years several studies have contributed significantly to a better understanding of the diversity of the extinct Holocene tortoises from the Bahamas archipelago (Franz and Franz 2009; Kehlmaier et al. 2017, 2021; Albury et al. 2018; Franz et al. 2020; Steadman et al. 2020). All native Bahamian tortoises vanished shortly after the arrival of prehistoric humans 900–700 years ago (Steadman et al. 2020). These tortoises were originally widely distributed in the archipelago and were large-bodied to giant-sized (Franz et al. 2020; Kehlmaier et al. 2021a). Tortoises on the individual islands differed in morphological characters so much that Steadman et al. (2020) suggested that up to seven distinct species could have existed there. Franz and Franz (2009) initially described a single species from the Bahamas, *C. alburyorum*. Later, based on morphological evidence, Franz et al. (2020) described two additional subspecies: *C. alburyorum keegani* and *C. a. sementis*. Subsequently, a comparison of near-complete mitochondrial genomes of 11 subfossil specimens from six islands by Kehlmaier et al. (2021a) revealed only two well-supported but weakly divergent clades and, in comparison to the divergences observed in other testudinids, suggested conspecificity of the specimens. Their results also indicated the possible recognition of only two subspecies of *C. alburyorum* rather than three.

A fossil-calibrated molecular clock indicated that the two Bahamian clades separated ca. 1.5 mya, which is too early for in-situ divergence, since high interglacial sea levels made the Bahamas only permanently hospitable for land vertebrates about 400,000 ybp. This suggests that the Bahamas tortoises colonized the islands twice from other landmasses, most likely the Greater Antilles (Kehlmaier et al. 2021a).

One of the two clades recovered by Kehlmaier et al. (2021a) included sequences from one specimen from Great Abaco and two from Crooked Island; the other clade contained sequences corresponding to the remaining tortoise material from Great Abaco and Crooked Island, as well as from Eleuthera, Mayaguana, Middle Caicos, and Grand Turk. Archeological evidence and radiocarbon dates of the studied material suggested that the occurrence of tortoises of both clades on Great Abaco and Crooked Island was the result of prehistoric human transport of tortoises between islands (Kehlmaier et al. 2021a).

DISTRIBUTION UPDATES

Specific distribution updates were recorded in the 2011, 2012, and 2014 checklists, but were not continued in the 2017 or current checklists, except as reflected in the revised maps and distributional information under each taxonomic entry.

Distribution Updates 2011

Dermatemys mawii: Honduras (?) deleted from the range as per CBFTT account (Vogt et al. 2011).

Kinosternon scorpioides albogulare: Population in San Andrés, Colombia, indicated as possible prehistoric or modern introduction, as per CBFTT accounts (Berry and Iverson 2011, Forero-Medina and Castaño-Mora 2011).

Chrysemys dorsalis: Apparently established in Florida (Krysko et al. 2011).

Graptemys pseudogeographica: Apparently established in Florida (Krysko et al. 2011).

Malaclemys terrapin and *M. t. centrata*: Bermuda added to native range (Parham et al. 2008).

Pseudemys nelsoni: Introduced to Tortola, British Virgin Islands (Owen et al. 2005).

Trachemys scripta scripta: Apparently established in Florida (Krysko et al. 2011).

Emys orbicularis orbicularis: Spain deleted from range; Spanish populations are attributable to subspecies *E. o. fritzjurgenobsti* and *E. o. galloitalica*, but not to the nominate subspecies.

Terrapene nelsoni: Chihuahua, Mexico (*T. n. klauberi*) and Jalisco, Mexico (*T. n. nelsoni*) added to the range, as per CBFTT account (Buskirk and Ponce-Campos 2011).

Batagur kachuga: Occurrence in Nepal confirmed.

Cuora bourreti: Occurrence in Laos indicated as uncertain.

Cuora picturata: Occurrence in central Vietnam confirmed (Ly et al. 2011).

Cuora zhoui: Vietnam added as uncertain occurrence.

Geoemyda spengleri: Laos added to the range (Stuart et al. 2011).

Mauremys caspica caspica: Israel, Jordan and Lebanon deleted from the range; populations there attributable to *M. rivulata*. Turkmenistan added to the range.

Mauremys caspica siebenrocki: Turkmenistan deleted from the range; population there attributable to *M. c. caspica*.

Sacalia bealei: Occurrence in Guangxi, China, indicated as uncertain.

Aldabrachelys/Dipsochelys gigantea/dussumieri hololissa: Cousine Island added to extirpated range, Round and Cousine islands added to introduced range, as per CBFTT account (Gerlach 2011b).

Stigmochelys pardalis: Sudan deleted from the range, and the new nation of South Sudan added to the range.

Pelodiscus parviformis: Vietnam added to the range.

Acanthochelys pallidipectoris: Mendoza, Argentina, corrected from native range to introduced, as per CBFTT account (Vinke et al. 2011).

South Sudan: This newly independent nation has seven taxa of freshwater turtles and tortoises: *Kinixys belliana belliana, Stigmochelys pardalis, Cyclanorbis elegans, Cyclanorbis senegalensis, Trionyx triunguis, Pelomedusa subrufa,* and *Pelusios adansonii*.

Distribution Updates 2012

Cuora bourreti: Occurrence in Laos was confirmed (Stuart et al. 2011), and possible occurrence in Cambodia was deleted from the checklist following consultation with range-state biologists.

Mauremys reevesii: Based on analysis of mitochondrial DNA, Suzuki et al. (2011) indicated that Japanese populations of *M. reevesii* were possibly derived from multiple historical introductions from nearby countries, and thus questioned its traditional status as a presumed native species. They noted the ability of *M. reevesii* to hybridize with native *M. japonica* as a threat, but also recognized these populations to be valuable in the context of depleted populations elsewhere in its range.

Kinixys: Ranges adjusted according to species ranges outlined by Kindler et al. (2012).

Nilssonia formosa: Liebing et al. (2012) referred to a record of *N. formosa* from Shuangbai (Yunnan, China), and photographs of a specimen from the Lancang River [= Mekong] in the Xishuanbanna region of Yunnan, as suggesting that the species has crossed the watershed divide into the Mekong River basin of Yunnan, China. However, given the substantial documented

trade volumes of live turtles from Myanmar into Yunnan and onwards, and the propensity of turtles to escape or be intentionally released by humans, combined with the great biogeographical barriers (despite their very close proximity) between the Salween, Mekong, and Yangtze, we consider it doubtful that these records represent natural occurrences.

Mesoclemmys heliostemma: Additional occurrences across the Amazon basin of Brazil were reported by Molina et al. (2012).

Italy (Sardinia): Vamberger et al. (2011) compared mitochondrial and nuclear DNA of *Testudo graeca* from Sardinia with that of *T. graeca* from North Africa, and concluded that the near-absence of differentiation from other *graeca* populations, and reduced variation within the Sardinian population, indicated prehistoric introduction into Sardinia by humans. They also reviewed recent studies of the other non-marine turtle species occurring in Sardinia and concluded that Sardinia's populations of *Testudo hermanni hermanni*, *T. marginata*, and *Emys orbicularis galloitalica* likely each represented prehistoric or early historic human introductions.

Latvia: Pupins and Pupina (2011) recorded introduced populations or individuals of *Trachemys scripta elegans*, *T. s. troostii*, *Mauremys caspica*, *M. rivulata*, *Testudo horsfieldii*, and *Pelodiscus sinensis* in Latvia. However, they did not document successful reproduction in the wild, and it remains uncertain whether these records represent established populations or isolated individuals.

South Korea: Chang et al. (2012) noted that the native softshells in South Korea are attributable to *Pelodiscus maackii*, and reported *P. sinensis*, *Trachemys scripta elegans*, *T. s. scripta*, *Pseudemys rubriventris*, and *Mauremys* [*Ocadia*] *sinensis* as introduced into the wild.

Distribution Updates 2014

Chelydra acutirostris: Páez et al. (2012) listed the occurrence of this species in the Colombian Departments of Caldas and Quindio, but did not list records from Atlántico, Bolivar, Magdalena, or Sucre.

Trachemys c. or *o. callirostris*: Páez et al. (2012) reported the occurrence of this taxon in the Colombian Department of Cundinamarca.

Trachemys d. decussata: Parham et al. (2013) documented the occurrence of this taxon in northwestern Jamaica, including hybridization with *T. terrapen*; whether the occurrence is native or introduced remains unknown.

Trachemys emolli or *grayi emolli*: McCranie et al. (2013) documented the occurrence of this taxon in Honduras, while Ibarra Portillo et al. (2009) documented its occurrence in eastern El Salvador.

Cuora amboinensis: Wangyal et al. (2012) reported this and four other species from southern Bhutan, the first turtles reported from that country.

Cuora mouhotii: Rahman (2012) reported the occurrence of this species in the southern Chittagong Hill Tracts of Bangladesh, Wangyal et al. (2012) reported it from southern Bhutan, and Ly et al. (2013) extended its range in southern Vietnam.

Cyclemys gemeli: Wangyal et al. (2012) reported this species from southern Bhutan.

Melanochelys tricarinata: Wangyal et al. (2012) reported this species from southern Bhutan.

Aldabrachelys gigantea: The historic and present distribution of native and introduced populations of the various morphotypes or subspecies (*gigantea*, *arnoldi*, and *hololissa*) of giant tortoises in the Seychelles (including all granitic and coralline islands) has been analyzed in detail and updated by Gerlach et al. (2013).

Centrochelys sulcata: Participants at the IUCN/TFTSG Sub-Saharan African Red List workshop in 2013 noted that *C. sulcata* occurs in Benin, Cameroon, and Togo, and may possibly occur in Djibouti, Somalia, Saudi Arabia, and Yemen. Its presence in Yemen and Saudi Arabia was also previously noted by Gasperetti et al. (1993).

Indotestudo elongata: Wangyal et al. (2012) reported this species from southern Bhutan.

Kinixys erosa: Participants at the Sub-Saharan African Red List workshop considered that *Kinixys erosa* certainly occurs in Benin and Togo, but is absent from Burkina Faso.

Kinixys homeana: In their CBFTT species account, Luiselli and Diagne (2013) noted that *K. homeana* occurs in the Central African Republic. They questioned its occurrence in Gabon and noted that it does not occur in Congo (ROC) and that old historical records from the eastern Congo (DRC) were likely based on misidentified *K. erosa*. These historical records need further evaluation.

Kinixys nogueyi: Participants at the Sub-Saharan African Red List workshop considered the distribution of *Kinixys nogueyi* to include the Central African Republic, but that the species does not range as far south as Equatorial Guinea or Gabon, and that records from Mauritania are likely historic, but that the species no longer occurs there.

Cyclanorbis senegalensis: Participants at the Sub-Saharan African Red List workshop considered that the occurrence of *C. senegalensis* is uncertain in Cameroon, Central African Republic, and Liberia, and that the species is likely extirpated from Mauritania.

Rafetus swinhoei: Wang et al. (2013) extended and defined the known recent historic range of this Critically Endangered species in the upper Red River of southern Yunnan, China.

Mesoclemmys dahli: Páez et al. (2012) and Forero-Medina et al. (2013, CBFTT account) documented the occurrence of *M. dahli* in the Colombian Department of Magdalena.

Mesoclemmys gibba: Páez et al. (2012) recorded the occurrence of *M. gibba* in the Colombian Departments of Arauca and Guaviare.

Mesoclemmys perplexa: Campos et al. (2011) recorded the occurrence of *M. perplexa* in the Brazilian State of Goiás.

Mesoclemmys vanderhaegei: Vinke et al. (2013) reviewed the distribution of *M. vanderhaegei* and concluded that there are no confirmed records for Bolivia.

Platemys platycephala: Páez et al. (2012) recorded the occurrence of *P. platycephala* in the Colombian Departments of Guainía, Guaviare, Meta, and Vichada.

Peltocephalus dumerilianus: Páez et al. (2012) reported *P. dumerilianus* to inhabit the Colombian Department of Guaviare.

Pelusios bechuanicus: In earlier versions of this checklist, we included Congo (DRC) as part of the range of *P. bechuanicus*. However, we have been unable to verify this occurrence, and consider that this was based on old literature records of *P. upembae*, which was originally described as a subspecies of *P. bechuanicus*.

Pelusios castaneus: Stuckas et al. (2013) questioned earlier records of occurrence of *P. castaneus* on Cape Verde and suggested that our recording of that presence on our previous checklists was outdated. We have investigated this further and agree with them. Although Boulenger (1906b) documented the collection of a specimen of "*Sternothaerus derbianus*" (= *P. castaneus*) from a "small island in Praja Bay, S. Jago" (= Santiago), Chevalier (1935) noted that the specimen was most likely introduced from West Africa and that no Caboverdians were aware of any freshwater turtles in the islands. However, Boulenger's record (mapped among others by Iverson 1992) led to the assumption for a long time that the species occurred in the islands; but surveys of the local herpetofauna have failed to record its presence (Schleich 1982, 1987, 1996b; Vasconcelos et al. 2013), even as an introduced population. The small island where it was originally collected housed a prison where turtles had evidently been released into a small pond at some point in the past. We therefore remove Cape Verde from the distribution of *P. castaneus*.

For this same species, we also question whether it occurs natively on São Tomé. Although it has been recorded from there, and specimens have been collected and genetically analyzed (Stuckas et al. 2013), the species does not occur on either nearby Príncipe or the other volcanic oceanic islands in the same archipelago (Manaças 1956; Jones 1994), nor is there any record of the species occurring on the nearby continental island of Bioko (Equatorial Guinea) off the coast of Cameroon. São Tomé was first settled by the Portuguese, who brought African slaves to the island, so it appears most likely that West African *P. castaneus* were introduced to São Tomé in conjunction with the slave trade. In fact, the genetic analysis by Stuckas et al. (2013) demonstrated that their São Tomé specimen was essentially indistinguishable from an Ivory Coast specimen, lending further strength to this theory.

Podocnemis erythrocephala: Páez et al. (2012) reported that *P. erythrocephala* occurs in the Colombian Department of Guaviare, and perhaps in Vichada.

Podocnemis lewyana: Páez et al. (2012) reported that the range of *P. lewyana* extends into the Colombian Department of Tolima.

Podocnemis sextuberculata: Páez et al. (2012) reported that *P. sextuberculata* occurs in the Colombian Departments of Caquetá and Putomayo.

Podocnemis vogli: Páez et al. (2012) reported that *P. vogli* occurs in the Colombian Department of Guaviare, but did not indicate occurrence in Boyacá.

LITERATURE CITED
(Divided into Three Separate Sections)

1. General Bibliographic Citations
2. *Chelonian Research Monographs*
Conservation Biology of Freshwater Turtles and Tortoises
(CBFTT Checklists and CBFTT Species Accounts)
3. IUCN Red List and Green List Assessments

Acuña-Mesén, R.A. 1993. Las Tortugas Continentales de Costa Rica. San José, Costa Rica: ICER, 58 pp.

Adler, K. 1962. A new name for a Chinese turtle, genus *Clemmys*. Natural History Bulletin of the Siam Society 20(2):135.

Adler, K. (Ed.). 1989. Contributions to the History of Herpetology. Contributions to Herpetology No. 5, 202 pp.

Adler, K. (Ed.). 2007. Contributions to the History of Herpetology. Volume 2. Society for the Study of Amphibians and Reptiles, 389 pp.

Adler, K. (Ed.). 2012. Contributions to the History of Herpetology. Volume 3. Society for the Study of Amphibians and Reptiles, 564 pp.

Adler, K. 2016. Theodore Cantor (1809–1860): pioneer scientific herpetologist in China. In: Cantor, T. Zoology of Chusan. Society for the Study of Amphibians and Reptiles, Facsimile Reprint, pp. v–xi.

Agarwal, A.K. 1987. Observations on the sexual dimorphism in Indian freshwater tortoise *Lissemys punctata punctata*. Geobios (Jodhpur) 14(6):277–280.

Agarwal, A.K., Wadhawan, B.S., and Lavania, R.K. 1986. Sexual dimorphism in Indian fresh water turtle *Kachuga tecta* (Gray). Bionature 6(2):94–97.

Agassiz, L. 1847 [“1846”]. Nomenclatoris Zoologici Index Universalis. Solothurn: Jent and Gassmann, 393 pp.

Agassiz, L. 1857a. Contributions to the Natural History of the United States of America. First Monograph. Vol. I. Part I. Essay on Classification. Part II. North American Testudinata. Boston: Little, Brown and Co., Vol. I, pp. 1–452.

Agassiz, L. 1857b. Contributions to the Natural History of the United States of America. First Monograph. Vol. II. Part III. Embryology of the Turtle. Boston: Little, Brown and Co., Vol. II, pp. 453–643.

Agha, M., Ennen, J.R., Nowakowski, A.J., Lovich, J.E., Sweat, S.C., and Todd, B.D. 2018. Macroecological patterns of sexual size dimorphism in turtles of the world. Journal of Evolutionary Biology 31(3):336–345.

Agostini, M.A.P., Batista, E.O., Dias Junior, W.R., Vogt, R.C., and Portelinha, T.C.G. 2023. New record of Spot-Legged Turtle, *Rhinoclemmys punctularia* (Daudin, 1801) (Reptilia, Testudines, Geoemydidae), from an Amazon ecotonal zone in the Tocantins–Araguaia river basin, Brazil. Check List 19(5):605–610.

Ahl, E. 1932. Beschreibung einer neuen Schildkröte aus Australien. Sitzungsberichte der Gesellschaft Naturforschender Freunde zu Berlin 1932(1/3):127–129.

Ahmed, M.F. and Das, A. 2009. Tortoises and freshwater turtles of Kaziranga National Park, Assam - diversity, distribution, conservation status. In: Vasudevan, K. (Ed.). Freshwater Turtles and Tortoises of India. ENVIS Bulletin: Wildlife and Protected Areas 12(1):49–51.

Ahsan, M.F. and Saeed, M.A. 1989. The Bostami turtle, *Trionyx nigricans* Anderson: population status, distribution, historical background and length-weight relationship. Journal of the Bombay Natural History Society 86:1–6.

Ahsan, M.F., Haque, M.N., and Fugler, C.E. 1991. Observations on *Aspideretes nigricans*, a semi-domesticated endemic turtle from eastern Bangladesh. Amphibia-Reptilia 12(2):131–136.

Akani, G.C., Eniang, E.A., Amadi, N., Dendi, D., Hema, E.M., Diagne, T., Ségniagbeto, G.H., Di Vittorio, M., Gbewaa, S.B., Pauwels, O.S.G., Chirio, L., and Luiselli, L. 2018. Macrohabitat and microhabitat usage by two softshell turtles (*Trionyx triunguis* and *Cyclanorbis senegalensis*) in west and central Africa. Herpetological Conservation and Biology 13(3):642–651.

Akçakaya, H.R., Bennett, E.L., Brooks, T.M., Grace, M.K., Heath, A., Hedges, S., Hilton-Taylor, C., Hoffmann, M., Keith, D.A., Long, B., Mallon, D.P., Meijaard, E., Milner-Gulland, E.J., Rodrigues, A.S.L., Rodriguez, J.P., Stephenson, P.J., Stuart, S.N., and Young, R.P. 2018. Quantifying species recovery and conservation success to develop an IUCN Green List of Species. Conservation Biology 32(5):1128–1138.

Alacs, E.A. 2008. Forensics, phylogeography and population genetics: a case study using the Australasian snake-necked turtle, *Chelodina rugosa*. Ph.D. Thesis, University of Canberra, Australia.

Albury, N.A., Franz, R., Rimoli, R., Lehman, P., and Rosenberger, A.L. 2018. Fossil land tortoises (Testudines: Testudinidae) from the Dominican Republic, West Indies, with a description of a new species. American Museum Novitates 3904:1–28.

Aldrovandi, U. 1637. De Quadrupedibus Digitatis Viviparis Libri Tres et de Quadrupedibus Digitatis Oviparis Libri Duo. Bononiae: Nicolai Tebaldini, 718 pp.

Alqahtani, A.R.M. 2017. Revision of geographical distribution of *Pelomedusa subrufa* in Saudi Arabia. King Khalid University Journal of Basic and Applied Sciences 3(2):26–28.

Alvarez del Toro, M. 1972. Los Reptiles de Chiapas, 2nd. Ed. Tuxtla Gutierrez, Mexico: Gobierno del Estado de Chiapas, 178 pp.

Álvarez-Varas, R., Rojas-Hernández, N., Heidemeyer, M., Riginos, C., Benítez, H.A., Araya-Donoso, R., Reséndiz, E., Lara-Uc, M., Godoy, D.A., Muñoz-Pérez, J.P., Alarcón-Ruales, D.E., Alfaro-Shigueto, J., Ortiz-Alvarez, C., Mangel, J.C., Vianna, J.A., and Véliz, D. 2021. Green, yellow or black? Genetic differentiation and adaptation signatures in a highly migratory marine turtle. Proceedings of the Royal Society B 288:20210754:1–10.

Ameghino, F. 1882. Catálogo explicativo de las colecciones de Antropología, prehistoria y de paleontología de Florentino Ameghino. Buenos Aires: Catálogo de la Sección de la Provincia de Buenos Aires en la Exposición Continental Sud Americana, Anexo A:35–42.

Anderson, C. 1925. Notes on the extinct chelonian *Meiolania*, with a record of a new occurrence. Records of the Australian Museum 14:223–242.

Anderson, J. 1871. Note on *Testudo phayrei*, Blyth. Proceedings of the Zoological Society of London 1871:425–428.

Anderson, J. 1872. On *Manouria* and *Scapia*, two genera of land-tortoises. Proceedings of the Zoological Society of London 1872:132–144.

Anderson, J. 1875. Description of some new Asiatic mammals and Chelonia. Annals and Magazine of Natural History (4)16:282–285.

Anderson, J. 1876. Note on the plastron of the Gangetic mud-turtle (*Emyda dura* of Buchanan-Hamilton). Journal of the Proceedings of the Linnean Society of Zoology 12:514–516.

Anderson, J. 1879 [“1878”]. Anatomical and Zoological Researches, Comprising an Account of the Zoological Results of the Two Expeditions to Western Yunnan in 1868 and 1875. London: Bernard Quaritch, Vol. I, 985 pp., Vol. II, 29 pp. + pls.

Andersson, L.G. 1900. Catalogue of Linnean type-specimens of Linnaeus’s Reptilia in the Royal Museum in Stockholm. Bihang till Kongl. Svenska Vetenskaps-Akademiens Handlingar 26(4:1):1–29.

Angielczyk, K.D. and Feldman, C.R. 2013. Are diminutive turtles miniaturized? The ontogeny of plastron shape in emydine turtles. Biological Journal of the Linnean Society 108:727–755.

Angielczyk, K.D., Burroughs, R.W., and Feldman, C.R. 2015. Do turtles follow the rules? Latitudinal gradients in species richness, body size, and geographic range area of the world’s turtles. Journal of Experimental Zoology 324B:270–294.

Annandale, N. 1906. *Testudo baluchiorum*, a new species. Journal and Proceedings of the Asiatic Society of Bengal n.s. 2(3):75–76.

Annandale, N. 1912a. The Indian mud–turtles (Trionychidae). Records of the Indian Museum 7(2):151–180.

Annandale, N. 1912b. The aquatic chelonia of the Mahanaddi and its tributaries. Records of the Indian Museum 7(3):261–266.

Annandale, N. 1913. The tortoises of Chota Nagpur. Records of the Indian Museum 9(5):63–78.

Annandale, N. 1915a. Notes on some Indian Chelonia. Records of the Indian Museum 11(11):189–195.

Annandale, N. 1915b. Herpetological notes and descriptions. Records of the Indian Museum 11(19):341–347.

Annandale, N. 1918. Chelonia and Batrachia of the Inlé Lake. Records of the Indian Museum 14:67–69.

Ansorge, H., Fritz, U., Terbish, K., and Shar, S. 2012. “*Agrionemys kazachstanica terbishi*” or the two-faced Mongolian steppe tortoise. Exploration into the Biological Resources of Mongolia 12:213–218.

Antenbrink-Vetter, S. and Vetter, H. 1998. Neuer Name für altbekannte Schildkröten. Schildkröten II/98:3–5.

Aoki, R. 1977. The occurrence of a short-necked chelid in the Palau Islands. Japanese Journal of Herpetology 7(2):32–33. [in Japanese]

Apodaca, J.J., Krohn, A.R., Collins, L., Godwin, J.C., Pearson, L., and Walde, A.D. 2023. Population structure and genetic differentiation in extant Alligator Snapping Turtles (genus *Macrochelys*) with implications for taxonomy and

conservation. Southeastern Naturalist 22(Special Issue 12):1–24.

Arakelyan, M., Türkozan, O., Hezaveh, N., and Parham, J.F. 2018. Ecomorphology of tortoises (*Testudo graeca* complex) from the Araks River valley. Russian Journal of Herpetology 25(4):245–252.

Arambourg, C. 1947. Contribution à l'étude géologique et paléontologique du bassin du lac Rodolphe et de la basse vallée de l'Omo. In: Arambourg, C. (Ed.). Mission scientifique de l'Omo. Vol. 1: Géologie et Anthropologie. Paris: Editions du Muséum, pp. 231–562.

Ardjima, L., Hema, E.M., Konate, S., Sirima, D., Kabre, B.G., Petrozzi, F., Fa, J.E., and Luiselli, L. 2020. Unleashing the potential of local captive populations for conservation – the case study of the African spurred tortoise. Acta Oecologica 105: doi.org/10.1016/j.actao.2020.103581.

Arnold, E.N. 1979. Indian Ocean giant tortoises: their systematics and island adaptations. Philosophical Transactions of the Royal Society of London 286B:127–145.

Arteaga, A. 2022. Brown Wood-Turtle (*Rhinoclemmys annulata*). In: Arteaga, A., Bustamante, L., Vieira, J., and Guayasamin, J.M. (Eds.). Reptiles of Ecuador: Life in the Middle of the World. doi: 10.47051/CNDL4838, www.reptilesofecuador.com, 10 pp.

Artner, H. 2003. Nomenklatur aktuell. Die rezenten Schildkrötenarten der Erde. Emys 10(6):iv–xxxviii.

Aryal, P.C., Dhamala, M.K., Bhurtel, B.P., Suwal, M.K., and Rijal, B. 2010. Turtles of Nepal: A Field Guide for Species Accounts and Distribution. Kathmandu, Nepal: Environmental Graduates in Himalaya (EGH), Resources Himalaya Foundation and Companions for Amphibians and Reptiles of Nepal (CARON), 33 pp.

Assis, C., Valadão, R.M., Teixeira De Mendonça, S.H.S., Peçanha, E.L.S., Costa, H., Novaes, C.M., Fonseca De Barros, T., Rodrigues, L.S., Gasparini, J.L., and Feio, R. 2024. Extensive sampling and citizen science expand the distribution of the threatened freshwater turtle *Ranacephala hogei* (Mertens, 1967). Anais da Academia Brasileira de Ciências 96(Suppl. 1):e20240484, 18 pp.

Attum, O., Baha el Din, S.M., Carranza, S., Earley, R., Arnold, E.N., and Kingsbury, B. 2007. An evaluation of the taxonomic validity of *Testudo werneri*. Amphibia-Reptilia 28(3):393–401.

Auer, M. 2011. Beobachtungen zu Vorkommen und Handel von Schildkröten in Nordlaos. Sauria 33(1):3–11.

Auer, M. and Herz, M. 2006. Anmerkung zur Maximalgrösse der Maurischen Landschildkröte *Testudo graeca* (Linnaeus, 1758) in Kleinasien. Sacalia 13(4):44–47.

Auer, M. and Herz, M. 2021. Chelonians in Georgia, a travel report, with notes on the distribution of tortoises in the Caucasus region. Radiata 30(2):4–21.

Auffenberg, W. 1958. Fossil turtles of the genus *Terrapene* in Florida. Bulletin of the Florida State Museum 3:53–92.

Auffenberg, W. 1962. A new species of *Geochelone* from the Pleistocene of Texas. Copeia 1962(3):627–636.

Auffenberg, W. 1963a. *Testudo hypselonota* Bourret referred to *Geochelone radiata* (Shaw). Journal of the Bombay Natural History Society 60:462–465.

Auffenberg, W. 1963b. Fossil testudinine turtles of Florida, genera *Geochelone* and *Floridemys*. Bulletin of the Florida State Museum, Biological Sciences 7(2):53–97.

Auffenberg, W. 1966. A new species of Pliocene tortoise, genus *Geochelone*, from Florida. Journal of Paleontology 40(4):877–882.

Auffenberg, W. 1967. Notes on West Indian tortoises. Herpetologica 23:34–44.

Auffenberg, W. 1974. Checklist of fossil land tortoises (Testudinidae). Bulletin of the Florida State Museum, Biological Sciences 18:121–251.

Auffenberg, W. 1988. A new species of *Geochelone* (Testudinata: Testudinidae) from the Pleistocene of Florida (U.S.A.). Acta Zoologica Cracoviensia 31(11–27):591–604.

Auliya, M. and Chen, P.N. 2024. A note on carapace size of the Malaysian giant turtle, *Orlitia borneensis* (Testudines: Geoemydidae) in Peninsular Malaysia, and the species' conservation status. Malayan Nature Journal 76(2):143–149.

Austin, J.J. and Arnold, E.N. 2001. Ancient mitochondrial DNA and morphology elucidate an extinct island radiation of Indian Ocean giant tortoises (*Cylindraspis*). Proceedings of the Royal Society of London 268B:2515–2523.

Austin, J.J., Arnold, E.N., and Bour, R. 2002. The provenance of type specimens of extinct Mascarene Island giant tortoises (*Cylindraspis*) revealed by ancient mitochondrial DNA sequences. Journal of Herpetology 36:280–285.

Austin, J.J., Arnold, E.N., and Bour, R. 2003. Was there a second adaptive radiation of giant tortoises in the Indian Ocean? Using mitochondrial DNA to investigate speciation and biogeography of *Aldabrachelys* (Reptilia, Testudinidae). Molecular Ecology 12:1415–1424.

Auth, D.L., Smith, H.M., Brown, B.C., and Lintz, D. 2000. A description of the Mexican amphibian and reptile collection of the Strecker Museum. Bulletin of the Chicago Herpetological Society 35(4):65–85.

Averill-Murray, R.C., Woodman, A.P., and Howland, J.M. 2002. Population ecology of the Sonoran desert tortoise in Arizona. In: Van Devender, T.R. (Ed.). The Sonoran Desert Tortoise: Natural History, Biology, and Conservation. Tucson: University of Arizona Press and Arizona–Sonora Desert Museum, pp. 109–134.

Avila-Pires, T.C.S., Hoogmoed, M.S., and da Rocha, W.A. 2010. Notes on the vertebrates of northern Pará, Brazil: a forgotten part of the Guianan Region, I. Herpetofauna. Boletim do Museu Paraense de História Natural e Ethnographia, Ciências Naturais 5:13–112.

Ayaz, D. and Budak, A. 2008. Distribution and morphology of *Mauremys rivulata* (Valenciennes, 1833) (Reptilia: Testudines: Geoemydidae) in the Lake District and Mediterranean region of Turkey. Turkish Journal of Zoology 32:137–145.

Ayaz, D., Bayrakci, Y., Çiçek, K., Ihlow, F., Tok, C.V., and Fritz, U. 2021. On the brink of extinction: results of a 20-year quest for Eiselt's Pond Turtle (*Emys orbicularis eiselti*) in southeastern Turkey. Chelonian Conservation and Biology 20(2): doi:10.2744/CCB-1505.1.

Baard, E.H.W. 1995. Growth, age at maturity and sexual dimorphism in the geometric tortoise, *Psammobates geometricus*. Journal of the Herpetological Association of Africa 44:10–15.

Babcock, H.L. 1919. The Turtles of New England. Monographs of the Boston Society of Natural History 8(3):323–431.

Babcock, H.L. 1937. A new subspecies of the red-bellied terrapin *Pseudemys rubriventris* (Le Conte). Occasional Papers of the Boston Society for Natural History 8:293.

Bachmayer, F., Brinkerink, J.P., and Symeonidis, N. 1975. Pleistozäne Schildkröten aus Höhlen der Insel Kreta. Annales Géologiques des Pays Helléniques 27:110–121.

Bager, A., Bach, G., Mascarenhas, C.S., Rosado, J.L.O., and Bernardo, M.D. 2003. Morphological characterization and observations relating to the ecology of *Hydromedusa tectifera* in the south of Rio Grande do Sul - Brazil. Annals of the 2003 Joint Meeting of Ichthyologists and Herpetologists, Manaus, Brazil. Abstract.

Bager, A., Lucas, P.S., Costa, A., Lima, J.C.S., and Silveira, M.L. 2016. Morphology and sexual dimorphism of *Acanthochelys spixii* (Testudines, Chelidae) in Brazil. Tropical Zoology 29:73–86.

Baha El Din, S.M. 2006. A guide to the amphibians and reptiles of Egypt. Cairo: The American University in Cairo Press, pp. 359.

Baha El Din, S.M., Attum, O., and Baha El Din, M. 2003. Status of *Testudo kleinmanni* and *T. werneri* in Egypt. Chelonian Conservation and Biology 4(3):648–655.

Baird, S.F. and Girard, C. 1852. Descriptions of new species of reptiles, collected by the U.S. Exploring Expedition under the command of Capt. Charles Wilkes, U.S.N. First Part.—Including the species from the western coast of America. Proceedings of the Acadademy of Natural Sciences of Philadelphia 1852:174–177.

Balestra, R.A.M., Heming, N.M., Carvalho, R.M.V., Silveira, A.L., Drummond, G.M., Ferreira, D.I., and Luz, V.L.F. 2008. Diversidade e abundância de quelônios na bacia do Rio São Francisco. Goiânia: Report to Ministério do Meio Ambiente, 44 pp.

Ballasina, D. 1995. Salviamo le Tartarughe! Carapax Centre: Edagricole, 288 pp.

Ballasina, D., Vandepitte, V., Mochi, E., and Fenwick, H. 2006. La nécessité de réintroduction de *Geochelone sulcata* nées en captivité: stratégies pour la gestion de groupes d'élevage en captivité. Chelonii 4:111.

Banning, G.H. 1933. Hancock Expedition to the Galapagos Islands, 1933: General Report. Bulletin of the Zoological Society of San Diego 10:1–31.

Barbour, T. 1935. A new *Pseudemys* from Cat Island, Bahamas. Occasional Papers Boston Society of Natural History 8:205–206.

Barbour, T. and Carr, A.F., Jr. 1938. Another Bahamian fresh-water tortoise. Proceedings of the New England Zoology Club 17:75–76.

Barbour, T. and Carr, A.F., Jr. 1940. Antillean terrapins. Memoirs of the Museum of Comparative Zoology 54(5):381–415.

Barbour, T. and Carr, A.F., Jr. 1941. Terrapin from Grand Cayman. Proceedings of the New England Zoology Club 18:57–60.

Barbour, T. and Loveridge, A. 1929. Typical reptiles and amphibians in the

Museum of Comparative Zoology. Bulletin of the Museum of Comparative Zoology 69:205–360.

Barbour, T. and Loveridge, A. 1946. First supplement to typical reptiles and amphibians. Bulletin of the Museum of Comparative Zoology 96:59–214.

Barko, V.A. and Briggler, J.T. 2006. Midland Smooth Softshell (*Apalone mutica*) and Spiny Softshell (*Apalone spinifera*) turtles in the Middle Mississippi River: habitat associations, population structure, and implications for conservation. Chelonian Conservation and Biology 5(2):225–231.

Barley, A.J., Spinks, P.Q., Thomson, R.C., and Shaffer, H.B. 2010. Fourteen nuclear genes provide phylogenetic resolution for difficult nodes in the turtle tree of life. Molecular Phylogenetics and Evolution: 55(3):1189–1194.

Barrio-Amorós, C.L. 2001. *Podocnemis unifilis* (Terecay). Maximum size. Herpetological Review 32(1):39.

Barrio-Amorós, C.L. and Manrique, R. 2006. Record de taille pour une matamata (*Chelus fimbriatus*), au Venezuela. Manouria 32:23–26.

Barrio-Amorós, C.L. and Narbaiza, I. 2008. Die Schildkröten des Bundesstaates Amazonas (Venezuela). Radiata 17(1):2–20.

Barth, D., Bernhard, D., Guicking, D., Stöck, M., and Fritz, U. 2003. Is *Chinemys megalocephala* Fang, 1934 a valid species? New insights based on mitochondrial DNA sequence data. Salamandra 38(2002)[2003]:233–244.

Barth, D., Bernhard, D., Fritzsch, G., and Fritz, U. 2004. The freshwater turtle genus *Mauremys* (Testudines, Geoemydidae) – a textbook example of an east-west disjunction or a taxonomic misconcept? Zoologica Scripta 33:213–221.

Bartlett, E. 1895a. Notes on tortoises, No. 2. Sarawak Gazette 25:29–30.

Bartlett, E. 1895b. Notes on tortoises, No. 3. Sarawak Gazette 25:83–84.

Bartlett, E. 1896. Notes on tortoises, No. 4. Sarawak Gazette 26:113.

Bartram, W. 1791. Travels through North and South Carolina, Georgia, east and west Florida, the Cherokee county, the executive territories of the Muscogulges, or Creek Confederacy, and the county of the Chactaws; containing an account of the soil and natural productions of those regions, together with observations on the manners of the Indians. Philadelphia: James and Johnson, 522 pp.

Baruah, C., Chetia, P., Bhuyan, S.K., and Sharma, D.K. 2012. Diversity of freshwater turtles and their possible conservation in and around the Orang National Park, Assam, India. Tigerpaper 39(1):24–30.

Baruah, C., Devi, P., and Sharma, D.K. 2016. Comparative morphometry and biogeography of the freshwater turtles of genus *Pangshura* (Testudines: Geoemydidae: *Pangshura*). International Journal of Pure and Applied Zoology 4:107–123.

Batistella, A.M. 2008. Biologia de *Trachemys adiutrix* (Vanzolini, 1995) (Testudines, Emydidae) no litoral do Nordeste-Brasil. Ph.D. Thesis, Universidade Federal do Amazonas, Manaus.

Batsch, A.J.G.C. 1788. Versuch einer Anleitung zur Kenntniss und Geschichte der Thiere und Mineralien. Erster Theil. Allgemeine Geschichte der Natur; besondre der Säugthiere, Vögel, Amphibien und Fische. Jena: Akademische Buchhandlung, 528 pp.

Batsch, A.J.G.C. 1796. Umriss der gesammten Naturgeschichte. Jena: Christian Ernst Gabler, Vol. 1, 287 pp.

Bauer, A.M. and Adler, K. 2001. The dating and correct citation of A.F.A. Wiegmann's "Amphibien" section of Meyen's Reise um die Erde, with a bibliography of Wiegmann's herpetological publications. Archives of Natural History 28(3):313–326.

Bauer, A.M. and Petit, R.E. 2004. On the herpetology of Perry's Arcana and two forgotten reptile names. Newsletter and Bulletin of the International Society for the History and Bibliography of Herpetology 5(1):9–17.

Bauer, A.M., DeBoer, J.C., and Taylor, D.J. 2017. Atlas of the reptiles of Libya. Proceedings of the California Academy of Sciences (4)64(8):155–318.

Baur, G. 1887. Osteologische Notizen über Reptilien. Fortsetzung II. Zoologischer Anzeiger 10(244):96–102.

Baur, G. 1888a. Osteologische Notizen über Reptilien. Fortsetzung III. Zoologischer Anzeiger 11(285):417–424.

Baur, G. 1888b. Osteologische Notizen über Reptilien. Fortsetzung IV. Zoologischer Anzeiger 11(291):592–597.

Baur, G. 1888c. Notes on the American Trionychidae. American Naturalist 22:1121–1122.

Baur, G. 1889. The gigantic land tortoises of the Galapagos Islands. American Naturalist 23:1039–1057.

Baur, G. 1890a. Two new species of tortoises from the south. Science 16(405):262–263.

Baur, G. 1890b. An apparently new species of *Chelys*. American Naturalist 24:967–968.

Baur, G. 1890c. On the classification of the Testudinata. American Naturalist 24:530–536.

Baur, G. 1891a. The very peculiar tortoise, *Carettochelys* Ramsay, from New Guinea. Science 17(426):190.

Baur, G. 1891c. On the relations of *Carettochelys*, Ramsay. American Naturalist 25:631–639.

Baur, G. 1892. Bemerkungen über verschiedene Arten von Schildkröten. Zoologischer Anzeiger 15:155–159.

Baur, G. 1893a. Notes on the classification and taxonomy of the Testudinata. Proceedings of the American Philosophical Society 31:210–225.

Baur, G. 1893b. Notes on the classification of the Cryptodira. American Naturalist 27:672–674.

Baur, G. 1893c. Two new species of North American Testudinata. American Naturalist 27:675–677.

Baur, G. 1896. Der Schädel einer neuen grossen Schildkröte (*Adelochelys*) aus dem zoologischen Museum in München. Anatomischer Anzeiger 12:314–319.

Baur, G. 1925. [*Kinosternon abaxillare*]. In: Stejneger, L. New species and subspecies of American turtles. Journal of the Washington Academy of Science 15:462–463. [p. 462]

Bechstein, J.M. 1800. Herrn De la Cepede's Naturgeschichte der Amphibien oder der eyerlegenden vierfüszigen Thiere und der Schlangen. Erster Band. Weimar: Industrie-Comptoir, 524 pp.

Bedriaga, J. von. 1881. Die Amphibien und Reptilien Griechenlands. (Fortsetzung). Bulletin de la Société Impériale des Naturalistes de Moscou 56(3-4):278–344.

Beheregaray, L.B., Ciofi, C., Caccone, A., Gibbs, J.P., and Powell, J.R. 2003. Genetic divergence, phylogeography and conservation units of giant tortoises from Santa Cruz and Pinzón, Galápagos Islands. Conservation Genetics 4:31–46.

Behn, F.D. 1760. Jakob Theodor Kleins Classification und Kurze Geschichte der Vierfüssigen Thiere (Transl.). Lübeck: Jonas Schmidt, 381 pp.

Bélanger, C. 1834. Voyage aux Indes-Orientales, par le Nord de l'Europe, les Provinces du Caucase, la Géorgie, l'Arménie et la Perse, suivi de Détails topographiques, statistiques et autres sur le Pégou, les Iles de Java, de Maurice et de Bourbon, sur le Cap de Bonne-Espérance et Sainte-Hélène, pendant les Années 1825, 1826, 1827, 1828 et 1829. Préface. Paris: A. Bertrand, pp. xvii–xxxix.

Bell, C.J. and Bauer, A.M. 2020. Constantine Rafinesque's herpetological observations along the Ohio River in 1818. Bibliotheca Herpetologica 14(4):19–31.

Bell, T. 1825a. A monograph of the tortoises having a moveable sternum, with remarks on their arrangement and affinities. Zoological Journal 2:299–310.

Bell, T. 1825b. [*Sternotherus*]. In: Gray, J.E. A synopsis of the genera of reptiles and amphibia, with a description of some new species. Annals of Philosophy (2)10:211.

Bell, T. 1826. Description of a new species of *Terrapene*; with further observations on *T. carolina* and *T. maculata*. Zoological Journal 2:484–486.

Bell, T. 1827. On two new genera of land tortoises. Transactions of the Linnean Society of London 15:392–401.

Bell, T. 1828a. Descriptions of three new species of land tortoises. Zoological Journal 3(11):419–421.

Bell, T. 1828b. On *Hydraspis*, a new genus of freshwater tortoises, of the family Emydidae. Zoological Journal 3(12):511–513.

Bell, T. 1828c. Characters of the order, families, and genera of the Testudinata. Zoological Journal 3(12):513–516.

Bell, T. 1834. A freshwater tortoise described as the type of a new genus, *Cyclemys*. Proceedings of the Zoological Society of London 1834:17.

Bell, T. 1835. A Monograph of the Testudinata. Part IV. London: Samuel Highley, unnumbered pages and plates. [published prior to May 1835]

Ben Kaddour, K., El Mouden, E.H., Slimani, T., Bonnet, X., and Lagarde, F. 2008. Sexual dimorphism in the Greek Tortoise: a test of the body shape hypothesis. Chelonian Conservation and Biology 7(1):21–27.

Bennett, E.T. 1836. [Footnote 2: *Testudo whitei*]. In: White, G. The Natural History and Antiquities of Selborne. London: J. and A. Arch, 640 pp. [pp. 360–361].

Beolens, B., Watkins, M., and Grayson, M. 2011. The Eponym Dictionary of Reptiles. Baltimore, MD: Johns Hopkins University Press, 296 pp.

Bergounioux, F.M. 1936. Monographie des chéloniens fossiles conservées au Laboratoire de Géologie de la Faculté des Sciences de Lyon. Travaux du Laboratoire de Géologie du Faculté des Sciences de Lyon 31(26):7–40.

Berlandier, J.L. 1850. Reptiles. In: Berlandier, L. and Chovel, R. Diario de Viage

de la Comision de Limites que puso el Gobierno de la Republica. Mexico: Juan R. Navarro, pp. 287–291.

Berlandier, J.L. 1980. Journey to Mexico during the years 1826 to 1834. Translated by S.M. Ohlendorff, J.M. Bigelow, and M.M. Standifer. Botanical notes by C.H. Muller and K.K. Muller. Denton, TX: The Texas State Historical Association. Two Volumes, 672 pp.

Berman, M.J. 1994. Preliminary report on a vertebrate assemblage excavated from the Three Dog site, San Salvador, Bahamas. In: Kass, L.B. (Ed.). Proceedings of the Fifth Symposium on the Natural History of the Bahamas. San Salvador, Bahamas: Bahamian Field Station, Ltd., pp. 5–13.

Berry, G., Brown, G.J., Haden, L., Jones, R.L., Pearson, L., and Selman, W. 2020. Chutes and ladders: drainage exchange of Map Turtles (genus *Graptemys*) across the Tennessee-Tombigbee Waterway in northeastern Mississippi. Chelonian Conservation and Biology 19(2):262–267.

Berry, J.F. 1978. Variation and systematics in the *Kinosternon scorpioides* and *K. leucostomum* complexes (Reptilia: Testudines: Kinosternidae) of Mexico and Central America. Ph.D. Thesis, University of Utah.

Berry, J.F. and Iverson, J.B. 1980. A new species of mud turtle, genus *Kinosternon*, from Oaxaca, Mexico. Journal of Herpetology 14(4):313–320.

Berry, J.F. and Iverson, J.B. 2001a. *Kinosternon leucostomum*. Catalogue of American Amphibians and Reptiles 724:1–8.

Berry, J.F. and Iverson, J.B. 2001b. *Kinosternon scorpioides*. Catalogue of American Amphibians and Reptiles 725:1–11.

Berry, J.F. and Legler, J.M. 1980. A new turtle (genus *Kinosternon*) from northwestern Mexico. Contributions in Science, Natural History Museum of Los Angeles County 325:1–12.

Berry, J.F. and Shine, R. 1980. Sexual size dimorphism and sexual selection in turtles (Order Testudines). Oecologia 44:185–191.

Berry, J.F., Seidel, M.E., and Iverson, J.B. 1996. [*Kinosternon chimalhuaca*]. In: Rogner, M. Schildkröten 2. Hürtgenwald: Heidi-Rogner-Verlag, 265 pp. [pp. 23–24].

Berry, J.F., Seidel, M.E., and Iverson, J.B. 1997. A new species of mud turtle (genus *Kinosternon*) from Jalisco and Colima, Mexico, with notes on its natural history. Chelonian Conservation and Biology 2(3):329–337.

Beshkov, V.A. 1997. Record-sized tortoises, *Testudo graeca ibera* and *Testudo hermanni boettgeri*, from Bulgaria. Chelonian Conservation and Biology 2(4):593–596.

Besnard, G., Thèves, C., Mata, X., Holota, H., Rakotozafy, L.M.A., and Pedrono, M. 2016. Shotgun sequencing of the mitochondrial genome of the Aldabra giant tortoise (*Aldabrachelys gigantea*). Mitochondrial DNA Part A: DNA Mapping, Sequencing, and Analysis; doi:10.3109/19401736.2015.1101554, 2 pp.

Beyer, G.E. 1900. Louisiana herpetology. Proceedings of the Louisiana Society of Naturalists 1897–1899:24–46.

Bhaskar, R. and Mohindra, V. 2019. Phylogenetic relationships among Indian freshwater turtles (family Trionychidae and Geoemydidae) with special reference to *Lissemys punctata*, inferred from mitochondrial cytochrome b gene sequences. Meta Gene 22:100610.

Bhullar, B.S. and Bever, G.S. 2009. An archosaur-like laterosphenoid in early turtles (Reptilia: Pantestudines). Breviora 518:1–11.

Bibron, G. and Bory de Saint-Vincent, J.B. 1833 ["1832"]. Vertébrés à sang froid. Reptiles et poissons. In: Bory de Saint-Vincent, J.B. (Ed.). Expédition Scientifique de Morée. Travaux de la Section des Sciences Physiques. Tome III, Première Partie. Zoologie - Première Section. Animaux vertébrés, Mollusques et Polypiers. Paris: F.G. Levrault, pp. 57–80.

Bickham, J.W. and Carr, J.L. 1983. Taxonomy and phylogeny of the higher categories of cryptodiran turtles based on a cladistic analysis of chromosomal data. Copeia 1983(4):918–932.

Bickham, J.W., Lamb, T., Minx, P., and Patton, J.C. 1996. Molecular systematics of the genus *Clemmys* and the intergeneric relationships of emydid turtles. Herpetologica 52:89–97.

Biello, R., Zampiglia, M., Corti, C., Deli, G., Biaggini, M., Crestanello, B., Delaugerre, M., Di Tizio, L., Leonetti Francesco, L., Stefano, C., Olivieri, O., Pellegrino, F., Romano, A., Sperone, E., Heidi, C.H., Trabalza-Marinucci, M., Bertorelle, G., and Canestrrelli, D. 2021. Mapping the geographic origin of captive and confiscated Hermann's Tortoises: a genetic toolkit for conservation and forensic analyses. Forensic Science International: Genetics 51:102447.

Blainville, H. de 1816. Prodrome d'une nouvelle distribution systématique du règne animal. Bulletin des Sciences par la Société Philomatique de Paris 1816:113–124 ["105–112, 121–124" in error].

Blamires, S.J., Spencer, R.-J., King, P., and Thompson, M.B. 2005. Population parameters and life table analysis of two co-existing freshwater turtles: are the Bellinger River turtle populations threatened? Wildlife Research 32:339–347.

Blanck, T. 2005. *Cuora yunnanensis* (Boulenger, 1906), the Yunnan box turtle, rediscovered after one-hundred years? Radiata 14(2):10–33.

Blanck, T. and Kremser, T. 2008. First record of *Ocadia sinensis* (Gray, 1834) on Lamma Island, Hong Kong, China, with comments on the distribution, habitats, vulnerability and captive propagation of the species in China. Radiata 17(4):2–18.

Blanck, T., McCord, W.P., and Le, M. 2006a. On the variability of *Cuora trifasciata* (Bell, 1825); the rediscovery of the type specimen, with descriptions of a new *Cuora* species and subspecies, and remarks on the distribution, habitat and vulnerability of these species (Reptilia: Testudines: Geoemydidae). Frankfurt: Edition Chimaira, 153 pp.

Blanck, T., Zhou, T., and McCord, W.P. 2006b. The Yunnan box turtle, *Cuora yunnanensis* (Boulenger 1906); historical background and an update on the morphology, distribution and vulnerabilities of the only known living specimens. Sacalia 13(4):14–35.

Blanck, T., Protiva, T., Zhou, T., Li, Y., Crow, P., and Tiedemann, R. 2017. New subspecies of *Cuora cyclornata* (Blanck, McCord & Le, 2006), *Cuora trifasciata* (Bell, 1825) and *Cuora aurocapitata* (Luo & Zong, 1988). Sichuan Journal of Zoology 36(4):368–385.

Blanck, T., Gaillard, D., Protiva, T., Wheatley, M., Shi, H., Liu, L., Ray, P.C., and Anders, B. 2023. A taxonomic hide and seek: phylogenetic and phylogeographic relationships in the Southeast Asian Box Turtle, *Cuora amboinensis* (Riche *in* Daudin, 1801). Russian Journal of Herpetology 30(6) (Supplement):1–52.

Blanford, W.T. 1870. Notes on some Reptilia and Amphibia from central India. Journal of the Asiatic Society of Bengal 39:335–376.

Blasus, R.E., Gerholdt, J., and Crawford, M. 2004. *Emydoidea blandingii* (Blanding's Turtle). Maximum size. Herpetological Review 35:54.

Bleeker, P. 1857a. Berigt omtrent eenige Reptilien van Sumatra, Borneo, Batjan en Boero. Natuurkundig Tijdschrift voor Nederlandsch-Indië 13:470–475.

Bleeker, P. 1857b. Opsomming der soorten van Reptilien, tot dus verre van het eiland Java bekend geworden. Natuurkundig Tijdschrift voor Nederlandsch-Indië 14:235–244.

Bloch, M.E. 1786. Nachricht von der Dosenschildkröte. Schriften der Berlinischen Gesellschaft der Naturforschender Freunde 7(1):131–134.

Bloxam, Q.M.C. and Hayes, K.T. 1991. Further field observations on the Malagasy flat-tailed tortoise *Pyxis planicauda*. Dodo 27:138–145.

Blumenbach, J.F. 1779. Handbuch der Naturgeschichte. Ed. 1. Part 1. Göttingen: J.C. Dieterich, 448 pp.

Blyth, E. 1854 ["1853"]. Notices and descriptions of various reptiles, new or little known. Journal of the Asiatic Society of Bengal 22(7):639–655.

Blyth, E. 1856. Report for October Meeting, 1855. Donations received during the last three months. Journal of the Asiatic Society of Bengal 24:711–723.

Blyth, E. 1859. Report of Curator, Zoological Department, for May, 1858. [4. A small collection from Major Berdmore, sent by him from the Sitang Valley, Pegu]. Journal of the Asiatic Society of Bengal 27:267–281.

Blyth, E. 1863. Report of the Curator, Zoological Department. [III. W.T. Blanford, Esq., of the Indian Geological Survey. A collection of sundries from different parts of Burma]. Journal of the Asiatic Society of Bengal 32:73–90.

Bocourt, M.F. 1868. Description de quelques chéloniens nouveaux appartenant à la faune Mexicaine. Annales des Sciences Naturelles, Zoologie et Paléontologie, Paris (5)10:121–122.

Bocourt, M.F. 1876a. Note sur quelques reptiles de l'Isthme de Tehuantepec (Mexique) donnés par M. Sumichrast au Muséum. Journal de Zoologie (Paris) 5:386–411.

Bocourt, M.F. 1876b. Addition. [*Emys grayi*, Duméril et Bocourt, devra prendre un nouveau nom, *Emys umbra*, Nob.]. In: Bocourt, M.F. Sur quelques reptiles de l'Isthme de Tehuantepec (Mexique) donnés par M. Sumichrast au Muséum. Extrait du Journal de Zoologie publié par M. Paul Gervais, t. V, 1876. Paris: Imprimerie Bouchard-Huzard, pp. 1–26. [p. 26].

Boddaert, P. 1770. Brief van de kraakbeenige schildpad. Epistola de testudine cartilaginea. Amsterdam: Kornelis van Tongerlo, 39 pp.

Bodenheimer, F.S. 1935. Animal Life in Palestine. Jerusalem: L. Mayer, 235 pp.

Boettger, O. 1893. Katalog der Reptilien-Sammlung im Museum der Senckenbergischen Naturforschenden Gesellschaft in Frankfurt am Main. I. Teil

(Rhynchocephalen, Schildkröten, Krokodile, Eidechsen, Chamäleons). Frankfurt: Knauer, 140 pp.

Boettger, O. 1894. Materialien zur herpetologischen Fauna von China III. Bericht über die Senckenbergische Naturforschende Gesellschaft, Frankfurt 25:129–152.

Bogert, C.M. 1943. A new box turtle from southeastern Sonora, Mexico. American Museum Novitates 1226:1–7.

Bogert, C.M. and Oliver, J.A. 1945. A preliminary analysis of the herpetofauna of Sonora. Bulletin of the American Museum of Natural History 83:297–426.

Bohlman, B., Barron, H., Albright, S., and Rogers, K. 2019. *Gopherus polyphemus* (Gopher Tortoise). Size. Herpetological Review 50(2):352–353.

Bohls, J. 1895. Bemerkung zur Eintheilung der Chelydidae. Zoologischer Anzeiger 18:51–53.

Bolten, A.B. 1999. Techniques for measuring sea turtles. In Eckert, K.L., Bjorndal, K.A., Abreu-Grobois, F.A., and Donnelly, M. (Eds.). Research and Management Techniques for the Conservation of Sea Turtles. Washington, DC: IUCN/SSC Marine Turtle Specialist Group Publication No. 4, pp. 110–114.

Bonaparte, C.L. 1830. Sulla Seconda Edizione del Regno Animale del Barone Cuvier. Osservazioni. Annali Storia Naturale di Bologna 4:1–172.

Bonaparte, C.L. 1831. Saggio di una Distribuzione Metodica degli Animali Vertebrati. Rome: Presso Antonio Boulzaler, 144 pp.

Bonaparte, C.L. 1836. Cheloniorum Tabula Analytica. Rome: 9 pp.

Bondarenko, D.A. and Dujsebayeva, T.N. 2012. Central Asian Turtle, *Agrionemys horsfieldii* (Gray, 1844), in Kazakhstan (its distribution, habitat division, and population density). Current Studies in Herpetology 12(1/2):3–26. [in Russian]

Bondarenko, D.A. and Peregontsev, E.A. 2017. Distribution of the Central Asian Tortoise *Agrionemys horsfieldii* (Gray, 1844) in Uzbekistan (range, regional and landscape distribution, populations density). Current Studies in Herpetology 17(3/4):124–146. [in Russian]

Bondarenko, D.A., Ergashev, U.H., and Najmudinov, T.A. 2014. Contemporary state of the Central Asian Tortoise *Agrionemys horsfieldii* (Gray, 1844) populations, in southern Tajikistan. Bulletin of Moscow Society of Naturalists 119(4):19–29. [in Russian]

Bonin, F., Devaux, B., and Dupré, A. 2006. Toutes les Tortues du Monde. Paris: Delachaux and Niestle, 416 pp.

Bonnaterre, P.-J. 1789. Tableau Encyclopédique et Méthodique des Trois Règnes de la Nature. Erpétologie. Paris: Panckoucke, Hôtel de Thou, 70 pp.

Bonnemains, J. and Bour, R. 1996. Les chéloniens de la collection LeSueur du Muséum d'Histoire naturelle du Havre. Bulletin Trimestriel de la Société Géologique de Normandie et des Amis du Muséum du Havre 83(3/4):5–45.

Bonnet, X., Lagarde, F., Henen, B.T., Corbin, J., Nagy, K.A., Naulleau, G., Balhoul, K., Chastel, O., Legrand, A., and Cambag, R. 2001. Sexual dimorphism in steppe tortoises (*Testudo horsfieldii*): influence of the environment and sexual selection on body shape and mobility. Biological Journal of the Linnean Society 72:357–372.

Bonnet, X., Delmas, V., El-Mouden, H., Slimani, T., Sterijovski, B., and Kuchling, G. 2010. Is sexual body shape dimorphism consistent in aquatic and terrestrial chelonians? Zoology 113:213–220.

Bontius, J. 1658. Historiae Naturalis et Medicae Indiae Orientalis. Libri Sex. In: Piso, W. De Indiae Utriusque re Naturali et Medica. Libri Quatuordecim. Amstelaedami (Amsterdam): Ludovicum et Danielem Elzevirios, 160 pp.

Bory de Saint-Vincent, J.B. 1804. Voyage dans les Quatre Principales Îles des Mers d'Afrique. Paris: F. Buisson, Tome 2, 430 pp., Planches, 56 pls.

Bory de Saint-Vincent, J.B. (Ed.). 1833. Expédition Scientifique de Morée. Travaux de la Section des Sciences Physiques. Zoologie. Paris: F.G. Levrault, planches, troisième série, pls. 6–17.

Bory de Saint-Vincent, J.B. (Ed.). 1835. Expédition Scientifique de Morée. Travaux de la Section des Sciences Physiques. Zoologie. Paris: F.G. Levrault, Atlas, pls. 6–17 [corrigenda].

Bory de Saint-Vincent, J.B. 1836. Expédition Scientifique de Morée. Section des Sciences Physiques. Tome Premier. Relation. Paris: F.G. Levrault, 472 pp.

Bouchard, C., Tessier, N., and Lapointe, F.-J. 2019. Watersheds influence the wood turtle's (*Glyptemys insculpta*) genetic structure. Conservation Genetics 20(3):653–664.

Boulaouad, B.A., Bakhouche, B., Friel, B., and Escoriza, D. 2023. African Spurred Tortoise *Centrochelys sulcata*: range extension to Algeria. Herpetological Bulletin 164:44–45.

Boulenger, G.A. 1886a. On the South-African tortoises allied to *Testudo geometrica*. Proceedings of the Zoological Society of London 1886:540–542.

Boulenger, G.A. 1886b. A synopsis of the reptiles and batrachians of the Province Rio Grande do Sul, Brazil. Annals and Magazine of Natural History (5)18:423–445.

Boulenger, G.A. 1887a. On a new family of pleurodiran turtles. Annals and Magazine of Natural History (5)19:170–172.

Boulenger, G.A. 1887b. On the systematic position of the genus *Miolania*, Owen (*Ceratochelys*, Huxley). Proceedings of the Zoological Society of London 1887:554–555.

Boulenger, G.A. 1888a. Description of a new land-tortoise from South Africa, from a specimen living in the Society's Gardens. Proceedings of the Zoological Society of London 1888:251.

Boulenger, G.A. 1888b. On the chelydoid chelonians of New Guinea. Annali del Museo Civico di Storia Naturale di Genova (2)6:449–452.

Boulenger, G.A. 1889. Catalogue of the Chelonians, Rhynchocephalians, and Crocodiles in the British Museum (Natural History). London: Trustees of the Museum, 311 pp.

Boulenger, G.A. 1891. On some chelonian remains preserved in the Museum of the Royal College of Surgeons. Proceedings of the Zoological Society of London 1891:4–8.

Boulenger, G.A. 1895a. Esplorazione del Giuba e dei suoi affluenti compiuta dal Cap. V. Bottego durante gli anni 1892–93 sotto gli auspicii della Società Geografica Italiana. Risultati Zoologici. Rettili e Batraci. Annali del Museo Civico di Storia Naturale di Genova (2)15:7–18.

Boulenger, G.A. 1895b. On the American box turtles. Annals and Magazine of Natural History (6)15:330–331.

Boulenger, G.A. 1897a. Description of a new genus and species of tortoises from Borneo. Annals and Magazine of Natural History (6)19:468–469.

Boulenger, G.A. 1897b. Description of a new tortoise of the genus *Sternothaerus*. Proceedings of the Zoological Society of London 1897:919.

Boulenger, G.A. 1902a. Descriptions of new batrachians and reptiles from north-western Ecuador. Annals and Magazine of Natural History (7)9:51–57.

Boulenger, G.A. 1902b. A list of the fishes, batrachians, and reptiles collected by Mr. J. ffolliott Darling in Mashonaland, with descriptions of new species. Proceedings of the Zoological Society of London 1902(2):13–18.

Boulenger, G.A. 1903a. Report on the batrachians and reptiles. In: Annandale, N. and Robinson, H.C. (Eds.). Fasciculi Malayenses: Anthropological and Zoological Results of an Expedition to Perak and the Siamese Malay States, 1901–1902. Zoology. Liverpool: University Press, pp. 131–170.

Boulenger, G.A. 1903b. On a collection of batrachians and reptiles from the interior of Cape Colony. Annals and Magazine of Natural History (7)12:215–217.

Boulenger, G.A. 1906a. Descriptions of new reptiles from Yunnan. Annals and Magazine of Natural History (7)17:567–568.

Boulenger, G.A. 1906b. Report on the reptiles collected by the late L. Fea in West Africa. Annali del Museo Civico di Storia Naturale di Genova (3)2(42):196–216.

Boulenger, G.A. 1907. A new tortoise from Travancore. Journal of the Bombay Natural History Society 17:560–561.

Boulenger, G.A. 1913. On a collection of batrachians and reptiles made by Dr. H.G.F. Spurrell, F.Z.S., in the Choco, Colombia. Proceedings of the Zoological Society of London 1913(4):1019–1038.

Boulenger, G.A. 1920. Une tortue extraordinaire: *Testudo loveridgii*, sp.n. Comptes Rendus de l'Académie des Sciences, Paris 170:263–266.

Boulenger, G.A. 1921. Description of a new land-tortoise from northern Persia. Journal of the Bombay Natural History Society 27:251–252.

Boundy, J. and Kennedy, C. 2006. Trapping survey results for the Alligator Snapping Turtle (*Macrochelys temminckii*) in southeastern Louisiana, with comments on exploitation. Chelonian Conservation and Biology 5(1):3–9.

Bour, R. 1973. Contribution à la connaissance de *Phrynops nasutus* (Schweigger: 1812) et *Phrynops tuberculatus* (Luederwaldt: 1926). Description d'une nouvelle sous-espèce originaire du Paraguay, *Phrynops tuberculatus vanderhaegei* (Testudinata – Pleurodira – Chelidae). Bulletin de la Société Zoologique de France 98(1):175–190.

Bour, R. 1978. Les tortues des Mascareignes; description d'une espèce nouvelle d'après un document (Mémoires de l'Académie) de 1737 dans lequel le crâne est figuré. Comptes Rendus de l'Académie des Sciences 287D:491–493.

Bour, R. 1979. Les tortues actuelles de Madagascar (République malgache): liste systématique et description de deux sous–espèces nouvelles (Reptilia–Testudines). Bulletin de la Société d'Etudes Scientifiques de l'Anjou n.s. 10(1978)[1979]:141–154.

Bour, R. 1980a. Essai sur la taxinomie des Testudinidae actuels (Reptilia, Chelonii). Bulletin du Muséum National d'Histoire Naturelle, Paris (4)2A:541–546.

Bour, R. 1980b. Position systematique de *Geoclemys palaeannamitica* Bourret, 1941 (Reptilia-Testudines-Emydidae). Amphibia-Reptilia 1(2):149–159.

Bour, R. 1981. Etude systematique du genre endemique malgache *Pyxis* Bell, 1827 (Reptilia, Chelonii). Bulletin Mensuel de la Société Linnéenne de Lyon 50:132–144, 154–176.

Bour, R. 1982a. Contribution à la connaisance des tortues terrestres des Seychelles: définition du genre endémique et description d'une espèce nouvelle probablement originaire des îles granitiques et au bord de l'extinction. Comptes Rendus de l'Académie des Sciences 295:117–122.

Bour, R. 1982b. Etude systematique du genre endemigue malgache *Pyxis* Bell, 1827 (Reptilia, Chelonii) (Note complémentaire). Bulletin Mensuel de la Société Linnéenne de Lyon 51:28–31.

Bour, R. 1982c. *Pelomedusa subrufa* (Lacepède, 1788), *Pelusios subniger* (Lacepède, 1788) (Reptilia, Chelonii) et le séjour de Philibert Commerson à Madagascar. Bulletin du Muséum National d'Histoire Naturelle, Paris (4)4A:531–539.

Bour, R. 1983. Trois populations endémiques du genre *Pelusios* (Reptilia, Chelonii, Pelomedusidae) aux îles Seychelles; relations avec les espèces africaines et malgaches. Bulletin du Muséum National d'Histoire Naturelle, Paris (4)5A:343–382.

Bour, R. 1984a. Note sur *Pelusios williamsi* Laurent, 1965 (Chelonii, Pelomedusinae). Revue Français Aquariologie 11:27–32.

Bour, R. 1984b. Taxonomy, history and geography of Seychelles land tortoises and fresh-water turtles. In: Stoddart, D.R. (Ed.). Biogeography and Ecology of the Seychelles Islands. The Hague: W. Junk, pp. 281–307.

Bour, R. and Dubois, A. 1984c. Comment on the application concerning *Sternotherus* Gray, 1825 and *Pelusios* Wagler, 1830. Bulletin of Zoological Nomenclature 41:198–205.

Bour, R. 1985 ["1984"]. Les tortues terrestres géantes des îles de l'Océan Indien occidental: données géographiques, taxinomiques et phylogénétiques. In: Broin, F. de. and Jiménez–Fuentes, E. (Eds.). Studia Geologica Salmanticensia Vol. Esp. 1. Studia Palaeocheloniologica 1:17–76.

Bour, R. 1986. Note sur *Pelusios adansonii* (Schweigger, 1812) et sur une nouvelle espèce affine du Kenya (Chelonii, Pelomedusidae). Studia Palaeocheloniologica 2:23–54.

Bour, R. 1987a ["1986"]. L'identité des Tortues terrestres européennes: spécimens-types et localités-types. Revue Française d'Aquariologie 13(4):111–122.

Bour, R. 1987b. Type-specimen of the alligator snapper, *Macroclemys temminckii* (Harlan, 1835). Journal of Herpetology 21(4):340–343.

Bour, R. 1988. Taxonomic and nomenclatural status of *Homopus signatus* (Gmelin, 1789): Reptilia - Chelonii. Journal of the Herpetology Association of Africa 35:1–6.

Bour, R. 1989 ["1988"]. Caractères diagnostiques offerts par le crâne des tortues terrestres du genre *Testudo*. Mésogée 48:13–19.

Bour, R. 1994. Recherches sur des animaux doublement disparus: les tortues géantes subfossiles de Madagascar. Mémoires et Travaux de l'Institut de Montpellier 19:1–253.

Bour, R. 1996 ["1995"]. Une nouvelle espèce de tortue terrestre dans le Péloponnèse (Grèce). Dumerilia 2(1995)[1996]:23–54.

Bour, R. 1998. Histoire du genre *Manouria* Gray, 1854, et des espèces actuelles incluses. Manouria 1(1):25–40.

Bour, R. 2000. Une nouvelle espèce de *Pelusios* du Gabon (Reptilia, Chelonii, Pelomedusidae). Manouria 3(8):1–32.

Bour, R. 2001. Gravures et lithographies anciennes figurant des tortues terrestres du genre *Testudo*. Chelonii 3:12–27.

Bour, R. 2003. Previously unrecognized original type specimens of American turtles collected by John Le Conte in 1828. Chelonian Conservation and Biology 4(3):537–547.

Bour, R. 2004a. Holotype descriptions: *Kinosternon cruentatum* Duméril et Bibron, 1851. Emys 11(3):32–34.

Bour, R. 2004b. Type descriptions: holotype of *Pelusios subniger* (Lacepède, 1788). Emys 11(5):32–33.

Bour, R. 2004c. Type descriptions: holotype of "La grande tortue des Indes" in Perrault, 1676 – *Cylindraspis indica* (Schneider, 1783). Emys 11(6):33–35.

Bour, R. 2004d. Deux nouvelles tortues terrestres au Maroc. Manouria 7(25):12–13.

Bour, R. 2005a. The type specimens of *Cuora galbinifrons* Bourret, 1939. Emys 12(1):33–36.

Bour, R. 2005b. The type specimens of *Emysaurus rossignonii* Bocourt, 1868. Emys 12(2):32–34.

Bour, R. 2005c. The type specimens of *Testudo planicauda* Grandidier, 1867 and *Testudo morondavaensis* Vuillemin, 1972. Emys 12(4):22–27.

Bour, R. 2005d. Type specimens of *Emys gibba* Scheigger, 1812, *Platemys miliusii* Duméril and Bibron, 1835, and other related nominal species. Emys 12(5):24–29.

Bour, R. 2005e. Type specimens of *Testudo rotunda* Latreille, 1801. Emys 12(6):23–27.

Bour, R. 2006a. Types of Podocnemidae in the Muséum National d'Histoire Naturelle. Emys 13(1):27–40.

Bour, R. 2006b. The holotype of *Chelonia agassizii* Bocourt, 1868. Emys 13(2):18–21.

Bour, R. 2006c. Identity of *Testudo gigantea* Schweigger, 1812 and rediscovery of the type specimen. Emys 13(4):12–23.

Bour, R. 2006d. Le genre *Kinixys* Bell: histoire nomenclaturale et taxinomique. Chéloniens 3:8–15.

Bour, R. 2006e. *Kinixys belliana domerguei* (Vuillemin, 1972), la Tortue à dos articulé de Madagascar: données historiques et ecologiques. Chéloniens 3:42–55.

Bour, R. 2007a. The plowshare tortoise: past, present, and uncertain future: the type of *Testudo yniphora* Vaillant, 1885, with a selected bibliography. Emys 14(1):33–46.

Bour, R. 2007b. The type specimens of *Rhinoclemmys areolata* (Duméril and Bibron, 1851), *R. pulcherrima incisa* (Bocourt, 1868), and *R. punctularia* (Daudin, 1801). Emys 14(2):28–34.

Bour, R. 2007c. The holotype of *Testudo cartilaginea* Boddaert, 1770, and the lectotype of *Trionyx javanicus* Geoffroy Saint-Hilaire, 1809. Emys 14(3):26–32.

Bour, R. 2007d. The holotype of *Emys nasuta* Schweigger, 1812. Emys 14(4):28–32.

Bour, R. 2008a. August Friedrich Schweigger (1783–1821). In: Bauer, A.M. (Ed.). The Life and Herpetological Contributions of August Friedrich Schweigger (1783–1821). Facsimile Reprints in Herpetology, Society for the Study of Amphibians and Reptiles, pp. 7–54.

Bour, R. 2008b. The type specimens of *Testudo angulata* Schweigger, 1812 and *Testudo bellii* Gray, 1828. Emys 15(1):28–34.

Bour, R. 2008c. Types of three species of sideneck turtle belonging to the genus *Phrynops* Wagler, 1830: *Emys geoffroana* Schweigger, 1812, *Platemys hilarii* Duméril & Bibron, 1835 and *Platemys waglerii* Duméril & Bibron, 1835. Emys 15(2):35–41.

Bour, R. 2008d. The holotypes of *Pentonyx gabonensis* A. Duméril, 1856 and *Cryptopodus aubryi* A. Duméril, 1856. Emys 15(3):41–44.

Bour, R. 2008e. Neotype of *Emys castanea* Schweigger, 1812 (Pelomedusidae). Emys 15(4):36–40.

Bour, R. 2009a. Type specimens of *Testudo subrufa* Lacepède, 1788 and *Emys olivacea* Schweigger, 1812. Emys 16(1)26–30.

Bour, R. 2009b. Type specimens of *Emys ocellata* Duméril and Bibron, 1835 with notes on the species of *Morenia* Gray, 1870. Emys 16(2):33–42.

Bour, R. 2009c. The types of *Tetraonyx longicollis* Lesson, 1831 and *Tetraonyx lessonii* Duméril and Bibron, 1835, and the confusing history of the generic names *Tetronyx* Lesson, 1832 and *Batagur* Gray, 1855. Emys 16(3):30–38.

Bour, R. 2009d. Holotype and type locality of *Sternothaerus niger* Duméril and Bibron, 1835. Emys 16(4):33–37.

Bour, R. 2010. *Trionyx spiniferus* LeSueur, 1827 and *Trionyx muticus* LeSueur, 1827. Type specimens and original illustrations. Emys 17(1):26–36.

Bour, R. 2013a. Actualités chez les tortues des Seychelles. Chéloniens 29:27–41.

Bour, R. 2013b. Le type de *Testudo sulcata* Miller, 1779. Chéloniens 31:14–27.

Bour, R. 2017. About the original description of Berlandier's tortoise, *Xerobates berlandieri* Agassiz, 1857 – notes on Jean Louis Berlandier. II. Bibliotheca Herpetologica 13(1-2):50–67.

Bour, R. and Dubois, A. 1983. Statut nomenclatural et spécimens-types d'*Emys pseudogeographica* Gray, 1831 et d'*Emys lesueuri* Gray, 1831 (Reptilia, Chelonii, Emydidae). Bulletin Mensuel de la Société Linnéenne de Lyon 52:42–46.

Bour, R. and Dubois, A. 1984a ["1983"]. Nomenclatural availability of *Testudo coriacea* Vandelli, 1761: a case against a rigid application of the rules to old, well-known zoological works. Journal of Herpetology 17(4)(1983):356–361.

Bour, R. and Dubois, A. 1984b. Nomenclature ordinale et familiale des tortues (Reptilia). In: Broin, F. de and Jiminez-Fuentes, E. (Eds.). Studia Geologica Salmanticensia, Vol. Esp. 1. Studia Palaeocheloniologica I:77–86.

Bour, R. and Henkel, M. 2012. Hybridation entre la tortue d'Aldabra *Dipsochelys*

dussumieri (Gray, 1831) et la tortue sillonnée *Centrochelys sulcata* (Miller, 1779). Chéloniens 26:36–41.

Bour, R. and Maran, J. 1999 ["1998"]. Taxinomie de *Mauremys leprosa* (Schweigger, 1812) dans le sud du Maroc: la "tortue aux yeux bleus" (Reptilia, Chelonii, Geoemydidae). Manouria 1(2)(1998)[1999]:22–52.

Bour, R. and Maran, J. 2003. Une nouvelle espèce de *Pelusios* de Côte d'Ivoire (Reptilia, Chelonii, Pelomedusidae). Manouria 6(21):24–43.

Bour, R. and Ohler, A. 2008. *Chersine* Merrem, 1820 and *Chersina* Gray, 1831: a nomenclatural survey. Zootaxa 1752:66–68.

Bour, R. and Pauler, I. 1987. Identité de *Phrynops vanderhaegei* Bour, 1973, et des espèces affines (Reptilia - Chelonii - Chelidae). Mésogée – Bulletin du Museum d'Histoire Naturelle de Marseille 47:3–23.

Bour, R. and Schmidtler, J. 2014. Les dessins et aquarelles de Nikolaus Michael Oppel. 2. Les tortues. München and Paris: Privately printed, 48 pp.

Bour, R. and Zaher, H. 2005. A new species of *Mesoclemmys*, from the open formations of northeastern Brazil (Chelonii, Chelidae). Papeis Avulsos de Zoologia 45:295–311.

Bour, R., Dubois, A., and Webb, R.G. 1995. Types of recent trionychid turtles in the Muséum national d'Histoire naturelle, Paris. Dumerilia 2:73–92.

Bour, R., Devaux, B., Pritchard, P.C.H., and Wood, R.C. 2001. Observations sur *Pelusios adansonii* au Sénégal. Manouria 4(11):23–32.

Bour, R., Pritchard, P., Cheke, A., Collie, J., Arnold, E.N., Meylan, P.A., Bury, R.B., Dodd, C.K., Jr., Kraus, O., McCarthy, C., Fleischer-Dogley, F., et al., Casale, P., Gaffney, E.S., Tatayah, V., Jones, C., Reynolds, R.P., Howell, K.M., Ng, P.K.L., Chellam, R., Palkovacs, E.P., and Gerlach, J. 2009. Comments on the proposed conservation of usage of *Testudo gigantea* Schweigger, 1812 (currently *Geochelone (Aldabrachelys) gigantea*; Reptilia, Testudines) (Case 3463; see BZN 66: 34–50, 80–87). Bulletin of Zoological Nomenclature 66(2):169–186.

Bour, R., Mourer-Chauviré, C., and Ribes, S. 2014. Morphology and palaeontological exploration (up to 2000) of the extinct tortoises of the Mascarene islands. In: Gerlach, J. (Ed.). Western Indian Ocean Tortoises: Ecology, Diversity, Evolution, Conservation, Palaeontology. Manchester, United Kingdom: Siri Scientific Press, pp. 121–202.

Bourque, J.R. 2012a. An extinct mud turtle of the *Kinosternon flavescens* group (Testudines, Kinosternidae) from the middle Miocene (late Barstovian) of New Mexico. Journal of Vertebrate Paleontology 32(1):68–81.

Bourque, J.R. 2012b. A fossil mud turtle (Testudines, Kinosternidae) from the early middle Miocene (early Barstovian) of New Mexico. Journal of Vertebrate Paleontology 32(4) 836–853.

Bourque, J.R. 2016. New mud turtles (Kinosternidae, Kinosternon) from the middle–late Miocene of the United States. Journal of Paleontology 89:821–844.

Bourque, J.R. and Schubert, B.W. 2015. Fossil musk turtles (Kinosternidae, *Sternotherus*) from the Miocene-early Pliocene (Hemphillean) of Tennessee and Florida. Journal of Vertebrate Paleontology 35, doi:10.1080/02724634.2014.885441.

Bourque, J.R., Hulbert, R.C., and Wood, A.R. 2012. Assessing species diversity and intrapecific variability in shield-tailed tortoises (Testudinidae, *Hesperotestudo*) spanning the Early Clarendonian through Late Rancholabrean of Florida (Abstract). Journal of Vertebrate Paleontology 32:67.

Bourquin, O. 1992. Biggest recorded Natal hinged tortoise. Tortoise Society of Natal Newsletter 3(3):7–8.

Bourret, R. 1939. Notes herpétologiques sur l'Indochine française. XVI. Tortues de la collection du Laboratoire des Sciences Naturelles de l'Université. Description d'une espèce nouvelle. Annexe au Bulletin Général de l'Instruction Publique 1939(6):1–34.

Bourret, R. 1940 ["1939"]. Notes herpétologiques sur l'Indochine française. XVIII. Reptiles et batraciens reçus au Laboratoire des Sciences Naturelles de l'Université au cours de l'année 1939. Descriptions de quatre espèces et d'une variété nouvelles. Annexe au Bulletin Général de l'Instruction Publique 1939(4)[1940]:1–40.

Bourret, R. 1941a. Note sur un crâne de tortue fossile. Comptes Rendus des Séances du Conseil de Recherches Scientifiques de l'Indochine 1940–1941. 1941(1):9–11.

Bourret, R. 1941b. Notes herpétologiques sur l'Indochine française. XXI. Reptiles et batraciens reçus au Laboratoire des Sciences Naturelles de l'Université au cours de l'année 1940. Description d'une espèce fossile nouvelle. Annexe au Bulletin Général de l'Instruction Publique 1941:1–16.

Bourret, R. 1941c. Les Tortues de l'Indochine. Notes Institut Océanographique de l'Indochine 38:1–235.

Bowen, B.W. and Karl, S.A. 2007. Population genetics and phylogeography of sea turtles. Molecular Ecology 16(23):4886–4907.

Bowen, B.W., Meylan, A.B., Ross, J.P., Limpus, C.J., Balazs, G.H., and Avise, J.C. 1992. Global population structure and natural history of the green turtle (*Chelonia mydas*) in terms of matriarchal phylogeny. Evolution 46:865-881.

Bowen, B.W., Nelson, W.S., and Avise, J.C. 1993. A molecular phylogeny for marine turtles: trait mapping, rate assessment, and conservation relevance. Proceedings of the National Academy of Science 90:5574–5577.

Boycott, R.C. 2001. The terrapins and tortoises (Chelonia: Pelomedusidae and Testudinidae) of Swaziland. Durban Museum Novitates 26: 25–37.

Boycott, R.C. and Bourquin, O. 2000. The Southern African Tortoise Book. A Guide to Southern African Tortoises, Terrapins and Turtles. Revised expanded edition. Pietermaritzburg: Interpak, 228 pp.

Bozhanski, A.T., Makeyev, V.M., Miroutenko, M.V., and Frolov, V.Y. 1996. Distribution, numbers and morphological characteristics of the Central Asian tortoise (*Agrionemys horsfieldi*) in Turkmenistan. In: Devaux, B. (Ed.). Proceedings – International Congress of Chelonian Conservation. Gonfaron, France: Editions SOPTOM, pp. 207–209.

Bradley, R. 1721. A Philosophical Account of the Works of Nature, Endeavouring to Set Forth the Several Gradations Remarkable in the Mineral, Vegetable, and Animal Parts of the Creation. London: W. Mears, 194 pp.

Brakels, P., Chea, S., and Jones, C. 2016. Range extension of *Cyclemys atripons* Iverson & McCord 1997 with the discovery of a population in Oddar Meanchey Province, northwestern Cambodia. Cambodian Journal of Natural History 2016(1):20–22.

Bramble, D.M. 1982. *Scaptochelys*: generic revision and evolution of gopher tortoises. Copeia 1982(4):852–867.

Bramble, D.M. and Hutchison, J.H. 2014. Morphology, taxonomy, and distribution of of North American tortoises. In: Rostal, D.C., McCoy, E.D., and Mushinsky, H.R. (Eds.). Biology and Conservation of North American Tortoises. Baltimore, Maryland: Johns Hopkins University Press, pp. 1–12.

Branch, W.R. 1989. *Homopus bergeri*, nama or Berger's padloper (English), Bergerse skilpad (Afrikaans). In: Swingland, I.R. and Klemens, M.W. (Eds.). The Conservation Biology of Tortoises. Occasional Papers of the IUCN Species Survival Commission No. 5, pp. 75–77.

Branch, W.R. 1992. *Homopus 'bergeri'* - a wrong name for a new tortoise from southern Namibia. Journal of the Herpetological Association of Africa 40:11.

Branch, W.R. 2007. A new species of tortoise of the genus *Homopus* (Chelonia: Testudinidae) from southern Namibia. African Journal of Herpetology 56:1–21.

Branch, W. 2008. Tortoises Terrapins and Turtles of Africa. Cape Town: Struik Publishers, 128 pp.

Brandt, J.F. 1857. Observationes quadam ad generis Trionychum species duas novas spectantes. Bulletin de la Classe Physico-mathématique de l'Académie Impériale des Sciences de St.-Pétersbourg 16:110–111.

Brandt, J.F. and Ratzeburg, J.T.C. 1829. Medizinische Zoologie oder getreue Darstellung und Beschreibung der Thiere. Erster Band. Berlin: A. Hirschwald, 198 pp.

Bressan, R.F. and Verrastro, L. 2020. Size + shape = morphological variation in a population of *Phrynops williamsi* from southern Brazil. 18th Annual Symposium on the Conservation and Biology of Tortoises and Freshwater Turtles, Program and Abstracts, p. 23.

Brimley, C.S. 1928. Two new terrapins of the genus *Pseudemys* from the southern states. Journal of the Elisha Mitchell Science Society 44:66–69.

Bringsøe, H. and Buskirk, J.R. 1998. Distribution of *Testudo kleinmanni* Lortet, 1883 and *Testudo graeca* Linnaeus, 1758 in the Negev desert, southern Israel (Reptilia: Testudines: Testudinidae). Faunistische Abhandlungen Staatliches Museum fur Tierkunde Dresden 21, Suppl. 4:23–30.

Bringsøe, H., Buskirk, J.R., and Willemsen, R.E. 2001. *Testudo marginata* Schoepff, 1792 – Breitrandschildkröte. In: Fritz, U. (Ed.). Handbuch der Reptilien und Amphibien Europas, Band 3/IIIA. Wiesbaden: Aula-Verlag, pp. 291–334.

Brito, E.S., Strüssmann, C., Kawashita-Ribeiro, R.A., Morais, D.H., Ávila, R.W., and Campos, V.A. 2012. New records and distribution extensions of three species of *Mesoclemmys* Gray, 1863 (Testudines: Chelidae) in Mato Grosso state, Brazil, with observations on terrestrial movements. Check List 8:294–297.

Brito, E.S., Valadão, R.M., Cunha, F.A.G., de Araújo, C.G., Viana, P.F., and Médice, I.F. 2019a. New records of *Mesoclemmys raniceps* (Testudines, Chelidae) for the states of Amazonas, Pará and Rondônia, North Brazil, including the Tocantins basin. Herpetology Notes 12:283–289.

Brito, E.S., Dorado-Rodrigues, T.F., and Valadão, R.M. 2019b. First record of *Mesoclemmys vanderhaegei* (Bour, 1973) (Testudines, Chelidae) for the north-northeast Atlantic Basin. Herpetology Notes 12:709–712.

Brito, E.S., Rodrigues, E.S., Ferrão, M., Ferreira, V.L., Tomas, W.M., and Strüssman, C. 2020. *Acanthochelys macrocephala* (Big-headed Pantanal Swamp Turtle). Population and movement. Herpetological Review 51:104–105.

Broadley, D.G. 1981a. A review of the genus *Pelusios* Wagler in southern Africa (Pleurodira: Pelomedusidae). Occasional Papers of the National Museums and Monuments of Rhodesia, B. Natural Sciences 6(9):633–686.

Broadley, D.G. 1981b. A review of the populations of *Kinixys* (Testudinidae) occurring in south-eastern Africa. Annals of the Cape Provincial Museums (Natural History) 13:195–216.

Broadley, D.G. 1989. *Geochelone pardalis*, leopard tortoise (English), bergskilpad (Afrikaans). In: Swingland, I.R. and Klemens, M.W. (Eds.). The Conservation Biology of Tortoises. Occasional Papers of the IUCN Species Survival Commission No. 5, pp. 43–46.

Broadley, D.G. 1992. The Savannah species of *Kinixys* (Testudinidae). The Journal of the Herpetological Association of Africa 40:12–13.

Broadley, D.G. 1993. A review of the southern African species of *Kinixys* Bell (Reptilia, Testudinidae). Annals of the Transvaal Museum 36(6):41–52.

Broin, F. de. 1969. Sur la présence d'une tortue, *Pelusios sinuatus* (A. Smith) au Villafranchien Inférieur du Tchad. Bulletin de la Société Géologique de France (7)11:909–916.

Broin, F. de. 1988. Les tortues et le Gondwana. Examen des rapports entre le fractionnement du Gondwana et la dispersion géographique des tortues pleurodires à partir du Crétacé. Studia Palaeocheloniologica 2:103–142.

Brongersma, L.D. 1961. Notes upon some sea turtles. Zoologische Verhandelingen, Rijksmuseum van Natuurlijke Historie Leiden 51:1–46.

Brongersma, L.D. 1972. European Atlantic turtles. Zoologische Verhandelingen, Rijksmuseum van Natuurlijke Historie Leiden 121:1–318.

Brongniart, A. 1800a. Essai d'une classification naturelle des reptiles. [1]. Magasin Encyclopédique, ou Journal des Sciences, des Lettres et des Arts (5)6[An 8]:184–201.

Brongniart, A. 1800b. Essai d'une classification naturelle des reptiles. [2]. Bulletin des Sciences par la Société Philomatique 3(2)[An 8]:81–82, 89–91.

Brongniart, A. 1805. Essai d'une classification naturelle des reptiles. [3]. Paris: Baudouin, Imprimeur de l'Institut National, 53 pp.

Brooks, R.J., Shilton, C.M., Brown, G.P., and Quinn, N.W.S. 1992. Body size, age distribution, and reproduction in a northern population of wood turtles (*Clemmys insculpta*). Canadian Journal of Zoology 70:462–469.

Brophy, T.R. 2002. Variation and systematics of the Malayan snail-eating turtle, *Malayemys subtrijuga* (Schlegel and Müller, 1844). Ph.D. Thesis, George Mason University, Fairfax, Virginia.

Brophy, T.R. 2004. Geographic variation and systematics in the south-east Asian turtles of the genus *Malayemys* (Testudines: Bataguridae). Hamadryad 29:63–79.

Brophy, T.R. and Ernst, C.H. 2004. Sexual dimorphism, allometry and vertebral scute morphology in *Notochelys platynota* (Gray, 1834). Hamadryad 29:80–88.

Brown, A.D., Temple-Miller, K., Roosenberg, W.M., and White, M.M. 2012. Mitochondrial DNA variation in the Ouachita map turtle. Copeia 2012:301–306.

Brown, A.E. 1908. Generic types of Nearctic Reptilia and Amphibia. Proceedings of the Academy of Natural Sciences, Philadelphia 60:112–127.

Brown, D.J., Mali, I., Jones, M.C., and Forstner, M.R.J. 2020. Morphometric data for five freshwater turtles in south, central, and west Texas. Data in Brief 29:105356, 5 pp.

Brown, G.J., Pearson, L., Berry, G., and Selman, W. 2020. Discovery of the Northern Map Turtle (*Graptemys geographica* LeSueur 1817) in far northeastern Mississippi. Chelonian Conservation and Biology 19(2):256–261.

Brown, P. 1776. New Illustrations of Zoology. London: B. White, 136 pp.

Brown, R.W. 1956. Composition of Scientific Words. Washington, DC: Smithsonian Institution Press, 882 pp.

Brown, T.W., Augustinus, E., Izaguirre, A., and Solis, J.M. 2021. Central American Snapping Turtle (Chelydridae, *Chelydra rossignonii*) on Utila Island, Honduras, demonstrates hurricanes are a likely past and future oversea dispersal pathway for species introduction in the Caribbean. Caribbean Journal of Science 51:30–36.

Browne, P. 1756. The Civil and Natural History of Jamaica. London: T. Osborne and J. Shipton, 503 pp.

Bruguière, J.G. 1792. Description d'une nouvelle espèce de tortue de Cayenne. Journal d'Histoire Naturelle, Paris 1(7):253–261.

Bruno, S. 1986. Guida a Tartarughe e Sauri d'Italia. Firenze: Giunti, 255 pp.

Bu, X., Wang, X., Cai, W., Xia, X., and Nie, L. 2019. Genetic diversity of the captive Chinese Pond Turtle (*Mauremys reevesii*) populations in China assessed by microsatellite markers. Journal of Plant and Animal Sciences 29:1160–1168.

Buckner, S.D., Franz, R., and Reynolds, R.G. 2012. Bahama Islands and Turks & Caicos Islands. In: Powell, R. and Henderson, R.W. (Eds.). Island lists of West Indian amphibians and reptiles. Bulletin of the Florida Museum of Natural History 51(2):93–110.

Buhlmann, K.A., Akre, T.S.B., Iverson, J.B., Karapatakis, D., Mittermeier, R.A., Georges, A., Rhodin, A.G.J., van Dijk, P.P., and Gibbons, J.W. 2009. A global analysis of tortoise and freshwater turtle distributions with identification of priority conservation areas. Chelonian Conservation and Biology 8(2):116–149.

Burbidge, A.A. 1967. The biology of south-western Australian tortoises. Ph.D. Thesis, University of Western Australia.

Burbidge, A.A., Kirsch, J.A.W., and Main, A.R. 1974. Relationships within the Chelidae (Testudines: Pleurodira) of Australia and New Guinea. Copeia 1974:392–409.

Burke, A.C., Anderson, M., Weld, A., and Gaffney, E.S. 1983. The reconstruction and casting of a large extinct turtle, *Meiolania*. Curator 26:5–25.

Burke, R.L., Leuteritz, T.E., and Wolf, A.E. 1996. Phylogenetic relationships of emydine turtles. Herpetologica 52:572–584.

Burleigh, R. and Arnold, E.N. 1986. Age and dietary differences of recently extinct Indian Ocean tortoises *Geochelone* sensu lato revealed by carbon isotope analysis. Proceedings of the Royal Society of London, Biological Sciences 227B:137–144.

Burleigh, R., Matthews, K., and Ambers, J. 1982. British Museum natural radiocarbon measurements XIV. Radiocarbon 24(3):229–261.

Burmeister, H. 1837. Handbuch der Naturgeschichte. Zweite Abtheilung: Zoologie. Berlin: Verlag Enslin, 858 pp.

Bury, R.B. 2017. Biogeography of Western Pond Turtles in the western Great Basin: dispersal across a Northwest Passage? Western Wildlife 4:72–80.

Buskirk, J.R. 1989. A third specimen and neotype of *Heosemys leytensis* (Chelonia: Emydidae). Copeia 1989:224–227.

Buskirk, J.R. 1993. Yucatan box turtle *Terrapene carolina yucatana*. Tortuga Gazette 29(5):1–4.

Butler, J.A. 2002. Population ecology, home range, and seasonal movements of the Carolina diamondback terrapin, *Malaclemys terrapin centrata*, in northeastern Florida. Final Report. Tallahassee, FL: Florida Fish and Wildlife Conservation Commission.

Butler, J.A., Seigel, R.A., and Mealey, B.K. 2006. *Malaclemys terrapin* – diamondback terrapin. In: Meylan, P.A. (Ed.). Biology and Conservation of Florida Turtles. Chelonian Research Monographs No. 3, pp. 279–295.

Butler, J.M., Dodd, C.K., Jr., Aresco, M., and Austin, J.D. 2011. Morphological and molecular evidence indicates that the Gulf Coast box turtle (*Terrapene carolina major*) is not a distinct evolutionary lineage in the Florida Panhandle. Biological Journal of the Linnean Society 102:889–901.

Caballero, S., Martínez, J.G., Morales-Betancourt, M.A., Bolaños, N., and Lasso, C.A. 2022. Genomic analyses of the scorpion mud turtle (*Kinosternon scorpioides*) (Linnaeus, 1766) in insular and continental Colombia: evidence for multiple conservation and taxonomic units. Frontiers in Conservation Science 3:938040:1–16.

Cabrera, M.R. 1998. Las Tortugas Continentales de Sudamerica Austral. Córdoba: BR Copias, pp. 108.

Cabrera, M.R. and Colantonio, S.E. 1997. Taxonomic revision of the South American subspecies of the turtle *Kinosternon scorpioides*. Journal of Herpetology 31: 507–513.

Cacciali, P., Scott, N.J., Aquino Ortíz, A.L., Fitzgerald, L.A., and Smith, P. 2016. The Reptiles of Paraguay: Literature, Distribution, and an Annotatd Taxonomic Checklist. Special Publication of the Museum of Southwestern Biology No. 11, 373 pp.

Caccone, A. 2020. Evolution and phylogenetics. In: Gibbs, J.P., Cayot, L.J., and Tapia Aguilera, W. (Eds.). Galapagos Giant Tortoises. London: Academic Press, pp. 117–138.

Caccone, A., Gibbs, J.P., Ketmaier, V., Suatoni, E., and Powell, J.R. 1999. Origin and evolutionary relationships of giant Galápagos tortoises. Proceedings of the National Academy of Sciences (PNAS) 96:13223–13228.

Caccone, A., Gentile, G., Gibbs, J.P., Fritts, T.H., Snell, H.L., Betts, J., and Powell, J.R. 2002. Phylogeography and history of giant Galápagos tortoises. Evolution 56(10):2052–2066.

Cadena, E.-A., Scheyer, T.M., Carrillo-Briceño, J.D., Sánchez, R., Aguilera-Socorro, O.A., Vanegas, A., Pardo, M., Hansen, D.M., and Sánchez-Villagra, M.R. 2020. The anatomy, paleobiology, and evolutionary relationships of the largest extinct side-necked turtle. Science Advances 6:eaay4593,1–13.

Cagle, F.R. 1953a. Two new subspecies of *Graptemys pseudogeographica*. Occasional Papers of the Museum of Zoology of the University of Michigan 546:1–17.

Cagle, F.R. 1953b. The status of the turtle *Graptemys oculifera* (Baur). Zoologica 38:137–144.

Cagle, F.R. 1954. Two new species of the genus *Graptemys*. Tulane Studies in Zoology 1:167–186.

Caldwell, D.K. 1962. Sea turtles in Baja California waters (with special reference to those of the Gulf of California), and the description of a new subspecies of northeastern Pacific green turtle. Contributions in Science, Natural History Museum of Los Angeles County 61:1–31.

Călinescu, R. 1931. Contributiuni sistematice si zoogeografice la studiul amphibiilor si reptilelor din Romania. Memoriile Sectiunii Sciintifice Academiei Romane, Bucaresti 3(7):119–291. [In Romanian].

Camp, C.D. 1986. *Sternotherus minor* (Loggerhead Musk Turtle). Size. Herpetological Review 17(4):89–91.

Camper, J.D. 2019. The Reptiles of South Carolina. Columbia, SC: University of South Carolina Press, 273 pp.

Campos, F.S., Moraes, R.L. de, and Pereira, C.S.A. 2011. New state record of *Mesoclemmys perplexa* Bour and Zaher, 2005 (Reptilia: Chelidae) in Brazil. Herpetology Notes 4:263–264.

Cann, J. 1997a. Georges short-necked turtle. Monitor (Journal of the Victorian Herpetological Society) 9(1):18–23, 31–32.

Cann, J. 1997b. The northern yellow-faced turtle. Monitor (Journal of the Victorian Herpetological Society) 9(1):24–29, 31–32, 34–35.

Cann, J. 1997c. Irwin's turtle. Monitor (Journal of the Victorian Herpetological Society) 9(1):36–40, 31–32.

Cann, J. 1997d. Kuchling's turtle. Monitor (Journal of the Victorian Herpetological Society) 9(1):41–44, 32.

Cann, J. 1998. Australian Freshwater Turtles. Singapore: Beaumont Publications, 292 pp.

Cann, J. and Legler, J.M. 1994. The Mary River tortoise: a new genus and species of short-necked chelid from Queensland, Australia (Testudines; Pleurodira). Chelonian Conservation and Biology 1(2):81–96.

Cann, J. and Sadlier, R. 2017. Freshwater Turtles of Australia. Rodeo, NM: ECO Wear and Publishing, 448 pp.

Cann, J., McCord, W.P., and Joseph-Ouni, M. 2003. [*Emydura macquarii emmotti* ssp. nov.] In: McCord, W.P., Cann, J., and Joseph-Ouni, M. A taxonomic assessment of *Emydura* (Testudines: Chelidae) with descriptions of new subspecies from Queensland, Australia. Reptilia (GB) (Barcelona) 27:60–61.

Cantor, T. 1842a. Zoology of Chusan, Part 1. Kolkata [Calcutta]: 32 pp.

Cantor, T. 1842b. General features of Chusan, with remarks on the flora and fauna of that island. Annals and Magazine of Natural History (1)9:265–278, 361–370, 481–493.

Cantor, T. 1847. Catalogue of reptiles inhabiting the Malayan peninsula and islands. Journal of the Asiatic Society of Bengal 16:607–656, 897–952, 1026–1078.

Capinha, C., Seebens, H., Cassey, P., García-Díaz, P., Lenzner, B., Mang, T., Moser, D., Pyšek, P., Rödder, D., Scalera, R., Winter, M., Dullinger, S., and Essl, F. 2017. Diversity, biogeography and the global flows of alien amphibians and reptiles. Diversity and Distributions 23:1313–1322.

Capocaccia, L. 1961. Catalogo dei tipi di rettili del Museo Civico di Storia Naturale di Genova. Annali del Museo Civico di Storia Naturale di Genova 72:86–111.

Cárdenas-Barrantes, M., Matta, N.E., Rodríguez-Fandiño, O.A., Fritz, U., and Vargas-Ramírez, M. 2024. Genetic structure and loss of genetic diversity in the Savannah Side-Necked Turtle *Podocnemis vogli* (Testudines: Podocnemidae). Chelonian Conservation and Biology 23:92–102.

Carlson, L.A. 1999. Aftermath of a feast: human colonization of the southern Bahamian Archipelago and its effects on the indigenous fauna. Ph.D. Thesis, University of Florida, Gainesville.

Carr, A.F., Jr. 1937a. A new turtle from Florida, with notes on *Pseudemys floridana mobiliensis* (Holbrook). Occasional Papers of the Museum of Zoology of the University of Michigan 348:1–7.

Carr, A.F., Jr. 1937b. The status of *Pseudemys scripta* and *Pseudemys troostii*. Herpetologica 1:75–77.

Carr, A.F., Jr. 1938a. *Pseudemys nelsoni*, a new turtle from Florida. Occasional Papers of the Boston Society of Natural History 8:305–310.

Carr, A.F., Jr. 1938b. A new subspecies of *Pseudemys floridana* with notes on the *floridana* complex. Copeia 1938(3):105–109.

Carr, A.F., Jr. 1942. A new *Pseudemys* from Sonora, Mexico. American Museum Novitates 1181:1–4.

Carr, A.F., Jr. 1952. Handbook of Turtles. The Turtles of the United States, Canada, and Baja California. Ithaca, NY: Cornell University Press, 542 pp.

Carr, A.F., Jr. and Hirth, H. 1962. The ecology and migrations of sea turtles. 5. Comparative features of isolated green turtle colonies. American Museum Novitates 2091:1–42.

Carr, A.F., Jr. and Marchand, L.J. 1942. A new turtle from the Chipola River, Florida. Proceedings of the New England Zoology Club 20:95–100.

Carr, J.L. and Mast, R.B. 1988. Natural history observations of *Kinosternon herrerai* (Testudines: Kinosternidae). Trianea (Acta cientifica y tecnologica INDERENA) 1:87–97.

Carr, J.L., Kessler, E.J., and Johnston, G.R. (Eds.). 2023. Biology and Conservation of Alligator Snapping Turtles. Southeastern Naturalist 22 (Special Issue 12): xvi + 456 pp.

Carretero, M.A., Znari, M., Harris, D.J., and Macé, J.C. 2005. Morphological divergence among populations of *Testudo graeca* from west-central Morocco. Animal Biology 55:259–279.

Carrillo de Espinoza, N. and Lamas, G. 1985. Un nuevo registro de tortuga terrestre para el Peru (Reptilia: Testudinata). Publicaciones del Museo de Historia Natural "Javier Prado" Ser. A, Zoologia 31:1–7.

Carter, H.J. 1852. Geology of the Island of Bombay. Journal of the Bombay Branch of the Royal Asiatic Society 4(16):161–215.

Carvalho, J.C. de M. (Ed.). 1972. Alexandre Rodrigues Ferreira. Viagem Filosófica pelas capitanias do Grão-Pará, Rio Negro, Mato Grosso e Cuiabá. Memórias Zoologia, Botânica. Rio de Janeiro, Brazil: Conselho Federal de Cultura, 246 pp.

Carvalho, V.T. de, Martínez, J.G., Hernández-Rangel, S.M., Astolfi-Filho, S., Vogt, R.C., Farias, I.P., and Hrbek, T. 2016. Giving IDs to turtles: SNP markers for assignment of individuals to lineages of the geographically structured *Phrynops geoffroanus* (Chelidae: Testudines). Conservation Genetics Resources, doi:10.1007/s12686-016-0626-8.

Carvalho, V.T. de, Vogt, R.C., Rojas, R.R., Nunes, M.D.S., Fraga, R.D., Ávila, R.W., Rhodin, A.G.J., Mittermeier, R.A., Hrbek, T., and Farias, I.P. 2022. Four in one: cryptic diversity in Geoffroy's Side-Necked Turtle *Phrynops geoffroanus* (Schweigger 1812) (Testudines: Pleurodira: Chelidae) in Brazil. Diversity 14(360):1–16.

Cassano, M.J. 2017. Restricción de la distribución de *Mesoclemmys vanderhaegei* (Bour, 1973) (Testudines: Chelidae) en Argentina. Cuadernos de Herpetología 31:25–27.

Castañeda Gaytán, G. 2020. Periodic review of *Terrapene coahuila*. CITES AC31 Doc. 41.3, 14 pp.

Castillo-Morales, C.A., Sáenz-Arroyo, A., Castellanos-Morales, G., and Ruíz-Montoya, L. 2023. Mitochondrial DNA and local ecological knowledge reveal two lineages of leatherback turtle on the beaches of Oaxaca, Mexico. Scientific Reports 13:8836, 10 pp.

Catesby, M. 1743. The Natural History of Carolina, Florida and the Bahama Islands. Vol. 2. London: 100 pp.

Catesby, M. 1771. The Natural History of Carolina, Florida and Bahama Islands. Folio 2. Edited by G. Edwards. London.

Ceballos, C.P. and Brand, W.A. 2014. Morphology and conservation of the MesoAmerican slider (*Trachemys venusta*, Emydidae) from the Atrato River basin, Colombia. Acta Biológica Colombiana 19(3):483–488.

Ceballos, C.P. and Iverson, J.B. 2014. Patterns of sexual size dimorphism in Chelonia: revisiting Kinosternidae. Biological Journal of the Linnean Society 111:806–809.

Ceballos, C.P., Adams, D.C., Iverson, J.B., and Valenzuela, N. 2013. Phylogenetic patterns of sexual size dimorphism in turtles and their implications for Rensch's Rule. Evolutionary Biology 40:194–208.

Ceballos, C.P., Zapata, D., Alvarado, C., and Rincón, E. 2016. Morphology, diet, and population structure of the Southern White-lipped Mud Turtle *Kinosternon leucostomum postinguinale* (Testudines: Kinosternidae) in the Nus River

drainage, Colombia. Journal of Herpetology 50:374–380.

Ceríaco, L.M.P. and Bauer, A.M. 2017. "*Testudo torticollis*," an unpublished name for the Matamata, *Chelus fimbriatus* (Schneider, 1783) (Testudines: Chelidae), with comments on early illustrations of the species. Bibliotheca Herpetologica 13:35–49.

Ceríaco, L.M.P. and Bour, R. 2012. Schweigger's (1812) chelonian types from the extinct eighteenth century Portuguese "Royal cabinet of natural history of Ajuda": some contributions for their identification and nomenclatural implications. Zootaxa 3395:18–32.

Ceselli, L. 1846. Sopra una tartaruga trovata alle Acque Caje di Viterbo: lettera al sig. Prof. Sebastiano Purgotti. Rome: Salviucci, 26 pp.

Chaianunporn, T., Mokarat, M., Ruangsakunporn, A., Kutanan, W., Lomthaisong, K., Chaianunporn, K., and Keithmaleesatti, S. 2024a. Distribution ranges of Snail-Eating Turtle species (*Malayemys* spp.) in northeastern and eastern Thailand. Chelonian Conservation and Biology 23:216–227.

Chaianunporn, T., Mokarat, M., Ruangsakunporn, A., Walalite, T., Kutanan, W., Lomthaisong, K., Chaianunporn, K., and Keithmaleesatti, S. 2024b. Genetic diversity of the Khorat snail-eating turtle (*Malayemys khoratensis*) in Thailand. PeerJ 12:e18699, 16 pp.

Chan-ard, T., Thirakhupt, K., and van Dijk, P.P. 1996. Observations on *Manouria impressa* at Phu Luang Wildlife Sanctuary, northeastern Thailand. Chelonian Conservation and Biology 2(1):109–113.

Chan-ard, T., Cota, M., Achachak, C., and Safuwong, M. 2012. Measurements of the Big-Headed Turtle (*Platysternon megacephalum* Gray, 1831) (Platysternidae, Testudines) from Phu Luang, Loei Province, Northeastern Thailand. Thailand Natural History Museum Journal 6(2):153–157.

Chandler, C.H. and Janzen, F.J. 2009. The phylogenetic position of the snapping turtles (Chelydridae) based on nucleotide sequence data. Copeia 2009:209–213.

Chang, M.H., Song, J.Y., and Koo, K.S. 2012.The status of distribution for native freshwater turtles in Korea, with remarks on taxonomic position. Korean Journal of Environmental Biology 30(2):151–155.

Chang, M.L.Y. 1957. [Testudoformes]. Science (Ko-xue) (China) 33(1):50. [in Chinese].

Chang, T.-H. 1929. Notes on an apparently new or rarely known hard-shelled turtle from Fuchow. Contributions of the Biological Laboratory of the Science Society of China, Nanking 5(1):1–5.

Charruau, P. and Morales-Garduza, M.A. 2022. Maximum size and mass, and sexual size dimorphism in the Furrowed Wood Turtle, *Rhinoclemmys areolata* (Testudines: Geoemydidae). Herpetology Notes 15:207–210.

Charruau, P., Escobedo-Galván, A.H., and Morales-Garduza, M.A. 2014. *Rhinoclemmys areolata* (Furrowed Wood Turtle). Maximum size and mass. Herpetological Review 45(3):487.

Cheke, A.S. and Hume, J.P. 2008. Lost Land of the Dodo: The Ecological history of the Mascarene Islands. New Haven, Connecticut: Yale University Press.

Cheke, A.S., Pedrono, M., Bour, R., Anderson, A., Griffiths, C., Iverson, J.B., Hume, J.P., and Walsh, M. 2017. Giant tortoises spread to western Indian Ocean islands by sea drift in pre-Holocene times, not by later human agency – response to Wilmé et al. (2016a). Journal of Biogeography 44:1426–1429.

Chemnitz, J.H. 1784. Neues systematisches Conchylien-Cabinet. Siebender Band. Nürnberg: Raspischen Buchhandlung, 356 pp.

Chen, C., Ji, L., Huang, G., Liu, X., Chen, H., Wang, Y., Yu, L., Liu, Y., Hong, X., Wei, C., Wu, C., Luo, L., Zhu, X., and Li, W. 2025. Comparative analysis of the mitochondrial genomes of the softshelled turtles *Palea steindachneri* and *Pelodiscus axenaria* and phylogenetic implications for Trionychia. Scientific Reports 15:7138, 16 pp.

Chen, H.G., Liu, W.B., and Zhang, X.J. 2005. Comparative analysis of mitochondrial DNA 12S rRNA region between *Pelodiscus sinensis* and *Pelodiscus axenaria* and their molecular marker for identification. Journal of Fisheries China 29:318–322.

Chen, H.G., Liu, W.B., Li, J.Z., and Zhang, X.J. 2006. Comparative analysis of mitochondrial DNA cytochrome b gene and their molecular identification markers in three species of soft-turtles. Shuisheng Shengwu Xuebao 30:380–385.

Chen, T.-H. and Lue, K.-Y. 2010. Population status and distribution of freshwater turtles in Taiwan. Oryx 44(2):261–266.

Chesi, F. 2009. Il registro fossile italiano dei cheloni. Ph.D. Thesis, Università di Firenze, Italy.

Chevalier, A. 1935. Les îles du Cap Vert. Géographie, biogéographie, agriculture. Flore de l'archipel. Revue de Botanique Appliquée et d'Agriculture Tropicale, Paris 15:733–1090.

Cheylan, M. 2001. *Testudo hermanni* Gmelin, 1789 - Griechische Landschildkröte. In: Fritz, U. (Ed.). Handbuch der Reptilien und Amphibien Europas, Band 3/IIIA. Wiesbaden: Aula–Verlag, pp. 179–289.

Chiari, Y. 2020. Morphology. In: Gibbs, J.P., Cayot, L.J., and Tapia Aguilera, W. (Eds.). Galapagos Giant Tortoises. London: Academic Press, pp. 139–156.

Chiari, Y., Thomas, M., Pedrono, M., and Vieites, D.R. 2005. Preliminary data on genetic differentiation within the Madagascar spider tortoise, *Pyxis arachnoides* (Bell, 1827). Salamandra 41:35–43.

Chiari, Y., Hyseni, C., Fritts, T.H., Glaberman, S., Marquez, C., Gibbs, J.P., Claude, J., and Caccone, A. 2009. Morphometrics parallel genetics in a newly discovered and endangered taxon of Galápagos tortoise. PLoS One 4(7):e6272.

Chiocchio, A., Zampiglia, M., Biaggini, M., Biello, R., Di Tizio, L., Leonetti, F.L., Olivieri, O., Sperone, E., Trabalza-Marinucci, M., Corti, C., and Canestrelli, D. 2022. Unveiling a hotspot of genetic diversity in southern Italy for the endangered Hermann's tortoise *Testudo hermanni*. BMC Ecology and Evolution 22:131, 10 pp.

Chirio, L. and Ineich, I. 2006. Biogeography of the reptiles of the Central African Republic. African Journal of Herpetology 55(1):23–59.

Chkhikvadze, V.M. 1970. [On the origin of the modern Palaearctic land tortoises]. Soobshcheniya Akademii Nauk Gruzinskoi SSR [Bulletin of the Academy of Sciences of Georgia] 57(1):245–247. [in Russian].

Chkhikvadze, V.M. 1983. [Les tortues fossiles du Caucase et du Nord de la Mer Noire]. Tbilisi: Metzniereba, 149 pp. [in Russian].

Chkhikvadze, V.M. 1987. [On systematic position of the USSR Far East softshelled turtle.]. Soobshcheniya Akademii Nauk Gruzinskoi SSR [Bulletin of the Academy of Sciences of Georgia] 128(3):609–611. [in Russian].

Chkhikvadze, V.M. 1988. O sistematicheskom polozhenii sobremennykh sukhoputnykh cherepakh srednei Azii i Kazakhstana. [Taxonomic status of modern land tortoise of Middle Asia and Kazakhstan]. Soobshcheniya Akademii Nauk Gruzinskoi SSR [Bulletin of the Academy of Sciences of Georgia] 14(2):110–114. [in Russian].

Chkhikvadze, V.M. 1989. Neogenovye cherepakhi SSSR. [Neogene turtles of the USSR]. Tbilisi: Metsniereba, 104 pp. [in Russian].

Chkhikvadze, V.M. 2008. [*Agrionemys bogdanovi*]. In: Chkhikvadze, V.M., Brushko, Z.K., and Kubykin, R.A. Краткий обзор систематики среднеазиатских черепах (Testudinidae: *Agrionemys*) и подвижные зоны панциря у этой группы черепах. [A brief overview of the systematics of the Central Asian tortoise (Testudinidae: *Agrionemys*) and mobile shell zone in this group of turtles]. Selevinia (Almaty) 2008:100–104. [in Russian].

Chkhikvadze, V.M. 2009. Среднеазиатская черепахиа в Монголии. [Central Asiatic tortoises in Mongolia]. Problems of Desert Development (Ashgabat) 2009(3/4):60–63. [in Russian].

Chkhikvadze, V.M. and Bakradze, M.A. 1991. [On the systematic position of the Recent land turtle from the Araxes Valley]. Trudy Tbilisskogo Gosudarstvennogo Universitet Tbilisi [Proceedings of Tbilisi University] 305:59–63. [in Russian].

Chkhikvadze, V.M. and Bakradze, M.A. 2002. Novyi podvid sukhoputnoi cherepakhi iz Dagestana. [A new subspecies of land tortoises of Daghestan]. Trudy Instituta Zoologii Akademia Nauk Gruzii [Proceedings of the Zoology Institute of the Georgia Academy of Sciences] 21:276–279. [in Russian].

Chkhikvadze, V.M. and Tuniyev, B.S. 1986. [On the taxonomic status of modern land tortoise of the western Transcaucasus]. Soobshcheniya Akademii Nauk Gruzinskoi SSR [Bulletin of the Academy of Sciences of Georgia] 124(3):617–620. [in Russian].

Chkhikvadze, V.M., Amiranashvili, N.G., and Ataev, C.A. 1990. Noviy podvid sukhoputnoi cherepakhi iz yugo–zapadnovo Turkmenistana. [A new subspecies of tortoise from the southwestern Turkmenistan]. Izvestiya Akademii Nauk Turkmenskoi SSR, Seriya Biologicheskie Nauki 1:72–75. [in Russian].

Chkhikvadze, V.M., Ataev, C.A., Shammakov, S., and Zatoka, A.L. 2009. [*Agrionemys kazachstanica kuznetzovi*]. In: Chkhikvadze, V.M., Ataev, C.A., and Shammakov, S. [New taxons of Central Asian tortoises (Testudinidae: *Agrionemys bogdanovi* and *A. kazachstanica kuznetzovi*)]. Problems of Desert Development (Ashgabat) 2009(1/2):49–54. [in Russian].

Chkhikvadze, V.M., Bondarenko, D.A., and Shammakov, S. 2010. Shell morphology of Central Asian tortoise (*Agrionemys horsfieldii*) in southeastern Turkmenistan and northern Iran, and taxonomic status of *Agrionemys* genus. Current Studies in Herpetology 10:40–46. [in Russian]

Chkhikvadze, V.M., Mazanaeva, L.F., and Shammakov, S.M. 2011. A short account

of a new species of land tortoise in Dagestan. Proceedings of the International Conference 'Biological Diversity and Conservation Problems of the Fauna of the Caucasus'. Yerevan, Armenia: National Academy of Sciences of the Republic of Armenia, pp. 336–340. [in Russian].

Chkhikvadze, V.M., Mazanaeva, L.F., and Kvachadze, T.O. 2013. [Terrestrial turtles of the Caucasus and North-West Iran]. Buletin ştiinţific–Revistă de Etnografie, Ştiinţele Naturii şi Muzeologie (Ştiinţele Naturii). Serie nouă 18(31):72–86. [in Russian]

Chu Cheong, L. 1990. Observations on the nesting population of leatherback turtles, *Dermochelys coriacea*, in Trinidad. Caribbean Marine Studies 1:48–53.

Çilingir, F.G., Hansen, D., Bunbury, N., Postma, E., Baxter, R., Turnbull, L., Ozgul, A., and Grossen, C. 2022a. Low-coverage reduced representation sequencing reveals subtle within-island genetic structure in Aldabra giant tortoises. Ecology and Evolution 12:e8739, 13 pp.

Çilingir, F.G., A'Bear, L., Hansen, D., Davis, L.R., Bunbury, N., Ozgul, A., Croll, D., and Grossen, C. 2022b. Chromosome-level genome assembly for the Aldabra giant tortoise enables insights into the genetic health of a threatened population. GigaScience 11:p.giac090.

Ciofi, C., Wilson, G.A., Beheregaray, L.B., Marquez, C., Gibbs, J.P., Tapia, W., Snell, H.L., Caccone, A., and Powell, J.R. 2006. Phylogeographic history and gene flow among giant Galápagos tortoises on southern Isabela Island. Genetics 172:1727–1744.

CITES. 2010. Interpretation and implementation of the Convention. Trade control and marking. Standard nomenclature. CITES Document CoP15 Doc. 35, 33 pp.

Clausen, C.J., Cohen, A.D., Emiliani, C., Holman, J.A., and Stipp, J.J. 1979. Little Salt Spring, Florida: a unique underwater site. Science 203:609–614.

Cliburn, J.W. 1971. The ranges of four species of *Graptemys* in Mississippi. Journal of the Mississippi Academy of Science 16:16–19.

Close, L.M. and Seigel, R.A. 1997. Differences in body size among populations of red-eared sliders (*Trachemys scripta elegans*) subjected to different levels of harvesting. Chelonian Conservation and Biology 2(4):563–566.

Clostio, R.W., Martinez, A.M., LeBlanc, K.E., and Anthony, N.M. 2012. Population genetic structure of a threatened tortoise across the south-eastern United States: implications for conservation management. Animal Conservation 15(6):613–625.

Cochran, D.M. 1961. Type specimens of reptiles and amphibians in the U.S. National Museum. Bulletin of the U.S. National Museum 220:1–291.

Cocteau, J.T. and Bibron, G. 1838. Reptilia. In: Sagra, D.R. de la. Historia Fisica, Politica y Natural de la Isla de Cuba. Vol. 4. Reptiles y Peces, pp. 1–143.

Cogger, H.G. 1979. Type specimens of reptiles and amphibians in the Australian Museum. Records of the Australian Museum 32:163–210.

Cogger, H.G. 2018. Reptiles and Amphibians of Australia. Updated Seventh Edition. Clayton South, Victoria: CSIRO Publishing, 1096 pp.

Cogger, H.G., Cameron, E.E., and Cogger, H.M. 1983. Zoological Catalogue of Australia. Volume 1. Amphibia and Reptilia. Canberra: Australian Government Publishing Service, 313 pp.

Coleman, A.T. 2020. Urban Turtle Project: using citizen science to document freshwater turtle communities and populations in Birmingham, Alabama, with focus on Alabama Map Turtles (*Graptemys pulchra*). Chelonian Conservation and Biology 19(2):283–290.

Collins, J.T., Collins, S.L., and Taggart, T.W. 2010. Amphibians, Reptiles, and Turtles in Kansas. Eagle Mountain, UT: Eagle Mountain Publishing, 312 pp.

Colombo, W.D., Justino, J.D.F., Barcelos, A.C., Vilaça, S.T., Pavanelli, L., and Vargas, S.M. 2024. Reassessing leatherback turtle lineages and unveiling the first evidence of nuclear mitochondrial DNA in sea turtles. Scientific Reports 14:31313, 8 pp.

Conant, R. 1958. A Field Guide to Reptiles and Amphibians of the United States and Canada East of the 100th Meridian. Boston: Houghton Mifflin Co., 366 pp.

Conant, R. and Goin, C.J. 1948. A new subspecies of soft-shelled turtle from the central United States, with comments on the application of the name *Amyda*. Occasional Papers of the Museum of Zoology, University of Michigan 510:1–19.

Conix, S., Garnett, S.T., Thiele, K.R., Christidis, L., van Dijk, P.P., Bánki, O.S., Barik, S.K., Buckeridge, J.S., Costello, M.J., Hobern, D., Kirk, P.M., Lien, A., Nikolaeva, S., Pyle, R.L., Thomson, S.A., Zhang, Z., and Zachos, F.E. 2021. Towards a global list of accepted species III. Independence and stakeholder inclusion. Organisms Diversity & Evolution: doi.org/10.1007/s13127-021-00496-x, 13 pp.

Contreras-Calvario, A.I., Mora-Reyes, A., Fuentes-Moreno, A., Arreortúa, M., Cázares-Hernández, E., Lara-Hernández, F.A., and López-Luna, M.A. 2024. New distributional records for an endangered turtle, *Claudius angustatus* (Reptilia, Testudines), and notes on its threats in Veracruz, Mexico. Revista Latinoamericana de Herpetología 7(2):23–28.

Converse, P.E., Kuchta, S.R., Hauswaldt, J.S., and Roosenburg, W.M. 2017. Turtle soup, Prohibition, and the population genetic structure of Diamondback Terrapins (*Malaclemys terrapin*). PLoS ONE 12(8):e0181898, 18 pp.

Cook, S., Abb, D., and Frair, W. 1972. A new record size box turtle. International Turtle and Tortoise Society Journal 6(3):9–17.

Cooper, J.G. 1861. New Californian animals. Proceedings of the California Academy of Sciences, San Francisco 2:118–123.

Cope, E.D. 1860. Notes and descriptions of foreign reptiles. Proceedings of the Academy of Natural Sciences, Philadelphia 11:294–297.

Cope, E.D. 1864. On the limits and relations of the Raniformes. Proceedings of the Academy of Natural Sciences, Philadelphia 16:181–183.

Cope, E.D. 1865. Third contribution to the herpetology of tropical America. Proceedings of the Academy of Natural Sciences, Philadelphia 17:185–198.

Cope, E.D. 1868. An examination of the Reptilia and Batrachia obtained by the Orton expedition to Equador and the upper Amazon, with notes on other species. Proceedings of the Academy of Natural Sciences, Philadelphia 20:96–140.

Cope, E.D. 1869 ["1868"]. On the origin of genera. Proceedings of the Academy of Natural Sciences, Philadelphia 20:242–300.

Cope, E.D. 1870a. Seventh contribution to the herpetology of tropical America. Proceedings of the American Philosophical Society 11(1869)[1870]:147–169.

Cope, E.D. 1870b. Synopsis of the extinct Batrachia, Reptilia and Aves of North America. Transactions of the American Philosophical Society, new series 14(1869)[1870]:1–252.

Cope, E.D. 1871. On the homologies of some of the cranial bones of the Reptilia, and on the systematic arrangement of the class. Proceedings of the American Association for the Advancement of Science 1870(19)[1871]:194–247.

Cope, E.D. 1872. Synopsis of the species of the Chelydrinae. Proceedings of the Academy of Natural Sciences, Philadelphia 1872:22–29.

Cope, E.D. 1875a. On the Batrachia and Reptilia of Costa Rica. In: Cope, E.D. (Ed.). On the Batrachia and Reptilia of Costa Rica with notes on the Herpetology and Ichthyology of Nicaragua and Peru. Journal of the Academy of Natural Sciences, Philadelphia (2)8(4)1875:93–154.

Cope, E.D. 1875b. Check-list of North American Batrachia and Reptilia; with a systematic list of the higher groups, and an essay on geographical distribution. Bulletin of the United States National Museum 1:1–104.

Cope, E.D. 1878. Description of new Vertebrata from the upper Tertiary formations of the West. Proceedings of the American Philosophical Society 17:219–231.

Cope, E.D. 1885. A contribution to the herpetology of Mexico. Proceedings of the American Philosophical Society 22:379–404.

Cope, E.D. 1887. Catalogue of Batrachia and Reptilia of Central America and Mexico. Bulletin of the U.S. National Museum 32:1–98.

Cope, E.D. 1892a. A contribution to the vertebrate paleontology of Texas. Proceedings of the American Philosophical Society 30:123–131.

Cope, E.D. 1892b. A contribution to the knowledge of the fauna of the Blanco beds of Texas. Proceedings of the Academy of Natural Sciences, Philadelphia 1892:226–229.

Cope, E.D. 1893. A preliminary report on the vertebrate paleontology of the Llano Estacado. Annual Reports of the Geological Survey of Texas 4:1–136.

Cope, E.D. 1895. Taylor on box turtles. American Naturalist 29:756–757.

Cope, E.D. 1899. Vertebrate remains from Port Kennedy bone deposit. Journal of the Academy of Natural Sciences, Philadelphia (2)11:193–267.

Cornalia, E. 1849. Vertebratorum Synopsis in Museo Mediolanense Extantium quae per novam orbem Cajetanus Osculati collegit annis 1846-47-1848 speciebus novis vel minus cognitis adjectis nec non descriptionibus atque iconibus illustratis. Modoetiae: Typographia Corbetta, 15 pp.

Cortés-Rodríguez, X., Borja-Martínez, G., and Vázquez-Domínguez, E. 2021. Striking habitat reduction, decreased genetic diversity, and imperilled conservation of natural populations of *Terrapene coahuila*. Freshwater Biology 66:842–858.

Costa, H.C., Rezende, D.T. de., Molina, F.B., Nascimento, L.B., Leite, F.S.F., and Fernandes, A.P.B. 2015. New distribution records and potentially suitable areas for the threatened Snake-Necked Turtle *Hydromedusa maximiliani* (Testudines: Chelidae). Chelonian Conservation and Biology 14(1):88–94.

Cotton, W.B. 1918. Hardwicke's Tortoise (*T. hardwickei*). In: Wallace, J. Proceedings of the meeting held on 16th April 1918. Journal of the Bombay Natural History Society 26:313–314.

Coutard, C. 2006. Note sur la maintenance et la reproduction en captivité de *Testudo marginata* Schoepff, 1795 et *Testudo weissingeri* (Bour, 1995). Manouria 9(31):18–26.

Coutinho, J.M. da Silva. 1868. Sur les tortues de l'Amazone. Bulletin de la Société Impériale Zoologique d'Acclimatation (2)5:147–166.

Covacevich, J. 1971. Amphibian and reptile type-specimens in the Queensland Museum. Memoirs of the Queensland Museum 16(1):49–67.

Cowan, C.F. 1969. Notes on Griffith's Animal Kingdom of Cuvier (1824-1835). Journal of the Society of Bibliography of Natural History 5:137–140.

Cox, M.J., van Dijk, P.P., Nabhithabhata, J., and Thirakhupt, K. 1998. A Photographic Guide to Snakes and Other Reptiles of Peninsular Malaysia, Singapore and Thailand. London: New Holland Publishers, 144 pp.

Cox, N., Young, B.E., Bowles, P., Fernandez, M., Marin, J., Rapacciuolo, G., Böhm, M., Brooks, T.M., Hedges, S.B., Hilton-Taylor, C., Hoffmann, M., Jenkins, R.K.B., Tognelli, M.F., Alexander, G.J., Allison, A., Ananjeva, N.B., Auliya, M., Avila, L.J., Chapple, D.G., Cisneros-Heredia, D.F., Cogger, H.G., Colli, G.R., De Silva, A., Eisemberg, C.C., Els, J., G., A.F., Grant, T.D., Hitchmough, R.A., Iskandar, D.T., Kidera, N., Martins, M., Meiri, S., Mitchell, N.J., Molur, S., Nogueira, C.D.C., Ortiz, J.C., Penner, J., Rhodin, A.G.J., Rivas, G., Rödel, M.-O., Roll, U., Sanders, K.L., Santos-Barrera, G., Shea, G.M., Spawls, S., Stuart, B.L., Tolley, K.A., Trape, J.-F., Vidal, M.A., Wagner, P., Wallace, B.P., and Xie, Y. 2021. Global reptile assessment shows commonality of tetrapod conservation needs. In review.

Cragin, F.W. 1894. Herpetological notes from Kansas and Texas. Colorado College Studies 5:37–39.

Crawford, N.G., Faircloth, B.C., McCormack, J.E., Brumfield, R.T., Winker, K., and Glenn, T.C. 2012. More than 1000 ultraconserved elements provide evidence that turtles are the sister group of archosaurs. Biology Letters 8(5):783–786.

Crawford, N.G., Parham, J.F., Sellas, A.B., Faircloth, B.C., Glenn, T.C., Papenfuss, T.J., Henderson, J.B., Hansen, M.H., and Simison, W.B. 2015. A phylogenomic analysis of turtles. Molecular Phylogenetics and Evolution 83:250–257.

Crippen, R.G. 1962. Holotype specimens of amphibians and reptiles in the Museum of Vertebrate Zoology, University of California, Berkeley. Herpetologica 18:187–194.

Crucitti, P. and Emiliani, F. 2012. Morphological variability of the Spurthighed Tortoise, *Testudo graeca*, in the Nemrut Volcano (Eastern Turkey) (Testudines: Testudinidae). Zoology in the Middle East 56:19–26.

Crumly, C.R. 1994. Phylogenetic systematics of North American tortoises (genus *Gopherus*): evidence of their classification. In: Bury, R.B. and Germano, D.J. (Eds.). Biology of North American Tortoises. U.S. Fish and Wildlife Research 12:7–32.

Cuadrado-Ríos, S., Vargas-Ramírez, M., Kehlmaier, C., and Fritz, U. 2025. Complete mitochondrial genome and phylogenetic analysis of the Atrato slider, *Trachemys medemi* (Testudines, Emydidae). ZooKeys 1224:253–260.

Cunha, F.A.G., Fernandes, T., Franco, J., and Vogt, R.C. 2019. Reproductive biology and hatchling morphology of the Amazon Toad-headed Turtle (*Mesoclemmys raniceps*) (Testudines: Chelidae), with notes on species morphology and taxonomy of the *Mesoclemmys* group. Chelonian Conservation and Biology 18(2):195–209.

Cunha, F.A.G., Fagundes, C.K., Brito, E.S., Vogt, R.C., Maffei, F., Pezzuti, J., Félix-Silva, D., Rojas-Runjaic, F.J.M., Lasso, C.A., Morales-Betancourt, M.A., Carvalho, V.T. de, Amaral, J.V. do, Balestra, R.A.M., Acácio, M., Malvasio, A., and Lustosa, A.P.G. 2021. Distribution of *Chelus fimbriata* and *Chelus orinocensis* (Testudines: Chelidae). Chelonian Conservation and Biology 20(1):109–115.

Cunha, F.A.G., Sampaio, I., Carneiro, J., and Vogt, R.C. 2021. A new species of Amazon freshwater toad-headed turtle in the genus *Mesoclemmys* (Testudines: Pleurodira: Chelidae) from Brazil. Chelonian Conservation and Biology 20(2):151–166.

Cunha, F.A.G., Sampaio, I., Carneiro, J., Vogt, R.C., Mittermeier, R.A., Rhodin, A.G.J., and Andrade, M.C. 2022. A new South American freshwater turtle of the genus *Mesoclemmys* from the Brazilian Amazon (Testudines: Pleurodira: Chelidae). Chelonian Conservation and Biology 21(2):158–180.

Cunha, F.A.G., Pinheiro, R.T., de Freitas, A.P., and Costa de Campos, C.E. 2025 ["2024"]. *Mesoclemmys nasuta* (Guyanan Toad-headed Turtle). Maximum size. Herpetological Review 55(2)[2024]:252.

Cunha, F.A.G., Da Silva, T.L., and Vasconcelos, V.S. 2025 ["2024"]. *Mesoclemmys raniceps* (Amazon Toad-headed Turtle). Phenotypic change. Herpetological Review 55(3)[2024]:426–428.

Cunha, O.R. da. 1970. Uma nova subespécie de quelônio, *Kinosternon scorpioides carajasensis* da Serra dos Carajás, Pará. Boletim do Museo Paraense Emílio Goeldi 73:1–11.

Cuvier, G.L.C.F.D. 1816 ["1817"]. Le Règne Animal Distribué d'après son Organisation, pour Servir de Base à l'Histoire Naturelle des Animaux et d'Introduction à l'Anatomie Comparée. [Edition 1]. Tome II. Contenant les Reptiles, les Poissons, les Mollusques et les Annélides. Paris: Deterville, 532 pp.

Cuvier, G.L.C.F.D. 1825 ["1824"]. Recherches sur les Ossemens Fossiles, où l'on rétablit les caractères de plusieurs animaux dont les révolutions du globe ont détruit les espèces. Nouvelle édition, entièrement refondue, et considérablement augmentée. Tome cinquième, IIe. partie, contenant les ossemens de reptiles et le résumé général. Paris: Dufour et d'Ocagne, 547 pp.

Cuvier, G.L.C.F.D. 1829. Le Règne Animal Distribué d'après son Organisation, pour Servir de Base à l'Histoire Naturelle des Animaux et d'Introduction à l'Anatomie Comparée. Nouvelle Édition, Revue et Augmentée [Edition 2]. Tome II. Paris: Deterville, 406 pp.

D'Amato, A.F. and Morato, S.A.A. 1991. Notas biológicas e localidades de registro de *Platemys spixii* (Duméril & Bibron, 1835) (Testudines: Chelidae) para o Estado do Paraná, Brasil. Acta Biologica Leopoldensia 13(2):119–130.

Dahl, G. and Medem, F. 1964. Informe sobre la fauna acuatica del rio Sinu. Parte II. Los reptiles acuaticos de la hoya del Sinu. Bogota: Corporacion Autonoma Regional de los Valles del Magdalena y del Sinu–CVM, Departamento de Investigaciones Ictiologicas y Faunisticas, pp. 110–152.

Dajčman, U., Hofmeyr, M.D., Anunciação, P.R., Ihlow, F., and Vamberger, M. 2021. Tortoise forensics: conservation genetics of the leopard tortoise *Stigmochelys pardalis* in southern Africa. Salamandra 57(1):139–145.

Dalman, J.W. 1820. Underrättelse om några sköldpadd-skal, som blifvit funna vid gräfning af Götha Canal. [Notification regarding some turtle shells, found while digging Götha Canal.] Kongliga Vetenskaps Academiens Handlingar, Stockholm (3)8:286–293. [in Swedish]

Daniels, S.R., Hofmeyr, M.D., Henen, B.T., and Crandall, K.A. 2007. Living with the genetic signature of Miocene induced change: evidence from the phylogeographic structure of the endemic angulate tortoise *Chersina angulata*. Molecular Phylogenetics and Evolution 45:915–926.

Daniels, S.R., Hofmeyr, M.D., Henen, B.T., and Baard, E.H.W. 2010. Systematics and phylogeography of a threatened tortoise, the speckled padloper. Animal Conservation 13:237–246.

Danilov, I.G. and Parham, J.F. 2006. A redescription of '*Plesiochelys*' *tatsuensis*, a turtle from the Late Jurassic of China, and its bearing on the antiquity of the crown clade Cryptodira. Journal of Vertebrate Paleontology 26:573–580.

Danilov, I.G., Cherepanov, G.O., and Vitek, N.S. 2013. Chelonological studies of L.I. Khosatzky with his annotated bibliography on turtles. Proceedings of the Zoological Institute Russian Academy of Sciences 317(4):382–425.

Dao, V.T. 1957. [Rapport sur les recherches zoologiques dans la région de Vinh-Linh (Province de Quang-Tri, centre Vietnam)]. Zoologicheskii Zhurnal 36(8):1209–1216. [in Russian].

Darevsky, I.S. and Mertens, R. 1973. Zwei unveröffentlichte Schildkrötentafeln von Pallas. Salamandra 9(3/4):99–102.

Das, A., Basu, D., Converse, L., and Choudhury, S.C. 2012. Herpetofauna of Katerniaghat Wildlife Sanctuary, Uttar Pradesh, India. Journal of Threatened Taxa 4(5):2553–2568.

Das, I. 1994. The identity of the Plio-Pleistocene turtle, *Geoemyda pilgrimi* Prasad and Satsangi, 1967 (Testudines: Cryptodira: Bataguridae). Hamadryad 19:41–46.

Das, I. 1995. Turtles and Tortoises of India. Bombay: Oxford University Press, 176 pp.

Das, I. 2001. Die Schildkröten des Indischen Subkontinents. Frankfurt am Main: Chimaira, pp. 181.

Das, I. 2005. Asian turtles in the collection of the Oxford University Museum, including a list of type specimens. Herpinstance 3:4–7.

Das, I. 2009. The discovery of Indian turtles, with notes on publications, type localities and type repositories. In: Vasudevan, K. (Ed.). Freshwater Turtles and Tortoises of India. ENVIS Bulletin: Wildlife and Protected Areas 12(1):1–14.

Das, I., Dattagupta, B., and Gayen, N.C. 1998. History and catalogue of reptile types in the collection of the Zoological Survey of India. Journal of South Asian Natural History 3:121–172.

Daudin, F.M. 1801. Histoire Naturelle, Générale et Particulière des Reptiles.

Tome Second. Paris: Imprimerie F. Dufart, 432 pp.

Daudin, F.M. 1802. Histoire Naturelle, Générale et Particulière des Reptiles. Tome Quatrième. Paris: Imprimerie F. Dufart, 397 pp.

Davenport, J., Holland, D.L., and East, J. 1990. Thermal and biochemical characteristics of the lipids of the leatherback turtle *Dermochelys coriacea*: evidence of endothermy. Journal of the Marine Biological Association of the United Kingdom 70(1):33–41.

David, A. 1875. Journal de mon Troisième Voyage d'Exploration dans l'Empire Chinois. Tome Second. Paris: Hachette, 348 pp.

David, P. 1994. Liste des reptiles actuels du monde. I. Chelonii. Dumerilia 1:7–127.

Davis, J.G., Lipps, G.J., Jr., Wynn, D., Armitage, B.J., Matson, T.O., Pfingsten, R.A., and Caldwell, C. (Eds.). 2021. Reptiles of Ohio. Ohio Biological Survey Bulletin, New Series 20(1) Part I, 405 pp.

Davy, C.M. and Murphy, R.W. 2014. Conservation genetics of the endangered Spotted Turtle (*Clemmys guttata*) illustrate the risks of "bottleneck tests". Canadian Journal of Zoology 92:149–162.

Davy, C.M., Bernardo, P.H., and Murphy, R.W. 2014. A Bayesian approach to conservation genetics of Blanding's turtle (*Emys blandingii*) in Ontario, Canada. Conservation Genetics 15:319–330.

Dawson, J.E., Gaillard, D., Gong, S., Lin, L., McCormack, T.E.M., Nanthavong, C., Nguyen, T.T., Nguyen, T.Q., Pham, T.V., and Shi, H. 2022. Geographical and elevational distributions of the Black-breasted Leaf Turtle, *Geoemyda spengleri* (Gmelin, 1789) (Testudines: Geoemydidae). Amphibian & Reptile Conservation 16(1):235–244(e312).

Dawson, J.E., Anders, B., Bettelheim, M., and Ihlow, F. 2025. *Damonia crassiceps* Gray, 1870, a junior synonym of *Mauremys nigricans* (Gray, 1834) (Testudines: Geoemydidae). Zootaxa 5609(3):445–450.

de Bruin, R.W.F. and Artner, H.G. 1999. On the turtles of Hainan Island, southern China. Chelonian Conservation and Biology 3(3):479–486.

de Rooij, N. 1915. The Reptiles of the Indo-Australian Archipelago. I. Lacertilia, Chelonia, Emydosauria. Leiden: E.J. Brill, 334 pp.

de Silva, A. 2003. The biology and status of the star tortoise in Sri Lanka. Gampola: Ministry of Environment and Natural Resources, AMP Print Shop, 100 pp.

de Silva, A., Wijeratne, R., Rodrigo, K., Hettiarachchi, H.A.H.R., and Prasad, G.A.T. 2017. The giant star tortoise (*Geochelone elegans*) of Lunugamwehera National Park, Sri Lanka. Wildlanka 5:31–35.

de Silva, A., Vamberger, M., Karunarathna, S., Ukuwela, K., Weerasinghe, K.G.S., Bandara, M.A.C.W., Silva, P.D., Gayan, A., and Ranasinghe, B. 2022. The last chance: temple and privately owned ponds as an alternative for conservation of fresh water terrapins in Sri Lanka: preliminary observations. Wildlanka 10:101–108.

De Vis, C.W. 1897. The extinct freshwater turtles of Queensland. Annals of the Queensland Museum 3:1–7.

Dean, R.H. and Bickham, J.W. 1983. *Staurotypus salvini*. Catalogue of American Amphibians and Reptiles 327:1–2.

deBraga, M. and Rieppel, O. 1997. Reptile phylogeny and the interrelationships of turtles. Zoological Journal of the Linnean Society 120:281–354.

Deepak, V. and Vasudevan, K. 2009. Endemic turtles of India. In: Vasudevan, K. (Ed.). Freshwater Turtles and Tortoises of India. ENVIS Bulletin: Wildlife and Protected Areas 12(1):25–42.

Degenhardt, W.G., Painter, C.W., and Price, A.H. 1996. Amphibians and Reptiles of New Mexico. Albuquerque: University of New Mexico Press, 431 pp.

de la Fuente, M., Lapparent de Broin, F. de, and Bianco, T.M. 2001. The oldest and first nearly complete skeleton of a chelid, of the *Hydromedusa* sub-group (Chelidae, Pleurodira), from the Upper Cretaceous of Patagonia. Bulletin de la Société Géologique de France 172(2):237–244.

del Barco, D.M. and Larriera, A. 1993. Sobre la validez de las subespecies de *Trachemys dorbignyi* y su distribución geographica. Revista Asociacion Ciencias Naturales Litoral 22(2)(1991):11–17.

Denzer, W. and Kaiser, H. 2023. Naming and gaming: the illicit taxonomic practice of 'nomenclatural harvesting' and how to avoid it. Journal of Zoology 320:161–168.

Denzer, W. and Kaiser, H. 2025. Nomenclature protects stability only when supported by good science: a response to Frétey's (2024) list of turtle genus-series nomina. Bionomina 41:20–24.

Deraniyagala, P.E.P. 1933. The loggerhead turtles (Carettidae) of Ceylon. Ceylon Journal of Science 18B:59–70.

Deraniyagala, P.E.P. 1934. Relationships among loggerhead turtles (Carettidae). Ceylon Journal of Science 18B:207–208.

Deraniyagala, P.E.P. 1939. The Tetrapod Reptiles of Ceylon. Volume 1. Testudinates and Crocodilians. London: Dulau Co., 412 pp.

Deraniyagala, P.E.P. 1948. Some scientific results of two visits to Africa. Spolia Zeylanica 25(2):1–42.

Deraniyagala, P.E.P. 1952. The loggerhead turtles (Carettinae) of Europe. Herpetologica 8:57–58.

Deraniyagala, P.E.P. 1953. A Colored Atlas of Some Vertebrates from Ceylon. Vol. 2. Tetrapod Reptilia. Ceylon Natural History Museum Publication, 101 pp.

DeSola, C.R. 1930. The Liebespiel of *Testudo vandenburghi*, a new name for the mid-Albemarle Island Galapagos tortoise. Copeia 1930(3):79–80.

Devaux, B. 2000. La Tortue qui Pleure – The Crying Tortoise, *Geochelone sulcata* (Miller, 1779). Chelonii 1:1–87.

Devaux, B. 2003. Namibie, Afrique du Sud; des tortues d'exception! La Tortue 63/64:40–56.

Devaux, B. 2007. La tortue géante de Seychelles, une survivante: *Dipsochelys elephantina* (Duméril et Bibron, 1835). Chelonii 5:1–120.

Devi, N.A., Eprilurahman, R., Yudha, D.S., Raharjo, S., As-Singkily, M., Gunalen, D., and Arida, E. 2021. Genetic diversity and species identity of the critically endangered Rote Island snake-necked turtle, *Chelodina mccordi* Rhodin, 1994. IOP Conference Series: Earth and Environmental Science 948: 012001, 10 pp.

Diesing, C.M. 1839. Neue Gattungen von Binnenwurmern nebst einem Nachtrage zur Monographie der Amphistomen. Annalen des Wiener Museums der Naturgeschichte 2:219–242.

Diesmos, A.C., Parham, J.F., Stuart, B.L., and Brown, R. 2005. The phylogenetic position of the recently rediscovered Philippine forest turtle (Bataguridae: *Heosemys leytensis*). Proceedings of the California Academy of Sciences 56:31–41.

Diesmos, A.C., Brown, R.M., Alcala, A.C., and Sison, R.V. 2008. Status and distribution of nonmarine turtles of the Philippines. Chelonian Conservation and Biology 7:157–177.

Dixon, J.R. and Soini, P. 1977. The reptiles of the upper Amazon basin, Iquitos region, Peru. II. Crocodilians, turtles and snakes. Milwaukee Public Museum Contributions in Biology and Geology 12:1–91.

Djurakic, M.R. and Milankov, V.R. 2019. Carapace shape variation of genetically divergent populations of *Testudo hermanni boettgeri* (Reptilia: Testudines). Archives of Biological Science 71:609–619.

Djurakic, M.R. and Milankov, V.R. 2020. The utility of plastron shape for uncovering cryptic diversity in Hermann's Tortoise. Journal of Zoology 310:145–157.

do Valle, M.F., Marques, R., and Tinôco, M.S. 2016. Filling gaps in the disjunct distribution of *Rhinoclemmys punctularia* (Daudin, 1801) (Testudines: Geoemydidae) and first record from Bahia state, northeast Brazil. Check List 12(4):1951, 4 pp.

Dodd, C.K., Jr. 1988. Synopsis of the biological data on the loggerhead sea turtle *Caretta caretta* (Linnaeus 1758). U.S. Fish and Wildlife Service Biological Reports 88(14):1–110.

Dodd, C.K., Jr. 1997. Population structure and the evolution of sexual size dimorphism and sex ratios in an insular population of Florida box turtles (*Terrapene carolina bauri*). Canadian Journal of Zoology 75:1495–1507.

Dodd, C.K., Jr. 2001. North American Box Turtles: A Natural History. Norman, OK: University of Oklahoma Press, 231 pp.

Dodd, C.K., Jr., Adler, K., Bour, R., Schaffer, C., Thomson, S., and Walsh, T.J. 2016. The etymology of terrapin and *Terrapene*. Herpetological Review 47(2):191–192.

Dollo, L. 1886. Première note sur les chéloniens du Bruxellien (Eocène moyen) de la Belgique. Bulletin du Musée Royal d'Histoire Naturelle de Belgique 4:75–96.

Dolph, E. 2017. Assessment of Painted Turtle size and age from long-term pond study. University of Nebraska - Lincoln , Environmental Studies Undergraduate Student Theses, No. 197.

Donndorff, J.A. 1798. Zoologische Beyträge zur XIII. Ausgabe des Linneischen Natursystems. Dritter Band. Amphibien und Fische. Leipzig: Weidmannschen Buchhandlung, 980 pp.

Donoso-Barros, R. 1965. Distribución de las tortugas en Sudamerica. Publicaciónes Ocasionales del Museo Nacional de Historia Natural, Santiago 8:1–14.

Dorado-Rodrigues, T.F., Brito, E.S., Silva, K.R., Valadão, R.M., Cunha, F.A.G., and Strüssmann, C. 2023. New records and distribution extension of *Acanthochelys macrocephala* (Rhodin, Mittermeier & McMorris 1984) in midwestern Brazil. Cuadernos de Herpetología 37(2):223–226.

Dornburg, A., Beaulieu, J.M., Oliver, J.C., and Near, T.J. 2011. Integrating fossil

preservation biases in the selection of calibrations for molecular divergence time estimation. Systematic Biology 60:519–527.

Drabeck, D.H., Chatfield, M.W.H., and Richard-Zawacki, C.L. 2014. The status of Louisiana's Diamondback Terrapin (*Malaclemys terrapin*) populations in the wake of the Deepwater Horizon oil spill: insights from population genetic and contaminant analysis. Journal of Herpetology 48:125–136.

Du Tertre, J.B. 1667. Histoire Générale des Antilles Habitées par les François. Tome II. Contenant l'Histoire Naturelle. Paris: Thomas Jolly, 539 pp.

Dubois, A. 2025. *Bionomina*, opinion papers and nomenclatural *Code*-compliance. Bionomina 41:1–19.

Dubois, A. and Bour, R. 2010a. The nomenclatural status of the nomina of amphibians and reptiles created by Garsault (1764), with a parsimonious solution to an old nomenclatural problem regarding the genus *Bufo* (Amphibia, Anura), comments on the taxonomy of this genus, and comments on some nomina created by Laurenti (1768). Zootaxa 2447:1–52.

Dubois, A. and Bour, R. 2010b. The distinction between family-series and class-series nomina in zoological nomenclature, with emphasis on the nomina created by Batsch (1788, 1789) and on the higher nomenclature of turtles. Bonn Zoological Bulletin 57(2):149–171.

Dubois, A. and Frétey, T. 2024. LTH 5. Literatura Taxonomica Herpetologica. 5. Year 1760. Bionomina 39:46–75.

Duchene, S., Frey, A., Alfaro-Núñez, A., Dutton, P.H., and Gilbert, M.T.P. 2012. Marine turtle mitogenome phylogenetics and evolution. Molecular Phylogenetics and Evolution 65:241–250.

Ducotterd, J.-M. 1997. Notes sur la taille record d'une *Testudo marginata*. CITS Bulletin 9:41–42.

Duellman, W.E. and Berg, B. 1962. Type specimens of amphibians and reptiles in the Museum of Natural History, the University of Kansas. University of Kansas Publications, Museum of Natural History 15(4):183–204.

Duerden, J.E. 1906. South African tortoises of the genus *Homopus*, with description of a new species. Records of the Albany Museum 1:405–411.

Dugès, A. 1888. Erpetología del Valle de México. La Naturaleza (2)1:97–146.

Duisebayeva, T., Brushko, Z., Saraev, F., and Bidashko, F. 2017. Distribution of the marsh turtle *Emys orbicularis* (Linnaeus, 1758) on the eastern periphery of the species (Kazakhstan). In: Spatial-Time Dynamics of Biota in the Aralo-Caspian Basin Ecosystem. Proceedings of the II International Conference, Orenburg, Russia, pp. 174–183. [in Russian]

Duméril, A.H.A. 1852. Description des reptiles nouveaux ou imparfaitement connus de la collection du Muséum d'Histoire Naturelle et remarques sur la classification et les caractères des reptiles. Premiere Mémoire. Ordre des chéloniens et premières familles de l'ordre des sauriens (crocodiliens et caméléoniens). Archives du Muséum d'Histoire Naturelle, Paris 6:209–264.

Duméril, A.H.A. 1855. Notice historique sur la ménagerie des reptiles du Muséum d'Histoire Naturelle et observations qui y ont été recueillies. Archives du Muséum d'Histoire Naturelle, Paris 7:193–320.

Duméril, A.H.A. 1856. Note sur les reptiles du Gabon. Revue et Magasin de Zoologie Pure et Appliquée, Paris (2)8:369–377, 417–424.

Duméril, A.H.A. 1861a. Reptiles et poissons de l'Afrique occidentale. Étude précédée de considérations générales sur leur distribution géographique. Archives du Muséum d'Histoire Naturelle, Paris 10:138–268.

Duméril, A.H.A. 1861b. Catalogue des poissons, reptiles de la collection du Muséum d'Histoire Naturelle de Paris. Archives du Muséum d'Histoire Naturelle, Paris 10:429–460.

Duméril, A.H.A. and Bocourt, M.F. 1870. Observations sur les reptiles et les batraciens de la Région Centrale de l'Amérique. Classe des reptiles. Ordre des chéloniens. In: Duméril, A.H.A., Bocourt, M.F., and Mocquart, F. Mission Scientifique au Mexique et dans l'Amérique Centrale. Recherches Zoologiques. Troisième Partie, Première Section. Etude sur les Reptiles. Paris: Imprimerie Impériale, pp. 1–32 (première livraison).

Duméril, A.M.C. 1805 ["1806"]. Zoologie Analytique, ou Méthode Naturelle de Classification des Animaux. Paris: Perronneau, 344 pp. [Published Nov 1805].

Duméril, A.M.C. and Bibron, G. 1834. Erpétologie Générale ou Histoire Naturelle Complète des Reptiles. Tome Premier. Paris: Roret, 439 pp.

Duméril, A.M.C. and Bibron, G. 1835. Erpétologie Générale ou Histoire Naturelle Complète des Reptiles. Tome Second. Paris: Roret, 680 pp.

Duméril, A.M.C. and Bibron, G. 1851. [*Emys areolata, Emys berardii, Cinosternon leucostomum, Cinosternon cruentatum*]. In: Duméril, A.M.C. and Duméril, A.H.A. Catalogue Méthodique de la Collection des Reptiles (Muséum d'Histoire Naturelle de Paris). Paris: Gide and Baudry, 224 pp.

Duméril, A.M.C. and Duméril, A.H.A. 1851. Catalogue Méthodique de la Collection des Reptiles (Muséum d'Histoire Naturelle de Paris). Paris: Gide and Baudry, 224 pp.

Dundee, H.A. and Rossman, D.A. 1989. The Amphibians and Reptiles of Louisiana. Baton Rouge: Louisiana State University Press, 300 pp.

Dunn, E.R. 1917. Reptile and amphibian collections from the North Carolina mountains, with especial reference to salamanders. Bulletin of the American Museum of Natural History 37:593–634.

Dunn, E.R. 1920. Note on *Melanemys* Shufeldt. Copeia 1920:7–8.

Dunn, E.R. 1930. A new *Geoemyda* from Costa Rica. Proceedings of the New England Zoological Club 12:31–34.

Dunn, E.R. and Stuart, L.C. 1951. Comments on some recent restrictions of type-localities of certain South and Central American amphibians and reptiles. Copeia 1951(1):55–61.

Dürigen, B. 1897. Deutschlands Amphibien und Reptilien. Magdeburg: Creutz, 676 pp.

Durell, B. and Keeley, T. 2019. Sexually dimorphic growth in the Western Swamp Tortoise, *Pseudemydura umbrina* (Testudines: Chelidae). Journal of Zoo and Aquarium Research 7(3):102–108.

Dutton, P.H., Roden, S.E., Stewart, K.R., LaCasella, E., Tiwari, M., Formia, A., Thomé, J.C., Livingstone, S.R., Eckert, S., Chacon-Chaverri, D., Rivalan, P., and Allman, P. 2013. Population stock structure of leatherback turtles (*Dermochelys coriacea*) in the Atlantic revealed using mtDNAand microsatellite markers. Conservation Genetics 14:625–636.

Duvernoy, G.L. 1838. Les Reptiles. Avec un Atlas. In: Le Règne Animal distribué d'après son organisation ... par Georges Cuvier. Paris: Fortin, Masson et Cie. Texte, 169 pp., Atlas, 42 plates.

Duysebaeva, T.N., Doronin, I.V., Malakhov, D.V., Kukushkin, O.V., and Bakiev, A.G. 2019. GIS analysis of the distribution and habitation conditions of *Emys orbicularis orbicularis* (Testudines, Emydidae): methodical aspects. Proceedings of the VII Congress of the Nikolsky Herpetological Society. University Proceedings, Volga Region, Natural Sciences 1:28–40. [in Russian]

Echelle, A.A., Hackler, J.C., Lack, J.B., Ballard, S.R., Roman, J., Fox, S.F., Leslie, D.M., Jr., and Van Den Bussche, R.A. 2009 [2010]. Conservation genetics of the alligator snapping turtle: cytonuclear evidence of range-wide bottleneck effects and unusually pronounced geographic structure. Conservation Genetics 11(4):1375–1387.

Echeverri-García, L.D.P., Carr, J.L., Garcés-Restrepo, M.F., Galvis Rizo, C.A., and Giraldo, A. 2012. *Rhinoclemmys melanosterna* (Gray 1861). In: Páez, V.P., Morales-Betancourt, M.A., Lasso, C.A., Castaño-Mora, O.V., and Bock, B.C. (Eds.). Biología y Conservación de las Tortugas Continentales de Colombia. Bogotá, Colombia: Instituto de Investigación de Recursos Biológicos Alexander von Humboldt, pp. 308–314.

Echternacht, A.C., Burton, F.J., and Blumenthal, J.M. 2011. The amphibians and reptiles of the Cayman Islands: conservation issues in the face of invasion. In: Hailey, A., Wilson, B.S., and Horrocks, J.A. (Eds.). Conservation of Caribbean Island Herpetofaunas. Volume 2. Regional Accounts of the West Indies. Brill, pp. 129–147.

Eckert, K.L. and Luginbuhl, C. 1988. Death of a giant. Marine Turtle Newsletter 43:2–3.

Eckert, K.L., Wallace, B.P., Frazier, J.G., Eckert, S.A., and Pritchard, P.C.H. 2012. Synopsis of the biological data on the leatherback sea turtle (*Dermochelys coriacea*). Washington, DC: U.S. Department of Interior, Fish and Wildlife Service, Biological Technical Publication BTP-R4015-2012, 160 pp.

Edwards, G. 1747. A Natural History of Birds. Most of which have not been figured or described, and others very little known, from obscure or too brief Descriptions without Figures, or from Figures very ill designed. Part II. London: College of Physicians, p. 70.

Edwards, G. 1751. A Natural History of Birds. The most of which have not hitherto been either figured or described, and the Rest, by Reason of Obscure, or too brief Descriptions without Figures, or of Figures very ill designed, are hitherto but little known. Part IV. London: Royal College of Physicians, plates 211–260.

Edwards, G. 1760. Gleanings of Natural History, Exhibiting Figures of Quadrupeds, Birds, Insects, Plants, etc., Most of which have not, till now, been either Figured or Described. Part II. London: College of Physicians, plates 261–310.

Edwards, D.L., Garrick, R.C., Tapia, W., and Caccone, A. 2014. Cryptic structure and niche divergence within threatened Galapagos giant tortoises from southern Isabela Island. Conservation Genetics 15:1357–1369.

Edwards, T., Berry, K.H., Inman, R.D., Esque, T.C., Nussear, K.E., Jones,

C.A., and Culver, M. 2015a. Testing taxon tenacity of tortoises: evidence for a geographical selection gradient at a secondary contact zone. Ecology and Evolution 5:2095–2114.

Edwards, T., Tollis, M., Hsieh, P., Gutenkunst, R.N., Liu, Z., Kusimi, K., Culver, M., and Murphy, R.W. 2015b. Assessing models of speciation under different biogeographic scenarios; an empirical study using multi-locus and RNA-seq analyses. Ecology and Evolution; doi:10.1002/ece3.1865.

Edwards, T., Karl, A.E., Vaughn, M., Rosen, P.C., Melendez Torres, C., and Murphy, R.W. 2016. The desert tortoise trichotomy: Mexico hosts a third, new sister species of tortoise in the *Gopherus morafkai – G. agassizii* group. ZooKeys 562:131–158.

Ehret, D.J., Bourque, J.R., and Hulbert, R.C., Jr. 2013. Case 3628. *Terrapene putnami* Hay, 1906 (Testudines, Emydidae): replacement of the holotype by designation of a neotype. Bulletin of Zoological Nomenclature 70(3):193–198.

Ehrhart, L.M. 1978. Threatened and endangered species of the Kennedy Space Center. Semi-Annual Report to National Aeronautics and Space Administration, John F. Kennedy Space Center, pp. 590–652.

Ehrhart, L.M. and Yoder, R.G. 1978. Marine turtles of Merritt Island National Wildlife Refuge, Kennedy Space Center, Florida. In: Henderson, G.E. Proceedings of the Florida and Interregional Conference on Sea Turtles, 24–25 July 1976, Jensen Beach, Florida. Florida Marine Resource Publications 33:25–30.

Eichwald, C.E. von. 1831. Zoologia Specialis quam Expositis Animalibus, tum Fossilibus Potissimum Rossiae in Universum, et Poloniae in Specie, in usum Lectionum Publicarum in Universitate Caesarea Vilnensi. Pars Posterior. Vilna: Josephi Zawadzki, 404 pp.

Eichwald, C.E. von. 1840. Fauna Caspio-Caucasia nonnullis observationibus novis. Fasciculus I. Petropoli (St. Petersburg): Litteris Typhographiae Diarii, 233 pp.

Eisemberg, C.C., Rose, M., Yaru, B., and Georges, A. 2011. Demonstrating decline of an iconic species under sustained indigenous harvest – the pig-nosed turtle (*Carettochelys insculpta*) in Papua New Guinea. Biological Conservation 144:2282–2288.

Ellis, R.J. and Georges, A. 2015. An annotated type catalogue of the turtles (Testudines: Pleurodira: Chelidae) in the collection of the Western Australian Museum. Records of the Western Australian Museum 30:52–60.

Emerson, B.C. and Faria, C.M.A. 2014. Fission or fusion in island taxa – serendipity, or something to be expected? Molecular Ecology 23:5132–5134.

Enge, K.M. and Foster, D.R. 1986. *Sternotherus minor minor* (Loggerhead Musk Turtle). Size. Herpetological Review 17(1):25.

Enge, K.M., Stevenson, D.J., Thomas, T.M., Johnston, G.R., Jensen, J.B., Stegenga, B.S., Chandler, H.C., and Moler, P.E. 2021. Distribution and relative abundance of the Suwannee Alligator Snapping Turtle (*Macrochelys suwanniensis*). Chelonian Conservation and Biology 20(2):184–199.

Engel, E., Gassert, F., Mestdagh, X., and Proess, R. 2018. Verbreitungsatlas der Reptilien des Grossherzogtums Luxemburg. Ferrantia 28:1–72.

Engstrom, T.N., Shaffer, H.B., and McCord, W.P. 2004. Multiple data sets, high homoplasy, and the phylogeny of softshell turtles (Testudines: Trionychidae). Systematic Biology 53:693–710.

Ennen, J.R., Lovich, J.E., Kreiser, B.R., Selman, W., and Qualls, C.P. 2010. Genetic and morphological variation between populations of the Pascagoula Map Turtle (*Graptemys gibbonsi*) in the Pearl and Pascagoula rivers with description of a new species. Chelonian Conservation and Biology 9(1):98–113.

Ennen, J.R., Kreiser, B.R., Qualls, C.P., Gaillard, D., Aresco, M., Birkhead, R., Tuberville, T., McCoy, E., Mushinsky, H., Henges, T., and Schrey, A. 2012. Mitochondrial DNA assessment of the phylogeography of the gopher tortoises. Journal of Fish and Wildlife Management 3:110–122.

Ennen, J.R., Kalis, M.E., Patterson, A.L., Kreiser, B.R., Lovich, J.E., Godwin, J., and Qualls, C.P. 2014. Clinal variation or validation of a subspecies? A case study of the *Graptemys nigrinoda* complex (Testudines: Emydidae). Biological Journal of the Linnean Society 111:810–822.

Ennen, J.R., Godwin, J., Lovich, J.E., Kreiser, B.R., Folt, B., and Hazzard, S. 2016. Interdrainage morphological and genetic differentiation in the Escambia Map Turtle, *Graptemys ernsti*. Herpetological Conservation and Biology 11:122–131.

Ennen, J.R., Agha, M., Sweat, S.C., Matamoros, W.A., Lovich, J.E., Rhodin, A.G.J., Iverson, J.B., and Hoagstrom, C.W. 2020. Turtle biogeography: global regionalization and conservation priorities. Biological Conservation 241:108323, pp.1–11.

Ennen, J.R., Agha, M., Sweat, S.C., Matamoros, W.A., Lovich, J.E., Iverson, J.B., Rhodin, A.G.J., Thomson, R.C., Shaffer, H.B., and Hoagstrom, C.W. 2021. A watershed moment: analysis of sub-basins refocuses the geography of turtle conservation across the globe. Biological Conservation 253: 108925: pp.1–9.

Ernst, C.H. 1967. Intergradation between the painted turtles *Chrysemys picta picta* and *Chrysemys picta dorsalis*. Copeia 1967(1):131–136.

Ernst, C.H. 1970. The status of the painted turtle, *Chrysemys picta*, in Tennessee and Kentucky. Journal of Herpetology 4:39–45.

Ernst, C.H. 1978. A revision of the Neotropical turtle genus *Callopsis* (Testudines: Emydidae: Batagurinae). Herpetologica 34(2):113–134.

Ernst, C.H. 1980a. *Rhinoclemmys areolata*. Catalogue of American Amphibians and Reptiles 251:1–2.

Ernst, C.H. 1980b. *Rhinoclemmys funerea*. Catalogue of American Amphibians and Reptiles 263:1–2.

Ernst, C.H. 1981a. *Rhinoclemmys pulcherrima*. Catalogue of American Amphibians and Reptiles 275:1–2.

Ernst, C.H. 1981b. *Phrynops gibbus*. Catalogue of American Amphibians and Reptiles 279:1–2.

Ernst, C.H. 1983. *Platemys spixii*. Catalogue of American Amphibians and Reptiles 326:1–2.

Ernst, C.H. 1984. Geographic variation in the neotropical turtle, *Platemys platycephala*. Journal of Herpetology 17(4)(1983)[1984]:345–355.

Ernst, C.H. 1987. *Platemys, Platemys platycephala*. Catalogue of American Amphibians and Reptiles 405:1–4.

Ernst, C.H. 1988. *Cuora mccordi*, a new Chinese box turtle from Guangxi Province. Proceedings of the Biological Society of Washington 101:466–470.

Ernst, C.H. 1990. *Pseudemys gorzugi*. Catalogue of American Amphibians and Reptiles 461:1–2.

Ernst, C.H. 2008. *Trachemys emolli*. Catalogue of American Amphibians and Reptiles 846:1–3.

Ernst, C.H. and Barbour, R.W. 1989. Turtles of the World. Washington, DC: Smithsonian Institution Press, 313 pp.

Ernst, C.H. and Bogadek, A. 2005. *Palea steindachneri* (Wattle-necked Softshell). Size. Herpetological Review 36(3):313.

Ernst, C.H. and Bury, R.B. 1977. *Clemmys muhlenbergii*. Catalogue of American Amphibians and Reptiles 204:1–2.

Ernst, C.H. and Bury, R.B. 1982. *Malaclemys, M. terrapin*. Catalogue of American Amphibians and Reptiles 299:1–4.

Ernst, C.H. and Ernst, E.M. 1971. The taxonomic status and zoogeography of the painted turtle, *Chrysemys picta*, in Pennsylvania. Herpetologica 27(4):390–396.

Ernst, C.H. and Hartsell, T.D. 2000. An earlier name for the mangrove diamondback terrapin, *Malaclemys terrapin rhizophorarum* (Reptilia: Testudines: Emydidae). Proceedings of the Biological Society of Washington 113:887–889.

Ernst, C.H. and Laemmerzahl, A. 2002. Geographic variation in the Asian big-headed turtle *Platysternon megacephalum* (Reptilia: Testudines: Platysternidae). Proceedings of the Biological Society of Washington 115:18–24.

Ernst, C.H. and Lovich, J.E. 1990. A new species of *Cuora* (Reptilia: Testudines: Emydidae) from the Ryukyu Islands. Proceedings of the Biological Society of Washington 103:26–34.

Ernst, C.H. and Lovich, J.E. 2009. Turtles of the United States and Canada. Second Edition. Baltimore: Johns Hopkins University Press, 827 pp.

Ernst, C.H. and McBreen, J.F. 1991. *Terrapene carolina*. Catalogue of American Amphibians and Reptiles 512:1–13.

Ernst, C.H. and McCord, W.P. 1987. Two new turtles from southeast Asia. Proceedings of the Biological Society of Washington 100:624–628.

Ernst, C.H., Laemmerzahl, A.F., and Creque, T.R. 2006a. A review of morphological and pattern variation in the painted turtle, *Chrysemys picta*, in Missouri, USA, with an alternate hypothesis of the origin of *Chrysemys picta marginata*. Herpetological Bulletin 95:6–15.

Ernst, C.H., Altenburg, R.G.M., and Barbour, R.W. 2006b. Turtles of the world. DVD-ROM for Windows and Macintosh, version 1.3.

Ernst, C.H., Laemmerzahl, A.F., and Lovich, J.E. 2008. A morphological review of the *Cuora flavomarginata* complex (Testudines: Geoemydidae). Proceedings of the Biological Society of Washington 121:391–397.

Ernst, C.H., Batistella, A.M., and Vogt, R.C. 2010. *Trachemys adiutrix*. Catalogue of American Amphibians and Reptiles 869:1–4.

Ernst, C.H., Laemmerzahl, A.F., and Lovich, J.E. 2011. Does the "*kamaroma*"-plastron pattern morph occur in both Philippine subspecies of the turtle *Cuora amboinensis*? Proceedings of the Biological Society of Washington 124(4):259–269.

Ernst, C.H., Orr, J.M., Laemmerzahl, A.F., and Creque, T.R. 2015. Variation and zoogeography of the turtle *Chrysemys picta* in Virginia, USA. Herpetological Bulletin 130:9–15.

Ernst, C.H., Laemmerzahl, A.F., and Lovich, J.E. 2016. A morphological review of subspecies of the Asian box turtle, *Cuora amboinensis* (Testudines, Geomydidae). Proceedings of the Biological Society of Washington 129:144–156.

Eschscholtz, J.F. von. 1829a [previously cited as 1829b]. Beschreibungen dreier neuer Meerschildkröten. Die Quatember, Mitau [Jelgava, Latvia] 1(1):10–18. [published Jan 1829].

Eschscholtz, J.F. von. 1829b [previously cited as 1829a]. Zoologischer Atlas, enthaltend Abbildungen und Beschreibungen neuer Thierarten, während des Flottcapitains von Kotzebue zweiter Reise um die Welt, auf der Russisch–Kaiserlichen Kriegsschlupp Predpriaetië in den Jahren 1823–1826. Berlin: G. Reimer, Erstes Heft, 17 pp. + 15 pl. [published after May 1829].

Espindola, S., Vázquez-Domínguez, E., Nakamura, M., Osorio-Olvera, L., Martínez-Meyer, E., Myers, E.A., Overcast, I., Reid, B.N., and Burbrink, F.T. 2022. Complex genetic patterns and distribution limits mediated by native congeners of the worldwide invasive red-eared slider turtle. Molecular Ecology 31(6):1766–1782.

Ettmar, S. 2019. Toad-headed Turtles of the Genus *Mesoclemmys*: Distribution, Natural History, Husbandry. Frankfurt am Main: Edition Chimaira, 204 pp.

Eustace, A., Esser, L.F., Mremi, R., Malonza, P.K., and Mwaya, R.T. 2021. Protected areas network is not adequate to protect a critically endangered East Africa chelonian: modelling distribution of pancake tortoise, *Malacochersus tornieri* under current and future climates. PLoS ONE 16(1): e0238669.

Ewert, M.A. 2005. *Sternotherus odoratus* (common musk turtle): size and reproduction. Herpetological Review 36:314.

Ewert, M.A. and Jackson, D.R. 2005. *Kinosternon baurii* (Striped Mud Turtle). Size and reproduction. Herpetological Review 36:442.

Ewert, M.A., Etchberger, C.R., and Nelson, C.E. 2004. Turtle sex-determining modes and TSD patterns, and some TSD pattern correlates. In: Valenzuela, N. and Lance, V.A. (Eds.). Temperature–Dependent Sex Determination in Vertebrates. Washington, DC: Smithsonian Books, pp. 21–32.

Ewert, M.A., Jackson, D.R., and Buhlmann, K.A. 2006. *Deirochelys reticularia* – chicken turtle. In: Meylan, P.A. (Ed.). Biology and Conservation of Florida Turtles. Chelonian Research Monographs 3:249–259.

Ewert, M.A., Pritchard, P.C.H., and Wallace, G.E. 2006. *Graptemys barbouri* – Barbour's map turtle. In: Meylan, P.A. (Ed.). Biology and Conservation of Florida Turtles. Chelonian Research Monographs 3:260–272.

Fachín-Terán, A., Vogt, R.C., and Thornbjarnarson, J.B. 2003. Estrutura populacional, razão sexual e abundância de *Podocnemis sextuberculata* (Testudines, Podocnemididae) na Reserva de Desenvolvimento Sustentável Mamirauá, Amazonas, Brasil. Phyllomedusa 2(1):43–63.

Fagundes, C.K., Bager, A., and Cechin, S.T.Z. 2010. *Trachemys dorbigni* in an anthropic environment in southern Brazil: I) Sexual size dimorphism and population estimates. Herpetological Journal 20:185–193.

Falconer, H. and Cautley, P.T. 1837. On additional fossil species of the order Quadrumana from the Siwalik Hills. Journal of the Asiatic Society of Bengal 6:354–360.

Fall, P.L., van Hengstum, P.J., Lavold-Foote, L., Donnelly, J.P., Albury, N.A., and Tamalavage, A.E. 2021. Human arrival and landscape dynamics in the northern Bahamas. PNAS 118(10):e2015764118, 7 pp.

Fan, T.H. 1931. Preliminary report of reptiles from Yaoshan, Kwangsi, China. Bulletin of the Department of Biology, College of Science, Sun Yatsen University No. 11, 154 pp.

Fang, P.W. 1934. Notes on some chelonians of China. Sinensia 4(7):145–200.

Farina, B.M., Godoy, P.L., Benson, R.B.J., Langer, M.C., and Ferreira, G.S. 2023. Turtle body size evolution is determined by lineage-specific specializations rather than global trends. Ecology and Evolution 13:e10201, 23 pp.

Farkas, B. 1992. Wiederentdeckung eines Exemplars von *Rafetus swinhoei* (Gray, 1873) im Naturhistorischen Museum Wien. Salamandra 28(2):145–152.

Farkas, B. 1994. Notes on type and type locality of the narrow-headed softshell turtle, *Chitra indica* (Gray, 1831) (Testudines, Trionychidae). Miscellanea Zoologica Hungarica 9:117–119.

Farkas, B. and Csorba, G. 1999. Notes on the distribution and maximum size of *Melanochelys trijuga indopeninsularis* in Nepal (Reptilia: Testudines: Bataguridae). Faunistische Abhandlungen, Staatliches Museum für Tierkunde Dresden 21:305–307.

Farkas, B. and Fritz, U. 1998. On the identity of *Rafetus swinhoei* (Gray, 1873) and *Pelochelys maculatus* (Heude, 1880) (Reptilia: Testudines: Trionychidae). Zoologische Abhandlungen, Staatliches Museum für Tierkunde Dresden 50(5):59–75.

Farkas, B. and Webb, R.G. 2003. *Rafetus leloii* Ha Dinh Duc, 2000 – an invalid species of softshell turtle from Hoan Kiem Lake, Hanoi, Vietnam (Reptilia, Testudines, Trionychidae). Zoologische Abhandlungen (Dresden) 53:107–112.

Farkas, B.L., Sasvari, L., and Buskirk, J.R. 1997. Maximum size of the Egyptian tortoise, *Testudo kleinmanni*. Chelonian Conservation and Biology 2(3):415.

Farkas, B., Le, M.D., and Nguyen, T.Q. 2011. *Rafetus vietnamensis* Le, Le, Tran, Phan, Phan, Tran, Pham, Nguyen, Nong, Phan, Dinh, Truong, and Ha, 2010—another invalid name for an invalid species of softshell turtle (Reptilia: Testudines: Trionychidae). Russian Journal of Herpetology 18(1):65–72.

Farkas, B., Ziegler, T., Pham, C.T., Ong, A.V., and Fritz, U. 2019. A new species of *Pelodiscus* from northeastern Indochina (Testudines, Trionychidae). ZooKeys 824:71–86.

Feldman, C.R. and Parham, J.F. 2002. A molecular phylogeny for emydine turtles: taxonomic revision and the evolution of shell kinesis. Molecular Phylogenetics and Evolution 22:388–398.

Feldman, C.R. and Parham, J.F. 2004. Molecular systematics of Old World stripe-necked turtles (Testudines: *Mauremys*). Asiatic Herpetological Research 10:28–37.

Fermin, P. 1765. Histoire Naturelle de la Hollande Equinoxiale, ou Description des Animaux, Plantes, Fruits, et Autres Curiosités Naturelles, qui se Trouvent dans la Colonie de Surinam. Amsterdam: M. Magerus, 240 pp.

Fernández, M.S. 1988. Las Testudinidae (Reptilia: Chelonii) argentinas: osteología, sistemática y distribución geográfica. Doctoral Thesis, Universidad Nacional de La Plata.

Ferrara, C.R., Fagundes, C.K., Morcattty, T.Q., and Vogt, R.C. 2017. Quelônios Amazônicos: Guia de Identificação e Distribuição. Manaus, Brazil: Wildlife Conservation Society, 180 pp.

Ferreira, G.S., Rincón, A.S., Solórzano, A., and Langer, M.C. 2016. Review of the fossil matamata turtles: earliest well-dated record and hypotheses on the origin of their present geographical distribution. The Science of Nature 103:28:1–12.

Ferreira, G.S., Bronzati, M., Langer, M.C., and Sterli, J. 2018. Phylogeny, biogeography and diversification patterns of side-necked turtles (Testudines: Pleurodira). Royal Society Open Science 5:171773.

Ferreira, J.B. 1897. Sobre alguns reptis ultimamente enviados a seccao zoologica do Museu de Lisboa. Jornal de Sciencias Mathematicas Physicas e Naturaes, Lisboa (2)5(18):111–116.

Ferri, V. and Diagne, T. 2013. New knowledge about the distribution of *Pelusios williamsii* Laurent, 1965 (Pelomedusidae) in Uganda. Proceedings ATTI II Italian Herpetological Congress, pp. 63–67.

Ferronato, B.O., Molina, F.B., Molina, F.C., Espinosa, R.A., and Morales, V.R. 2011. New locality records for chelonians (Testudines: Chelidae, Podocnemidae, Testudinidae) from Departamento de Pasco, Peru. Herpetology Notes 4:219–224.

Fielder, D. 2013. Ancient phenotypes revealed through present day species—a morphological analysis of Australia's Saw-Shelled Turtles including the Threatened *Myuchelys bellii* (Testudines: Chelidae). Chelonian Conservation and Biology 12(1):101–111.

Fielder, D., Vernes, K., Alacs, E., and Georges, A. 2012. Mitochondrial variation among Australian freshwater turtles (genus *Myuchelys*), with special reference to the Endangered *M. bellii*. Endangered Species Research 17:63–71.

Figgener, C., Bernardo, J., and Plotkin, P.T. 2022. Marine turtles are only minimally sexually size dimorphic, a pattern that is distinct from most nonmarine aquatic turtles. Ecology and Evolution 12:e8963, 10 pp.

Firdous, F. 1989. Male leatherback strands in Karachi. Marine Turtle Newsletter 47:14–15.

Fischer, J. von. 1872. *Staurotypus marmoratus* n. sp. Archiv für Naturgeschichte 38:265–272.

Fitzinger, L.J. 1826. Neue Classification der Reptilien, nach ihren Natürlichen Verwandtschaften nebst einer Verwandtschafts–Tafel und einem Verzeichnisse der Reptilien–Sammlung des k.k. Zoologischen Museum zu Wien. Wien: J.G. Hübner Verlagen, 66 pp.

Fitzinger, L.J. 1835. Entwurf einer systematischen Anordnung der Schildkröten nach den Grundsätzen der natürlichen Methode. Annalen des Wiener Museums der Naturgeschichte 1:105–128.

Fitzinger, L.J. 1843. Systema Reptilium. Fasciculus Primus: Amblyglossae.

Vindobona [Vienna]: Braumüller und Seidel, 106 pp.

Fitzinger, L.J. 1853. Versuch einer Geschichte der Menagerien des Österreichisch-Kaiserlichen Hofes. Wien: Kaiserlich-Königliche Hof- und Staatsdruckerei in Commission bei W. Braumüller, 198 pp.

FitzSimons, V.F.M. 1932. Preliminary descriptions of new forms of South African Reptilia and Amphibia, from the Vernay-Lang Kalahari Expedition, 1930. Annals of the Transvaal Museum 15:35–40.

FitzSimons, V.F.M. 1937. Notes on the reptiles and amphibians collected and described from South Africa by Andrew Smith. Annals of the Transvaal Museum 17:259–274.

FitzSimons, V.F.M. 1938. Transvaal Museum Expedition to South-West Africa and Little Namaqualand, May to August 1937. Reptiles and Amphibians. Annals of the Transvaal Museum 19:153–209.

Fleming, J. 1822. The Philosophy of Zoology; or a General View of the Structure, Functions, and Classification of Animals. Vol. II. Edinburgh: Archibald Constable and Co., 618 pp.

Fleming, J. 1828. A History of British Animals. Edinburgh: Archibald Constable and Co., 568 pp.

Flores-Villela, O.A. 1993. Herpetofauna Mexicana: lista anotada de las especies de anfibios y reptiles de Mexico, cambios taxonomicos recientes, y nuevas especies. Carnegie Museum of Natural History Special Publications 17:1–73.

Flores-Villela, O., Adler, K., and Eimermacher, T.G. 2016. Identity of three new sea turtles named by J. Friedrich Eschscholtz. Chelonian Conservation and Biology 15(1):157–162.

Folkerts, G.W. and Mount, R.H. 1969. A new subspecies of the turtle *Graptemys nigrinoda* Cagle. Copeia 1969(4):677–682.

Folt, B. 2020. Population demography of *Rhinoclemmys funerea* (Black River Turtle) at a protected forest reserve in Costa Rica. Herpetological Conservation and Biology 15(3):611–619.

Folt, B. and Guyer, G. 2015. Evaluating recent taxonomic changes for alligator snapping turtles (Testudines: Chelydridae). Zootaxa 3947(3):447–450.

Fong, J.J. and Chen, T.H. 2010. DNA evidence for the hybridization of wild turtles in Taiwan: possible genetic pollution from trade animals. Conservation Genetics 11:2061–2066.

Fong, J.J. and Qiao, G.X. 2010. New localities of endangered Chinese turtles from museum specimens and the practical and ethical challenges using and reporting natural history collection data. Zootaxa 2393:59–68.

Fong, J.J., Parham, J.F., and Fu, J. 2002. A reassessment of the distribution of *Cuora flavomarginata* Gray 1863 on mainland China. Russian Journal of Herpetology 9:9–14.

Fong, J.J., Parham, J.F., Shi, H., Stuart, B.L., and Carter, R.L. 2007. A genetic survey of heavily exploited, endangered turtles: caveats on the conservation value of trade animals. Animal Conservation 10:452–460.

Fong, J.J., Stuart, B.L., McCormack, T.E.M., and Parham, J.F. 2019. First genetic data of the critically endangered Vietnamese Pond Turtle (*Mauremys annamensis*) from known-locality specimens. Current Herpetology 38:140–152.

Fontaine, M.C. 2017. A genomic perspective is needed for the re-evaluation of species boundaries, evolutionary trajectories, and conservation strategies of the Galápagos giant tortoises. Peer Community in Evolutionary Biology, 100031, doi:10.24072/pci.evolbiol.100031.

Forero-Medina, G., Ferrara, C.R., Vogt, R.C., Fagundes, C.K., Balestra, R.A.M., Andrade, P.C.M., Lacava, R., Bernhard, R., Lipman, A.J., Lenz, A.J., Ferrer, A., Calle, A., Aponte, A.F., Calle-Rendón, B.R., Camilo, C.S., Perrone, E., Miraña, E., Cunha, F.A.G., Loja, E., Del Rio, J., Vera Fernandez, J.L., Hernández, O.E., Aguila, R.D., Pino, R., Cueva, R., Martinez, S., Bernardes, V.C.D., Sainz, L., and Horne, B.D. 2019. On the future of the giant South American river turtle *Podocnemis expansa*. Oryx, doi:https://doi.org/10.1017/S0030605318001370, 8 pp.

Forskål, P. [Forsskål, P.]. 1775. Descriptiones Animalium: Avium, Amphibiorum, Piscium, Insectorum, Vermium; quae in Itinere Orientali Observavit. Post mortem auctoris edidit Carsten Niebuhr. Hauniae [Copenhagen]: Mölleri, 164 pp.

Forstner, M.R.J., Dixon, J.R., Guerra, T.M., Winters, J.M., Stuart, J.N., and Davis, S.K. 2014. Status of U.S. populations of the Big Bend Slider (*Trachemys gaigeae*). In: Hoyt, C.A. and J. Karges, A. (Eds). Proceedings of the Sixth Symposium on the Natural Resources of the Chihuahuan Desert Region. Fort Davis, TX: Chihuahuan Desert Research Institute, pp. 335–367.

Forth, G. 2017. Ethnographic reports of freshwater turtles on Flores Island, Indonesia: possibilities of an undocumented chelonian species. Herpetological Review 48:304–310.

Forth, G. 2018. Further evidence for undocumented freshwater turtles on Flores Island, Indonesia. Herpetological Review 49:19–21.

Fowler, H.W. 1906. Some cold–blooded vertebrates of the Florida Keys. Proceedings of the Academy of Natural Sciences, Philadelphia 58:77–113.

França, M.A.G. and Langer, M.C. 2006. Phylogenetic relationships of the Bauru Group turtles (Late Cretaceous of south-central Brazil). Revista Brasileira de Paleontologia 9(3):365–373.

Franz, R. 2014. The fossil record of North American tortoises. In: Rostal, D.C., McCoy, E.D., and Mushinsky, H.R. (Eds.). Biology and Conservation of North American Tortoises. Baltimore, Maryland: Johns Hopkins University Press, pp. 13–24.

Franz, R. and Franz, S.E. 2009. A new fossil land tortoise in the genus *Chelonoidis* (Testudines: Testudinidae) from the northern Bahamas, with an osteological assessment of other Neotropical tortoises. Bulletin of the Florida Museum of Natural History 49(1):1–44.

Franz, R. and Woods, C.A. 1983. A fossil tortoise from Hispaniola. Journal of Herpetology 17(1):79–81.

Franz, R., Carlson, L.A., Owen, R.D., and Steadman, D. 2001. Fossil tortoises from the Turks and Caicos Islands, B.W.I. In: Clark-Simpson, C.A. and Smith, G.W. (Eds.). Proceedings of the 8th Symposium on the Natural History of the Bahamas, pp. 27–31.

Franz, R., Albury, N.A., and Steadman, D.W. 2020. Extinct tortoises from the Turks and Caicos Islands. Bulletin of the Florida Museum of Natural History 58(1):1–38.

Franzen, M. and Glaw, F. 2007. Type catalogue of reptiles in the Zoologische Staatssammlung München. Spixiana 30(2):201–274.

Frazier, J. 2006. A neotype for the Aldabra tortoise, *Testudo gigantea* Schweigger, 1812. Herpetological Review 37:275–280.

Frazier, J. 2008. Case 3463: *Testudo gigantea* Schweigger, 1812 (Reptilia, Testudines): proposed conservation of usage. Bulletin of Zoological Nomenclature 65(2):82.

Frazier, J. 2009. Case 3463: *Testudo gigantea* Schweigger, 1812 (currently *Geochelone (Aldabrachelys) gigantea*; Reptilia, Testudines): proposed conservation of usage of the specific name by maintenance of a designated neotype, and suppression of *Testudo dussumieri* Gray, 1831 (currently *Dipsochelys dussumieri*). Bulletin of Zoological Nomenclature 66(1):34–50.

Frazier, J. 2020. The Galapagos: Island home of giant tortoises. In: Gibbs, J.P., Cayot, L.J., and Tapia Aguilera, W. (Eds.). Galapagos Giant Tortoises. London: Academic Press, pp. 3–21.

Frazier, J. and Matyot, P. 2010. On the identity of Monsieur Dussumier's Dutch tortoise and the lectotype of *Testudo dussumieri* Gray, 1831. Zootaxa 2665: 29–50.

Freedberg, S. and Myers, E.M. 2012. Cytonuclear equilibrium following interspecific introgression in a turtle lacking sex chromosomes. Biological Journal of the Linnean Society 106:405–417.

Freeman, A.B., Eisemberg, C., and Stoetzel, H. 2018. Habitat use and movements in an upland population of Johnstone River Snapping Turtles, *Elseya irwini*. Herpetological Conservation and Biology 13(2):464–472.

Freiberg, M.A. 1936. Una nueva tortuga del norte Argentino. Physis 12:169–171.

Freiberg, M.A. 1945. Una nueva especie de tortuga del genero *Platemys* Wagler. Physis 20:19–23.

Freiberg, M.A. 1947. El alotipo de la tortuga *Platemys pallidipectoris* Freiberg. Physis 20:112–114.

Freiberg, M.A. 1969. Una nueva subespecie de *Pseudemys dorbignyi* (Duméril et Bibron) (Reptilia, Chelonia, Emydidae). Physis 28:299–314.

Freiberg, M.A. 1973. Dos nuevas tortugas terrestres de Argentina. Boletín de la Sociedad de Biología de Concepción 46:81–93.

Fretey, J. and Bour, R. 1980. Redecouverte du type de *Dermochelys coriacea* (Vandelli) (Testudinata, Dermochelyidae). Bollettino di Zoologia 47:193–205.

Fretey, J., Hoogmoed, M.S., and Lescure, J. 1977. Etude taxinomique de *Rhinoclemmys punctularia punctularia* (Daudin) (Testudinata, Emydidae). Zoologische Mededelingen 52:63–80.

Frétey, T. 2022. The authorship and date of the familial nomen Dermochelyidae (Chelonii). Bionomina 31:46–69.

Frétey, T. 2024. Etymology of the nomina of chelonians of the world. 1. Genus-series nomina. Bionomina 40:1–116.

Fritts, T.H. 1983. Morphometrics of Galapagos tortoises: evolutionary implications. In: Bowman, R.I. and Leviton, A.E. (Eds.). Patterns of Evolution in Galapagos Organisms. San Francisco: American Association for the

Advancement of Science, pp. 107–122.

Fritz, U. 1989. Zur innerartlichen Variabilität von *Emys orbicularis* (Linnaeus, 1758). 1. Eine neue Unterart der Europäischen Sumpfschildkröte aus Kleinasien *Emys orbicularis luteofusca* subsp. nov. Salamandra 25:143–168.

Fritz, U. 1992. Zur innerartlichen Variabilität von *Emys orbicularis* (Linnaeus, 1758). 2. Variabilität in Osteuropa und Redefinition von *Emys orbicularis orbicularis* (Linnaeus, 1758) und *E. o. hellenica* (Valenciennes, 1832) (Reptilia, Testudines: Emydidae). Zoologische Abhandlungen, Staatliches Museum für Tierkunde Dresden 47:37–77.

Fritz, U. 1993. Zur innerartlichen Variabilität von *Emys orbicularis* (Linnaeus, 1758). 3. Zwei neue Unterarten von der Iberischen Halbinsel und aus Nordafrika, *Emys orbicularis fritzjuergenobsti* subsp. nov. und *E. o. occidentalis* subsp. nov. (Reptilia, Testudines: Emydidae). Zoologische Abhandlungen, Staatliches Museum für Tierkunde Dresden 47:131–155.

Fritz, U. 1994. Zur innerartlichen Variabilität von *Emys orbicularis* (Linnaeus, 1758). 4. Variabilität und Zoogeographie im pontokaspischen Gebiet mit Beschreibung von drei neuen Unterarten (Reptilia: Testudines: Emydidae). Zoologische Abhandlungen, Staatliches Museum für Tierkunde Dresden 48:53–93.

Fritz, U. 1995. Zur innerartlichen Variabilität von *Emys orbicularis* (Linnaeus, 1758). 5a. Taxonomie in Mittel–Westeuropa, auf Korsika, Sardinien, der Apenninen–Halbinsel und Sizilien und Unterartengrupppen von *E. orbicularis* (Reptilia: Testudines: Emydidae). Zoologische Abhandlungen, Staatliches Museum für Tierkunde Dresden 48:185–242.

Fritz, U. 1998. Introduction to zoogeography and subspecific differentiation in *Emys orbicularis* (Linnaeus, 1758). In: Fritz, U., Joger, U., Podloucky, R., and Servan, J. (Eds.). Proceedings of the EMYS Symposium Dresden 96. Mertensiella 10:1–27.

Fritz, U. 2001. *Emys orbicularis* (Linnaeus, 1758) - Europäische Sumpfschildkröte. In: Fritz, U. (Ed.). Handbuch der Reptilien und Amphibien Europas, Band 3/IIIA. Wiesbaden: Aula-Verlag, pp. 343–515.

Fritz, U. and Bininda-Emonds, O.R.P. 2007. When genes meet nomenclature: tortoise phylogeny and the shifting generic concepts of *Testudo* and *Geochelone*. Zoology 110:298–307.

Fritz, U. and Buskirk, J.R. 1997. The occurrence of *Testudo kleinmanni* in Tripolitania. Casopis Národniho Muzea Rada Prirodovedná 167:143–144.

Fritz, U. and Havaš, P. 2006. Checklist of Chelonians of the World, at the request of the CITES Nomenclature Committee and the German Agency for Nature Conservation. Dresden: German Federal Ministry of Environment, Nature Conservation and Nuclear Safety and Museum of Zoology, 230 pp.

Fritz, U. and Havaš, P. 2007. Checklist of chelonians of the world. Vertebrate Zoology 57:149–368.

Fritz, U. and Havaš, P. 2013. Order Testudines: 2013 update. Zootaxa 3703:12–14.

Fritz, U. and Havaš, P. 2014. On the reclassification of Box Turtles (*Terrapene*): a response to Martin et al. (2014). Zootaxa 3835:295–298.

Fritz, U. and Kraus, O. 2008. Comments on "*Chersine* Merrem, 1820 and *Chersina* Gray, 1831: a nomenclatural survey by Bour and Ohler, Zootaxa, 1752: 66–68". Zootaxa 1893:65–68.

Fritz, U. and Obst, F.J. 1996. Zur Kenntnis der Celebes-Erdschildkröte, *Heosemys yuwonoi* (McCord, Iverson and Boeadi, 1995). Herpetofauna 18(102):27–34.

Fritz, U. and Pauler, I. 1992. *Phrynops chacoensis* spec. nov. (Reptilia, Chelidae), eine neue Krötenkopfschildkröte. Mitteilung des Zoologischen Museums Berlin 68:299–307.

Fritz, U. and Pauler, I. 1999. *Phrynops chacoensis* Fritz & Pauler, 1992, ein Juniorsynonym von *Platemys macrocephala* Rhodin, Mittermeier & McMorris, 1984. Salamandra 35:53–56.

Fritz, U. and Wischuf, T. 1997. Zur Systematik westasiatisch–südosteuropäischer Bachschildkröten (Gattung *Mauremys*) (Reptilia: Testudines: Bataguridae). Zoologische Abhandlungen, Staatliches Museum für Tierkunde Dresden 49:223–260.

Fritz, U. and Ziegler, T. 1999. Contribution to the knowledge of *Cyclemys tcheponensis* (Bourret, 1939) and the distribution of *Cyclemys* in the Indochinese region (Reptilia: Testudines: Bataguridae). Revue française d'Aquariologie 26:71–78.

Fritz, U., Obst, F.-J., and Günther, R. 1994. Kritischer Typen-Katalog der Schildkrötensammlung (Reptilia: Testudines) des Zoologischen Museums Berlin. Mitteilung des Zoologischen Museums Berlin 70:157–175.

Fritz, U., Keller, C., and Budde, M. 1996. Eine neue Unterart der Europäischen Sumpfschildkröte aus Südwestspanien, *Emys orbicularis hispanica* subsp. nov. Salamandra 32:129–152.

Fritz, U., Gaulke, M., and Lehr, E. 1997. Revision der südostasiatischen Dornschildkröten-Gattung *Cyclemys* Bell, 1834, mit Beschreibung einer neuen Art. Salamandra 33:183–212.

Fritz, U., Andreas, B., and Lehr, E. 1998a. Eine neue Unterart der Dreikiel–Scharnierschildkröte, *Pyxidea mouhotii* (Gray, 1862) (Reptilia: Testudines: Bataguridae). Zoologische Abhandlungen, Staatliches Museum für Tierkunde Dresden 50:33–43.

Fritz, U., Baran, I., Budak, A., and Amthauer, E. 1998b. Some notes on the morphology of *Emys orbicularis* in Anatolia, especially on *E. o. luteofusca* and *E. o. colchica*, with the description of a new subspecies from southeastern Turkey. In: Fritz, U., Joger, U., Podloucky, R., and Servan, J. (Eds.). Proceedings of the EMYS Symposium Dresden 96. Mertensiella 10:103–122.

Fritz, U., Niekisch, M., and Ziegler, T. 1999. Rediscovery of the holotype of *Geoemyda tcheponensis* Bourret, 1939 (Reptilia: Testudines: Bataguridae). Zoologische Abhandlungen, Staatliches Museum für Tierkunde Dresden 50:243–247.

Fritz, U., Ziegler, T., Herrmann, H.W., and Lehr, E. 2002. Intergradation between subspecies of *Cuora galbinifrons* Bourret, 1939 and *Pyxidea mouhotii* (Gray, 1862) in southern North Vietnam (Reptilia: Testudines: Geoemydidae). Faunistische Abhandlungen Staatliches Museum für Tierkunde Dresden 23:59–74.

Fritz, U., Guicking, D., Lenk, P., Joger, U., and Wink, M. 2004. When turtle distribution tells European history: mtDNA haplotypes of *Emys orbicularis* reflect in Germany former division by the Iron Curtain. Biologia 59 (Suppl. 14):19–25.

Fritz, U., Fattizzo, T., Guicking, D., Tripepi, S., Pennisi, M.G., Lenk, P., Joger, U., and Wink, M. 2005a. A new cryptic species of pond turtle from southern Italy, the hottest spot in the range of the genus *Emys* (Reptilia, Testudines, Emydidae). Zoologica Scripta 34:351–371.

Fritz, U., Široký, P., Kami, H., and Wink, M. 2005b. Environmentally caused dwarfism or a valid species—is *Testudo weissingeri* Bour, 1996 a distinct evolutionary lineage? New evidence from mitochondrial and nuclear genomic markers. Molecular Phylogenetics and Evolution 37:389–401.

Fritz, U., Fritzsch, G., Lehr, E., Ducotterd, J.-M., and Müller, A. 2005c. The Atlas Mountains, not the Strait of Gibraltar, as a biogeographic barrier for *Mauremys leprosa* (Reptilia: Testudines). Salamandra 41:97–106.

Fritz, U., Barata, M., Busack, S.D., Fritzsch, G., and Castillo, R. 2006a. Impact of mountain chains, sea straits and peripheral populations on genetic and taxonomic structure of a freshwater turtle, *Mauremys leprosa* (Reptilia, Testudines, Geoemydidae). Zoologica Scripta 35:97–108.

Fritz, U., Auer, M., Bertolero, A., Cheylan, M., Fattizzo, T., Hundsdörfer, A.K., Sampayo, M.M., Pretus, J.L., Široký, P., and Wink, M. 2006b. A rangewide phylogeography of Hermann's tortoise, *Testudo hermanni* (Reptilia: Testudines: Testudinidae): implications for taxonomy. Zoologica Scripta 35:531–543.

Fritz, U., Petzold, A., and Auer, M. 2006c. Osteology in the *Cuora galbinifrons* complex suggests conspecifity of *C. bourreti* and *C. galbinifrons*, with notes on shell osteology and phalangeal formulae within the Geoemydidae. Amphibia-Reptilia 27:195–205.

Fritz, U., d'Angelo, S., Pennisi, M.G., and Lo Valvo, M. 2006d. Variation of Sicilian pond turtles, *Emys trinacris* - what makes a species cryptic? Amphibia-Reptilia 27:513–529.

Fritz, U., Hundsdörfer, A.K., Široký, P., Auer, M., Kami, H., Lehmann, J., Mazanaeva, L.F., Türkozan, O., and Wink, M. 2007a. Phenotypic plasticity leads to incongruence between morphology-based taxonomy and genetic differentiation in western Palearctic tortoises (*Testudo graeca* complex; Testudines, Testudinidae). Amphibia-Reptilia 28:97–121.

Fritz, U., Guicking, D., Kami, H., Arakelyan, M., Auer, M., Ayaz, D., Ayres Fernández, C., Bakiev, A.G., Celani, A., Džukic, G., Fahd, S., Havaš, P., Joger, U., Khabibullin, V.F., Mazanaeva, L.F., Široký, P., Tripepi, S., Valdeón Vélez, A., Velo Antón, G., Wink, M. 2007b. Mitochondrial phylogeography of European pond turtles (*Emys orbicularis, Emys trinacris*) – an update. Amphibia-Reptilia 28:418–426.

Fritz, U., Ayaz, D., Buschbom, J., Kami, H.G., Mazanaeva, L.F., Aloufi, A.A., Auer, M., Rifai, L., Silic, T., and Hundsdörfer, A.K. 2008a. Go east: phylogeographies of *Mauremys caspica* and *M. rivulata* – discordance of morphology, mitochondrial and nuclear genomic markers and rare hybridization. Journal of Evolutionary Biology 21:527–540.

Fritz, U., Guicking, D., Auer, M., Sommer, R.S.; Wink, M., and Hundsdörfer, A.K. 2008b. Diversity of the Southeast Asian leaf turtle genus *Cyclemys*: how

many leaves on its tree of life? Zoologica Scripta 37:367–390.

Fritz, U., Auer, M., Chirikova, M.A., Duysebayeva, T.N., Eremchenko, V.K., Kami, H.G., Kashkarov, R.D., Masroor, R., Moodley, Y., Pindrani, A., Široký, P., and Hundsdörfer, A.K. 2009a. Mitochondrial diversity of the widespread Central Asian steppe tortoise (*Testudo horsfieldii* Gray, 1844): implications for taxonomy and relocation of confiscated tortoises. Amphibia-Reptilia 30:245–257.

Fritz, U., Ayaz, D., Hundsdörfer, A.K., Kotenko, T., Guicking, D., Wink, M., Tok, C.V., Çiçek, K., and Buschbom, J. 2009b. Mitochondrial diversity of European pond turtles (*Emys orbicularis*) in Anatolia and the Ponto-Caspian Region: multiple old refuges, hotspot of extant diversification and critically endangered endemics. Organisms Diversity and Evolution 9:100–114.

Fritz, U., Harris, D.J., Fahd, S., Rouag, R., Graciá Martínez, E., Giménez Casalduero, A., Široký, P., Kalboussi, M., Jdeidi, T.B., and Hundsdörfer, A.K. 2009c. Mitochondrial phylogeography of *Testudo graeca* in the Western Mediterranean: old complex divergence in North Africa and recent arrival in Europe. Amphibia-Reptilia 30:63–80.

Fritz, U., Daniels, S.R., Hofmeyr, M.D., González, J., Barrio-Amorós, C.L., Široký, P., Hundsdörfer, A.K., and Stuckas, H. 2010a. Mitochondrial phylogeography and subspecies of the wide-ranging sub-Saharan leopard tortoise *Stigmochelys pardalis* (Testudines: Testudinidae) – a case study for the pitfalls of pseudogenes and GenBank sequences. Journal of Zoological Systematics and Evolutionary Research 48:348–359.

Fritz, U., Gong, S., Auer, M., Kuchling, G., Schneeweiss, N., and Hundsdörfer, A.K. 2010b. The world's economically most important chelonians represent a diverse species complex (Testudines: Trionychidae: *Pelodiscus*). Organisms, Diversity and Evolution 10:227–242.

Fritz, U., Branch, W.R., Hofmeyr, M.D., Maran, J., Prokop, H., Schleicher, A., Široký, P., Stuckas, H., Vargas-Ramírez, M., Vences M., and Hundsdörfer, A.K. 2011a. Molecular phylogeny of African hinged and helmeted terrapins (Testudines: Pelomedusidae: *Pelusios* and *Pelomedusa*). Zoologica Scripta 40:115–125.

Fritz, U., Schmidt, C., and Ernst, C.H. 2011b. Competing generic concepts for Blanding's, Pacific and European pond turtles (*Emydoidea*, *Actinemys* and *Emys*) – which is best? Zootaxa 2791:41–53.

Fritz, U., Stuckas, H., Vargas-Ramírez, M., Hundsdörfer, A.K., Maran, J., and Päckert, M. 2011c. Molecular phylogeny of Central and South American slider turtles: implications for biogeography and systematics (Testudines: Emydidae: *Trachemys*). Journal of Zoological Systematics and Evolutionary Research 50:125–136.

Fritz, U., Alcalde, L., Ramírez-Vargas, M., Goode, E.V., Fabius-Turoblin, D.U., and Praschag, P. 2012a. Northern genetic richness and southern purity, but just one species in the *Chelonoidis chilensis* complex. Zoologica Scripta 41:220–232.

Fritz, U., Vargas-Ramírez, M., and Široký, P. 2012b. Phylogenetic position of *Pelusios williamsi* and a critique of current GenBank procedures (Reptilia: Testudines: Pelomedusidae). Amphibia-Reptilia 33:150–154.

Fritz, U., Branch, W.R., Gehring, P.S., Harvey, J., Kindler, C., Meyer, L., Du Preez, L., Široký, P., Vieites, D.R., and Vences, M. 2012c [2013]. Weak divergence among African, Malagasy and Seychellois hinged terrapins (*Pelusios castanoides*, *P. subniger*) and evidence for human-mediated oversea dispersal. Organisms, Diversity and Evolution 13:215–224.

Fritz, U., Gemel, R., Kehlmaier, C., Vamberger, M., and Praschag, P. 2014a. Phylogeography of the Asian softshell turtle *Amyda cartilaginea* (Boddaert, 1770): evidence for a species complex. Vertebrate Zoology 64:229–243.

Fritz, U., Petzold, A., Kehlmaier, C., Kindler, C., Campbell, P., Hofmeyr, M.D., and Branch, W.R. 2014b. Disentangling the *Pelomedusa* complex using type specimens and historical DNA (Testudines: Pelomedusidae). Zootaxa 3795:501–522.

Fritz, U., Mazuch, T., Petzold, A., and Prokop, H. 2015a. Coloration and pattern of hatchlings of six *Pelomedusa* species. Salamandra 51:277–282.

Fritz, U., Kehlmaier, C., Mazuch, T., Hofmeyr, M.D., du Preez, L., Vamberger, M., and Vörös, J. 2015b. Important new records of *Pelomedusa* species for South Africa and Ethiopia. Vertebrate Zoology 65:383–389.

Fritz, U., Tolley, K.A., Vamberger, M., and Ihlow, F. 2022. Phylogeny and phylogeography of chelonians from sub-Saharan Africa—a review of current knowledge in tribute to Margaretha D. Hofmeyr. Vertebrate Zoology 72:951–969.

Fritz, U., Kehlmaier, C., Scott, R.J., Fournier, R., Mccranie, J.R., and Gallego-García, N. 2023. Central American *Trachemys* revisited: new sampling questions current understanding of taxonomy and distribution (Testudines: Emydidae). Vertebrate Zoology 73:513–523.

Fritz, U., Herrmann, H.-W., Rosen, P.C., Auer, M., Vargas-Ramírez, M., and Kehlmaier, C. 2024. *Trachemys* in Mexico and beyond: beautiful turtles, taxonomic nightmare, and a mitochondrial poltergeist (Testudines: Emydidae). Vertebrate Zoology 74:435–452.

Fry, D.B. 1915. On a new *Chelodina* from Australia, with a key to the genus. Proceedings of the Royal Society of Queensland 27:88–90.

Fry, E. 1850. Remarks on the morphology of the vertebrate skeleton. Proceedings of the Zoological Society of London 1850:15–22.

Gable, S.M., Byars, M.I., Literman, R., and Tollis, M. 2022. A genomic perspective on the evolutionary diversification of turtles. Systematic Biology 71(6):1331–1347.

Gadow, H. 1894. On the remains of some gigantic land-tortoises, and of an extinct lizard, recently discovered in Mauritius. Transactions of the Zoological Society of London 13(8):313–324.

Gadow, H. 1905. Distribution of Mexican amphibians and reptiles. Proceedings of the Zoological Society of London 2:191–244.

Gaffney, E.S. 1975. A taxonomic revision of the Jurassic turtles *Portlandemys* and *Plesiochelys*. American Museum Novitates 2574:1–19.

Gaffney, E.S. 1977. The side-necked turtle family Chelidae: a theory of relationships using shared derived characters. American Museum Novitates 2620:1–28.

Gaffney, E.S. 1981. A review of the fossil turtles of Australia. American Museum Novitates 2720:1–38.

Gaffney, E.S. 1983. The cranial morphology of the extinct horned turtle, *Meiolania platyceps*, from the Pleistocene of Lord Howe Island, Australia. Bulletin of the American Museum of Natural History 175(4):326–479.

Gaffney, E.S. 1985. The cervical and caudal vertebrae of the cryptodiran turtle, *Meiolania platyceps*, from the Pleistocene of Lord Howe Island, Australia. American Museum Novitates 2805:1–29.

Gaffney, E.S. 1996. The postcranial morphology of *Meiolania platyceps* and a review of the Meiolaniidae. Bulletin of the American Museum of Natural History 229:1–166.

Gaffney, E.S. and Jenkins, F.A. 2010. The cranial morphology of *Kayentachelys*, an Early Jurassic cryptodire, and the early history of turtles. Acta Zoologica 91:335–368.

Gaffney, E.S. and Meylan, P.A. 1988. A phylogeny of turtles. In: Benton, M.J. (Ed.). The Phylogeny and Classification of the Tetrapods, Volume I: Amphibians, Reptiles, Birds. Systematics Association Special Volume 35A:157–219.

Gaffney, E.S., Balouet, J.C., and Broin, F. de. 1984. New occurrences of extinct meiolaniid turtles in New Caledonia. American Museum Novitates 2800:1–6.

Gaffney, E.S., Tong, H., and Meylan, P.A. 2006. Evolution of the side-necked turtles: the families Bothremydidae, Euraxemydidae, and Araripemydidae. Bulletin of the American Museum of Natural History 300:1–698.

Gaffney, E.S., Rich, T.H., Vickers-Rich, P., Constantine, A., Vacca, R., and Kool, L. 2007. *Chubutemys*, a new eucryptodiran turtle from the early Cretaceous of Argentina, and the relationships of the Meiolaniidae. American Museum Novitates 3599:1–35.

Gaffney, E.S., Meylan, P.A., Wood, R.C., Simons, E., and Almeida Campos, D. de. 2011. Evolution of the side-necked turtles: the family Podocnemididae. Bulletin of the American Museum of Natural History 350:1–237.

Gaillard, D., Ennen, J.R., Kreiser, B.R., Qualls, C.P., Sweat, S.C., Birkhead, R., Tuberville, T.D., Aresco, M., McCoy, E.D., Mushinsky, H.R., and Hentges, T.W. 2017. Range-wide and regional patterns of population structure and genetic diversity in the gopher tortoise. Journal of Fish and Wildlife Management 8(2):497–512.

Gallego-García, N. and Castaño-Mora, O.V. 2008. Ecology and status of the Magdalena River turtle, *Podocnemis lewyana*, a Colombian endemic. Chelonian Conservation and Biology 7(1): 37–44.

Gallego-García, N., Vargas-Ramírez, M., Forero-Medina, G., and Caballero, S. 2018. Genetic evidence of fragmented populations and inbreeding in the Colombian endemic Dahl's toad-headed turtle (*Mesoclemmys dahli*). Conservation Genetics 19:221–233.

Gallego-García, N., Forera-Medina, G., Vargas-Ramírez, M., Caballero, S., and Shaffer, H.B. 2019. Landscape genomic signatures indicate reduced gene flow and forest-associated adaptive divergence in an endangered Neotropical turtle. Molecular Ecology 28:2757–2771.

Gallego-García, N., Ihlow, F., Ettmar, S., Iverson, J.B., and Fritz, U. 2023a.

Where to set the bar? Recent descriptions inflate species number in South American toad-headed turtles (*Mesoclemmys*). Zootaxa 5263(4):566–574.

Gallego-García, N., Vargas-Ramírez, M., and Shaffer, H.B. 2023b. The importance of cryptic diversity in the conservation of wide-ranging species: the red-footed tortoise *Chelonoidis carbonarius* in Colombia. Molecular Ecology 00:1–15. https://doi.org/10.1111/mec.17052.

Gaos, A.R., Lewison, R.L., Liles, M.J., Gadea, V., Altamirano, E., Henríquez, A.V., Torres, P., Urteaga, J., Vallejo, F., Baquero, A., LeMarie, C., Muñoz, J.P., Chaves, J.A., Hart, C.E., Peña de Niz, A., Chácon, D., Fonseca, L., Otterstrom, S., Yañez, I.L., LaCasella, E.L., Frey, A., Jensen, M.P., and Dutton, P.H. 2016. Hawksbill turtle terra incognita: conservation genetics of eastern Pacific rookeries. Ecology and Evolution 6:1251–1264.

Garbin, R.C., Karlguth, D.H., Fernandes, D.S., and Pinto, R.R. 2016. Morphological variation in the Brazilian Radiated Swamp Turtle *Acanthochelys radiolata* (Mikan, 1820) (Testudines: Chelidae). Zootaxa 4105:45–64.

Garbin, R.C., Bandyopadhyay, S., and Joyce, W.G. 2020. A taxonomic revision of geoemydid turtles from Siwalik-age of India and Pakistan. European Journal of Taxonomy 652:1–67.

García-Díaz, P., Ross, J.V., Ayres, C., and Cassey, P. 2015. Understanding the biological invasion risk posed by the global wildlife trade: propagule pressure drives the introduction and establishment of Nearctic turtles. Global Change Biology 21:1078–1091.

Garman, S. 1880. On certain species of Chelonioidae. Bulletin of the Museum of Comparative Zoology 6:123–126.

Garman, S. 1884. The reptiles of Bermuda. In: Jones, J.M. and Goode, G.B. (Eds.). Contributions to the Natural History of the Bermudas. Bulletin of the U.S. National Museum 25:285–303.

Garman, S. 1891. On a tortoise found in Florida and Cuba, *Cinosternum baurii*. Bulletin of the Essex Institute 23:141–144.

Garman, S. 1917. The Galapagos tortoises. Memoirs of the Museum of Comparative Zoology 30(4):261–296.

Garrick, R.C., Benavides, E., Russello, M.A., Gibbs, J.P., Poulakakis, N., Dion, K.B., Hyseni, C., Kajdacsi, B., Márquez, L., Baran, S., Ciofi, C., Tapia, W., and Caccone., A. 2012. Genetic rediscovery of an 'extinct' Galápagos giant tortoise species. Current Biology 22:R10–R11.

Garrick, R.C., Benavides, E., Russello, M.A., Hyseni, C., Edwards, D.L., Gibbs, J.P., Tapia, W., Ciofi, C., and Caccone, A. 2014. Lineage fusion in Galápagos giant tortoises. Molecular Ecology 23:5276–5290.

Garrick, R.C., Kajdacsi, B., Russello, M.A., Benavides, E., Hyseni, C., Gibbs, J.P., Tapia, W., and Caccone, A. 2015. Naturally rare versus newly rare: demographic inferences on two timescales inform conservation of Galápagos giant tortoises. Ecology and Evolution 5:676–694.

Garrido, J.R., Butterfield, T., Scoville, A.G., Macip-Ríos, R., and Beck, D.D. 2021. Comparing semi-urban and forest populations of the Jalisco Mud Turtle (*Kinosternon chimalhuaca*) in Mexico. Herpetological Review 52:725–729.

Garsault, F.A. de. 1764. Les Figures des plantes et animaux d'usage en médecine, décrits dans la Matière Médicale de Mr. Geoffroy Médecin. Paris: 20 pp., 87 pl.

Gasperetti, J., Stimson, A.F., Miller, J.D., Ross, J.P., and Gasperetti, P.R. 1993. Turtles of Arabia. Fauna of Saudi Arabia 13:170–367.

Gaughran, S.J., Gray, R., Ochoa, A., Jones, M., Fusco, N., Miller, J.M., Poulakakis, N., De Queiroz, K., Caccone, A., and Jensen, E.L. 2025. Whole-genome sequencing confirms multiple species of Galapagos giant tortoises. Evolution 79(2):296–308.

Gaur, A., Reddy, A., Annapoorni, S., Satyarebala, B., and Shivaji, S. 2006. The origin of Indian Star tortoises (*Geochelone elegans*) based on nuclear and mitochondrial DNA analysis: a story of rescue and repatriation. Conservation Genetics 2006:231–240.

Ge, Y., Wei, Y., Chek, S.N., and Gong, S. 2018. Diversity and conservation of amphibians and reptiles in Macau, China. Chinese Journal of Wildlife 39(3):639–645.

Gemel, R. and Grillitsch, H. 2008. *Hydrochelys picta* Wagler, 1821: a junior synonym of *Chrysemys picta picta* (Schneider, 1783). Herpetozoa 20(3/4):187–189.

Gemel, R., Gassner, G., and Schweiger, S. 2019. Katalog der Typen der Herpetologischen Sammlung des Naturhistorischen Museums Wien – 2018. Annalen des Naturhistorischen Museums in Wien, B 121:33–248.

Geoffroy Saint-Hilaire, E. 1809a. Mémoire sur les tortues molles. Nouveau Bulletin des Sciences, par la Société Philomatique de Paris 1(22):363–367.

Geoffroy Saint-Hilaire, E. 1809b. Mémoire sur les tortues molles, nouveau genre sous le nom de *Trionyx*, et sur la formation des carapaces. Annales du Muséum d'Histoire Naturelle de Paris 14:1–20.

Georges, A. 1985. Reproduction and reduced body size of reptiles in unproductive insular environments. In: Grigg, G., Shine, R., and Ehmann, H. (Eds.). Biology of Australasian Frogs and Reptiles. Sydney: Royal Zoological Society of New South Wales, pp. 311–318.

Georges, A. 2018. Book review: *Freshwater Turtles of Australia* by John Cann and Ross Sadlier (2017). Australian Journal of Zoology 66:84–87.

Georges, A. 2021. Genetic analysis of species identity of turtles from the Roper River collected as part of the Beetaloo Basin baseline surveys. Report to the Research Institute for the Environment and Livelihoods (RIEL), Charles Darwin University. Biomatix Pty., Ltd.

Georges, A. and Adams, M. 1992. A phylogeny for Australian chelid turtles based on allozyme electrophoresis. Australian Journal of Zoology 40:453–476.

Georges, A. and Adams, M. 1996. Electrophoretic delineation of species boundaries within the short-necked chelid turtles of Australia. Zoological Journal of the Linnean Society 118:241–260.

Georges, A. and Kennett, R.M. 1989. Dry-season distribution and ecology of *Carettochelys insculpta* (Chelonia: Carettochelyidae) in Kakadu National Park, Northern Australia. Australian Wildlife Research 16:323–335.

Georges, A. and Thomson, S. 2006. Evolution and zoogeography of Australian freshwater turtles. In: Merrick, J.R., Archer, M., Hickey, G.M., and Lee, M.S.Y. (Eds.). Evolution and Biogeography of Australasian Vertebrates. Sydney: Australian Scientific Publishing, pp. 291–308.

Georges, A. and Thomson, S. 2010. Diversity of Australasian freshwater turtles, with an annotated synonymy and keys to species. Zootaxa 2496:1–37.

Georges, A., Birrell, J., Saint, K., McCord, W.P., and Donnellan, S. 1998. A phylogeny for side-necked turtles (Chelonia: Pleurodira) based on mitochondrial and nuclear gene sequence variation. Biological Journal of the Linnean Society 67:213–246.

Georges, A., Doody, S., Young, J., and Cann, J. 2000. The Australian Pig-Nosed Turtle. Canberra: Robey, 37 pp.

Georges, A., Adams, M., and McCord, W. 2002. Electrophoretic delineation of species boundaries within the genus *Chelodina* (Testudines: Chelidae) of Australia, New Guinea and Indonesia. Zoological Journal of the Linnean Society 134:401–421.

Georges, A., Guarino, F., and Bito, B. 2006. Freshwater turtles of the TransFly region of Papua New Guinea – notes on diversity, distribution, reproduction, harvest and trade. Wildlife Research 33:373–384.

Georges, A., Zhang, X., Unmack, P., Reid, B.N., Le, M., and McCord, W.P. 2014. Contemporary genetic structure of an endemic freshwater turtle reflects Miocene orogenesis of New Guinea. Biological Journal of the Linnean Society 111(1):192–208.

Georges, A., Gruber, B., Pauly, G.B., White, D., Adams, M., Young, M.J., Kilian, A., Zhang, X., Shaffer, H.B., and Unmack, P.J. 2018a. Genomewide SNP markers breathe new life into phylogeography and species delimitation for the problematic short-necked turtles (Chelidae: *Emydura*) of eastern Australia. Molecular Ecology 27:5195–5213.

Georges, A., Spencer, R.-J., Killian, A., and Zhang, X. 2018b. Assault from all sides: hybridization and introgression threaten the already critically endangered *Myuchelys georgesi* (Chelonia: Chelidae). Endangered Species Research 37:239–247.

Georges, A., Wedd, D., and Dostine, P. 2021. Genetic analysis of species identity of turtles from the Roper River collected as part of the Beetaloo Basin Geological and Bioregional Assessment (GBA). Report to the Research Institute for the Environment and Livelihoods (RIEL), Charles Darwin University, in support of the Beetaloo GBA Program, 15 pp.

Gerholdt, J.E. and Oldfield, B. 1987. *Chelydra serpentina serpentina* (Common Snapping Turtle). Size. Herpetological Review 18(4):73.

Gerlach, J. 2001. Tortoise phylogeny and the '*Geochelone*' problem. Phelsuma 9a:1–24.

Gerlach, J. 2004. Giant Tortoises of the Indian Ocean. The genus *Dipsochelys* inhabiting the Seychelles Islands and the extinct giants of Madagascar and the Mascarenes. Frankfurt: Chimaira, 208 pp.

Gerlach, J. 2011a. Development of distinct morphotypes in captive Seychelles–Aldabra giant tortoises. Chelonian Conservation and Biology 10(1):102–112.

Gerlach, J. and Bour, R. 2003. Morphology of hatchling giant tortoises. Radiata 12:11–12.

Gerlach, J. and Canning, K.L. 1996. The Seychelles giant tortoise, its rediscovery

and prospects for conservation. In: Devaux, B. (Ed.). Proceedings – International Congress of Chelonian Conservation. Gonfaron, France: Editions SOPTOM, pp. 133–135.

Gerlach, J. and Canning, K.L. 1998. Taxonomy of Indian Ocean giant tortoises (*Dipsochelys*). Chelonian Conservation and Biology 3(1):3–19.

Gerlach, J. and Canning, L. 2001. Range contractions in the critically endangered Seychelles terrapins (*Pelusios* spp.). Oryx 35:313–321.

Gerlach, J. and Paquette, S.R. 2014. Evolution of the tortoises of the western Indian Ocean. In: Gerlach, J. (Ed.). Western Indian Ocean Tortoises: Ecology, Diversity, Evolution, Conservation, Paleontology. Manchester, United Kingdom: Siri Scientific Press, pp. 9–30.

Gerlach, J., Rocamora, G., Gane, J., Jolliffe, K., and Vanherck, L. 2013. Giant tortoise distribution and abundance in the Seychelles Islands: past, present, and future. Chelonian Conservation and Biology 12(1):70–83.

Germano, D.J. and Riedle, J.D. 2015. Population structure, growth, survivorship, and reproduction of *Actinemys marmorata* from a high elevation site in the Tehachapi Mountains, California. Herpetologica 71:102–109.

Gervais, P. 1843. Dictionnaire Universal d'Histoire Naturelle. Vol. 3, p. 457.

Gesner, C. 1558. Historiae Animalium, Liber IIII. Piscium et Aquatilium Animantium Natura. Tiguri: Christoph. Froschoverum, 1297 pp.

Gibbons, J.W. and Lovich, J.E. 1990. Sexual dimorphism in turtles with emphasis on the slider turtle (*Trachemys scripta*). Herpetological Monographs 4:1–29.

Gibbons, J.W., Novak, S.S., and Ernst, C.H. 1988. *Chelydra serpentina*. Catalogue of American Amphibians and Reptiles 420:1–4.

Gibbons, J.W., Lovich, J.E., Tucker, A.D., FitzSimmons, N.N., and Greene, J.L. 2001. Demographic and ecological factors affecting conservation and management of the diamondback terrapin (*Malaclemys terrapin*) in South Carolina. Chelonian Conservation and Biology 4(1):66–74.

Gibbs, J.P., Breisch, A.R., Ducey, P.K., Johnson, G., Behler, J., and Bothner, R. 2007. The Amphibians and Reptiles of New York State: Identification, Natural History, and Conservation. New York: Oxford University Press, 422 pp.

Gibbs, J.P., Cayot, L.J., and Tapia Aguilera, W. (Eds.). 2020. Galapagos Giant Tortoises. London: Academic Press, 518 pp.

Gidis, M., Spinks, P.Q., Çevik, E., Kaska, Y., and Shaffer, H.B. 2011. Shallow genetic divergence indicates a Congo–Nile riverine connection for the softshell turtle *Trionyx triunguis*. Conservation Genetics 12:589–594.

Giebel, C.G. 1866a. *Cistudo anhaltina* n. sp. aus der Latdorfer Braunkohle. Zeitschrift für die Gesammten Naturwissenschaften 27:1–11.

Giebel, C.G. 1866b. Die Schildkröten der Insel Banka. Zeitschrift für die Gesammten Naturwissenschaften 27:11–21.

Gillette, D.D. 1977. Catalogue of type specimens of fossil vertebrates, Academy of Natural Sciences, Philadelphia. Part IV: Reptilia, Amphibia, and Tracks. Proceedings of the Academy of Natural Sciences of Philadelphia 129:101–111.

Gilmore, C.W. 1923. A new fossil turtle, *Kinosternon arizonense*, from Arizona. Proceedings of the United States National Museum 62:1–8.

Gilmore, C.W. 1927. On fossil turtles from the Pleistocene of Florida. Proceedings of the United States National Museum 71:1–10.

Giraldo, A., Carr, J.L., and Garcés Restrepo, M.F. 2012. *Rhinoclemmys annulata* (Gray 1860). In: Páez, V.P., Morales-Betancourt, M.A., Lasso, C.A., Castaño-Mora, O.V., and Bock, B.C. (Eds.). Biología y Conservación de las Tortugas Continentales de Colombia. Bogotá, Colombia: Instituto de Investigación de Recursos Biológicos Alexander von Humboldt, pp. 300–304.

Girard, C. 1858. United States Exploring Expedition during the years 1838, 1839, 1840, 1841, 1842. Under the command of Charles Wilkes, U.S.N. Volume XX. Herpetology. Philadelphia: J.B. Lippincott, 496 pp.

Girondot, M. and Fretey, J. 1996. Leatherback turtles, *Dermochelys coriacea*, nesting in French Guiana, 1978–1995. Chelonian Conservation and Biology 2(2):204–208.

Girondot, M., Mourrain, B., Chevallier, D., and Godfrey, M.H. 2021. Maturity of a giant: age and size reaction norm for sexual maturity for Atlantic leatherback turtles. Marine Ecology 2021; 00:e12631, 11 pp.

Gistel, J. 1848. Naturgeschichte des Thierreichs für höhere Schulen bearbeitet durch Johannes Gistel...mit einem Atlas von 32 Tafeln (darstelland 617 illuminierte Figuren) und mehrern dem Texte eingedruckten Xylographien. Stuttgart: 216 pp.

Gistel, J. 1868. Die Lurche Europa's. Ein Beitrag zur Lehre von der geographischen Verbreitung derselben. In: Gistel, J. Blicke in das Leben der Natur und des Menschen. Leipzig: Verlag Ed. Wartig, pp. 144–167.

Glass, B.P. and Hartweg, N. 1951. *Kinosternon murrayi*, a new musk turtle of the *hirtipes* group from Texas. Copeia 1951(1):50–52.

Glauert, L. 1923. A new freshwater tortoise from the Murchison River. Journal of the Royal Society of Western Australia 9:53–56.

Glauert, L. 1954. A new swamp tortoise from the Swan River district. Western Australian Naturalist 4:125–127.

Glorioso, B.M., Selman, W., Kreiser, B.R., and Ford, A. 2024. Hiding in plain sight: federally protected Ringed Map Turtles (*Graptemys oculifera*) found in a new river system. Herpetological Conservation and Biology 19(1):96–105.

Gmelin, J.F. 1789 ["1788"]. Caroli a Linné, Systema Naturae per Regna Tria Naturae secundum Classes, Ordines, Genera, Species, cum Characteribus, Differentiis, Synonymis, Locis. Ed. 13. Tom. I. Pars III. Leipzig: G.E. Beer, Ed. 13, 1(3):1033–1516.

Gmelin, S.G. 1774. Reise durch Russland zur Untersuchung der drey Natur-Reiche. Dritter Theil. Reise durch das nordliche Persien, in den Jahren 1770, 1771, bis im April 1772. St. Petersburg: Kayserliche Academie der Wissenschaften, 508 pp.

Gmira, S. 1993. Une nouvelle espéce de tortue Testudininei (*Testudo kenitrensis* n. sp.) de l'inter Amirien-Tensiftien de Kénitra (Maroc). Comptes Rendus de l'Académie des Sciences Série II 316:701–707.

Gmira, S. 1995. Étude des Chéloniens Fossiles du Maroc. Paris: Cahier de Paléontologie, 140 pp.

Gnetneva, A.N. 2020. Systematics and distribution of turtles of the genus *Agrionemys*. Dissertation, Zoological Institute of the Russian Academy of Sciences, St. Petersburg.

Godwin, J.C., Lovich, J.E., Ennen, J.R., Kreiser, B.R., Folt, B., and Lechowicz, C. 2014. Hybridization of two megacephalic map turtles (Testudines: Emydidae: *Graptemys*) in the Choctawhatchee River drainage of Alabama and Florida. Copeia 2014:725–742.

Goetz, M. 2007. Husbandry and breeding of the Spiny Turtle *Heosemys spinosa* (Gray, 1931 [sic]) at the Durrell Wildlife Conservation Trust. Radiata 16(2):2–15.

Göldi, E.A. 1886. Ueber eine vermuthlich neue Schildkröte der Gattung *Podocnemis* vom Rio Negro und über die Chelonier des Amazonas-Gebietes im Allgemeinen. St. Gallische Naturwissenschaftliche Gesellschaft, Bericht über die Thäligheit 1884–1885[1886]:273–280.

Gong, S., Shi, H., Xu, R., Wang, Z., Xu, J., and Zhang, Y. 2006. A survey of freshwater turtles in Jianfengling Nature Reserve, Hainan Province, China. Chinese Journal of Zoology 41(1):80–83.

Gong, S., Shi, H., Mo, Y., Auer, M., Vargas-Ramírez, M., Hundsdörfer, A.K., and Fritz, U. 2009a. Phylogeography of the endangered black-breasted leaf turtle (*Geoemyda spengleri*) and conservation implications for other chelonians. Amphibia-Reptilia 30:57–62.

Gong, S., Shi, H., Fong, J.J., and Lau, M. 2009b. Recent records of freshwater turtles from Guangdong, China. Turtle and Tortoise Newsletter 13:24–27.

Gong, S., Ge, Y., Chen, G., Liu, L., Chen, Y., and Lau, M. 2012. Diversity and conservation of reptiles in Dinghushan National Nature Reserve, Guangdong, China. Sichuan Journal of Zoology 31(3):483–487.

Gong, S., Vamberger, M., Auer, M., Praschag, P., and Fritz, U. 2018. Millennium-old farm breeding of Chinese softshell turtles (*Pelodiscus* spp.) results in massive erosion of biodiversity. The Science of Nature 105(34):1–10.

Gong, S., Fritz, U., Vamberger, M., Gao, Y., and Farkas, B. 2022. Disentangling the *Pelodiscus axenaria* complex, with the description of a new Chinese species and neotype designation for *P. axenaria* (Zhou, Zhang & Fang, 1991). Zootaxa 5125(2):131–143.

Gong, S., Suwannapoom, C., Le, M., Nguyen, T.Q., Ge, Y., Wei, Y., and Gao, Y. 2023. Genomic analyses reveal three phylogenetic species and their evolutionary histories in the big-headed turtle. iScience 26:107343, 15 pp.

Gong, S., Li, J., Fritz, U., and Farkas, B. 2025. When light casts long shadows: the increasingly dark truth about *Pelodiscus axenaria* (Zhou, Zhang & Fang, 1991). Zootaxa 5604(1):94–100.

Gong, Y.A., Peng, L.F., Huang, S., Lin, Y.F., Huang, R.Y., Xu, Y.H., Yang, D.C., and Nie, L.W. 2021. A new species of the Genus *Pelodiscus* Fitzinger, 1835 (Testudines: Trionychidae) from Huangshan, Anhui, China. Zootaxa 5060(1):137–145.

Gonzales, L. and Reichle, S. 2002. Singularidades de la herpetofauna de la serranía Santiago de Chiquitos, Santa Cruz - Bolivia. Revista Boliviana de Ecología 11:77–85.

González-Porter, G.P., Hailer, F., Flores-Villela, O.A., Garcia-Anleu, R., and Maldonado, J.E. 2011. Patterns of genetic diversity in the critically endangered Central American river turtle: human influence since the Mayan

age? Conservation Genetics 12:1229–1242.

González-Porter, G.P., Maldonado, J.E., Flores-Villela, O., Vogt, R.C., Janke, A., Fleischer, R.C., and Hailer, F. 2013. Cryptic population structuring and the role of the Isthmus of Tehuantepec as a gene flow barrier in the Critically Endangered Central American River Turtle. PLoS ONE 8(9):e71668.

Goode, J. 1967. Freshwater Tortoises of Australia and New Guinea (in the Family Chelidae). Melbourne: Lansdowne Press, 154 pp.

Gosnell, J.S., Rivera, G., and Blob, R.W. 2009. A phylogenetic analysis of sexual size dimorphism in turtles. Herpetologica 65:70–81.

Graciá, E., Vargas-Ramírez, M., Delfino, M., Anadón, J.D., Giménez, A., Fahd, S., Corti, C., Jdeidi, T.B., and Fritz, U. 2017a. Expansion after expansion: dissecting the phylogeography of the widely distributed spur-thighed tortoise, *Testudo graeca* (Testudines: Testudinidae). Biological Journal of the Linnean Society 121:641–654.

Graciá, E., Rodríguez-Caro, R.C., Andreu, A.C., Fritz, U., Giménez, A., and Botella, F. 2017b. Human-mediated secondary contact of two tortoise lineages results in sex-biased introgression. Scientific Reports 7(4019):1–11.

Graham, T.E. 1991. *Pseudemys rubriventris*. Catalogue of American Amphibians and Reptiles 510:1–4.

Graham, T.E. 1995. Habitat use and population parameters of the spotted turtle, *Clemmys guttata*, a Species of Special Concern in Massachusetts. Chelonian Conservation and Biology 1(3):207–214.

Gramentz, D. 1998. Zur Morphologie und Merkmalsvariation von *Cycloderma aubryi* (Duméril, 1856). Salamandra 34(4):333–348.

Gramentz, D. 2008. African Flapshell Turtles—The Genera *Cyclanorbis* and *Cycloderma*. Frankfurt: Edition Chimaira, 191 pp.

Gramentz, D. 2011. *Lissemys punctata*: The Indian Flapshell Turtle. Frankfurt: Edition Chimaira, 280 pp.

Grandidier, A. 1867. Liste des reptiles nouveaux découverts, en 1866, sur la côte sud-ouest de Madagascar. Revue et Magazine de Zoologie, Paris (2)19:223–234.

Grandidier, A. 1868. Sur les découvertes zoologiques faites récemment à Madagascar. Annales des Sciences Naturelles. Zoologie et Paléontologie, Paris (5)10:375–378.

Grandidier, A. 1869. Descriptions d'un Rhinolophe et d'une Tortue de Madagascar. Revue et Magazine de Zoologie, Paris (2)21:257–258.

Gravenhorst, J.L.C. 1829. Deliciae Musei Zoologici Vratislaviensis. Fasciculus primus, continens chelonios et batrachia. Lipsiae: Leopoldi Vossii, 106 pp.

Gray, J.E. 1825. A synopsis of the genera of reptiles and amphibia, with a description of some new species. Annals of Philosophy (2)10:193–217.

Gray, J.E. 1828. Spicilegia Zoologica; or original figures and short systematic descriptions of new and unfigured animals. Part I. London: Richard Taylor, 8 pp.

Gray, J.E. 1830a. Illustrations of Indian Zoology, chiefly selected from the collection of Major-General Hardwicke. Vol. I, Part 1, pl. 77. London: Treuttel, Wurtz, Treuttel, Jun. and Richter. [Published Jan 1830].

Gray, J.E. 1830b. Illustrations of Indian Zoology, chiefly selected from the collection of Major-General Hardwicke. Vol. I, Part 2, pl. 72. London: Treuttel, Wurtz, Treuttel, Jun. and Richter. [Published Mar 1830].

Gray, J.E. 1830c. [*Emys occipitatis; Emys ornata*]. In: Griffith E. and Pidgeon, E. The Class Reptilia arranged by the Baron Cuvier, with specific descriptions. In: Griffith, E. (Ed.). The Animal Kingdom Arranged in Conformity with its Organization, by the Baron Cuvier, with Additional Descriptions of all the Species Hitherto Named, and of many not before Noticed. Vol. 9. Reptilia. London: Whittaker, Treacher, and Co., pp. 75–76. [Part 25, published Sep 1830].

Gray, J.E. 1830d. Illustrations of Indian Zoology, chiefly selected from the collection of Major-General Hardwicke. Vol. I, Part 4, pls. 75, 78. London: Treuttel, Wurtz, Treuttel, Jun. and Richter. [Published Oct 1830].

Gray, J.E. 1830e. A Synopsis of the Species of the Class Reptilia. In: Griffith E. and Pidgeon, E. The Class Reptilia arranged by the Baron Cuvier, with specific descriptions. In: Griffith, E. (Ed.). The Animal Kingdom Arranged in Conformity with its Organization, by the Baron Cuvier, with Additional Descriptions of all the Species Hitherto Named, and of many not before Noticed. Vol. 9. Reptilia. Supplement. London: Whittaker, Treacher, and Co., 110 pp. [Part 26, published Dec 1830].

Gray, J.E. 1831a. Illustrations of Indian Zoology, chiefly selected from the collection of Major-General Hardwicke. Vol. I, Part 5, pl. 74. London: Treuttel, Wurtz, Treuttel, Jun. and Richter. [Published Jan 1831].

Gray, J.E. 1831b. Illustrations of Indian Zoology, chiefly selected from the collection of Major-General Hardwicke. Vol. I, Part 6, pl. 76. London: Treuttel, Wurtz, Treuttel, Jun. and Richter. [Published Apr 1831].

Gray, J.E. 1831c. A specimen of a tortoise regarded as the type of a new genus in the family Emydidae. Proceedings of the Zoological Society of London 1831(1):106–107. [Published May 1831].

Gray, J.E. 1831d. Synopsis Reptilium; or Short Descriptions of the Species of Reptiles. Part I.—Cataphracta. Tortoises, Crocodiles, and Enaliosaurians. London: Treuttel, Wurz, and Co., 85 pp. [Published May 1831].

Gray, J.E. 1831e. Illustrations of Indian Zoology, chiefly selected from the collection of Major-General Hardwicke. Vol. I, Part 7, pls. 73, 79. London: Treuttel, Wurtz, Treuttel, Jun. and Richter. [Published Jul 1831].

Gray, J.E. 1831f. Illustrations of Indian Zoology, chiefly selected from the collection of Major-General Hardwicke. Vol. I, Part 8, pl. 80. London: Treuttel, Wurtz, Treuttel, Jun. and Richter. [Published Oct 1831].

Gray, J.E. 1832a. Illustrations of Indian Zoology, chiefly selected from the collection of Major-General Hardwicke. Vol. I, Part 10, Direction. London: Treuttel, Wurtz, Treuttel, Jun. and Richter. [Published Apr 1832].

Gray, J.E. 1832b. Illustrations of Indian Zoology, chiefly selected from the collection of Major-General Hardwicke. Vol. II, Part 11, pl. 60. London: Adolphus Richter and Co. [Published Jul 1832].

Gray, J.E. 1834a. Characters of several new species of freshwater tortoises (*Emys*) from India and China. Proceedings of the Zoological Society of London 1834(2):53–54.

Gray, J.E. 1834b. Characters of two new genera of reptiles (*Geoemyda* and *Gehyra*). Proceedings of the Zoological Society of London 1834(2):99–101.

Gray, J.E. 1835. Illustrations of Indian Zoology, chiefly selected from the collection of Major-General Hardwicke. Vol. II, Parts 19-20, pls. 57, 65-66, Direction. London: Adolphus Richter and Co. [Published Feb 1835].

Gray, J.E. 1841. A catalogue of the species of reptiles and amphibia hitherto described as inhabiting Australia, with a description of some new species from Western Australia, and some remarks on their geographical distribution. In: Grey, G. Journals of Two Expeditions of Discovery in Northwest and Western Australia. London: T. and W. Boone, Vol. 2. Appendix E, pp. 422–449.

Gray, J.E. 1842. Description of some hitherto unrecorded species of Australian reptiles and batrachians. Zoological Miscellany 2:51–57.

Gray, J.E. 1844. Catalogue of the Tortoises, Crocodiles, and Amphisbaenians in the Collection of the British Museum. London: Edward Newman, 80 pp.

Gray, J.E. 1847. Description of a new genus of Emydae. Proceedings of the Zoological Society of London 1847(15):55–56.

Gray, J.E. 1849. Description of a new species of box tortoise from Mexico. Proceedings of the Zoological Society of London 1849:16–17.

Gray, J.E. 1854a. Description of a new genus and some new species of tortoises. Proceedings of the Zoological Society of London 1852[1854]:133–135.

Gray, J.E. 1854b. Description of a new species of tortoise (*Testudo planiceps*), from the Galapagos Islands. Proceedings of the Zoological Society of London 1853[1854]:12–13.

Gray, J.E. 1856a. On some new species of freshwater tortoises from North America, Ceylon and Australia, in the collection of the British Museum. Proceedings of the Zoological Society of London 1855[1856](23):197–202. [Published Feb 1856].

Gray, J.E. 1856b ["1855"]. Catalogue of Shield Reptiles in the Collection of the British Museum. Part I. Testudinata (Tortoises). London: British Museum, 79 pp. [Published Mar 1856].

Gray, J.E. 1857. Description of a new species of *Chelodina* from Australia. Proceedings of the Zoological Society of London 1856[1857]:369–371.

Gray, J.E. 1859. Description of a new species of freshwater tortoise from Siam. Proceedings of the Zoological Society of London 1859(27):478–479.

Gray, J.E. 1860a. Description of a soft tortoise (*Aspidochelys livingstonii*) from the Zambesi, sent to the British Museum by Dr. Livingstone. Proceedings of the Zoological Society of London 1860(28):5–6.

Gray, J.E. 1860b. Description of a new species of *Geoclemmys* from Ecuador. Proceedings of the Zoological Society of London 1860(28):231–232.

Gray, J.E. 1860c. Description of a new species of *Emys* lately living in the gardens of the Zoological Society. Proceedings of the Zoological Society of London 1860(28):232–233.

Gray, J.E. 1860d. On some new species of Mammalia and tortoises from Cambojia. Annals and Magazine of Natural History (3)6:217–218.

Gray, J.E. 1861a. Description of a soft tortoise from Camboja. Proceedings of the Zoological Society of London 1861:41–42.

Gray, J.E. 1861b. On a new species of water–tortoise (*Geoclemmys melanosterna*)

from Darien. Proceedings of the Zoological Society of London 1861:204–205.

Gray, J.E. 1862a. Notice of a new species of *Cyclemys* from the Lao Mountains, in Siam. Annals and Magazine of Natural History (3)10:157.

Gray, J.E. 1862b. Notice of two new species of *Batagur* in the collection of the British Museum. Proceedings of the Zoological Society of London 1862:264–265.

Gray, J.E. 1862c. Notice of a new species of *Dogania* from Asia. Proceedings of the Zoological Society of London 1862:265–266.

Gray, J.E. 1863a. On the species of *Chelymys* from Australia; with the description of a new species. Annals and Magazine of Natural History (3)12:98–99.

Gray, J.E. 1863b. Notice of a new species of *Pelomedusa* from Natal. Annals and Magazine of Natural History (3)12:99–100.

Gray, J.E. 1863c. Notes on American Emydidae, and Professor Agassiz's observations on my catalogue of them. Annals and Magazine of Natural History (3)12:176–183.

Gray, J.E. 1863d. Notice of a new species of *Kinixys* and other tortoises from central Africa. Annals and Magazine of Natural History (3)12:381–382.

Gray, J.E. 1863e. Observations on the box tortoises, with the descriptions of three new Asiatic species. Proceedings of the Zoological Society of London 1863:173–179.

Gray, J.E. 1863f. On the species of the genus *Sternothaerus*, with some observations on *Kinixys*. Proceedings of the Zoological Society of London 1863:192–197.

Gray, J.E. 1863g. Notice of a new species of *Batagur* from north-western India. Proceedings of the Zoological Society of London 1863:253.

Gray, J.E. 1863h. Description of a new *Geoclemys* lately living in the Gardens of the Zoological Society. Proceedings of the Zoological Society of London 1863:254–255.

Gray, J.E. 1864a. Notes on certain species of tortoises from the Asiatic Islands transmitted to the British Museum by Dr. Bleeker. Proceedings of the Zoological Society of London 1864:11–13.

Gray, J.E. 1864b. Revision of the species of Trionychidae found in Asia and Africa, with the descriptions of some new species. Proceedings of the Zoological Society of London 1864:76–98.

Gray, J.E. 1864c. Description of a new species of *Staurotypus* (*S. salvinii*) from Guatemala. Proceedings of the Zoological Society of London 1864:127–128.

Gray, J.E. 1864d. On the genera of Chelydidae and the characters furnished by the study of their skulls. Proceedings of the Zoological Society of London 1864:128–135.

Gray, J.E. 1865a. Notice of a new genus and species of the family Trionychidae from Western Africa. Annals and Magazine of Natural History (3)16:204–206.

Gray, J.E. 1865b. On the development of the sternal callosities in *Cyclanosteus senegalensis*, and on the synonyms of *Cyclanosteus* and its allied genera. Proceedings of the Zoological Society of London 1865:422–428.

Gray, J.E. 1867. Description of a new Australian tortoise (*Elseya latisternum*). Annals and Magazine of Natural History (3)20:43–45.

Gray, J.E. 1868. Notice of *Hydraspis gordoni*, a new species from Trinidad, living in the gardens of the Society. Proceedings of the Zoological Society of London 1868:563–564.

Gray, J.E. 1869a. Notes on the families and genera of tortoises (Testudinata), and on the characters afforded by the study of their skulls. Proceedings of the Zoological Society of London 1869:165–225.

Gray, J.E. 1869b. Description of *Mauremys laniaria*, a new freshwater tortoise. Proceedings of the Zoological Society of London 1869:499–500.

Gray, J.E. 1869c. Description of *Emys flavipes*. Proceedings of the Zoological Society of London 1869:643–644.

Gray, J.E. 1870a. Notice of a new Chilian tortoise (*Testudo chilensis*). Annals and Magazine of Natural History (4)6:190.

Gray, J.E. 1870b. Notes on tortoises in the British Museum, with descriptions of some new species. Proceedings of the Zoological Society of London 1870:653–659.

Gray, J.E. 1870c. Supplement to the Catalogue of Shield Reptiles in the Collection of the British Museum. Part I. Testudinata (Tortoises). London: British Museum, 120 pp.

Gray, J.E. 1870d. Notes on three species of tortoises living in the Society's gardens. Proceedings of the Zoological Society of London 1870:706–708.

Gray, J.E. 1870e. On the family Dermatemydae, and a description of a living species in the gardens of the Society. Proceedings of the Zoological Society of London 1870:711–716.

Gray, J.E. 1870f. Notes on *Bartlettia*, a new species of freshwater tortoises belonging to the family Peltocephalidae. Proceedings of the Zoological Society of London 1870:718–721.

Gray, J.E. 1871a. On *Euchelymys*, a new genus and two new species of Australian freshwater tortoises. Annals and Magazine of Natural History (4)8:117–118.

Gray, J.E. 1871b. Notes on Australian freshwater tortoises. [1]. Annals and Magazine of Natural History (4)8:291–292.

Gray, J.E. 1871c. Notes on Australian freshwater tortoises. [2]. Annals and Magazine of Natural History (4)8:366.

Gray, J.E. 1871d. *Damonia oblonga*, a new species of freshwater tortoise. Annals and Magazine of Natural History (4)8:367.

Gray, J.E. 1872a. Notes on the mud-tortoises of India (*Trionyx*, Geoffroy). Annals and Magazine of Natural History (4)10:326–340.

Gray, J.E. 1872b. On *Spatulemys lasalae*, a new genus of Hydraspidae from Rio Parana, Corrientes. Annals and Magazine of Natural History (4)10:463.

Gray, J.E. 1872c. Appendix to the Catalogue of Shield Reptiles in the Collection of the British Museum. Part I. Testudinata (Tortoises). London: British Museum, 28 pp.

Gray, J.E. 1872d. On the genus *Chelymys* and its allies from Australia. Proceedings of the Zoological Society of London 1872:504–514.

Gray, J.E. 1873a. Notes on tortoises. Annals and Magazine of Natural History (4)11:143–149.

Gray, J.E. 1873b. On a new freshwater tortoise from Borneo (*Orlitia borneensis*). Annals and Magazine of Natural History (4)11:156–157.

Gray, J.E. 1873c. Observations on chelonians, with descriptions of new genera and species. Annals and Magazine of Natural History (4)11:289–308.

Gray, J.E. 1873d. Notes on the family Chelydradae. Annals and Magazine of Natural History (4)12:66–70.

Gray, J.E. 1873e. *Damonia unicolor*, a new species of water-tortoise from China, sent by Mr. Swinhoe. Annals and Magazine of Natural History (4)12:77–78.

Gray, J.E. 1873f. Notes on the tortoises of the 'Zoology of Mexico' of MM. A. Duméril and Bocourt. Annals and Magazine of Natural History (4)12:109–114.

Gray, J.E. 1873g. Notes on Chinese mud-tortoises (Trionychidae), with the description of a new species sent to the British Museum by Mr. Swinhoe, and observations on the male organ of this family. Annals and Magazine of Natural History (4)12:156–161.

Gray, J.E. 1873h. Notes on mud-tortoises (*Trionyx*, Geoffroy), and on the skulls of the different kinds. Proceedings of the Zoological Society of London 1873:38–72.

Gray, J.E. 1873i. Notes on the genera of turtles (Oiacopodes), and especially on their skeletons and skulls. Proceedings of the Zoological Society of London 1873:395–411.

Gray, J.E. 1873j. Hand-List of the Specimens of Shield Reptiles in the British Museum. London: British Museum, 124 pp.

Gray, J.E. 1873k ["1872"]. List of the Books, Memoirs, and Miscellaneous Papers by Dr. John Edward Gray, F.R.S., with a few Historical Notes. London: Private Distribution, 55 pp.

Gray, J.E. 1874 ["1873"]. On the skulls and alveolar surfaces of land-tortoises (Testudinata). Proceedings of the Zoological Society of London 1873[1874]:722–728.

Gregory, S.M.S. 2010. The two 'editions' of Duméril's Zoologie analytique, and the potential confusion caused by Froriep's translation Analytische Zoologie. Zoological Bibliography 1(1):6–8.

Grew, N. 1681. Musaeum Regalis Societatis, or a Catalogue and Description of the Natural and Artificial Rarities Belonging to the Royal Society and Preserved at Gresham Colledge. London: W. Rawlins, 386 pp.

Griffith, E. and Pidgeon, E. 1830. The Class Reptilia Arranged by the Baron Cuvier, with Specific Descriptions. In: Griffith, E. (Ed.). The Animal Kingdom Arranged in Conformity with its Organization, by the Baron Cuvier, with Additional Descriptions of all the Species Hitherto Named, and of many not before Noticed. Vol. 9. Reptilia. London: Whittaker, Treacher, and Co., 481 pp. [Part 25, pp. 1–192, published Sep 1830].

Gronovius, L.T. 1756. Amphibiorum Animalium Historia Zoologica. In: Museum Ichthyologicum, sistens Piscium Indigenorum & Quorumdam Exoticorum... in Museo Laurentii Theodori Gronovii... et Amphibiorum Animalium Historia Zoologica. Tomus secundus. Leiden, T. Haak, pp. 47–88.

Gronovius, L.T. 1763. Zoophylacium Gronovianum, exhibens animalia. Fasciculus primus. Leiden: pp.1–136.

Grossmann, W. and Grychta, U. 1998. Bemerkungen zur Grösse der

Malayen-Weichschildkröte *Dogania subplana* (Geoffroy, 1809). Sauria (Berlin) 20(2):11–13.

Groves, J.D. 1983. Taxonomic status and zoogeography of the painted turtle *Chrysemys picta* (Testudines: Emydidae), in Maryland. American Midland Naturalist 109:274–279.

Grünewald, F. 2008. Observations on the South-American Snapping Turtle *Chelydra acutirostris* Peters, 1862 in Ecuador. Emys 15(3):33–40.

Güçlü, Ö., Ulger, C., Türkozan, O., Gemel, R., Reimann, M., Levy, Y., Ergene, S., Uçar, A.H., and Aymak, C. 2009. First assessment of mitochondrial DNA diversity in the endangered Nile Softshell Turtle, *Trionyx triunguis*, in the Mediterranean. Chelonian Conservation and Biology 8(2):222–226.

Guedes, P., dos Santos, Y., Matilde, E., Alves, J., Rato, C., and Rocha, R. 2023. Phylogenetic relationships of the West African mud turtle (*Pelusios castaneus*) on the islands of São Tomé and Príncipe, West Central Africa. Amphibia-Reptilia 2023; doi:10.1163/15685381-bja10145, 7 pp.

Guérin, F.E. 1829. Iconographie du Règne Animal de G. Cuvier. I. Planches des animaux vertébrés. Paris: J.B. Baillibre, Reptiles, planches 1–30.

Guicking, D., Fritz, U., Wink, M., and Lehr, E. 2002. New data on the diversity of the Southeast Asian leaf turtle genus *Cyclemys* Bell, 1834. Molecular results (Reptilia: Testudines: Geoemydidae). Faunistische Abhandlungen Staatliches Museum für Tierkunde Dresden 23:75–86.

Guillon, J.M., Guéry, L., Hulin, V., and Girondot, M. 2012. A large phylogeny of turtles (Testudines) using molecular data. Contributions to Zoology 81(3):147–158.

Guyer, C., Bailey, M.A., and Mount, R.H. 2015. Turtles of Alabama. Tuscaloosa, AL: University of Alabama Press, 288 pp.

Günther, A.C.L.G. 1860. On the reptiles of Siam. Proceedings of the Zoological Society of London 1860:113–117.

Günther, A.C.L.G. 1864. The Reptiles of British India. London: Ray Society, Robert Hardwicke, 452 pp.

Günther, A.C.L.G. 1869. Report on two collections of Indian reptiles. Proceedings of the Zoological Society of London 1869:500–507.

Günther, A.C.L.G. 1873a. Preliminary notice of some extinct tortoises from the islands of Rodriguez and Mauritius. Annals and Magazine of Natural History (4)11:397.

Günther, A.C.L.G. 1873b. Reptilia. The Zoological Record 8:77–88.

Günther, A.C.L.G. 1874. Descriptions of the living and extinct races of gigantic land-tortoises. Parts I and II. Introduction, and the tortoises of the Galapagos Islands. [Abstract]. Proceedings of the Royal Society of London 153:421–422.

Günther, A.C.L.G. 1875a. Descriptions of the living and extinct races of gigantic land–tortoises. Parts I and II. Introduction, and the tortoises of the Galapagos Islands. Philosophical Transactions of the Royal Society of London 165:251–284.

Günther, A.C.L.G. 1875b. The gigantic land tortoises of the Mascarene and Galapagos Islands. Nature, London (1875):238–239, 259–261, 296–297.

Günther, A.C.L.G. 1877. The Gigantic Land-Tortoises (Living and Extinct) in the Collection of the British Museum. London: Taylor and Francis, 96 pp.

Günther, A.C.L.G. 1882. Description of a new species of tortoise (*Geoemyda impressa*) from Siam. Proceedings of the Zoological Society of London 1882:343–346.

Günther, A.C.L.G. 1884. Contribution to our knowledge of *Hydromedusa*, a genus of South-American freshwater turtles. Annals and Magazine of Natural History (5)14:421–425.

Günther, A.C.L.G. 1885. Reptilia and Batrachia. In: Godman, F.D. and Salvin, O. (Eds.). Biologia Centrali-Americana. London: R.H. Porter, 326 pp. [parts 37–38; pp. 1–24].

Guo, C.-W., Nie, L.-W., and Wang, M. 1997. The karyotypes and NORs of two species of *Chinemys*. In: Zhao, E. (Ed.). Chinese Chelonian Research. Chinese Society for the Study of Amphibians and Reptiles, Herpetological Series No. 9, Sichuan Journal of Zoology 15 (Supplement):97–104.

Gutsche, A. 2016. Arend Friedrich August Wiegmann (1802–1841) und sein Beitrag zur Herpetologie der Neotropis am Zoologischen Museum Berlin. Mertensiella 23:156–169.

Gutsche, A. and McCranie, J.R. 2016. Johann Georg Wagler and the "Natürliches System der Amphibien." Bibliotheca Herpetologica 12:41–49.

Guyer, C., Bailey, M.A., and Mount, R.H. 2015. Turtles of Alabama. Tuscaloosa, AL: University of Alabama Press, 267 pp.

Ha, D.D. 1995. Are Hoan Kiem tortoises an unknown species? Vietnam News (Hanoi) [Newspaper] 31 December 1995, p. 4.

Ha, D.D. 2000. Rua Ho Guom, loai rua moi cho khoa hoc. [Turtles in Hoan Kiem Lake, new species for science]. Khao co Hoc [Archaeology Magazine], Vietnam 4:104–111. [in Vietnamese]

Halámková, L., Schulte, J.A., II, and Langen, T.A. 2013. Patterns of sexual size dimorphism in Chelonia. Biological Journal of the Linnean Society 108(2):396–419.

Halk, J.H. 1986. Life history notes. *Trionyx spiniferus* (spiny softshell turtle). Size. Herpetological Review 17(3):65.

Hallermann, J. 1998. Annotated catalogue of the type specimens of the herpetological collection in the Zoological Museum of the University of Hamburg. Mitteilungen aus dem Hamburgischen Zoologischen Museum und Institut 95:197–223.

Hallowell, E. 1839. Description of a species of land tortoise, from Africa. Journal of the Academy of Natural Sciences, Philadelphia 8:161–169.

Hallowell, E. 1844. Description of new species of reptiles from Africa. Proceedings of the Academy of Natural Sciences, Philadelphia 2(5):118–120.

Hallowell, E. 1854. Descriptions of some new reptiles from California. Proceedings of the Academy of Natural Sciences, Philadelphia 7(3):91–97.

Hamann, M., Schäuble, C.S., Emerick, S.P., Limpus, D.J., and Limpus, C.J. 2008. Freshwater turtle populations in the Burnett River. Memoirs of the Queensland Museum 52:221–232.

Hansen, D.M., Donlan, C.J., Griffiths, C.J., and Campbell, K.J. 2010. Ecological history and latent conservation potential: large and giant tortoises as a model for taxon substitutions. Ecography 33:272–284.

Hansen, D.M., Austin, J.J., Baxter, R.H., de Boer, E.J., Falcon, W., Norder, S.J., Rijsdijk, K.F., Thebaud, C., Bunbury, N.J., and Warren, B.H. 2017. Origins of endemic island tortoises in the western Indian Ocean: a critique of the human-translocation hypothesis. Journal of Biogeography 44:1430–1435.

Hardwicke, T. 1835. Reptiles of India, China, etc. Vol. I. Bequeathed by Maj. Gen. Hardwicke. Unpublished volume of drawings in the British Museum (Natural History).

Harfush-Meléndez, M. and Buskirk, J.R. 2008. New distributional data on the Tehuantepec Slider, *Trachemys grayi*, in Oaxaca, Mexico. Chelonian Conservation and Biology 7(2):274–276.

Harlan, R. 1826 ["1827"]. Description of a land tortoise, from the Gallapagos Islands, commonly known as the "elephant tortoise." Journal of the Academy of Natural Sciences, Philadelphia 5:284–292.

Harlan, R. 1835. Genera of North American Reptilia, and a synopsis of the species. In: Harlan, R. Medical and Physical Researches: or Original Memoirs in Medicine, Surgery, Physiology, Geology, Zoology, and Comparative Anatomy. Philadelphia: L.R. Bailey, pp. 84–163.

Harlan, R. 1837. Description of a new species of fresh water tortoise, inhabiting the Columbia River. American Journal of Science 31:382–383.

Harper, F. 1940. Some works of Bartram, Daudin, Latreille, and Sonnini, and their bearing upon North American herpetological nomenclature. American Midland Naturalist 23:692–723.

Harris, D.J., Znari, M., Mace, J.C., and Carretero, M.A. 2003. Genetic variation in *Testudo graeca* from Morocco estimated using 12S rRNA DNA sequencing. Revista Española de Herpetología 16:5–9.

Hart, K.M. and McIvor, C.C. 2008. Demography and ecology of Mangrove Diamondback Terrapins in a wilderness area of Everglades National Park, Florida, USA. Copeia 2008(1):200–208.

Hart, K.M., Hunter, M.E., and King, T.L. 2014. Regional differentiation among populations of the Diamondback Terrapin (*Malaclemys terrapin*). Conservation Genetics 15:593–603.

Hartman, W.L. 1958. Intergradation between two subspecies of painted turtle, genus *Chrysemys*. Copeia 1958(4):261–265.

Hartweg, N. 1934. Description of a new kinosternid from Yucatan. Occasional Papers of the Museum of Zoology, University of Michigan 277:1–2.

Hartweg, N. 1938. *Kinosternon flavescens stejnegeri*, a new turtle from northern Mexico. Occasional Papers of the Museum of Zoology, University of Michigan 371:1–5.

Hartweg, N. 1939. A new American *Pseudemys*. Occasional Papers of the Museum of Zoology, University of Michigan 397:1–4.

Hastings, A.K., Krigbaum, J., Steadman, D.W., and Albury, N.A. 2014. Domination by reptiles in a terrestrial food web of the Bahamas prior to human occupation. Journal of Herpetology 48(3):380–388.

Hawkins, S., Worthy, T.H., Bedford, S., Spriggs, M., Clark, G., Irwin, G., Best, S., and Kirch, P. 2016. Ancient tortoise hunting in the southwest Pacific. Scientific Reports 6(38317), doi:10.1038/srep38317, 6 pp.

Hawlitschek, O., Ramírez Garrido, S., and Glaw, F. 2017. How marine currents influenced the widespread natural overseas dispersal of reptiles in the Western Indian Ocean region. Journal of Biogeography 44:1435–1440.

Haworth, A.H. 1825. A binary arrangement of the class Amphibia. Philosophical Magazine and Journal (1)65:372–373.

Hawthorne, W.H. and Johnston, G.R. 2024. *Pseudemys peninsularis* (Peninsula Cooter). Maximum male size. Herpetological Review 55(1):88.

Hay, O.P. 1892. The Batrachians and Reptiles of the State of Indiana. Annual Report of the Indiana Department of Geology and Natural Resources 17:412–602.

Hay, O.P. 1902. Descriptions of two new species of extinct tortoise, one new. Proceedings of the Academy of Natural Sciences, Philadelphia 54:383–388.

Hay, O.P. 1903. Two new species of fossil turtles from Oregon. University of California Publications, Bulletin of the Department of Geological Sciences 3:237–241.

Hay, O.P. 1904. On the existing genera of the Trionychidae. Proceedings of the American Philosophical Society 42:268–274.

Hay, O.P. 1906. Descriptions of two new genera (*Echmatemys* and *Xenochelys*) and two new species (*Xenochelys formosa* and *Terrapene putnami*) of fossil turtles. Bulletin of the American Museum of Natural History 22(3):27–31.

Hay, O.P. 1907. Description of seven new species of turtles from Tertiary of the United States. Bulletin of the American Museum of Natural History 23:847–863.

Hay, O.P. 1908a. On three existing species of sea-turtles, one of them (*Caretta remivaga*) new. Proceedings of the United States National Museum 34:183–198.

Hay, O.P. 1908b. The Fossil Turtles of North America. Carnegie Institution of Washington, Publication 75:1–568.

Hay, O.P. 1908c. Descriptions of five species of North American fossil turtles, four of which are new. Proceedings of the United States National Museum 35:161–169.

Hay, O.P. 1916a. Descriptions of some Floridian fossil vertebrates, belonging mostly to the Pleistocene. Annual Report of the Florida State Geological Survey 8:39–76.

Hay, O.P. 1916b. Descriptions of some fossil vertebrates found in Texas. Bulletin of the University of Texas 71:3–24.

Hay, O.P. 1921. Descriptions of some Pleistocene Vertebrata found in the United States. Proceedings of the U.S. National Museum 58:83–146.

Hay, O.P. 1924. The Pleistocene of the middle region of North America and its vertebrated animals. Carnegie Institution of Washington, Publication 322:1–374.

Hay, W.P. 1905 ["1904"]. A revision of *Malaclemmys*, a genus of turtles. Bulletin of the U.S. Bureau of Fisheries 24:1–19. [Published Feb 1905].

Haynes, D. and McKown, R.R. 1974. A new species of map turtle (Genus *Graptemys*) from the Guadalupe River system in Texas. Tulane Studies in Zoology and Botany 18(4):143–152.

He, J., Zhou, T., Rao, D.-Q., and Zhang, Y.-P. 2007. [Studies on the molecular identification and phylogeny of *Cuora yunnanensis*.] Chinese Science Bulletin 52(17):2085–2088. [In Chinese]

Hemprich, W. 1820. Grundriss der Naturgeschichte für höhere Lehranstalten. Berlin: August Rucker, 432 pp.

Henderson, J.R. 1912. Preliminary note on a new tortoise from South India. Records of the Indian Museum, Calcutta 7(3):217–218.

Hendrickson, J.R. 1965. Reptiles of the Galapagos. Pacific Discovery 18(5):28–36.

Hennessy, E. 2015. The molecular turn in conservation: genetics, pristine nature, and the rediscovery of an extinct species of Galápagos giant tortoise. Annals of the Association of American Geographers 105:87–104.

Herman, J.S., McGowan, R.Y., and Swinney, G.N. 1990. Catalogue of the type specimens of recent vertebrates in the National Museums of Scotland. National Museums of Scotland Information Series 4:1–34.

Hermann, J. 1793. [*Testudo graja*]. In: Schoepff, J.D. Historia Testudinum Iconibus Illustrata. Erlangae: Ioannis Iacobi Palm, 136 pp. [p. 52].

Hermann, J. 1804. Observationes Zoologicae. Opus posthumum edidit Fridericus Ludovicus Hammer. Argentorati [Strasbourg]: A. Koenig, 332 pp.

Herrera, A.L. 1901. Nouvelle nomenclature des êtres organisés et des minéraux. Mexico: Gouvernement Federal, 88 pp.

Hervet, S. 2004. A new genus of 'Ptychogasteridae' (Chelonii, Testudinoidea) from the Geiseltal (Lucetian of Germany). Comptes Rendus Palevol 3:125–132.

Herz, M. and Rudolphi, M. 2006. The chelonians of the Kükürt Gölü and the Dalyan Delta, Turkey – a snapshot. Radiata 15(3):19–29.

Heude, P.M. 1880. Mémoire sur les *Trionyx*. Mémoires concernant l'Histoire Naturelle de l'Empire Chinois 1(1):1–38.

Hewitt, J. 1927. Further descriptions of reptiles and batrachians from South Africa. Records of the Albany Museum 3:371–415.

Hewitt, J. 1931. Descriptions of some African tortoises. Annals of the Natal Museum 6:461–506.

Hewitt, J. 1933a. Descriptions of some new reptiles and a frog from Rhodesia. Occasional Papers of the Rhodesian Museums 1(2):45–50.

Hewitt, J. 1933b. On the Cape species and subspecies of the genus *Chersinella* Gray. Part I. Annals of the Natal Museum 7(2):255–293.

Hewitt, J. 1934. On the Cape species and subspecies of the genus *Chersinella* Gray. Part II. Annals of the Natal Museum 7(3):303–349.

Hewitt, J. 1935. Some new forms of batrachians and reptiles from South Africa. Records of the Albany Museum 4:283–357.

Hibbard, C.W. 1944. A new land-tortoise, *Testudo riggsi*, from the middle Pliocene of Seward County, Kansas. University of Kansas Science Bulletin 30:71–76.

Highfield, A.C. 1990. Tortoises of North Africa; taxonomy, nomenclature, phylogeny and evolution with notes on field studies in Tunisia. Journal of Chelonian Herpetology 1(2):1–56.

Highfield, A.C. and Martin, J. 1989a. A revision of the Testudines of North Africa, Asia and Europe. Genus: *Testudo*. Journal of Chelonian Herpetology 1(1):1–12.

Highfield, A.C. and Martin, J. 1989b. *Testudo whitei* Bennett 1836 - new light on an old carapace - Gilbert White's Selborne tortoise re-discovered. Journal of Chelonian Herpetology 1(1):13–22.

Highfield, A.C. and Martin, J. 1989c. Description of a miniature tortoise *Testudo flavominimaralis* n. species from North Africa. In: Highfield, A.C. Introduction to a Conservation Project for the North African Tortoise. London: Tortoise Trust, no pagination [pp. 9–12].

Hirayama, R., Kaneko, N., and Okazaki, H. 2007. *Ocadia nipponica*, a new species of aquatic turtle (Testudines: Testudinoidea: Geoemydidae) from the Middle Pleistocene of Chiba Prefecture, central Japan. Paleontological Research 11:1–19.

Hirth, H.F. 1980. *Chelonia mydas*. Catalogue of American Amphibians and Reptiles 249:1–4.

Hirth, H.F. 1982. Weight and length relationships of some adult marine turtles. Bulletin of Marine Science 32(1):336–341.

Hirth, H.F. 1997. Synopsis of biological data on the green turtle *Chelonia mydas* (Linnaeus 1758). U.S. Department of the Interior Biological Report 97(1), pp. 1–120.

Hitschfeld, E., Auer, M., and Fritz, U. 2008. Phalangeal formulae and ontogenetic variation of carpal morphology in *Testudo horsfieldii* and *T. hermanni*. Amphibia-Reptilia 29:93–99.

Hmar, G.Z., Lalbiakzuala, Lalmuansanga, Zote, D., Vanlalhruaia, Ramengmawii, H.B., Das, K.C., and Lalremsanga, H.T. 2020. A first distribution record of the Indian Peacock Softshell Turtle *Nilssonia hurum* (Gray, 1830) (Reptilia: Testudines: Trionychidae) from Mizoram, India. Journal of Threatened Taxa 12:17036–17040.

Hmar, G.Z., Decemson, H., Malsawmdawngliana, F., Mathipi, V., and Lalremsanga, H.T. 2024. On the occurrence of the Myanmar Brown Leaf Turtle, *Cyclemys fusca* (Testudines: Geoemydidae), from Manipur, India. Reptiles and Amphibians 31:e20816, 4 pp.

Hodges, K., Donnellan, S., and Georges, A. 2014. Phylogeography of the Australian freshwater turtle *Chelodina expansa* reveals complex relationships among inland and coastal bioregions. Biological Journal of the Linnean Society 111:789–805.

Hodges, K., Donnellan, S., and Georges, A. 2015. Significant genetic structure despite high vagility revealed through mitochondrial phylogeography of an Australian freshwater turtle (*Chelodina longicollis*). Marine and Freshwater Research; doi:10.1071/MF14102, 12 pp.

Hoffmann, C.K. 1890. Schildkröten. In: Bronn, H.G. (Ed.). Klassen und Ordnungen des Thier–Reichs. Vol. 6, Part 3, Reptilien, Vol. 1. Leipzig: Winter'sche Verlagshandlung, 442 pp.

Hoffmann, M., Hilton-Taylor, C., Angulo, A., Böhm, M., Brooks, T.M., Butchart, S.H.M., Carpenter, K.E., Chanson, J., Collen, B., Cox, N.A., Darwall, W.R.T., Dulvy, N.K., Harrison, L.R., Katariya, V., Pollock, C.M., Quader, S., Richman, N.I., Rodrigues, A.S.L., Tognelli, M.F., Vié, J.-C., Aguiar, J.M., Allen, D.J., Allen, G.R., Amori, G., Ananjeva, N.B., Andreone, F., Andrew, P., Aquino Ortiz, A.L., Baillie, J.E.M., Baldi, R., Bell, B.D., Biju, S.D., Bird, J.P., Black-Decima, P., Blanc, J.J., Bolaños, F., Bolivar-G., W., Burfield, I.J., Burton, J.A., Capper, D.R., Castro, F., Catullo, G., Cavanagh, R.D., Channing, A., Chao, N.L., Chenery, A.M., Chiozza, F., Clausnitzer,

V., Collar, N.J., Collett, L.C., Collette, B.B., Cortez Fernandez, C.F., Craig, M.T., Crosby, M.J., Cumberlidge, N., Cuttelod, A., Derocher, A.E., Diesmos, A.C., Donaldson, J.S., Duckworth, J.W., Dutson, G., Dutta, S.K., Emslie, R.H., Farjon, A., Fowler, S., Freyhof, J., Garshelis, D.L., Gerlach, J., Gower, D.J., Grant, T.D., Hammerson, G.A., Harris, R.B., Heaney, L.R., Hedges, S.B., Hero, J.-M., Hughes, B., Hussain, S.A., Icochea M., J., Inger, R.F., Ishii, N., Iskandar, D.T., Jenkins, R.K.B., Kaneko, Y., Kottelat, M., Kovacs, K.M., Kuzmin, S.L., Marca, E.L., Lamoreux, J.F., Lau, M.W.N., Lavilla, E.O., Leus, K., Lewison, R.L., Lichtenstein, G., Livingstone, S.R., Lukoschek, V., Mallon, D.P., McGowan, P.J.K., McIvor, A., Moehlman, P.D., Molur, S., Alonso, A.M., Musick, J.A., Nowell, K., Nussbaum, R.A., Olech, W., Orlov, N.L., Papenfuss, T.J., Parra-Olea, G., Perrin, W.F., Polidoro, B.A., Pourkazemi, M., Racey, P.A., Ragle, J.S., Ram, M., Rathbun, G., Reynolds, R.P., Rhodin, A.G.J., Richards, S.J., Rodríguez, L.O., Ron, S.R., Rondinini, C., Rylands, A.B., Sadovy de Mitcheson, Y., Sanciangco, J.C., Sanders, K.L., Santos-Barrera, G., Schipper, J., Self-Sullivan, C., Shi, Y., Shoemaker, A., Short, F.T., Sillero-Zubiri, C., Silvano, D.L., Smith, K.G., Smith, A.T., Snoeks, J., Stattersfield, A.J., Symes, A.J., Taber, A.B., Talukdar, B.K., Temple, H.J., Timmins, R., Tobias, J.A., Tsytsulina, K., Tweddle, D., Ubeda, C., Valenti, S.V., van Dijk, P.P., Veiga, L.M., Veloso, A., Wege, D.C., Wilkinson, M., Williamson, E.A., Xie, F., Young, B.E., Akçakaya, H.R., Bennun, L., Blackburn, T.M., Boitani, L., Dublin, H.T., Fonseca, G.A.B. da, Gascon, C., Lacher, T.E., Jr., Mace, G.M., Mainka, S.A., McNeely, J.A., Mittermeier, R.A., Reid, G.M., Rodriguez, J.P., Rosenberg, A.A., Samways, M.J., Smart, J., Stein, B.A., and Stuart, S.N. 2010. The impact of conservation on the status of the world's vertebrates. Science 330:1503–1509.

Hofmeyr, M.D. and Branch, W.R. 2018. The padloper's tortuous path (Chelonia: Testudinidae): two genera, not one. African Journal of Herpetology 2018: https://doi.org/10.1080/21564574.2017.1398187.

Hofmeyr, M.D., Vamberger, M., Branch, W., Schleicher, A., and Daniels, S.R. 2017. Tortoise (Reptilia, Testudinidae) radiations in Southern Africa from the Eocene to the present. Zoologica Scripta 46:389–400.

Hofmeyr, M.D., Ihlow, F., Fouche, P., and Daniels, S.R. 2020. Niche divergence corresponds to genetic differentiation within the Parrot-beaked Tortoise *Homopus areolatus* (Reptilia: Tesudinidae), endemic to South Africa. Zoological Journal of the Linnean Society 190:1256–1273.

Holbrook, J.E. 1836. North American Herpetology; or, a Description of the Reptiles Inhabiting the United States. Ed. 1, Vol. 1. Philadelphia: J. Dobson, 120 pp.

Holbrook, J.E. 1838a. North American Herpetology; or, a Description of the Reptiles Inhabiting the United States. Ed. 1, Vol. 2. Philadelphia: J. Dobson, 125 pp.

Holbrook, J.E. 1838b. North American Herpetology; or, a Description of the Reptiles Inhabiting the United States. Ed. 1, Vol. 3. Philadelphia: J. Dobson, 122 pp.

Holbrook, J.E. 1840. North American Herpetology; or, a Description of the Reptiles Inhabiting the United States. Ed. 1, Vol. 4. Philadelphia: J. Dobson, 126 pp.

Holley, J.A., Sterli, J., and Basso, N.G. 2020. Dating the origin and diversification of Pan-Chelidae (Testudines, Pleurodira) under multiple molecular clock approaches. Contributions to Zoology 89:146–174.

Holm, Å. 1957. Specimina Linnaeana. I Uppsala bevarade zoologiska samlingar från Linnés tid. Uppsala Universitets Årsskrift 1957(6):1–68. [in Swedish]

Holman, J.A. 1972. Amphibians and reptiles. In: Skinner, M.F. and Hibbard, C.W. (Eds.). Early Pleistocene pre-glacial and glacial rocks and faunas of north-central Nebraska. Bulletin of the American Museum of Natural History 148(1):55–148.

Holman, J.A. 1978. The Late Pleistocene herpetofauna of Devil's Den sinkhole, Levy County, Florida. Herpetologica 34(2):228–237.

Holman, J.A. and Clausen, C.J. 1984. Fossil vertebrates associated with Paleo-Indian artifact at Little Salt Spring, Florida. Journal of Vertebrate Paleontology 4:146–154.

Honda, M., Yasukawa, Y., Hirayama, R., and Ota, H. 2002a. Phylogenetic relationships of the Asian box turtles of the genus *Cuora* sensu lato (Reptilia: Bataguridae) inferred from mitochondrial DNA sequences. Zoological Science 19:1305–1312.

Honda, M., Yasukawa, Y., and Ota, H. 2002b. Phylogeny of Eurasian freshwater turtles of the genus *Mauremys* Gray 1869 (Testudines), with special reference to a close affinity of *Mauremys japonica* with *Chinemys reevesii*. Journal of Zoological Systematics and Evolutionary Research 40:195–200.

Hong, X., Zhu, X., Chen, C., Cai, X., Li, Y., and Li, X. 2022. Reproductive characteristics of the Burmese Narrow-headed Softshell Turtle, *Chitra vandijki*, in captivity. Amphibian & Reptile Conservation 16(2):62–68(e316).

Hoogmoed, M.S. and Crumly, C.R. 1984. Land tortoise types in the Rijksmuseum van Natuurlijke Historie with comments on nomenclature and systematics (Reptilia: Testudines: Testudinidae). Zoologische Mededelingen, Leiden 58:241–259.

Hoogmoed, M.S. and Gruber, U. 1983. Spix and Wagler type specimens of reptiles and amphibians in the Natural History Musea in Munich (Germany) and Leiden (The Netherlands). Spixiana Supplement 9:319–415.

Hoogmoed, M.S., Gassó Miracle, M.E., and van den Hoek Ostende, L.W. 2010. Type specimens of recent and fossil Testudines and Crocodylia in the collections of the Netherlands Centre for Biodiversity Naturalis, Leiden, the Netherlands. Zoologische Mededelingen, Leiden 84(8):159–199.

Hossain, M.L. and Sarker, M.S.U. 1995. Observations on the narrow-headed softshell turtle (*Chitra indica*) in Bangladesh. Journal of the Bombay Natural History Society 92:423–426.

Hou, X. and Shi, H. 2025. Amur Softshell Turtle (*Pelodiscus maackii*) population size, structure, and spatial distribution. Animals 15:255, 15 pp.

Hou, X., Lin, L., Li, M., Jiang, H., and Shi, H. 2023. A new species of the genus *Pelodiscus* (Testudines, Trionychia) from Jiangxi China. International Journal of Zoology and Animal Biology 6(1): doi:10.23880/izab-16000444, 7 pp. (online preprint, withdrawn prior to actual publication).

Houttuyn, M. 1764. Natuurlyke Historie of uitvoerige beschryving der Dieren, Planten en Mineraalen, volgens het samenstel van den Heer Linnaeus. Met naauwkeurige Afbeeldingen. Eerste deels, zesde stuk. Dieren van beiderley leven. Amsterdam: F. Houttuyn, 558 pp.

Høybye-Mortensen, K. 2004. The tortoise *Manouria emys emys*: behaviour and habitat in the wild. Masters Thesis, University of Southern Denmark.

Hsü, H.F. 1930. Preliminary note on a new variety of *Cyclemys flavomarginata* from China. Contributions from the Biological Laboratory of the Science Society of China, Zoological Series 6(1):1–7.

Htun, S. and Platt, S.G. 2016. A field observation and significant range extension of *Manouria impressa* in Myanmar. Asian Herpetological Research 7(4):295–297.

Hu, Q., Huang, C., Xu, S., Zhang, Q., Ma, N., and Zhong, H. 2013. Primary phylogenies of *Mauremys guangxiensis* and *Mauremys iversoni* inferred from DNA sequences of mitochondrial ND4 gene and nuclear c-mos gene. Sichuan Journal of Zoology 32(2):180–186.

Hubrecht, A.A.W. 1881. On certain tortoises in the collections of the Leyden Museum. Notes from the Leyden Museum 3:41–50.

Huebinger, R.M., Bickham, J.W., Rhodin, A.G.J., and Mittermeier, R.A. 2013. Mitochondrial DNA corroborates taxonomy of the South American chelid turtles of the genera *Platemys* and *Acanthochelys*. Chelonian Conservation and Biology 12(1):168–171.

Humboldt, A. de. 1819a. Voyage aux Régions Équinoxiales du Nouveau Continent, fait en 1799, 1800, 1801, 1802, 1803 et 1804, par Al. de Humboldt et A. Bonpland. Tome Second. Paris: N. Maze, 381 pp.

Humboldt, A. de. 1819b. Personal Narrative of Travels to the Equinoctial Regions of the New Continent, during the years 1799–1804, by Alexandre de Humboldt and Aimé Bonpland. Vol. IV. [Translated by H.M. Williams]. London: Longman, Hurst, Rees, Orme, and Brown.

Humboldt, A. von. 1820. Reise in die Aequinoctial-Gegenden des neuen Continents in den Jahren 1799, 1800, 1801, 1802, 1803, und 1804. Verfasst von Alexander von Humboldt und A. Bonpland. Dritter Theil. Stüttgart und Tübingen: J.G. Cotta'schen Buchhandlung.

Hümmel, K. 1929. Die fossilen weichschildkröten (Trionychia). Eine morphologisch-systematische und stammesgeschichtliche Studie. Geologische und Palæontologische Abhandlungen 16:359–487.

Hurd, L.E., Smedes, G.W., and Dean, T.A. 1979. An ecological study of a natural population of diamondback terrapins (*Malaclemys t. terrapin*) in a Delaware salt marsh. Estuaries 2:28–33.

Hurtado-Gómez, J.P., Vargas-Ramírez, M., Iverson, J.B., Joyce, W., McCranie, J., Paetzold, C., and Fritz, U. 2024. Diversity and biogeography of South American mud turtles elucidated by multilocus DNA sequencing (Testudines: Kinosternidae). Molecular Phylogenetics and Evolution 197:108083:1–19.

Hurter, J. 1911. Herpetology of Missouri. Transactions of the Academy of Sciences, St. Louis 20:59–274.

Husson, A.M. and Holthuis, L.B. 1955. The dates of publication of "Verhandelingen over de Natuurlijke Geschiedenis der Nederlandsche Overzeesche Bezittingen" edited by C.J. Temminck. Zoologische Mededelingen, Leiden

34(2):17–24.

Hutchison, J.H. 1991. Early Kinosterninae (Reptilia: Testudines) and their phylogenetic significance. Journal of Vertebrate Paleontology 11:145–167.

Hutchison, J.H. 2006. *Bridgeremys* (Geoemydidae, Testudines) a new genus from the middle Eocene of North America. Fossil Turtle Research 1:63–83.

Hutchison, J.H. and Weems, R.E. 1998. Paleocene turtle remains from South Carolina. Transactions of the American Philosophical Society 88:165–195.

Huxley, T.H. 1887. Preliminary note on the fossil remains of a chelonian reptile, *Ceratochelys sthenurus*, from Lord Howe's Island, Australia. Proceedings of the Royal Society 42:232–238.

Ibarra Portillo, R., Henriquez, V., and Greenbaum, E. 2009. Geographic distribution. *Trachemys emolli* (Moll's Slider). Herpetological Review 40(1):111.

ICZN [International Commission on Zoological Nomenclature]. 1922. Opinion 72. Herrera's zoological formulae. Smithsonian Miscellaneous Collections 73(1):19–22.

ICZN [International Commission on Zoological Nomenclature]. 1926. Opinion 92. Sixteen generic names of Pisces, Amphibia, and Reptilia placed in the Official List of Generic Names. Smithsonian Miscellaneous Collections, 73(4): 3–4.

ICZN [International Commission on Zoological Nomenclature]. 1944. Opinion 184. On the status of names first published in volumes 1 to 11 of Martini (F.H.W.) and Chemnitz (J.H.), Neues systematisches Conchylien-Cabinet, Nürnberg, 1769-1795. Opinions and Declarations Rendered by the International Commission on Zoological Nomenclature, Vol. 3, Part 3, pp. 25–36.

ICZN [International Commission on Zoological Nomenclature]. 1954. Direction 1. Addition to the Official Lists and Official Indexes of certain scientific names and of the titles of certain books dealt with in Opinions 182 to 194. Opinions and Declarations Rendered by the International Commission on Zoological Nomenclature, Vol. 3, Part 30, pp. 401–416.

ICZN [International Commission on Zoological Nomenclature]. 1956a. Direction 32. Addition to the Official List of Works Approved as Available for Zoological Nomenclature of the titles of certain works dealt with in Opinions rendered in the period up to the end of 1936 and to the Official Index of Rejected and Invalid Works in Zoological Nomenclature of certain works similarly dealt with in the same period and matters incidental thereto. Opinions and Declarations Rendered by the International Commission on Zoological Nomenclature, Vol. 1, Sec. C, pp. 307–328.

ICZN [International Commission on Zoological Nomenclature]. 1956b. Direction 56. Completion and in certain cases correction of entries relating to the names of genera belonging to the Classes Pisces, Amphibia and Reptilia made in the Official List of Generic Names in Zoology in the period up to the end of 1936. Opinions and Declarations Rendered by the International Commission on Zoological Nomenclature, Vol. 1, Sec. D, pp. 337–364.

ICZN [International Commission on Zoological Nomenclature]. 1963. Opinion 660. Suppression under the plenary powers of seven specific names of turtles (Reptilia, Testudines). Bulletin of Zoological Nomenclature 20:187–190.

ICZN [International Commission on Zoological Nomenclature]. 1982. Opinion 1236. *Trionyx steindachneri* Siebenrock, 1906 (Reptilia, Testudines): conserved. Bulletin of Zoological Nomenclature 39:258–259.

ICZN [International Commission on Zoological Nomenclature]. 1984. Opinion 1280. Rafinesque, C.S., 1822 "On the turtles of the United States": suppressed. Bulletin of Zoological Nomenclature 41:221–222.

ICZN [International Commission on Zoological Nomenclature]. 1985a. Opinion 1309. *Geoemyda* Gray, 1834, and *Rhinoclemmys* Fitzinger, 1835 (Reptilia, Testudines): conserved. Bulletin of Zoological Nomenclature 42:152–153.

ICZN [International Commission on Zoological Nomenclature]. 1985b. Opinion 1313. *Testudo scripta* Schoepff, 1792 and *Emys cataspila* Günther, 1885 (Reptilia, Testudines): conserved. Bulletin of Zoological Nomenclature 42:160–161.

ICZN [International Commission on Zoological Nomenclature]. 1985c. Opinion 1343. *Kinosternon alamosae* Berry and Legler, 1980 and *Kinosternon oaxacae* Berry and Iverson, 1980 (Reptilia, Testudines): conserved. Bulletin of Zoological Nomenclature 42:266–268.

ICZN [International Commission on Zoological Nomenclature]. 1987a. Opinion 1463. De Lacepède, 1788–1789, Histoire Naturelle des Serpens and later editions: rejected as a non-binomial work. Bulletin of Zoological Nomenclature 44(4):265–267.

ICZN [International Commission on Zoological Nomenclature]. 1987b. Official Lists and Indexes of Names and Works in Zoology. London: The International Commission of Zoological Nomenclature, 366 pp.

ICZN [International Commission on Zoological Nomenclature]. 1989. Opinion 1534. *Sternotherus* Gray, 1825 and *Pelusios* Wagler, 1830 (Reptilia, Testudines): conserved. Bulletin of Zoological Nomenclature 46:81–82.

ICZN [International Commission on Zoological Nomenclature]. 1991. Opinion 1659. *Trionyx sinensis* Wiegmann, 1834 (Reptilia, Testudines): specific name conserved. Bulletin of Zoological Nomenclature 48(3):276.

ICZN [International Commission on Zoological Nomenclature]. 1995. Opinion 1800. *Emys* Duméril, 1806 (Reptilia, Testudines): conserved. Bulletin of Zoological Nomenclature 52(1):111–112.

ICZN [International Commission on Zoological Nomenclature]. 1999. International Code of Zoological Nomenclature. Fourth Edition. London: International Trust for Zoological Nomenclature, 306 pp.

ICZN [International Commission on Zoological Nomenclature]. 2005a. Opinion 2104 (Case 3226). Lacepède, B.G.É. de la V., 1788, Histoire Naturelle de Quadrupèdes Ovipares: rejected as a non-binomial work. Bulletin of Zoological Nomenclature 62(1):55.

ICZN [International Commission on Zoological Nomenclature]. 2005b. Opinion 2119 (Case 3277). *Chitra chitra* Nutaphand, 1986 (Reptilia, Testudines): specific name given precedence over *Chitra selenkae* Jaekel, 1911. Bulletin of Zoological Nomenclature 62(2):118–119.

ICZN [International Commission on Zoological Nomenclature]. 2012. Amendment of Articles 8, 9, 10, 21, and 78 of the International Code of Zoological Nomenclature to expand and refine methods of publication. ZooKeys 219:1–10.

ICZN [International Commission on Zoological Nomenclature]. 2013a. Opinion 2315 (Case 3351). *Chelodina rugosa* Ogilby, 1890 (currently *Macrochelodina rugosa*; Reptilia, Testudines): precedence not granted over *Chelodina oblonga* Gray, 1841. Bulletin of Zoological Nomenclature 70(1):57–60.

ICZN [International Commission on Zoological Nomenclature]. 2013b. Opinion 2316 (Case 3463). *Testudo gigantea* Schweigger, 1812 (currently *Geochelone (Aldabrachelys) gigantea*; Reptilia, Testudines): usage of the specific name conserved by maintenance of a designated neotype, and suppression of *Testudo dussumieri* Gray, 1831 (currently *Dipsochelys dussumieri*). Bulletin of Zoological Nomenclature 70(1):61–65.

ICZN [International Commission on Zoological Nomenclature]. 2019. Opinion 2431 (Case 3587) – *Podocnemis unifilis* Troschel, 1848 (Reptilia, Testudines): precedence given over *Emys cayennensis* Schweigger, 1812. Bulletin of Zoological Nomenclature 76:77–78.

ICZN [International Commission on Zoological Nomenclature]. 2021. Opinion 2468 (Case 3601) – *Spracklandus* Hoser, 2009 (Reptilia, Serpentes, Elapidae) and Australasian Journal of Herpetology issues 1–24: confirmation of availability declined; Appendix A (Code of Ethics): not adopted as a formal criterion for ruling on Cases. Bulletin of Zoological Nomenclature 78:42–45.

Iftime, A. 2002. *Testudo hermanni* Gmelin, 1789 in Dobroudja (SE Romania), with comments on conservation. Herpetozoa 15(3/4):183–186.

Ihering, H. von. 1926. [*Hydraspis lutzi*]. In: Luederwaldt, H. Os chelonios brasileiros. Revista Museo Paulista 14:403–470. [p. 441].

Ihlow, F., Ahmadzadeh, F., Ghaffari, H., Taskavak, E., Hartmann, T., Etzbauer, C., and Rödder, D. 2014. Assessment of genetic structure, habitat suitability and effectiveness of reserves for future conservation planning of the Euphrates soft-shelled turtle *Rafetus euphraticus* (Daudin, 1802). Aquatic Conservation: Marine and Freshwater Ecosystems 24:831–840.

Ihlow, F., Vamberger, M., Flecks, M., Hartmann, T., Cota, M., Makchai, S., Meewattana, P., Dawson, J.E., Kheng, L., Rödder, D., and Fritz, U. 2016. Integrative taxonomy of Southeast Asian snail-eating turtles (Geoemydidae: *Malayemys*) reveals a new species and mitochondrial introgression. PLoS ONE 11:e0153108, 26 pp.

Ihlow, F., Farooq, H.M., Gvožďík, V., Hofmeyr, M.D., Conradie, W., Campbell, P.D., Harvey, J., Verburgt, L., and Fritz, U. 2019. Geographic range extension of Speke's Hinge-back Tortoise *Kinixys spekii* Gray, 1863. Amphibian and Reptile Conservation 13(2):61–67.

Ihlow, F., Van Huyssteen, R., Vamberger, M., Cory-Toussaint, D., Hofmeyr, M.D., and Fritz, U. 2020. Geographic range extension for the Lobatse Hinge-back Tortoise, *Kinixys lobatsiana* (Power, 1927), with first records from the Soutpansberg region. Amphibian and Reptile Conservation 14(1):132–139.

Ihlow, F., Spitzweg, C., Vamberger, M., Augustine, L., Hagen, C., Davis, A., Leprince, B., Wagner, P., Pham Van, T., and Fritz, U. 2023. Molecular differentiation and conservation of the Indochinese box turtles *Cuora galbinifrons*, *Cuora bourreti*, and *Cuora picturata*. Salamandra 59(4):336–346.

Ihlow, F., Spitzweg, C., Flecks, M., Poyarkov, N.A., Mohapatra, P.P., Deepak, V., and Fritz, U. 2024. Unexpected lack of genetic and morphological divergence in a widespread tortoise – phylogeography of *Indotestudo elongata*. Salamandra 60:183–194.

Iskandar, D.T. 2000. Turtles and Crocodiles of Insular Southeast Asia and New Guinea. Bandung, Indonesia: PALMedia Citra, 191 pp.

Iskandar, D.T. 2004. On the giant Javanese softshelled turtles (Trionychidae). Hamadryad 28:128–130.

Itescu, Y., Karraker, N.E., Raia, P., Pritchard, P.C.H., and Meiri, S. 2014. Is the island rule general? Turtles disagree. Global Ecology and Biogeography 23:689–700.

Iturralde-Vinent, M.A. 2006. Meso-Cenozoic Caribbean paleogeography: implications for the historical biogeography of the region. International Geology Review 48(9):791–827.

IUCN [International Union for the Conservation of Nature]. 2001. IUCN Red List Categories and Criteria: Version 3.1. Gland, Switzerland: IUCN Species Survival Commission, 30 pp.

IUCN [International Union for the Conservation of Nature]. 2013. Guidelines for Reintroductions and Other Conservation Translocations. Version 1.0. Gland, Switzerland: IUCN Species Survival Commission, 57 pp.

IUCN [International Union for the Conservation of Nature]. 2016. Guidelines for Using the IUCN Red List Categories and Criteria. Version 12. Prepared by the Standards and Petitions Subcommittee. Gland, Switzerland: IUCN Species Survival Commission, 101 pp.

IUCN [International Union for the Conservation of Nature]. 2022. Guidelines for Using the IUCN Red List Categories and Criteria. Version 15. Prepared by the Standards and Petitions Committee. https://www.iucnredlist.org/documents/RedListGuidelines.pdf.

Iverson, J.B. 1976. The genus *Kinosternon* in Belize (Testudines: Kinosternidae). Herpetologica 32:258–262.

Iverson, J. B. 1977. Geographic variation in the musk turtle, *Sternotherus minor*. Copeia 1977(3):502–517.

Iverson, J.B. 1978. Distributional problems of the genus *Kinosternon* in the American southwest. Copeia 1978(3):476–479.

Iverson, J.B. 1979a. On the validity of *Kinosternon arizonense* Gilmore. Copeia 1979(1):175–177.

Iverson, J.B. 1979b. A taxonomic reappraisal of the yellow mud turtle, *Kinosternon flavescens* (Testudines: Kinosternidae). Copeia 1979(2):212–225.

Iverson, J.B. 1980. *Kinosternon acutum*. Catalogue of American Amphibians and Reptiles 261:1–2.

Iverson, J.B. 1981. Biosystematics of the *Kinosternon hirtipes* species group (Testudines: Kinosternidae). Tulane Studies in Zoology and Botany 23:1–74.

Iverson, J.B. 1983. *Staurotypus triporcatus*. Catalogue of American Amphibians and Reptiles 328:1–2.

Iverson, J.B. 1986. Notes on the natural history of the Oaxaca mud turtle, *Kinosternon oaxacae*. Journal of Herpetology 20(1):119–123.

Iverson, J.B. 1989. The Arizona mud turtle, *Kinosternon flavescens arizonense* (Kinosternidae), in Arizona and Sonora. Southwestern Naturalist 34:356–368.

Iverson, J.B. 1991. Phylogenetic hypotheses for the evolution of modern kinosternine turtles. Herpetological Monographs 4:1–27.

Iverson, J.B. 1992. A Revised Checklist with Distribution Maps of the Turtles of the World. Richmond, IN: Privately printed, 363 pp.

Iverson, J.B. 1998. Molecules, morphology, and mud turtle phylogenetics (family Kinosternidae). Chelonian Conservation and Biology 3(1):113–117.

Iverson, J.B. 2002. Reproduction in female Razorback Musk Turtles (*Sternotherus carinatus*: Kinosternidae). Southwestern Naturalist 47(2):215–224.

Iverson, J.B. 2008. *Kinosternon scorpioides abaxillare* (Central Chiapas mud turtle): size, growth, and reproduction. Herpetological Review 39:217–218.

Iverson, J.B. 2010. Reproduction in the Red-cheeked Mud Turtle (*Kinosternon scorpioides cruentatum*) in southeastern Mexico and Belize, with comparisons across the species range. Chelonian Conservation and Biology 9(2):250–261.

Iverson, J.B. 2022a. A review of chelonian type specimens (order Testudines). Megataxa 7(1):1–85.

Iverson, J.B. 2022b. *Kinosternon steindachneri* (Florida Mud Turtle). Reproductive output. Herpetological Review 53:311–312.

Iverson, J.B. and Berry, J.F. 1980. *Claudius, C. angustatus*. Catalogue of American Amphibians and Reptiles 236:1–2.

Iverson, J.B. and Berry, J.F. 2024. Morphometric variation in the Red-cheeked Mud Turtle (*Kinosternon cruentatum*) and its taxonomic implications. Chelonian Conservation and Biology 23:13–22.

Iverson, J.B. and Lewis, E.L. 2018. How to measure a turtle. Herpetological Review 49:453–460.

Iverson, J.B. and McCord, W.P. 1989. The proper taxonomic allocations of *Emys nigricans* Gray, *Emys muticus* Cantor, and *Geoclemys kwangtungensis* Pope. Amphibia-Reptilia 10(1):23–33.

Iverson, J.B. and McCord, W.P. 1992a. A new Chinese eyed turtle of the genus *Sacalia* (Batagurinae: Testudines). Proceedings of the Biological Society of Washington 105(3):426–432.

Iverson, J.B. and McCord, W.P. 1992b. A new subspecies of *Cuora galbinifrons* (Testudines: Batagurinae) from Hainan Island, China. Proceedings of the Biological Society of Washington 105(3):433–439.

Iverson, J.B. and McCord, W.P. 1994. Variation in east Asian turtles of the genus *Mauremys* (Bataguridae; Testudines). Journal of Herpetology 28(2):178–187.

Iverson, J.B. and McCord, W.P. 1997a. Redescription of the Arakan forest turtle *Geoemyda depressa* Anderson 1875 (Testudines: Bataguridae). Chelonian Conservation and Biology 2(3):384–389.

Iverson, J.B. and McCord, W.P. 1997b. A new species of *Cyclemys* (Testudines: Bataguridae) from southeast Asia. Proceedings of the Biological Society of Washington 110(4):629–639.

Iverson, J.B. and McCord, W.P. 2006. Intraspecific variation in the Giant Asian Pond Turtle, *Heosemys grandis* (Gray, 1860). Hamadryad 30(1):124–130.

Iverson, J.B. and Vogt, R.C. 2002. *Peltocephalus, P. dumerilianus*. Catalogue of American Amphibians and Reptiles 744:1–4.

Iverson, J.B, Ernst, C.H., Gotte, S., and Lovich, J.E. 1989. The validity of *Chinemys megalocephala* (Testudines: Batagurinae). Copeia 1989:494–498.

Iverson, J.B., Higgins, H., Sirulnik, A., and Griffiths, C. 1997. Local and geographic variation in the reproductive biology of the snapping turtle (*Chelydra serpentina*). Herpetologica 53:96–117.

Iverson, J.B., Kimerling, J., Kiester, A.R., Hughes, L.E., and Nicolello, J. 2001a. Turtles of the World. [http://emys.geo.orst.edu].

Iverson, J.B., Thomson, S.A., and Georges, A. 2001b. Validity of taxonomic changes for turtles proposed by Wells and Wellington. Journal of Herpetology 35:361–368.

Iverson, J.B., Spinks, P.Q., Shaffer, H.B., McCord, W.P., and Das, I. 2001c. Phylogenetic relationships among the Asian tortoises of the genus *Indotestudo* (Reptilia: Testudines: Testudinidae). Hamadryad 26(2):272–275.

Iverson, J.B., Brown, R.M., Akre, T.S., Near, T.J., Le, M., Thomson, R.C., and Starkey, D.E. 2007. In search of the tree of life for turtles. In: Shaffer, H.B., FitzSimmons, N.N., Georges, A., and Rhodin, A.G.J. (Eds.). Defining Turtle Diversity: Proceedings of a Workshop on Genetics, Ethics, and Taxonomy of Freshwater Turtles and Tortoises. Chelonian Research Monographs 4:85–106.

Iverson, J.B., Meylan, P.A., and Seidel, M.E. 2008. Testudines—Turtles. In: Crother, B.I. (Ed.). Scientific and standard English names of amphibians and reptiles of North America north of Mexico, with comments regarding confidence in our understanding. Sixth Edition. Society for the Study of Amphibians and Reptiles, Herpetological Circulars No. 37, pp. 67–74.

Iverson, J.B., Le, M., and Ingram, C. 2013. Molecular phylogenetics of the mud and musk turtle family Kinosternidae. Molecular Phylogenetics and Evolution 69(3):929–939.

Iverson, J.B., Schneider, L., and Vogt, R.C. 2017. *Podocnemis sextuberculata*. Catalogue of American Amphibians and Reptiles 913:1–24.

Iverson, J.B., Lemos-Espinal, J., and Smith, G.R. 2018. *Kinosternon durangoense* (Durango Mud Turtle). Life history. Herpetological Review 49:109.

Ives, I.E., Platt, S.G., Tasirin, J.S., Hunowu, I., Siwu, S., and Rainwater, T.R. 2008. Field surveys, natural history observations, and comments on the exploitation and conservation of *Indotestudo forstenii*, *Leucocephalon yuwonoi*, and *Cuora amboinensis* in Sulawesi, Indonesia. Chelonian Conservation and Biology 7(2):240–248.

Jackson, C.G., Jr. 1964. The status of *Deirochelys floridana* Hay with comments on the fossil history of the genus. Tulane Studies in Geology 2(3):103–106.

Jackson, C.G., Jr. 1974. The status of *Trachemys jarmani* Hay with clarification of the fossil record of *Deirochelys*. Copeia 1974:536–537.

Jackson, C.G., Jr., Trotter, M.W., and Allen, W. 1980. A gigantic specimen of *Gopherus agassizii* (Testudinidae). Southwestern Association of Naturalists 25(3):430–432.

Jackson, D.R. 1995. Systematics of the *Pseudemys concinna–floridana* complex (Testudines: Emydidae): an alternative interpretation. Chelonian Conservation

and Biology 1(4):329–333.

Jackson, D.R. 2006. *Pseudemys concinna* – River Cooter. In: Meylan, P.A. (Ed.). Biology and Conservation of Florida Turtles. Chelonian Research Monographs 3:325–337.

Jackson, D.R. and Brechtel, B. 2006. *Terrapene, T. carolina, T. c. bauri, T. c. major* (American Box Turtles, Eastern Box Turtle, Florida Box Turtle, Gulf Coast Box Turtle). Maximum size. Herpetological Review 37(3):342–343.

Jackson, D.R. and Walker, R.N. 1997. Reproduction in the Suwannee cooter, *Pseudemys concinna suwanniensis*. Bulletin of the Florida Museum of Natural History 41(2):69–167.

Jackson, J.T., Starkey, D.E., Guthrie, R.W., and Forstner, M.R.J. 2008. A mitochondrial DNA phylogeny of extant species of the genus *Trachemys* with resulting taxonomic implications. Chelonian Conservation and Biology 7(1):131–135.

Jackson, T.G., Jr., Nelson, D.H., and Morris, A.B. 2012. Phylogenetic relationships in the North American genus *Pseudemys* (Emydidae) inferred from two mitochondrial genes. Southeastern Naturalist 11(2):297–310.

Jadhav, T.D., Sawant, N.S., and Shyama, S.K. 2018. Diversity and distribution of freshwater turtles (Reptilia: Testudines) in Goa, India. Journal of Threatened Taxa 10(9):12194–12202.

Jaeger, E.C. 1955. A Source-book of Biological Names and Terms. Third Edition. Springfield, IL: Charles C. Thomas, 323 pp.

Jaekel, O. 1911. Die fossilen Schildkrötenreste von Trinil. In: Selenka, M.L. and Blanckenhorn, M. (Eds.). Die Pithecanthropus-Schichten auf Java. Geologische und Paläontologische Ergebnisse der Trinil-Expedition (1907 und 1908). Leipzig: Wilhelm Engelmann, pp. 75–81.

Jaffe, A.L., Slater, G.J., and Alfaro, M.E. 2011. The evolution of island gigantism and body size variation in tortoises and turtles. Biology Letters 7:558–561.

Jäger, G. von. 1861. Bemerkungen ueber die Sumpfschildkroete (*Emys europaea*) in fossilem Zustande. Bulletin de la Société Impériale des Naturalistes de Moscou 34:190–200.

Jain, A., Cavada-Blanco, F., Palot, M.J., Das, S., Deepak, V., and Das, A. 2021. A review of distribution of Cantor's Giant Softshell Turtle in India along with multiple notable new records from Kerala State. Chelonian Conservation and Biology 20(1):125–132.

Jarocki, F.P.N. 1822. Zoologia czyly Zwierzetopismo ogolne podlug Naynowszego Systematu. Vol. 3. Warsaw. [in Polish].

Jaruthanin, K. 2002. Talui thin pla hayak. [Salween: the dangerous river]. Fish Zone, Bangkok 3(19):27–40. [in Thai].

Javanbakht, H., Ihlow, F., Jablonski, D., Široky, P., Fritz, U., Rödder, D., Sharifi, M., and Mikulícek, P. 2017. Genetic diversity and Quaternary range dynamics in Iranian and Transcaucasian tortoises. Biological Journal of the Linnean Society 121:627–640.

Jensen, E.L., Govindarajulu, P., and Russello, M.A. 2014a. When the shoe doesn't fit: applying conservation unit concepts to western painted turtles at their northern periphery. Conservation Genetics 15:261–274.

Jensen, E.L., Govindarajulu, P., Madsen, J., and Russello, M.A. 2014b. Extirpation by introgression? Genetic evidence reveals hybridization between introduced *Chrysemys picta* and endangered Western Painted Turtles (*C. p. bellii*) in British Columbia. Herpetological Conservation and Biology 9:342−353.

Jensen, E.L., Govindarajulu, P., and Russello, M.A. 2015a. Genetic assessment of taxonomic uncertainty in painted turtles. Journal of Herpetology 49:314–324.

Jensen, E.L., Tapia, W., Caccone, A., and Russello, M.A. 2015b. Genetics of a head-start program to guide conservation of an endangered Galápagos tortoise (*Chelonoidis ephippium*). Conservation Genetics 16:823–832.

Jensen, E.L., Quinzin, M.C., Miller, J.M., Russello, M.A., Garrick, R.C., Edwards, D.L., Glaberman, S., Chiari, Y., Poulakakis, N., Tapia, W., Gibbs, J.P., and Caccone, A. 2022a. A new lineage of Galapagos giant tortoises identified from museum samples. Heredity 128:261–270.

Jensen, E.L., Gaughran, S.J., Fusco, N.A., Poulakakis, N., Tapia, W., Sevilla, C., Malaga, J., Mariani, C., Gibbs, J.P., and Caccone, A. 2022b. The Galapagos giant tortoise *Chelonoidis phantasticus* is not extinct. Communications Biology 5(546):1–8.

Jensen, M.P., FitzSimmons, N.N., Bourjea, J., Hamabata, T., Reece, J., and Dutton, P.H. 2019. The evolutionary history and global phylogeography of the Green Turtle (*Chelonia mydas*). Journal of Biogeography 46:860–870.

Jerdon, T.C. 1870. Notes on Indian herpetology. Proceedings of the Asiatic Society of Bengal 1870(3):66–85.

Jesu, R., Piombo, R., Salvidio, S., Lamagni, L., Ortale, S., and Genta, P. 2004. Un nuovo taxon di testuggine palustre endemico della Liguria occidentale: *Emys orbicularis ingauna* n. ssp. (Reptilia, Emydidae). Annali del Museo Civico di Storia Naturale "G. Doria" 96:133–192. [in Italian].

Johnston, C.S. 1937. Osteology of *Bysmachelys canyonensis*, a new turtle from the Pliocene of Texas. Journal of Geology 45:439–447.

Johnston, G.R., Suarez, E., Mitchell, J.C., Shemitz, G.A., Butt, P.L., and Kaunert, M. 2012. Population ecology of the snapping turtle (*Chelydra serpentina osceola*) in a northern Florida river. Bulletin of the Florida Museum of Natural History 51(4):243–256.

Johnston, G.R., Thomas, T.M., Suarez, E., Lau, A., and Mitchell, J.C. 2015. Population structure and body size of the Suwannee Alligator Snapping Turtle (*Macrochelys suwanniensis*) in Northern Florida. Chelonian Conservation and Biology 14(1):73–81.

Johnston, G.R., Mitchell, J.C., Geiger, J.S., Casteel, J.L., and Jones, C.S. 2019. *Kinosternon baurii* (Striped Mud Turtle). Body size variation. Herpetological Review 50:772–773.

Johnston, G.R., Geiger, J.S., Thomas, T.M., Enge, K.M., Suarez, E., and Davis, B. 2023. Maximum body size of the Suwannee Alligator Snapping Turtle (*Macrochelys suwanniensis*). Southeastern Naturalist 22(Special Issue 12):418–428.

Johnston, G.R., Eaton, P., Casteel, J., Ehlers, A., Geiger, J.S., Godwin, C.D., and Hootman, T. 2024a ("2023"). *Pseudemys concinna suwanniensis* (Suwannee Cooter). Maximum size. Herpetological Review 54(3):453–454.

Johnston, G.R., Munscher, E.C., Geiger, J.S., Godwin, C.D., and Cole, S. 2024b ("2023"). *Chelydra serpentina* (Snapping Turtle). Maximum size. Herpetological Review 54(4):642–643.

Johnston, G.R., Snipes, S., Salvatico, N., Titterington, K., Hussain, L., and Hootman, T. 2024c. *Trachemys scripta scripta* (Yellow-bellied Slider). Maximum male size. Herpetological Review 55(1):89.

Jones, B. 2017. A few bad scientists are threatening to topple taxonomy. Smithsonian Magazine, https://www.smithsonianmag.com/science-nature/the-big-ugly-problem-heart-of-taxonomy-180964629/

Jones, P.J. 1994. Biodiversity in the Gulf of Guinea: an overview. Biodiversity and Conservation 3:772–785.

Joos, J., Kirchner, M., Vamberger, M., Kaviani, M., Rahimibashar, M.R., Fritz, U., and Müller, J. 2017. Climate and patterns of body size variation in the European pond turtle, *Emys orbicularis*. Biological Journal of the Linnean Society 122:351–365.

Joos, J., Pimiento, C., Miles, D.B., and Müller, J. 2022. Quaternary megafauna extinctions altered body size distribution in tortoises. Proceedings of the Royal Society B 289:20221947, 10 pp.

Jordan, M.A., Mumaw, V., Millspaw, N., Mockford, S.W., and Janzen, F.J. 2019. Range-wide phylogeography of Blanding's Turtle [*Emys* (= *Emydoidea*) *blandingii*]. Conservation Genetics 20(3):419–430.

Joseph-Ouni, M. and McCord, W.P. 2019a. A new species of *Elseya* (Testudines: Chelidae) from West Papua Province, Indonesia. The Batagur Monographs 1:21–41.

Joseph-Ouni, M. and McCord, W.P. 2019b. A new species of 'dwarf' *Elseya* (Testudines: Chelidae) in the northern mountains. The Batagur Monographs 1:43–54.

Joseph-Ouni, M. and McCord, W.P. 2022a. A new species of cryptic *Chelodina* (Testudines: Chelidae) from the Aramia River region, Papua New Guinea. The Batagur Monographs 4:7–26.

Joseph-Ouni, M. and McCord, W.P. 2022b. A new species of *Elseya* (Testudines: Chelidae) from Nabire, Indonesia. The Batagur Monographs 4:27–39.

Joseph-Ouni, M. and McCord, W.P. 2022c. A new species of *Elseya* (Testudines: Chelidae) from the Merauke River region, Indonesia. The Batagur Monographs 4:40–55.

Joseph-Ouni, M. and McCord, W.P. 2023a. A strange new subspecies of *Elseya caelatus* (Testudines: Chelidae) from the Berau Gulf region, New Guinea. The Batagur Monographs 8:7–27.

Joseph-Ouni, M. and McCord, W.P. 2023b. Morphological revision of the *Elseya (Hanwarachelys)* complex (Testudines: Chelidae) of southern New Guinea. The Batagur Monographs 8:85–151.

Joseph-Ouni, M. and McCord, W.P. 2025. 'Rediscovery' of the type series of *Elseya caelatus berau* (Testudines: Chelidae) - registration and current images. Chelonological Contributions 4:1–6.

Joseph-Ouni, M., McCord, W.P., Cann, J., and Smales, I. 2019a. A new species of *Macrochelodina* (Testudines: Chelidae) from the Northern Territory, Australia. The Batagur Monographs 1:5–20.

Joseph-Ouni, M., McCord, W.P., Cann, J., Smales, I., and Freeman, A. 2019b. A new subspecies of *Emydura subglobosa* (Testudines: Chelidae) from the Jardine River, Queensland, Australia. The Batagur Monographs 2:53–70.

Joseph-Ouni, M., McCord, W.P., Cann, J., Smales, I., Freeman, A., Sadlier, R., Couper, P., White, A., and Amey, A. 2020. The relics of Riversleigh: re-examination of the fossil record of *Elseya* (Testudines: Chelidae) with description of a new extant species from the Gulf of Carpentaria drainages, Queensland, Australia. The Batagur Monographs 3:7–69.

Joseph-Ouni, M., McCord, W.P., and Dwyer, Q. 2022a. Solomons' treasure: a new and endemic freshwater turtle fauna (Testudines: Chelidae). The Batagur Monographs 4:56–81.

Joseph-Ouni, M., Cann, J., and McCord, W.P. 2022b. The morphological identity of *Elseya dentata* (Testudines: Chelidae), northern Australia. The Batagur Monographs 5:7–34.

Joseph-Ouni, M., McCord, W.P., and Cann, J. 2022c. A new species of *Elseya* (Testudines: Chelidae), from the rivers of the Kimberley, Western Australia. The Batagur Monographs 5:35–54.

Joseph-Ouni, M., Cann, J., and McCord, W.P. 2022d. Strangers in the river: first documented sympatry of *Elseya* lineages (Testudines: Chelidae) in Australia. The Batagur Monographs 5:55–92.

Joseph-Ouni, M., McCord, W.P., and Cann, J. 2022e. An illustrated guide to external morphological character states for application to the species of the genus *Elseya* (Testudines: Chelidae). The Batagur Monographs 5:93–118.

Joseph-Ouni, M., White, A., Smales, I., Sadlier, R., Cann, J., McCord, W.P., Couper, P., Amey, A., and Freeman, A. 2022f. Further evidence in support of the recognition of the freshwater turtle *Elseya oneiros* (Testudines: Chelidae) from the Nicholson and Gregory rivers of northern Queensland. Memoirs of the Queensland Museum – Nature 63:93–112.

Joseph-Ouni, M., Cann, J., Turner, G.S., Freeman, A., and McCord, W.P. 2023a. The Johnstone River Snapping Turtle *Elseya stirlingi* (Testudines: Chelidae): taxonomy, morphology, history and distribution. The Batagur Monographs 8:28–84.

Joseph-Ouni, M., Cann, J., McCord, W.P., and Frewer, J. 2023b. Discovery of extant *Elseya (Elseya) lavarackorum* and confirmation of two lineages of 'living fossil' in the Nicholson-Gregory drainages, with a visual guide to scutation and osteological suture characters in northern Australian *Elseya* (Testudines: Chelidae). The Batagur Monographs 10:7–121.

Jourdan, C. 1862. Extraits des procès verbaux des séances de la Société impériale d'Agriculture et d'Industrie de Lyon, séance du 21 mars. Annales des Sciences Physiques et Naturelles de la Société d'Agriculture et d'Industrie de Lyon 6:32–33.

Joyce, W.G. 2007. Phylogenetic relationships of Mesozoic turtles. Bulletin of the Peabody Museum of Natural History 48(1):3–102.

Joyce, W.G. and Bourque, J.R. 2016. A review of the fossil record of turtles of the clade Pan-Kinosternoidea. Bulletin of the Peabody Museum of Natural History 57:57–95.

Joyce, W.G., Parham, J.F., and Gauthier, J.A. 2004. Developing a protocol for the conversion of rank-based taxon names to phylogenetically defined clade names, as exemplified by turtles. Journal of Paleontology 78:989–1013.

Joyce, W.G., Petričević, A., Lyson, T.R., and Czaplewski, N.J. 2012. A new Box Turtle from the Miocene/Pliocene boundary (Latest Hemphillian) of Oklahoma and a refined chronology of box turtle diversification. Journal of Paleontology 86(1):177–190.

Joyce, W.G., Parham, J.F., Lyson, T.R., Warnock, R.C.M., and Donoghue, P.C.J. 2013. A divergence dating analysis of turtles using fossil calibrations: an example of best practices. Journal of Paleontology 87:612–634.

Joyce, W.G., Parham, J.F., Anquetin, J., Claude, J., Danilov, I.G., Iverson, J.B., et al. 2020. Testudines. In: de Queiroz, K., Cantino, P.D., and Gauthier, J.A. (Eds.). Phylonyms – A Companion to the PhyloCode. Boca Raton: CRC Press, pp. 1049–1051.

Joyce, W.G., Anquetin, J., Cadena, E.A., Claude, J., Danilov, I.G., Evers, S.W., Ferreira, G.S., Gentry, A.D., Georgalis, G.L., Lyson, T.R., Pérez-García, A., Rabi, M., Sterli, J., Vitek, N.S., and Parham, J.F. 2021. A nomenclature for fossil and living turtles using phylogenetically defined clade names. Swiss Journal of Palaeontology 140:5, 45 pp., doi.org/10.1186/s13358-020-00211-x.

Kabisch, K. 1997. Ein ungewöhnlich grosser Knochenpanzer von *Chelus fimbriatus* (Schneider, 1783). Sauria, Berlin 19(1):45–46.

Kagayama, S., Ogano, D., Taniguchi, M., Mine, K., Ueno, S., Takahashi, H., Kamezaki, N., and Hasegawa, M. 2020. Species distribution modeling provides new insights into different spatial distribution patterns among native and alien freshwater turtles in Japan. Current Herpetology 39(2):147–159.

Kaiser, H. 2014. Best practices in herpetological taxonomy: errata and addenda. Herpetological Review 45:257–268.

Kaiser, H., Crother, B.I., Kelly, C.M.R., Luiselli, L., O'Shea, M., Ota, H., Passos, P., Schleip, W.D., and Wüster, W. 2013. Best practices: in the 21st Century, taxonomic decisions in herpetology are acceptable only when supported by a body of evidence and published via peer-review. Herpetological Review 44:8–23.

Kanberg, H. 1924. Über eine neue Schildkröte aus Kamerun. Zoologischer Anzeiger 60:195–197.

Kanberg, H. 1930. Eine neue Schildkröte aus Costa Rica. Zoologischer Anzeiger 88:161–162.

Karl, H.-V. 1987. Revision der fossilen Schildkröten (Reptilia, Testudines) von Jawa. Gothaer Museumsheft 14:37–44.

Karl, H.-V. 1995. Über einen fossilen Schildkrötenrest (Reptilia, Testudines) von Kuba. Mitteilungen aus dem Geologisch-Paläontologischen Institut der Universität Hamburg 78:129–137.

Karl, H.-V. and Paust, E. 2014. Die Geschichte der Europäischen Sumpfschildkröte in Deutschland 2: Checkliste zur prähistorischen Verbreitung der Europäischen Sumpfschildkröte (*Emys orbicularis* L., 1758) in Thüringen. Mainzer Naturwissenschaftliches Archiv 51:145–165.

Karl, H.-V. and Paust, E. 2024. Die Geschichte der Europäischen Sumpfschildkröte in Deutschland 3: Checkliste zur prähistorischen Verbreitung der Europäischen Sumpfschildkröte (*Emys orbicularis* L., 1758) in Sachsen-Anhalt und Sachsen (Testudines: Emydidae). Mainzer Naturwissenschaftliches Archiv 61:53–83.

Karl, H.-V. and Tichy, G. 1999. Die Taxonomie von *Homopus bergeri* Lindholm 1906 (Testudines: Testudinoidea). Mauritiana 17:277–284.

Karl, H.-V. and Tichy, G. 2002. Erstnachweis von *Testudo hermanni* Gmelin 1789 aus dem Pleistozän Zentraleuropas nördlich der Alpen (Testudines; Testudinidae). Revue de Paléobiologie 21:781–787.

Karl, S.A. and Bowen, B.W. 1999. Evolutionary significant units versus geopolitical taxonomy: molecular systematics of an endangered sea turtle (genus *Chelonia*). Conservation Biology 13:990–999.

Kehlmaier, C., Barlow, A., Hastings, A.K., Vamberger, M., Paijmans, J.L.A., Steadman, D.W., Albury, N.A., Franz, R., Hofreiter, M., and Fritz, U. 2017. Tropical ancient DNA reveals relationships of the extinct Bahamian giant tortoise *Chelonoidis alburyorum*. Proceedings of the Royal Society B 284(20162235):1–8.

Kehlmaier, C., Zhang, X., Georges, A., Campbell, P.D., Thomson, S., and Fritz, U. 2019a. Mitogenomics of historical type specimens of Australasian turtles: clarification of taxonomic confusion and old mitochondrial introgression. Scientific Reports 9(5841):1–12.

Kehlmaier, C., Graciá, E., Campbell, P.D., Hofmeyr, M.D., Schweiger, S., Martínez-Silvestre, A., Joyce, W., and Fritz, U. 2019b. Ancient mitogenomics clarifies radiation of extinct Mascarene giant tortoises (*Cylindraspis* spp.). Scientific Reports 9(17487):1–8.

Kehlmaier, C., Albury, N.A., Steadman, D.W., Graciá, E., Franz, R., and Fritz, U. 2021a. Ancient mitogenomics elucidates diversity of extinct West Indian tortoises. Scientific Reports 11:3224, 9 pp.

Kehlmaier, C., López-Jurado, L.F., Hernández-Acosta, N., Mateo-Miras, A., and Fritz, U. 2021b. "Ancient DNA" reveals that the scientific name for an extinct tortoise from Cape Verde refers to an extant South American species. Scientific Reports 11:97064, 7 pp.

Kehlmaier, C., Graciá, E., Ali, J.R., Campbell, P.D., Chapman, S.D., Deepak, V., Ihlow, F., Jalil, N.-E., Pierre-Huyet, L., Samonds, K.E., Vences, M., and Fritz, U. 2023. Ancient DNA elucidates the lost world of western Indian Ocean giant tortoises and reveals a new extinct species from Madagascar. Science Advances 9:eabq2574, 13 pp.

Kehlmaier, C., Fritz, U., and Kuchling, G. 2024a. Show me your true face: how many *Emydura* species occur in the Mitchell River Drainage, Kimberley, Australia? Salamandra 60:51–59.

Kehlmaier, C., Kuchling, G., Shea, G., Georges, A., Zhang, X., Fritz, U., and Thomson, S. 2024b. History and historical DNA: identity of *Chelodina intergularis* Fry, 1915 and type localities of *C. intergularis* and *C. rugosa* Ogilby, 1890. Zootaxa 5448(4):531–555.

Kehlmaier, C., Fritz, U., and Kuchling, G. 2025. The taxonomic quagmire of northern Australian snakenecked turtles (Testudines: Chelidae): *Chelodina*

kuchlingi—extinct or hiding in plain sight? Vertebrate Zoology 75:127–145.

Keller, C. and Busack, S.D. 2001. *Mauremys leprosa* (Schweigger, 1812) - Maurische Bachschildkröte. In: Fritz, U. (Ed.). Handbuch der Reptilien und Amphibien Europas, Band 3/IIIA. Wiesbaden: Aula-Verlag, pp. 57– 87.

Keller, C., Andreu, A.C., and Ramo, C. 1998. Aspects of the population structure of *Emys orbicularis hispanica* from southwestern Spain. In: Fritz, U.; Joger, U.; Podloucky, R.; and Servan, J. (Eds.). Proceedings of the EMYS Symposium Dresden 96. Mertensiella 10:147–158.

Keller, C., Villamarín, F., Bernhard, R., and da Silva, D.F. 2016. Checklist of chelonians from the upper Madeira River and the lower Madeira-Purus interfluvium (Brazilian Amazon), including a range expansion for *Podocnemis sextuberculata* Cornalia, 1849. Check List 12(4):1937(1–32).

Khabibullin, V. 2004. Distribution of *Emys orbicularis* in South Urals, Russia. Biologia, Bratislava 59(Suppl.14):27–32.

Khan, M.S. 2006. The Amphibians and Reptiles of Pakistan. Malabar, FL: Krieger Publishing, 311 pp.

Khan, M.Z., Safi, A., Fatima, F., Ghalib, S.A., Hashmi, M.U.A., Khan, I.S., Siddiqui, S., Zehra, A., and Hussain, B. 2015. An evaluation of distribution, status and abundance of freshwater turtles in the selected areas of Sindh and Khyber Pakhtunkhwa Provinces of Pakistan. Canadian Journal of Pure and Applied Sciences 9:3201–3219.

Khin Myo Myo and Platt, S.G. 2016. *Indotestudo elongata* (Elongated Tortoise). Maximum body size. Herpetological Review 47(2):285–286.

Khosatzky, L.I. 1987. Mesontology–a particular direction in the study of evolution. Ezhegodnik Vsesoyuznogo Paleontologicheskogo Obshchestva 30:50–66. [In Russian]

Khosatzky, L.I. and Młynarski, M. 1966. *Agrionemys*—nouveau genre de tortues terrestres (Testudinidae). Bulletin de l'Académie Polonaise des Sciences. Cl. 2. Série des Sciences Biologiques 14:123–125.

Kiester, A.R. 2019. The last of the phantastica? A tortoise lost and found. The Tortoise 2(4):55–62.

Killebrew, F.C. and Porter, D. 1989a. *Graptemys caglei* (Cagle's Map Turtle). Size maximum. Herpetological Review 20(3):70.

Killebrew, F.C. and Porter, D. 1989b. *Pseudemys texana* (Texas River Cooter). Size maximum. Herpetological Review 20(3):70.

Killebrew, F.C. and Porter, D. 1990. *Graptemys caglei* (Cagle's Map Turtle). Size. Herpetological Review 21(4):92.

Kimble, S.J.A., Rhodes, O.E., Jr., and Williams, R.N. 2014. Unexpectedly low rangewide population genetic structure of the imperiled eastern box turtle *Terrapene c. carolina*. PLoS ONE 9:e92274.

Kinaston, R., Buckley, H., Valentin, F., Bedford, S., Spriggs, M., Hawkins, S., and Herrscher, E. 2014. Lapita diet in Remote Oceania: new stable isotope evidence from the 3000-year-old Teouma site, Efate Island, Vanuatu. PLOS ONE 9(3)e90376:1–18.

Kindler, C., Branch, W.R., Hofmeyr, M.D., Maran, J., Široký, P., Vences, M., Harvey, J., Hauswaldt, J.S., Schleicher, A., Stuckas, H., and Fritz, U. 2012. Molecular phylogeny of African hinge-back tortoises (*Kinixys*): implications for phylogeography and taxonomy (Testudines: Testudinidae). Journal of Zoological Systematics and Evolutionary Research 50(3):192–201.

Kindler, C., Moosig, M., Branch, W.R., Harvey, J., Kehlmaier, C., Nagys, Z.T., Prokop, H., Široký, P., and Fritz, U. 2016. Comparative phylogeographies of six species of hinged terrapins (*Pelusios* spp.) reveal discordant patterns and unexpected differentiation in the *P. castaneus* / *P. chapini* complex and *P. rhodesianus*. Biological Journal of the Linnaean Society 117:305–321.

King, F.W. and Burke, R.L. 1989. Crocodilian, Tuatara, and Turtle Species of the World. A Taxonomic and Geographic Reference. Washington, DC: Association of Systematics Collections, 216 pp.

Kirsch, J.A.W. and Mayer, G.C. 1998. The platypus is not a rodent: DNA hybridization, amniote phylogeny and the palimpsest theory. Philosophical Transactions of the Royal Society of London 353(B):1221–1237.

Kitana, N. 1997. Sexual dimorphism and annual reproductive cycle of the Common Asiatic Softshell Turtle *Amyda cartilaginea*. Masters Thesis, Chulalongkorn University, Bangkok, Thailand.

Kitimasak, W., Thirakhupt, K., Boonyaratpalin, S., and Moll, D.L. 2005. Distribution and population status of the Narrow-Headed Softshell Turtle *Chitra* spp. in Thailand. Natural History Journal of Chulalongkorn University 5(1):31–42.

Kizirian, D.A., King, W.K., and Dixon, J.R. 1990. *Graptemys versa* (Texas Map Turtle). Size maximum and diet. Herpetological Review 21(3):60.

Klein, J.T. 1751. Quadrupedum Dispositio Brevisque Historia Naturalis. Lipsiae: B.C. Breitkopfium, 127 pp.

Klein, J.T. 1760. [Testudinata]. In: Behn, F.D. (Transl.). Jakob Theodor Kleins Classification und Kurze Geschichte der Vierfüssigen Thiere. Lübeck: Jonas Schmidt, 381 pp.

Klemens, M.W. 1993. Amphibians and Reptiles of Connecticut and Adjacent Regions. State Geological and Natural History Survey of Connecticut, Bulletin No. 12, 318 pp.

Kluge, A. 1984. Type-specimens of reptiles in the University of Michigan Museum of Zoology. Miscellaneous Publications, Museum of Zoology, University of Michigan 167:1–85.

Knauss, G.E., Joyce, W.G., Lyson, T.R., and Pearson, D. 2011. A new kinosternoid from the Late Cretaceous Hell Creek Formation of North Dakota and Montana and the origin of the *Dermatemys mawii* lineage. Palaeontologische Zeitschrift 85:125–142.

Koné, D., Diarrassouba, A., Diagne, T., and Luiselli, L. 2024. On the occurrence of *Pelusios gabonensis* in the Taï National Park (Côte d'Ivoire). African Journal of Ecology 62:e13320, 3 pp.

Koshikawa, A. 1982. Three new species of reptiles from Hainan Island, Guangdong Province. Translation and introduction. Smithsonian Herpetological Information Service 53:1–9.

Kot, C.Y., Fujioka, E., Dimatteo, A.D., Bandimere, A., Wallace, B.P., Hutchinson, B.J., Cleary, J., Halpin, P.N., and Mast, R.B. 2023. The State of the World's Sea Turtles Online Database: Data provided by the SWOT Team and hosted on OBIS-SEAMAP. Oceanic Society, Conservation International, IUCN Marine Turtle Specialist Group (MTSG), and Marine Geospatial Ecology Lab, Duke University. http://seamap.env.duke.edu/swot.

Kou, Z. 1989. *Cyclemys* from Yunnan, a description of a new species and a new record to China (Testudinata: Emydidae). In: Matsui, M., Hikida, T. and Goris, R.C. (Eds.). Current Herpetology in East Asia. Proceedings of the Second Japan–China Herpetological Symposium. Herpetological Society of Japan 1989:193–197.

Kraus, F. 2009. Alien Reptiles and Amphibians. A Scientific Compendium and Analysis. Springer Verlag, 563 pp.

Krefft, G. 1876. Notes on Australian animals in New Guinea with description of a new species of fresh water tortoise belonging to the genus *Euchelymys* (Gray). Annali del Museo Civico di Storia Naturale Giacomo Doria (1)8:390–394.

Krell, F.-T. 2021. Suppressing works of contemporary authors using the Code's publication requirements is neither easy nor advisable. Bulletin of Zoological Nomenclature 78:61–67.

Krenz, J.G., Naylor, G.J.P., Shaffer, H.B., and Janzen, F.J. 2005. Molecular phylogenetics and evolution of turtles. Molecular Phylogenetics and Evolution 37:178–191.

Krysko, K.L., Burgess, J.P., Rochford, M.R., Gillette, C.R., Cueva, D., Enge, K.M., Somma, L.A., Stabile, J.L., Smith, D.C., Wasilewski, J.A., Kieckhefer, G., III, Granatosky, M.C., and Nielsen, S.V. 2011. Verified non-indigenous amphibians and reptiles in Florida from 1863 through 2010: outlining the invasion process and identifying invasion pathways and stages. Zootaxa 3028:1–64.

Krysko, K.L., Enge, K.M., and Moler, P.E. 2019. Amphibians and Reptiles of Florida. Gainesville: University of Florida Press, 706 pp.

Kuchling, G. 1988. Gonadal cycles of the Western Australian long-necked turtles *Chelodina oblonga* and *Chelodina steindachneri* (Chelonia: Chelidae). Records of the Western Australian Museum 14:189–198.

Kuchling, G. 2010. Taxonomy and nomenclature of the longneck turtle (genus *Chelodina*) from south-western Australia. Records of the Western Australian Museum 25:449–454.

Kuchling, G. 2020. Revised type locality and distribution of the Data Deficient *Chelodina kuchlingi* and a review of its status as a distinct species. Chelonian Conservation and Biology 19(1):48–57.

Kuchling, G., Rhodin, A.G.J., Ibarrondo, B.R., and Trainor, C.R. 2007. A new subspecies of the snakeneck turtle *Chelodina mccordi* from Timor-Leste (East Timor) (Testudines: Chelidae). Chelonian Conservation and Biology 6(2):213–222.

Kuhl, H. 1820. Beiträge zur Kenntniss der Amphibien. In: Kuhl, H. Beiträge zur Zoologie und vergleichenden Anatomie. Erste Abtheilung. Beiträge zur Zoologie. Frankfurt: Hermannschen Buchhandlung, pp. 75–132.

Kuhn, O. 1960. Die Familien der fossilen Amphibien und Reptilien. Bericht der Naturforschenden Gesellschaft in Bamberg 37:20–52.

Kuhn, O. 1964. Testudines. In: Westphal, F. (Ed.). Fossilium Catalogus. I:

Animalia, Pars 107. Gravenhage: W. Junk, 299 pp.

Kundu, S., Laskar, B.A., Venkataraman, K., Banerjee, D., and Kumar, V. 2016. DNA barcoding of *Nilssonia* congeners corroborates existence of wild *N. nigricans* in northeast India. Mitochondrial DNA Part A: Mapping, Sequencing and Analysis 27:2753–2756.

Kundu, S., Kumar, V., Murthy, B.H.C.K., and Chandra, K. 2018a. Chelonian types of National Zoological Collections. Kolkata: Zoological Survey of India, 70 pp.

Kundu, S., Kumar, V., Laskar, B.A., Tyagi, K., and Chandra, K. 2018b. Morphology and genetic variation in the endangered tortoise *Manouria emys*: distinct lineages or plastron anomalies? Mitochondrial DNA B: Resources 3:166–170.

Kundu, S., Kumar, V., Tyagi, K., Chakraborty, R., Singha, D., Rahaman, I., Pakrashi, A., and Chandra, K. 2018c. Complete mitochondrial genome of Black Soft-shell Turtle (*Nilssonia nigricans*) and comparative analysis with other Trionychidae. Scientific Reports 8:17378, 11 pp.

Kundu, S., Kumar, V., Tyagi, K., and Chandra, K. 2020. The complete mitochondrial genome of the endangered Assam Roofed Turtle, *Pangshura sylhetensis* (Testudines: Geoemydidae): genomic features and phylogeny. PLOS ONE, doi.org/10.1371/journal.pone.0225233.

Kundu, S., Mukherjee, T., Kim, A.R., Lee, S.-R., Mukherjee, A., Jung, W.-K., and Kim, H.-W. 2023. Mitochondrial DNA and distribution modelling evidenced the lost genetic diversity and wild-residence of Star Tortoise, *Geochelone elegans* (Testudines: Testudinidae) in India. Animals 13:150, doi.org/10.3390/ani13010150, 16 pp.

Kunz, T.S., Ghizoni-Jr., I.R., Cherem, J.J., Bressan, R.F., Leonardi, S.B., and Zanotelli, J.C. 2018. New records, threats and conservation of *Phrynops williamsi* (Testudines: Chelidae) in southern Brazil. Herpetology Notes 11:147–152.

Kurck, C. 1917. Den forntida utbredningen af kärrsköldpaddan *Emys orbicularis* (Lin.) i Sverige, Danmark och angränsande länder. [The prehistoric distribution of the pond turtle *Emys orbicularis* (Lin.) in Sweden, Denmark and adjacent countries]. Lunds Universitets Årsskrift (N.F.) (2)13(9):1–129. [in Swedish]

Kusrini, M.D., Mardiastuti, A., Darmawan, B., Mediyansyah, and Muin, A. 2009. Survey on harvest and trade of Asiatic softshell turtle *Amyda cartilaginea* in East Kalimantan, Indonesia. Bogor: Nature Harmony, 11 pp.

Kuzmin, S.L. 2002. The Turtles of Russia and Other Ex-Soviet Republics (Former Soviet Union). Frankfurt am Main: Edition Chimaira, 159 pp.

Lacepède, B.G.E. de. 1788. Histoire Naturelle des Quadrupèdes Ovipares et des Serpens. Tome Premier. Paris: Hôtel de Thou, 651 pp. [octavo edition, 8º].

Lamb, T., Avise, J.C., and Gibbons, J.W. 1989. Phylogeographic patterns in mitochondrial DNA of the desert tortoise (*Xerobates agassizii*), and evolutionary relationships among the North American gopher tortoises. Evolution 43:76–87.

Lambert, M.R.K. 1993. On growth, sexual dimorphism, and the general ecology of the African spurred tortoise *Geochelone sulcata*, in Mali. Chelonian Conservation and Biology 1:37–46.

Lambert, M.R.K. 1995. On geographical size variation, growth, and sexual dimorphism of the leopard tortoise, *Geochelone pardalis*, in Somaliland. Chelonian Conservation and Biology 1:269–278.

Lambert, M.R.K., Campbell, K.L.I., and Kabigumila, J.D. 1998. On growth and morphometrics of leopard tortoises, *Geochelone pardalis*, in Serengeti National Park, Tanzania, with observations on effects of bushfires and latitudinal variation in populations of eastern Africa. Chelonian Conservation and Biology 3(1):46–57.

Lapparent de Broin, F. de. 2000a. Les chéloniens de Sansan. Mémoires du Muséum national d'Histoire Naturelle 183:219–261.

Lapparent de Broin, F. de. 2000b. African chelonians from the Jurassic to the present. A preliminary catalog of the African fossil chelonians. Palaeontologica Africana 36:43–82.

Lapparent de Broin, F. de, Bour, R., Parham, J.F., and Perälä, J. 2006a. *Eurotestudo*, a new genus for the species *Testudo hermanni* Gmelin, 1789 (Chelonii, Testudinidae). Comptes Rendus Palevol 5:803–811.

Lapparent de Broin, F. de, Bour, R., and Perälä, J. 2006b. Morphological definition of *Eurotestudo* (Testudinidae, Chelonii): second part. *Eurotestudo* (Chelonii, Testudinidae), définition, approche morphologique. Deuxième partie. Annales de Paléontologie 92:325–357.

Largen, M. and Spawls, S. 2010. The Amphibians and Reptiles of Ethiopia and Eritrea. Frankfurt: Edition Chimaira, 693 pp.

Lataste, F. 1881. Diagnose d'une nouvelle tortue *Testudo graeca bettai*, n. sbsp. Le Naturaliste (1)3:396.

Lataste, F. 1886. Description d'une tortue nouvelle du Haut Sénégal (*Homopus nogueyi*). Le Naturaliste (2)8:286–287.

Lataste, F. 1888. Description d'une tortue nouvelle originaire du Haut-Sénégal (*Cinixys dorri*, n. sp.). Le Naturaliste (2)10:164–166.

Latreille, P.A. 1800. Histoire naturelle des Salamandres de France, précédée d'un tableau méthodique des autres Reptiles indigènes. Paris: Imprimerie de Crapelet, 61 pp.

Latreille, P.A. 1801. Histoire Naturelle des Reptiles. In: Sonnini, C.S. and Latreille, P.A. Histoire Naturelle des Reptiles, avec figures dessinées d'après nature. Tome Premier. Première Partie. Quadrupèdes et Bipèdes Ovipares. Paris: Imprimerie de Crapelet, 280 pp.

Latreille, P.A. 1825. Familles Naturelles du Règne Animal, exposées succinctement et dans un ordre analytique, avec l'indication de leurs genres. Paris: Baillière, 570 pp.

Laurent, R.F. 1956. Contribution à l'herpétologie de la région des grands lacs de l'Afrique centrale. I. Généralités. II. Chéloniens. III. Ophidiens. Annales du Musée Royal du Congo Belge, Tervuren, Sciences Zoologiques 48:5–390.

Laurent, R.F. 1965. A contribution to the knowledge of the genus *Pelusios* (Wagler). Annales du Musée Royal de l'Afrique Centrale, Sciences Zoologiques, Tervuren 135:1–33.

Laurin, M. and Reisz, R.R. 1995. A reevaluation of early amniote phylogeny. Zoological Journal of the Linnean Society 113:165–223.

Le, M. 2007. Conservation of turtles in Vietnam: a survey of Cat Tien National Park. Oryx 41(4):544–547.

Le, M. and McCord, W.P. 2008. Phylogenetic relationships and biogeographical history of the genus *Rhinoclemmys* Fitzinger, 1835 and the monophyly of the turtle family Geoemydidae (Testudines: Testudinoidea). Zoological Journal of the Linnean Society 153:751–767.

Le, M. and Pritchard, P.C.H. 2009. Genetic variability of the critically endangered softshell turtle, *Rafetus swinhoei*: a preliminary report. Proceedings of the First Vietnamese National Symposium on Reptiles and Amphibians, pp. 84–92.

Le, M., Raxworthy, C.J., McCord, W.P., and Mertz, L. 2006. A molecular phylogeny of tortoises (Testudines: Testudinidae) based on mitochondrial and nuclear genes. Molecular Phylogenetics and Evolution 40:517–531.

Le, M., McCord, W.P., and Iverson, J.B. 2007. On the paraphyly of the genus *Kachuga* (Testudines: Geoemydidae). Molecular Phylogenetics and Evolution 45:398–404.

Le, M., Reid, B.N., McCord, W.P., Naro-Maciel, E., Raxworthy, C.J., Amato, G., and Georges, A. 2013. Resolving the phylogenetic history of the short-necked turtles, genera *Elseya* and *Myuchelys* (Testudines: Chelidae) from Australia and New Guinea. Molecular Phylogenetics and Evolution 68:251–258.

Le, M., Duong, H.T., Dinh, L.D., Nguyen, T.Q., Pritchard, P.C.H., and McCormack, T. 2014. A phylogeny of softshell turtles (Testudines: Trionychidae) with reference to the taxonomic status of the critically endangered, giant softshell turtle, *Rafetus swinhoei*. Organisms, Diversity and Evolution 14:279-293.

Le, M.D., McCormack, T.E.M., Hoang, H.V., Duong, H.T., Nguyen, T.Q., Ziegler, T., Nguyen, H.D., and Ngo, H.T. 2020. Threats from wildlife trade: the importance of genetic data in safeguarding the endangered Four-eyed Turtle (*Sacalia quadriocellata*). Nature Conservation 41:91–111.

Le, T.B., Le, Q.H., Tran, M.L., Phan, T.H., Phan, M.T., Tran, T.T.H., Pham, T.T., Nguyen, D.T., Nong, V.H., Phan, V.C., Dinh, D.K., Truong, N.H., and Ha, D.D. 2010. Comparative morphological and DNA analysis of specimens of giant freshwater soft-shelled turtle in Vietnam related to Hoan Kiem turtle. Tạp Chí Công Nghệ Sinh Học [Journal of Biotechnology, Vietnamese Academy of Science and Technology] 8(3A):949–954.

Le Conte, J. 1830. Description of the species of North American tortoises. Annals of the Lyceum of Natural History, New York 3:91–131.

Le Conte, J. 1854. Description of four new species of *Kinosternum*. Proceedings of the Academy of Natural Sciences, Philadelphia 7:180–190.

Le Conte, J. 1860. Description of two new species of tortoises. Proceedings of the Academy of Natural Sciences, Philadelphia 11(1859)[1860]:4–7.

Le Duc, O., Van, T.P., Leprince, B., Bordes, C., Tuan, A.N., Benansio, J.S., Pacini, N., Luu, V.Q., and Luiselli, L. 2020. Fishers, dams, and the potential survival of the world's rarest turtle, *Rafetus swinhoei*, in two river basins in northern Vietnam. Aquatic Conservation: Marine and Freshwater Ecosystems 2020, doi:10.1002/aqc.3317, 14 pp.

Lee, D.S. and Carey, E. 2001. Conservation concerns facing the Inagua slider. Turtle and Tortoise Newsletter 3:7–9.

Lee, D.S. and Ross, J.P. 2001. The Cat Island turtle: a reptile of problematic origin; including a bibliography of the genus *Trachemys* in the West Indies

region. In: Clark-Simpson, C.A. and Smith, G.W. (Eds.). Proceedings of the 8th Symposium on the Natural History of the Bahamas. San Salvador, Bahamas: Gerace Research Center, pp. 36–47.

Legler, J.M. 1959. A new tortoise, genus *Gopherus*, from north-central Mexico. University of Kansas Publications of the Museum of Natural History 11(5):335–343.

Legler, J.M. 1960. A new subspecies of slider turtle (*Pseudemys scripta*) from Coahuila, Mexico. University of Kansas Publications of the Museum of Natural History 13(3):73–84.

Legler, J.M. 1965. A new species of turtle, genus *Kinosternon*, from Central America. University of Kansas Publications of the Museum of Natural History 15(13):617–625.

Legler, J.M. 1981. The taxonomy, distribution, and ecology of Australian freshwater turtles (Testudines: Pleurodira: Chelidae). National Geographic Society Research Reports 13:391–404.

Legler, J.M. 1990. The genus *Pseudemys* in Mesoamerica: taxonomy, distribution and origins. In: Gibbons, J.W. (Ed.). Life History and Ecology of the Slider Turtle. Washington, DC: Smithsonian Institution Press, pp. 82–105.

Legler, J.M. and Cann, J. 1980. A new genus and species of chelid turtle from Queensland, Australia. Contributions in Science, Natural History Museum of Los Angeles County 324:1–18.

Legler, J.M. and Vogt, R.C. 2013. The Turtles of Mexico: Land and Freshwater Forms. Berkeley: University of California Press, 416 pp.

Legler, J.M. and Webb, R.G. 1970. A new slider turtle (*Pseudemys scripta*) from Sonora, Mexico. Herpetologica 26(2):157–168.

Legler, J.M., Smith, H.M., and Smith, R.B. 1980. *Testudo scripta* Schoepff, 1792: *Emys cataspila* Günther, 1885: proposed conservation (Reptilia, Testudines). Bulletin of Zoological Nomenclature 37:240–246.

Lehr, E., Fritz, U., and Obst, F.J. 1998. *Cuora galbinifrons picturata* subsp. nov., eine neue Unterart der Hinterindischen Scharnierschildkröte. Herpetofauna 20(113):5–11.

Leidy, J. 1868a. Notice of some vertebrate remains from Harden Co., Texas. Proceedings of the Academy of Natural Sciences, Philadelphia 1868:174–176.

Leidy, J. 1868b. Notice of some vertebrate remains from the West Indian islands. Proceedings of the Academy of Natural Sciences of Philadelphia 1868:178–180.

Leidy, J. 1877. Description of vertebrate remains, chiefly from the phosphate beds of South Carolina. Journal of the Academy of Natural Sciences of Philadelphia (2)8:209–261.

Leidy, J. 1889a. Fossil vertebrates from Florida. Proceedings of the Academy of Natural Sciences, Philadelphia 1889:96–97.

Leidy, J. 1889b. Description of vertebrate remains from Peace Creek, Florida. Transactions of the Wagner Free Institute of Science of Philadelphia 2:19–31.

Lemos-Espinal, J.A., Smith, H.M., Dixon, J.R., and Cruz, A. 2015. Amphibians and Reptiles of Sonora, Chihuahua and Coahuila, Mexico. Tlalpan, Mexico: CONABIO, Vol. II, 668 pp.

Lemos-Espinal, J.A., Smith, G.R., Woolrich-Piña, G.A., and Cruz, A. 2017. Amphibians and reptiles of the state of Chihuahua, Mexico, with comparisons with adjoining states. ZooKeys 658:105–130.

Lemos-Espinal, J.A., Smith, G.R., and Valdez-Lares, R. 2019. Amphibians and Reptiles of Durango, Mexico. Rodeo, NM: ECO Herpetological Publishing.

Lenk, P., Fritz, U., Joger, U., and Wink, M. 1999. Mitochondrial phylogeography of the European pond turtle, *Emys orbicularis* (Linnaeus 1758). Molecular Ecology 8:1911–1922.

Leschenault de la Tour, J.-B. 1812. [*Emys couro*]. In: Schweigger, A.F. Prodromus monographiae Cheloniorum. Königsberger Archiv für Naturwissenschaft und Mathematik 1:271–368, 406–462 [p.315].

Lescure, J. and Fretey, J. 1976 ["1975"]. Etude taxinomique de *Phrynops (Batrachemys) nasutus* (Schweigger) (Testudinata, Chelidae). Bulletin du Muséum Nationale d'Histoire Naturelle, Paris, Zoologie 239(1975):1317–1328.

Lesson, R.P. 1830. Centurie Zoologique, ou Choix d'Animaux Rares, Nouveaux ou Imparfaitement Connus. Paris: F.G. Levrault, 235 pp.

Lesson, R.P. 1831a. Catalogue des Reptiles qui font partie d'une Collection zoologique recueillie dans l'Inde continentale ou en Afrique, et apportée en France par M. Lamare-Piquot. Bulletin des Sciences Naturelles et Géologiques, Paris 25(2):119–123.

Lesson, R.P. 1831b ["1834"]. Reptiles. In: Bélanger, C. (Ed.). Voyage aux Indes-Orientales, par le Nord de l'Europe, les Provinces du Caucase, la Géorgie, l'Arménie et la Perse, suivi de Détails topographiques, statistiques et autres sur le Pégou, les Iles de Java, de Maurice et de Bourbon, sur le Cap de Bonne-Espérance et Sainte-Hélène, pendant les Années 1825, 1826, 1827, 1828 et 1829. Zoologie. Paris: A. Bertrand, pp. 289–336.

Lesson, R.P. 1832. Illustrations de Zoologie, ou Recueil de figures d'Animaux peintes d'après nature. Pl. 7. Paris: A. Bertrand, 60 pls.

LeSueur, C.A. 1817. An account of an American species of tortoise, not noticed in the systems. Journal of the Academy of Natural Sciences, Philadelphia 1:86–88.

LeSueur, C.A. 1827. Note sur deux espèces de tortues, du genre *Trionyx* de M. Geoffroy-Saint-Hilaire. Mémoires du Muséum d'Histoire Naturelle, Paris 15:257–268.

Leuteritz, T.E.J. and Ganz, D.T. 2013. Sexual dimorphism in Radiated Tortoises (*Astrochelys radiata*). In: Castellano, C.M., Rhodin, A.G.J., Ogle, M., Mittermeier, R.A., Randriamahazo, H., Hudson, R., and Lewis, R.E. (Eds.). Turtles on the Brink in Madagascar: Proceedings of Two Workshops on the Status, Conservation, and Biology of Malagasy Tortoises and Freshwater Turtles. Chelonian Research Monographs 6:105–112.

Li, J., Lu, Y., Zan, J., and Nie, L. 2017. Complete mitochondrial genome of the *Cyclemys pulchristriata* (Chelonia: Geoemydidae). Mitochondrial DNA Part B: Resources 2:403–404.

Li, M.-M., Luo, P.-Z., Chen, D.-H., Shi, H.-T., and Lin, L. 2023. Genetic diversity analysis of Pan's Box Turtle *Cuora pani* based on DNA molecular markers. Chinese Journal of Zoology 58(1):126–134. [in Chinese]

Li, W., Zhang, X.C., Zhao, J., Shi, Y., and Zhu, X.P. 2015. Complete mitochondrial genome of *Cuora trifasciata* (Chinese three-striped box turtle), and a comparative analysis with other box turtles. Gene 555:169–177.

Li, Z.C., Xiao, Z., and Liu, S.R. 2011. Amphibians and Reptiles of Guangdong. Guangzhou: Guangdong Science and Technology Press. [in Chinese]

Li, Z.X. 1958. Report on the investigation of reptiles of Hainan Island. Chinese Journal of Zoology 2(4):234–239.

Liaw, Y.-L. and Tsai, C.-H. 2022. Taxonomic revision of *Chinemys pani* (Testudines: Geoemydidae) from the Pleistocene of Taiwan and its implications of conservation paleobiology. Anatomical Record 306:1501–1507.

Lichtenstein, H. and von Martens, C.E. 1856. Nomenclator reptilium et amphibiorum Musei Zoologici Berolinensis. Namenverzeichniss der in der zoologischen Sammlung der Königlichen Universität zu Berlin ausgestellten Arten von Reptilien und Amphibien nach ihren Ordnungen, Familien und Gattungen. Berlin: Königlichen Akademie der Wissenschaften, 48 pp.

Lidth de Jeude, T.W. van. 1893. On a new species of the genus *Testudo*. Notes from the Leyden Museum 15:312–313.

Lieb, C.S. 2023. Type specimens, type localities, synonymies, and authors/collectors of amphibians and reptiles described from Texas. In: Schmidly, D.J., Bradley, R.D., Bradley, L.C., and Yancey, F.D., II (Eds.). Taxonomic Catalogs for the Recent Terrestrial Vertebrates (Species and Subspecies) Described from Texas. Special Publications, Museum of Texas Tech University, No. 77, pp. 213–378.

Liebing, N., Praschag, P., Gosh, R., Vasudevan, K., Rashid, S.M.A., Rao, D.Q., Stuckas, H., and Fritz, U. 2012. Molecular phylogeny of the softshell turtle genus *Nilssonia* revisited, with first records of *N. formosa* for China and wild-living *N. nigricans* for Bangladesh. Vertebrate Zoology 62:261–272.

Lien, A., Conix, S., Zachos, F.E., Christidis, L., van Dijk, P.P., Bánki, O.S., Barik, S.K., Buckeridge, J.S., Costello, M.J., Hobern, D., Montgomery, N., Nikolaeva, S., Pyle, R.L., Thiele, K.R., Thomson, S.A., Zhang, Z., and Garnett, S.T. 2021. Towards a global list of accepted species IV. Overcoming fragmentation in the governance of taxonomic lists. Organisms Diversity & Evolution: doi.org/10.1007/s13127-021-00499-8, 11 pp.

Lim, B.L. and Das, I. 1999. Turtles of Borneo and Peninsular Malaysia. Kota Kinabalu: Natural History Publications (Borneo), 151 pp.

Limpus, C. 1985. A study of the loggerhead sea turtle, *Caretta caretta*, in eastern Australia. Ph.D. Thesis, University of Queensland, Brisbane.

Limpus, C. 2008. Freshwater turtles in the Mary River: review of biological data for turtles in the Mary River, with emphasis on *Elusor macrurus* and *Elseya albagula*. Brisbane: Queensland Government, 84 pp.

Lin, L., Sun, L., Wang, W., and Shi, H. 2018a. Taxonomic status and nomenclature of Four Eye-spotted Turtle from Hainan Island. Sichuan Journal of Zoology 37:435–438.

Lin, L., Hu, Q., Fong, J.J., Yang, J., Chen, Z., Zhou, F., Wang, J., Xiao, F., and Shi, H. 2018b. Reproductive ecology of the endangered Beal's-eyed turtle, *Sacalia bealei*. PeerJ 6:e4997.

Lin, L., Chen, H., Wang, Z., Gaillard, D., Zhai, X., and Shi, H. 2020. Characterization and comparison of mitogenomes of three 'eyed' turtles *Sacalia* spp.

Mitochondrial DNA Part B 5:3206–3208.

Lin, L., Chen, H.-Q., Gaillard, D., Shi, H.-T., and Luo, S.-J. 2022. Phylogeography of the endangered "eyed" turtles (genus *Sacalia*) and the discovery of a lineage derived from natural interspecific hybridization. Ecology and Evolution 12:e9545, 17 pp.

Lindeman, P.V. 2003. Sexual difference in habitat use of Texas map turtles (Emydidae: *Graptemys versa*) and its relationship to size dimorphism and diet. Canadian Journal of Zoology 81:1185–1191.

Lindeman, P.V. 2009. On the type locality and type specimen of *Testudo geographica* LeSueur 1817. Chelonian Conservation and Biology 8(1):95–98.

Lindeman, P.V. 2013. The Map Turtle and Sawback Atlas: Ecology, Evolution, Distribution, and Conservation. Norman, OK: University of Oklahoma Press, 460 pp.

Lindeman, P.V., Louque, I., Huntzinger, C., Lyons, E., Shively, S.H., and Selman, W. 2015. Eye color and chin pattern in the turtle *Graptemys pseudogeographica* in the Calcasieu River drainage of Louisiana, with comparison to adjacent drainages. Herpetological Review 46:179–185.

Lindholm, W.A. 1906. Beschreibung einer neuen Schildkrötenart aus Deutsch-Sudwestafrika nebst Bemerkungen über die Gattung *Homopus* D. et B. Jahrbücher des Nassauischen Vereins für Naturkunde, Wiesbaden 59:345–351.

Lindholm, W.A. 1929. Revidiertes Verzeichnis der Gattungen der rezenten Schildkröten nebst Notizen zur Nomenklatur einiger Arten. Zoologischer Anzeiger 81:275–295.

Lindholm, W.A. 1931. Über eine angebliche *Testudo*-Art aus Südchina. Zoologischer Anzeiger 97:27–30.

Link, H.F. 1807. Beschreibung der Naturalien-Sammlung der Universität zu Rostock, Volume 2. Rostock: Adlers Erben, 100 pp.

Linnaeus, C. 1745. Amphibia Gyllenborgiana, Dissertatione Academica, descripta et publico examini subjecta a Barth. Rudolpho Hast. Upsaliae: 34 pp. Reprinted in Linnaeus (1749a).

Linnaeus, C. 1746. Museum Adolpho-Fridericianum, publico bonorum examini submittit Laurentius Balk. Holmiae [Stockholm]: Laurentii Salvii, 48 pp. Reprinted in Linnaeus (1749b).

Linnaeus, C. 1749a. Amphibia Gyllenborgiana, descripta a Barth. Rudolpho Hast. Amoenitates Academicae 1:107–140. Reprint of Linnaeus (1745).

Linnaeus, C. 1749b. Museum Adolpho-Fridericianum, propositum a Laurent. Balk. (Museum Principis). Amoenitates Academicae 1:277–327. Reprint of Linnaeus (1746).

Linnaeus, C. 1754. Museum S:ae R:ae M:tis Adolphi Friderici Regis Svecorum, Gothorum, Vandalorumque, in quo Animalia Rarioria Imprimis, et Exotica: Quadrupedia, Aves, Amphibia, Pisces, Insecta, Vermes Describuntur et Determinantur. Tomus I. Holmiae [Stockholm]: Typographia Regia, 102 pp, 33 pls.

Linnaeus, C. 1758. Systema Naturae, per Regna Tria Naturae, secundum Classes, Ordines, Genera, Species, cum Characteribus, Differentiis, Synonymis, Locis. Tomus I. Editio Decima, Reformata. [10th Ed.]. Holmiae [Stockholm]: Laurentii Salvii, 824 pp.

Linnaeus, C. 1764. Museum S:ae R:ae M:tis Adolphi Friderici Regis Svecorum, Gothorum, Vandalorumque, in quo Animalia Rariora Imprimis et Exotica: Aves, Amphibia, Pisces Describuntur. Tomi Secundi Prodromus. Holmiae [Stockholm]: Laur. Salvii, 110 pp. [written in 1754]

Linnaeus, C. 1766. Systema Naturae. Editio Duodecima, Reformata. Tomus I, Pars I, Regnum Animale. [12th Ed.]. Holmia [Stockholm]: Laurentii Salvii, 532 pp.

Linné, C. von, Jr. and Thunberg, C.P. 1780. Inventeringen på Academiska i Horto Botanico varande Museum. Unpublished manuscript, Uppsala University Zoological Museum. Reproduced in Holm (1957:10–18). [in Swedish]

Linthoi, N. and Sharma, D.K. 2009. Turtles and tortoises in Manipur. In: Vasudevan, K. (Ed.). Freshwater Turtles and Tortoises of India. ENVIS Bulletin: Wildlife and Protected Areas 12(1):49–51.

Liu, J. 2019. Fernandina found: a Q&A with Forrest Galante of *Extinct or Alive*. The Tortoise 2(4):63–73.

Liu, J.-Y., Fu, W., Ma, Y.-H., Zhou, T., Blanck, T., Zhou, G.-J., Xiao, Y.-M., and Mo, X.-Y. 2024. Shedding light into the confusing history of *Pelodiscus axenaria* (Zhou, Zhang et Fang, 1991), clarifying and correcting its taxonomic position within the *P. axenaria* complex with the description of yet another *Pelodiscus* species. Russian Journal of Herpetology 31(6):337–350.

Liu, X., Li, W., Ye, Z., Zhu, Y., Hong, X., and Zhu, X. 2019. Morphological characterization and phylogenetic relationships of Indochinese box turtles – the *Cuora galbinifrons* complex. Ecology and Evolution 9:13030–13042.

Li Vigni, F. (Ed.). 2013. A Life for Reptiles and Amphibians, Volume 1. A Collection of 55 Interviews on "How to become a Herpetologist." Frankfurt: Edition Chimaira, 495 pp.

Loc-Barragán, J.A. and Iverson, J.B. 2025 ["2024"]. *Kinosternon cora* (Cora Mud Turtle). Distribution and morphometrics. Herpetological Review 55(2) [2024]:249–250.

Loc-Barragán, J.A., Reyes-Velasco, J., Woolrich-Piña, G.A., Grünwald, C.I., Venegas de Anaya, M., Rangel-Mendoza, J.A., and López-Luna, M.A. 2020. A new species of mud turtle of genus *Kinosternon* (Testudines: Kinosternidae) from the Pacific coastal plain of northwestern Mexico. Zootaxa 4885(4):509–529.

Loehr, V.J.T., Henen, B.T., and Hofmeyr, M.D. 2006. Shell characteristics and sexual dimorphism in the Namaqualand speckled padloper, *Homopus signatus signatus*. African Journal of Herpetology 55:1–11.

Loire, E. and Galtier, N. 2017. Lacking conservation genomics in the giant Galápagos tortoise. bioRxiv 101980, doi:10.1101/101980, 14 pp.

Loire, E., Chiari, Y., Bernard, A., Cahais, V., Romiguier, J., Nabholz, B., Lourenço, J.M., and Galtier, N. 2013. Population genomics of the endangered Galapagos tortoise. Genome Biology 14:R136.

Lönnberg, E. 1896. Linnean type-specimens of birds, reptiles, batrachians and fishes in the Zoological Museum of the R. University in Upsala. Bihang Svenska Vetenskaps Akademien Handlingar 22(4):1–45.

Loomis, F.B. 1927. A giant tortoise from Florida. American Journal of Science (5)13:435–439.

López-Jurado, L.F., Mateo, J.A., and García-Márquez, M. 1998. La tortuga fósil de la Isla de Sal (Archipiélago de Cabo Verde). Revista Española de Herpetología 12:111–114.

López-Luna, M.A., Cupul-Magaña, F.G., Escobedo-Galván, A.H., González-Hernández, A.J., Centenero-Alcala, E., Rangel-Mendoza, J.A., Ramírez-Ramírez, M.M., and Cazares-Hernández, E. 2018. A distinctive new species of mud turtle from western Mexico. Chelonian Conservation and Biology 17(1):2–13.

López-Luna, M.A., Escobedo-Galván, A.H., and Cupul-Magaña, F.G. 2020. Expansion of the geographic scope and the maximum size of *Kinosternon chimalhuaca* (Testudines: Kinosternidae). Acta Biologia Colombiana 25:431–433.

López-Luna, M.A., Venegas-Anaya, M., Cupul-Magaña, F.G., Rangel-Mendoza, J.A., and Escobedo-Galván, A.H. 2021. Mitochondrial DNA data support the recognition of the mud turtle, *Kinosternon vogti* (Cryptodira: Kinosternidae). Chelonian Conservation and Biology 20(1):97–102.

Lortet, L. 1883. Poissons et reptiles du lac de Tibériade et de quelques autres parties de la Syrie. Archives du Muséum d'Histoire Naturelle de Lyon 3:99–194.

Loulida, S., Naimi, M., Znari, M., and Bendami, S. 2019. Tolerance to salinity and dehydration in the Sahara Desert blue-eyed turtle, *Mauremys leprosa saharica* (Testudines: Geoemydidae) from a brackish pond in the Lower Draa basin, southern Morocco. African Journal of Herpetology 68:58–76.

Loulida, S., Znari, M., Naimi, M., and Bendami, S. 2021. Population structure and body size of the Sahara blue-eyed pond turtle *Mauremys leprosa saharica*, from an isolated pond in the Lower Draa Basin, southern Morocco. Environmental Science and Pollution Research, https://doi.org/10.1007/s11356-020-12331-5, 8 pp.

Lourenço, J.M., Claude, J., Galtier, N., and Chiari, Y. 2012. Dating cryptodiran nodes: origin and diversification of the turtle superfamily Testudinoidea. Molecular Phylogenetics and Evolution 62(1):496–507.

Loveridge, A. 1923. Notes on East African tortoises collected 1921–1923, with the description of a new species of soft land tortoise. Proceedings of the Zoological Society of London 1923:923–933.

Loveridge, A. 1934. Australian reptiles in the Museum of Comparative Zoology, Cambridge, Massachusetts. Bulletin of the Museum of Comparative Zoology 77:243–383.

Loveridge, A. 1935. Scientific results of an expedition to rain forest regions in eastern Africa. I. New reptiles and amphibians from East Africa. Bulletin of the Museum of Comparative Zoology 79:1–19.

Loveridge, A. and Williams, E.E. 1957. Revision of the African tortoises and turtles of the suborder Cryptodira. Bulletin of the Museum of Comparative Zoology 115(6):163–557.

Lovich, J.E. and Gibbons, J.W. 1992. A review of techniques for quantifying sexual size dimorphism. Growth, Development and Aging 56:269–281.

Lovich, J.E. and Hart, K.M. 2018. Taxonomy: a history of controversy and uncertainty. In: Roosenburg, W.M. and Kennedy, V.S. (Eds.). Ecology and

Conservation of the Diamond-Backed Terrapin. Baltimore: Johns Hopkins University Press, pp. 37–50.

Lovich, J.E. and McCoy, C.J. 1992. Review of the *Graptemys pulchra* group (Reptilia: Testudines: Emydidae), with descriptions of two new species. Annals of the Carnegie Museum 61(4):293–315.

Lovich, J.E., Ernst, C.H., and McBreen, J.F. 1990. Growth, maturity, and sexual dimorphism in the wood turtle, *Clemmys insculpta*. Canadian Journal of Zoology 68:672–677.

Lovich, J.E., Znari, M., Baamrane, M.A.A., Naimi, M., and Mostalih, A. 2010. Biphasic geographic variation in sexual size dimorphism of turtle (*Mauremys leprosa*) populations along an environmental gradient in Morocco. Chelonian Conservation and Biology 9(1):45–53.

Lovich, J.E., Gibbons, J.W., and Agha, M. 2014a. Does the timing of attainment of maturity influence sexual size dimorphism and adult sex ratio in turtles? Biological Journal of the Linnean Society 112:142–149.

Lovich, J.E., LaRue, C.T., Drost, C.A., and Arundel, T.R. 2014b. Traditional cultural use as a tool for inferring biogeography and provenance: a case study involving Painted Turtles (*Chrysemys picta*) and Hopi Native American culture in Arizona, USA. Copeia 2014:21–220.

Lubcke, G.M. and Wilson, D.S. 2007. Variation in shell morphology of the Western Pond Turtle (*Actinemys marmorata* Baird and Girard) from three aquatic habitats in northern California. Journal of Herpetology 41(1):107–114.

Luederwaldt, H. 1926. Os chelonios brasileiros. Revista Museo Paulista 14:403–470.

Luiselli, L., Demaya, G.S., Benansio, J.S., Lado, T.F., and Jubarah, S. 2022. Nubian flapshell turtle found in northern Uganda. Oryx 56(1):10.

Luján, A.H, Delfino, M., Robles, J.M., and Alba, D.M. 2016. The Miocene tortoise *Testudo catalaunica* Bataller, 1926, and a revised phylogeny of extinct species of genus *Testudo* (Testudines: Testudinidae). Zoological Journal of the Linnean Society 178:312–342.

Luo, B. and Zong, Y. 1988. A new species of *Cuora—Cuora aurocapitata*. Acta Herpetologica Sinica 3:13–15.

Ly, T., Hoang, H.D., and Stuart, B.L. 2011. Market turtle mystery solved in Vietnam. Biological Conservation 144:1767–1771.

Ly, T., Hoang, H.D., and Stuart, B.L. 2013. Occurrence of the Endangered Keeled Box Turtle, *Cuora mouhotii*, in southern Vietnam. Chelonian Conservation and Biology 12(1):184–187.

Lydekker, R. 1880. A sketch of the history of the fossil Vertebrata of India. Journal of the Asiatic Society of Bengal 49(2):8–40.

Lydekker, R. 1885. Indian Tertiary and post-Tertiary Vertebrata. Siwalik and Nerbada Chelonia. Memoirs of the Geological Survey of India, Palaeontologia Indica (10)3:155–208.

Lydekker, R. 1886. On a new emydine chelonian from the Pliocene of India. Quarterly Journal of the Geological Society of London 42:540–541.

Lydekker, R. 1889. Catalogue of the Fossil Reptilia and Amphibia in the British Museum. Part III. Chelonia. London: British Museum of Natural History, 239 pp.

Lyons, J.A., Natusch, D.J.D., and Shepherd, C.R. 2013. The harvest of freshwater turtles (Chelidae) from Papua, Indonesia, for the international pet trade. Oryx 47(2):298–302.

MacCulloch, R.D. and Secoy, D.M. 1983. Demography, growth, and food of western painted turtles, *Chrysemys picta bellii* (Gray), from southern Saskatchewan. Canadian Journal of Zoology 61:1499–1509.

Macip-Ríos, R., Arias Cisneros, M.d.L., Aguilar-Miguel, X.S., and Casas-Andreu, G. 2009. Population ecology and reproduction of the Mexican Mud Turtle (*Kinosternon integrum*) in Tonatico, Estado de México. Western North American Naturalist 69(4):501–510.

Macip-Ríos, R., Jones, M.T., Willey, L.W., Akre, T.S., González-Akre, E. and Díaz-Gamboa, L.F. 2018. Population structure and natural history of Creaser's Mud Turtle (*Kinosternon creaseri*) in Central Yucatán. Herpetological Conservation and Biology 13:366–372.

Madella-Auricchio, C.R., Auricchio, P., and Soares, E.S. 2017. Reptile species composition in the Middle Gurguéia and comparison with inventories in the eastern Parnaíba River basin, state of Piauí, Brazil. Papéis Avulsos de Zoologia 57:375–386.

Maffei, F., do Nascimento, B.T.M., Moya, G.M., and Donatelli, R.J. 2016. New distribution records of *Mesoclemmys vanderhaegei* (Testudines: Chelidae) from southeastern Brazil, including observations on reproduction. Journal of Threatened Taxa 8(11):9322–9326.

Mahé, J. and Sourdat, M. 1972. Sur l'extinction des Vertébrés subfossiles et l'aridification du climat dans le Sud-Ouest de Madagascar. Description des gisements. Datations absolues. Bulletin de la Société Géologique de France (7)14:295–309.

Maier, C.R. and Sonneman, T. 2018. *Chelodina burrungandjii* (Sandstone Snake-necked Turtle). Maximum size. Herpetological Review 49:521–522. [= *Chelodina walloyarrina*].

Malnate, E.V. 1971. A catalog of primary types in the herpetological collections of the Academy of Natural Sciences, Philadelphia (ANSP). Proceedings of the Academy of Natural Sciences, Philadelphia 123:345–375.

Manaças, S. 1956. Anfíbios e répteis das ilhas de São Tomé e do Príncipe e do ilhéu das Rolas. Conferência Internacional dos Africanistas Ocidentais, Lisboa, Vol. 4, pp. 179–192.

Mandimbihasina, A.R., Frasier, C.L., Hagenson, R.A., Robertson, B.a.D., Engberg, S.E., Lewis, R.E., Woolaver, L.G., Razafimahatratra, E., Rabetafika, L.L., and Louis, E.E., Jr. 2020. Conservation genetics of Madagascar's critically endangered Ploughshare Tortoise (*Astrochelys yniphora*). Conservation Genetics 21:109–121.

Mann, M.R. 2007. A taxonomic study of the morphological variation and intergradation of *Chrysemys picta* (Schneider) (Emydidae, Testudines) in West Virginia. M.S. Thesis, Marshall University, Huntington, West Virginia.

Maran, J. 1996. L'émyde lépreuse, *Mauremys leprosa* (Schweigger, 1812). CITS Bulletin 7:16–43.

Maran, J. 2006a. Beobachtungen an den kontinentalen Schildkröten Gabuns. Emys 13(1):5–26.

Maran, J. 2006b. Retour au Bresil. La Tortue 74:58–75.

Maran, J. 2006c. Observations on Gabonese chelonians. In: Artner, H., Farkas, B., and Loehr, V. (Eds.). Turtles: Proceedings: International Turtle and Tortoise Symposium Vienna 2002. Frankfurt: Edition Chimaira, pp. 351–373.

Marchand, K.A., Stulberg, A., Somers, C.M., and Poulin, R.G. 2015. *Chrysemys picta bellii* (Western Painted Turtle). Record carapace length. Herpetological Review 46:617.

Mardiastuti, A. 2008. Harvest sustainability of Asiatic softshell turtle *Amyda cartilaginea* in Indonesia. Report to the CITES Management Authority of Indonesia, 13 pp.

Marggraf, G. 1648. Historiae Rerum Naturalium Brasiliae, Libri Octo. In: Piso, W. Historia Naturalis Brasiliae. Antwerp: Joannes de Laet, 293 pp.

Márquez, R. 1990. Sea turtles of the world. An annotated and illustrated catalogue of sea turtle species known to date. FAO Fisheries Synopsis 11(125):1–81.

Marrone, F., Sacco, F., Arizza, V., and Arculeo, M. 2016. Amendment of the type locality of the endemic Sicilian pond turtle *Emys trinacris* Fritz et al. 2005, with some notes on the highest altitude reached by the species (Testudines, Emydidae). Acta Herpetologica 11(1):59–61.

Martin, B.T., Bernstein, N.P., Birkhead, R.D., Koukl, J.F., Mussmann, S.M., and Placyk, J.S. 2013. Sequence-based molecular phylogenetics and phylogeography of the American box turtles (*Terrapene* spp.) with support from DNA barcoding. Molecular Phylogenetics and Evolution 68(1):119–134.

Martin, B.T., Bernstein, N.P., Birkhead, R.D., Koukl, J.F., Mussmann, S.M., and Placyk, J.S., Jr. 2014. On the reclassification of the *Terrapene* (Testudines: Emydidae): a response to Fritz & Havaš. Zootaxa 3835:292–294.

Martin, B.T., Douglas, M.R., Chafin, T.K., Placyk, J.S., Birkhead, R.D., Phillips, C.A., and Douglas, M.E. 2020. Contrasting signatures of introgression in North American box turtles (*Terrapene* spp.) contact zones. Molecular Ecology 29:4186–4202.

Martin, B.T., Chafin, T.K., Douglas, M.R., Placyk, J.S., Birkhead, R.D., Phillips, C.A., and Douglas, M.E. 2021. The choices we make and the impacts they have: machine learning and species delimitation in North American box turtles (*Terrapene* spp.). Molecular Ecology Resources; doi:10.1111/1755-0998.13350, 17 pp.

Martin, T.E., Guillemin, M., Nivet-Mazerolles, V., Landsmann, C., Dubos, J., Eudeline, R., and Stroud, J.T. 2017. The herpetofauna of central Uzbekistan. Amphibian & Reptile Conservation 11(1):93–107 (e140).

Martínez del Mármol, G.M., Harris, D.J., Geniez, P., de Pous, P., and Salvi, D. 2019. Amphibians and Reptiles of Morocco. Frankfurt: Edition Chimaira, 478 pp.

Marx, H. 1958. Catalogue of type specimens of reptiles and amphibians in Chicago Natural History Museum. Fieldiana: Zoology 36:407–496.

Mashkaryan, V., Vamberger, M., Arakelyan, M., Hezaveh, N., Carretero, M.A., Corti, C., Harris, D.J., and Fritz, U. 2013. Gene flow among deeply

divergent mtDNA lineages of *Testudo graeca* (Linnaeus, 1758) in Transcaucasia. Amphibia-Reptilia 34:337–351.

Maslin, T.P. 1959. An annotated check list of the amphibians and reptiles of Colorado. University of Colorado Studies Series in Biology 6:1–98.

Mayer, F.J.C. 1849. System des Thier-Reiches oder Eintheilung der Thiere nach einem Principe. Verhandlungen des Naturhistorischen Vereins der Preussischen Rheinlande und Westphalens 6:169–210.

Mayer, R. 1992. Europäische Landschildkröten. Leben – Haltung – Zucht. Kempten: Agrar Verlag Allgäu, 127 pp.

Mazuch, T., Trailin, V., Fritz, U., and Vamberger, M. 2016. Senegal Flapshell Turtle (*Cyclanorbis senegalensis*) in Ethiopia (Testudines: Trionychidae). Amphibian and Reptile Conservation 10(2):1–5.

McAllister, C.T., Forstner, M.R.J., and Fuller, J.P. 2007. Second report of the southern painted turtle, *Chrysemys dorsalis* (Testudines: Emydidae), from Texas, with comments on its genetic relationship to other populations. Texas Journal of Science 59(2):155–160.

McClain, C.R., Balk, M.A., Benfield, M.C., Branch, T.A., Chen, C., Cosgrove, J., Dove, A.D.M., Gaskins, L.C., Helm, R.R., Hochberg, F.G., Lee, F.B., Marshall, A., Mcmurray, S.E., Schanche, C., Stone, S.N., and Thaler, A.D. 2015. Sizing ocean giants: patterns of intraspecific size variation in marine megafauna. PeerJ 3:e715; DOI 10.7717/peerj.715.

McClusky, E.M., Mockford, S.W., Sands, K., Herman, T.B, Johnson, G., and Gonser, R.A. 2016. Population genetic structure of Blanding's turtles (*Emydoidea blandingii*) in New York. Journal of Herpetology 50:70–76.

McCord, R.D. 2016. What is *Kinosternon arizonense*? Historical Biology 28:310–315.

McCord, W.P. 1998 ["1997"]. *Mauremys pritchardi*, a new batagurid turtle from Myanmar and Yunnan, China. Chelonian Conservation and Biology 2(4):555–562.

McCord, W.P. and Iverson, J.B. 1991. A new box turtle of the genus *Cuora* (Testudines: Emydidae) with taxonomic notes and a key to the species. Herpetologica 47(4):407–420.

McCord, W.P. and Iverson, J.B. 1992. A new species of *Ocadia* (Testudines: Bataguridae) from Hainan Island, China. Proceedings of the Biological Society of Washington 105(1):13–18.

McCord, W.P. and Iverson, J.B. 1994. A new species of *Ocadia* (Testudines: Batagurinae) from southwestern China. Proceedings of the Biological Society of Washington 107(1):52–59.

McCord, W.P. and Joseph-Ouni, M. 2007a. A new species of *Chelodina* (Testudines: Chelidae) from southwestern New Guinea (Papua, Indonesia). Reptilia (GB) (Barcelona) 52:47–52.

McCord, W.P. and Joseph-Ouni, M. 2007b. A new genus of Australian longneck turtle (Testudines: Chelidae) and a new species of *Macrochelodina* from the Kimberley region of Western Australia (Australia). Reptilia (GB) (Barcelona) 55:56–64.

McCord, W.P. and Philippen, H.-D. 1998. A new subspecies of box turtle, *Cuora amboinensis lineata*, from northern Myanmar (Burma), with remarks on the distribution and geographic variation of the species. Reptile Hobbyist 1998(March):51–58.

McCord, W.P. and Pritchard, P.C.H. 2003a ["2002"]. A review of the softshell turtles of the genus *Chitra*, with the description of new taxa from Myanmar and Indonesia (Java). Hamadryad 27(1)[2003]:11–56.

McCord, W.P. and Pritchard, P.C.H. 2003b. *Chitra chitra* Nutaphand, 1986 (Reptilia, Testudines): proposed precedence of the specific name over that of *Chitra selenkae* Jaekel, 1911. Bulletin of Zoological Nomenclature 60(3):208–210.

McCord, W.P. and Thomson, S.A. 2002. A new species of *Chelodina* (Testudines: Pleurodira: Chelidae) from northern Australia. Journal of Herpetology 36(2):255–267.

McCord, W.P., Iverson, J.B., and Boeadi. 1995. A new batagurid turtle from northern Sulawesi, Indonesia. Chelonian Conservation and Biology 1(4):311–316.

McCord, W.P., Iverson, J.B., Spinks, P.Q., and Shaffer, H.B. 2000. A new genus of geoemydid turtle from Asia. Hamadryad 25(2):86–90.

McCord, W.P., Joseph-Ouni, M., and Lamar, W.W. 2001. A taxonomic reevaluation of *Phrynops* (Testudines: Chelidae) with the description of two new genera and a new species of *Batrachemys*. Revista de Biologia Tropical 49(2):715–764.

McCord, W.P., Cann, J., and Joseph-Ouni, M. 2003. A taxonomic assessment of *Emydura* (Testudines: Chelidae) with descriptions of new subspecies from Queensland, Australia. Reptilia (GB) (Barcelona) 27:59–63.

McCord, W.P., Joseph-Ouni, M., and Tabaka, C. 2005. Chelonian illustrations #18: African hinge-back tortoises. Reptilia (GB) (Barcelona) 38:71–74.

McCord, W.P., Joseph-Ouni, M., and Hagen, C. 2007a. A new species of *Chelodina* (Testudines: Chelidae) from eastern Timor Island (East Timor). Reptilia (GB) (Barcelona) 52:53–57.

McCord, W.P., Joseph-Ouni, M., and Hagen, C. 2007b. A new subspecies of *Chelodina mccordi* (Testudines: Chelidae) from eastern Rote Island, Indonesia. Reptilia (GB) (Barcelona) 52:58–61.

McCord, W.P., Joseph-Ouni, M., Hagen, C., and Blanck, T. 2010. Three new subspecies of *Trachemys venusta* (Testudines: Emydidae) from Honduras, northern Yucatán (Mexico), and Pacific Coastal Panama. Reptilia (GB) (Barcelona) 71:39–49.

McCoy, C.J. and Vogt, R.C. 1985. *Pseudemys alabamensis*. Catalogue of American Amphibians and Reptiles 371:1–2.

McCoy, E.D., Aguirre, G., Kazmaier, R.T., and Tracy, C.R. 2014. Demography of North American tortoises. In: Rostal, D.C., McCoy, E.D., and Mushinsky, H.R. (Eds.). Biology and Conservation of North American Tortoises. Baltimore, MD: Johns Hopkins University Press, pp. 134–142.

McCranie, J.R. 2018. The lizards, crocodiles, and turtles of Honduras. Systematics, distribution, and conservation. Bulletin of the Museum of Comparative Zoology, Special Publications Series No. 2, 646 pp.

McCranie, J.R., Köhler, F., Gutsche, A., and Valdés Orellana, L. 2013. *Trachemys grayi emolli* (Testudines, Emydidae) in Honduras and its systematic relationships based on mitochondrial DNA. Zoosystematics and Evolution 89(1):21–29.

McCulloch, A.R. 1908. A new genus and species of turtle, from north Australia. Records of the Australian Museum 7:126–128.

McDowell, S.B. 1964. Partition of the genus *Clemmys* and related problems in the taxonomy of the aquatic Testudinidae. Proceedings of the Zoological Society of London 143:239–279.

McGaugh, S.E. 2008. Color variation among habitat types in the spiny softshell turtles (Trionychidae: *Apalone*) of Cuatrociénegas, Coahuila, Mexico. Journal of Herpetology 42(2):347–353.

McGaugh, S.E. 2012. Comparative population genetics of aquatic turtles in the desert. Conservation Genetics 13:1561–1576.

McGaugh, S.E. and Janzen, F.J. 2008. The status of *Apalone atra* populations in Cuatro Ciénegas, Coahuila, Mexico: preliminary data. Chelonian Conservation and Biology 7(1):88–95.

McGaugh, S.E., Eckerman, C.M., and Janzen, F.J. 2008. Molecular phylogeography of *Apalone spinifera* (Reptilia, Trionychidae). Zoologica Scripta 37(3):289–304.

McKnight, D.T., Georges, A., Guarino, F., and Bower, D.S. 2023. The ecology and morphology of Australia's desert turtle (*Emydura macquarii emmotti*). Austral Ecology 48:1657–1680.

Medem, F. 1958. Informe sobre reptiles Colombianos (II). El conocimiento actual sobre la distribucion geografica de las Testudinata en Colombia. Boletin del Museo de Ciencias Naturales 2–3:13–45.

Medem, F. 1969 ["1968"]. El desarrollo de la herpetología en Colombia. Revista de la Academia Colombiana de Ciencias Exactas, Físicas y Naturales 13(50):149–199.

Medem, F. 1977. Contribucion al conocimiento sobre la taxonomia, distribucion geografica y ecologia de la tortuga "bache" (*Chelydra serpentina acutirostris*). Caldasia 12(56):41–101.

Mehnert, E. 1890. Untersuchungen über die Entwicklung des Beckengürtels der *Emys lutaria taurica*. Morphologische Jahrbucher 16:537–571.

Meier, H. and Schaefer, L. 2003. Experiences and observations of *Chelus fimbriatus*, the maintenance in terrarium and successful reproduction. Radiata 12(2):3–20.

Meiri, S., Itescu, Y., Shacham, H., and Werner, Y.L. 2011. What tortoise is *Testudo floweri* from the Negev, named by Bodenheimer (1935)? Israel Journal of Ecology and Evolution 57:261.

Mendes-Pinto, T.J., de Souza, S.M., Vogt, R.C., and Bernhard, R. 2011. First record of *Platemys platycephala melanonota* Ernst, 1984 (Reptilia, Testudines, Chelidae) for the Brazilian Amazon. Revista de Ciências Ambientais, Canoas 5(2):103–107.

Merchán Fornelino, M. 2002. Estudio biométrico de juveniles de tortuga "jicotea" (*Trachemys scripta emolli*) en Costa Rica. Revista Española de Herpetologia 16:11–17.

Merchán Fornelino, M. 2003. Contribución al conocimiento de la biología de la tortuga negra (*Rhinoclemmys funerea*) y la tortuga roja (*R. pulcherrima manni*)

en Costa Rica. Ph.D. Thesis, Universidad Complutense de Madrid, Spain.
Merrem, B. 1820. Versuch eines Systems der Amphibien. Tentamen Systematis Amphibiorum. Marburg: J.C. Krieger, 191 pp.
Mertens, R. 1937. Bemerkungen über die Rassen von *Pelomedusa subrufa* (La Cepède). Zoologischer Anzeiger 117:139–142.
Mertens, R. 1946. Uber einige mediterrane Schildkröten-Rassen. Senckenbergiana 27:111–118.
Mertens, R. 1949. *Medaestia* Wussow, 1916, a synonym of *Testudo* Linnaeus, 1758. Copeia 1949(3):232.
Mertens, R. 1954. Zur Kenntnis der Schildkrötenfauna Venezuelas. Senckenbergiana Biologica 35(1/2):3–7.
Mertens, R. 1967a. Bemerkenswerte Süsswasserschildkröten aus Brasilien. Senckenbergiana Biologica 48:71–82.
Mertens, R. 1967b. Die herpetologische Sektion des Natur-Museums und Forschungs-Institutes Senckenberg in Frankfurt a.M. nebst einem Verzeichnis ihrer Typen. Senckenbergiana Biologica 48(Supplement A):1–106.
Mertens, R. 1969a. Eine neue Rasse der Dachschildkröte, *Kachuga tecta*. Senckenbergiana Biologica 50:23–30.
Mertens, R. 1969b. Eine neue Halswender-Schildkröte aus Peru. Senckenbergiana Biologica 50:132.
Mertens, R. 1970. Zur Kenntnis von *Phrynops nasutus*. Senckenbergiana Biologica 51:17–20.
Mertens, R. and Müller, L. 1928. Liste der Amphibien und Reptilian Europas. Abhandlungen Senckenbergische Naturforschende Gesellschaft 41:1–62.
Mertens, R. and Wermuth, H. 1955. Die rezenten Schildkröten, Krokodile und Brückenechsen. Eine kritische Liste der heute lebenden Arten und Rassen. Zoologische Jahrbücher 83:323–440.
Mertens, R. and Wermuth, H. 1961. Proposed use of the plenary powers to suppress eight specific names of turtles (Reptilia, Testudines). Bulletin of Zoological Nomenclature 18:211–213.
Métrailler, S. and Le Gratiet, G. 1996. Tortues continentales de Guyane française. Continental Turtles of French Guiana. Martigny, Switzerland: Pillet SA, 127 pp.
Meuschen, F.C. 1778. Museum Gronovianum sive Index Rerum Naturalium. Lugduni Batavorum: Th. Haak and J. Meerburg, 251 pp.
Meyer, A.B. 1874. Eine Mittheilung über die von mir auf Neu-Guinea und den Inseln Jobi, Mysore und Mafoor im Jahre 1873 gesammelten Amphibien. Monatsberichte der Akademie der Wissenschaften zu Berlin 39:128–140.
Meyer, A.B. 1887. Verzeichniss der von mir in den Jahren 1870-1873 im Ostindischen Archipel gesammelten Reptilien und Batrachier. Abhandlungen und Berichte des Königl. Zoologischen und Anthropologisch-Ethnographischen Museums zu Dresden 1887(2):1–16.
Meyer, F.A.A. 1790. Kurze Beschreibungen neuer Thiere. Ausgezogen aus dem Leipziger Naturhistorischen Magazin. Magazin für Thiergeschichte, Thieranatomie und Thierarzenkunde, Göttingen 1:80–83.
Meyer, H. von. 1835. Mittheilungen an Professor Bronn gerichtet. Neues Jahrbuch für Mineralogie, Geognosie, Geologie und Petrefaktenkunde, Stuttgart 1835:63–69.
Meylan, P.A. 1987. The phylogenetic relationships of soft-shelled turtles (Family Trionychidae). Bulletin of the American Museum of Natural History 186:1–101.
Meylan, P.A. 1996. Skeletal morphology and relationships of the early cretaceous side-necked turtle, *Araripemys barretoi* (Testudines: Pelomedusoides: Araripemydidae), from the Santana formation of Brazil. Journal of Vertebrate Paleontology 16(1):20–33.
Meylan, P.A. and Sterrer, W. 2000. *Hesperotestudo* (Testudines: Testudinidae) from the Pleistocene of Bermuda, with comments on the phylogenetic position of the genus. Zoological Journal of the Linnean Society 128:51–76.
Michael, D. and Lindermeyer, D.G. 2010. Reptiles of the NSW Murray Catchment: A Guide to Their Identification, Ecology and Conservation. Canberra: CSIRO Publishing.
Michahelles, C. 1829. Commentatio de speciebus aut rarioribus, aut novis cheloniorum Europam meridionalem inhabitantibus. Isis von Oken 22:1295–1300.
Michels, J. and Vargas-Ramírez, M. 2018. Red-headed Amazon River Turtles in Venezuela and Colombia: population separation and connection along the famous route of Alexander von Humboldt. Zoology 130:67–78.
Mifsud, D.A. and Stapleton, M.M. 2014. *Kinixys* Conservation Blueprint: A Comprehensive Assessment to Ensure the Future of the Genus. Herpetological Resource and Management Technical Publication 2014, 134 pp.
Mikan, J.C. 1820. Delectus Florae et Faunae Brasiliensis. Fasciculus Primus. Vindobonae: 6 pp., 6 pls.
Mikan, J.C. 1825. Delectus Florae et Faunae Brasiliensis. Fasciculus Quartus. Vindobonae: 6 pp., 6 pls.
Mikuliček, P., Jandzik, D., Fritz, U., Schneider, C., and Široký, P. 2013. AFLP analysis shows high incongruence between genetic differentiation and morphology-based taxonomy in a widely distributed tortoise. Biological Journal of the Linnean Society 108(1):151–160.
Miller, J.F. 1779. *Testudo sulcata*, pl. 26. In: Miller, J.F. 1776–1784. Icones Animalium et Plantarum. (Various subjects of Natural History, Wherein are Delineated Birds, Animals and Many Curious Plants). London, 10 pp., 60 pls.
Miller, J.F. and Shaw, G. 1796. Cimelia Physica. Figures of Rare or Curious Quadrupeds, Birds, &c. Together with Several of the Most Elegant Plants, with Descriptions. London: Benjamin and John White and John Sewell, 106 pp., 60 pls.
Miller, J.M., Quinzin, M.C., Edwards, D.L., Eaton, D.A.R., Jensen, E.L., Russello, M.A., Gibbs, J.P., Tapia, W., Rueda, D., and Caccone, A. 2018. Genome-wide assessment of diversity and divergence among extant Galapagos giant tortoise species. Journal of Heredity 109(6):611–619.
Milne-Edwards, A. 1868. Sur des découvertes zoologiques faites récemment à Madagascar par M. Alfred Grandidier. Comptes Rendus Hebdomadaires des Séances de l'Académie des Sciences, Paris 67:1165–1167.
Milne-Edwards, A. 1875. Nouveaux documents sur l'époque de la disparition de la Faune ancienne de l'île Rodrigue. Annales des Sciences Naturelles, Zoologie (6)2(4):1–20.
Milstead, W.W. 1956. Fossil turtles of Friesenhahn Cave, Texas, with the description of a new species of *Testudo*. Copeia 1956(3):162–171.
Milstead, W.W. 1967. Fossil box turtles (*Terrapene*) from central North America, and box turtles of eastern Mexico. Copeia 1967(1):168–179.
Milstead, W.W. 1969. Studies on the evolution of the box turtles (genus *Terrapene*). Bulletin of the Florida State Museum, Biological Sciences 14(1):1–113.
Mimila-Manzur, F.D., Ramírez-Bautista, A., Macip-Ríos, R., and Díaz-Marín, C.A. 2023. Reproductive ecology and natural history of *Kinosternon herrerai* (Testudines: Kinosternidae) at the center of its distribution. Phyllomedusa 22(2):171–179.
Minx, P. 1996. Phylogenetic relationships among the box turtles, genus *Terrapene*. Herpetologica 52(4):584–597.
Mital, A., Gunia, L., Tachung, B., and Das, A. 2020. The impressed tortoise (*Manouria impressa*) in India – extended range and natural history notes. Herpetological Bulletin 153:40–43.
Mitchell, J.C. 1994. The Reptiles of Virginia. Washington, DC: Smithsonian Institution Press, 352 pp.
Mitchell, J.C., Johnston, G.R., and Munscher, E.C. 2017. *Trachemys scripta scripta* (Yellow-bellied Slider). Maximum Size. Herpetological Review 48(3):624–625.
Mitrus, S. and Zemanek, M. 2000. Distribution and biology of *Emys orbicularis* (L) in Poland. Stapfia 69, zugleich Kataloge des OÖ. Landesmuseums, Neue Folge Nr. 149, pp. 107–118.
Mittermeier, R.A. and Wilson, R.A. 1974. Redescription of *Podocnemis erythrocephala* (Spix, 1824), an Amazonian pelomedusid turtle. Papéis Avulsos de Zoologia, São Paulo 28(8):147–162.
Mittermeier, R.A., Rhodin, A.G.J., Medem, F., Soini, P., Hoogmoed, M.S., and Carrillo de Espinoza, N. 1978. Distribution of the South American chelid turtle *Phrynops gibbus*, with observations on habitat and reproduction. Herpetologica 34:94–100.
Mittermeier, R.A., van Dijk, P.P., Rhodin, A.G.J., and Nash, S.D. 2015. Turtle hotspots: an analysis of the occurrence of tortoises and freshwater turtles in Biodiversity Hotspots, High-Biodiversity Wilderness Areas, and Turtle Priority Areas. Chelonian Conservation and Biology 14(1):2–10.
Mittleman, M.B. 1945. Type localities of two American turtles. Copeia 1945(3):171.
Mittleman, M.B. 1947. The allocation of *Testudo rugosa* Shaw. Herpetologica 3:173–176.
Młynarski, M. 1976. Handbuch der Palaeoherpetologie. Part 7. Testudines. Stuttgart, Germany: Fischer Verlag, 130 pp.
Modesto, S.P. and Anderson, J.S. 2004. The phylogenetic definition of Reptilia. Systematic Biology 53(5):815–821.
Mojsisovics, A. von. 1889. Zoogeographische Notizen über Süd-Ungarn aus den Jahren 1886–1888. III. Nachtrag zur "Fauna von Béllye und Darda". Mitteilungen des Naturwissenschaftlichen Vereines für Steiermark, Graz

25(1888)[1889]:233–269.

Moler, P.E. 2006a. *Apalone mutica calvata* – Gulf Coast smooth softshell turtle. In: Meylan, P.A. (Ed.). Biology and Conservation of Florida Turtles. Chelonian Research Monographs No. 3, pp. 169–172.

Moler, P.E. 2006b. *Apalone spinifera aspera* – Gulf Coast spiny softshell turtle. In: Meylan, P.A. (Ed.). Biology and Conservation of Florida Turtles. Chelonian Research Monographs No. 3, pp. 173–177.

Molfetti, E., Torres Vilaça, S., Georges, J.Y., Plot, V., Delcroix, E., Le Scao, R., Lavergne, A., Barrioz, S., Rodrigues dos Santos, F., and de Thoisy, B. 2013. Recent demographic history and present fine-scale structure in the Northwest Atlantic Leatherback (*Dermochelys coriacea*) turtle population. PLOS ONE 8:e58061.

Molina, F.B., Machado, F.A., and Zaher, H. 2012. Taxonomic validity of *Mesoclemmys heliostemma* (McCord, Joseph-Ouni and Lamar, 2001) (Testudines, Chelidae) inferred from morphological analysis. Zootaxa 3575:63–77.

Molina-Serrano, K.M., Aguirre-Toledo, H.D., Meza-Inostroza, P.M., Castro-Bastidas, H.A., Bucio-Pacheco, M., and Iverson, J.B. 2025. Southern range extension of *Kinosternon sonoriense* Le Conte, 1984 (Testudines: Kinosternidae). Zootaxa 5604(1):87–93.

Moll, D. 1994. The ecology of sea beach nesting in slider turtles (*Trachemys scripta venusta*) from Caribbean Costa Rica. Chelonian Conservation and Biology 1:107–116.

Moll, E.O. 1987. Survey of the freshwater turtles of India. Part II: The genus *Kachuga*. Journal of the Bombay Natural History Society 84:7–25.

Moll, E.O. 1989. *Geochelone elegans*, Indian star tortoise. In: Swingland, I.R. and Klemens, M.W. (Eds.). The Conservation Biology of Tortoises. Occasional Papers of the IUCN Species Survival Commission No. 5, pp. 113–114.

Moll, E.O. and Vijaya, J. 1986. Distributional records for some Indian turtles. Journal of the Bombay Natural History Society 83:57–62.

Moodie, K.B. and Van Devender, T.R. 1979. Extinction and extirpation in the herpetofauna of the southern high plains with emphasis on *Geochelone wilsoni* (Testudinidae). Herpetologica 35:198–206.

Mooser, O. 1972. A new species of Pleistocene fossil tortoise, genus *Gopherus*, from Aguascalientes, Aguascalientes, Mexico. Southwestern Naturalist 17(1):61–65.

Mooser, O. 1980. Pleistocene fossil turtles from Aguascalientes, state of Aguascalientes. Revista Mexicana de Ciencias Geológicas 4(1):63–66.

Moraru, V., Buhaciuc-Ionita, E., Măntoiu, D.Ș., Gavril, V.D., Popescu-Mirceni, R., and Strugariu, A. 2016. The spur-thighed tortoise (*Testudo graeca ibera*) in Romania: new locality records suggest a more optimistic situation. North-Western Journal of Zoology 12(2):396–400.

Moreira, G.M. 2002. Distribuição, status populacional e conservação do cágado *Phrynops hogei* (Mertens, 1967) (Testudinae: Chelidae) no rio Carangola. Masters Thesis, Universidade Federal de Minas Gerais, Brazil.

Moreno, N., Heaton, A., Bruening, K., Milligan, E., Nelson, D., Glaberman, S., and Chiari, Y. 2022. Hybridization and low genetic diversity in the endangered Alabama red-bellied turtle (*Pseudemys alabamensis*). Ecology and Evolution 12:e8964, 15 pp.

Morgan, G.S. 1993. Quaternary land vertebrates of Jamaica. Geological Society of America Memoirs 182:417–442.

Mosimann, J.E. and Rabb, G.B. 1953. A new subspecies of the turtle *Geoemyda rubida* (Cope) from western Mexico. Occasional Papers of the Museum of Zoology, University of Michigan 548:1–7.

Mount, R.H. 1975. The Reptiles and Amphibians of Alabama. Auburn, AL: Auburn University Agricultural Experiment Station, 347 pp.

Mrykalo, R., Juszli, G., Mezza, G., and Southerland, J. 2016. *Gopherus polyphemus* (Gopher Tortoise). Size. Herpetological Review 47(2):284–285.

Mu, C. and Li, P. 2023. Assessing the invasion risk of *Chelydra serpentina* in China under current and future climate change scenarios. Frontiers in Ecology and Evolution 11:1277058, 11 pp., doi: 10.3389/fevo.2023.1277058.

Mukherjee, D., Nixon, A.M.A., and Bhupathy, S. 2006. Observations on the morphometry of two subspecies of *Melanochelys trijuga* from the Western Ghats, Southwestern India. Turtle and Tortoise Newsletter 9:7–9.

Müller, L. 1935. Uber eine neue *Podocnemis*-Art (*Podocnemis vogli*) aus Venezuela nebst ergänzenden Bemerkungen über die systematischen Merkmale der ihr nächstverwandten Arten. Zoologischer Anzeiger 110(5/6):97–109.

Müller, L. 1936. Beiträge zur Kenntnis der Schildkrötenfauna von Mexiko. Zoologischer Anzeiger 113:97–114.

Müller, L. 1940. Über *Pseudemys callirostris* (Gray). Ibero–amerikanische Studien, Hamburg 13:108–125.

Müller, L. and Hellmich, W. 1936. Amphibien und Reptilien. I. Teil: Amphibia, Chelonia, Loricata. Wissenschaftliche Ergebnisse der Deutschen Gran Chaco Expedition. Stuttgart: Verlag Von Strecker und Schroder, pp. 96–108.

Munscher, E. and Franklin, C.J. 2019. *Pseudemys texana* (Texas River Cooter). Maximum size. Herpetological Review 50:128.

Munthe, H. 1895. Om fyndet af ett benredskap i Ancyluslera nära Norsholm i Östergötland. Öfversigt af Kongliga Vetenskaps-Akademiens Förhandlingar 52(3):151–177. [in Swedish]

Murphy, J.C. 2018. Arizona's Amphibians and Reptiles. A Natural History and Field Guide. Second Edition. Privately Printed: Book Services, 333 pp.

Murphy, R.W. 2014. Systematics of extant North American tortoises. In: Rostal, D.C., McCoy, E.D., and Mushinsky, H.R. (Eds.). Biology and Conservation of North American Tortoises. Baltimore, Maryland: Johns Hopkins University Press, pp. 25–29.

Murphy, R.W., Berry, K.H., Edwards, T., Leviton, A.E., Lathrop, A., and Riedle, J.D. 2011. The dazed and confused identity of Agassiz's land tortoise, *Gopherus agassizii* (Testudines, Testudinidae) with the description of a new species, and its consequences for conservation. ZooKeys 113:39–71.

Murray, C.M., McMahan, C.D., Dobie, J.L., and Guyer, C. 2014. Cranial variation amongst independent lineages of the alligator snapping turtle (*Macrochelys temminckii*). Journal of Zoological Systematics and Evolutionary Research 52: 305–311.

Murray, J.A. 1884a. The Vertebrate Zoology of Sind; with descriptions of all the known species of mammals, birds, and reptiles inhabiting the province; observations on their habits; tables of their geographical distribution in Persia, Beloochistan, and Afghanistan; Punjab, North-West provinces, and the peninsula of India generally. London: Richardson and Co., 424 pp.

Murray, J.A. 1884b. Additions to the reptilian fauna of Sind. Annals and Magazine of Natural History (5)14:106–111.

Murthy, H.C.K. and Das, I. 2009. The turtle collection of the Zoological Survey of India, Kolkata, India. In: Vasudevan, K. (Ed.). Freshwater Turtles and Tortoises of India. ENVIS Bulletin: Wildlife and Protected Areas 12(1):15–24.

Mushinsky, H.R., Mccoy, E.D., Berish, J.E., Ashton, R.E., Jr., and Wilson, D.S. 2006. *Gopherus polyphemus* – Gopher Tortoise. In: Meylan, P.A. (Ed.). Biology and Conservation of Florida Turtles. Chelonian Research Monographs 3:350–375.

Myers, E.M. 2008. Post-orbital color pattern variation and the evolution of a radiation of turtles (*Graptemys*). Ph.D. Thesis, Iowa State University, Ames.

Nagy, Z.T., Kielgast, J., Moosig, M., Vamberger, M., and Fritz, U. 2015. Another candidate species of *Pelomedusa* (Testudines: Pelomedusidae) from the Democratic Republic of the Congo? Salamandra 51:212–214.

Naish, D. 2025. Taxonomic trolling. The Biologist: Royal Society of Biology 72(2):26–29.

Nardo, G.D. 1864. Sopra una nuovo rarissima specie di cheloniano pescato alle nostre spiaggie. Atti del Reale Istituto Veneto di Scienze, Lettere ed Arti (3)9:1418–1423. [in Italian].

Naro-Maciel, E., Le, M., FitzSimmons, N.N., and Amato, G. 2008. Evolutionary relationships of marine turtles: a molecular phylogeny based on nuclear and mitochondrial genes. Molecular Phylogenetics and Evolution 49:659–662.

Naro-Maciel, E., Reid, B.N., Alter, S.E., Amato, G., Bjorndal, K.A., Bolten, A.B., Martin, M., Nairn, C.J., Shamblin, B., and Pineda-Catalan, O. 2014. From refugia to rookeries: phylogeography of Atlantic green turtles. Journal of Experimental Marine Biology and Ecology 461:306–316.

Nath, B. 1959. Animal remains of the 12th Century A.D. from Sarnath, Uttar Pradesh, India. Journal of the Zoological Society of India 10:165–175.

Neill, W.T. 1951. The taxonomy of North American soft-shelled turtles, genus *Amyda*. Publications of the Research Division of Ross Allen's Reptile Institute 1(2):7–24.

Neill, W.T. 1965. New and noteworthy amphibians and reptiles from British Honduras. Bulletin of the Florida State Museum, Biological Sciences 9:77–130.

Neill, W.T. and Allen, E.R. 1959. Studies on the amphibians and reptiles of British Honduras. Publications of the Research Division of Ross Allen's Reptile Institute 2(1):1–76.

Nekrasova, O., Yanish, Y., Tytar, V., and Pupins, M. 2019. GIS-modeling of the range shifts of the sub-fossil and extant European Pond Turtle (*Emys orbicularis*) in eastern Europe in Holocene. Diversity 11(8):121, doi:10.3390/d11080121.

Newsome, T.M., Wolf, C., Nimmo, D.G., Kopf, R.K., Ritchie, E.G., Smith, F.A., and Ripple, W.J. 2020. Constraints on vertebrate range size predict extinction

risk. Glbal Ecology and Biogeography 29:76–86.

Ngo, H.T., McCormack, T.E.M., Hoang, H.V., Nguyen, T.T., Tapley, B., Le, M.H., Le, D.T., Nguyen, T.T., Trinh, H.L.T., Ziegler, T., Nguyen, T.Q., and Le, M.D. 2023. Genetic diversity of the critically endangered Big-headed Turtle (*Platysternon megacephalum*) based on wild and traded samples: implications for conservation. Diversity 15:958, 13 pp.

Ngo, H.T., McCormack, T.E.M., Hoang, H.V., Nguyen, L.T., Nguyen, T.T., Nguyen, V.T.H., Do, H.T., Trinh, H.L.T., Ziegler, T., Blair, M.E., and Le, M.D. 2025. Introgressive hybridization and a natural barrier: molecular analyses lay out a conservation blueprint for the critically endangered Vietnamese Pond Turtle (*Mauremys annamensis*). Diversity 17:260, 19 pp.

Nguyen, L.T., Lam, N.Q., Carney, J., Hoang, H.V., McCormack, T.E.M., Nguyen, T.T., and Nguyen, S.N. 2020. First record of Western Black-Bridged Leaf Turtle, *Cyclemys atripons* Iverson & McCord, 1997 (Testudines, Geoemydidae), in Vietnam. Check List 16(3):571–577.

Nguyen, V.S., Ho, T.C., and Nguyen, Q.T. 2009. Herpetofauna of Vietnam. Frankfurt: Edition Chimaira.

Nikolsky, A.M. 1896. Diagnosis Reptilium et Amphibiorum novorum in Persia orientali a N. Zarudny Collectorum. Annuaire du Musée Zoologique de l'Académie Impériale des Sciences de St. Pétersbourg 4:369–372.

Nikolsky, A.M. 1897. Les reptiles, amphibiens et poissons recueillis par Mr. N. Zaroudny dans la Perse orientale. Annuaire du Musée Zoologique de l'Academie Imperiale des Sciences de St. Petersbourg 2:306–348.

Nikolsky, A.M. 1899. Herpetologica Turanica. Moscow: Friedlander, 84 pp.

Nikolsky, A.M. 1915. [Faune de la Russie et des Pays Limitrophes. Reptiles (Reptilia). Volume I. Chelonia et Sauria.] Petrograd: 532 pp. [in Russian].

Nilsson, S. 1841. Beskrifning öfver en i Skåne funnen fossil sköldpadda, jemförd med andra i Svensk jord funna qvarlefvor af samma djurordning. [Description of a fossil turtle found in Skåne, compared with other Swedish remains of the same order of animals]. Kongliga Svenska Vetenskaps-Academiens Handlingar 1839[1841]:194–211. [in Swedish].

Nori, J., Tessarolo, G., Ficetola, G.F., Loyola, R., Di Cola, V., and Leynaud, G. 2017. Buying environmental problems: the invasive potential of imported freshwater turtles in Argentina. Aquatic Conservation: Marine and Freshwater Ecosystems 27:685–691.

Nowak-Kemp, M. and Fritz, U. 2010. Chelonian type specimens at the Oxford University Museum. Zootaxa 2604:1–19.

Nutaphand, W. 1979. The Turtles of Thailand. Bangkok: Siamfarm Zoological Garden, 222 pp.

Nutaphand, W. 1986. [Manlai, the world's largest soft-shelled turtle]. Thai Zoological Magazine 1(4):64–70. [in Thai].

Nutaphand, W. 1990. [Softshelled turtles]. Thai Zoological Magazine 5(56):93–104. [in Thai].

Obst, F.J. 1976. Über den Holotypus von *Platemys novae guineae* Meyer, 1874 (Reptilia, Chelonia). Zoologische Abhandlungen, Staatliches Museum für Tierkunde Dresden 34(3):43–46.

Obst, F.J. 1983. Beitrag zur Kenntnis der Landschildkröten-Gattung *Manouria* Gray, 1852 (Reptilia, Testudines, Testudinata). Zoologische Abhandlungen, Staatliches Museum für Tierkunde Dresden 38(15):247–256.

Obst, F.J. and Reimann, M. 1994. Bemerkenswerte Variabilität bei *Cuora galbinifrons* Bourret, 1939, mit Beschreibung einer neuen geographischen Unterart: *Cuora galbinifrons bourreti* subsp. nov. (Reptilia: Testudines: Cryptodira: Bataguridae). Zoologische Abhandlungen, Staatliches Museum für Tierkunde Dresden 48(7):125–137.

Oelrich, T.M. 1953. A new boxturtle from the Pleistocene of southwestern Kansas. Copeia 1953(1):33–38.

Oelrich, T.M. 1957. The status of the Upper Pliocene turtle, *Testudo turgida* Cope. Journal of Paleontology 31:228–241.

Ogilby, J.D. 1890. Description of a new Australian tortoise. Records of the Australian Museum 1:56–59.

Ogilby, J.D. 1905. Catalogue of the Emydosaurian and Testudinian reptiles of New Guinea. Proceedings of the Royal Society of Queensland 19(1):1–31.

Oh, H.S., Park, S.M., and Han, S.H. 2017. Mitochondrial haplotype distribution and phylogenetic relationship of an endangered species Reeve's turtle (*Mauremys reevesii*) in East Asia. Journal of Asia-Pacific Biodiversity 10:27–31.

Okamoto, K. and Kamezaki, N. 2014. Morphological variation in *Chelonia mydas* (Linnaeus, 1758) from the coastal water of Japan, with special reference to the turtles allied to *Chelonia mydas agassizii* Bocourt, 1868. Current Herpetology 33:46–56.

Okayama, T., Díaz-Fernández, R., Baba, Y., Halim, M., Abe, O., Azeno, N., and Koike, H. 1999. Genetic diversity of the hawksbill turtle in the Indo-Pacific and Caribbean regions. Chelonian Conservation and Biology 3(2):362–367.

Olearius, A. 1674. Gottorffische Künst-Kammer: worinnen allerhand ungemeine Sachen, so theils die Natur, theils künstliche Hände hervorgebracht und bereitet, vor diesem aus allen vier Theilen der Welt zusammen getragen und vor einigen Jahren beschrieben, auch mit behörigen Kupffern gezieret. Schleswig: Gottfriedt Schultzen, 80 pp.

Oliveira, J.C.F., Castro, T.M., Silva-Soares, T., and Rocha, C.F.D. 2019. First-order effects of fire and prolonged-drought effects on an undescribed semi-aquatic turtle in Atlantic rainforest in southeastern Brazil. Journal of Coastal Conservation 23:367–372.

Olivier, G.A. 1807. Voyage dans l'Empire Othoman, l'Égypte et la Perse. Tome VI. Paris: H. Agasse, 522 pp.

Olson, S.L. 2015. More on the status of *Testudo nigra* Quoy and Gaimard and *Testudo nigrita* Duméril and Bibron as *nomina dubia* for Galapagos tortoises (Testudines: Testudinidae). Proceedings of the Biological Society of Washington 128(4):204–208.

Olson, S.L. 2017. The early scientific history of Galapagos tortoises. Archives of Natural History 44(2):241–258.

Olson, S.L. and David, N. 2014. The gender of the tortoise genus *Chelonoidis* Fitzinger, 1835 (Testudines: Testudinidae). Proceedings of the Biological Society of Washington 126:393–394.

Olson, S.L. and Humphrey, J.R. 2017. The island of origin of Richard Harlan's Galapagos tortoise *Testudo elephantopus*. Archives of Natural History 44:110–117.

Oppel, M. 1811. Die Ordnungen, Familien und Gattungen der Reptilien als Prodrom einer Naturgeschichte derselben. München: J. Lindauer, 86 pp.

Ortiz, J.C. and Nuñez, H. 1986. Catalogo critico de los tipos de reptiles conservados en el Museo Nacional de Historia Natural de Santiago, Chile. Publicación Ocasional Museo Nacional de Historia Natural 43:5–23.

Osbeck, P. 1757. Dagbok Öfwer en Ostindisk Resa Åren 1750, 1751, 1752. Med Anmärkningar uti Naturkunnigheten, Främmande Folkslags Språk, Seder, Hushållning, m.m. Stockholm: L.L. Grefing, 376 pp. [in Swedish]

Osborn, H.F. 1929. Biographical memoir of Edward Drinker Cope, 1840–1897. National Academy of Sciences, Biographical Memoirs 13(3):125–317.

Osculati, G. 1850. Esplorazione delle regioni equatoriali lungo il Napo ed il fiume delle Amazzoni, frammento di un viaggio fatto nelle due Americhe negli anni 1846-1847-1848. Milano: Tipografia Bernardoni, 320 pp.

Ottley, J.R. and Velázques Solis, V.M. 1989. An extant, indigenous tortoise population in Baja California Sur, Mexico, with the description of a new species of *Xerobates* (Testudines: Testudinidae). Great Basin Naturalist 49:496–502.

Otsuka, H. and Takahashi, A. 2000. Pleistocene vertebrate faunas in the Ryukyu Islands: their migration and extinction. Tropics 10:25–40.

Otsuka, H., Nakamura, T., and Ota, T. 2008. ^{14}C ages of vertebrate fossil beds in the Ryukyu Islands, south Japan. Summaries of Researches Using AMS at Nagoya University 19:135–153.

Ouwens, P.A. 1914. List of Dutch East Indian chelonians in the Buitenzorg Zoological Museum. Contributions à la Faune des Indes Néerlandaises, Buitenzorg 1:29–32.

Owen, J., Perry, G., Lazell, J., and Petrovic, C. 2005. *Pseudemys nelsoni* (Florida red-bellied turtle). Distribution. Herpetological Review 36:466.

Owen, R. 1853. Descriptive Catalogue of the Osteological Series Contained in the Museum of the Royal College of Surgeons of England. Vol I. Pisces, Reptilia, Aves, Marsupialia. London: Taylor and Francis, 350 pp.

Owen, R. 1886. Description of fossil remains of two specimens of a Megalanian genus (*Meiolania*, Ow.), from Lord Howe's Island. Proceedings of the Royal Society of London 1886(40):315–316.

Páez, V.P., Morales-Betancourt, M.A., Lasso, C.A., Castaño-Mora, O.V., and Bock, B.C. (Eds.). 2012. Biología y Conservación de las Tortugas Continentales de Colombia. Bogotá, Colombia: Serie Editorial Recursos, Hidrobiológicos y Pesqueros Continentales de Colombia. Instituto de Investigación de Recursos Biológicos Alexander von Humboldt, 528 pp.

Palacios, C., Urrutia, C., Knapp, N., Quintana, M.F., Bertolero, A., Simon, G., Du Preez, L., and Verneau, O. 2015. Demographic structure and genetic diversity of *Mauremys leprosa* in its northern range reveal new populations and mixed origins. Salamandra 51:221–230.

Palkovacs, E.P., Gerlach, J., and Caccone, A. 2002. The evolutionary origin of Indian Ocean tortoises (*Dipsochelys*). Molecular Phylogenetics and Evolution

24:216–227.

Palkovacs, E.P., Marschner, M., Ciofi, C., Gerlach, J., and Caccone, A. 2003. Are the native giant tortoises from the Seychelles really extinct? A genetic perspective based on mtDNA and microsatellite data. Molecular Ecology 12:1403–1413.

Pallas, P.S. 1814. Zoographia Rosso-Asiatica. III. Animalia Monocardia seu Frigidi Sanguinis Imperii Rosso-Asiatici. Petropolis: Officina Caes. Academiae Scientiarum, 428 pp.

Palmer, W.M. and Braswell, A.L. 1995. Reptiles of North Carolina. Chapel Hill, NC: University of North Carolina Press, 423 pp.

Palupcíková, K., Somerová, B., Protiva, T., Rehák, I., Velensky, P., Hulva, P., Gunalen, D., and Frynta, D. 2012. Genetic and shell-shape analyses of *Orlitia borneensis* (Testudines: Geoemydidae) reveal limited divergence among founders of the European zoo population. Zootaxa 3280:56–66.

Paolillo O., A. 1985. Description of a new subspecies of the turtle *Rhinoclemmys punctularia* (Daudin) (Testudines: Emydidae) from southern Venezuela. Amphibia-Reptilia 6(3):293–305.

Parenzan, P. 1932. Revisione delle specie del Gen. *Testudo* della Balcania. Atti del Reale Istituto Veneto di Scienze, Lettere ed Arti 91(11):1149–1169.

Parham, J.F. 2008. Rediscovery of an "extinct" Galapagos tortoise. Proceedings of the National Academy of Sciences (PNAS) 105:15227–15228.

Parham, J.F. and Fastovsky, D.E. 1998 ["1997"]. The phylogeny of cheloniid sea turtles revisited. Chelonian Conservation and Biology 2(4):548–554.

Parham, J.F. and Feldman, C.R. 2002. Generic revisions of emydine turtles. Turtle and Tortoise Newsletter 6:28–30.

Parham, J.F. and Li, D. 1999. A new locality for *Cuora pani* Song 1984 with comments on its known range. Asiatic Herpetological Research 8:111–113.

Parham, J.F. and Pyenson, N.D. 2010. New sea turtle from the Miocene of Peru and the iterative evolution of feeding ecomorphologies since the Cretaceous. Journal of Paleontology 84(2):231–247.

Parham, J.F. and Zug, G.R. 1996. *Chelonia agassizii* – valid or not? Marine Turtle Newsletter 72:2–5.

Parham, J.F., Simison, W.B., Kozak, K.H., Feldman, C.R., and Shi, H. 2001. New Chinese turtles: endangered or invalid? A reassessment of two species using mitochondrial DNA, allozyme electrophoresis and known-locality specimens. Animal Conservation 4:357–367.

Parham, J.F., Stuart, B.L., Bour, R., and Fritz, U. 2004. Evolutionary distinctiveness of the extinct Yunnan box turtle revealed by DNA from an old museum specimen. Proceedings of the Royal Society Series B: Biology Letters 271(1556[S6]):391–394.

Parham, J.F., Feldman, C.R., and Boore, J.L. 2006a. The complete mitochondrial genome of the enigmatic bigheaded turtle (*Platysternon*): description of unusual genomic features and the reconciliation of phylogenetic hypotheses based on mitochondrial and nuclear DNA. BMC Evolutionary Biology 6(11):1–11.

Parham, J.F., Macey, J.R., Papenfuss, T.J., Feldman, C.R., Türkozan, O., Polymeni, R., and Boore, J.L. 2006b. The phylogeny of Mediterranean tortoises and their close relatives based on complete mitochondrial genome sequences from museum specimens. Molecular Phylogenetics and Evolution 38:50–64.

Parham, J.F., Türkozan, O., Stuart, B.L., Arakelyan, M., Shafei, S., Macey, J.R., and Papenfuss, T.J. 2006c. Genetic evidence for premature taxonomic inflation in Middle Eastern tortoises. Proceedings of the California Academy of Sciences 57(3):955–963.

Parham, J.F., Outerbridge, M.E., Stuart, B.L., Wingate, D.B., Erlenkeuser, H., and Papenfuss, T.J. 2008. Introduced delicacy or native species? A natural origin of Bermudian terrapins supported by fossil and genetic data. Biology Letters 4:216–219.

Parham, J.F., Stuart, B.L., Danilov, I.G., and Ananjeva, N.B. 2012. A genetic characterization of CITES-listed Iranian tortoises (*Testudo graeca*) through the sequencing of topotypic samples and a 19th century holotype. Herpetological Journal 22:73–78.

Parham, J.F., Papenfuss, T.J., van Dijk, P.P., Wilson, B.S., Marte, C., Schettino, L.R., and Simison, W.B. 2013. Genetic introgression and hybridization in Antillean freshwater turtles (*Trachemys*) revealed by coalescent analyses of mitochondrial and cloned nuclear markers. Molecular Phylogenetics and Evolution 67:176–187.

Parham, J.F., Papenfuss, T.J., Buskirk, J.R., Parra-Olea, G., Chen, J.Y., and Simison, W.B. 2015. *Trachemys ornata* or not *ornata*: reassessment of a taxonomic revision for Mexican *Trachemys*. Proceeding of the California Academy of Science 4:359–367.

Parham, J.F., Papenfuss, T.J., Sellas, A.B., Stuart, B.L., and Simison, W.B. 2020. Genetic variation and admixture of red-eared sliders (*Trachemys scripta elegans*) in the USA. Molecular Phylogenetics and Evolution 145:106722, 9 pp.

Pavlov, P.A. 1932. Materials for the study of fauna of northern China, Manchuria and Mongolia. Reptilia and Amphibia. Part 1. Chelonia. Publications du Musée Hoang ho Pai ho de Tien Tsin 13:1–37.

Pavlov, P.A. 1933. Reptilia and Amphibia collected in 1932 by the staff of the Hoang ho Pai ho Museum. Publications du Musée Hoang ho Pai ho de Tien Tsin 23:1–12.

Pawar, S.S. and Choudhury, B.C. 2000. An inventory of chelonians from Mizoram, north-east India: new records and some observations on threats. Hamadryad 25(2):144–158.

Pedall, I., Fritz, U., Stuckas, H., Valdeón, A., and Wink, M. 2011. Gene flow across secondary contact zones of the *Emys orbicularis* complex in the Western Mediterranean and evidence for extinction and re-introduction of pond turtles on Corsica and Sardinia (Testudines: Emydidae). Journal of Zoological Systematics and Evolutionary Research 49(1):44–57.

Pedrono, M. 2008. The Tortoises and Turtles of Madagascar. Kota Kinabalu: Natural History Publications (Borneo), 147 pp.

Pedrono, M. and Clausen, A. 2018. Twilight of the Angonoka: Biology and Conservation of the World's Rarest Tortoise. Borneo: Natural History Publications.

Pedrono, M. and Markwell, T. 2001. Maximum size and mass of the ploughshare tortoise, *Geochelone yniphora*. Chelonian Conservation and Biology 4(1):190.

Pedrono, M. and Smith, L.L. 2013. Overview of the natural history of Madagascar's endemic tortoises and freshwater turtles: essential components for effective conservation. In: Castellano, C.M., Rhodin, A.G.J., Ogle, M., Mittermeier, R.A., Randriamahazo, H., Hudson, R., and Lewis, R.E. (Eds.). Turtles on the Brink in Madagascar: Proceedings of Two Workshops on the Status, Conservation, and Biology of Malagasy Tortoises and Freshwater Turtles. Chelonian Research Monographs 6:59–66.

Pennant, T. 1771. An account of two new tortoises; in a letter to Matthew Maty, M.D. Philosophical Transactions of the Royal Society of London 61:266–273.

Pennant, T. 1801. [*Testudo tuberculata*]. In: Schoepff, J.D. 1801. Historia Testudinum Iconibus Illustrata. Erlangae: Ioannis Iacobi Palm, 136 pp. [p. 123].

Perälä, J. 1996. Etelä-Turkin maakilpikonnista: morfologisia ja ekologisia eroja (*Testudo ibera* Pallas 1814 & *Testudo ibera anamurensis* Weissinger 1987) sekä uuden maakilpikonnalajin kuvaus. [in Finnish]. Appendix: Tortoises in southern Turkey: morphological and ecological differences (*Testudo ibera* Pallas 1814 & *Testudo ibera anamurensis* Weissinger 1987) and the description of a new tortoise species. In: Perälä, J., Vikberg, J., and Kanza, M. (Eds.). Herpetokongressi I – The Official Congress Publication, Herpetological Society of Finland, pp. 14–29; Appendix, pp. 1–21.

Perälä, J. 2001. A new species of *Testudo* (Testudines: Testudinidae) from the Middle East, with implications for conservation. Journal of Herpetology 35(4):567–582.

Perälä, J. 2002a. The genus *Testudo* (Testudines: Testudinidae): phylogenetic inferences. Chelonii 3:32–39.

Perälä, J. 2002b. Biodiversity in relatively neglected taxa of *Testudo* L., 1758 S. L. Chelonii 3:40–53.

Perälä, J. 2002c. Morphological variation among Middle Eastern *Testudo graeca* L., 1758 (sensu lato), with a focus on taxonomy. Chelonii 3:78–108.

Perälä, J. 2006. Assessment of the threatened status of *Testudo werneri* Perälä, 2001 (Testudines: Testudinidae) for the IUCN Red List. Chelonian Conservation and Biology 5(1):57–66.

Perälä, J. and Bour, R. 2004. Neotype of *Testudo terrestris* Forsskål, 1775 (Testudines, Testudinidae). Asiatic Herpetological Research 10:114–119.

Peregontsev, E. 2002. Horsfield tortoise (*Testudo (Agrionemys) horsfieldi* Gray, 1844) in Uzbekistan - its current status, conservation and sustainable use of the species. Proceedings of the International Congress on the Genus Testudo. Chelonii 3:293–299.

Pereira, A.G., Sterli, J., Moreira, F.R.R., and Schrago, C.G. 2017. Multilocus phylogeny and statistical biogeography clarify the evolutionary history of major lineages of turtles. Molecular Phylogenetics and Evolution 113:59–66.

Pereira, L.N., Santos, D.L., Vasconcelos, T.S., and Oda, F.H. 2013. Filling gaps on the distribution of *Rhinoclemmys punctularia* (Daudin, 1801) (Testudines: Geoemydidae) in the state of Maranhão, Brazil. Check List 9(1):146–147.

Perez, M., Leblois, R., Livoreil, B., Bour, R., Lambourdiere, J., Samadi, S., and Boisselier, M.-C. 2012. Effects of landscape features and demographic history on the genetic structure of *Testudo marginata* populations in the

southern Peloponnese and Sardinia. Biological Journal of the Linnean Society 105(3):591–606.

Perez, M., Livoreil, B., Mantovani, S., Boisselier, M.-C., Crestanello, B., Abdelkrim, J., Bonillo, C., Goutner, V., Lambourdière, J., Pierpaoli, M., Sterijovski, B., Tomovic, L., Vilaça, S.T., Mazzotti, S., and Bertorelle, G. 2014. Genetic variation and population structure in the Endangered Hermann's Tortoise: the roles of geography and human-mediated processes. Journal of Heredity 105(1):70–81.

Perrault, C. 1676. Description anatomique d'une grande Tortuë des Indes. Suites des Mémoires pour servir à l'Histoire Naturelle des Animaux 2:192–205.

Perry, G. 1810. Arcana; or the Museum of Natural History: Containing the Most Recent Discovered Objects. London: James Stratford, unpaginated text, plate 33 [*Testudo panama*]. [Complete work with 84 plates with unnumbered text issued in several parts: plates 1–48 in 1810, 49–84 in 1811].

Peters, J.A. 1952. Catalogue of type specimens in the herpetological collections of the University of Michigan Museum of Zoology. Occasional Papers of the Museum of Zoology University of Michigan 529:1–55.

Peters, W.K.H. 1848. Ueber eigenthümliche Moschusdrüsen bei Schildkröten. Archiv für Anatomie, Physiologie und Wissenschaftliche Medicin 1848:492–496.

Peters, W.K.H. 1854. Übersicht der auf seiner Reise nach Mossambique beobachteten Schildkröten. Bericht über die Bekanntmachung geeigneten Verhandlungen der Königlich-Preussischen Akademie der Wissenschaften zu Berlin 1854:215–216.

Peters, W.K.H. 1862. Über einen neuen *Phyllodactylus* aus Guayaquil. Monatsberichte der Königlichen Akademie der Wissenschaften zu Berlin 1862:626–627.

Peters, W.K.H. 1864. Eine neue Art der Baumvipern, *Atheris polylepis*, aus Liberia. Monatsberichte der Königlichen Akademie der Wissenschaften zu Berlin 1864:642–645.

Peters, W.K.H. 1866. Eine vorläufige Übersicht der aus dem Nachlass des Baron Carl von der Decken stammenden und auf seiner ostafrikanischen Reise gesammelten Säugethiere und Amphibien. Monatsberichte der Königlichen Akademie der Wissenschaften zu Berlin 1866:884–894.

Peters, W.K.H. 1868. Über eine neue Nagergattung, *Chiropodomys pencillatus*, so wie über neue oder weniger bekannte Amphibien und Fische. Monatsberichte der Königlichen Akademie der Wissenschaften zu Berlin 1868:448–453.

Peters, W.K.H. 1870. *Platemys tuberosa*, eine neue Art von Schildkröten aus British-Guiana. Monatsberichte der Königlichen Akademie der Wissenschaften zu Berlin 1870:311–313.

Peters, W.K.H. 1873. Über eine neue Schildkrötenart, *Cinosternon effeldtii* und einige andere neue oder weniger bekannte Amphibien. Monatsberichte der Königlichen Akademie der Wissenschaften zu Berlin 1873:603–618.

Peters, W.K.H. 1874 ["1875"]. Über neue Amphibien (*Gymnopis, Siphonops, Polypedates, Rhacophorus, Hyla, Cyclodus, Euprepes, Clemmys*). Monatsberichte der Königlichen Akademie der Wissenschaften zu Berlin 1874:616–624.

Petit, F.P. de. 1737. Description anatomique des yeux de la grenouille et de la tortue. Memoires de l'Academie Royale des Sciences, Paris 1737:199–237.

Petrozzi, F., Hema, E.M., Ségniagbeto, G.H., Amadi, N., Akani, G.C., Burke, R.L., Chirio, L., and Luiselli, L. 2019. Correlates of African Spurred Tortoise, *Centrochelys sulcata*, occurrence in the West African Sahel. Chelonian Conservation and Biology 18(1):19–23.

Petzold, A., Vargas-Ramírez, M., Kehlmaier, C., Vamberger, M., Branch, W.R., Du Preez, L., Hofmeyr, M.D., Meyer, L., Schleicher, A., Široký, P., and Fritz, U. 2014. A revision of African helmeted terrapins (Testudines: Pelomedusidae: *Pelomedusa*), with descriptions of six new species. Zootaxa 3795:523–548.

Pham, V.T., Vu, T.T., Dawson, J.E., Bui, T.T., and Leprince, B. 2018. Natural history observations on the endangered turtle *Geoemyda spengleri* in Tay Yen Tu Nature Reserve (Vietnam), with notes on other sympatric species. Herpetological Bulletin 146:1–7.

Philippen, H.-D. and Grossmann, P. 1990. Eine neue Schlangenhalsschildkröte von Neuguinea: *Chelodina reimanni* sp. n. (Reptilia, Testudines, Pleurodira: Chelidae). Zoologische Abhandlungen, Staatliches Museum Tierkunde Dresden 46(5):95–102.

Philippi, R.A. 1887. Vorlaufige Nachricht über die chilenischen Seeschildkröten und einige Fische der chilenischen Küste. Zoologischer Garten 28:84–88.

Philippi, R.A. 1899. Las tortugas chilenas. Anales de Universidad de Chile 104:727–736.

Phillips, C.A., Dimmick, W.W., and Carr, J.L. 1996. Conservation genetics of the common snapping turtle (*Chelydra serpentina*). Conservation Biology 10:397–405.

Phillips, K.F., Martin, K.R., Stahelin, G.D., Savage, A.E., and Mansfield, K.L. 2022. Genetic variation among sea turtle life stages and species suggests connectivity among ocean basins. Ecology and Evolution 12:e9426, 26 pp.

Pictet, F.J. 1853. Traité de Paléontologie, ou Histoire Naturelle des Animaux Fossiles dans leurs rapports Zoologiques et Géologiques. Seconde Édition. Tome Premier. Paris: J.-B. Baillière, 584 pp.

Pieh, A. 2001. *Testudo graeca soussensis*, eine neue Unterart der Maurischen Landschildkröte aus dem Sousstal (Südwest-Marokko). Salamandra 36(4):209–222.

Pieh, A. and Perälä, J. 2002. Variabilität von *Testudo graeca* Linnaeus, 1758 im östlichen Nordafrika mit Beschreibung eines neuen Taxons von der Cyrenaika (Nordostlibyen). Herpetozoa 15(1/2):3–28.

Pieh, A. and Perälä, J. 2004. Variabilität der Maurischen Landschildkröten (*Testudo graeca* Linnaeus, 1758 – Komplex) im zentralen und norwestlichen Marokko mit Beschreibung zweier neuer Taxa (Testudines: Testudinidae). Herpetozoa 17(1/2):19–47.

Pieh, A., Fritz, U., and Berglas, R. 2002. New data on morphology, distribution and nomenclature of *Testudo graeca armeniaca* Chkhikvadze & Bakradze, 1991 (Reptilia: Testudines: Testudinidae). Faunistische Abhandlungen Staatliches Museum für Tierkunde Dresden 22(21):329–345.

Pierlioni, A. 2019. Raksasa, the Indonesian giant – a new record size Spiny Turtle (*Heosemys spinosa*). Radiata 28(1):66–67.

Ping, C. 1930. Notes on the shell of a land tortoise from the ancient ruins of Annyang. Bulletin of the Fan Memorial Institute of Biology 1(13):217–226.

Pipatsawasdikul, K., Voris, H.K., and Thirakhupt, K. 2010. Distribution of the Big-Headed Turtle (*Platysternon megacephalum*, Gray 1831) in Thailand. Zoological Studies 49(5):640–650.

Piso, W. 1658. Historiae Naturalis et Medicae Indiae Occidentalis. Libri Quinque. In: Piso, W. De Indiae Utriusque re Naturali et Medica. Libri Quatuordecim. Amstelaedami (Amsterdam): Ludovicum et Danielem Elzevirios, 327 pp.

Platt, S.G., Win Ko Ko, Lay Lay Khaing, Khin Myo Myo, Tint Lwin, Thanda Swe, Kalyar, and Rainwater, T.R. 2005. Noteworthy records and exploitation of chelonians from the Ayeyarwady, Chindwin, and Dokhtawady rivers, Myanmar. Chelonian Conservation and Biology 4:942–948.

Platt, S.G., Heng, S., Long, K., Holloway, R., Stuart, B.L., and Rainwater, T.R. 2008. Biodiversity, exploitation, and conservation of turtles in the Tonle Sap Biosphere Reserve, Cambodia, with notes on reproductive ecology of *Malayemys subtrijuga*. Chelonian Conservation and Biology 7(2):195–204.

Platt, S.G., Khin Myo Myo, Win Ko Ko, Aung Maung, and Rainwater, T.R. 2010. Field observations and conservation of *Heosemys depressa* in the Rakhine Yoma Elephant Range of western Myanmar. Chelonian Conservation and Biology 9(1):114–119.

Platt, S.G., Platt, K., Khin Myo Myo, Kyaw Moe, Me Me Soe, That Zaw Naing, Naing Lin, and Rainwater, T.R. 2013. Noteworthy records of chelonians from the Chindwin River Basin and Naga Hills of western Myanmar. Herpetological Conservation and Biology 8(2):335−350.

Platt, S.G., Zug, G.R., Platt, K., Win Ko Ko, Khin Myo Myo, Me Me Soe, Tint Lwin, Myo Min Win, Swann Htet Naing Aung, Nay Win Kyaw, Htun Thu, Kyaw Thu Zaw Wint, van Dijk, P.P., Horne, B.D., and Rainwater, T.R. 2018. Field records of turtles, snakes and lizards in Myanmar (2009–2017) with natural history observations and notes on folk herpetological knowledge. Natural History Bulletin of the Siam Society 63:67–114.

Platt, S.G., Tint Lwin, Platt, K., Elsey, R.M., and Rainwater, T.R. 2019a. *Geochelone platynota* (Burmese Star Tortoise). Maximum body size and giantism. Herpetological Review 50(1):121–123.

Platt, S.G., Platt, K., Win Ko Ko, Khin Myo Myo, Me Me Soe, and Rainwater, T.R. 2019b. *Batagur baska* (Northern River Terrapin). Body size, growth, and conservation status. Herpetological Review 50(2):309–310.

Platt, S.G., Tint Lwin, Myo Min Win, Platt, K., Reh, B., Haislip, N.A., and Rainwater, T.R. 2019c. *Batagur trivittata* (Burmese Roofed Turtle). Sexual size dimorphism. Herpetological Review 50(3):553–555.

Platt, S.G., Haislip, N.A., and Rainwater, T.R. 2024 ("2023"). *Kinosternon hirtipes* (Rough-footed Mud Turtle). Demography, sex ratio, and sexual size dimorphism. Herpetological Review 54(3):448–450.

Plieninger, T. 1847. Verzeichnis der Reptilien Württembergs. Jahreshefte des Vereins für Vaterländische Naturkunde, Württemberg 3:194–208.

Pope, C.H. 1934. A new emydid turtle of the genus *Geoclemys* from Kwangtung Province, China. American Museum Novitates 691:1–2.

Pope, C.H. 1935. The Reptiles of China. New York: American Museum of Natural

History, Natural History of Central Asia, Vol. X, 604 pp.

Pope, C.H. 1939. Turtles of the United States and Canada. New York: Knopf, 343 pp.

Porter, D. 1815. Journal of a Cruise made to the Pacific Ocean by Captain David Porter, in the United States Frigate *Essex*, in the years 1812, 1813, 1814. Philadelphia: Bradford and Inskeep, 2 volumes.

Portis, A. 1890. I Rettili Pliocenici del Valdarno Superiore e di Alcune Altre Localitá Plioceniche di Toscana. Firenze: Le Monnier, 32 pp.

Pöschel, J., Heltai, B., Graciá, E., Quintana, M.F., Velon-Antón, G.V., Arribas, O., Valdeón, A., Wink, M., Fritz, U., and Vamberger, M. 2018. Complex hybridization patterns in European Pond Turtles (*Emys orbicularis*) in the Pyrenean Region. Science Reports 8:15925.

Poterala, S.F., Rattay, P., Johnson, A.S., Buijlayyil, M.S.A., Kiraz, A., and Ajaj, A.M.S. 2024. *Testudo graeca tripolitania*, a new taxon of spur-thighed tortoise found in Libya at risk of exploitation for the international pet trade. Basic and Applied Herpetology 38:111–125.

Pough, F.H. and Pough, M.B. 1968. The systematic status of painted turtles (*Chrysemys*) in the northeastern United States. Copeia 1968(3):612–618.

Poulakakis, N., Glaberman, S., Russello, M., Beheregaray, L.B., Ciofi, C., Powell, J.R., and Caccone, A. 2008. Historical DNA analysis reveals living descendants of an extinct species of Galápagos tortoise. Proceedings of the National Academy of Sciences (PNAS) 105:15464–15469.

Poulakakis, N., Russello, M., Geist, D., and Caccone, A. 2012. Unravelling the peculiarities of island life: vicariance, dispersal, and the diversification of the extinct and extant giant Galápagos tortoises. Molecular Ecology 21:160–173.

Poulakakis, N., Edwards, D.L., Chiari, Y., Garrick, R.C., Russello, M.A., Benavides, E., Watkins-Colwell, G.J., Glaberman, S., Tapia, W., Gibbs, J.P., Cayot, L.J., and Caccone, A. 2015. Description of a new Galapagos giant tortoise species (*Chelonoidis*; Testudines: Testudinidae) from Cerro Fatal on Santa Cruz Island. PLoS ONE 10:e0138779; 18 pp.

Poulakakis, N., Miller, J.M., Jensen, E.L., Beheregaray, L.B., Russello, M.A., Glaberman, S., Boore, J., and Caccone, A. 2020. Colonization history of Galapagos giant tortoises: insights from mitogenomes support the progression rule. Journal of Zoological Systematics and Evolutionary Research 58:1262–1275.

Powell, R., Conant, R., and Collins, J.T. 2016. Peterson Field Guide to the Reptiles and Amphibians of the Eastern and Central North America. Fourth Edition. Boston, MA: Houghton Mifflin Harcourt, 494 pp.

Power, J.H. 1927. On the herpetological fauna of the Lobatsi-Linokana Area. Part I. Transactions of the Royal Society of South Africa 14:405–422.

Power, J.H. 1932. *Testudo verreauxii* Smith: a study in variation. South African Journal of Science 29:466–472.

Power, T. and Gilhen, J. 2018. Status, distribution, and nesting ecology of Snapping Turtle (*Chelydra serpentina*) on Cape Breton Island, Nova Scotia, Canada. Canadian Field-Naturalist 132:8–17.

Pradhan, A., Chettri, N., and Sengupta, S. 2021. First record of the Assam Leaf Turtle *Cyclemys gemeli* (Fritz et al. 2008) (Reptilia: Testudines: Geoemydidae) from the Darjeeling-Sikkim Himalaya, India. Journal of Threatened Taxa 12:16909–16911.

Prasad, K.N. and Satsangi, P.P. 1967. On a new chelonian from the Siwalik beds of Himachal Pradesh. Records of the Geological Survey of India 95:533–536.

Praschag, P. and Gemel, R. 2002. Identity of the black softshell turtle *Aspideretes nigricans* (Anderson, 1875), with remarks on related species. Faunistische Abhandlungen Staatliches Museum für Tierkunde Dresden 23:87–116.

Praschag, P. and Gemel, R. 2022. A new subspecies of the Asiatic softshell turtle *Amyda ornata* (Gray, 1861) from its northern distribution range. Herpetozoa 35:81–93.

Praschag, P., Schmidt, C., Fritzsch, G., Müller, A., Gemel, R., and Fritz, U. 2006. *Geoemyda silvatica*, an enigmatic turtle of the Geoemydidae (Reptilia: Testudines), represents a distinct genus. Organisms, Diversity, and Evolution 6:151–162.

Praschag, P., Hundsdörfer, A.K., Reza, A.H.M.A., and Fritz, U. 2007a. Genetic evidence for wild-living *Aspideretes nigricans* and a molecular phylogeny of South Asian softshell turtles (Reptilia: Trionychidae: *Aspideretes, Nilssonia*). Zoologica Scripta 36(4):301–310.

Praschag, P., Hundsdörfer, A.K., and Fritz, U. 2007b. Phylogeny and taxonomy of endangered South and South-east Asian freshwater turtles elucidated by mtDNA sequence variation (Testudines: Geoemydidae: *Batagur, Callagur, Hardella, Kachuga, Pangshura*). Zoologica Scripta 36(5):429–442.

Praschag, P., Sommer, R.S., McCarthy, C., Gemel, R., and Fritz, U. 2008. Naming one of the world's rarest chelonians, the southern *Batagur*. Zootaxa 1758:61–68.

Praschag, P., Holloway, R., Georges, A., Päckert, M., Hundsdörfer, A.K., and Fritz, U. 2009a. A new subspecies of *Batagur affinis* (Cantor, 1847), one of the world's most critically endangered chelonians (Testudines: Geoemydidae). Zootaxa 2233:57–68.

Praschag, P., Hundsdörfer, A.K., and Fritz, U. 2009b. Further specimens and phylogenetic position of the recently described leaf turtle species *Cyclemys gemeli* (Testudines: Geoemydidae). Zootaxa 2008:29–37.

Praschag, P., Stuckas, H., Päckert, M., Maran, J., and Fritz, U. 2011. Mitochondrial DNA sequences suggest a revised taxonomy of Asian flapshell turtles (*Lissemys* Smith, 1931) and the validity of previously unrecognized taxa (Testudines: Trionychidae). Vertebrate Zoology 61(1):147–160.

Praschag, P., Ihlow, F., Flecks, M., Vamberger, M., and Fritz, U. 2017. Diversity of North American map and sawback turtles (Testudines: Emydidae: *Graptemys*). Zoologica Scripta 2017(00), doi:10.1111/zsc.12249, 8 pp.

Praschag, P., Rai, T.P., and Schleich, H. 2022. First record of the critically endangered Black softshell turtle, *Nilssonia nigricans* (Anderson, 1875) for Nepal. ARCO Veröffentlichungen – Arco-Nepal Newsletter 24:15–22.

Premkishore, G. and Chandran, M.R. 1996. Nesting studies of two freshwater turtles (*Lissemys punctata punctata* and *Melanochelys trijuga trijuga*) of Tamil Nadu, India, in the context of their conservation. Annales des Sciences Naturelles – Zoologie et Biologie Animale 17:99–104.

Preston, R.E. 1979. Late Pleistocene cold-blooded vertebrate faunas from the mid-Continental United States. I. Reptilia, Testudines, Crocodilia. University of Michigan Museum of Paleontology, Papers on Paleontology, No. 19, 53 pp.

Preston, R.E. and McCoy, C.J. 1971. The status of *Emys twentei* (Reptilia: Testudinidae) based on new fossil records from Kansas and Oklahoma. Journal of Herpetology 5:23–30.

Pritchard, P.C.H. 1967. Living Turtles of the World. Jersey City: TFH Publ., 288 pp.

Pritchard, P.C.H. 1971a. A further report on Galapagos tortoises. Herpetological Review 3(1):25–27.

Pritchard, P.C.H. 1971b. Galapagos tortoises, 1971. Herpetological Review 3(3):49–51.

Pritchard, P.C.H. 1979. Encyclopedia of Turtles. Neptune, NJ: TFH Publications, 895 pp.

Pritchard, P.C.H. 1980a. *Dermochelys coriacea*. Catalogue of American Amphibians and Reptiles 238:1–4.

Pritchard, P.C.H. 1980b. Record size turtles from Florida and South Amerika [sic]. Chelonologica 1(3):113–123.

Pritchard, P.C.H. 1984. Piscivory in turtles, and evolution of the long-necked Chelidae. In: Ferguson, M.W. (Ed.). The Structure, Development and Evolution of Reptiles. Symposia of the Zoological Society of London 52:87–110.

Pritchard, P.C.H. 1989. The Alligator Snapping Turtle: Biology and Conservation. Milwaukee: Milwaukee Public Museum, 104 pp.

Pritchard, P.C.H. 1990. Turtles of the World (book review). Copeia 1990: 62–67.

Pritchard, P.C.H. 1994. Comments on *Chinemys palaeannamitica* and certain other chelonian taxa based upon material from archaeological sites. Chelonian Conservation and Biology 1(2):131–137.

Pritchard, P.C.H. 1996. The Galápagos Tortoises: Nomenclatural and Survival Status. Chelonian Research Monographs 1:1–85.

Pritchard, P.C.H. 2000. *Indotestudo travancorica*...a valid species of tortoise? Reptile and Amphibian Hobbyist 5(6):18–28.

Pritchard, P.C.H. 2001. Observations on body size, sympatry, and niche divergence in softshell turtles (Trionychidae). Chelonian Conservation and Biology 4(1):5–27.

Pritchard, P.C.H. 2005. The Pinta Tortoise: Globalization and the Extinction of Island Species. Worldviews for the 21st Century: A Monograph Series 3(2):1–46.

Pritchard, P.C.H. 2012. *Rafetus*: The Curve of Extinction. The Story of the Giant Softshell Turtle of the Yangtze and Red Rivers. Ada, OK: Living Art Publishing, 173 pp.

Pritchard, P.C.H. and McCord, W.P. 1991. A new emydid turtle from China. Herpetologica 47(2):139–147.

Pritchard, P.C.H. and Mortimer, J.A. 1999. Taxonomy, external morphology, and species identification. In: Eckert, K.L., Bjorndal, K.A., Abreu–Grobois, F.A., and Donnelly, M. (Eds.). Research and Management Techniques for the Conservation of Sea Turtles. IUCN/SSC Marine Turtle Specialist Group, Publ. No. 4, pp. 21–38.

Pritchard, P.C.H. and Pronek, N. 1982. Request for suppression of *Kinosternon alamose* and *K. oaxacae* Pritchard, 1979 (Reptilia, Testudines). Bulletin of Zoological Nomenclature 39(3):212–213.

Pritchard, P.C.H. and Trebbau, P. 1984. The Turtles of Venezuela. Society for the Study of Amphibians and Reptiles, Contributions in Herpetology No. 2, 403 pp.

Prokop, H. 2010. Breeding of the endemic Turkana Mud Turtle, *Pelusios broadleyi* Bour, 1986. Radiata 19(2):2–27.

Protiva, T., Gunalen, D., Bauerová, A., Palupčíková, K., Somerová, B., Frýdlová, P., Jančúchová-Lásková, J., Šimková, O., Frynta, D., and Rehák, I. 2016. Shell shape and genetic variability of Southeast Asian Box Turtles (*Cuora amboinensis*) from Borneo and Sumatra. Vertebrate Zoology 66:387–396.

Pupins, M. and Pupina, A. 2011. First records of 5 allochthonous species and subspecies of turtles (*Trachemys scripta troostii, Mauremys caspica, Mauremys rivulata, Pelodiscus sinensis, Testudo horsfieldii*) and new records of subspecies *Trachemys scripta elegans* in Latvia. Management of Biological Invasions 2:69–81.

Pyle, R.L., Barik, S.K., Christidis, L., Conix, S., Costello, M.J., van Dijk, P.P., Garnett, S.T., Hobern, D., Kirk, P.M., Lien, A.M., Orrell, T.M., Remsen, D., Thomson, S.A., Wambiji, N., Zachos, F.E., Zhang, Z.Q., and Thiele, K.R. 2021. Towards a global list of accepted species V. The devil is in the detail. Organisms Diversity & Evolution 21:657–675.

Qin, J.Q. 1992 ["1991"]. A new species of genus *Clemmys*: *C. guangxiensis*. In: Qian, Y.M., Zhao, E.M., and Zhao, K.T. (Eds.). Animal Science Research. A volume issued to celebrate the 90th birthday of Mangven L.Y. Chang. China Forestry Press, Beijing (12)2:60–62.

Quoy, J.-R.-C. and Gaimard, J.-P. 1824a. Description d'une nouvelle espèce de tortue et de trois espèces nouvelles de scinques. Bulletin des Sciences Naturelles et de Géologie, Paris 1:90–91.

Quoy, J.-R.-C. and Gaimard, J.-P. 1824b. Sous-genre Tortue de Terre–*Testudo* Brongn. Tortue Noire–*Testudo nigra* N. In: Freycinet, M.L. de. Voyage Autour du Monde, Entrepris par le Ministère et conformément aux instructions de s. exc. M. le Vicomte du Bouchage, Secrétaire d'Etat au Département de la Marine, Exécuté sur les corvettes de S.M. l'Uranie et la Physicienne, pendant les années 1817–1820. Zoologie. Paris: Pillet Aîné, pp. 174–175.

Rabi, M., Worthy, T., Hawkins, S., Bedford, S., and Spriggs, M. 2019. Neolithic human-induced extinction of previously unrecognized giant testudinid tortoises endemic to Melanesia. Journal of Vertebrate Paleontology, Program and Abstracts, 2019, p. 175.

Raemy, M., Fritz, U., Cheylan, M., and Ursenbacher, S. 2017. Hybridisation between turtle subspecies: a case study with the European pond turtle (*Emys orbicularis*). Conservation Genetics 18:287–296.

Rafinesque, C.S. 1814. Prodromo di erpetologia Siciliana. Specchio delle Scienze, Palermo 2(9):65–67, 102–104. [in Italian]

Rafinesque, C.S. 1815. Analyse de la Nature ou Tableau de l'Univers et des Corps Organisés. Palermo: 223 pp.

Rafinesque, C.S. 1817. Tracts of C.S. Rafinesque. In: Rafinesque, C.S. Florula Ludoviciana; or a Flora of the State of Louisiana. New York: C. Wiley and Co., pp. 166–172.

Rafinesque, C.S. 1818. Discoveries in natural history, made during a journey through the western region of the United States. The American Monthly Magazine and Critical Review 3(5):354–356.

Rafinesque, C.S. 1822. On the turtles of the United States. Kentucky Gazette (n.s.1)36(no.21, May 23):3.

Rafinesque, C.S. 1832. Description of two new genera of soft shell turtles of North America. Atlantic Journal and Friend of Knowledge 1:64–65.

Rahman, S.C. 2012. Keeled box turtle. The Daily Star, Dhaka: January 3, 2012, http://archive.thedailystar.net/newDesign/news-details.php?nid=216819.

Rahman, S.C., Rashid, S.M.A., Datta, R., Mro, P., and Roy, C.J. 2015. Status, exploitation, and conservation of freshwater turtles and tortoises in Chittagong Hill Tracts, Bangladesh. Chelonian Conservation and Biology 14:130–135.

Rai, K.R. 2004. Ecological distribution of *Cyclemys oldhamii* (Gray 1863) from Nepal. Our Nature 2:7–12.

Rai, K.R., Kästle, W., and Schleich, H.H. 2002. *Pangshura tentoria circumdata* Mertens, 1969. In: Schleich, H.H. and Kästle, W. (Eds.). Amphibians and Reptiles of Nepal. Ruggell, Germany: A.R.G. Gantner Verlag, pp. 547–549.

Ramírez-González, C.G. and Canseco-Márquez, L. 2015. *Chelydra rossignonii*, confirmación de su presencia en el estado de Oaxaca, México. Revista Mexicana de Biodiversidad 86:832–834.

Ramsay, E.P. 1886. On a new genus and species of fresh water tortoise from the Fly River, New Guinea. Proceedings of the Linnaean Society of New South Wales (2)1:158–162.

Ranjbar, N., Malekian, M., Ashrafzadeh, M.R., and Hemami, M.R. 2022. Phylogeographic and phenotypic divergence between two subspecies of *Testudo graeca* (*T. g. buxtoni* and *T. g. zarudnyi*) across their contact zone in Iran. Scientific Reports 12:13579, 15 pp.

Ranzani, C. 1832. De testudine coriacea marina. Bononiae [Bologna]: Tiocchi, 11 pp.

Rashid, S.M.A. and Swingland, I.R. 1997. On the ecology of some freshwater turtles in Bangladesh. In: Van Abbema, J. (Ed.). Proceedings: Conservation, Restoration, and Management of Tortoises and Turtles – An International Conference. N.Y. Turtle and Tortoise Society, pp. 225–242.

Ray, J. 1693. Synopsis Methodica Animalium Quadrupedum et Serpentini Generis. Londini (London): Smith and Walford, 336 pp.

Reed, K.M., Hanks, B.G., Bickham, J.W., Rhodin, A.G.J., Greenbaum, I.F., Mittermeier, R.A., and Fedullo, L.P. 1991. Cytogenetic analysis of the pleurodine turtle *Phrynops hogei* and its taxonomic implications. Amphibia-Reptilia 12(2):203–212.

Regis, K.W. and Meik, J.M. 2017. Allometry of sexual size dimorphism in turtles: a comparison of mass and length data. PeerJ 5:e2914; doi:10.7717/peerj.2914.

Reid, B.N., Le, M., McCord, W.P., Iverson J.B., Georges, A., Bergmann, T., Amato, G., Desalle, R., and Naro-Maciel, E. 2011. Comparing and combining distance-based and character-based approaches for barcoding turtles. Molecular Ecology Resources 11:956–967.

Reimann, M. 1979. [*Geoemyda trijuga wiroti*, *Testudo nutapundi*]. In: Nutaphand, W. The Turtles of Thailand. Bangkok: Siamfarm Zoological Garden, pp. 177–178, 193–195.

Renner, S.S. 2016. A return to Linnaeus's focus on diagnosis, not description: the use of DNA characters in the formal naming of species. Systematic Biology 65(6):1085–1095.

Reshetnikov, A.N., Zibrova, M.G., Ayaz, D., Bhattarai, S., Borodin, O.V., Borzée, A., Brejcha, J., Çiçek, K., Dimaki, M., Doronin, I.V., Drobenkov, S.M., Gichikhanova, U.A., Gladkova, A.Y., Gordeev, D.A., Ioannidis, Y., Ilyukh, M.P., Lnteresova, E.A., Jadhav, T.D., Karabanov, D.P., Khabibullin, V.F., Khabilov, T.K., Khan, M.M.H., Kidov, A.A., Klimov, A.S., Kochetkov, D.N., Kolbintsev, V.G., Kuzmin, S.L., Lotiev, K.Y., Louppova, N.E., Lvov, V.D., Lyapkov, S.M., Martynenko, I.M., Maslova, I.V., Masroor, R., Mazanaeva, L.F., Milko, D.A., Milto, K.D., Mozaffari, O., Nguyen, T.Q., Novitsky, R.V., Petrovskiy, A.B., Prelovskiy, V.A., Serbin, V.V., Shi, H.-T., Skalon, N.V., Struijk, R.P.J.H., Taniguchi, M., Tarkhnishvili, D., Tsurkan, V.F., Tyutenkov, O.Y., Ushakov, M.V., Vekhov, D.A., Xiao, F., Yakimov, A.V., Yakovleva, T.I., Yang, P., Zeleev, D.F., and Petrosyan, V.G. 2023. Rarely naturalized, but widespread and even invasive: the paradox of a popular pet terrapin expansion in Eurasia. NeoBiota 81:91–127.

Retzius, A.J. 1792. [*Testudo tricarinata, Testudo scabra*]. In: Schoepff, J.D. 1792. Historia Testudinum Iconibus Illustrata. Erlangae: Ioannis Iacobi Palm, 136 pp. [pp. 9–16].

Reuss, A. 1844. [*Emys lyrae*]. In: Gray, J.E. Catalogue of the Tortoises, Crocodiles, and Amphisbaenians in the Collection of the British Museum, p.39.

Reyes-Grajales, E., Macip-Ríos, R., Iverson, J.B., and Matamoros, W.A. 2021. Population ecology and morphology of the Central Chiapas Mud Turtle (*Kinosternon abaxillare*). Chelonian Conservation and Biology 20(1):18–26.

Reynolds, L.N. 2022. Geographical Distributions: Pelomedusidae, *Pelusios gabonensis* Duméril, 1856, Forest Hinged Terrapin. African Herp News 79:45–48.

Reynolds, R.P., Gotte, S.W., and Ernst, C.H. 2007. Catalog of type specimens of recent Crocodilia and Testudines in the National Museum of Natural History, Smithsonian Institution. Smithsonian Contributions to Zoology 626:1–56.

Reynoso, V.H. and Montellano-Ballesteros, M. 2004. A new giant turtle of the genus *Gopherus* (Chelonia: Testudinidae) from the Pleistocene of Tamaulipas, México, and a review of the phylogeny and biogeography of gopher tortoises. Journal of Vertebrate Paleontology 24(4):822–837.

Rezazadeh, E., Alucheh, R.M., and Kami, H.G. 2014. A preliminary study on the Mediterranean spur-thighed tortoise *Testudo graeca* Linnaeus, 1758 from northwestern Iran. Herpetology Notes 7:127–133.

Rhodin, A.G.J. 1993. The turtles of Mount Desert Island, Maine: summary of research results, 1991-1992, with analysis of painted turtle (*Chrysemys picta*) systematics and natural history in New England. Report to Maine Department of Inland Fisheries and Wildlife and National Park Service, Acadia National Park, 51 pp.

Rhodin, A.G.J. 1994a. Chelid turtles of the Australasian Archipelago: I. A new species of *Chelodina* from southeastern Papua New Guinea. Breviora 497:1–36.

Rhodin, A.G.J. 1994b. Chelid turtles of the Australasian Archipelago: II. A new species of *Chelodina* from Roti Island, Indonesia. Breviora 498:1–31.

Rhodin, A.G.J. 2006. Turtles and humans in Florida and the World: a global perspective on diversity, threats, and economic development. In: Meylan, P.A. (Ed.). Biology and Conservation of Florida Turtles. Chelonian Research Monographs 3:18–27.

Rhodin, A.G.J. and Butler, B.O. 1997. The painted turtles (*Chrysemys picta*) of New England: taxonomy, morphometrics, and reproduction. In: Tyning, T.F. (Ed.). Status and Conservation of Turtles of the Northeastern United States. Lanesboro, MN: Serpent's Tale, pp. 34–40.

Rhodin, A.G.J. and Carr, J.L. 2009. A quarter millenium of uses and misuses of the turtle name *Testudo scabra*: identification of the type specimens of *T. scabra* Linnaeus 1758 (= *Rhinoclemmys punctularia*) and *T. scripta* Thunberg *in* Schoepff 1792 (= *Trachemys scripta scripta*). Zootaxa 2226:1–18.

Rhodin, A.G.J. and Genorupa, V.R. 2000. Conservation status of freshwater turtles in Papua New Guinea. In: van Dijk, P.P., Stuart, B.L., and Rhodin, A.G.J. (Eds.). Asian Turtle Trade: Proceedings of a Workshop on Conservation and Trade of Freshwater Turtles and Tortoises in Asia. Chelonian Research Monographs 2:129–136.

Rhodin, A.G.J. and Mittelhauser, G.H. 1994. Maximum size and clutch size records for eastern painted turtles, *Chrysemys picta picta*, from mid-coastal Maine. Chelonian Conservation and Biology 1(2):148–150.

Rhodin, A.G.J. and Mittermeier, R.A. 1976. *Chelodina parkeri*, a new species of chelid turtle from New Guinea, with a discussion of *Chelodina siebenrocki* Werner, 1901. Bulletin of the Museum of Comparative Zoology 147(11):465–488.

Rhodin, A.G.J. and Mittermeier, R.A. 1983. Description of *Phrynops williamsi*, a new species of chelid turtle of the South American *P. geoffroanus* complex. In: Rhodin, A.G.J. and Miyata, K. (Eds.). Advances in Herpetology and Evolutionary Biology. Essays in Honor of Ernest E. Williams. Cambridge, MA: Museum of Comparative Zoology, pp. 58–73.

Rhodin, A.G.J. and Smith, H.M. 1982. The original authorship and type specimen of *Dermochelys coriacea*. Journal of Herpetology 16(3):316–317.

Rhodin, A.G.J., Mittermeier, R.A., and Rocha e Silva, R.D. 1982. Distribution and taxonomic status of *Phrynops hogei*, a rare chelid turtle from southeastern Brazil. Copeia 1982(1):179–181.

Rhodin, A.G.J., Mittermeier, R.A., and McMorris, J.R. 1984a. *Platemys macrocephala*, a new species of chelid turtle from central Bolivia and the Pantanal region of Brazil. Herpetologica 40(1):38–46.

Rhodin, A.G.J., Rocha e Silva, R.D., and Mittermeier, R.A. 1984b. Distribution of the South American chelid turtles *Platemys radiolata* and *P. spixii*. Copeia 1984(3):780–786.

Rhodin, A.G.J., Mittermeier, R.A., and Buskirk, J.R. 1988. *Phrynops williamsi*. Catalogue of American Amphibians and Reptiles 439:1–2.

Rhodin, A.G.J., van Dijk, P.P., and Parham, J.F. 2008. Turtles of the world: annotated checklist of taxonomy and synonymy. In: Rhodin, A.G.J., Pritchard, P.C.H., van Dijk, P.P., Saumure, R.A., Buhlmann, K.A., and Iverson, J.B. (Eds.). Conservation Biology of Freshwater Turtles and Tortoises: A Compilation Project of the IUCN/SSC Tortoise and Freshwater Turtle Specialist Group. Chelonian Research Monographs 5(1):000.1–000.38.

Rhodin, A.G.J., Kaiser, H., van Dijk, P.P., Wüster, W., O'Shea, M., Archer, M., Auliya, M., Boitani, L., Bour, R., Clausnitzer, V., Contreras-Macbeath, T., Crother, B.I., Daza, J.M., Driscoll, C.A., Flores-Villela, O., Frazier, J., Fritz, U., Gardner, A., Gascon, C., Georges, A., Glaw, F., Grazziotin, F.G., Groves, C.P., Haszprunar, G., Havaš, P., Hero, J.M., Hoffmann, M., Hoogmoed, M.S., Horne, B.D., Iverson, J.B., Jäch, M., Jenkins, C.L., Jenkins, R.K.B., Kiester, A.R., Keogh, J.S., Lacher, T.E., Jr., Lovich, J.E., Luiselli, L., Mahler, D.L., Mallon, D., Mast, R., Mcdiarmid, R.W., Measey, J., Mittermeier, R.A., Molur, S., Mossbrugger, V., Murphy, R., Naish, D., Niekisch, M., Ota, H., Parham, J.F., Parr, M.J., Pilcher, N.J., Pine, R.H., Rylands, A.B., Sanderson, J.G., Savage, J., Schleip, W., Scrocchi, G.J., Shaffer, H.B., Smith, E.N., Sprackland, R., Stuart, S.N., Vetter, H., Vitt, L.J., Waller, T., Webb, G., Wilson, E.O., Zaher, H., and Thomson, S. 2015. Comment on *Spracklandus* Hoser, 2009 (Reptilia, Serpentes, Elapidae): request for confirmation of the availability of the generic name and for the nomenclatural validation of the journal in which it was published. Bulletin of Zoological Nomenclature 72(1):65–78.

Rhodin, A.G.J., Quinn, H.R., Mittermeier, R.A., Heard, N., Launay, F., and Al Mubarak, R. 2018a. Two million-dollar turtle funds: Turtle Conservation Fund (TCF) and Mohamed bin Zayed Species Conservation Fund (MBZ). Turtle Survival 2017:54–55.

Rhodin, A.G.J., Stanford, C.B., van Dijk, P.P., Eisemberg, C., Luiselli, L., Mittermeier, R.A., Hudson, R., Horne, B.D., Goode, E.V., Kuchling, G., Walde, A., Baard, E.H.W., Berry, K.H., Bertolero, A., Blanck, T.E.G., Bour, R., Buhlmann, K.A., Cayot, L.J., Collett, S., Currylow, A., Das, I., Diagne, T., Ennen, J.R., Forero-Medina, G., Frankel, M.G., Fritz, U., García, G., Gibbons, J.W., Gibbons, P.M., Gong, S., Guntoro, J., Hofmeyr, M.D., Iverson, J.B., Kiester, A.R., Lau, M., Lawson, D.P., Lovich, J.E., Moll, E.O., Páez, V.P., Palomo-Ramos, R., Platt, K., Platt, S.G., Pritchard, P.C.H., Quinn, H.R., Rahman, S.C., Randrianjafizanaka, S.T., Schaffer, J., Selman, W., Shaffer, H.B., Sharma, D.S.K., Shi, H., Singh, S., Spencer, R., Stannard, K., Sutcliffe, S., Thomson, S., and Vogt, R.C. 2018b. Global conservation status of turtles and tortoises (Order Testudines). Chelonian Conservation and Biology 17(2):135–161.

Rhymer, J.M. and Simberloff, D. 1996. Extinction by hybridization and introgression. Annual Review of Ecology and Systematics 27:83–109.

Riabinin, A.N. 1918. [Sur les tortues des dépôts méotiens de Bessarabie]. Trudy Geol. Min. Muz. Petra Velikago Imper. Akad. Nauk 1(1915–1918):1–16. [in Russian]

Riche, M. 1801. [*Testudo amboinensis*]. In: Daudin, F.M. Histoire Naturelle, Générale et Particulière, des Reptiles. Tome Second. Paris: Dufart, 432 pp. [pp. 309–312].

Ríos-Solís, J.A., Lavariega, M.C., García-Padilla, E., and Mata-Silva, V. 2021. Noteworthy records of freshwater turtles in Oaxaca, Mexico. Revista Latinoamericana de Herpetología 4(2):184–191.

Ripple, W.J., Wolf, C., Newsome, T.M., Hoffmann, M., Wirsing, A.J., and McCauley, D.J. 2017. Extinction risk is most acute for the world's largest and smallest vertebrates. PNAS 114(40):10683.

Ritgen, F.A. 1828. Versuch einer Natürlichen Eintheilung der Amphibien. Nova Acta Physico-Medica Academiae Caesareae Leopoldino-Carolinae Naturae Curiosorum 14:245–284.

Rivas, G., McCord, W.P., Barros, T.R., and Barrio-Amorós, C.L. 2007. *Rhinoclemmys diademata* (Mertens, 1954) or "Galapago de Maracaibo" (Testudines; Geoemydidae): an unprotected turtle in the Maracaibo Basin, Venezuela. Radiata 16(1):16–23.

Rivas, G.A., Barros, T.R., Molina, F.D.B., Trebbau, P., and Pritchard, P.C.H. 2015. The presence of *Mesoclemmys raniceps* and *Mesoclemmys nasuta* in Venezuela and comments on the Type Locality of *Hydraspis maculata* (Chelidae). Chelonian Conservation and Biology 14(1):104–107.

Rivers, J.J. 1889. Description of a new turtle from the Sacramento River, belonging to the family of Trionychidae. Proceedings of the California Academy of Sciences (2)2:333–336.

Rochebrune, A.T. de. 1884. Faune de la Sénégambie. Reptiles. Paris: O. Dom, 221 pp.

Rodrigues, J.F.M. and Lima-Ribeiro, M.S. 2018. Predicting where species could go: climate is more important than dispersal for explaining the distribution of a South American turtle. Hydrobiologia 808:343–352.

Rodrigues, J.F.M., Coelho, M.T.P., and Diniz-Filho, J.A.F. 2016. Exploring intraspecific climatic niche conservatism to better understand species invasion: the case of *Trachemys dorbigni* (Testudines, Emydidae). Hybrobiologia 779:127–134.

Rodrigues, J.F.M., Olalla-Tárraga, M.A., Iverson, J.B., and Diniz-Filho, J.A.F. 2018. Temperature is the main correlate of the global biogeography of turtle body size. Global Ecology and Biogeography 27:429–438.

Rodríguez-Robles, J.A., Good, D.A., and Wake, D.B. 2003. Brief history of herpetology in the Museum of Vertebrate Zoology, University of California, Berkeley, with a list of type specimens of Recent amphibians and reptiles. University of California Publications in Zoology 131:1–119.

Rodríguez Schettino, L., Mancina, C.A., and Rivalta González, V. 2013. Reptiles of Cuba: checklist and geographic distributions. Smithsonian Herpetological Information Service 144:1–96.

Rogner, M. 1996. Schildkröten 2. Hürtgenwald: Heidi-Rogner-Verlag, 265 pp.

Rogner, M., Iverson, J.B., Berry, J.F., Seidel, M.E., and Rhodin, A.G.J. 2013. Case 3625. *Kinosternon chimalhuaca* Berry, Seidel, & Iverson *in* Rogner, 1996 (Reptilia, Testudines): proposed confirmation of the publication date. Bulletin of Zoological Nomenclature 70(3):190–192.

Rogner, M., Iverson, J.B., Berry, J.F., Seidel, M.E., and Rhodin, A.G.J. 2016.

Comments on the proposed suppression *Kinosternon chimalhuaca* Rogner, 1996 (Reptilia, Testudines): (Case 3625). Bulletin of Zoological Nomenclature 73(1):59–60.

Rohilla, M.S., Rao, R.J., and Tiwari, P.K. 2009. Enzyme polymorphism in Indian freshwater soft shell turtle *Lissemys punctata*. Acta Herpetologica 4(1):15–28.

Roman, J., Santhuff, S.D., Moler, P.E., and Bowen, B.W. 1999. Population structure and cryptic evolutionary units in the alligator snapping turtle. Conservation Biology 13(1):135–142.

Román-Palacios, C. and Wiens, J.J. 2018. The tortoise and the finch: testing for island effects on diversification using two iconic Galápagos radiations. Journal of Biogeography 45:1701–1712.

Rook, L., Croitor, R., Delfino, M., Ferretti, M.P., Gallai, G., and Pavia, M. 2013. The Upper Valdarno Plio-Pleistocene vertebrate record: an historical overview, with notes on palaeobiology and stratigraphic significance of some important taxa. Italian Journal of Geosciences 132:104–125.

Rosales-Martínez, C.S., Escobedo-Galván, A.H., and Cupul-Magaña, F.G. 2021. Distribución *Rhinoclemmys rubida perixantha*. Revista Latinoamericana de Herpetologia 4(2):234–235.

Rose, F.L. and Judd, F.W. 2014. The Texas Tortoise: A Natural History. Norman, OK: Univeristy of Oklahoma Press, 189 pp.

Rosen, P.C., Hermann, H.-W., and Melendez, C. 2006. Phylogeography of the Southwestern mud turtle (*Kinosternon sonoriense* Leconte). Unpublished final report to Arizona Game and Fish Department and CONABIO, Mexico, 20 pp.

Rosenbaum, D., Rudolph, D.C., Saenz, D., Fitzgerald, L.A., Nelson, R.E., Collins, C.S., Hibbitts, T.J., Maxey, R.W., Crump, P., and Schalk, C.M. 2023. Distribution and demography of the Alligator Snapping Turtle (*Macrochelys temminckii*) in Texas: a 20-year perspective. Southeastern Naturalist 22(Special Issue 12):197–220.

Ross, W. and Macartney, J. 1802. Lectures on Comparative Anatomy, translated from the French of G. Cuvier. Volume 1. London, Oriental Press, 710 pp.

Rostal, D.C., McCoy, E.D., and Mushinsky, H.R. (Eds.). 2014. Biology and Conservation of North American Tortoises. Baltimore, Maryland: Johns Hopkins University Press, 182 pp.

Rothschild, W. 1901. On a new land-tortoise from the Galapagos Islands. Novitates Zoologicae 8:372.

Rothschild, W. 1902. Description of a new species of gigantic land-tortoise from the Galapagos Islands. Novitates Zoologicae 9:619.

Rothschild, W. 1903. Description of a new species of gigantic land tortoise from Indefatigable Island. Novitates Zoologicae 10:119.

Rothschild, W. 1906. A new species of giant tortoise. Novitates Zoologicae 13:753–754.

Rothschild, W. 1915. On the gigantic land tortoises of the Seychelles and Aldabra-Madagascar group with some notes on certain forms of the Mascarene group. Novitates Zoologicae 22:418–442.

Rouot, S. and Maurer, M. 2023. Monograph and natural history of the Ryukyu black breasted leaf turtle, *Geoemyda japonica* Fan, 1932. Radiata 32(2):24–38.

Roux-Esteve, R. 1979. Liste des amphibiens et reptiles des collections du Muséum National d'Histoire Naturelle de Paris, recoltés par LeSueur (1778–1846). Bulletin Trimestriel de la Société Géologique de Normandie et des Amis du Muséum du Havre 66(3):25–29.

Rowe, J.W. 1997. Growth rate, body size, sexual dimorphism and morphometric variation in four populations of painted turtles (*Chrysemys picta bellii*) from Nebraska. American Midland Naturalist 138:174–188.

Rudolphi, M. and Weser, R. 2000. Zucht der F2 Generation bei der Riesenerdschildkröte *Heosemys grandis* (Gray, 1860). Sauria 22:17–24.

Rueda-Almonacid, J.V., Carr, J.L., Mittermeier, R.A., Rodríguez-Mahecha, J.V., Mast, R.B., Vogt, R.C., Rhodin, A.G.J., de la Ossa-Velásquez, J., Rueda, J.N., and Mittermeier, C.G. 2007. Las Tortugas y los Cocodrilianos de los Países Andinos del Trópico. Bogotá, Colombia: Editorial Panamericana, Formas e Impresos, Serie de Guías Tropicales de Campo No. 6, Conservación Internacional, 538 pp.

Ruiz de Xelva, M. 1801. [*Testudo bispinosa*]. In: Daudin, F.M. Histoire Naturelle, Generale et Particuliere, des Reptiles. Tome Second. Paris: Dufart, 432 pp. [pp. 94–97].

Rummler, H.-J. and Fritz, U. 1991. Geographische Variabilität der Amboina-Scharnierschildkröte *Cuora amboinensis* (Daudin, 1802), mit Beschreibung einer neuen Unterart, *C. a. kamaroma* subsp. nov. Salamandra 27(1):17–45.

Rüppell, E. 1835. Neue Wirbelthiere zu der Fauna von Abyssinien gehörig. Amphibien. Frankfurt: Siegmund Schmerber, 18 pp.

Rüppell, E. 1845. Beschreibung und Abbildung einer neuen Art von Landschildkröten, zur Gattung *Kinyxis* gehörig. Museum Senckenbergianum 3:223–228.

Russello, M.A., Glaberman, S., Gibbs, J.P., Marquez, C., Powell, J.R., and Caccone, A. 2005. A cryptic taxon of Galápagos tortoise in conservation peril. Biological Letters 1:287–290.

Russello, M.A., Beheregaray, L.B., Gibbs, J.P., Fritts, T., Havill, N., Powell, J.R., and Caccone, A. 2007. Lonesome George is not alone among Galápagos tortoises. Current Biology 17(9):R317–R318.

Rust, H.T., Mertens, R., and Müller, L. 1934. Systematische Liste der Lebenden Schildkröten. Blätter für Aquarien- und Terrarien-Kunde 45:42–45, 59–67.

Sabaj, M.H. 2019. Standard symbolic codes for institutional resource collections in herpetology and ichthyology: an online reference. Version 7.1. Washington, DC: American Society of Ichthyologists and Herpetologists, https://asih.org/sites/default/files/2019-04/Sabaj_2019_ASIH_Symbolic_Codes_v7.1.pdf.

Sadlier, R. and Cann, J. 2018. Response to Arthur Georges' review of *Freshwater Turtles of Australia* (2017). Australian Journal of Zoology 66:88–91.

Safi, A. and Khan, M.Z. 2014. Distribution and current population status of freshwater turtles of District Charsadda of Khyber Pakhtunkhwa, Pakistan. Journal of Zoology Studies 1:31–38.

Said-Aliev, S.A. 1979. Zemnovodnje i presmykajuszcziesja Tadzhikistana. [Amphibians and reptiles of Tajikistan]. Donish: Dushanbe, 145 pp.

Sajdak, R.A. and Molina, F.D.B. 1991. *Geochelone denticulata* (Yellowfoot Tortoise). Maximum size. Herpetological Review 22(3):98.

Salleh, M.H.M., Esa, Y., and Pau, S.N. 2023. Conservation genetics of the critically endangered Southern River Terrapin (*Batagur affinis*) in Malaysia: genetic diversity and novel subspecies distribution ranges. Biology 12(520):1–22, doi.org/10.3390/biology12040520.

Salvator, L. 1897. Die Balearen, geschildert in Wort und Bild. Zweiter Band. Würzburg and Leipzig: Hofbuchhandlung Leo Woerl, 452 pp.

Sampedro Marín, A.C. 2012. *Trachemys decussata* Gray, 1831 – Jicotea, Cuban Slider. In: González Alonso, H., Rodríguez Schettino, L., Rodríguez, A., Mancina, C.A., and Ramos García, I. (Eds.). 2012. Libro Rojo de los Vertebrados de Cuba. La Habana: Editorial Academia, Editorial Academia, pp. 196–200.

Sanches, T.M. and Bellini, C. 2002. *Chelonia mydas* (Green Sea Turtle). Adult male size. Herpetological Review 33(3):199–200.

Sánchez, J., Alcalde, L., and Bolzán, A.D. 2015. First evidence of chromosomal variation within *Chelonoidis chilensis* (Testudines: Testudinidae). Herpetological Journal 25:83–89.

Sánchez, J., Holley, J.A., Poljak, S., Bolzán, A.D., and Bravi, C.M. 2017. Phylogenetic and divergence time analysis of the *Chelonoidis chilensis* complex (Testudines: Testudinidae). Zootaxa 4320(3):487–504.

Sánchez, R.M., Semeñiuk, M.B., Cassano, M.J., Alcalde, L., Leynaud, G.C., and Moreno, L. 2019. Review of chelid and emydid turtle distributions in southern South America with emphasis on extralimital populations and new records for Argentina. Herpetological Journal 29:219–229.

Sánchez-Vialas, A., Calvo-Revuelta, M., and de la Riva, I. 2022. Synopsis of the terrestrial reptiles of Equatorial Guinea. Zootaxa 5202(1):001–197.

Sánchez-Villagra, M.R., Pritchard, P.C.H., Paolillo, A., and Linares, O.J. 1995. Geographic variation in the matamata turtle, *Chelus fimbriatus*, with observations on its shell morphology and morphometry. Chelonian Conservation and Biology 1(4):293–300.

Santana, D.O., Faria, R.G., and Mesquita, D.O. 2023. Ecology of *Phrynops geoffroanus* and *Mesoclemmys tuberculata* (Testudines: Chelidae) in climatically distinct areas of the semi-arid Caatinga and Atlantic Forest in northeast Brazil. Chelonian Conservatioon and Biology 22:89–102.

Santilli, J. and Rollinson, N. 2018. Toward a general explanation for latitudinal clines in body size among chelonians. Biological Journal of the Linnean Society 124:381–393.

Santos, R.C.d., Viana, M.d.N.S., Monjeló, L.A.d.S., Andrade, P.C.M., Pantoja-Lima, J., Oliveira, P.H.G., Vogt, R.C., Pezzuti, J.C.B., Sites, J.W., Jr., Hrbek, T., and Farias, I.P. 2016. Testing the effects of barriers on the genetic connectivity in *Podocnemis erythrocephala* (Red-Headed Amazon River Turtle): implications for management and conservation. Chelonian Conservation and Biology 15:12–22.

Sauzier, T. 1892. Tortue de terre gigantesque à l'île Maurice. La Nature, Revue des Sciences et de leurs Applications aux Arts et à l'Industrie, Paris 20:395–398.

Sauzier, T. 1899. Notes sur l'origine de la Tortue terrestre geante *T. hololissa*, Gunther. Bulletin de la Société Zoologique de France 24:138–142.

Savage, J.M. 1953. Remarks on the Indo-Chinese turtle *Annamemys merkleni*;

with special reference to the status of *Cyclemys annamensis*. Annals and Magazine of Natural History (12)6:468–472.

Savage, J.M. 2002. The Amphibians and Reptiles of Costa Rica: A Herpetofauna between Two Continents, between Two Seas. Chicago: The University of Chicago Press, 934 pp.

Savage, J.M. 2003. Lacepède, B.G.E. de la V., 1788, Histoire Naturelle des Quadrupèdes Ovipares: proposed rejection as a non-binominal work. Bulletin of Zoological Nomenclature 60:138–140.

Savage, J.M. 2020. Nicolaus Michael Oppel and his classification of amphibians and reptiles, with remarks on other 18th and early 19th century classification schemes. Herpetological Review 51:510–516.

Savage, J.M. 2021. The *nomen oblitum* rule, Article 23.9 of the International Code of Zoological Nomenclature (1999): the case of the turtle name *Xerobates berlandieri* Agassiz, 1857. Herpetological Review 52(1):61–62.

Sawyer, F.C. 1953. The dates of issue of J.E. Gray's "Illustrations of Indian Zoology" (London, 1830-1835). Journal of the Society of Bibliography of Natural History 3(1):48–55.

Say, T. 1825. On the fresh water and land tortoises of the United States. Journal of the Academy of Natural Sciences, Philadelphia 4(2):203–219, 412 [errata].

Schilde, M. 2004. Asiatische Sumpfschildkröte. Die Familie Geoemydidae in Südostasien, China und Japan. Natur und Tier Verlag, 189 pp.

Schinz, H.R. 1833. Naturgeschichte und Abbildungen der Reptilien. Leipzig: Weidmann, 240 pp.

Schlegel, H. 1844. Abbildungen neuer oder unvollständig bekannter Amphibien, nach der Natur oder dem Leben entworfen, herausgegeben und mit einem erläuternden Texte begleitet. Düsseldorf: Arnz, 141 pp.

Schlegel, H. and Müller, S. 1840. Over de Schildpadden van den Indischen Archipel, en beschrijving eener nieuwe soort van Sumatra. In: Temminck, C.J. (Ed.). Verhandelingen over de Natuurlijke Geschiedenis der Nederlandsche Overzeesche Bezittingen, 1839–44. Part 3. Zoologie, Schildpadden. Leiden: Luchtmans and van der Hoek, plate 4. [in Dutch] [published 24 April 1840; see Husson and Holthuis 1955]

Schlegel, H. and Müller, S. 1845 ["1844"]. Over de Schildpadden van den Indischen Archipel, en beschrijving eener nieuwe soort van Sumatra. In: Temminck, C.J. (Ed.). Verhandelingen over de Natuurlijke Geschiedenis der Nederlandsche Overzeesche Bezittingen, 1839–44. Part 3. Zoologie, Schildpadden. Leiden: Luchtmans and van der Hoek, pp. 29–36. [in Dutch] [published 26 June 1845; see Husson and Holthuis 1955]

Schleich, H.-H. 1982. Vorlaufige mitteilung zur herpetofauna der Kapverden. Courier Forschunginstitut Senckenberg 52:245–248.

Schleich, H.-H. 1987. Herpetofauna Caboverdiana. Spixiana (München), Supplement 12:1–75.

Schleich, H.-H. 1996a. Beitrag zur Systematik des Formenkreises von *Mauremys leprosa* (Schweigger) in Marokko. Teil I. Spixiana Suppl. 22:29–59.

Schleich, H.-H. 1996b. Lista Vermelha para os Répteis (Reptilia). In: Leyens, T. and Lobin, W. (Eds.). Primeira Lista Vermelha de Cabo Verde. Frankfurt: Courier Forschungsinstitut Senckenberg, pp. 122–125.

Schleich, H.-H. and Gruber, U. 1984. Eine neue Grosskopfschildkröte, *Platysternon megacephalum tristernalis* nov. ssp., aus Yünnan, China. Spixiana 7:67–73.

Schleich, H.-H. and Kästle, W. (Eds.). 2002. Amphibians and Reptiles of Nepal. Biology, Systematics, Field Guide. Koenigstein: Koeltz Scientific Books, pp. 1201.

Schleich, H.-H., Kästle, W., and Kabisch, K. 1996. Amphibians and Reptiles of North Africa. Koenigstein: Koeltz Scientific Books, 630 pp.

Schleicher, A. 2004. Experiences with keeping and breeding the Namibian Cape Tortoise Homopus sp. ("*Homopus bergeri*"), in a Namibian outdoor enclosure. Radiata 13(4):3–12.

Schlüter, A. and Hallermann, J. 1997. The type specimens in the herpetological collection of the Staatliches Museum für Naturkunde in Stuttgart. Stuttgarter Beiträge zur Naturkunde Serie A (553):1–15.

Schmid, K. 1819. Naturhistorische Beschreibung der Amphibien. Systematisch bearbeitet zum gemeinnützigen Gebrauche. München: Kunst-Anstalt bey der Feyertags-Schule, 95 pp.

Schmidt, K.P. 1919. Contributions to the herpetology of the Belgian Congo based on the collection of the American Congo Expedition, 1909-1915. Part I. Turtles, crocodiles, lizards, and chameleons. Bulletin of the American Museum of Natural History 39(2):385–624.

Schmidt, K.P. 1925. New reptiles and a new salamander from China. American Museum Novitates 157:1–5.

Schmidt, K.P. 1928. Amphibians and land reptiles of Porto Rico, with a list of those reported from the Virgin Islands. In: Scientific Survey of Porto Rico and the Virgin Islands. New York Academy of Science, Vol. 10, 160 pp.

Schmidt, K.P. 1941. The amphibians and reptiles of British Honduras. Zoological Series of the Field Museum of Natural History 22(8):473–510.

Schmidt, K.P. 1946. Turtles collected by the Smithsonian Biological Survey of the Panamá Canal Zone. Smithsonian Miscellaneous Collections 106(8):1–9.

Schmidt, K.P. 1947. A new kinosternid turtle from Colombia. Fieldiana Zoology 31(13):109–112.

Schmidt, K.P. 1953. A Check List of North American Amphibians and Reptiles. Sixth edition. Chicago: University of Chicago Press, 280 pp.

Schmidt, K.P. and Owens, D.W. 1944. Amphibians and reptiles of northern Coahuila, Mexico. Field Museum of Natural History Zoology 29:97–115.

Schmidtler, J.F., Rösler, H., Schmitz, A., and Ernst, N. 2014. Die Illustrationen amerikanischer Amphibien und Reptilien in den Nürnberger Prachtwerken von Johann Wolf (1818/1822): "Abbildungen und Beschreibungen merkwürdiger naturgeschichtlicher Gegenstände" und Johann Wagler (1821): "Die Amphibien. Heft I". Mertensiella 21:2–15

Schnee, P. 1900. Uber eine Sammlung südbrasilianischer Reptilien und Amphibien, nebst Beschreibung einer neuen Schildkröte (*Platemys werneri*). Zoologischer Anzeiger 23(622):461–464.

Schneider, C. and Schneider, W. 2008. The Egyptian Tortoise, *Testudo kleinmanni* Lortet, 1883 in Libya. Salamandra 44:141–152.

Schneider, J.G. 1783. Allgemeine Naturgeschichte der Schildkröten, nebst einem systematischen Verzeichnisse der einzelnen Arten und zwey Kupfern. Leipzig: J.G. Müller, 364 pp.

Schneider, J.G. 1784. Sammlung vermischter Abhandlungen zur Aufklärung der Zoologie und der Handlungsgeschichte. IV. Beiträge zu der Naturgeschichte der Schildkröten. Berlin: J.F. Unger, pp. 304–317.

Schneider, J.G. 1787. Erster Beitrag zur Naturgeschichte der Schildkröten. Leipzig: J.G. Müller, 16 pp.

Schneider, J.G. 1789. Zweiter Beitrag zur Naturgeschichte der Schildkröten. Leipzig: J.G. Müller, 32 pp.

Schneider, J.G. 1792. Beschreibung und Abbildung einer neuen Art von Wasserschildkröte nebst Bestimmungen einiger bisher wenig bekannten fremden Arten. Schriften der Gesellschaft Naturforschender Freunde zu Berlin 10:259–284.

Schneider, L., Iverson, J.B., and Vogt, R.C. 2012. *Podocnemis unifilis*. Catalogue of American Amphibians and Reptiles 890.1–33.

Schoepff, J.D. 1792. Historia Testudinum Iconibus Illustrata. Fasciculus I et II Erlangae: Ioannis Iacobi Palm, pp. 1–32, pls. I–X.

Schoepff, J.D. 1793a. Historia Testudinum Iconibus Illustrata. Fasciculus III et IV. Erlangae: Joannis Jacobi Palm, pp. 33–80, pls. XI–XVI, XVIIB–XX.

Schoepff, J.D. 1793b. Naturgeschichte der Schildkröten, mit Abbildungen erläutert. Dritter und vierter Heft. Erlangen: Johann Jacob Palm, pp. 33–88, pls. XI–XX.

Schoepff, J.D. 1795. Historia Testudinum Iconibus Illustrata. Fasciculus V. Erlangae: Joannis Jacobi Palm, pp. 81–112, pls. XVII, XXI–XXV.

Schoepff, J.D. 1801. Historia Testudinum Iconibus Illustrata. Fasciculus V. Erlangae: Joannis Jacobi Palm, pp. 113–136, pls. XXVI–XXXI.

Schreiber, E. 1875. Herpetologia Europaea: eine systematische Bearbeitung der Amphibien und Reptilien welche bisher in Europa aufgefunden sind. Braunschweig: F. Vieweg und Sohn, 639 pp.

Schreiber, E. 1912. Herpetologia Europaea: eine systematische Bearbeitung der Amphibien und Reptilien welche bisher in Europa aufgefunden sind. Jena: Gustav Fischer, 960 pp.

Schuchert, C. 1897. A synopsis of American fossil Brachiopoda, including bibliography and synonymy. Bulletin of the U.S. Geological Survey No. 87, 464 pp.

Schwartz, A. 1955. The diamondback terrapins (*Malaclemys terrapin*) of peninsular Florida. Proceedings of the Biological Society of Washington 68:157–164.

Schwartz, A. 1956a. Geographic variation in the chicken turtle *Deirochelys reticularia* Latreille. Fieldiana Zoology 34:461–503.

Schwartz, A. 1956b. The relationships and nomenclature of the soft-shelled turtles (genus *Trionyx*) of the southeastern United States. Charleston Museum Leaflets 26:1–21.

Schweiger, M. and Gemel, R. 2020. Where do you come from, stranger? A scientific-historical digression with discussion on nomenclature and taxonomy of *Testudo graeca* Linnaeus, 1758. Herpetozoa 33:31–38.

Schweigger, A.F. 1812. Prodromus monographiae Cheloniorum. Königsberger Archiv für Naturwissenschaft und Mathematik 1:271–368, 406–462.

Sclater, P.L. 1858. Bibliographical notice. Contributions to the Natural History of the United States of America. By Louis Agassiz. Annals and Magazine of Natural History (3)1:289–294.

Sclater, P.L. 1870. Remarks on the animals lately described by Dr. Gray as *Testudo chilensis* and *Ateles bartlettii*. Annals and Magazine of Natural History (4)6:470–473.

Scott, P.A. and Rissler, L.J. 2015. Integrating dynamic occupancy modeling and genetics to infer the status of the imperiled flattened musk turtle. Biological Conservation 192:294–303.

Scott, P.A., Glenn, T.C., and Rissler, L.J. 2018. Resolving taxonomic turbulence and uncovering cryptic diversity in the musk turtles (*Sternotherus*) using robust demographic modeling. Molecular Phylogenetics and Evolution 120:1–15.

Scott, P.A., Allison, L.J., Field, K.J., Averill-Murray, R.C., and Shaffer, H.B. 2020. Individual heterozygosity predicts translocation success in threatened desert tortoises. Science 370(6520):1086–1089.

Seateun, S., Karraker, N.E., Stuart, B.L., and Aowphol, A. 2019. Population demography of Oldham's leaf turtle (*Cyclemys oldhamii*) in protected and disturbed habitats in Thailand. PeerJ 7:e7196.

Seba, A. 1734. Locupletissimi Rerum Naturalium Thesauri Accurata Descriptio, et Iconibus Artificiosissimis Expressio, per Universam Physices Historiam. Tomus I. Amstelaedami [Amsterdam]: J. Wetstenium, Gul. Smith, and Janssonio Waesbergios, 178 pp.

Seddon, J., Georges, A., Baverstock, P., and McCord, W. 1997. Phylogenetic relationships of chelid turtles (Pleurodira: Chelidae) based on mitochondrial 12S rRNA gene sequence variation. Molecular Phylogenetics and Evolution 7:55–61.

Seeliger, L.M. 1945. Variation in the Pacific mud turtle. Copeia 1945(3):150–159.

Segniagbeto, G.H., Afiademagmo, K., Akani, G.C., Petrozzi, F., and Luiselli, L. 2015. Sex-ratio, size structure and morphometrics of turtle populations from Togo, West Africa. Herpetozoa 28:29–38.

Seligmann, J.M. 1764. Die See Schild-kröte. In: Seligmann, J.M. Sammlung verschiedener ausländischer und seltener Vögel, und eineger anderer Seltenheiten der Natur, worinnen ein jeder derselben nicht nur auf das genaueste beschreiben, sondern auch in einer richtigen und sauber illuminirten Abbildungen. Sechster Theil. Nürnberg: J.J. Fleischmann, pl. CI.

Seligmann, J.M. 1773. Die kleine Morast-Schildkröte. In: Seligmann, J.M. Sammlung verschiedener ausländischer und seltener Vögel, und eineger anderer Seltenheiten der Natur, in richtigen und sauber illuminirten Abbildungen. Achter Theil. Nürnberg: Seligmanns Verlegern, pl. LXXVII.

Seidel, M.E. 1988. Revision of the West Indian emydid turtles (Testudines). American Museum Novitates 2918:1–41.

Seidel, M.E. 1989. *Trachemys dorbigni*. Catalogue of American Amphibians and Reptiles 486:1–3.

Seidel, M.E. 1994. Morphometric analysis and taxonomy of cooter and red-bellied turtles in the North American genus *Pseudemys* (Emydidae). Chelonian Conservation and Biology 1(2):117–130.

Seidel, M.E. 1995. How many species of cooter turtles and where is the scientific evidence? – A reply to Jackson. Chelonian Conservation and Biology 1(4):333–336.

Seidel, M.E. 1996. Current status of biogeography of the West Indian turtles in the genus *Trachemys* (Emydidae). In: Powell, R. and Henderson, R.W. (Eds.). Contributions to West Indian Herpetology: A Tribute to Albert Schwartz. SSAR Contributions to Herpetology 12:169–174.

Seidel, M.E. 2002. Taxonomic observations on extant species and subspecies of slider turtles, genus *Trachemys*. Journal of Herpetology 36:285–292.

Seidel, M.E. 2010. *Trachemys nebulosa*. Catalogue of American Amphibians and Reptiles 870:1–5.

Seidel, M.E. and Ernst, C.H. 2017. A systematic review of the turtle family Emydidae. Vertebrate Zoology 67(1):1–122.

Seidel, M.E. and Inchaustegui Miranda, S.J. 1984. Status of the trachemyd turtles (Testudines: Emydidae) on Hispaniola. Journal of Herpetology 18(4):468–479.

Seidel, M.E., Iverson, J.B., and Adkins, M.D. 1986. Biochemical comparisons and phylogenetic relationships in the family Kinosternidae (Testudines). Copeia 1986(2):285–294.

Selman, W. 2017. Diagnostic trait variability in the imperiled freshwater turtle, *Graptemys flavimaculata*. Herpetologica 73(2):105–112.

Selman, W. 2024. Distribution and abundance of *Graptemys oculifera* (Ringed Sawback) and *Graptemys pearlensis* (Pearl Map Turtle) in the Pearl River system of Louisiana, USA. Herpetological Conservation and Biology 19(1):129–143.

Selman, W., Kreiser, B., and Qualls, C. 2013. Conservation genetics of the yellow-blotched sawback *Graptemys flavimaculata* (Testudines: Emydidae). Conservation Genetics 14:1193–1203.

Seminoff, J.A., Allen, C.D., Balazs, G.H., Dutton, P.H., Eguchi, T., Haas, H.L., Hargrove, S.A., Jensen, M.P., Klemm, D.L., Lauritsen, A.M., MacPherson, S.L., Opay, P., Possardt, E.E., Pultz, S.L., Seney, E.E., Van Houtan, K.S., and Waples, R.S. 2015. Status review of the Green Turtle (*Chelonia mydas*) under the U.S. Endangered Species Act. NOAA Technical Memorandum, NOAA NMFS-SWFSC-539, 571 pp.

Senneke, D. 2006. Husbandry and breeding of *Indotestudo elongata* in a temperate environment. In: Artner, H., Farkas, B., and Loehr, V. (Eds.). Turtles: Proceedings: International Turtle and Tortoise Symposium Vienna 2002. Frankfurt: Edition Chimaira, pp. 96–101.

Serb, J.M., Phillips, C.A., and Iverson, J.B. 2001. Molecular phylogeny and biogeography of *Kinosternon flavescens* based on complete mitochondrial control region sequences. Molecular Phylogenetics and Evolution 18:149–162.

Sethuraman, A., McGaugh, S.E., Becker, M.L., Chandler, C.H., Christiansen, J.L., Hayden, S., LeClere, A., Monson-Miller, J., Myers, E.M., Paitz, R.T., Refsnider, J.M., VanDeWalle, T.J., and Janzen, F.J. 2014. Population genetics of Blanding's turtle (*Emys blandingii*) in the midwestern United States. Conservation Genetics 15:61–73.

Sethy, C., Samantasinghar, L., and Pramanik, D.S. 2015. Occurrence and habitat type of seven non marine chelonian species in some parts of Odisha, India. European Journal of Experimental Biology 5(10):6–14.

Seymour Sewell, R.B. and Guha, B.S. 1931. Zoological remains. In: Marshall, J. (Ed.). Mohenjo-Daro and the Indus Civilization, Volume II. London: Arthur Probsthain, pp. 649–673.

Shaffer, H.B., Meylan, P., and McKnight, M.L. 1997. Test of turtle phylogeny: molecular, morphological, and paleontological approaches. Systematic Biology 46:235–268.

Shaffer, H.B., Starkey, D.E., and Fujita, M.K. 2008. Molecular insights into the systematics of the snapping turtles (Chelydridae). In: Steyermark, A.C., Finkler, M.S., and Brooks, R.J. (Eds.). Biology of the Snapping Turtle (*Chelydra serpentina*). Baltimore: Johns Hopkins University Press, pp. 44–49.

Shamblin, B.M., Bjorndal, K.A., Bolten, A.B., Hillis-Starr, Z.M., Lundgren, I., Naro-Maciel, E., and Nairn, C.J. 2012. Mitogenomic sequences better resolve stock structure of southern Greater Caribbean green turtle rookeries. Molecular Ecology 21:2330–2340.

Shamblin, B.M, Bolten, A.B., Abreu-Grobois, F.A., Bjorndal, K.A., Cardona, L., Carreras, C., Clusa, M., Monzón-Argüello, C., Nairn, C.J., Nielsen, J.T., Nel, R., Soares, L.S., Stewart, K.R., Vilaça, S.T., Türkozan, O., Yilmaz, C., and Dutton, P.H. 2014. Geographic patterns of genetic variation in a broadly distributed marine vertebrate: new insights into the Loggerhead Turtle stock structure from expanded mitochondrial DNA sequences. PLoS One 9:e85956.

Shamblin, B.M., Godfrey, M.H., Pate, S.M., Thompson, W.P., Sutton, H., Altman, J., Fair, K., McClay, J., Wilson, A.M., Milligan, B., Stetzar, E.J., and Nairn, C.J. 2018. Green Turtles nesting at their northern range limit in the United States represent a distinct subpopulation. Chelonian Conservation and Biology 17:314–319.

Shammakov, S. 1981. [Reptiles of the Plains of Turkmenistan.] Ashgabat: Ylym, 308 pp. [in Russian]

Shaw, G. 1793. Naturalist's Miscellany. Vol. 4. London: Frederick P. Nodder, 156 pp.

Shaw, G. 1794. Zoology of New Holland. Vol. I. London: J. Davis, 33 pp.

Shaw, G. 1802. General Zoology, or Systematic Natural History. Volume III, Part I, Amphibia. London: G. Kearsley, 312 pp.

Shaw, G. and Nodder, E. 1810. *Testudo nilotica*. Naturalist's Miscellany 21:179–184.

Shea, G.M. and Sadlier, R.A. 1999. A catalogue of the non-fossil amphibian and reptile type specimens in the collection of the Australian Museum: types currently, previously and purportedly present. Technical Reports of the Australian Museum 15:1–91.

Shea, G., Thomson, S., and Georges, A. 2020. The identity of *Chelodina oblonga* Gray 1841 (Testudines: Chelidae) reassessed. Zootaxa 4779(3):419–437.

Shepherd, L. 2023. Taxonomic vandalism: discovering new species is hard, but only if you're a decent scientist. https://abitofbio.medium.com/

Sherborn, C.D. and Woodward, A.S. 1901. Dates of publication of the zoological and botanical portions of some French voyages. Annals and Magazine of

Natural History (7)8:161–164, 333–336.

Shi, H., Parham, J.F., Simison, W.B., Wang, J., Gong, S., and Fu, B. 2005. A report on the hybridization between two species of threatened Asian box turtles (Testudines: *Cuora*) in the wild on Hainan Island (China) with comments on the origin of '*serrata*'-like turtles. Amphibia-Reptilia 26:377–381.

Shi, H., Fong, J.J., Parham, J.F., Pang, J., Wang, J., Hong, M., and Zhang, Y.-P. 2008. Mitochondrial variation of the "eyed" turtles (*Sacalia*) based on known-locality and trade specimens. Molecular Phylogenetics and Evolution 49:1025–1029.

Shoemaker, K.T. and Gibbs, J.P. 2013. Genetic connectivity among populations of the threatened bog turtle (*Glyptemys muhlenbergii*) and the need for a regional approach to turtle conservation. Copeia 2013(2):324–331.

Shrestha, T.K. 1997. Status, biology, conservation, and management of tortoises and turtles in the Himalayan foothills of Nepal. In: Van Abbema, J. (Ed.). Proceedings: Conservation, Restoration, and Management of Tortoises and Turtles – An International Conference. N.Y. Turtle and Tortoise Society, pp. 278–286.

Shufeldt, R.W. 1919. Observation on the chelonians of North America. IV. Aquatic Life 1919["1918(September)"]:155–157.

Siciliano, S., de Moura, J.F., Tavares, D.C., Silva de Amorim, C.E., and Matias, C.A.R. 2014. On the intriguing occurrence of *Rhinoclemmys punctularia* (Daudin, 1801) in coastal plains of eastern Rio de Janeiro, Brazil. Herpetology Notes 7:667–671.

Siders, Z.A., Stratmann, T.A., Turner Tomaszewicz, C.N., Walde, A.D., and Munscher, E.C. 2023. Somatic growth and maturity for four species of River Cooter including *Pseudemys concinna suwanniensis, P. nelsoni, P. peninsularis*, and *P. texana*. Biology 12:965, 19 pp. doi.org/10.3390/biology12.

Siebold, P.F. de. 1826. De Historiae Naturalis in Japonia Statu, nec non de augmento emolumentisque in decursu perscrutationum expectandis. Dissertatio, cui accedunt Spicilegia Faunae Japonicae. Wirceburgi: Car. Phil. Bonitas, 20 pp.

Siebenrock, F. 1901. Beschreibung einer neuen Schildkrötengattung aus der Familie Chelydidae von Australien: *Pseudemydura*. Anzeiger der Kaiserlichen Akademie der Wissenschaften in Wien (Mathemathisch-Naturwissenschaftliche Klasse) 38(22):248–250.

Siebenrock, F. 1902a. Eine neue Schildkröte aus Madagascar (nach Gerrard). Zoologischer Anzeiger 25(1901)[1902]:6–8.

Siebenrock, F. 1902b. Uber zwei seltene Schildkröten der herpetologischen Sammlung des Wiener Museums. Anzeiger der Kaiserlichen Akademie der Wissenschaften in Wien (Mathemathisch-Naturwissenschaftliche Klasse) 1902(2):11–13.

Siebenrock, F. 1902c. Zur Systematik der Schildkrötenfamilie Trionychidae Bell, nebst der Beschreibung einer neuen *Cyclanorbis* Art. Sitzungsberichte der Kaiserlichen Akademie der Wissenschaften in Wien (Mathemathisch-Naturwissenschaftliche Klasse) 91:807–846.

Siebenrock, F. 1903a. Schildkröten des östlichen Hinterindien. Sitzungsberichte der Kaiserlichen Akademie der Wissenschaften in Wien (Mathemathisch-Naturwissenschaftliche Klasse) 112:333–353.

Siebenrock, F. 1903b. Über zwei seltene und eine neue Schildkröte des Berliner Museums. Sitzungsberichte der Kaiserlichen Akademie der Wissenschaften in Wien (Mathemathisch-Naturwissenschaftliche Klasse) 112:439–445.

Siebenrock, F. 1904a. Eine neue *Testudo*-Art der *geometrica*-Gruppe aus Süd-Afrika. Anzeiger der Kaiserlichen Akademie der Wissenschaften in Wien (Mathemathisch-Naturwissenschaftliche Klasse) 41:194–195.

Siebenrock, F. 1904b. Schildkröten von Brasilien. Denkschriften der Kaiserlichen Akademie der Wissenschaften in Wien (Mathemathisch-Naturwissenschaftliche Klasse) 76:1–28.

Siebenrock, F. 1906a. Zur Kenntnis der Schildkrötenfauna der Insel Hainan. Zoologischer Anzeiger 30:578–586.

Siebenrock, F. 1906b. Eine neue *Cinosternum*-Art aus Florida. Zoologischer Anzeiger 30:727–728.

Siebenrock, F. 1906c. Schildkröten von Ostafrika und Madagaskar. In: Voeltzkow, A. Reise in Ost-Afrika in den Jahren 1903–1905 mit Mitteln der Hermann und Elise geb. Heckmann-Wentzel-Stiftung. Wissenschaftliche Ergebnisse, Systematischen Arbeiten, Stuttgart 2:1–40.

Siebenrock, F. 1907. Die Schildkrötenfamilie Cinosternidae m. Monographisch bearbeitet. Sitzungsberichte der Kaiserlichen Akademie der Wissenschaften in Wien (Mathemathisch-Naturwissenschaftliche Klasse) (1)116:527–599.

Siebenrock, F. 1909a. Synopsis der rezenten Schildkröten, mit Berücksichtigung der in historischer Zeit ausgestorbenen Arten. Zoologische Jahrbücher Supplement 10(3):427–618.

Siebenrock, F. 1909b. Über die Berechtigung der Selbständigkeit von *Sternotherus nigricans seychellensis* Siebenrock. Zoologischer Anzeiger 34:359–362.

Siebenrock, F. 1914. Eine neue *Chelodina* Art aus Westaustralien. Anzeiger der Kaiserlichen Akademie der Wissenschaften in Wien (Mathemathisch-Naturwissenschaftliche Klasse) 51(17):386–387.

Silva, A., Rocha, S., Gerlach, J., Rocamora, G., Dufrenne, A., and Harris, D.J. 2010. Assessment of mtDNA genetic diversity within the terrapins *Pelusios subniger* and *Pelusios castanoides* across the Seychelles islands. Amphibia-Reptilia 31:583–588.

Simison, W.B., Sellas, A.B., Feldheim, K.A., and Parham, J.F. 2013. Isolation and characterization of microsatellite markers for identifying hybridization and genetic pollution associated with red-eared slider turtles (*Trachemys scripta elegans*). Conservation Genetics Resources 5(4):1139–1140.

Simpson, G.G. 1937. New reptiles from the Eocene of South America. American Museum Novitates 927:1–3.

Široký, P. and Fritz, U. 2007. Is *Testudo werneri* a distinct species? Biologia (Bratislava) Section Zoology 62(2):1–4.

Široký, P., Stuchlik, S., and Moravec, J. 2004. Current situation and Pleistocene, Holocene, and historic records of *Emys orbicularis* in the Czech Republic. In: Fritz, U. and Havas, P. (Eds.). Proceedings of the 3rd International Symposium on *Emys orbicularis*. Biologia (Bratislava) 59(Suppl. 14):73–78.

Slevin, J.R. and Leviton, A.E. 1956. Holotype specimens of reptiles and amphibians in the collection of the California Academy of Sciences. Proceedings of the California Academy of Sciences (4)28:529–560.

Smales, I., McCord, W.P., Cann, J., and Joseph-Ouni, M. 2019a. New Guinean *Emydura* (Testudines: Pleurodira: Chelidae) with the description of a new species. The Batagur Monographs 2:7–51.

Smales, I., Sadlier, R., and Campbell, P.D. 2019b. *Elseya latisternum* Gray 1867 (Testudines: Chelidae): the identity of type specimens and a review of nineteenth century specimens in the Natural History Museum, London. Herpetological Review 50(4):699–708.

Smith, A. 1838. Illustrations of the Zoology of South Africa; consisting chiefly of Figures and Descriptions of the Objects of Natural History collected during an Expedition into the Interior of South Africa, in the years 1834, 1835, and 1836. Part No. 1. London: Smith, Elder and Co., 23 pp., 10 pls. [Reptilia, plate 1].

Smith, A. 1839a. Illustrations of the Zoology of South Africa; consisting chiefly of Figures and Descriptions of the Objects of Natural History collected during an Expedition into the Interior of South Africa, in the years 1834, 1835, and 1836. Part No. 6. London: Smith, Elder and Co., 23 pp., 10 pls. [Reptilia, plate 6].

Smith, A. 1839b. Illustrations of the Zoology of South Africa; consisting chiefly of Figures and Descriptions of the Objects of Natural History collected during an Expedition into the Interior of South Africa, in the years 1834, 1835, and 1836. Part No. 8. London: Smith, Elder and Co., 23 pp., 10 pls. [Reptilia, plate 8].

Smith, H., Galicki, S., and Selman, W. 2020. Three's company: observations of a nonnative Map Turtle (*Graptemys pseudogeographica*) occurring syntopically with two endemic *Graptemys* in the Pearl River, Mississippi. Chelonian Conservation and Biology 19(2):268–276.

Smith, H.M. and Brandon, R.A. 1968. Data nova herpetologica Mexicana. Transactions of the Kansas Academy of Science 71:49–61.

Smith, H.M. and Chiszar, D. 2006. Dilemma of name-recognition: why and when to use new combinations of scientific names. Herpetological Conservation and Biology 1(1):6–8.

Smith, H.M. and Glass, B.P. 1947. A new musk turtle from southeastern United States. Journal of the Washington Academy of Sciences 37:22–24.

Smith, H.M. and Ramsey, L.W. 1952. A new turtle from Texas. Wasmann Journal of Biology 10(1):45–54.

Smith, H.M. and Rhodin, A.G.J. 1986. Authorship of the scientific name of the leatherback sea turtle. Journal of Herpetology 20(3):450–451.

Smith, H.M. and Smith, R.B. 1980 ["1979"]. Synopsis of the Herpetofauna of Mexico, Vol. VI. Guide to Mexican Turtles. Bibliographic Addendum III. North Bennington, Vermont: John Johnson, 1044 pp.

Smith, H.M. and Taylor, E.H. 1950a. Type localities of Mexican reptiles and amphibians. University of Kansas Science Bulletin 33(8):313–380.

Smith, H.M. and Taylor, E.H. 1950b. An annotated checklist and key to the reptiles of Mexico exclusive of the snakes. United States National Museum Bulletin 199:1–253.

Smith, H.M., Langebartel, D.A., and Williams, K.L. 1964. Herpetological type-specimens in the University of Illinois Museum of Natural History. Illinois Biological Monographs 32:1–80.

Smith, H.M., Chiszar, D., and Smith, R.B. 1980a. Rafinesque, 1822, "On the turtles of the United States" (Reptilia, Testudines): proposed suppression by use of the plenary powers. Bulletin of Zoological Nomenclature 37:53–56.

Smith, H.M., Smith, R.B., and Chiszar, D. 1980b. *Sternotherus* Gray, 1825, correct spelling; and *Pelusios* Wagler, 1830, proposed conservation (Reptilia, Testudines). Bulletin of Zoological Nomenclature 37:124–128.

Smith, H.M., Ernst, C.H., and Smith, R.B. 1980c. *Geoemyda* Gray, 1834, and *Rhinoclemmys* Fitzinger, 1835 (Reptilia, Testudines): proposed conservation. Bulletin of Zoological Nomenclature 37:233–239.

Smith, H.M., Smith, R.B., and Chiszar, D. 1983. *Chelydra osceola* Stejneger, 1918 (Reptilia, Testudines): proposed conservation by use of the Plenary Powers. Bulletin of Zoological Nomenclature 40:225–227.

Smith, H.M., Humphrey, R., and Chiszar, D. 1996. A range extension for the box turtle *Terrapene yucatana*. Bulletin Maryland Herpetological Society 32:14–15.

Smith, M.A. 1916. A list of the crocodiles, tortoises, turtles and lizards at present known to inhabit Siam. Journal of the Natural History Society of Siam 2(1):48–57.

Smith, M.A. 1931. The Fauna of British India, including Ceylon and Burma. Reptilia and Amphibia. Vol. I. Loricata, Testudines. London: Taylor and Francis, 185 pp.

Smith, P.W. 1951. A new frog and a new turtle from the western Illinois sand prairies. Bulletin of the Chicago Academy of Sciences 9(10):189–199.

Somerová, B., Rehák, I., Velensky, P., Palupcíkova, K., Protiva, T., and Frynta, D. 2015. Haplotype variation in founders of the *Mauremys annamensis* population kept in European zoos. Acta Herpetologica 10:7–15.

Sommer, R.S., Lindqvist, C., Persson, A., Bringsøe, H., Rhodin, A.G.J., Schneeweiss, N., Siroky, P., Bachmann, L., and Fritz, U. 2009. Unexpected early extinction of the European pond turtle (*Emys orbicularis*) in Sweden and climatic impact on its Holocene range. Molecular Ecology 18:1252–1262.

Song, M.-T. 1984. [A new species of the turtle genus *Cuora* (Testudoformes: Testudinidae)]. Acta Zootaxonomica Sinica 9(3):330–332. [in Chinese]

Song, M.-T. 2001. Translation: A new species of the turtle genus *Cuora* (Testudoformes: Testudinidae). Asiatic Herpetological Research 9:142–144.

Sonnini, C.S. and Latreille, P.A. 1801. Histoire Naturelle des Reptiles, avec figures dessinées d'après nature. Tome Premier. Première Partie. Quadrupèdes et Bipèdes Ovipares. Paris: Deterville, 280 pp.

Sousa, B.M.D., Nascimento, A.E.R.D., Gomides, S.C., Rios, C.H.V., Hudson, A.D.A., and Novelli, I.A. 2010. Reptiles in fragments of Cerrado and Atlantic Forest at the Campo das Vertentes, Minas Gerais State, southeastern Brazil. Biota Neotropica 10(2):129–138.

Souza, F.L. and Abe, A.S. 1997. Population structure, activity, and conservation of the neotropical freshwater turtle, *Hydromedusa maximiliani*, in Brazil. Chelonian Conservation and Biology 2(4):521–525.

Souza, F.L. and Abe, A.S. 2001. Population structure and reproductive aspects of the freshwater turtle, *Phrynops geoffroanus*, inhabiting an urban river in southeastern Brazil. Studies on Neotropical Fauna and Environment 36:57–62.

Souza, F.L., Rodrigues, J., Olalla-Tárraga, M., Diniz-Filho, J., Martinez, P., and Sawaya, R.J. 2019. Niche divergence and diversification in South American freshwater turtles of the genus *Acanthochelys* (Chelidae). Amphibia-Reptilia 40:475–485.

Spawls, S., Howell, K., Drewes, R., and Ashe, J. 2002. A Field Guide to the Reptiles of East Africa: Kenya, Tanzania, Uganda, Rwanda and Burundi. London and San Diego: Academic Press, 543 pp.

Spencer, R. 2019. Book review: *Freshwater Turtles of Australia*. John Cann and Ross Sadlier. 2017. Australian Zoologist 40(2):326–327.

Spencer, R.J., Georges, A., Lim, D., Welsh, M., Reid, A.M. 2014. The risk of inter-specific competition in Australian short-necked turtles. Ecological Research 29(4):767–777.

Speybroeck, J., Beukema, W., Dufresnes, C., Fritz, U., Jablonski, D., Lymberakis, P., Martínez-Solano, I., Razzetti, E., Vamberger, M., Vences, M., Vörös, J., and Crochet, P.-A. 2020. Species list of the European herpetofauna – 2020 update by the Taxonomic Committee of the Societas Europaea Herpetologica. Amphibia-Reptilia 41(2):139–189.

Spinks, P.Q. and Shaffer, H.B. 2005. Range-wide molecular analysis of the western pond turtle (*Emys marmorata*): cryptic variation, isolation by distance, and their conservation implications. Molecular Ecology 14:2047–2064.

Spinks, P.Q. and Shaffer, H.B. 2007. Conservation phylogenetics of the Asian box turtles (Geoemydidae, *Cuora*): mitochondrial introgression, numts, and inferences from multiple nuclear loci. Conservation Genetics 8:641–657.

Spinks, P.Q. and Shaffer, H.B. 2009. Conflicting mitochondrial and nuclear phylogenies for the widely disjunct *Emys* (Testudines: Emydidae) species complex, and what they tell us about biogeography and hybridization. Systematic Biology 58(1):1–20.

Spinks, P.Q., Shaffer, H.B., Iverson, J.B., and McCord, W.P. 2004. Phylogenetic hypotheses for the turtle family Geoemydidae. Molecular Phylogenetics and Evolution 32:164–182.

Spinks, P.Q., Thomson, R.C., and Shaffer, H.B. 2009a. A reassessment of *Cuora cyclornata* Blanck, McCord and Le, 2006 (Testudines, Geoemydidae) and a plea for taxonomic stability. Zootaxa 2018:58–68.

Spinks, P.Q., Thomson, R.C., Lovely, G.A., and Shaffer, H.B. 2009b. Assessing what is needed to resolve a molecular phylogeny: simulations and empirical data from emydid turtles. BMC Evolutionary Biology 9:56.

Spinks, P.Q., Thomson, R.C., and Shaffer, H.B. 2010. Nuclear gene phylogeography reveals the historical legacy of an ancient inland sea on lineages of the western pond turtle, in California. Molecular Ecology 19:542–556.

Spinks, P.Q., Thomson, R.C., Zhang, Y., Che, J., Wu, Y., and Shaffer, H.B. 2012a. Species boundaries and phylogenetic relationships in the critically endangered Asian box turtle genus *Cuora*. Molecular Phylogenetics and Evolution 63(3):656–667.

Spinks, P.Q., Thomson, R.C., Hughes, B., Moxley, B., Brown, R., Diesmos, A., and Shaffer, H.B. 2012b. Cryptic variation and the tragedy of unrecognized taxa: the case of international trade in the spiny turtle *Heosemys spinosa* (Testudines: Geoemydidae). Zoological Journal of the Linnean Society 164:811–824.

Spinks, P.Q., Thomson, R.C., Pauly, G.B., Newman, C.E., Mount, G., and Shaffer, H.B. 2013. Misleading phylogenetic inferences based on single-exemplar sampling in the turtle genus *Pseudemys*. Molecular Phylogenetics and Evolution 68:269–281.

Spinks, P.Q., Thomson, R.C., and Shaffer, H.B. 2014a. The advantages of going large: genome-wide SNPs clarify the complex population history and systematics of the threatened western pond turtle. Molecular Ecology 23:2228–2241.

Spinks, P.Q., Thomson, R.C., Gidis, M., and Shaffer, H.B. 2014b. Multilocus phylogeny of the New-World mud turtles (Kinosternidae) supports the traditional classification of the group. Molecular Phylogenetics and Evolution 76:254–260.

Spinks, P.Q., Georges, A., and Shaffer, H.B. 2015. Phylogenetic uncertainty and taxonomic re-revisions: an example from the Australian Short-necked Turtles (Testudines: Chelidae). Copeia 103(3):536–540.

Spinks, P.Q., Thomson, R.C., McCartney-Melstad, E., and Shaffer, H.B. 2016. Phylogeny and temporal diversification of the New World pond turtles (Emydidae). Molecular Phylogenetics and Evolution 103:85–97.

Spitzweg, C., Hofmeyr, M.D., Fritz, U., and Vamberger, M. 2019. Leopard tortoises in southern Africa have greater genetic diversity in the north than in the south (Testudinidae). Zoologica Scripta 48:57–68.

Spitzweg, C., Vamberger, M., Ihlow, F., Fritz, U., and Hofmyer, M.D. 2020. How many species of Angulate Tortoises occur in southern Africa? (Testudines: Testudinidae: *Chersina*). Zoologica Scripta 49:412–426.

Spix, J.B. 1824. Animalia Nova sive Species Novae Testudinum et Ranarum. Monachii: 53 pp.

Squire, E.G. 1870. Honduras: Descriptive, Historical, and Statistical. London: Trübner and Co., 268 pp.

Srinivasulu, C. and Kumar, G.C. 2022. A checklist of herpetofauna of Telangana state, India. Journal of Threatened Taxa 14(6):21266–21281.

Starkey, D.E., Shaffer, H.B., Burke, R.L., Forstner, M.R.J., Iverson, J.B., Janzen, F.J., Rhodin, A.G.J., and Ultsch, G.R. 2003. Molecular systematics, phylogeography, and the effects of Pleistocene glaciation in the painted turtle (*Chrysemys picta*) complex. Evolution 57(1):119–128.

Steadman, D.W., Albury, N.A., Maillis, P., Mead, J.I., Slapcinsky, J., Krysko, K.L., Singleton, H.M., and Franklin, J. 2014. Late-Holocene faunal and landscape change in the Bahamas. The Holocene 24(2):220–230.

Steadman, D.W., Albury, N.A., Carlson, L.A., Franz, R., LeFebvre, M.J., Kakuk, B., and Keegan, W.F. 2020. The paleoecology and extinction of endemic tortoises in the Bahamian Archipelago. The Holocene 30(3):420–427.

Stebbins, R.C. 2003. A Field Guide to Western Reptiles and Amphibians. Third Edition. Boston: Houghton Mifflin Co., 533 pp.

Stejneger, L. 1902. Some generic names of turtles. Proceedings of the Biological Society of Washington 15:235–238.

Stejneger, L. 1907. Herpetology of Japan and adjacent territory. Smithsonian Institution United States National Museum Bulletin 58:1–577.

Stejneger, L. 1909. Generic names of some Chelyid turtles. Proceedings of the

Biological Society of Washington 22:125–127.

Stejneger, L. 1918. Description of a new lizard and a new snapping turtle from Florida. Proceedings of the Biological Society of Washington 31:89–92.

Stejneger, L. 1925. New species and subspecies of American turtles. Journal of the Washington Academy of Science 15:462–463.

Stejneger, L. 1933. Description of a new box turtle from Mexico. Proceedings of the Biological Society of Washington 46:119–120.

Stejneger, L. 1938. Restitution of the name *Ptychemys hoyi* Agassiz for a western river tortoise. Proceedings of the Biological Society of Washington 51:173–175.

Stejneger, L. 1941. Notes on Mexican turtles of the genus *Kinosternon*. Proceedings of the United States National Museum 90:457–459.

Stejneger, L. 1944. Notes on the American soft-shelled turtles with special reference to *Amyda agassizii*. Bulletin of the Museum of Comparative Zoology 94(1):1–75.

Stejneger, L. and Barbour, T. 1917. A Checklist of North American Amphibians and Reptiles. Cambridge: Harvard University Press, 125 pp.

Stephens, P.R. 1998. Variation in the cranial osteological morphology of turtles in the genus *Graptemys* (Reptilia; Anapsida; Testudines; Cryptodira; Emydidae; Deirochelyinae). Masters Thesis, University of South Alabama, Mobile.

Stephens, P.R. and Wiens, J.J. 2003. Ecological diversification and phylogeny of emydid turtles. Biological Journal of the Linnean Society 79:577–610.

Stephens, P.R. and Wiens, J.J. 2009. Evolution of sexual size dimorphisms in emydid turtles: ecological dimorphism, Rensch's Rule, and sympatric divergence. Evolution 63:910–925.

Stephenson, P.J., Grace, M.K., Akçakaya, H.R., Rodrigues, A.S.L., Long, B., Mallon, D.P., Meijaard, E., Rodriguez, J.P., Young, R.P., Brooks, T.M., and Hilton-Taylor, C. 2019. Defining the indigenous ranges of species to account for geographic and taxonomic variation in the history of human impacts: reply to Sanderson 2019. Conservation Biology 33(5):1211–1213.

Sterli, J. 2010. Phylogenetic relationships among extinct and extant turtles: the position of Pleurodira and the effects of the fossils on rooting crown-group turtles. Contributions to Zoology 79 (3): 93–106.

Sterli, J. 2015. A review of the fossil record of Gondwanan turtles of the clade Meiolaniformes. Bulletin of the Peabody Museum of Natural History 56(1):21–45.

Sterli, J. and de la Fuente, M.S. 2013. New evidence from the Palaeocene of Patagonia (Argentina) on the evolution and palaeo-biogeography of Meiolaniformes (Testudinata, new taxon name). Journal of Systematic Palaeontology 11(7):835–852.

Stobaeus, K. 1730. Descriptio Testudinis americanae terrestris; forte Jaboti Brasiliensibus, cagado de terra Lusitanis dictae, Marcgravii. Acta Literaria et Scientarum Sveciae 3:58–62.

Storr, G. 1978. Taxonomic notes on the reptiles of the Shark Bay Region, Western Australia. Records of the Western Australia Museum. 6(3):303–318.

Strain, W.S. 1966. Blancan mammalian fauna and Pleistocene formations, Hudspeth County, Texas. Bulletin of the Texas Memorial Museum 10:1–55.

Strauch, A. 1862. Chelonologische studien, mit besonderer Beziehung auf die Schildkrötensammlung der kaiserlichen Akademie der Wissenschaften zu St. Petersburg. Mémoires de l'Académie Impériale des Sciences de St.-Pétersbourg (7)5(7):1–196.

Strauch, A. 1865. Die Vertheilung der Schildkröten über den Erdball. Ein zoogeographischer Versuch. Mémoires de l'Académie Impériale des Sciences de St.-Pétersbourg (7)8(13):1–207.

Strauch, A. 1890. Bemerkungen über die Schildkrötensammlung im Zoologischen Museum der kaiserlichen Akademie der Wissenschaften zu St. Petersburg. Mémoires de l'Académie Impériale des Sciences de St.-Pétersbourg (7)38(2):1–127.

Struijk, R.P.J.H., McCormack, T.E.M., Nguyen, T.T., Pham, T.V., Stumpel, J.B.G., Wang, J., and Auer, M. 2016. Intergradation between *Cuora mouhotii mouhotii* (Gray, 1862) and *Cuora mouhotii obsti* Fritz, Andreas & Lehr, 1998 with notes on the species' geographical distribution and phenotypic variation. Sauria 38(1):31–47.

Stuart, B.L. and Fritz, U. 2008. Historical DNA from museum type specimens clarifies diversity of Asian leaf turtles (*Cyclemys*). Biological Journal of the Linnaean Society 94:131–141.

Stuart, B.L. and Parham, J.F. 2004. Molecular phylogeny of the critically endangered Indochinese box turtle (*Cuora galbinifrons*). Molecular Phylogenetics and Evolution 31:164–177.

Stuart, B.L. and Parham, J.F. 2007. Recent hybrid origin of three rare Chinese turtles. Conservation Genetics 8:169–175.

Stuart, B.L. and Platt, S.G. 2004. Recent records of turtles and tortoises from Laos, Cambodia, and Vietnam. Asiatic Herpetological Research 10:129–150.

Stuart, B.L. and Thorbjarnarson, J. 2003. Biological prioritization of Asian countries for turtle conservation. Chelonian Conservation and Biology 4(3):642–647.

Stuart, B.L., Hallam, C.D., Sayavong, S., Nanthavong, C., Sayaleng, S., Vongsa, O., and Robichaud, W.G. 2011. Two additions to the turtle fauna of Laos. Chelonian Conservation and Biology 10(1):113–116.

Stuart, L.C. 1963. A checklist of the herpetofauna of Guatemala. Miscellaneous Publications, Museum of Zoology, University of Michigan 122:1–150.

Stuckas, H. and Fritz, U. 2011. Identity of *Pelodiscus sinensis* revealed by DNA sequences of an approximately 180-year-old type specimen and a taxonomic reappraisal of *Pelodiscus* species (Testudines: Trionychidae). Journal of Zoological Systematics and Evolutionary Research 49:335–339.

Stuckas, H., Gemel, R., and Fritz, U. 2013. One extinct turtle species less: *Pelusios seychellensis* is not extinct, it never existed. PLoS ONE 8(4):e57116.

Stuckas, H., Velo-Antón, G., Fahd, S., Kalboussi, M., Rouag, R., Arculeo, M., Marrone, F., Sacco, F., Vamberger, M., and Fritz, U. 2014. Where are you from, stranger? The enigmatic biogeography of North African pond turtles (*Emys orbicularis*). Organisms, Diversity and Evolution 14:295–306.

Sturm, J. 1828. Deutschlands Fauna in Abbildungen nach der Natur mit Beschreibungen. III. Abtheilung. Die Amphibien. Nürnberg: Privately printed.

Suckow, G.A. 1798. Anfangsgründe der theoretischen und angewandten Naturgeschichte der Thiere. Dritter Theil. Von den Amphibien. Leipzig: Weidmannischen Buchhandlung, 298 pp.

Sumontha, M., Brophy, T.R., Kunya, K., Wiboonatthapol, S., and Pauwels, O.S.G. 2016. A new snail-eating turtle of the genus *Malayemys* Lindholm, 1931 (Geoemydidae) from Thailand and Laos. Taprobanica 8(1):1–9.

Sung, Y.-H., Hau, B.C.H., and Karraker, N.E. 2014. Reproduction of endangered Big-headed Turtle, *Platysternon megacephalum* (Reptilia: Testudines: Platysternidae). Acta Herpetologica 9(2):243–247.

Sutton, K.K. and Christiansen, J.L. 1999. The habitat and distribution of the stinkpot, *Sternotherus odoratus*, in Iowa. Journal of the Iowa Academy of Science 106(3):63-65.

Suzuki, D. and Hikida, T. 2011. Mitochondrial phylogeography of the Japanese pond turtle, *Mauremys japonica* (Testudines, Geoemydidae). Journal of Zoological Systematics and Evolutionary Research 49:141–147.

Suzuki, D. and Hikida, T. 2014. Taxonomic status of the soft-shell turtles populations in Japan: a molecular approach. Current Herpetology 33:171–179.

Suzuki, D., Ota, H., Oh, H.S., and Hikida, T. 2011. Origin of Japanese populations of Reeves' Pond Turtle, *Mauremys reevesii* (Reptilia: Geoemydidae), as inferred by a molecular approach. Chelonian Conservation and Biology 10(2):237–249.

Suzuki, D., Yabe, T., and Hikida, T. 2014. Hybridization between *Mauremys japonica* and *Mauremys reevesii* inferred by nuclear and mitochondrial DNA analyses. Journal of Herpetology 48:445–454.

Swainson, W. 1839. On the natural history and classification of fishes, amphibians, and reptiles. Vol. II. In: Lardner, D. (Ed.). The Cabinet Cyclopaedia. Natural History. London: Longman, 452 pp.

Swinhoe, R. 1863. A list of the Formosan reptiles; with notes on a few of the species, and some remarks on a fish (*Orthagoriscus*, sp.). Annals and Magazine of Natural History 3(12):219–226.

Taffortet, J. 1726. Relation de l'Isle Rodrigue. Paris: Manuscript, Archives Nationales, Fonds des Colonies, Correspondence de l'Île de France, C4, 12(48), Ministère de la Marine 29(1).

Takahashi, A. 2012. On *Manouria oyamai*. In: Yamasaki, S. and Fujita, M. (Eds.). [Pictorial Publication for the 100th Anniversary Special Exhibition of Seiho Oyama's Birth: Passion for Discoveries]. Naha: Okinawa Prefectural Museum & Art Museum, pp. 64–68. [in Japanese]

Takahashi, A., Otsuka, H., and Hirayama, R. 2003. A new species of the genus *Manouria* (Testudines: Testudinidae) from the Upper Pleistocene of the Ryukyu Islands, Japan. Paleontological Research 7(3):195–217.

Takahashi, A., Otsuka, H., and Ota, H. 2008. Systematic review of Late Pleistocene turtles (Reptilia: Chelonii) from the Ryukyu Archipelago, Japan, with special reference to paleogeographical implications. Pacific Science 62(3):395–402.

Takahashi, A., Hirayama, R., Alcala, A.C., Carretero, M.A., Danilov, I.G., Ernst, C.H., Honegger, R., Lovich, J.E., Malonza, P.K., Moravec, J., Penny,

M., Praschag, P., Široký, P., Spinks, P.Q., Griffiths, C., Hansen, D., Bauerfeld, K., Glaw, F., Fong, J.F., Fritz, U., Kraus, O., Hailey, A., Vences, M., Wanless, R.M., Williams, E.H., Jr., Nieves-Rivera, A.M., Bunkley-Williams, L., Grimm, U., Iverson, J.B., Mortimer, J.A., Buskirk, J., Morgan, J., Schmidt, F., and Miller, J. 2009. Comments on the proposed conservation of usage of *Testudo gigantea* Schweigger, 1812 (currently *Geochelone (Aldabrachelys) gigantea*; Reptilia, Testudines) (Case 3463; see BZN 66: 34–50, 80–87, 169–186). Bulletin of Zoological Nomenclature 66(3):274–290.

Takahashi, A., Hirayama, R., and Otsuka, H. 2018. Systematic revision of *Manouria oyamai* (Testudines, Testudinidae), based on new material from the Upper Pleistocene of Okinawajima Island, the Ryukyu Archipelago, Japan, and its paleogeographic implications. Journal of Vertebrate Paleontology 38(2):e1427594.

Tamayo, J.L. 1962. Geografia General de Mexico. Tomo III. Geografia Biologica y Humana. Mexico: Instituto Mexicano de Investigaciones Económicas, 633 pp.

Tang, Y. 1997. [Research on a new species of *Pelodiscus*, Trionychidae in China]. Zoological Research, Kunming 18(1):13–17. [in Chinese].

Tao, H.J. 1985. New fossil turtles, *Chinemys pani* n. sp. (Testudinidae) from the Chi-Ting Formation (Pleistocene), Tainan District, Taiwan Island. Journal of the Taiwan Museum 38(1):43–52.

Tao, H.J. 1986. Report of a new fossil soft-shelled turtle, *Trionyx liupani* from Taiwan, with comparative study to the living species, *Trionyx sinensis* (Wiegmann). Journal of the Taiwan Museum 39(2):21–41.

Tao, H.J. 1988. New fossil turtle, *Ocadia sinensis changwui* n. subsp., from Late Pleistocene, Taiwan Strait. Acta Zoologica Taiwanica 2:229–240.

Tapia, W. 2019. The most unexpected discovery of my life: a live tortoise on Fernandina Island. Galapagos Conservancy: https://www.galapagos.org/blog/2019-fernandina-tortoise-discovery/.

Tapia, W., Sevilla, A., Málaga, J., and Gibbs, J.P. 2020. Tortoise populations after 60 years of conservation. In: Gibbs, J.P., Cayot, L.J., and Tapia Aguilera, W. (Eds.). Galapagos Giant Tortoises. London: Academic Press, pp. 401–432.

Taskavak, E. 1998. Comparative morphology of the Euphrates soft-shelled turtle, *Rafetus euphraticus* (Daudin, 1802) (Reptilia, Testudines) in southeastern Anatolia. Amphibia-Reptilia 19(3):281–291.

Taskavak, E. and Akcinar, S.C. 2009. Marine records of the Nile soft-shelled turtle, *Trionyx triunguis* from Turkey. Marine Biodiversity Records 2:1–4.

Taylor, D.B. 1991. *Clemmys guttata* (spotted turtle). Size maxima. Herpetological Review 22(2):55.

Taylor, E.H. 1920. Philippine turtles. Philippine Journal of Science, Manila 16(2):111–144.

Taylor, E.H. 1943. An extinct turtle of the genus *Emys* from the Pleistocene of Kansas. University of Kansas Science Bulletin 29(II)(3):249–254.

Taylor, W.E. 1895. The box turtles of North America. Proceedings of the United States National Museum 17:573–588.

TCC [Turtle Conservation Coalition: Rhodin, A.G.J., Walde, A.D., Horne, B.D., van Dijk, P.P., Blanck, T., and Hudson, R. (Eds.)]. 2011. Turtles in Trouble: The World's 25+ Most Endangered Tortoises and Freshwater Turtles–2011. Lunenburg, MA: IUCN/SSC Tortoise and Freshwater Turtle Specialist Group, Turtle Conservation Fund, Turtle Survival Alliance, Turtle Conservancy, Chelonian Research Foundation, Conservation International, Wildlife Conservation Society, and San Diego Zoo Global, 54 pp.

TCC [Turtle Conservation Coalition: Stanford, C.B., Rhodin, A.G.J., van Dijk, P.P., Horne, B.D., Blanck, T., Goode, E.V., Hudson, R., Mittermeier, R.A., Currylow, A., Eisemberg, C., Frankel, M., Georges, A., Gibbons, P.M., Juvik, J.O., Kuchling, G., Luiselli, L., Shi, H., Singh, S., and Walde, A. (Eds.)]. 2018. Turtles in Trouble: The World's 25+ Most Endangered Tortoises and Freshwater Turtles–2018. Ojai, CA: IUCN SSC Tortoise and Freshwater Turtle Specialist Group, Turtle Conservancy, Turtle Survival Alliance, Turtle Conservation Fund, Chelonian Research Foundation, Conservation International, Wildlife Conservation Society, and Global Wildlife Conservation, 80 pp.

TCC [Turtle Conservation Coalition: Stanford, C.B., Rhodin, A.G.J., van Dijk, P.P., Blanck, T., Goode, E.V., Hudson, R., Mittermeier, R.A., and Páez, V.P. (Eds.)]. 2025. Turtles in Trouble: The World's Most Endangered Tortoises and Freshwater Turtles–2025. Ojai, CA: IUCN SSC Tortoise and Freshwater Turtle Specialist Group, Turtle Conservancy, Turtle Survival Alliance, Turtle Conservation Fund, Re:wild, and Chelonian Research Foundation.

TCF [Turtle Conservation Fund: Buhlmann, K.A., Hudson, R., and Rhodin, A.G.J.]. 2002. A Global Action Plan for Conservation of Tortoises and Freshwater Turtles. Strategy and Funding Prospectus 2002–2007. Washington, DC: Conservation International and Chelonian Research Foundation, 30 pp.

TCF [Turtle Conservation Fund]. 2003. Top 25 Turtles on Death Row: New List Spotlights Most Endangered Turtles and Action Plan to Save Them. Washington, DC: Conservation International and Turtle Conservation Fund. https://turtleconservationfund.org/wp-content/uploads/2008/02/top25.pdf.

TCF [Turtle Conservation Fund: Rhodin, A.G.J., Quinn, H.R., Goode, E.V., Hudson, R., Mittermeier, R.A., and van Dijk, P.P.]. 2019. Turtle Conservation Fund: A Partnership Coalition of Leading Turtle Conservation Organizations and Individuals—Summary Activity Report 2002–2018. Lunenburg, MA and Ojai, CA: Chelonian Research Foundation and Turtle Conservancy, 54 pp.

Temminck, C.J. and Schlegel, H. 1834. Reptilia. I. Les Chéloniens. In: Siebold, P.F. de. Fauna Japonica, sive Descriptio animalium, quae in itinere per Japoniam, jussu et auspiciis superiorum, qui summum in India Batava Imperium tenent, suscepto, annis 1823-1830 colleget, notis observationibus et adumbrationibus illustravit. Vol. III. Lugduni Batavorum [Leiden]: J.G. La Lau, pp. 1–80, pls.I–IX.

Temminck, C.J. and Schlegel, H. 1838. Reptilia. III. Explication des planches de sauriens et de batraciens. In: Siebold, P.F. de. Fauna Japonica, sive Descriptio animalium, quae in itinere per Japoniam, jussu et auspiciis superiorum, qui summum in India Batava Imperium tenent, suscepto, annis 1823-1830 colleget, notis observationibus et adumbrationibus illustravit. Vol. III. Lugduni Batavorum [Leiden]: J.G. La Lau, pp. 136–140.

Tesche, M.R. and Hodges, K.E. 2015. Rethinking biogeographic patterns: high local variation in relation to latitudinal clines for a widely distributed species. Oecologia 179:139–149.

Tewari, B.S. and Badam, G.L. 1969. A new species of fossil turtle from the Upper Siwaliks of Pinjore, India. Palaeontologica 12(4):555–558.

TEWG [Turtle Extinctions Working Group: Rhodin, A.G.J., Thomson, S., Georgalis, G.L., Karl, H.-V., Danilov, I.G., Takahashi, A., de la Fuente, M.S., Bourque, J.R., Delfino, M., Bour, R., Iverson, J.B., Shaffer, H.B., and van Dijk, P.P.]. 2015. Turtles and tortoises of the world during the rise and global spread of humanity: first checklist and review of extinct Pleistocene and Holocene chelonians. Chelonian Research Monographs 5(8):000e.1–66.

TFTSG [IUCN/SSC Tortoise and Freshwater Turtle Specialist Group]. 2007. Turtles in Trouble: the World's Top 25 Most Endangered Tortoises and Freshwater Turtles – 2007. https://iucn-tftsg.org/wp-content/uploads/file/Top%2025/Turtles-in-Trouble-Top_25_2007.pdf.

Thapa, A., Pradhan, P., Thakur, M., Joshi, B.D., Sharma, L.K., and Chandra, K. 2021. A new elevational and locality record of the Assam Leaf Turtle, *Cyclemys gemeli* (Geoemydidae), from the Darjeeling Hills in the Central Himalayan Biogeographic Province. Reptiles & Amphibians 28(2):326–328.

Theobald, W., Jr. 1856. Another rich collection of sundries, from Mergui and the valley of the Tenasserim river. Journal of the Asiatic Society of Bengal 24(1855):711–721.

Theobald, W., Jr. 1860. On the Tertiary and alluvial deposits of the central portion of the Nerbudda Valley. Memoirs of the Geological Survey of India 2:279–298.

Theobald, W., Jr. 1868a. Catalogue of Reptiles in the Museum of the Asiatic Society of Bengal. Journal of the Asiatic Society, Extra Number, 88 pp.

Theobald, W., Jr. 1868b. Catalogue of the reptiles of British Birma, embracing the provinces of Pegu, Martaban, and Tenasserim; with descriptions of new or little-known species. Journal of the Linnean Society of Zoology 10:4–67.

Theobald, W., Jr. 1874. Observations on some Indian and Burmese species of *Trionyx*. Proceedings of the Asiatic Society of Bengal 1874:75–86.

Theobald, W., Jr. 1875. Observations on some Indian and Burmese species of *Trionyx*, with a rectification of their synonymy and a description of two new species. Proceedings of the Asiatic Society of Bengal 1875:170–180.

Theobald, W., Jr. 1876. Descriptive Catalogue of the Reptiles of British India. Calcutta: Thacher, Spink and Co., 238 pp.

Theobald, W., Jr. 1877. Description of a new Emydine from the upper Tertiaries of the northern Punjab. Records of the Geological Survey of India 10:43–45.

Thiele, K.R., Conix, S., Pyle, R.L., Barik, S.K., Christidis, L., Costello, M.J., van Dijk, P.P., Kirk, P., Lien, A., Thomson, S.A., Zachos, F.E., Zhang, Z., and Garnett, S.T. 2021. Towards a global list of accepted species I. Why taxonomists sometimes disagree, and why this matters. Organisms Diversity & Evolution: doi.org/10.1007/s13127-021-00495-y, 8 pp.

Thieme, U. 2005. Notes on the East Cuban Slider, *Trachemys decussata decussata* (Gray, 1831). Radiata 14(4):29–32.

Thirakhupt, K. and van Dijk, P.P. 1995. Species diversity and conservation of

turtles of western Thailand. Natural History Bulletin of the Siam Society 42(2)(1994)[1995]:207–259.

Thomas, R.B. and Jansen, K.P. 2006. *Pseudemys floridana* – Florida cooter. In: Meylan, P.A. (Ed.). Biology and Conservation of Florida Turtles. Chelonian Research Monographs 3:338–347.

Thomas, T.M., Granatosky, M.C., Bourque, J.R., Krysko, K.L., Moler, P.E., Gamble, T., Suarez, E., Leone, E., Enge, K.M., and Roman, J. 2014. Taxonomic assessment of Alligator Snapping Turtles (Chelydridae: *Macrochelys*), with the description of two new species from the southeastern United States. Zootaxa 3786(2):141–165.

Thomson, R.C., Spinks, P.Q., and Shaffer, H.B. 2018. Molecular phylogeny and divergence of the map turtles (Emydidae: *Graptemys*). Molecular Phylogenetics and Evolution 121:61–70.

Thomson, R.C., Spinks, P.Q., and Shaffer, H.B. 2021. A global phylogeny of turtles reveals a burst of climate-associated diversification on continental margins. Proceedings of the National Academy of Sciences (PNAS) 118:7, e2012215118, 10 pp.

Thomson, S. 2000a. On the identification of the holotype of *Chelodina oblonga* (Testudinata: Chelidae) with a discussion of the taxonomic implications. Chelonian Conservation and Biology 3:745–749.

Thomson, S.A. 2000b. A revision of the fossil chelid turtles (Pleurodira) described by C.W. DeVis (1897). Memoirs of the Queensland Museum 43:593–598.

Thomson, S. 2006. ICZN Case 3351. *Chelodina rugosa* Ogilby, 1890 (currently *Macrochelodina rugosa*; Reptilia, Testudines): proposed precedence over *Chelodina oblonga* Gray, 1841. Bulletin of Zoological Nomenclature 63(3):187–193.

Thomson, S. 2007. Comment on the proposed precedence of *Chelodina rugosa* Ogilby, 1890 (currently *Macrochelodina rugosa*; Reptilia, Testudines) over *Chelodina oblonga* Gray, 1841. Bulletin of Zoological Nomenclature 64:127–128.

Thomson, S. and Georges, A. 2009. *Myuchelys* gen. nov.—a new genus for *Elseya latisternum* and related forms of Australian freshwater turtle (Testudines: Pleurodira: Chelidae). Zootaxa 2053:32–42.

Thomson, S. and Georges, A. 2016. A new species of freshwater turtle of the genus *Elseya* (Testudinata: Pleurodira: Chelidae) from the Northern Territory of Australia. Zootaxa 4061(1):18–28.

Thomson, S.A. and Lambertz, M. 2017. On the nomenclatural status of the recently described Snail-Eating Turtle from Southeast Asia (Testudines, Geoemydidae): *Malayemys khoratensis* Ihlow et al., 2016 vs. *Malayemys isan* Sumontha et al., 2016. Chelonian Conservation and Biology 16(2):239–245.

Thomson, S., White, A., and Georges, A. 1997. Re-evaluation of *Emydura lavarackorum*: identification of a living fossil. Memoirs of the Queensland Museum 42(1):327–336.

Thomson, S., Kennett, R., and Georges, A. 2000. A new species of long-necked turtle (Testudines: Chelidae) from the Arnhem Land Plateau, Northern Territory, Australia. Chelonian Conservation and Biology 3(4):675–685.

Thomson, S., Georges, A., and Limpus, C.J. 2006. A new species of freshwater turtle in the genus *Elseya* (Testudines: Chelidae) from central coastal Queensland, Australia. Chelonian Conservation and Biology 5(1):74–86.

Thomson, S., Amepou, Y., Anamiato, J., and Georges, A. 2015. A new species and subgenus of *Elseya* (Testudines: Pleurodira: Chelidae) from New Guinea. Zootaxa 4006(1):59–82.

Thomson, S.A., Pyle, R.L., Ahyong, S.T., Alonso-Zarazaga, M., Ammirati, J., Araya, J.F., Ascher, J.S., Audisio, T.L., Azevedo-Santos, V.M., Bailly, N., Baker, W.J., Balke, M., Barclay, M.V.L., Barrett, R.L., Benine, R.C., Bickerstaff, J.R.M., Bouchard, P., Bour, R., Bourgoin, T., Boyko, C.B., Breure, A.S.H., Brothers, D.J., Byng, J.W., Campbell, D., Ceríaco, L.M.P., Cernák, I., Cerretti, P., Chang, C., Cho, S., Copus, J.M., Costello, M.J., Cseh, A., Csuzdi, C., Culham, A., D'elía, G., D'udekem D'acoz, C., Daneliya, M.E., Dekker, R., Dickinson, E.C., Dickinson, T.A., van Dijk, P.P., Dijkstra, K.B., Dima, B., Dmitriev, D.A., Duistermaat, L., Dumbacher, J.P., Eiserhardt, W.L., Ekrem, T., Evenhuis, N.L., Faille, A., Fernández-Triana, J.L., Fiesler, E., Fishbein, M., Fordham, B.G., Freitas, A.V.L., Friol, N.R., Fritz, U., Frøslev, T., Funk, V.A., Gaimari, S.D., Garbino, G.S.T., Garraffoni, A.R.S., Geml, J., Gill, A.C., Gray, A., Grazziotin, F.G., Greenslade, P., Gutiérrez, E.E., Harvey, M.S., Hazevoet, C.J., He, K., He, X., Helfer, S., Helgen, K.M., Van Heteren, A.H., Hita Garcia, F., Holstein, N., Horváth, M.K., Hovenkamp, P.H., Hwang, W.S., Hyvönen, J., Islam, M.B., Iverson, J.B., Ivie, M.A., Jaafar, Z., Jackson, M.D., Jayat, J.P., Johnson, N.F., Kaiser, H., Klitgård, B.B., Knapp, D.G., Kojima, J., Kõljalg, U., Kontschán, J., Krell, F.-T., Krisai-Greilhuber, I., Kullander, S., Latella, L., Lattke, J.E., Lencioni, V., Lewis, G.P., Lhano, M.G., Lujan, Nathan K., Luksenburg, J.A., Mariaux, J., Marinho-Filho, J., Marshall, C.J., Mate, J.F., Mcdonough, M.M., Michel, E., Miranda, V.F.O., Mitroiu, M.D., Molinari, J., Monks, S., Moore, A.J., Moratelli, R., Murányi, D., Nakano, T., Nikolaeva, S., Noyes, J., Ohl, M., Oleas, N.H., Orrell, T., Páll-Gergely, B., Pape, T., Papp, V., Parenti, L.R., Patterson, D., Pavlinov, I.Y., Pine, R.H., Poczai, P., Prado, J., Prathapan, D., Rabeler, R.K., Randall, J.E., Rheindt, F.E., Rhodin, A.G.J., Rodríguez, S.M., Rogers, D.C., Roque, F. De O., Rowe, K.C., Ruedas, L.A., Salazar-Bravo, J., Salvador, R.B., Sangster, G., Sarmiento, C.E., Schigel, D.S., Schmidt, S., Schueler, F.W., Segers, H., Snow, N., Souza-Dias, P.G.B., Stals, R., Stenroos, S., Stone, R.D., Sturm, C.F., Stys, P., Teta, P., Thomas, D.C., Timm, R.M., Tindall, B.J., Todd, J.A., Triebel, D., Valdecasas, A.G., Vizzini, A., Vorontsova, M.S., De Vos, J.M., Wagner, P., Watling, L., Weakley, A., Welter-Schultes, F., Whitmore, D., Wilding, N., Will, K., Williams, J., Wilson, K., Winston, J.E., Wüster, W., Yanega, D., Yeates, D.K., Zaher, H., Zhang, G., Zhang, Z.Q., Zhou, H.Z. 2018. Taxonomy based on science is necessary for global conservation. PLoS Biology 16(3):e2005075.

Thomson, S.A., Thiele, K., Conix, S., Christidis, L., Costello, M.J., Hobern, D., Nikolaeva, S., Pyle, R.L., van Dijk, P.P., Weaver, H., Zachos, F.E., Zhang, Z.Q., and Garnett, S.T. 2021. Towards a global list of accepted species II. Consequences of inadequate taxonomic list governance. Organisms Diversity & Evolution 21:623–630.

Thomson, S.A., Friol, N.R., White, A., Wedd, D., and Georges, A. 2023. The Australian gulf snapping turtle *Elseya lavarackorum* (Testudines: Chelidae) revisited—is the late Pleistocene fossil species extant? Vertebrate Zoology 73:237–256.

Thorbjarnarson, J.B., Pérez, N., and Escalona, T. 1997. Biology and conservation of aquatic turtles in the Cinaruco-Capanaparo National Park, Venezuela. In: Van Abbema, J. (Ed.). Proceedings: Conservation, Restoration, and Management of Tortoises and Turtles – An International Conference. N.Y. Turtle and Tortoise Society, pp. 109–112.

Thunberg, C.P. 1787. Beskrifning på trenne sköld-paddor. [Description of three turtles]. Kongliga Vetenskaps Academiens Nya Handlingar, Stockholm (2)8:178–180. [in Swedish]

Thunberg, C.P. 1788. Resa uti Europa, Africa, Asia, förrättad Åren 1770-1779. Första Delen, innehållande Resan til Södra Europa och Goda Hoppets Udde i Africa, Åren 1770, 1771, 1772, 1773. Upsala: Joh. Edman, 389 pp. [in Swedish]

Thunberg, C.P. 1792. [*Testudo scripta*]. In: Schoepff, J.D. Historia Testudinum Iconibus Illustrata. Erlangae: Ioannis Iacobi Palm, 136 pp. [pp. 16–17].

Thunberg, C.P. 1808. Museum Naturalium Academiae Upsaliensis. Appendic. XV, publicae censurae committit Jon Jacobi. Donation Thunbergian. Append. XV. Upsaliae: Edmannianis, 7 pp.

Thunberg, C.P. 1810. Museum Naturalium Academiae Upsaliensis, Partem XXVII, publicae censurae committit Ericus Hasselhun. Donationis Reg. Gustavi IV:ti Adolphi. Continuatio IV:ta. Upsaliae: Edmannianis, 8 pp.

Thunberg, C.P. 1812. [*Testudo discolor*]. In: Schweigger, A.F. Prodromus monographiae Cheloniorum. Königsberger Archiv für Naturwissenschaftliche und Mathematik 1:271–368, 406–458. [p. 302].

Thunberg, C.P. 1828. General-Catalogue på Kongl. Akademiens i Upsala, Natural-Samlingar. Unpublished manuscript, Uppsala University Zoological Museum, 311 pp. [in Swedish]

Tiedemann, F. and Häupl, M. 1980. Typenkatalog der Herpetologischen Sammlung. Teil II: Reptilia. Kataloge der wissenschaftlichen Sammlungen des Naturhistorischen Museums in Wien, Band 4, Vertebrata, Heft 2:1–79.

Tiedemann, F., Häupl, M., and Grillitsch, H. 1994. Katalog der Typen der herpetologischen Sammlung nach dem stand vom 1. Jänner 1994. Teil II. Reptilia. Kataloge der wissenschaftlichen Sammlungen des Naturhistorischen Museums in Wien 10 (Vertebrata 4):1–102.

Tietz, B., Penner, J., and Vamberger, M. 2023. Chelonian challenge: three alien species from North America are moving their reproductive boundaries in Central Europe. NeoBiota 82:1–21.

Tinkle, D.W. 1958. The systematics and ecology of the *Sternothaerus carinatus* complex (Testudinata, Chelydridae). Tulane Studies in Zoology 6:1–56.

Tinkle, D.W. and Webb, R.G. 1955. A new species of *Sternotherus* with a discussion of the *Sternotherus carinatus* complex. Tulane Studies in Zoology 3(3):53–67.

Tirant, G. 1884. Notes sur les reptiles de la Cochinchine et du Cambodge.

Excursions et Reconnaissances 8(19):147–168.

Todd, E.V., Blair, D., Farley, S., Farrington, L., FitzSimmons, N.N., Georges, A., Limpus, C.J., and Jerry, D.R. 2013. Contemporary genetic structure reflects historical drainage isolation in an Australian snapping turtle, *Elseya albagula*. Zoological Journal of the Linnean Society 169:200–214.

Todd, E.V., Blair, D., and Jerry, D.R. 2014a. Influence of drainage divides versus arid corridors on genetic structure and demography of a widespread freshwater turtle, *Emydura macquarii krefftii*, from Australia. Ecology and Evolution 4:606–622.

Todd, E.V., Blair, D., Georges, A., Lukoschek, V., and Jerry, D.R. 2014b. A biogeographical history and timeline for the evolution of Australian snapping turtles (*Elseya*: Chelidae) in Australia and New Guinea. Journal of Biogeography 41(5):905–918.

Toledo, M.A., de la Concha, A., and de Lucas, J.L.R. 2021. Space use, distribution and conservation of the Spanish pond turtle (*Mauremys leprosa* Schweigger, 1812). A review (Chordata, Geoemydidae). Boletín de la Real Sociedad Española de Historia Natural 115:95–110.

Tomović, L., Ajtić, R., and Golubović, A. 2019. New records of *Testudo graeca* (Pallas, 1814) in Republic of Serbia. Bulletin of the Natural History Museum 12:209–215.

Townsend, C.H. 1925. The Galapagos tortoises in their relation to the whaling industry: a study of old logbooks. Zoologica 4(3):55–135.

Trape, J.-F., Trape, S., and Chirio, L. 2012. Lézards, Crocodiles et Tortues d'Afrique Occidentale et du Sahara. Marseille: IRD éditions, 503 pp.

Trebbau, P. and Pritchard, P.C.H. 2016. Venezuela y sus Tortugas. Caracas: Oscar Todtmann Editores, 184 pp.

Troost, G. 1835. [*Chelonura temminckii*]. In: Harlan, R. Genera of North American Reptilia, and a synopsis of the species. In: Harlan, R. Medical and Physical Researches; or Original Memoirs in Medicine, Surgery, Physiology, Geology, Zoology, and Comparative Anatomy. Philadelphia: Bailey, 653 pp. [pp. 157–158].

Troschel, F.H. 1848. Amphibien. In: Schomburgk, R. Reisen in Britisch-Guiana in den Jahren 1840–1844. Dritter Theil. Versuch einer Fauna und Flora von Britisch-Guiana. Leipzig: Verlagsbuchhandlung J.J. Weber, pp. 645–661.

True, F.W. 1882. On the North American land tortoises of the genus *Xerobates*. Proceedings of the United States National Museum 1881(1882):434–449.

Trutnau, L. 1994. Terraristik. Stuttgart: Eugen Ulmer Verlag, 248 pp.

Tschudi, J.J. von. 1846. Untersuchungen über die Fauna Peruana. Herpetologie. St. Gallen: Scheitlin and Zollikofer, 80 pp.

TTWG [Turtle Taxonomy Working Group: Bickham, J.W., Parham, J.F., Philippen, H.D., Rhodin, A.G.J., Shaffer, H.B., Spinks, P.Q., and van Dijk, P.P.]. 2007a. Turtle taxonomy: methodology, recommendations, and guidelines. In: Shaffer, H.B., FitzSimmons, N.N., Georges, A., and Rhodin, A.G.J. (Eds.). Defining Turtle Diversity: Proceedings of a Workshop on Genetics, Ethics, and Taxonomy of Freshwater Turtles and Tortoises. Chelonian Research Monographs 4:73–84.

TTWG [Turtle Taxonomy Working Group: Bickham, J.W., Iverson, J.B., Parham, J.F., Philippen, H.D., Rhodin, A.G.J., Shaffer, H.B., Spinks, P.Q., and van Dijk, P.P.]. 2007b. An annotated list of modern turtle terminal taxa with comments on areas of taxonomic instability and recent change. In: Shaffer, H.B., FitzSimmons, N.N., Georges, A., and Rhodin, A.G.J. (Eds.). Defining Turtle Diversity: Proceedings of a Workshop on Genetics, Ethics, and Taxonomy of Freshwater Turtles and Tortoises. Chelonian Research Monographs 4:173–199.

TTWG [Turtle Taxonomy Working Group: Rhodin, A.G.J., Parham, J.F., van Dijk, P.P., and Iverson, J.B.]. 2009. Turtles of the world: annotated checklist of taxonomy and synonymy, 2009 update, with conservation status summary. Chelonian Research Monographs 5(2):000.39–84.

TTWG [Turtle Taxonomy Working Group: Rhodin, A.G.J., van Dijk, P.P., Iverson, J.B., and Shaffer, H.B.]. 2010. Turtles of the world, 2010 update: annotated checklist of taxonomy, synonymy, distribution, and conservation status. Chelonian Research Monographs 5(3):000.85–164.

TTWG [Turtle Taxonomy Working Group: van Dijk, P.P., Iverson, J.B., Shaffer, H.B., Bour, R., and Rhodin, A.G.J.]. 2011. Turtles of the world, 2011 update: annotated checklist of taxonomy, synonymy, distribution, and conservation status. Chelonian Research Monographs 5(4):000.165–242.

TTWG [Turtle Taxonomy Working Group: van Dijk, P.P., Iverson, J.B., Shaffer, H.B., Bour, R., and Rhodin, A.G.J.]. 2012. Turtles of the world, 2012 update: annotated checklist of taxonomy, synonymy, distribution, and conservation status. Chelonian Research Monographs 5(5):000.243–328.

TTWG [Turtle Taxonomy Working Group: van Dijk, P.P., Iverson, J.B., Rhodin, A.G.J., Shaffer, H.B., and Bour, R.]. 2014. Turtles of the world, 7th edition: annotated checklist of taxonomy, synonymy, distribution with maps, and conservation status. Chelonian Research Monographs 5(7):000.329–479.

TTWG [Turtle Taxonomy Working Group: Rhodin, A.G.J., Iverson, J.B., Bour, R., Fritz, U., Georges, A., Shaffer, H.B., and van Dijk, P.P.]. 2017. Turtles of the World: Annotated Checklist and Atlas of Taxonomy, Synonymy, Distribution, and Conservation Status (8th Ed.). Chelonian Research Monographs 7:1–292.

TTWG [Turtle Taxonomy Working Group: Rhodin, A.G.J., Iverson, J.B., Bour, R., Fritz, U., Georges, A., Shaffer, H.B., and van Dijk, P.P.]. 2021. Turtles of the World: Annotated Checklist and Atlas of Taxonomy, Synonymy, Distribution, and Conservation Status (9th Ed.). Chelonian Research Monographs 8:1–472.

Tuberville, T.D., Buhlmann, K.A., Bjorkland, R.K., and Booher, D. 2005. Ecology of the Jamaican Slider Turtle (*Trachemys terrapen*), with implications for conservation and management. Chelonian Conservation and Biology 4(4):908–915.

Tucker, A.D. and Frazer, N.B. 1991. Reproductive variation in leatherback turtles, *Dermochelys coriacea*, at Culebra National Wildlife Refuge, Puerto Rico. Herpetologica 47(1):115–124.

Tucker, J.K., Lamer, J.T., Dolan, C.R., and Dustman, E.A. 2006. Chelonian species: record carapace lengths for Illinois. Herpetological Review 37: 453–455.

Türkozan, O., Kiremit, F., Parham, J.F., Olgun, K., and Taskavak, E. 2010. A quantitative reassessment of morphology-based taxonomic schemes for Turkish tortoises (*Testudo graeca*). Amphibia-Reptilia 31:69–83.

Türkozan, O., Kiremit, F., Lavin, B.R., Bardakci, F., and Parham, J.F. 2018. Morphological and mitochondrial variation of spur-thighed tortoises, *Testudo graeca*, in Turkey. Herpetological Journal 28:1–9.

Turnbull, L.A., Ozgul, A., Accouche, W., Baxter, R., ChongSeng, L., Currie, J.C., Doak, N., Hansen, D.M., Pistorius, P., Richards, H., van de Crommenacker, J., von Brandis, R., Fleischer-Dogley, F., and Bunbury, N. 2015. Persistence of distinctive morphotypes in the native range of the CITES-listed Aldabra giant tortoise. Ecology and Evolution 5:5499–5508.

Turner, D.S., Van Devender, T.R., Hale, S.F., Zach, R., Martínez, R., Van Devender, R.W., Salazar-Martínez, J.A., Molina-Padilla, G., Sherbrooke, W.C., Aguilar-Morales, C., Valdez-Villavicencio, J.H., Peralta-García, A., Minter, S.L., Silva-Kurumiya, H., Villa, R.A., Yanes-Arvayo, G., Montañez-Armenta, M.D.L.P., and Paholski, C. 2022. Amphibians and reptiles of Rancho Las Playitas area, Sonora, Mexico. Sonoran Herpetologist 35(2):62–71.

Turvey, S.T., Almonte, J., Hansford, J., Scofield, R.P., Brocca, J.L., and Chapman, S.D. 2017. A new species of extinct Late Quaternary giant tortoise from Hispaniola. Zootaxa 4277(1):1–16.

Uetz, P., Cherikh, S., Shea, G., Ineich, I., Campbell, P.D., Doronin, I.V., Rosado, J., Wynn, A., Tighe, K.A., McDiarmid, R., Lee, J.L., Köhler, G., Ellis, R., Doughty, P., Raxworthy, C.J., Scheinberg, L., Resetar, A., Sabaj, M., Schneider, G., Franzen, M., Glaw, F., Böhme, W., Schweiger, S., Gemel, R., Couper, P., Amey, A., Dondorp, E., Ofer, G., Meiri, S., and Wallach, V. 2019. A global catalog of primary reptile type specimens. Zootaxa 4695(5):438–450.

Uhrin, M., Havaš, P., Minařik, M., Kodejš, K., Bugoš, I., Danko, S., Husák, T., Koleska, D., and Jablonski, D. 2016. Distribution updates to amphibian and reptile fauna for the Republic of Macedonia. Herpetology Notes 9:201–220.

Ultsch, G.R., LeBerte, C.M., and McAlpine, D.F. 2000. *Chrysemys picta picta* (eastern painted turtle). Maximum size. Herpetological Review 31:103–104.

Ultsch, G.R., Ward, G.M., LeBerte, C.M., Kuhajda, B.R., and Stewart, E.R. 2001. Intergradation and origins of subspecies of the turtle *Chrysemys picta*: morphological comparisons. Canadian Journal of Zoology 79:485–498.

Umulisa, C., Djurakic, M., Bantlin, D., Müller, A., and Vamberger, M. 2025. An integrative approach of taxonomic, morphometric and ecological research to solve the status of cryptic chelonian taxa—a case study with *Pelomedusa kobe* in Rwanda. African Journal of Ecology 63:e70006, 13 pp.

UNEP-WCMC. 2017. Report on species/country combinations selected for review by the Animals Committee following CoP16. Annex 1 to CITES document AC29 Doc.13.2. https://cites.org/sites/default/files/eng/com/ac/29/E-AC29-13-02-A1.pdf.

Ureña-Aranda, C.A. and Espinosa de los Monteros, A. 2012. The genetic crisis

of the Mexican Bolson Tortoise (*Gopherus flavomarginatus*: Testudinidae). Amphibia-Reptilia 33:45–53.

USFWS [U.S. Fish and Wildlife Service]. 1970. Conservation of endangered species and other fish or wildlife. Federal Register 35:8491–8498.

USFWS [U.S. Fish and Wildlife Service]. 1973. Conservation of endangered species and other fish or wildlife. Amendments to lists of endangered fish and wildlife. Federal Register 38:14678.

USFWS [U.S. Fish and Wildlife Service]. 1976. Endangered status for 159 taxa of animals. Federal Register 41:24062–24067.

USFWS [U.S. Fish and Wildlife Service]. 1978. Listing and protecting Loggerhead Sea Turtles as "Threatened Species" and populations of Green and Olive Ridley Sea Turtles as Threatened Species or "Endangered Species." Federal Register 43:32800–32811.

USFWS [U.S. Fish and Wildlife Service]. 1979. Listing of the Bolson Tortoise as an endangered species. Federal Register 44:23062–23064.

USFWS [U.S. Fish and Wildlife Service]. 1980. Endangered and threatened wildlife and plants; listing as endangered with critical habitat for the Plymouth Red-bellied Turtle in Massachusetts. Federal Register 45:21828–21833.

USFWS [U.S. Fish and Wildlife Service]. 1983. Endangered and threatened wildlife and plants; listing of 17 species of foreign reptiles as endangered or threatened species. Federal Register 48:28460–28464.

USFWS [U.S. Fish and Wildlife Service]. 1986. Endangered and threatened wildlife and plants; determination of threatened status for the Ringed Sawback Turtle (*Graptemys oculifera*). Federal Register 51:45907–45910.

USFWS [U.S. Fish and Wildlife Service]. 1987a. Endangered and threatened wildlife and plants; determination of threatened status for the Flattened Musk Turtle (*Sternotherus depressus*). Federal Register 52:22418–22430.

USFWS [U.S. Fish and Wildlife Service]. 1987b. Endangered and threatened wildlife and plants; determination of endangered status for the Alabama Red-bellied Turtle. Federal Register 52:22939–22943.

USFWS [U.S. Fish and Wildlife Service]. 1987c. Endangered and threatened wildlife and plants; determination of threatened status for the Gopher Tortoise (*Gopherus polyphemus*). Federal Register 52:25376–25380.

USFWS [U.S. Fish and Wildlife Service]. 1990. Endangered and threatened wildlife and plants; determination of threatened status for the Mojave population of the Desert Tortoise. Federal Register 55:12178–12191.

USFWS [U.S. Fish and Wildlife Service]. 1991a. Endangered and threatened wildlife and plants: threatened status for the Yellow-blotched Map Turtle, *Graptemys flavimaculata*. Federal Register 56:1459–1463.

USFWS [U.S. Fish and Wildlife Service]. 1991b. Endangered and threatened wildlife and plants; endangered status for six foreign reptiles. Federal Register 56:49649–49653.

USFWS [U.S. Fish and Wildlife Service]. 1997. Endangered and threatened wildlife and plants; final rule to list the northern population of the Bog Turtle as threatened and the southern population as threatened due to similarity of appearance. Federal Register 62:59605–59623.

USFWS [U.S. Fish and Wildlife Service]. 2017a. Species Status Assessment report for the Sonoyta Mud Turtle. Version 2.0. Albuquerque, NM: USFWS, 135 pp.

USFWS [U.S. Fish and Wildlife Service]. 2017b. Endangered and threatened wildlife and plants; endangered species status for Sonoyta Mud Turtle. Federal Register 82:43897–43907.

USFWS [U.S. Fish and Wildlife Service]. 2017c. Species Status Assessment report for the Barbour's Map Turtle (*Graptemys barbouri*). Atlanta, GA: USFWS, 83 pp.

USFWS [U.S. Fish and Wildlife Service]. 2021a. Species Status Assessment report for the Massachusetts population of the northern Red-bellied Cooter (*Pseudemys rubriventris*), Version 1.0. Hadley, MA: USFWS, 109 pp.

USFWS [U.S. Fish and Wildlife Service]. 2021b. Species Status Assessment (SSA) report for Rio Grande Cooter (*Pseudemys gorzugi*). Version 1.0. Austin, TX: USFWS, 166 pp.

USFWS [U.S. Fish and Wildlife Service]. 2021c. Species Status Assessment report for the Alligator Snapping Turtle (*Macrochelys temminckii*). Version 1.2. Atlanta, GA: USFWS, 218 pp.

USFWS [U.S. Fish and Wildlife Service]. 2021d. Species Status Assessment Report for the Pascagoula Map Turtle (*Graptemys gibbonsi*). Version 1.1. Atlanta, GA: USFWS, 143 pp.

USFWS [U.S. Fish and Wildlife Service]. 2021e. Species Status Assessment report for the Gopher Tortoise (*Gopherus polyphemus*). Version 0.4. Atlanta, GA: USFWS, 288 pp.

USFWS [U.S. Fish and Wildlife Service]. 2021f. Species Status Assessment for the Sonoran Desert Tortoise (*Gopherus morafkai*). Version 2.1. Albuquerque, NM: USFWS, 183 pp.

USFWS [U.S. Fish and Wildlife Service]. 2022a. Species Status Assessment for the Egyptian Tortoise (*Testudo kleinmanni*). Version 1.1. Falls Church, VA: USFWS, 71 pp.

USFWS [U.S. Fish and Wildlife Service]. 2022b. Species Status Assessment report for the Suwannee Alligator Snapping Turtle (*Macrochelys suwanniensis*). Version 1.2. Atlanta, GA: USFWS, 112 pp.

USFWS [U.S. Fish and Wildlife Service]. 2022c. Interim Species Status Assessment/biological report for the northern population of the Bog Turtle (*Glyptemys muhlenbergii*). Version 1. Cortland, NY: USFWS, 141 pp.

USFWS [U.S. Fish and Wildlife Service]. 2023a. Endangered and threatened wildlife and plants; threatened species status with Section 4(d) Rule for Egyptian Tortoise. Federal Register 88:19004–19017.

USFWS [U.S. Fish and Wildlife Service]. 2023b. Species Status Assessment Report for the Pearl River Map Turtle (*Graptemys pearlensis*). Version 1.2. Atlanta, GA: USFWS, 152 pp.

USFWS [U.S. Fish and Wildlife Service]. 2023c. Species Status Assessment report for Northwestern Pond Turtle (*Actinemys marmorata*) and Southwestern Pond Turtle (*Actinemys pallida*). Version 1.1. Ventura, CA: USFWS, 183 pp.

USFWS [U.S. Fish and Wildlife Service]. 2023d. Endangered and threatened wildlife and plants; threatened species status with Section 4(d) Rule for the Northwestern Pond Turtle and Southwestern Pond Turtle. Federal Register 88:68370–68399.

USFWS [U.S. Fish and Wildlife Service]. 2024a. Endangered and threatened wildlife and plants; threatened status for the Suwannee Alligator Snapping Turtle with a section 4(d) rule. Federal Register 89:53507–53528.

USFWS [U.S. Fish and Wildlife Service]. 2024b. Endangered and threatened wildlife and plants; threatened species status for Pearl River Map Turtle with section 4(d) rule; and threatened species status for Alabama Map Turtle, Barbour's Map Turtle, Escambia Map Turtle, and Pascagoula Map Turtle due to similarity of appearance with section 4(d) rule. Federal Register 89: 57206–57236.

Vaillant, L. 1885a. Sur une tortue terrestre d'espèce nouvelle, rapportée par M. Humblot au Muséum d'Histoire Naturelle. Comptes Rendus de l'Academie des Sciences de Paris 101(6):440–441.

Vaillant, L. 1885b. Remarques complémentaires sur les tortues gigantesque de Madagascar. Comptes Rendus Hebdomadaires des Séances de l'Académie des Sciences, Paris 100:874–877.

Vaillant, L. 1889. Description d'une tortue terrestre d'especе nouvelle (*Testudo yniphora*). Nouvelles Archives du Muséum d'Histoire Naturelle (3)1:161–167.

Vaillant, L. 1894. Nouvelle espèce du genre *Geoemyda* trouvée au Tonkin par S.A. le Prince Henri d'Orléans. Bulletin de la Société Philomatique de Paris (8)6:68–69.

Vaillant, L. 1898. Dessins inédits de chéloniens tirés des manuscrits de Commerson. Bulletin du Musée National d'Histoire Naturelle, Paris 4:133–139.

Vaillant, L. 1911. Chéloniens et batracien urodèle, recueillis par M. le Dr. Rivet. In: Mission du Service Géographique de l'Armée pour la mesure d'un Arc de Méridien Équatorial en Amérique du Sud sous le contrôle Scientifique de l'Académie des Sciences, 1899-1906. Tome 9. Zoologie. Fascicule 2. Reptiles – Poissons – Batraciens. Paris: Ministère de l'Instruction Publique, pp. 45–60.

Vaillant, L. and Grandidier, G. 1910. Histoire naturelle des Reptiles. Première partie: Crocodiles et Tortues. In: Grandidier, A. and Grandidier, G. (Eds.). Histoire Physique, Naturelle et Politique de Madagascar. Vol. 17. Paris: Hachette, 86 pp.

Valakos, E.D., Pafilis, P., Sotiropoulos, K., Lymberakis, P., Maragou, P., and Foufopoulos, J. 2008. The Amphibians and Reptiles of Greece. Frankfurt: Edition Chimaira, 463 pp.

Valdez-Villavicencio, J.H., Peralta-García, A., and Guillen-González, J.A. 2016a. Nueva población de la tortuga de poza del suroeste *Emys pallida* en el Desierto Central de Baja California, México. Revista Mexicana de Biodiversidad 87:264–266.

Valdez-Villavicencio, J.H., Peralta-García, A., Galina-Tessaro, P., and Hollingsworth, B.D. 2016b. Notes on the reproduction of the Southwestern Pond Turtle *Emys pallida* in Baja California, México. Revista Mexicana de Herpetología 2(1):36–39.

Valenciennes, A. 1833. [*Chelonia pelasgorum, Emys hellenica, Emys iberica,*

Emys rivulata]. In: Bory de Saint-Vincent, J.B. (Ed.). Expédition Scientifique de Morée. Travaux de la Section des Sciences Physiques. Zoologie. Paris: F.G. Levrault, planches, troisième série, pls. 6–17.

Valenti, P., Vlachos, E., Kehlmaier, C., Fritz, U., Georgalis, G.L., Luján, À.H., Miccichè, R., Sineo, L., and Delfino, M. 2022. The last of the large-sized tortoises of the Mediterranean islands. Zoological Journal of the Linnean Society 196:1704–1717.

Valenzuela, A., Cau, M.A., and Alcover, J.A. 2016. Archaeological evidence for the introduction of *Emys orbicularis* (Testudines: Emydidae) in the Balearic Islands. Amphibia-Reptilia 37:229–236.

Vamberger, M. and Fritz, U. 2018. Big data can cause big mistakes: using the Societas Europaea Herpetologica atlas by Sillero et al. (2014), the distribution of *Emys orbicularis* will be misunderstood. Biologia 73:281–283.

Vamberger, M., Corti, C., Stuckas, H., and Fritz, U. 2011. Is the imperilled spur-thighed tortoise (*Testudo graeca*) native in Sardinia? Implications from population genetics and for conservation. Amphibia-Reptilia 32:9–25.

Vamberger, M., Stuckas, H., Ayaz, D., Graciá, E., Aloufi, A.A., Els, J., Mazanaeva, L.F., Kami, H.G., and Fritz, U. 2013. Conservation genetics and phylogeography of the poorly known Middle Eastern terrapin *Mauremys caspica* (Testudines: Geoemydidae). Organisms Diversity & Evolution 13(1):77–85.

Vamberger, M., Stuckas, H., Ayaz, D., Lymberakis, P., Široký, P., and Fritz, U. 2014. Massive transoceanic gene flow in a freshwater turtle (Testudines: Geoemydidae: *Mauremys rivulata*). Zoologica Scripta 43:313–322.

Vamberger, M., Stuckas, H., Sacco, F., D'Angelo, S., Arculeo, M., Cheylan, M., Corti, C., Lo Valvo, M., Marrone, F., Wink, M., and Fritz, U. 2015. Differences in gene flow in twofold secondary contact zone of pond turtles in southern Italy (Testudines: Emydidae: *Emys orbicularis galloitalica, E. o. hellenica, E. trinacris*). Zoologica Scripta 44:233–249.

Vamberger, M., Stuckas, H., Vargas-Ramírez, M., Kehlmaier, C., Ayaz, D., Aloufi, A.A., Lymberakis, P., Široký, P., and Fritz, U. 2017a. Unexpected hybridization patterns in Near Eastern terrapins (*Mauremys caspica, M. rivulata*) indicate ancient gene flow across the Fertile Crescent. Zoologica Scripta 46:401–413.

Vamberger, M., Durkin, L., Kim, C., Handschuh, M., Seng, R., and Fritz, U. 2017b. The leaf turtle population of Phnom Kulen National Park (northwestern Cambodia) has genetic and morphological signatures of hybridization. Journal of Zoological Systematics and Evolutionary Research 55:167–174.

Vamberger, M., Hofmeyr, M.D., Ihlow, F., and Fritz, U. 2018. In quest of contact: phylogeography of helmeted terrapins (*Pelomedusa galeata, P. subrufa* sensu stricto). PeerJ 6:e4901; doi 10.7717/peerj.4901, 21 pp.

Vamberger, M., Hofmeyr, M.D., Cook, C.A., Netherlands, E.C., and Fritz, U. 2019a. Phylogeography of the East African Serrated Hinged Terrapin *Pelusios sinuatus* (Smith, 1838) and resurrection of *Sternothaerus bottegi* Boulenger, 1895 as a subspecies of *P. sinuatus*. Amphibian and Reptile Conservation 13(2)[Special Section]:42–56 (e184).

Vamberger, M., Anunciação, P.R., Hofmeyr, M.D., and Fritz, U. 2019b. Mind the gap—is the distribution range of Pelomedusa galeata really disjunct in western South Africa? Amphibian & Reptile Conservation 13(2) [Special Section]:57–60 (e185).

Vamberger, M., Ihlow, F., Asztalos, M., Dawson, J.E., Jasinski, S.E., Praschag, P., and Fritz, U. 2020a. So different, yet so alike: North American slider turtles (*Trachemys scripta*). Vertebrate Zoology 70:87–96.

Vamberger, M., Spitzweg, C., De Silva, A., Masroor, R., Praschag, P., and Fritz, U. 2020b. Already too late? Massive trade in Indian Star Tortoises (*Geochelone elegans*) might have wiped out its phylogenetic differentiation. Amphibia-Reptilia 41:133–138.

Van Denburgh, J. 1895. A review of the herpetology of lower California. Part I – Reptiles. Proceedings of the California Academy of Sciences (2)5:77–162.

Van Denburgh, J. 1907. Expedition of the California Academy of Sciences to the Galapagos Islands, 1905–1906. I. Preliminary descriptions of four new races of gigantic land tortoises from the Galapagos Islands. Proceedings of the California Academy of Sciences (4)1:1–6.

Van Denburgh, J. 1914. Expedition of the California Academy of Sciences to the Galapagos Islands, 1905–1906. X. The gigantic land tortoises of the Galapagos Archipelago. Proceedings of the California Academy of Sciences (4)2(1):203–374.

Van Devender, T.R., Ferguson, G.M., Villa, R.A., and Turner, D.S. 2024. The Box Turtles (Terrapene, Emydidae) of Sonora, Mexico. Sonoran Herpetologist 37(1):12–18.

van der Kuyl, A.C., Ballasina, D.L.P., Dekker, J.T., Maas, J., Willemsen, R.E., and Goudsmit, J. 2002. Phylogenetic relationships among the species of the genus *Testudo* (Testudines: Testudinidae) inferred from mitochondrial 12S rRNA gene sequences. Molecular Phylogenetics and Evolution 22: 174–183.

van der Kuyl, A.C., Ballasina, D.L.P., and Zorgdrager, F. 2005. Mitochondrial haplotype diversity in the tortoise species *Testudo graeca* from North Africa and the Middle East. BMC Evolutionary Biology 5:1–8.

van Loben Sels, R.C., Congdon, J.D., and Austin, J.T. 1997. Life history and ecology of the Sonoran mud turtle (*Kinosternon sonoriense*) in southeastern Arizona: a preliminary report. Chelonian Conservation and Biology 2:338–344.

Van-Ernest, H. 1801. [*Testudo rugosa, Testudo melanocephala*]. In: Daudin, F.M. Histoire Naturelle, Générale et Particulière des Reptiles. Tome Second. Paris: Dufart, 432 pp. [pp. 37–38, 128].

Vandellius, D. 1761. Epistola de Holothurio, et Testudine Coriacea ad Celeberrimum Carolum Linnaeum. Patavii [Padova]: Conzatti, 12 pp.

Vandewege, M.W., Gutierrez, J., Davis, D.R., Forstner, M.R.J., and Mali, I. 2024. Patterns of genetic divergence in the Rio Grande cooter (*Pseudemys gorzugi*), a riverine turtle inhabiting an arid and anthropogenically modified system. Journal of Heredity 115:253–261.

Vanzolini, P.E. 1981. The scientific and political contexts of the Bavarian expedition to Brasil. In: Spix, J.B. and Wagler, J.G. Herpetology of Brazil. Society for the Study of Amphibia and Reptilia, Facsimile Reproductions in Herpetology, pp. ix–xxix.

Vanzolini, P.E. 1994. On the distribution of certain South American turtles (Testudines: Testudinidae and Chelidae). Smithsonian Herpetological Information Service 97:1–10.

Vanzolini, P.E. 1995. A new species of turtle, genus *Trachemys*, from the State of Maranhão, Brazil (Testudines, Emydidae). Revista Brasileira de Biologia 55(1):111–125.

Vargas-Ramírez, M., Castaño-Mora, O.V., and Fritz, U. 2008. Molecular phylogeny and divergence times of ancient South American and Malagasy river turtles (Testudines: Pleurodira: Podocnemididae). Organisms, Diversity and Evolution 8:388–398.

Vargas-Ramírez, M., Maran, J., and Fritz, U. 2010a. Red- and yellow-footed tortoises, *Chelonoidis carbonaria* and *C. denticulata* (Reptilia: Testudines: Testudinidae), in South American savannahs and forests: do their phylogeographies reflect distinct habitats? Organisms, Diversity and Evolution 10:161–172.

Vargas-Ramírez, M., Vences, M., Branch, W.R., Daniels, S.R., Glaw, F., Hofmeyr, M.D., Kuchling, G., Maran, J., Papenfuss, T.J., Široky, P., Vieites, D.R., and Fritz, U. 2010b. Deep genealogical lineages in the widely distributed African helmeted terrapin: evidence from mitochondrial and nuclear DNA (Testudines: Pelomedusidae: *Pelomedusa subrufa*). Molecular Phylogenetics and Evolution 56(1):428–440.

Vargas-Ramírez, M., Michels, J., Castaño-Mora, O.V., Cárdenas-Arevalo, G., Gallego-García, N., and Fritz, U. 2012a. Weak genetic divergence between the two South American toad-headed turtles *Mesoclemmys dahli* and *M. zuliae* (Testudines: Pleurodira: Chelidae). Amphibia-Reptilia 33:373–385.

Vargas-Ramírez, M., Stuckas, H., Castaño-Mora, O.V., and Fritz, U. 2012b. Extremely low genetic diversity and weak population differentiation in the endangered Colombian river turtle *Podocnemis lewyana* (Testudines: Podocnemididae). Conservation Genetics 13:65–77.

Vargas-Ramírez, M., Carr, J.L., and Fritz, U. 2013. Complex phylogeography in *Rhinoclemmys melanosterna*: conflicting mitochondrial and nuclear evidence suggests past hybridization (Testudines: Geoemydidae). Zootaxa 3670(2):238–254.

Vargas-Ramírez, M., del Valle, C., Ceballos, C.P., and Fritz, U. 2017. *Trachemys medemi* n. sp. from northwestern Colombia turns the biogeography of South American slider turtles upside down. Journal of Zoological Systematics and Evolutionary Research 55:326–339.

Vargas-Ramírez, M., Caballero, S., Morales-Betancourt, M.A., Lasso, C.A., Amaya, L., Martínez, J.G., Viana, M.d.N.S., Vogt, R.C., Farias, I.P., Hrbek, T., Campbell, P.D., and Fritz, U. 2020. Genomic analyses reveal two species of the matamata (Testudines: Chelidae: *Chelus* spp.) and clarify their phylogeography. Molecular Phylogenetics and Evolution 148:106823, 17 pp.

Vasconcelos, R., Brito, J.C., Carranza, S., and Harris, D.J. 2013. Review of the distribution and conservation status of the terrestrial reptiles of the Cape Verde Islands. Oryx 47(1):77–87.

Vasilyev, V.A., Bondarenko, D.A., Peregontsev, E.A., Voronov, A.S., Ryskov, A.P., and Semenova, S.K. 2008. Polymorphism of the 12S rRNA gene and phylogeography of the Central Asian tortoises *Agrionemys horsfieldii* Gray, 1844. Russian Journal of Genetics 44:682–685.

Vasilyev, V.A., Korsunenko, A.V., Pereshkolnik, S.L., Mazanaeva, L.F., Bannikova, A.A., Bondarenko, D.A., Peregontsev, E.A., and Semyenova, S.K. 2014. Differentiation of tortoises of the genera *Testudo* and *Agrionemys* (Testudinidae) based on the polymorphism of nuclear and mitochondrial markers. Russian Journal of Genetics 50:1060–1074.

Vásquez-Cruz, V., Cazares-Hernández, E., Reynoso-Martínez, A., Kelly-Hernández, A., Fuentes-Moreno, A., and Lara-Hernández, F.A. 2021. New distributional records of freshwater turtles from west-central Veracruz, Mexico. Reptiles & Amphibians 28(1):146–151.

Vecchioni, L., Marrone, F., Arculeo, M., Fritz, U., and Vamberger, M. 2020. Stand out from the crowd: small-scale genetic structuring in the endemic Sicilian Pond Turtle. Diversity 12(343): doi:10.3390/d12090343.

Velo-Antón, G. and Pinya, S. 2015. El galápago leproso (*Mauremys leprosa*) en la península ibérica e islas Baleares. Boletín de la Asociación Herpetológica Española 26:39–42.

Velo-Antón, G., García-París, M., and Cordero Rivera, A. 2008. Patterns of nuclear and mitochondrial DNA variation in Iberian populations of *Emys orbicularis* (Emydidae): conservation implications. Conservation Genetics 9:1263–1274.

Velo-Antón, G., Wink, M., Schneeweiss, N., and Fritz, U. 2011. Native or not? Tracing the origin of wild-caught and captive freshwater turtles in a threatened and widely distributed species (*Emys orbicularis*). Conservation Genetics 12:583–588.

Venzmer, G. 1920. Beobachtungen an der iberischen und an der kaspischen Schildkröte in Cilicien. Zoologischer Anzeiger 51:285–302.

Vergara-Ríos, D., Montes-Correa, A.C., Jiménez-Bolaño, J.D., Saboyá-Acosta, L., and Renjifo, J.M. 2015. Record of the largest size for a male of Magdalena River Turtle, *Podocnemis lewyana* Duméril, 1852 (Testudines: Podocnemididae). Herpetology Notes 8:335–337.

Veríssimo, J., Znari, M., Stuckas, H., Fritz, U., Pereira, P., Teixeira, J., Arculeo, M., Marrone, F., Sacco, F., Naimi, M., Kehlmaier, C., and Velo-Antón, G. 2016. Pleistocene diversification in Morocco and recent demographic expansion in the Mediterranean Pond Turtle *Mauremys leprosa*. Biological Journal of the Linnean Society 119:943–959.

Vetter, H. 2004. Turtles of the World, Vol. 2: North America. Frankfurt: Edition Chimaira, 127 pp.

Vetter, H. and van Dijk, P.P. 2006. Turtles of the World, Vol. 4: East and South Asia. Frankfurt: Edition Chimaira, 160 pp.

Viana, M.d.N.S., Oliveira, J.d.A., Agostini, M.A.P., Erickson, J., Matias De Morais, G., Monjeló, L.A.d.S., Andrade, P.C.M., Félix-Silva, D., Junior, W.P.d.O., Sites, J.W., Jr., Vogt, R.C., Hrbek, T., and Farias, I.P. 2017. Population genetics structure of the threatened Amazon River Turtle, *Podocnemis sextuberculata*. Chelonian Conservation and Biology 16:128–138.

Vilaça, S.T., Vargas, S.M., Lara-Ruiz, P., Molfetti, E., Reis, E.C., Lôbo-Hajdu, G., Soares, L.S., and Santos, F.R. 2012. Nuclear markers reveal a complex introgression pattern among marine turtle species on the Brazilian coast. Molecular Ecology 21(17):4300–4312.

Vilaça, S.T., Hahn, A.T., Naro-Maciel, E., Abreu-Grobois, F.A., Bowen, B.W., Castilhos, J.C., Ciofi, C., FitzSimmons, N.N., Jensen, M.P., Formia, A., Limpus, C.J., Natali, C., Soares, L.S., De Thoisy, B., Whiting, S.D., and Bonatto, S.L. 2022. Global phylogeography of ridley sea turtles (*Lepidochelys* spp.): evolution, demography, connectivity, and conservation. Conservation Genetics 23:995–1010.

Vilaró, J. 1867a. Notas. In: Poey, F. Repertorio Fisico-Natural de la Isla de Cuba 2(5):119–122.

Vilaró, J. 1867b. Nota sobre las jicoteas cubanas. In: Poey, F. Repertorio Fisico-Natural de la Isla de Cuba 2(9):204.

Viloria, N.A. and Forti, M. 2015 ["2012"]. Abundancia, estructura de tallas y proporción de sexos de *Podocnemis vogli* (Müller, 1935) en el hato Masaguaral, estado Guárico, Venezuela. Memoria de la Fundación La Salle de Ciencias Naturales 72(177–178) (2012):21–33.

Vinke, T. and Vinke, S. 1999. Ein wahrer Gigant – Rekordgrösse für die Köhlerschildkröte *Geochelone carbonaria* (Spix, 1824). Emys 6(2):27–29.

Vinke, S., Vetter, H., Vinke, T., and Vetter, S. 2008. South American Tortoises – *Chelonoidis carbonaria, C. denticulata* and *C. chilensis*. Frankfurt: Chimaira Verlag, Chelonian Library 3, 360 pp.

Vinke, T., Vinke, S., and Köhler, G. 2013. What is known about *Mesoclemmys vanderhaegei* (Bour, 1973): a systematic review of the available literature. Paraquaria Natural 1(2):21–31.

Viñola-López, L.W. and Almonte, J.N. 2022. Revision of the fossil land tortoises (Testudines: Testudinidae) from Hispaniola with the description of a new species. Novitates Caribaea 20:11–29.

Vitek, N.S. 2018. Delineating modern variation from extinct morphology in the fossil record using shells of the Eastern Box Turtle (*Terrapene carolina*). PLos One 13(3):e0193437.

Vlachos, E. 2015. Forming taxon names from Greek words. Bionomina 9: 1–26.

Vlachos, E. 2018. A review of the fossil record of North American turtles of the clade Pan-Testudinoidea. Bulletin of the Peabody Museum of Natural History 59(1):3–94.

Vlachos, E. 2022. The fossil record of turtles from the Pleistocene of Crete (Greece). Comptes Rendus Palevol 21(35):771–799.

Vlachos, E. and Rabi, M. 2018. Total evidence analysis and body size evolution of extant and extinct tortoises (Testudines: Cryptodira: Pan-Testudinidae). Cladistics 34(6):652–683.

Vogt, R.C. 1980. Natural history of the map turtles *Graptemys pseudogeographica* and *Graptemys ouachitensis* in Wisconsin. Tulane Studies in Zoology and Botany 22:17–48.

Vogt, R.C. 1995. *Graptemys pseudogeographica*. Catalogue of American Amphibians and Reptiles 604:1–6.

Vogt, R.C., Thomson, S.A., Rhodin, A.G.J., Pritchard, P.C.H., Mittermeier, R.A., and Baggi, N. 2013. Case 3587. *Podocnemis unifilis* Troschel, 1848 (Reptilia,Testudines): proposed precedence over *Emys cayennensis* Schweigger, 1812. Bulletin of Zoological Nomenclature 70(1):33–39.

Vogt, R.C., Oliveira, S.M., and Vieira, S.C. 2020. *Mesoclemmys raniceps* (Amazon Toad-headed Turtle). Coloration. Herpetological Review 51:317.

Vogt, T. 1911. Reptilien und Amphibien aus Neu-Guinea. Sitzungsberichte der Gesellschaft der Naturforschender Freunde, Berlin 9:410–414.

Vuillemin, S. 1972a. Note sur *Testudo morondavaensis* n. sp. Annales de l'Université de Madagascar, Série Sciences de la Nature et Mathématiques 9:127–134.

Vuillemin, S. 1972b. Note sur *Madakinixys domerguei* n. gen. n. sp. (Testudinidae). Annales de l'Université de Madagascar, Série Sciences de la Nature et Mathématiques 9:169–182.

Vuillemin, S. and Domergue, C. 1972. Contribution à la faune de Madagascar: description de *Pyxoides brygooi* gen. et sp. nov. (Testudinidae). Annales de l'Université de Madagascar, Série Sciences de la Nature et Mathématiques 9:193–200.

Vyas, R. 2011. Record size of Indian Star Tortoise, *Geochelone elegans* (Schoepff, 1795). Russian Journal of Herpetology 18:47–50.

Vyas, R. 2017. A Northern River Terrapin (*Batagur baska*) from Kutch, with comments on the species' distribution in western India and Pakistan. IRCF Reptiles and Amphibians 24(2):128–131.

Wagler, J.G. 1821. Die Amphibien (Lieferung 1). Nürnberg: J.B. Geyer, 12 pp., 6 plates.

Wagler, J.G. 1828. Vorläufige Uebersicht des Gerüstes, so wie Ankündigung seines Systema Amphibiorum. Isis von Oken 21(8):859–863.

Wagler, J.G. 1830a. Descriptiones et Icones Amphibiorum. Tres partes cum XXXVI tabulis. Part II. Monachii [München]: J.G. Cottae, pls. XIII-XXIV.

Wagler, J.G. 1830b. Natürliches System der Amphibien, mit vorangehender Classification der Säugthiere und Vögel. Ein Beitrag zur vergleichenden Zoologie. München: J.G. Cotta'schen Buchhandlung, 354 pp., pls. 1–2.

Wagler, J.G. 1830c. Natürliches System der Amphibien, mit vorangehender Classification der Säugthiere und Vögel. Tafeln. München: J.G. Cotta'schen Buchhandlung. Erstes Heft, pls. I–VII.

Wagler, J.G. 1833. Descriptiones et Icones Amphibiorum. Tres partes cum XXXVI tabulis. Part III. Monachii [München]: J.G. Cottae, pls. XXV–XXXVI.

Wagner, P., Bauer, A.M., Leviton, A.E., Wilms, T.M., and Böhme, W. 2016. A checklist of the amphibians and reptiles of Afghanistan: exploring herpetodiversity using biodiversity archives. Proceedings of the California Academy of Sciences (4)63:457–565.

Walbaum, J.J. 1782. Chelonographia oder Beschreibung einiger Schildkröten. Lubeck: J.F. Gleditsch, 132 pp.

Walbaum, J.J. 1785. Beschreibung der Spenglerischen Schildkröte. Schriften der Berliner Gesellschaft Naturforschender Freunde 6:122–131.

Walker, D. and Avise, J.C. 1998. Principles of phylogeography as illustrated by freshwater and terrestrial turtles in the southeastern United States. Annual Review of Ecology and Systematics 29:23–58.

Walker, D., Burke, V.J., Barak, I., and Avise, J.C. 1995. A comparison of mtDNA restriction sites vs. control region sequences on phylogeographic assessment of the musk turtle (*Sternotherus minor*). Molecular Ecology 4:365–373.

Walker, D., Orti, G., and Avise, J.C. 1998. Phylogenetic distinctiveness of a threatened aquatic turtle (*Sternotherus depressus*). Conservation Biology 12:639–645.

Wallace, B.P., Posnik, Z.A., Hurley, B.J., Dimatteo, A.D., Bandimere, A., Rodriguez, I., Maxwell, S.M., Meyer, L., Brenner, H., Jensen, M.P., Lacasella, E., Shamblin, B.M., Abreu-Grobois, F.A., Stewart, K.R., Dutton, P.H., Barrios-Garrido, H., Dalleau, M., Dell'amico, F., Eckert, K.L., FitzSimmons, N.N., Garcia-Cruz, M., Hays, G.C., Kelez, S., Lagueux, C.J., Madden Hof, C.A., Marco, A., Martins, S.L.T., Mobaraki, A., Mortimer, J.A., Nel, R., Phillott, A.D., Pilcher, N.J., Putman, N.F., Rees, A.F., Rguez-Baron, J.M., Seminoff, J.A., Swaminathan, A., Turkozan, O., Vargas, S.M., Vernet, P.D., Vilaça, S., Whiting, S.D., Hutchinson, B.J., Casale, P., and Mast, R.B. 2023. Marine turtle regional management units 2.0: an updated framework for conservation and research of wide-ranging megafauna species. Endangered Species Research 52:209–223.

Wallin, L. 1977. The Linnean type-specimen of *Testudo geometrica*. Zoon 5:77–78.

Wallin, L. 1985. A survey of Linnaeus's material of *Chelone mydas*, *Caretta caretta* and *Eretmochelys imbricata* (Reptilia, Cheloniidae). Zoological Journal of the Linnean Society 85:121–130.

Wallin, L. 2001. Uppsala University, Museum of Evolution, Zoology section (UUZM): Catalogue of type specimens. 4. Linnaean specimens. Uppsala, Sweden: Uppsala University, 12 pp.

Wang, J., Shi, H.-T., Wen, C., and Han, L.-X. 2013. Habitat selection and conservation suggestions for the Yangtze Giant Softshell Turtle (*Rafetus swinhoei*) in the Upper Red River, China. Chelonian Conservation and Biology 12(1):177–184.

Wang, K., Lyu, Z.T., Wang, J., Qi, S., Che, J. 2022. The updated checklist and zoogeographic division of the reptilian fauna of Yunnan Province, China. Biodiversity Science 30(4):21326, 31 pp, doi: 10.17520/biods.2021326.

Wang, L., Zhou,X., Nie, L., Xia, X., Liu, L., Jiang, Y., Huang, Z., and Jing, W. 2012. The complete mitochondrial genome sequences of *Chelodina rugosa* and *Chelus fimbriata* (Pleurodira: Chelidae): implications of a common absence of initiation sites (OL) in pleurodiran turtles. Molecular Biology Reports 39(3):2097–2107.

Wangyal, J.T., Wangchuk, D., and Das, I. 2012. First report of turtles from the Himalayan Kingdom of Bhutan. Chelonian Conservation and Biology 11(2):268–272.

Ward, J.P. 1980. Comparative cranial morphology of the freshwater turtle subfamily Emydinae: an analysis of the feeding mechanisms and systematics. Ph.D. Thesis, North Carolina State University, Raleigh.

Ward, J.P. 1984. Relationships of chrysemyd turtles of North America (Testudines: Emydinae). Special Publications of the Museum of Texas Tech University 21:1–50.

Ward, R., Babitzke, J.B., and Killebrew, F.C. 2013. Genetic population structure of Cagle's Map Turtle (*Graptemys caglei*) in the Guadalupe and San Marcos Rivers of Texas – a landscape perspective. Copeia 2013:723–728.

Wariss, M., Isaac, V.J., and Pezzuti, J.C.B. 2012. Habitat use, size structure and sex ratio of the spot-legged turtle, *Rhinoclemmys punctularia punctularia* (Testudines: Geoemydidae), in Algodoal-Maiandeua Island, Pará, Brazil. Revista Biología Tropical 60:413–424.

Waters, J.H. 1964. Subspecific intergradation in the Nantucket Island, Massachusetts, population of the turtle *Chrysemys picta*. Copeia 1964(3):550–553.

Waters, J.H. 1969. Additional observations of southeastern Massachusetts insular and mainland populations of painted turtles, *Chrysemys picta*. Copeia 1969(1):179–182.

Weaver, S., McGaugh, S.E., Kono, T.J.Y., Macip-Rios, R., and Gluesenkamp, A.G. 2022. Assessing genomic and ecological differentiation among subspecies of the Rough-Footed Mud Turtle, *Kinosternon hirtipes*. Journal of Heredity 113:538–551.

Webb, R.G. 1959. Description of a new softshell turtle from the southeastern United States. University of Kansas Publications, Museum of Natural History 11(9):517–525.

Webb, R.G. 1960. Type and type locality of the Gulf Coast spiny softshell turtle, *Trionyx spinifer asper* (Agassiz). Breviora 129:1–8.

Webb, R.G. 1962. North American Recent soft-shelled turtles (family Trionychidae). University of Kansas Publications, Museum of Natural History 13:429–611.

Webb, R.G. 1973. *Trionyx spiniferus*. Catalogue of American Amphibians and Reptiles 140:1–4.

Webb, R.G. 1975a. Types of two names of African softshell turtles of the genus *Cyclanorbis* (Testudines: Trionychidae). Herpetologica 31(3):348–350.

Webb, R.G. 1975b. Taxonomic status of *Aspidonectes californiana* Rivers, 1889 (Testudines, Trionychidae). Copeia 1975(4):771–773.

Webb, R.G. 1978. *Trionyx steindachneri* Siebenrock, 1906: proposed validation under the plenary powers (Reptilia, Testudines). Bulletin of Zoological Nomenclature 35:47–48.

Webb, R.G. 1980a. The identity of *Testudo punctata* Lacepède, 1788 (Testudines, Trionychidae). Bulletin du Musée National d'Histoire Naturelle, Paris (4):2A:547–557.

Webb, R.G. 1980b. Gray, Hardwicke, Buchanan-Hamilton, and drawings of Indian softshell turtles (family Trionychidae). Amphibia-Reptilia 1:61–74.

Webb, R.G. 1982. Taxonomic notes concerning the trionychid turtle *Lissemys punctata* (Lacepède). Amphibia-Reptilia 3(2/3):179–184.

Webb, R.G. 1985. Taxonomic status of *Testudo rostrata* Thunberg, 1787 (Testudines, Trionychidae). Herpetologica 41(1):84–88.

Webb, R.G. 1990. *Trionyx sinensis* Wiegmann, 1834 (Reptilia, Testudines): proposed conservation of the specific name. Bulletin of Zoological Nomenclature 47(2):122–123.

Webb, R.G. 1993. *Emys* Duméril, 1806 (Reptilia, Testudines): proposed conservation. Bulletin of Zoological Nomenclature 50(3):224–227.

Webb, R.G. 1995a. Redescription and neotype designation of *Pelochelys bibroni* from southern New Guinea (Testudines: Trionychidae). Chelonian Conservation and Biology 1(4):301–310.

Webb, R.G. 1995b. The date of publication of Gray's Catalogue of Shield Reptiles. Chelonian Conservation and Biology 1(4):322–323.

Webb, R.G. 2003 ["2002"]. Observations on the giant softshell turtle, *Pelochelys cantorii*, with description of a new species. Hamadryad 27(1)(2002) [2003]:99–107.

Webb, R.G. and Legler, J.M. 1960. A new softshell turtle (genus *Trionyx*) from Coahuila, Mexico. University of Kansas Science Bulletin 40(2):21–30.

Webb, R.G. and van Dijk, P.P. 2004. Comments on the Narrow-headed Softshell Turtle (*Chitra chitra*) (Testudines, Trionychidae). Hamadryad 29(1):94–100.

Weisrock, D.W. and Janzen, F.J. 2000. Comparative molecular phylogeography of North American softshell turtles (*Apalone*): implications for regional and wide-scale historical evolutionary forces. Molecular Phylogenetics and Evolution 14:152–164.

Weissinger, H. 1987. *Testudo graeca anamurensis* ssp. nov. aus Kleinasien. ÖGH-Nachrichten, Wien 10/11:14–18.

Weitzman, C.L., Hagerty, B.E., Sandmeier, F.C., and Tracy, C.R. 2021. Desert Tortoises in Zion National Park represent a natural extension of their range. Chelonian Conservation and Biology 20(1):91–96.

Wells, R.W. 2002a. A new subspecies of *Carettochelys* (Reptilia: Carettochelydidae) from northern Australia – *Carettochelys insculpta canni* ssp. nov. Australian Biodiversity Record 2002(1):1–7.

Wells, R.W. 2002b. Taxonomic notes on some Australian freshwater turtles of the genera *Chelodina* and *Elseya* (Reptilia: Chelidae). Australian Biodiversity Record 2002(2):1–30.

Wells, R.W. 2007a. Some taxonomic and nomenclatural considerations on the Class Reptilia in Australia. Notes on the recently described freshwater turtle *Chelodina canni* McCord and Thomson, 2002 and a redescription of *Chelodina rankini* Wells and Wellington, 1985. Australian Biodiversity Record 2007(1):1–5.

Wells, R.W. 2007b. Some taxonomic and nomenclatural considerations on the Class Reptilia in Australia. Some comments on the *Elseya dentata* (Gray, 1863) complex with redescriptions of the Johnstone River snapping turtle, *Elseya stirlingi* Wells and Wellington, 1985 and the Alligator Rivers snapping turtle, *Elseya jukesi* Wells 2002. Australian Biodiversity Record 2007(2):1–12.

Wells, R.W. 2007c. Some taxonomic and nomenclatural considerations on the Class Reptilia in Australia. A new genus of the family Chelidae from eastern Australia. Australian Biodiversity Record 2007(3):1–13.

Wells, R.W. 2009. Some taxonomic and nomenclatural considerations on the

Class Reptilia in Australia. A new species of freshwater turtle in the Genus *Wollumbinia* Wells 2007 (Reptilia: Chelidae) from eastern Australia. Australian Biodiversity Record 2009(1):1–12.

Wells, R.W. and Wellington, C.R. 1985. A classification of the Amphibia and Reptilia of Australia. Australian Journal of Herpetology, Supp. Ser. 1:1–61.

Wermuth, H. 1952. *Testudo hermanni robertmertensi* n. subsp. und ihr Vorkommen in Spanien. Senckenbergiana 33:157–164.

Wermuth, H. 1956. Versuch der Deutung einiger bisher übersehener Schildkröten-Namen. Zoologische Beiträge, Berlin (N.F.) 2(2-3):399–423.

Wermuth, H. 1958. Status und Nomenklatur der Maurischen Landschildkröte, *Testudo graeca*, in SW-Asien und NO-Afrika. Senckenbergiana Biologica 39:149–153.

Wermuth, H. 1965. Zum Status von *Testudo hypselonota* Bourret. Israel Journal of Zoology 14:277–285.

Wermuth, H. 1969. Eine neue Grosskopfschildkröte, *Platysternon megacephalum vogeli* n. ssp. Aquarien und Terrarien Zeitschrift 22(12):372–374.

Wermuth, H. and Mertens, R. 1961. Schildkröten. Krokodile. Brückenechsen. Jena: Gustav Fischer Verlag, 422 pp.

Wermuth, H. and Mertens, R. 1977. Liste der rezenten Amphibien und Reptilien: Testudines, Crocodylia, Rhynchocephalia. Tierreich 100:1–174.

Werner, F. 1897. Die Reptilien und Amphibien Oesterreich-Ungarns und der Occupationsländer. Wien: A. Pichler's Witwe und Sohn, 160 pp.

Werner, F. 1899. Beitrage zur Kenntniss der Reptilien- und Batrachier fauna der Balkanhalbinsel. Wissenschaften und Mitteilungen aus Bosnien und der Hercegovina 6:817–841.

Werner, F. 1901a. Ueber Reptilien und Batrachier aus Ecuador und Neu-Guinea. Verhandlungen der Zoologisch-Botanischen Gesellschaft, Wien 51:593–603.

Werner, F. 1901b. Neue Reptilien des Königsberger zoologischen Museums. Zoologischer Anzeiger 24:297–301.

Werner, Y.L., Korolker, N., Sion, G., and Göçmen, B. 2016. Bergmann's and Rensch's rules and the spur-thighed tortoise (*Testudo graeca*). Biological Journal of the Linnean Society 117(4):796–811.

Whitaker, N. and Vijaya, J. 2009. Biology of the Forest Cane Turtle, *Vijayachelys silvatica*, in South India. Chelonian Conservation and Biology 8(2):109–115.

White, A.W. and Archer, M. 1994. *Emydura lavarackorum*, a new Pleistocene turtle (Pleurodira: Chelidae) from fluviatile deposits at Riversleigh, northwestern Queensland. Records of the South Australian Museum 27(2):159–167.

White, A.W., Worthy, T.H., Hawkins, S., Bedford, S., and Spriggs, M. 2010. Megafaunal meiolaniid horned turtles survived until early human settlement in Vanuatu, Southwest Pacific. Proceedings of the National Academy of Sciences (PNAS) 107(41):15512–15516.

White, G. 1836. The Natural History and Antiquities of Selborne. Edited by E.T. Bennett. London: J. and A. Arch, 640 pp.

White, M. 2015. Largest Tortoise. https://www.guinnessworldrecords.com/news/60at60/2015/8/2002-largest-tortoise-392870.

Whittock, P.A., Pendoley, K.L., and Hamann, M. 2014. Inter-nesting distribution of flatback turtles *Natator depressus* and industrial development in Western Australia. Endangered Species Research 26:25–38.

Wied-Neuwied, M. zu. 1825a. Beitrage zur Naturgeschichte von Brasilien. I. Band. Weimar: Landes, 614 pp.

Wied-Neuwied, M. zu. 1825b. Abbildungen zur Naturgeschichte Brasiliens. Recueil de planches coloriées d'animaux du Brésil. Neunte Lieferung. Weimar: Landes-Industrie-Comptoir, 6 pp.

Wied-Neuwied, M. zu. 1839. Reise in das innere Nord-America in den Jahren 1832 bis 1834. Erster Band. Coblenz: J. Hoelscher, 653 pp.

Wied-Neuwied, M. zu. 1865. Verzeichniss der Reptilien, welche auf einer Reise im nördlichen America beobachtet wurden. Nova Acta Physico-Medica Academiae Caesareae Leopoldino-Carolinae Naturae Curiosorum 32:1–143.

Wiedemann, C.R.W. 1802. Anatomische Beschreibung der Schildkröten überhaupt und der getäfelten Schildkröte (*T. tessellata* Schneid. *T. tabulata* Walbaum) insbesondere. Archiv für Zoologie und Zootomie 2(2):177–210.

Wiegmann, A.F.A. 1828. Beyträge zur Amphibienkunde. Isis von Oken 21(3):364–383.

Wiegmann, A.F.A. 1834 ["1835"]. Beiträge zur Zoologie, gesammelt auf einer Reise um die Erde von Dr. F.J.F. Meyen. Siebente Abhandlung. Amphibien. Nova Acta Physico-Medica Academiae Caesareae Leopoldino-Carolinae Naturae Curiosorum 17:183–268.

Wieland, G.R. 1923. A new Parana pleurodiran. American Journal of Science (5)5:1–14.

Wiens, J.J., Kuczynski, C.A., and Stephens, P.A. 2010. Discordant mitochondrial and nuclear gene phylogenies in emydid turtles: implications for speciation and conservation. Biological Journal of the Linnean Society 99:445–461.

Williams, E.E. 1950a. Variation and selection in the cervical central articulations of living turtles. Bulletin of the American Museum of Natural History 94:505–562.

Williams, E.E. 1950b. *Testudo cubensis* and the evolution of Western Hemisphere tortoises. Bulletin of the American Museum of Natural History 95:1–36.

Williams, E.E. 1952. A new fossil tortoise from Mona Island, West Indies, and a tentative arrangement of the tortoises of the world. Bulletin of the American Museum of Natural History 99:545–560.

Williams, E.E. 1954. A key and description of the living species of the genus *Podocnemis* (sensu Boulenger) (Testudines, Pelomedusidae). Bulletin of the Museum of Comparative Zoology 111(8):279–295.

Williams, E.E. 1960. Two species of tortoises in northern South America. Breviora 120:1–13.

Wilmé, L., Waeber, P.O., and Ganzhorn, J.U. 2017. Human translocation as an alternative hypothesis to explain the presence of giant tortoises on remote islands in the south-western Indian Ocean. Journal of Biogeography 44:1–7.

Wilson, D.S., Mushinsky, H.R., and McCoy, E.D. 2006. *Kinosternon baurii* – striped mud turtle. In: Meylan, P.A. (Ed.). Biology and Conservation of Florida Turtles. Chelonian Research Monographs 3:180–188.

Wilson, S. and Swann, G. 2013. Complete Guide to the Reptiles of Australia. Chatswood, Australia: New Holland.

Wink, M., Guicking, D., and Fritz, U. 2001. Molecular evidence for hybrid origin of *Mauremys iversoni* Pritchard and McCord, 1991, and *Mauremys pritchardi* McCord, 1997 (Reptilia: Testudines: Bataguridae). Zoologische Abhandlungen, Staatliches Museum für Tierkunde, Dresden 51:41–49.

Wischuf, T. and Fritz, U. 1996. Eine neue Unterart der Bachschildkröte (*Mauremys caspica ventrimaculata* subsp. nov.) aus dem Iranischen Hochland. Salamandra 32(2):113–122.

Wischuf, T. and Fritz, U. 1997. [*Mauremys caspica siebenrocki*]. In: Fritz, U. and Wischuf, T. Zur Systematik westasiatisch–südosteuropäischer Bachschildkröten (Gattung *Mauremys*) (Reptilia: Testudines: Bataguridae). Zoologische Abhandlungen, Staatliches Museum für Tierkunde Dresden 49(13):223–260. [pp. 240–243].

Witzell, W.N. 1983. Synopsis of biological data on the hawksbill turtle *Eretmochelys imbricata* (Linnaeus, 1766). FAO Fisheries Synopsis 137:1–78.

Wolak, M.E., Gilchrist, G.W., Ruzicka, V.A., Nally, D.M., and Chambers, R.M. 2010. A contemporary, sex-limited change in body size of an estuarine turtle in response to commercial fishing. Conservation Biology 24:1268–1277.

Wong, R.A., Fong, J.J., and Papenfuss, T.J. 2010. Phylogeography of the African helmeted terrapin, *Pelomedusa subrufa*: genetic structure, dispersal, and human introduction. Proceedings of the California Academy of Sciences (4)61:575–585.

Wood, R.C. 1976. *Stupendemys geographicus*, the world's largest turtle. Breviora 436:1–31.

Worm, O. 1655. Museum Wormianum, seu Historia Rerum Rariorum. Lugdunum Batavorum: Johannem Elsevirium, 389 pp.

Worrell, E. 1970. Reptiles of Australia. Second Edition. Sydney: Angus and Robertson, 169 pp.

Worthy, T.H., Anderson, A.J., and Molnar, R.E. 1999. Megafaunal expression in a land without mammals – the first fossil faunas from terrestrial deposits in Fiji (Vertebrata: Amphibia, Reptilia, Aves). Senckenbergiana Biologica 79(2):237–242.

Wright, K.M. and Andrews, J.S. 2002. Painted turtles (*Chrysemys picta*) of Vermont: an examination of phenotypic variation and intergradation. Northeastern Naturalist 9(4):363–380.

Wussow, W. 1916. Meine Erfahrungen mit *Testudo horsfieldi*. Wochenschrift für Aquarien- und Terrarienkunde 13:169–172.

Wüster, W., Thomson, S.A., O'Shea, M., and Kaiser, H. 2021. Confronting taxonomic vandalism in biology: conscientious community self-organization can preserve nomenclatural stability. Biological Journal of the Linnean Society 133:645–670; https://doi.org/10.1093/biolinnean/blab009.

Xiao, F., Bu, R., Lin, L., Mueti, J., Wang, J., Ye, Z., and Shi, H. 2021. A survey of freshwater turtles in Diaoluoshan Nature Reserve with conservation implications for the endangered Big-Headed Turtle. Chelonian Conservation and Biology 20(1):139–143.

Xiaoyou, H., Xiaodan, C., Chen, C., Xiaoli, L., Jian, Z., Quanbo, Q., and Xinping, Z. 2019. Conservation status of the Asian Giant Softshell Turtle (*Pelochelys*

cantorii) in China. Chelonian Conservation and Biology 18(1):68–74.

Xiong, G., Wang, X.-Q., Zhou, X.-W., Zeng, D., Chen, Z.-N., Wang, P., and Kang, L. 2019. Genetic variation in the Chinese soft-shell turtles (*Pelodiscus* spp.) revealed by sequences of mitochondrial cytochrome b gene. Mitochondrial DNA Part A 30(8):874-879.

Yadava, M.R. and Prasad, B. 1982. Observations on the breeding biology of the Indian tropical pond turtle, *Lissemys punctata granosa* (Schoepff). Indian Journal of Zootomy 23(1):51–56.

Yakovleva, I.D. 1961. Opredelitel Presmykayushchikhsya Kirgizii. [Guide to Reptiles of Kirgizia.]. Frunze: Academy of Sciences Kirgiz SSR, 110 pp. [in Russian].

Yang, J., Ge, Y., Wei, Y., Duan, H., and Gong, S. 2018. Diversity and conservation of amphibians and reptiles in Guangdong Nanxiong Xiaoliukeng-Qingzhangshan Provincial Nature Reserve, Guangdong, China. Ecological Science 37(2):35–42.

Yang, P., Tang, Y., Ding, L., Guo, X., and Wang, Y. 2011. Validity of *Pelodiscus parviformis* (Testudines: Trionychidae) inferred from molecular and morphological analyses. Asian Herpetological Research 2(1):21–29.

Yasukawa, Y. and Ota, H. 1999. Geographic variation and biogeography of the geoemydine turtles (Testudines: Bataguridae) of the Ryukyu Archipelago, Japan. In: Ota, H. (Ed.). Tropical Island Herpetofauna: Origin, Current Diversity, and Conservation. Amsterdam: Elsevier, pp. 271–297.

Yasukawa, Y., Ota, H., and Hikida, T. 1992. Taxonomic re-evaluation of the two subspecies of *Geoemyda spengleri* (Gmelin, 1789) (Reptilia: Emydidae). Japanese Journal of Herpetology 14(3):143–159.

Yasukawa, Y., Ota, H., and Iverson, J.B. 1996. Geographic variation and sexual size dimorphism in *Mauremys mutica* (Cantor, 1842) (Reptilia: Bataguridae), with description of a new subspecies from the southern Ryukyus, Japan. Zoological Science (Japan) 13:303–317.

Yazarloo, M., Kami, H.G., and Bagherian Yazdi, A. 2017. Sexual dimorphism and morphometric study of Caspian pond turtle, *Mauremys caspica* (Testudines: Geoemydidae) in Golestan Province, southeast of the Caspian Sea. Caspian Journal of Environmental Science 15:321–334.

Yeh, H.-K. 1961. The first discovery of a box-turtle in China. Vertebrata Palasiatica 5:58–64.

Young, M.J., Unmack, P.J., Gruber, B., Gleeson, D.M., Eisemberg, C.C., Amepou, Y., Ibana, D., and Georges, A. 2025. Historical biogeography and genetic status of the enigmatic Pig-nosed Turtle (*Carettochelys insculpta*) within the Australo-Papuan region. Diversity and Distributions 31:e70015, 19 pp.

Young-Valencia, K., Fernando Ortega, A., and Botero-Botero, Á. 2014. Densidad y estructura de las poblaciones de tortuga pímpano (*Chelydra acutirostris* Peters 1862) (Chelydridae) en las quebradas Cajones y Los Coclí, departamento del Quindío, Colombia. Revista Biodiversidad Neotropical 4(2):149–161.

Yu, X.M., Lin, Y.F., Peng, L.F., Zhang, Y., Lu, S.Q., and Huang, S. 2019. The complete mitochondrial genome of *Pelodiscus axenaria* (Testudines: Trionychidae). Mitochondrial DNA Part B 4(2):2154–2155.

Yudha, D.S., Eprilurahman, R., Irwanjasmoro, and Supramono, Y. 2019. Survei awal analisa habitat ditemukannya Labi-Labi Bintang (*Chitra chitra*) di Sungai Sempor, Sleman, DIY. Warta Herpetofauna 11(1):25–33. [in Indonesian]

Zangerl, R. 1947. Redescription of *Taphrosphys olssoni*, a fossil turtle from Peru. Fieldiana (Geology) 10:29–40.

Zangerl, R. 1948. The vertebrate fauna of the Selma formation of Alabama. Part II. The Pleurodiran turtles. Fieldiana: Geology Memoirs 3(2):19–56.

Zangerl, R. 1958. Die oligozänen Meerschildkröten von Glarus. Schweizerische Paläontologische Abhandlungen 73:1–56.

Zangerl, R. and Medem, F. 1958. A new species of chelid turtle, *Phrynops (Batrachemys) dahli*, from Colombia. Bulletin of the Museum of Comparative Zoology 119:375–390.

Zangerl, R. and Turnbull, W.D. 1955. *Procolpochelys grandaeva* (Leidy), an early carettine sea turtle. Fieldiana Zoology 37:345–382.

Zappalorti, R.T. 2023. The Bog Turtle: Natural History and Conservation. Rodeo, NM: ECO Publishing, 188 pp.

Zappalorti, R.T. and Iverson, J.B. 2006. *Sternotherus minor* – loggerhead musk turtle. In: Meylan, P.A. (Ed.). Biology and Conservation of Florida Turtles. Chelonian Research Monographs 3:197–206.

Zavialov, E.V., Tabachishin, V.G., and Shlyakhtin, G.V. 2003. Recent distribution habitat of reptiles in the North of Low Volga region. Modern Herpetology 2:52–67. [in Russian]

Zenboudji, S., Cheylan, M., Arnal, V., Bertolero, A., Leblois, R., Astruc, G., Bertorelle, G., Pretus, J.L., Lo Valvo, M., Sotgiu, G., and Montgelard, C. 2016. Conservation of the endangered Mediterranean tortoise *Testudo hermanni hermanni*: the contribution of population genetics and historical demography. Biological Conservation 195:279–291.

Zhang, M. 1984. [A new species of *Pelochelys* from Zhejiang, with subfossil description]. Acta Herpetologica Sinica 3(4):71–76. [in Chinese]

Zhang, M., Zong, Y., and Ma, J. 1998. Fauna Sinica. Reptilia Vol. 1. General Accounts of Reptilia. Testudoformes and Crocodiliformes. Beijing: Science Press, 213 pp.

Zhang, X., Unmack, P.J., Kuchling, G., Wang, Y., and Georges, A. 2017. Resolution of the enigmatic phylogenetic relationship of the critically endangered Western Swamp Tortoise *Pseudemydura umbrina* (Pleurodira: Chelidae) using a complete mitochondrial genome. Molecular Phylogenetics and Evolution; doi:10.1016/j.ympev.2017.07.019.

Zhao, E. 1990. [*Cuora zhoui*]. In: Zhao, E., Zhou, T., and Ye, P. [A new Chinese box turtle (Testudinata: Emydidae)–*Cuora zhoui*]. In: Zhao, E. (Ed.). From Water Onto Land. Chinese Society for the Study of Amphibians and Reptiles, Beijing, pp. 213–216. [p. 213]. [in Chinese]

Zhao, E. and Adler, K. 1993. Herpetology of China. Society for the Study of Amphibians and Reptiles, Contributions to Herpetology, No. 10, 522 pp.

Zhao, E., Zhou, J., and Zhou, T. (Eds.) 1997. Chinese Chelonian Research. Chinese Society for the Study of Amphibians and Reptiles, Herpetological Series No. 9, Sichuan Journal of Zoology 15 (Supplement), 159 pp.

Zhao, J., Li, W., Zhang, D., Wen, P., and Zhu, X. 2016a. The mitochondrial genomes of three lineages of Asian yellow pond turtle *Mauremys mutica*. Mitochondrial DNA Part A: DNA Mapping, Sequencing, and Analysis 27:2466–2467.

Zhao, J., Li, W., Wen, P., Zhang, D., and Zhu, X. 2016b. Genetic diversity and relationship of *Mauremys mutica* and *M. annamensis* assessed by DNA barcoding sequences. Mitochondrial DNA Part A: DNA Mapping, Sequencing, and Analysis 27:3507–3510.

Zhao, Z., Heideman, N., Grobler, P., Jordaan, A., Bester, P., and Hofmeyr, M.D. 2020a. Unraveling the diversification and systematic puzzle of the highly polymorphic *Psammobates tentorius* (Bell, 1828) complex (Reptilia: Testudinidae) through phylogenetic analyses and species delimitation approaches. Journal of Zoological Systematics and Evolutionary Research 58(1):308–326.

Zhao, Z., Heideman, N., and Hofmeyr, M.D. 2020b. Codon-based analysis of selection pressure and genetic structure in the *Psammobates tentorius* (Bell, 1828) species complex, and phylogeny inferred from both codons and amino acid sequences. African Journal of Ecology 2020; https://doi.org/10.1111/aje.12840, 13 pp.

Zhao, Z., Heideman, N., Oosthuizen, J., and Hofmeyr, M.D. 2020c. Testing alternative phylogenetic hypotheses for the Tent Tortoise species complex (Reptilia, Testudinidae) using multiple data types and methods. bioRxiv preprint, doi:10.1101/2020.11.02.364745, 65 pp.

Zhao, Z., Heideman, N., Bester, P., Jordaan, A., and Hofmeyr, M.D. 2020d. Climatic and topographic changes since the Miocene influenced the diversification and biogeography of the tent tortoise (*Psammobates tentorius*) species complex in southern Africa. BMC Evolutionary Biology 20(153):1–33.

Zhao, Z., Oosthuizen, J., and Heideman, N. 2021. How many species does the *Psammobates tentorius* (tent tortoise) species complex (Reptilia, Testudinidae) comprise? A taxonomic solution potentially applicable to species complexes. Journal of Zoological Systematics and Evolutionary Research 2021, doi:10.1111/jzs.12525, 23 pp.

Zheng, C., Nie, L., Wang, J., Zhou, H., Hou, H., Wang, H., and Liu, J. 2013. Recombination and evolution of duplicate control regions in the mitochondrial genome of the Asian Big-headed Turtle, *Platysternon megacephalum*. PLoS ONE 8(12): e82854.

Zhou, G., Zhang, X., and Fang, Z. 1991. Bulletin of a new species *Trionyx*. Acta Scientiarum Naturalium Universitatis Normalis Hunanensis, Hunan Changsha 14(4):379–382. [in Chinese].

Zhou, H., Jiang, Y., Nie, L., Yin, H., Li, H., Dong, X., Zhao, F., Zhang, H., Pu, Y., Huang, Z., Song, J., and Sun, E. 2015. The historical speciation of *Mauremys* sensu lato: ancestral area reconstruction and interspecific gene flow level assessment provide new insights. PLoS One 10:e0144711.

Zhou, T. 2005. Discovery of a living male Yunnan box turtle, *Cuora yunnanensis* Boulenger, 1906. Sichuan Journal of Zoology 24(3):345–346.

Zhou, T. and Li, P. 2013. Chinese Tortoise and Turtle Classification – Primary Colour Encyclopedia. Beijing: China Agriculture Press, 286 pp. [in Chinese].

Zhou, T. and Zhao, E. 2004. On the occurrence of living *Cuora yunnanensis* since fifty-eight years, and its description. Sichuan Journal of Zoology 23:325–327.

Zittel, K.A. 1889. Handbuch der Palaeontologie. I. Abtheilung Palaeozoologie. III. Band. Pisces, Amphibia, Reptilia, Aves. München and Leipzig: R. Oldenbourg Verlag, 900 pp.

Zong, Y. and Pan, L. 1989. Studies on the genus *Cuora* of the Testudoformes. In: Matsui, M., Hikida, T., and Goris, R.C. (Eds.). Current Herpetology in East Asia. Proceedings of the second Japan-China Herpetological Symposium, Kyoto, July 1988. Herpetological Society of Japan 1989:198.

Zug, G.R. 1977. The Matamata (Testudines: Chelidae) is *Chelus* not *Chelys*! Herpetologica 33:53–54.

Zug, G.R. 2022. Amphibians and Reptiles of Myanmar: Checklists and Keys. I. Amphibians, Crocodilians, and Turtles. Smithsonian Contributions to Zoology 653:1–113.

Zug, G.R. and Schwartz, A. 1971. *Deirochelys, D. reticularia*. Catalogue of American Amphibians and Reptiles 107:1–3.

Zug, G.R., Ernst, C.H., and Wilson, R.V. 1998. *Lepidochelys olivacea*. Catalogue of American Amphibians and Reptiles 653:1–13.

Zug, G.R., Beaver, K., Bjorndal, K.A., Crumly, C.R., Das, I., Diamond, A.W., Hambler, C., Leuteritz, T., Shah, N.J., Murphy, J.B., Swingland, I.R., Bourn, D., Türkozan, O., Savage, J.M., Caccone, G., Parham, J.F., and Rhodin, A.G.J. 2009. Comments on the proposed conservation of usage of *Testudo gigantea* Schweigger, 1812 (currently *Geochelone (Aldabrachelys) gigantea*; Reptilia, Testudines) (Case 3463; see BZN 66: 34). Bulletin of Zoological Nomenclature 66(1):80–87.

CHELONIAN RESEARCH MONOGRAPHS
Conservation Biology of Freshwater Turtles and Tortoises (CBFTT)
(2008–2025; Chronological)

All CBFTT publications (11 checklists and 128 species accounts covering 173 taxa) are available online (**www.iucn-tftsg.org/**) as open-access downloadable doi-designated pdf's.

Rhodin, A.G.J., Pritchard, P.C.H., van Dijk, P.P., Saumure, R.A., Buhlmann, K.A., Iverson, J.B., and Mittermeier, R.A. (Eds.). 2008–2016. Conservation Biology of Freshwater Turtles and Tortoises: A Compilation Project of the IUCN/SSC Tortoise and Freshwater Turtle Specialist Group. Chelonian Research Monographs, No. 5 (Installments 1–9a). Lunenburg, Massachusetts: Chelonian Research Foundation.

Rhodin, A.G.J., Iverson, J.B., van Dijk, P.P., Saumure, R.A., Buhlmann, K.A., Pritchard, P.C.H., and Mittermeier, R.A. (Eds.). 2016. Conservation Biology of Freshwater Turtles and Tortoises: A Compilation Project of the IUCN/SSC Tortoise and Freshwater Turtle Specialist Group. Chelonian Research Monographs, No. 5 (Installment 9b). Lunenburg, Massachusetts: Chelonian Research Foundation.

Rhodin, A.G.J., Iverson, J.B., van Dijk, P.P., Saumure, R.A., Buhlmann, K.A., Pritchard, P.C.H., and Mittermeier, R.A. (Eds.). 2017. Conservation Biology of Freshwater Turtles and Tortoises: A Compilation Project of the IUCN/SSC Tortoise and Freshwater Turtle Specialist Group. Chelonian Research Monographs, No. 7. In: Chelonian Research Monographs, No. 5 (Supplement No. 1). Lunenburg, Massachusetts: Chelonian Research Foundation and Turtle Conservancy.

Rhodin, A.G.J., Iverson, J.B., van Dijk, P.P., Buhlmann, K.A., Pritchard, P.C.H., and Mittermeier, R.A. (Eds.). 2017–2018. Conservation Biology of Freshwater Turtles and Tortoises: A Compilation Project of the IUCN/SSC Tortoise and Freshwater Turtle Specialist Group. Chelonian Research Monographs, No. 5 (Installments 10–11). Lunenburg, Massachusetts: Chelonian Research Foundation and Turtle Conservancy.

Rhodin, A.G.J., Iverson, J.B., van Dijk, P.P., Stanford, C.B., Goode, E.V., Buhlmann, K.A., Pritchard, P.C.H., and Mittermeier, R.A. (Eds.). 2018–2019. Conservation Biology of Freshwater Turtles and Tortoises: A Compilation Project of the IUCN/SSC Tortoise and Freshwater Turtle Specialist Group. Chelonian Research Monographs, No. 5 (Installments 12–13). Lunenburg, Massachusetts: Chelonian Research Foundation and Turtle Conservancy.

Rhodin, A.G.J., Iverson, J.B., van Dijk, P.P., Stanford, C.B., Goode, E.V., Buhlmann, K.A., and Mittermeier, R.A. (Eds.). 2020–2025. Conservation Biology of Freshwater Turtles and Tortoises: A Compilation Project of the IUCN/SSC Tortoise and Freshwater Turtle Specialist Group. Chelonian Research Monographs, No. 5 (Installments 14–19). Arlington, Vermont: Chelonian Research Foundation and Turtle Conservancy.

CBFTT Checklists
(2007–2025, Chronological)

All checklists are available online as open-access downloadable doi-designated pdf's at **www.iucn-tftsg.org/checklist/**.

TTWG [Turtle Taxonomy Working Group: Bickham, J.W., Iverson, J.B., Parham, J.F., Philippen, H.D., Rhodin, A.G.J., Shaffer, H.B., Spinks, P.Q., and van Dijk, P.P.]. 2007b. An annotated list of modern turtle terminal taxa with comments on areas of taxonomic instability and recent change. Chelonian Research Monographs 4:173–199.

Rhodin, A.G.J., van Dijk, P.P., and Parham, J.F. 2008. Turtles of the world: annotated checklist of taxonomy and synonymy. Chelonian Research Monographs 5(1):000.1–38.

TTWG [Turtle Taxonomy Working Group: Rhodin, A.G.J., Parham, J.F., van Dijk, P.P., and Iverson, J.B.]. 2009. Turtles of the world: annotated checklist of taxonomy and synonymy, 2009 update, with conservation status summary. Chelonian Research Monographs 5(2):000.39–84.

TTWG [Turtle Taxonomy Working Group: Rhodin, A.G.J., van Dijk, P.P., Iverson, J.B., and Shaffer, H.B.]. 2010. Turtles of the world, 2010

update: annotated checklist of taxonomy, synonymy, distribution, and conservation status. Chelonian Research Monographs 5(3):000.85–164.

TTWG [Turtle Taxonomy Working Group: van Dijk, P.P., Iverson, J.B., Shaffer, H.B., Bour, R., and Rhodin, A.G.J.]. 2011. Turtles of the world, 2011 update: annotated checklist of taxonomy, synonymy, distribution, and conservation status. Chelonian Research Monographs 5(4):000.165–242.

TTWG [Turtle Taxonomy Working Group: van Dijk, P.P., Iverson, J.B., Shaffer, H.B., Bour, R., and Rhodin, A.G.J.]. 2012. Turtles of the world, 2012 update: annotated checklist of taxonomy, synonymy, distribution, and conservation status. Chelonian Research Monographs 5(5):000.243–328.

TTWG [Turtle Taxonomy Working Group: van Dijk, P.P., Iverson, J.B., Rhodin, A.G.J., Shaffer, H.B., and Bour, R.]. 2014. Turtles of the world, 7th edition: annotated checklist of taxonomy, synonymy, distribution with maps, and conservation status. Chelonian Research Monographs 5(7):000.329–479.

TEWG [Turtle Extinctions Working Group: Rhodin, A.G.J., Thomson, S., Georgalis, G.L., Karl, H.-V., Danilov, I.G., Takahashi, A., de la Fuente, M.S., Bourque, J.R., Delfino, M., Bour, R., Iverson, J.B., Shaffer, H.B., and van Dijk, P.P.]. 2015. Turtles and tortoises of the world during the rise and global spread of humanity: first checklist and review of extinct Pleistocene and Holocene chelonians. Chelonian Research Monographs 5(8):000e.1–66.

TTWG [Turtle Taxonomy Working Group: Rhodin, A.G.J., Iverson, J.B., Bour, R., Fritz, U., Georges, A., Shaffer, H.B., and van Dijk, P.P.]. 2017. Turtles of the World: Annotated Checklist and Atlas of Taxonomy, Synonymy, Distribution, and Conservation Status (8th Ed.). Chelonian Research Monographs 7:1–292.

TTWG [Turtle Taxonomy Working Group: Rhodin, A.G.J., Iverson, J.B., Bour, R., Fritz, U., Georges, A., Shaffer, H.B., and van Dijk, P.P.]. 2021. Turtles of the World: Annotated Checklist and Atlas of Taxonomy, Synonymy, Distribution, and Conservation Status (9th Ed.). Chelonian Research Monographs 8:1–472.

TTWG [Turtle Taxonomy Working Group: Rhodin, A.G.J., Iverson, J.B., Fritz, U., Gallego-García, N., Georges, A., Shaffer, H.B., and van Dijk, P.P.]. 2025. Turtles of the World: Annotated Checklist and Atlas of Taxonomy, Synonymy, Distribution, and Conservation Status (10th Ed.). Chelonian Research Monographs 10:1–575.

CBFTT Species Accounts (2008–2025)

From 2008 to date, 267 authors have produced the following 128 species accounts covering 173 taxa of turtles and tortoises in the *Conservation Biology of Freshwater Turtles and Tortoises* (CBFTT) series published in 19 installments of *Chelonian Research Monographs* No. 5. All are available for open-access download at **www.iucn-tftsg.org/cbftt/**.

Alcalde, L., Sánchez, R.M., and Pritchard, P.C.H. 2021. *Hydromedusa tectifera* Cope 1870 – South American Snake-necked Turtle, Argentine Snake-necked Turtle, Tortuga Cuello de Vibora, Cágado Pescoço de Cobra. Chelonian Research Monographs 5(15):113.1–17.

Anders, B. and Iverson, J.B. 2012. *Mauremys nigricans* (Gray 1834) – Red-necked Pond Turtle, Chinese Red-necked Turtle, Kwangtung River Turtle, Black-necked Pond Turtle. Chelonian Research Monographs 5(5):068.1–9.

Auliya, M., van Dijk, P.P., Moll, E.O., and Meylan, P.A. 2016. *Amyda cartilaginea* (Boddaert 1770) – Asiatic Softshell Turtle, Southeast Asian Softshell Turtle. Chelonian Research Monographs 5(9a):092.1–17.

Baker, P.J., Diagne, T., and Luiselli, L. 2015. *Cyclanorbis elegans* (Gray 1869) – Nubian Flapshell Turtle. Chelonian Research Monographs 5(8):089.1–7.

Berry, J.F. and Iverson, J.B. 2011. *Kinosternon scorpioides* (Linnaeus 1766) – Scorpion Mud Turtle. Chelonian Research Monographs 5(4):063.1–15.

Berry, K.H. and Murphy, R.W. 2019. *Gopherus agassizii* (Cooper 1861) – Mojave Desert Tortoise, Agassiz's Desert Tortoise. Chelonian Research Monographs 5(13):109.1–45.

Bertolero, A. and Busack, S.D. 2017. *Mauremys leprosa* (Schoepff *in* Schweigger 1812) – Eastern Mud Turtle. Chelonian Research Monographs 5(10):102.1–19.

Bertolero, A., Cheylan, M., Hailey, A., Livoreil, B., and Willemsen, R.E. 2011. *Testudo hermanni* (Gmelin 1789) – Hermann's Tortoise. Chelonian Research Monographs 5(4):059.1–20.

Bhupathy, S., Webb, R.G, and Praschag, P. 2014. *Lissemys punctata* (Bonnaterre 1789) – Indian Flapshell Turtle. Chelonian Research Monographs 5(7):076.1–12.

Blanck, T., Iverson, J.B., Zhou, T., Meier, E., and McCord, W.P. 2024. *Cuora zhoui* Zhao *in* Zhao, Zhou, and Ye 1990 – Zhou's Box Turtle. Chelonian Research Monographs 5(18):127.1–16.

Blankenship, E.L., Butterfield, B.P., and Godwin, J.C. 2008. *Graptemys nigrinoda* Cagle 1954 – Black-knobbed Map Turtle, Black-knobbed Sawback. Chelonian Research Monographs 5(1):005.1–6.

Bock, B.C., Páez, V.P., and Daza, J.M. 2010. *Trachemys callirostris* (Gray 1856) – Colombian Slider, Jicotea, Hicotea, Galapago, Morrocoy de Agua. Chelonian Research Monographs 5(3):042.1–9.

Bour, R. 2008. *Pelusios adansonii* (Schweigger 1812) – Adanson's Mud Turtle. Chelonian Research Monographs 5(1):017.1–4.

Bour, R. and Gerlach, J. 2008. *Pelusios seychellensis* (Siebenrock 1906) – Seychelles Mud Turtle. Chelonian Research Monographs 5(1):018.1–3.

Bour, R., Luiselli, L., Petrozzi, F., Segniagbeto, G.H., and Chirio, L. 2016. *Pelusios castaneus* (Schweigger 1812) – West African Mud Turtle, Swamp Terrapin. Chelonian Research Monographs 5(9a):095.1–11.

Bower, D.S. and Hodges, K.M. 2014. *Chelodina expansa* Gray 1857 – Broad-Shelled Turtle, Giant Snake-Necked Turtle. Chelonian Research Monographs 5(7):071.1–8.

Boycott, R.C. and Bourquin, O. 2008. *Pelomedusa subrufa* (Lacepède 1788) – Helmeted Turtle, Helmeted Terrapin. Chelonian Research Monographs 5(1):007.1–6.

Broadley, D.G. and Boycott, R.C. 2008. *Pelusios rhodesianus* Hewitt 1927 – Variable Mud Turtle, Variable Hinged Terrapin. Chelonian Research Monographs 5(1):004.1–3.

Broadley, D.G. and Boycott, R.C. 2009. *Pelusios sinuatus* (Smith 1838) – Serrated Hinged Terrapin. Chelonian Research Monographs 5(2):036.1–5.

Broadley, D.G. and Sachsse, W. 2011. *Cycloderma frenatum* Peters 1854 – Zambezi Flapshell Turtle, Nkhasi. Chelonian Research Monographs 5(4):055.1–5.

Buhlmann, K.A., Gibbons, J.W., and Jackson, D.R. 2008. *Deirochelys reticularia* (Latreille 1801) – Chicken Turtle. Chelonian Research Monographs 5(1):014.1–6.

Bury, R.B. and Germano, D.J. 2008. *Actinemys marmorata* (Baird and Girard 1852) – Western Pond Turtle, Pacific Pond Turtle. Chelonian Research Monographs 5(1):001.1–9.

Buskirk, J.R. and Ponce-Campos, P. 2011. *Terrapene nelsoni* Stejneger 1925 – Spotted Box Turtle, Tortuga de Chispitas, Tortuga de Monte. Chelonian Research Monographs 5(4):060.1–9.

Cann, J., Spencer, R.-J., Welsh, M., and Georges, A. 2015. *Myuchelys georgesi* (Cann 1997) – Bellinger River Turtle. Chelonian Research Monographs 5(8):091.1–9.

Carr, J.L. and Giraldo, A. 2009. *Rhinoclemmys nasuta* (Boulenger 1902) – Large-nosed Wood Turtle, Chocoan River Turtle. Chelonian Research Monographs 5(2):034.1–6.

Carr, J.L., Moll, D., Giraldo, A., and Garcés-Restrepo, M.F. 2023. *Rhinoclemmys annulata* (Gray 1860) – Brown Wood Turtle, Montañero, Bambera. Chelonian Research Monographs 5(17):123.1–16.

Cerdá-Ardura, A., Soberón-Mobarak, F., McGaugh, S.E., and Vogt, R.C. 2008. *Apalone spinifera atra* (Webb and Legler 1960) – Black Spiny Softshell Turtle, Cuatrociénegas Softshell, Tortuga Concha Blanda, Tortuga Negra de Cuatrociénegas. Chelonian Research Monographs 5(1):021.1–4.

Congdon, J.D., Graham, T.E., Herman, T.B., Lang, J.W., Pappas, M.J., and Brecke, B.J. 2008. *Emydoidea blandingii* (Holbrook 1838) – Blanding's Turtle. Chelonian Research Monographs 5(1):015.1–12.

Das, I. 2008. *Pelochelys cantorii* Gray 1864 – Asian Giant Softshell Turtle. Chelonian Research Monographs 5(1):011.1–6.

Das, I. 2009. *Melanochelys tricarinata* (Blyth 1856) – Tricarinate Hill Turtle, Three-keeled Land Turtle. Chelonian Research Monographs 5(2):025.1–5.

Das, I. 2010. *Morenia ocellata* (Duméril and Bibron 1835) – Burmese Eyed Turtle. Chelonian Research Monographs 5(3):044.1–5.

Das, I. and Bhupathy, S. 2009a. *Hardella thurjii* (Gray 1831) – Crowned River Turtle. Chelonian Research Monographs 5(2):023.1–6.

Das, I. and Bhupathy, S. 2009b. *Melanochelys trijuga* (Schweigger 1812) – Indian Black Turtle. Chelonian Research Monographs 5(2):038.1–9.

Das, I. and Bhupathy, S. 2010. *Geoclemys hamiltonii* (Gray 1830) – Spotted Pond Turtle, Black Pond Turtle. Chelonian Research Monographs 5(3):043.1–6.

Das, I. and Sengupta, S. 2010. *Morenia petersi* Anderson 1879 – Indian Eyed Turtle. Chelonian Research Monographs 5(3):045.1–5.

Das, I. and Singh, S. 2009. *Chitra indica* (Gray 1830) – Narrow-headed Softshell Turtle. Chelonian Research Monographs 5(2):027.1–7.

Das, I., Basu, D., and Singh, S. 2010. *Nilssonia hurum* (Gray 1830) – Indian Peacock Softshell Turtle. Chelonian Research Monographs 5(3):048.1–6.

Das, I., Sengupta, S., and Praschag, P. 2010. *Pangshura sylhetensis* Jerdon 1870 – Assam Roofed Turtle. Chelonian Research Monographs 5(3):046.1–6.

Das, I., Sirsi, S., Vasudevan, K., and Murthy, B.H.C.K. 2014. *Nilssonia leithii* (Gray 1872) – Leith's Softshell Turtle. Chelonian Research Monographs 5(7):075.1–5.

Das, I., McCormack, T.E.M., van Dijk, P.P., Hoang, H.V., and Struijk, R.P.J.H. 2016. *Cuora mouhotii* (Gray 1862) – Keeled Box Turtle. Chelonian Research Monographs 5(9b):099.1–12.

Dawson, J.E., Ihlow, F., Ettmar, S., van Dijk, P.P., and Thirakhupt, K. 2018. *Malayemys macrocephala* (Gray 1859) – Malayan Snail-eating Turtle, Rice-field Terrapin. Chelonian Research Monographs 5(12):108.1–16.

Dawson, J.E., Ihlow, F., and Platt, S.G. 2020. *Malayemys subtrijuga* (Schlegel and Müller 1845) – Mekong Snail-Eating Turtle. Chelonian Research Monographs 5(14):111.1–24.

D'Cruze, N., Mookerjee, A., Vyas, R., Macdonald, D.W., and de Silva, A. 2018. *Geochelone elegans* (Schoepff 1795) – Indian Star Tortoise, Star Tortoise. Chelonian Research Monographs 5(12):106.1–13.

Deepak, V., Ramesh, M., Bhupathy, S., and Vasudevan, K. 2011. *Indotestudo travancorica* (Boulenger 1907) – Travancore Tortoise. Chelonian Research Monographs 5(4):054.1–6.

Deepak, V., Praschag, P., and Vasudevan, K. 2014. *Vijayachelys silvatica* (Henderson 1912) – Cochin Forest Cane Turtle. Chelonian Research Monographs 5(7):078.1–7.

Diesmos, A.C., Buskirk, J.R., Schoppe, S., Diesmos, M.L.L., Sy, E.Y., and Brown, R.M. 2012. *Siebenrockiella leytensis* (Taylor 1920) – Palawan Forest Turtle, Philippine Forest Turtle. Chelonian Research Monographs 5(5):066.1–9.

Dodd, C.K., Jr. 2008. *Sternotherus depressus* Tinkle and Webb 1955 – Flattened Musk Turtle. Chelonian Research Monographs 5(1):013.1–7.

Els, J., Alqahtani, A.R., Fritz, U., Shobrak, M., Busais, S.M., and Burriel-Carranza, B. 2022. *Pelomedusa barbata* Petzold, Vargas-Ramírez, Kehlmaier, Vamberger, Branch, Du Preez, Hofmeyr, Meyer, Schleicher, Široký, and Fritz 2014 – Arabian Helmeted Turtle, Arabian Marsh Turtle, Arabian Mud Turtle. Chelonian Research Monographs 5(16):118.1–10.

Ennen, J.R., Lovich, J.E., and Jones, R.L. 2016. *Graptemys pearlensis* Ennen, Lovich, Kreiser, Selman, and Qualls 2010 – Pearl River Map Turtle. Chelonian Research Monographs 5(9a):094.1–8.

Escoriza, D., Díaz-Paniagua, C., Andreu, A., and Ben Hassine, J. 2022. *Testudo graeca* Linnaeus 1758 (Western Subspecies Clade: *Testudo g. graeca, T. g. cyrenaica, T. g. marokkensis, T. g. nabeulensis, T. g. whitei*) – Mediterranean Spur-thighed Tortoise, Moorish Tortoise, Libyan Tortoise, Moroccan Tortoise, Tunisian Tortoise, Souss Valley Tortoise. Chelonian Research Monographs 5(16):117.1–18.

Escoriza, D., Tzoras, E., and Bringsøe, H. 2023. *Testudo marginata* Schoepff 1793 – Marginated Tortoise, Kraspedochelóna, Breshka Malore, Testuggine Marginata. Chelonian Research Monographs 5(17):122.1–25

Escoriza, D., Tzoras, E., Bringsøe, H., Iverson, J.B., and Rhodin, A.G.J. In review. Case 3881 – *Testudo marginata weissingeri* Trutnau, 1994 (Reptilia, Testudines): proposed suppression in favor of *Testudo weissingeri* Bour, 1996. Bulletin of Zoological Nomenclature.

Fielder, D., Chessman, B., and Georges, A. 2015. *Myuchelys bellii* (Gray 1844) – Western Saw-shelled Turtle, Bell's Turtle. Chelonian Research Monographs 5(8):088.1–7.

Forero-Medina, G. and Castaño-Mora, O.V. 2011. *Kinosternon scorpioides albogulare* (Duméril and Bocourt 1870) – White-throated Mud Turtle, Swanka Turtle. Chelonian Research Monographs 5(4):064.1–5.

Forero-Medina, G., Castaño-Mora, O.V., Cárdenas-Arevalo, G., and Medina-Rangel, G.F. 2013. *Mesoclemmys dahli* (Zangerl and Medem 1958) – Dahl's Toad-Headed Turtle, Carranchina, Tortuga Montañera. Chelonian Research Monographs 5(6):069.1–8.

Freeman, A. and Cann, J. 2014. *Myuchelys latisternum* (Gray 1867) – Sawshelled Turtle, Saw-Shell Turtle. Chelonian Research Monographs 5(7):073.1–8.

Freeman, A., Thomson, S., and Cann, J. 2014. *Elseya lavarackorum* (White and Archer 1994) – Gulf Snapping Turtle, Gulf Snapper, Riversleigh Snapping Turtle, Lavarack's Turtle. Chelonian Research Monographs 5(7):082.1–10.

Georges, A., Doody, J.S., Eisemberg, C., Alacs, E.A., and Rose, M. 2008. *Carettochelys insculpta* Ramsay 1886 – Pig-nosed Turtle, Fly River Turtle. Chelonian Research Monographs 5(1):009.1–17.

Gerlach, J. 2008a. *Pelusios castanoides intergularis* Bour 1983 – Seychelles Yellow-bellied Mud Turtle, Seychelles Chestnut-bellied Terrapin. Chelonian Research Monographs 5(1):010.1–4.

Gerlach, J. 2008b. *Pelusios subniger parietalis* Bour 1983 – Seychelles Black Mud Turtle. Chelonian Research Monographs 5(1):016.1–4.

Gerlach, J. 2009. *Aldabrachelys arnoldi* (Bour 1982) – Arnold's Giant Tortoise. Chelonian Research Monographs 5(2):028.1–5.

Gerlach, J. 2011b. *Aldabrachelys hololissa* (Günther 1877) – Seychelles Giant Tortoise. Chelonian Research Monographs 5(4):061.1–5.

Hagen, C., Platt, S.G., and Innis, C.J. 2009. *Leucocephalon yuwonoi* (McCord, Iverson, and Boeadi 1995) – Sulawesi Forest Turtle, Kura-kura Sulawesi. Chelonian Research Monographs 5(2):039.1–7.

Hailey, A., Broadley, D.G., Boycott, R.C., and Spawls, S. 2021. *Kinixys spekii* Gray 1863 – Speke's Hinge-back Tortoise, Speke's Hinged Tortoise. Chelonian Research Monographs 5(15):115.1–14.

Hofmeyr, M.D. 2009. *Chersina angulata* (Schweigger 1812) – Angulate Tortoise, South African Bowsprit Tortoise. Chelonian Research Monographs 5(2):030.1–6.

Howeth, J.G. and Brown, W.S. 2011. *Terrapene coahuila* Schmidt and Owens 1944 – Coahuilan Box Turtle. Chelonian Research Monographs 5(4):049.1–13.

Hughes, D.F., Allender, M.C., Bernstein, N.P., Iverson, J.B., Kolthoff, C., Martin, B.T., Meshaka, W.E., Jr., and Reed, B.M. 2024. *Terrapene ornata* (Agassiz 1857) – Ornate Box Turtle, Plains Box Turtle, Western Box Turtle, Desert Box Turtle, Tortuga de Caja Ornamentada. Chelonian Research Monographs 5(18):126.1–40.

Ihlow, F., Dawson, J.E., Hartmann, T., and Som, S. 2016. *Indotestudo elongata* (Blyth 1854) – Elongated Tortoise, Yellow-headed Tortoise, Yellow Tortoise. Chelonian Research Monographs 5(9a):096.1–14.

Iverson, J.B. and Christiansen, J.L. 2023. *Kinosternon flavescens* (Agassiz 1857) – Yellow Mud Turtle, Casquito Amarillo. Chelonian Research Monographs 5(17):121.1–26.

Iverson, J.B. and Vogt, R.C. 2011. *Kinosternon acutum* Gray 1831 – Tabasco Mud Turtle, Montera, Chechagua de Monte. Chelonian Research Monographs 5(4):062.1–6.

Iverson, J.B., Carr, J.L., Castaño-Mora, O.V., Galvis-Rizo, C.A., Rentería-Moreno, L.E., and Forero-Medina, G. 2012. *Kinosternon dunni* Schmidt 1947 – Dunn's Mud Turtle, Cabeza de Trozo. Chelonian Research Monographs 5(5):067.1–5.

Jackson, D.R. 2010. *Pseudemys nelsoni* Carr 1938 – Florida Red-bellied Turtle. Chelonian Research Monographs 5(3):041.1–8.

Johnston, G.R., Dodd, C.K., Jr., and Wilson, D.S. 2021. *Kinosternon baurii* (Garman 1891) – Striped Mud Turtle. Chelonian Research Monographs 5(15):116.1–18.

Jones, R.L. and Selman, W. 2009. *Graptemys oculifera* (Baur 1890) – Ringed Map Turtle, Ringed Sawback. Chelonian Research Monographs 5(2):033.1–8.

Kennett, R., Roe, J., Hodges, K., and Georges, A. 2009. *Chelodina*

longicollis (Shaw 1784) – Eastern Long-necked Turtle, Common Long-necked Turtle, Common Snake-necked Turtle. Chelonian Research Monographs 5(2):031.1–8.

Kennett, R., Fordham, D.A., Alacs, E., Corey, B., and Georges, A. 2014. *Chelodina oblonga* Gray 1841 – Northern Snake-Necked Turtle. Chelonian Research Monographs 5(7):077.1–13.

Kiester, A.R. and Willey, L.L. 2015. *Terrapene carolina* (Linnaeus 1758) – Eastern Box Turtle, Common Box Turtle. Chelonian Research Monographs 5(8):085.1–25.

Leary, C.J., Dobie, J.L., Mann, T.M., Floyd, P.S., and Nelson, D.H. 2008. *Pseudemys alabamensis* Baur 1893 – Alabama Red-bellied Cooter, Alabama Red-bellied Turtle. Chelonian Research Monographs 5(1):019.1–9.

Lindeman, P.V. 2008. *Sternotherus carinatus* Gray 1856 – Razorback Musk Turtle, Razor-backed Musk Turtle. Chelonian Research Monographs 5(1):012.1–6.

Lindeman, P.V. and Vogt, R.C. 2024. *Graptemys pseudogeographica* (Gray 1831) – False Map Turtle, Mississippi Map Turtle. Chelonian Research Monographs 5(18):124.1–19.

Lindeman, P.V., Stuart, J.N., and Killebrew, F.C. 2016. *Graptemys versa* Stejneger 1925 – Texas Map Turtle. Chelonian Research Monographs 5(9a):093.1–10.

Lovich, J.E., Godwin, J.C., and McCoy, C.J. 2011. *Graptemys ernsti* Lovich and McCoy 1992 – Escambia Map Turtle. Chelonian Research Monographs 5(4):051.1–6.

Lovich, J.E., Godwin, J.C., and McCoy, C.J. 2014. *Graptemys pulchra* Baur 1893 – Alabama Map Turtle. Chelonian Research Monographs 5(7):072.1–6.

Lovich, J.E., Selman, W., and McCoy, C.J. 2009. *Graptemys gibbonsi* Lovich and McCoy 1992 – Pascagoula Map Turtle, Pearl River Map Turtle, Gibbons' Map Turtle. Chelonian Research Monographs 5(2):029.1–8.

Lovich, J.E., Yasukawa, Y., and Ota, H. 2011. *Mauremys reevesii* (Gray 1831) – Reeves' Turtle, Chinese Three-keeled Pond Turtle. Chelonian Research Monographs 5(4):050.1–10.

Luiselli, L. and Diagne, T. 2013. *Kinixys homeana* Bell 1827 – Home's Hinge-Back Tortoise. Chelonian Research Monographs 5(6):070.1–10.

Luiselli, L. and Diagne, T. 2014. *Kinixys erosa* (Schweigger 1812) – Forest Hinge-back Tortoise, Serrated Hinge-back Tortoise, Serrated Hinged Tortoise. Chelonian Research Monographs 5(7):084.1–13.

Luiselli, L., Bour, R., Petrozzi, F., Akani, G.C., and Segniagbeto, G.H. 2018. *Pelusios niger* (Duméril and Bibron 1835) – West African Black Mud Turtle. Chelonian Research Monographs 5(12):105.1–8.

Magnusson, W.E. and Vogt, R.C. 2014. *Rhinemys rufipes* (Spix 1824) – Red Side-necked Turtle, Red-footed Sideneck Turtle, Perema. Chelonian Research Monographs 5(7):079.1–7.

Mantziou, G. and Rifai, L. 2014. *Mauremys rivulata* (Valenciennes *in* Bory de Saint-Vincent 1833) – Western Caspian Turtle, Balkan Terrapin. Chelonian Research Monographs 5(7):080.1–9.

Marques, T.S., Böhm, S., Brito, E.S., Cabrera, M.R., and Verdade, L.M. 2014. *Mesoclemmys vanderhaegei* (Bour 1973) – Vanderhaege's Toad-headed Turtle, Karumbé-hy. Chelonian Research Monographs 5(7):083.1–8.

McCormack, T.E.M., Dawson, J.E., Hendrie, D.B., Ewert, M.A., Iverson, J.B., Hatcher, R.E., and Goode, J.M. 2014. *Mauremys annamensis* (Siebenrock 1903) – Vietnamese Pond Turtle, Annam Pond Turtle, Rùa Trung Bộ. Chelonian Research Monographs 5(7):081.1–14.

McGovern, P., Diagne, T., Diagne, L., Luiselli, L., and Meylan, P.A. 2021. *Cyclanorbis senegalensis* (Duméril and Bibron 1835) – Sahelian Flapshell Turtle, Senegal Flapshell Turtle. Chelonian Research Monographs 5(15):114.1–10.

Meshaka, W.E., Jr., Gibbons, J.W., Hughes, D.F., Klemens, M.W., and Iverson, J.B. 2017. *Kinosternon subrubrum* (Bonnaterre 1789) – Eastern Mud Turtle. Chelonian Research Monographs 5(10):101.1–16.

Mittermeier, R.A., Vogt, R.C., Bernhard, R., and Ferrara, C.R. 2015. *Podocnemis erythrocephala* (Spix 1824) – Red-headed Amazon River Turtle, Irapuca. Chelonian Research Monographs 5(8):087.1–10.

Moll, E.O., Platt, K., Platt, S.G., Praschag, P., and van Dijk, P.P. 2009. *Batagur baska* (Gray 1830) – Northern River Terrapin. Chelonian Research Monographs 5(2):037.1–10.

Moll, E.O., Platt, S.G., Chan, E.H., Horne, B.D., Platt, K., Praschag, P., Chen, P.N., and van Dijk, P.P. 2015. *Batagur affinis* (Cantor 1847) – Southern River Terrapin, Tuntong. Chelonian Research Monographs 5(8):090.1–17.

Mwaya, R.T., Moll, D., Malonza, P.K., and Ngwava, J.M. 2018. *Malacochersus tornieri* (Siebenrock 1903) – Pancake Tortoise, Tornier's Tortoise, Soft-shelled Tortoise, Crevice Tortoise, Kobe Ya Mawe, Kobe Kama Chapati. Chelonian Research Monographs 5(12):107.1–15.

Ota, H., Yasukawa, Y., Fu, J., and Chen, T.H. 2009. *Cuora flavomarginata* (Gray 1863) – Yellow-margined Box Turtle. Chelonian Research Monographs 5(2):035.1–10.

Ottonello, D., D'Angelo, S., Marrone, F., Oneto, F., Spadola, F., Zuffi, M.A.L., and Fritz, U. 2021. *Emys trinacris* Fritz, Fattizzo, Guicking, Tripepi, Pennisi, Lenk, Joger, and Wink 2005 – Sicilian Pond Turtle, Testuggine Palustre Siciliana. Chelonian Research Monographs 5(15):112.1–13.

Páez, V.P., Restrepo, A., Vargas-Ramírez, M., and Bock, B.C. 2009. *Podocnemis lewyana* (Duméril 1852) – Magdalena River Turtle. Chelonian Research Monographs 5(2):024.1–6.

Petrozzi, F., Hema, E.M., Demaya, G.S., Benansio, J.S., Eniang, E.A., Diagne, T., Segniagbeto, G.H., and Luiselli, L. 2020. *Centrochelys sulcata* (Miller 1779) – African Spurred Tortoise, Grooved Tortoise, Sahel Tortoise, Tortue Sillonnée. Chelonian Research Monographs 5(14):110.1–16.

Pierce, L.J.S., Stuart, J.N., Ward, J.P., and Painter, C.W. 2016. *Pseudemys gorzugi* Ward 1984 – Rio Grande Cooter, Western River Cooter, Tortuga de Oreja Amarilla, Jicotéa del Río Bravo. Chelonian Research Monographs 5(9b):100.1–12.

Platt, S.G., Thanda Swe, Win Ko Ko, Platt, K., Khin Myo Myo, Rainwater, T.R., and Emmett, D. 2011. *Geochelone platynota* (Blyth 1863) – Burmese Star Tortoise, Kye Leik. Chelonian Research Monographs 5(4):057.1–9.

Platt, S.G., Platt, K., Win Ko Ko, and Rainwater, T.R. 2014. *Chitra vandijki* McCord and Pritchard 2003 – Burmese Narrow-Headed Softshell Turtle. Chelonian Research Monographs 5(7):074.1–7.

Pritchard, P.C.H. 2008. *Chelus fimbriata* (Schneider 1783) – Matamata Turtle. Chelonian Research Monographs 5(1):020.1–10.

Reyes-Grajales, E., Iverson, J.B., Walde, A.D., and Akre, T.S. 2025. *Kinosternon abaxillare* Baur *in* Stejneger 1925 – Central Chiapas Mud Turtle, Casquito Pardo. Chelonian Research Monographs 5(19):128.1–13.

Rhodin, A.G.J., Ibarrondo, B.R., and Kuchling, G. 2008. *Chelodina mccordi* Rhodin 1994 – Roti Island Snake-necked Turtle, McCord's Snake-necked Turtle, Kura-kura Rote. Chelonian Research Monographs 5(1):008.1–8.

Rhodin, A.G.J., Métrailler, S., Vinke, T., Vinke, S., Artner, H., and Mittermeier, R.A. 2009. *Acanthochelys macrocephala* (Rhodin, Mittermeier, and McMorris 1984) – Big-headed Pantanal Swamp Turtle, Pantanal Swamp Turtle. Chelonian Research Monographs 5(2):040.1–8.

Santana, D.O., Marques, T.S., Vieira, G.H.C., Moura, G.J.B., Faria, R.G., and Mesquita, D.O. 2016. *Mesoclemmys tuberculata* (Luederwaldt 1926) – Tuberculate Toad-headed Turtle. Chelonian Research Monographs 5(9b):097.1–8.

Schoppe, S. and Das, I. 2011. *Cuora amboinensis* (Riche *in* Daudin 1801) – Southeast Asian Box Turtle. Chelonian Research Monographs 5(4):053.1–13.

Selman, W. and Jones, R.L. 2011. *Graptemys flavimaculata* Cagle 1954 – Yellow-blotched Sawback, Yellow-blotched Map Turtle. Chelonian Research Monographs 5(4):052.1–11.

Souza, F.L. and Martins, F.I. 2009. *Hydromedusa maximiliani* (Mikan 1825) – Maximilian's Snake-necked Turtle, Brazilian Snake-necked Turtle. Chelonian Research Monographs 5(2):026.1–6.

Stanford, C.B., Wanchai, P., Schaffer, C., Schaffer, R., and Thirakhupt, K. 2015. *Manouria emys* (Schlegel and Müller 1840) – Asian Giant Tortoise, Giant Asian Forest Tortoise. Chelonian Research Monographs 5(8):086.1–9.

Stone, P.A., Congdon, J.D., Stone, M.E.B., Stuart, J.N., Iverson, J.B., and Rosen, P.C. 2022. *Kinosternon sonoriense* (LeConte 1854) – Sonora Mud Turtle, Desert Mud Turtle, Sonoyta Mud Turtle, Casquito de Sonora.

Chelonian Research Monographs 5(16):119.1–22.

Stuart, J.N. and Ward, J.P. 2009. *Trachemys gaigeae* (Hartweg 1939) – Big Bend Slider, Mexican Plateau Slider, Jicotea de la Meseta Mexicana. Chelonian Research Monographs 5(2):032.1–12.

Taskavak, E., Atatür, M.K., Ghaffari, H., and Meylan, P.A. 2016. *Rafetus euphraticus* (Daudin 1801) – Euphrates Softshell Turtle. Chelonian Research Monographs 5(9b):098.1–11.

Thomson, S., Kennett, R., Tucker, A., FitzSimmons, N.N., Featherston, P., Alacs, E.A., and Georges, A. 2011. *Chelodina burrungandjii* Thomson, Kennett, and Georges 2000 – Sandstone Snake-Necked Turtle. Chelonian Research Monographs 5(4):056.1–7.

Türkozan, O., Javanbakht, H., Mazanaeva, L., Meiri, S., Kornilev, Y.V., Tzoras, E., Popgeorgiev, G., Shanas, U., and Escoriza, D. 2023. *Testudo graeca* Linnaeus 1758 (Eastern Subspecies Clades: *Testudo g. armeniaca, Testudo g. buxtoni, Testudo g. ibera, Testudo g. terrestris, Testudo g. zarudnyi*) – Armenian Tortoise, Zagros Tortoise, Anatolian Tortoise, Levantine Tortoise, Kerman Tortoise. Chelonian Research Monographs 5(17):120.1–33.

Vamberger, M., Busack, S.D., Ayaz, D., Fritz, U., and Mazanaeva, L.F. 2024. *Mauremys caspica* (Gmelin 1774) – Caspian Turtle, Caspian Terrapin, West Asian Stripe-necked Terrapin. Chelonian Research Monographs 5(18):125.1–15.

Vinke, T., Vinke, S., Richard, E., Cabrera, M.R., Paszko, L., Marano, P., and Métrailler, S. 2011. *Acanthochelys pallidipectoris* (Freiberg 1945) – Chaco Side-necked Turtle. Chelonian Research Monographs 5(4):065.1–7.

Vogt, R.C. 2018. *Graptemys ouachitensis* Cagle 1953 – Ouachita Map Turtle. Chelonian Research Monographs 5(11):103.1–13.

Vogt, R.C., Platt, S.G., and Rainwater, T.R. 2009. *Rhinoclemmys areolata* (Duméril and Bibron 1851) – Furrowed Wood Turtle, Black-bellied Turtle, Mojena. Chelonian Research Monographs 5(2):022.1–7.

Vogt, R.C., Polisar, J.R., Moll, D., and Gonzalez-Porter, G. 2011. *Dermatemys mawii* Gray 1847 – Central American River Turtle, Tortuga Blanca, Hickatee. Chelonian Research Monographs 5(4):058.1–12.

Vogt, R.C., Bulté, G., and Iverson, J.B. 2018. *Graptemys geographica* (LeSueur 1817) – Northern Map Turtle, Common Map Turtle. Chelonian Research Monographs 5(11):104.1–18.

Ward, J.P. and Jackson, D.R. 2008. *Pseudemys concinna* (Le Conte 1830) – River Cooter. Chelonian Research Monographs 5(1):006.1–7.

Yasukawa, Y. and Ota, H. 2008. *Geoemyda japonica* Fan 1931 – Ryukyu Black-breasted Leaf Turtle, Okinawa Black-breasted Leaf Turtle. Chelonian Research Monographs 5(1):002.1–6.

Yasukawa, Y. and Ota, H. 2010. *Geoemyda spengleri* (Gmelin 1789) – Black-breasted Leaf Turtle. Chelonian Research Monographs 5(3):047.1–6.

Yasukawa, Y., Yabe, T., and Ota, H. 2008. *Mauremys japonica* (Temminck and Schlegel 1835) – Japanese Pond Turtle. Chelonian Research Monographs 5(1):003.1–6.

IUCN RED LIST AND GREEN LIST ASSESSMENTS (1996–2025)

From 1996 to date, the following members and associated colleagues and working groups of the IUCN Tortoise and Freshwater Turtle Specialist Group (TFTSG) and the IUCN Marine Turtle Specialist Group (MTSG) have produced the following 505 IUCN Red List assessments and 3 IUCN Green List assessments. We acknowledge and thank them all for their invaluable and important contributions.

Abreu-Grobois, A. and Plotkin, P. (IUCN SSC Marine Turtle Specialist Group). 2008. *Lepidochelys olivacea*. The IUCN Red List of Threatened Species 2008: e.T11534A3292503.

Ahmed, M.F. and Singh, S. 2021. *Morenia petersi*. The IUCN Red List of Threatened Species 2021: e.T13874A544248.

Ahmed, M.F., Horne, B.D., Li, P., Platt, K., Rahman, S., and Wang, L. 2020. *Cuora mouhotii*. The IUCN Red List of Threatened Species 2020: e.T163414A1006285.

Ahmed, M.F., Praschag, P., de Silva, A., Das, I., Singh, S., and de Silva, P. 2020. *Melanochelys trijuga*. The IUCN Red List of Threatened Species 2020: e.T13039A511745.

Ahmed, M.F., Praschag, P., and Singh, S. 2021. *Hardella thurjii*. The IUCN Red List of Threatened Species 2021: e.T9696A3152073.

Ahmed, M.F., Choudhury, B.C., Das, I., and Singh, S. 2021. *Nilssonia gangetica*. The IUCN Red List of Threatened Species 2021: e.T39618A2930943.

Ahmed, M.F., Praschag, P., Choudhury, B.C., and Singh, S. 2021. *Pangshura smithii*. The IUCN Red List of Threatened Species 2021: e.T39554A2929235.

Ahmed, M.F., Praschag, P., and Singh, S. 2021. *Pangshura tecta*. The IUCN Red List of Threatened Species 2021: e.T46370A3005714.

Alcalde, L., Fagundes, C.K., and Carreira, S. In press. *Hydromedusa tectifera*. The IUCN Red List of Threatened Species.

Alcalde, L., Carreira, S., and Fagundes, C.K. In press. *Phrynops hilarii*. The IUCN Red List of Threatened Species.

Amepou, Y., Georges, A., Thomson, S., Eisemberg, C., and Rhodin, A.G.J. 2020. *Elseya rhodini*. The IUCN Red List of Threatened Species 2020: e.T90377834A90377837.

Andrade, P.C.M., Rojas-Runjaic, F.J.M., Ferrara, C.R., Bernhard, R., and Vogt, R.C. In press. *Podocnemis erythrocephala*. The IUCN Red List of Threatened Species.

ARASG [Australasian Reptile and Amphibian Specialist Group]. 1996. *Rheodytes leukops*. The IUCN Red List of Threatened Species 1996: e.T19483A8904917.

Asmat, G.S.M. 2002. *Nilssonia nigricans*. The IUCN Red List of Threatened Species 2002: e.T2173A9314968.

As-singkily, M., Eisemberg, C., Horne, B.D., Kuchling, G., and Rhodin, A.G.J. 2019. *Chelodina mccordi*. The IUCN Red List of Threatened Species 2019: e.T123814489A123814575.

As-singkily, M., Guntoro, J., Kusrini, M.D., and Schoppe, S. 2021. *Cyclemys dentata*. The IUCN Red List of Threatened Species 2021: e.T195849722A2929066.

As-singkily, M., Guntoro, J., Kusrini, M.D., and Hamidy, A. 2021. *Cyclemys enigmatica*. The IUCN Red List of Threatened Species 2021: e.T170505A1315402.

ATTWG [Asian Turtle Trade Working Group]. 2000. *Amyda cartilaginea*. The IUCN Red List of Threatened Species 2000: e.T1181A3309466.

ATTWG [Asian Turtle Trade Working Group]. 2000. *Batagur baska*. The IUCN Red List of Threatened Species 2000: e.T2614A9461838.

ATTWG [Asian Turtle Trade Working Group]. 2000. *Batagur borneoensis*. The IUCN Red List of Threatened Species 2000: e.T163458A5608163.

ATTWG [Asian Turtle Trade Working Group]. 2000. *Batagur dhongoka*. The IUCN Red List of Threatened Species 2000: e.T10953A3230309.

ATTWG [Asian Turtle Trade Working Group]. 2000. *Batagur kachuga*. The IUCN Red List of Threatened Species 2000: e.T10949A3229147.

ATTWG [Asian Turtle Trade Working Group]. 2000. *Batagur trivittata*. The IUCN Red List of Threatened Species 2000: e.T10952A3230190.

ATTWG [Asian Turtle Trade Working Group]. 2000. *Carettochelys insculpta*. The IUCN Red List of Threatened Species 2000: e.T3898A10155584.

ATTWG [Asian Turtle Trade Working Group]. 2000. *Chelodina mccordi*. The

IUCN Red List of Threatened Species 2000: e.T4606A11027770.
ATTWG [Asian Turtle Trade Working Group]. 2000. *Chelodina novaeguineae*. The IUCN Red List of Threatened Species 2000: e.T46580A11061655.
ATTWG [Asian Turtle Trade Working Group]. 2000. *Chelodina oblonga*. The IUCN Red List of Threatened Species 1996: e.T4607A11032585.
ATTWG [Asian Turtle Trade Working Group]. 2000. *Chelodina parkeri*. The IUCN Red List of Threatened Species 2000: e.T4608A11028994.
ATTWG [Asian Turtle Trade Working Group]. 2000. *Chelodina pritchardi*. The IUCN Red List of Threatened Species 2000: e.T4609A11030109.
ATTWG [Asian Turtle Trade Working Group]. 2000. *Chelodina reimanni*. The IUCN Red List of Threatened Species 2000: e.T4610A11032759.
ATTWG [Asian Turtle Trade Working Group]. 2000. *Chelodina rugosa* ssp. *siebenrocki*. The IUCN Red List of Threatened Species 2000: e.T39622A10252168.
ATTWG [Asian Turtle Trade Working Group]. 2000. *Chitra chitra*. The IUCN Red List of Threatened Species 2000: e.T4695A11088507.
ATTWG [Asian Turtle Trade Working Group]. 2000. *Chitra indica*. The IUCN Red List of Threatened Species 2000: e.T4696A11088615.
ATTWG [Asian Turtle Trade Working Group]. 2000. *Cuora amboinensis*. The IUCN Red List of Threatened Species 2000: e.T5958A11953035.
ATTWG [Asian Turtle Trade Working Group]. 2000. *Cuora aurocapitata*. The IUCN Red List of Threatened Species 2000: e.T5959A11956075.
ATTWG [Asian Turtle Trade Working Group]. 2000. *Cuora flavomarginata*. The IUCN Red List of Threatened Species 2000: e.T5960A11965283.
ATTWG [Asian Turtle Trade Working Group]. 2000. *Cuora galbinifrons*. The IUCN Red List of Threatened Species 2000: e.T5955A11962273.
ATTWG [Asian Turtle Trade Working Group]. 2000. *Cuora mccordi*. The IUCN Red List of Threatened Species 2000: e.T5961A11967952.
ATTWG [Asian Turtle Trade Working Group]. 2000. *Cuora mouhotii*. The IUCN Red List of Threatened Species 2000: e.T163414A5604435.
ATTWG [Asian Turtle Trade Working Group]. 2000. *Cuora pani*. The IUCN Red List of Threatened Species 2000: e.T5956A11949874.
ATTWG [Asian Turtle Trade Working Group]. 2000. *Cuora trifasciata*. The IUCN Red List of Threatened Species 2000: e.T5962A11969807.
ATTWG [Asian Turtle Trade Working Group]. 2000. *Cuora yunnanensis*. The IUCN Red List of Threatened Species 2000: e.T5957A11964344.
ATTWG [Asian Turtle Trade Working Group]. 2000. *Cuora zhoui*. The IUCN Red List of Threatened Species 2000: e.T5963A11972677.
ATTWG [Asian Turtle Trade Working Group]. 2000. *Cyclemys dentata*. The IUCN Red List of Threatened Species 2000: e.T39551A10247518.
ATTWG [Asian Turtle Trade Working Group]. 2000. *Dogania subplana*. The IUCN Red List of Threatened Species 2000: e.T46578A11061078.
ATTWG [Asian Turtle Trade Working Group]. 2000. *Elseya branderhorsti*. The IUCN Red List of Threatened Species 2000: e.T39623A10252261.
ATTWG [Asian Turtle Trade Working Group]. 2000. *Elseya novaeguineae*. The IUCN Red List of Threatened Species 2000: e.T46581A11061765.
ATTWG [Asian Turtle Trade Working Group]. 2000. *Emydura subglobosa*. The IUCN Red List of Threatened Species 2000: e.T46582A11061866.
ATTWG [Asian Turtle Trade Working Group]. 2000. *Geochelone elegans*. The IUCN Red List of Threatened Species 2000: e.T39430A10237876.
ATTWG [Asian Turtle Trade Working Group]. 2000. *Geochelone platynota*. The IUCN Red List of Threatened Species 2000: e.T9013A12950329.
ATTWG [Asian Turtle Trade Working Group]. 2000. *Geoclemys hamiltonii*. The IUCN Red List of Threatened Species 2000: e.T9029A12951978.
ATTWG [Asian Turtle Trade Working Group]. 2000. *Geoemyda japonica*. The IUCN Red List of Threatened Species 2000: e.T9042A12952738.
ATTWG [Asian Turtle Trade Working Group]. 2000. *Geoemyda spengleri*. The IUCN Red List of Threatened Species 2000: e.T39552A10247618.
ATTWG [Asian Turtle Trade Working Group]. 2000. *Hardella thurjii*. The IUCN Red List of Threatened Species 2000: e.T9696A13009662.
ATTWG [Asian Turtle Trade Working Group]. 2000. *Heosemys annandalii*. The IUCN Red List of Threatened Species 2000: e.T10041A3156321.
ATTWG [Asian Turtle Trade Working Group]. 2000. *Heosemys depressa*. The IUCN Red List of Threatened Species 2000: e.T39596A10244762.
ATTWG [Asian Turtle Trade Working Group]. 2000. *Heosemys grandis*. The IUCN Red List of Threatened Species 2000: e.T9943A13027989.
ATTWG [Asian Turtle Trade Working Group]. 2000. *Heosemys spinosa*. The IUCN Red List of Threatened Species 2000: e.T9942A13027850.
ATTWG [Asian Turtle Trade Working Group]. 2000. *Indotestudo elongata*. The IUCN Red List of Threatened Species 2000: e.T10824A3219349.
ATTWG [Asian Turtle Trade Working Group]. 2000. *Indotestudo forstenii*. The IUCN Red List of Threatened Species 2000: e.T10825A3219524.
ATTWG [Asian Turtle Trade Working Group]. 2000. *Indotestudo travancorica*. The IUCN Red List of Threatened Species 2000: e.T39548A10247310.
ATTWG [Asian Turtle Trade Working Group]. 2000. *Leucocephalon yuwonoi*. The IUCN Red List of Threatened Species 2000: e.T40761A10363014.
ATTWG [Asian Turtle Trade Working Group]. 2000. *Lissemys punctata*. The IUCN Red List of Threatened Species 2000: e.T46579A11061187.
ATTWG [Asian Turtle Trade Working Group]. 2000. *Lissemys scutata*. The IUCN Red List of Threatened Species 2000: e.T12124A3323090.
ATTWG [Asian Turtle Trade Working Group]. 2000. *Malayemys subtrijuga*. The IUCN Red List of Threatened Species 2000: e.T39555A10248240.
ATTWG [Asian Turtle Trade Working Group]. 2000. *Manouria emys*. The IUCN Red List of Threatened Species 2000: e.T12774A3380771.
ATTWG [Asian Turtle Trade Working Group]. 2000. *Manouria impressa*. The IUCN Red List of Threatened Species 2000: e.T12775A3380977.
ATTWG [Asian Turtle Trade Working Group]. 2000. *Mauremys annamensis*. The IUCN Red List of Threatened Species 2000: e.T12876A3393755.
ATTWG [Asian Turtle Trade Working Group]. 2000. *Mauremys japonica*. The IUCN Red List of Threatened Species 2000: e.T39612A10251032.
ATTWG [Asian Turtle Trade Working Group]. 2000. *Mauremys mutica*. The IUCN Red List of Threatened Species 2000: e.T39613A10251132.
ATTWG [Asian Turtle Trade Working Group]. 2000. *Mauremys nigricans*. The IUCN Red List of Threatened Species 2000: e.T4656A11059086.
ATTWG [Asian Turtle Trade Working Group]. 2000. *Mauremys sinensis*. The IUCN Red List of Threatened Species 2000: e.T15026A4488909.
ATTWG [Asian Turtle Trade Working Group]. 2000. *Melanochelys tricarinata*. The IUCN Red List of Threatened Species 2000: e.T13038A3406791.
ATTWG [Asian Turtle Trade Working Group]. 2000. *Melanochelys trijuga*. The IUCN Red List of Threatened Species 2000: e.T13039A3406900.
ATTWG [Asian Turtle Trade Working Group]. 2000. *Morenia ocellata*. The IUCN Red List of Threatened Species 2000: e.T13873A4359698.
ATTWG [Asian Turtle Trade Working Group]. 2000. *Morenia petersi*. The IUCN Red List of Threatened Species 2000: e.T13874A4359795.
ATTWG [Asian Turtle Trade Working Group]. 2000. *Nilssonia formosa*. The IUCN Red List of Threatened Species 2000: e.T14765A4460404.
ATTWG [Asian Turtle Trade Working Group]. 2000. *Nilssonia gangetica*. The IUCN Red List of Threatened Species 2000: e.T39618A10251627.
ATTWG [Asian Turtle Trade Working Group]. 2000. *Nilssonia hurum*. The IUCN Red List of Threatened Species 2000: e.T39619A10251753.
ATTWG [Asian Turtle Trade Working Group]. 2000. *Nilssonia leithii*. The IUCN Red List of Threatened Species 2000: e.T2174A9315657.
ATTWG [Asian Turtle Trade Working Group]. 2000. *Notochelys platynota*. The IUCN Red List of Threatened Species 2000: e.T14856A4465399.
ATTWG [Asian Turtle Trade Working Group]. 2000. *Orlitia borneensis*. The IUCN Red List of Threatened Species 2000: e.T15509A4698687.
ATTWG [Asian Turtle Trade Working Group]. 2000. *Palea steindachneri*. The IUCN Red List of Threatened Species 2000: e.T15918A5320047.
ATTWG [Asian Turtle Trade Working Group]. 2000. *Pangshura smithii*. The IUCN Red List of Threatened Species 2000: e.T39554A10248159.
ATTWG [Asian Turtle Trade Working Group]. 2000. *Pangshura sylhetensis*. The IUCN Red List of Threatened Species 2000: e.T10950A3229978.
ATTWG [Asian Turtle Trade Working Group]. 2000. *Pangshura tecta*. The IUCN Red List of Threatened Species 2000: e.T46370A11046458.
ATTWG [Asian Turtle Trade Working Group]. 2000. *Pangshura tentoria*. The IUCN Red List of Threatened Species 2000: e.T46577A11060648.
ATTWG [Asian Turtle Trade Working Group]. 2000. *Pelochelys bibroni*. The IUCN Red List of Threatened Species 2000: e.T16503A97400832.
ATTWG [Asian Turtle Trade Working Group]. 2000. *Pelochelys cantorii*. The IUCN Red List of Threatened Species 2000: e.T16502A5964413.
ATTWG [Asian Turtle Trade Working Group]. 2000. *Pelodiscus sinensis*. The IUCN Red List of Threatened Species 2000: e.T39620A10251914.
ATTWG [Asian Turtle Trade Working Group]. 2000. *Platysternon megacephalum*. The IUCN Red List of Threatened Species 2000: e.T17585A7142204.
ATTWG [Asian Turtle Trade Working Group]. 2000. *Rafetus swinhoei*. The IUCN Red List of Threatened Species 2000: e.T39621A10252043.

ATTWG [Asian Turtle Trade Working Group]. 2000. *Sacalia bealei*. The IUCN Red List of Threatened Species 2000: e.T19796A9016239.

ATTWG [Asian Turtle Trade Working Group]. 2000. *Sacalia quadriocellata*. The IUCN Red List of Threatened Species 2000: e.T19797A9016329.

ATTWG [Asian Turtle Trade Working Group]. 2000. *Siebenrockiella crassicollis*. The IUCN Red List of Threatened Species 2000: e.T39616A10251374.

ATTWG [Asian Turtle Trade Working Group]. 2000. *Siebenrockiella leytensis*. The IUCN Red List of Threatened Species 2000: e.T39599A10245269.

ATTWG [Asian Turtle Trade Working Group]. 2000. *Vijayachelys silvatica*. The IUCN Red List of Threatened Species 2000: e.T9038A12952577.

Averill-Murray, R.C., Rosen, P.C., Jones, C.A., Jones, T.R., Lara-Resendiz, R.A., Edwards, T., Karl, A., and Berry, K.H. 2023. *Gopherus morafkai*. The IUCN Red List of Threatened Species 2023: e.T97246109A97246177.

Baillie, J. and Groombridge, B. 1996. The 1996 IUCN Red List of Threatened Animals. Gland, Switzerland: IUCN, 368 pp.

Baker, P.J., Kabigumila, J., Leuteritz, T., Hofmeyr, M., and Ngwava, J.M. 2015. *Stigmochelys pardalis*. The IUCN Red List of Threatened Species 2015: e.T163449A1009442.

Baker, P.J., Luiselli, L., and Diagne, T. 2016. *Cyclanorbis elegans*. The IUCN Red List of Threatened Species 2016: e.T6004A3086539.

Barreto, L.N., Sousa Ribeiro, L.E.d., and Bock, B.C. In press. *Trachemys adiutrix*. The IUCN Red List of Threatened Species.

Behler, J.L. 1996. IUCN Red List Assessments for turtles and tortoises, including Least Concern species. Report from the IUCN SSC Tortoise and Freshwater Turtle Specialist Group to the IUCN Red List Programme.

Berry, K.H., Allison, L.J., McLuckie, A.M., Vaughn, M., and Murphy, R.W. 2021. *Gopherus agassizii*. The IUCN Red List of Threatened Species 2021: e.T97246272A3150871.

Bock, B.C., Páez, V.P., Carvalho, V.T. de, and Rhodin, A.G.J. In press. *Mesoclemmys perplexa*. The IUCN Red List of Threatened Species.

Bock, B.C., Barros, T.R., Guada, H.J., Rivas, G., Hernández, O., Páez, V.P., Rojas-Runjaic, F.J.M., Camargo Siliet, E., and Cortés-Duque, J. In press. *Trachemys callirostris*. The IUCN Red List of Threatened Species.

Bock, B.C., Cupul-Magaña, F.G., Páez, V.P., Reyes-Grajales, E., and van Dijk, P.P. In press. *Trachemys venusta*. The IUCN Red List of Threatened Species.

Branch, W.R. 1996. *Homopus signatus*. The IUCN Red List of Threatened Species 1996: e.T10241A3184678.

Branch, W.R. 1996. *Homopus solus*. The IUCN Red List of Threatened Species 1996: e.T10238A3184592.

Branch, W.R. 2018. *Chersobius solus* (amended version of 1996 assessment). The IUCN Red List of Threatened Species 2018: e.T10238A125807053.

Brito, E.S., Valadão, R.M., Cunha, F.A.G., Oliveira, J.C.F., Hernández, O., Ferrara, C.R., Andrade, P.C.M., and Carr, J.L. 2025. *Rhinoclemmys punctularia*. The IUCN Red List of Threatened Species 2025: e.T170513A275854981.

Brito, E.S., Valadão, R.M., Cunha, F., Vinke, S., Vinke, T., Cabrera, M.R., Marques, T.S., Carvajal-Bacarreza, P., and Strüssmann, C. In press. *Mesoclemmys vanderhaegei*. The IUCN Red List of Threatened Species.

Broderick, A.C., Türkozan, O., Demetropoulos, S., Mastrogiacomo, A., Demetropoulos, A., Levy, Y., and Leader, N. 2023. *Chelonia mydas* (Mediterranean subpopulation). The IUCN Red List of Threatened Species 2023: e.T4616A83319449.

Broderick, A.C., Türkozan, O., Demetropoulos, S., Mastrogiacomo, A., Demetropoulos, A., Levy, Y., and Leader, N. 2024. *Chelonia mydas* (Mediterranean subpopulation) (amended version of 2023 assessment). The IUCN Red List of Threatened Species 2024: e.T4616A259027885.

Brown, G.J. and Welc, M. 2025. *Sternotherus peltifer*. The IUCN Red List of Threatened Species 2025: e.T218364295A218364006.

Brown, G.J., Welc, M., and van Dijk, P.P. 2025. *Sternotherus minor*. The IUCN Red List of Threatened Species 2025: e.T218363617A218363714.

Brown, G.J., Iverson, J.B., and Welc, M. 2025. *Sternotherus odoratus*. The IUCN Red List of Threatened Species 2025: e.T163450A161721733.

Butterfield, T.G. and Matías-Palafox, M.L. 2025. *Kinosternon chimalhuaca*. The IUCN Red List of Threatened Species 2025: e.T63667A161720838.

Butterfield, T.G., Monsiváis-Molina, A., Díaz-Gamboa, L.F., Willey, L.L., Jones, M.T., and Platt, S.G. In press. *Rhinoclemmys areolata*. The IUCN Red List of Threatened Species.

Butterfield, T.G., Iverson, J.B., and Cupul-Magaña, F.G. In press. *Rhinoclemmys pulcherrima*. The IUCN Red List of Threatened Species.

Butterfield, T.G., Espinosa, S., Matias-Palafox, S., Domínguéz-Pompa, A., Monsiváis-Molina, A., García-Caballero, F., and Uc-Uc, M. In press. *Terrapene nelsoni*. The IUCN Red List of Threatened Species.

Caccone, A., Cayot, L.J., Gibbs, J.P., and Tapia, W. 2017. *Chelonoidis becki*. The IUCN Red List of Threatened Species 2017: e.T9018A82426296.

Caccone, A., Cayot, L.J., Gibbs, J.P., and Tapia, W. 2017. *Chelonoidis chathamensis*. The IUCN Red List of Threatened Species 2017: e.T9019A82688009.

Carr, J.L. and Giraldo, A. In press. *Rhinoclemmys annulata*. The IUCN Red List of Threatened Species.

Carr, J.L., Riedle, D.J., Munscher, E., Pearson, L.S., Kessler, E.J., and Dreslik, M.J. 2025. *Macrochelys temminckii*. The IUCN Red List of Threatened Species 2025: e.T232775771A507158.

Carr, J.L., Garcés-Restrepo, M.F., and Giraldo, A. In press. *Rhinoclemmys nasuta*. The IUCN Red List of Threatened Species.

Casale, P. and Tucker, A.D. 2017. *Caretta caretta* (amended version of 2015 assessment). The IUCN Red List of Threatened Species 2017: e.T3897A119333622.

Castañeda Gaytán, G., Stanford, C.B., Valenzuela-Ceballos, S., and Rhodin, A.G.J. In press. *Terrapene coahuila*. The IUCN Red List of Threatened Species.

Castañeda Gaytán, G., García-Velez, C., Fernández-Badillo, L., Valencia-Herverth, R., and Stanford, C.B. In press. *Terrapene mexicana*. The IUCN Red List of Threatened Species.

Cayot, L.J., Gibbs, J.P., Tapia, W., and Caccone, A. 2016. *Chelonoidis abingdonii*. The IUCN Red List of Threatened Species 2016: e.T9017A65487433.

Cayot, L.J., Gibbs, J.P., Tapia, W., and Caccone, A. 2016. *Chelonoidis darwini*. The IUCN Red List of Threatened Species 2016: e.T9020A82689845.

Cayot, L.J., Gibbs, J.P., Tapia, W., and Caccone, A. 2017. *Chelonoidis donfaustoi*. The IUCN Red List of Threatened Species 2017: e.T90377132A90377135.

Cayot, L.J., Gibbs, J.P., Tapia, W., and Caccone, A. 2017. *Chelonoidis duncanensis*. The IUCN Red List of Threatened Species 2017: e.T9021A3149054.

Cayot, L.J., Gibbs, J.P., Tapia, W., and Caccone, A. 2017. *Chelonoidis hoodensis*. The IUCN Red List of Threatened Species 2017: e.T9024A82777079.

Cayot, L.J., Gibbs, J.P., Tapia, W., and Caccone, A. 2017. *Chelonoidis porteri*. The IUCN Red List of Threatened Species 2017: e.T9026A82777132.

Cayot, L.J., Gibbs, J.P., Tapia, W., and Caccone, A. 2018. *Chelonoidis guntheri*. The IUCN Red List of Threatened Species 2018: e.T9022A116505802.

Cayot, L.J., Gibbs, J.P., Tapia, W., and Caccone, A. 2018. *Chelonoidis microphyes*. The IUCN Red List of Threatened Species 2018: e.T9025A116506297.

Cayot, L.J., Gibbs, J.P., Tapia, W., and Caccone, A. 2018. *Chelonoidis vandenburghi*. The IUCN Red List of Threatened Species 2018: e.T9027A116506698.

Cayot, L.J., Gibbs, J.P., Tapia, W., and Caccone, A. 2018. *Chelonoidis vicina*. The IUCN Red List of Threatened Species 2018: e.T9028A82532040.

Ceballos, C.P., Vinke, S., Vinke, T., Rojano Bolaño, C., Morcatty, T.Q., Escalona, T., and Gallego-García, N. In press. *Chelonoidis carbonarius*. The IUCN Red List of Threatened Species.

Chaloupka, M.Y. and Pilcher, N.J. 2019. *Chelonia mydas* (Hawaiian subpopulation). The IUCN Red List of Threatened Species 2019: e.T16285718A142098300.

Chirio, L., Diagne, T., and Pauwels, O.S.G. 2017. *Cycloderma aubryi*. The IUCN Red List of Threatened Species 2017: e.T163448A1009393.

Choudhury, B.C. and Praschag, P. 2021. *Vijayachelys silvatica*. The IUCN Red List of Threatened Species 2021: e.T9038A3149346.

Choudhury, B.C., Cota, M., McCormack, T., Platt, K., Das, I., Ahmed, M.F., Timmins, R.J., Rahman, S., and Singh, S. 2019. *Manouria emys*. The IUCN Red List of Threatened Species 2019: e.T12774A152052098.

Choudhury, B.C., de Silva, A., and Shepherd, C. 2020. *Geochelone elegans*. The IUCN Red List of Threatened Species 2020: e.T39430A123815345.

Choudhury, B.C., Das, I., Horne, B.D., Li, P., McCormack, T., Praschag, P., Rao, D.-Q., and Wang, L. 2021. *Pelochelys cantorii*. The IUCN Red List of Threatened Species 2021: e.T135458600A1076984.

Choudhury, B.C., Das, I., Praschag, P., and Singh, S. 2021. *Pangshura tentoria*. The IUCN Red List of Threatened Species 2021: e.T46577A3008697.

Cota, M. 2021. *Malayemys khoratensis*. The IUCN Red List of Threatened Species 2021: e.T123770233A123770237.

Cota, M. 2021. *Malayemys macrocephala*. The IUCN Red List of Threatened Species 2021: e.T170509A1315502.

Cota, M., Guntoro, J., As-singkily, M., Hamidy, A., and Kusrini, M.D. 2019. *Chitra chitra*. The IUCN Red List of Threatened Species 2019: e.T4695A152049778.

Cota, M., Hoang, H., Horne, B.D., Kusrini, M.D., McCormack, T., Platt, K., Schoppe, S., and Shepherd, C. 2020. *Cuora amboinensis*. The IUCN Red List of Threatened Species 2020: e.T5958A3078812.

Cota, M., Guntoro, J., Horne, B.D., Kusrini, M.D., Krishnasamy, K., and Shepherd, C. 2021. *Heosemys spinosa*. The IUCN Red List of Threatened Species 2021: e.T9942A3152508.

Cota, M., Hamidy, A., Platt, K., Kusrini, M.D., Guntoro, J., Shepherd, C., and Schoppe, S. 2021. *Dogania subplana*. The IUCN Red List of Threatened Species 2021: e.T46578A3008869.

Cota, M., Horne, B.D., McCormack, T., and Timmins, R.J. 2021. *Heosemys annandalii*. The IUCN Red List of Threatened Species 2021: e.T10041A495907.

Cota, M., Horne, B.D., and Shepherd, C. 2021. *Heosemys grandis*. The IUCN Red List of Threatened Species 2021: e.T9943A3152603.

Cota, M., Li, P., Platt, K., Rao, D.-Q., Stanford, C.B., Wang, L., and Wanchai, P. 2021. *Manouria impressa*. The IUCN Red List of Threatened Species 2021: e.T12775A508518.

Cox, N.A. and Temple, H.J. 2009. European Red List of Reptiles. Luxembourg: Office for Official Publications of the European Communities.

Cunha, F.A.G., Brito, E.S., and Valadão, R.M. In press. *Mesoclemmys nasuta*. The IUCN Red List of Threatened Species.

Cunha, F.A.G., Vasconcelos, V., Viana, P., Brito, E.S., and Valadão, R.M. In press. *Mesoclemmys raniceps*. The IUCN Red List of Threatened Species.

Cunha, F.A.G., Brito, E.S., Valadão, R.M., Oliveira, M., and Vasconcelos, V. In press. *Mesoclemmys wermuthi*. The IUCN Red List of Threatened Species.

Cunha, F.A.G., Brito, E.S., Valadão, R.M., and Rhodin, A.G.J. In press. *Platemys platycephala*. The IUCN Red List of Threatened Species.

Cupul-Magaña, F.G., Butterfield, T., Gregory, T., Iverson, J.B., Macip-Ríos, R., and López-Luna, M.A. 2022. *Kinosternon vogti*. The IUCN Red List of Threatened Species 2022:e.T215164369A215164374.

Cupul-Magaña, F.G., Flores-Guerrero, U.S., Nolasco-Luna, J.R., and Escobedo-Galván, A.H. In press. *Trachemys nebulosa*. The IUCN Red List of Threatened Species.

Cupul-Magaña, F.G., Flores-Guerrero, U.S., Nolasco-Luna, J.R., and Escobedo-Galván, A.H. In press. *Trachemys ornata*. The IUCN Red List of Threatened Species.

D'Cruze, N., Choudhury, B.C., and Mookerjee, A. 2016. *Geochelone elegans*. The IUCN Red List of Threatened Species 2016: e.T39430A2926441.

Das, I., Choudhury, B.C., Praschag, P., Ahmed, M.F., and Singh, S. 2019. *Batagur dhongoka*. The IUCN Red List of Threatened Species 2019: e.T10953A152042542.

Das, I., Choudhury, B.C., Ahmed, M.F., Praschag, P., and Singh, S. 2021. *Nilssonia hurum*. The IUCN Red List of Threatened Species 2021: e.T39619A2931203.

Diagne, T., Luiselli, L., Trape, J.-F., Rödel, M.-O., Baker, P.J., Chirio, L., Petrozzi, F., and Segniagbeto, G. 2016. *Cyclanorbis senegalensis*. The IUCN Red List of Threatened Species 2016: e.T6005A96447114.

Díaz-Gamboa, L.F., Butterfield, T., Jones, M.T., and Willey, L.L. 2025. *Kinosternon creaseri*. The IUCN Red List of Threatened Species 2025: e.T11006A161718337.

Doody, J.S., Gillingwater, S.D., Mali, I., Forstner, M.R.J., van Dijk, P.P., and Rhodin, A.G.J. 2024. *Apalone spinifera*. The IUCN Red List of Threatened Species 2024: e.T163451A161722123.

Drummond, G.M., Coutinho, M.E., and Vogt, R.C. 2016. *Mesoclemmys hogei*. The IUCN Red List of Threatened Species 2016: e.T17081A1316719.

Edwards, T., Karl, A.E., Rosen, P.C., Vaughn, M., Melendez Torres, C., Kiester, A.R., and Goode, E.V. 2018. *Gopherus evgoodei*. The IUCN Red List of Threatened Species 2018: e.T90377823A90377826.

Eisemberg, C., van Dijk, P.P., Georges, A., and Amepou, Y. 2018. *Carettochelys insculpta*. The IUCN Red List of Threatened Species 2018: e.T3898A2884984.

ERASG [European Reptile and Amphibian Specialist Group]. 1996. *Rafetus euphraticus*. The IUCN Red List of Threatened Species 1996: e.T19070A8809367.

ERASG [European Reptile and Amphibian Specialist Group]. 1996. *Testudo graeca* ssp. *nikolskii*. The IUCN Red List of Threatened Species 1996: e.T21654A9307594.

ERASG [European Reptile and Amphibian Specialist Group]. 1996. *Testudo hermanni* ssp. *hermanni*. The IUCN Red List of Threatened Species 1996: e.T21650A9306703.

ERASG [European Reptile and Amphibian Specialist Group]. 1996. *Trionyx triunguis* (Mediterranean subpopulation). The IUCN Red List of Threatened Species 1996: e.T22200A9364253.

Escalona, T., Norris, D., de Thoisy, B., Carvajal-Bacarreza, P., Cueva, R., Conway-Gómez, K., Andrade, P., Ouboter, P.E., and Acebey-Quiroga, S. In press. *Podocnemis unifilis*. The IUCN Red List of Threatened Species.

Ettmar, S. In press. *Mesoclemmys gibba*. The IUCN Red List of Threatened Species.

Fagundes, C.K., Alcalde, L., Carreira, S., Fallabrino, A., Estrades, A., Florencia David, M., Famelli, S., and Lustosa, A.P. In press. *Trachemys dorbigni*. The IUCN Red List of Threatened Species.

Famelli, S., Souza, F.L., Valadão, R.M., and Brito, E.S. In press. *Hydromedusa maximiliani*. The IUCN Red List of Threatened Species.

Ferrara, C.R., Fagundes, C.K., Forero-Medina, G., Brito, E.S., Miorando, P.S., Lacava, R., Gomes Lustosa, A.P., Hernández, O., Cueva, R., Millar, N., Dewynter, M., Páez, V.P., Carvajal, P., and Domínguez, A.L. In press. *Podocnemis expansa*. The IUCN Red List of Threatened Species.

Ferrara, C.R., Brito, E.S., Forero-Medina, G., Cunha, F., and Vogt, R.C. In press. *Rhinemys rufipes*. The IUCN Red List of Threatened Species.

Folt, B., van Dijk, P.P., Rhodin, A.G.J., and Such, R. In press. *Rhinoclemmys funerea*. The IUCN Red List of Threatened Species.

Fong, J., Hoang, H., Horne, B.D., Li, P., McCormack, T., Rao, D.-Q., Timmins, R.J., and Wang, L. 2020. *Cuora trifasciata*. The IUCN Red List of Threatened Species 2020: e.T123768950A123769768.

Fong, J., Hoang, H., Li, P., McCormack, T., Rao, D.-Q., Timmins, R.J., and Wang, L. 2020. *Geoemyda spengleri*. The IUCN Red List of Threatened Species 2020: e.T39552A2929166.

Fong, J., Hoang, H., Kuchling, G., Li, P., McCormack, T., Rao, D.-Q., Timmins, R.J., and Wang, L. 2021. *Rafetus swinhoei*. The IUCN Red List of Threatened Species 2021: e.T39621A2931537.

Fong, J., Hoang, H., Li, P., McCormack, T., Rao, D.-Q., Timmins, R.J., and Wang, L. 2021. *Palea steindachneri*. The IUCN Red List of Threatened Species 2021: e.T15918A794203.

Fong, J., Hoang, H., Li, P., McCormack, T., Rao, D.-Q., Shi, H., and Wang, L. 2021. *Sacalia quadriocellata*. The IUCN Red List of Threatened Species 2021: e.T19797A2507554.

Fong, J., Hoang, H., Li, P., McCormack, T., Rao, D.-Q., Timmins, R.J., Wang, L., and van Dijk, P.P. 2021. *Platysternon megacephalum*. The IUCN Red List of Threatened Species 2021: e.T17585A1423706.

Fong, J., Hoang, H., Li, P., McCormack, T., Rao, D.-Q., and Wang, L. 2021. *Mauremys mutica*. The IUCN Red List of Threatened Species 2021: e.T39613A2930788.

Forero Medina, G., Rentería-Moreno, E., Iverson, J.B., and Echavarría, M. In press. *Kinosternon dunni*. The IUCN Red List of Threatened Species.

Forero Medina, G., Iverson, J.B., Hurtado-Gómez, J.P., López-Luna, M.A., van Dijk, P.P., and Rhodin, A.G.J. In press. *Kinosternon scorpioides*. The IUCN Red List of Threatened Species.

Forstner, M.R.J. and Mali, I. In press. *Graptemys caglei*. The IUCN Red List of Threatened Species.

Frost, D., Hammerson, G., and Gadsden, H. 2007. *Kinosternon alamosae*. The IUCN Red List of Threatened Species 2007: e.T63665A12694308.

Frost, D., Hammerson, G., and Gadsden, H. 2007. *Kinosternon arizonense*. The IUCN Red List of Threatened Species 2007: e.T63666A12694945.

Frost, D., Hammerson, G., and Gadsden, H. 2007. *Trachemys ornata*. The IUCN Red List of Threatened Species 2007: e.T63661A12704799.

Frost, D., Hammerson, G., and Gadsden, H. 2007. *Trachemys yaquia*. The IUCN Red List of Threatened Species 2007: e.T63663A12693995.

Gallego-García, N. and Forero-Medina, G. In press. *Mesoclemmys dahli*. The IUCN Red List of Threatened Species.

Gentil, E., de Medeiros, L.A., Morales-Betancourt, M.A., Maffei, F., and Da Silveira, R. In press. *Peltocephalus dumerilianus*. The IUCN Red List of Threatened Species.

Georges, A., Rhodin, A.G.J., Eisemberg, C., and Amepou, Y. 2020. *Pelochelys bibroni*. The IUCN Red List of Threatened Species 2020: e.T16503A1077288.

Gerlach, J. 2003. *Pelusios castanoides* ssp. *intergularis*. The IUCN Red List of

Threatened Species 2003: e.T41603A10504198.

Gerlach, J. 2003. *Pelusios seychellensis*. The IUCN Red List of Threatened Species 2003: e.T16527A5998317.

Gerlach, J. 2003. *Pelusios subniger* ssp. *parietalis*. The IUCN Red List of Threatened Species 2003: e.T41604A10504742.

Ghaffari, H., Taskavak, E., Turkozan, O., and Mobaraki, A. 2017. *Rafetus euphraticus*. The IUCN Red List of Threatened Species 2017: e.T19070A1956551.

Godwin, C.D. 2025. *Apalone mutica*. The IUCN Red List of Threatened Species 2025: e.T165596A250316650.

Godwin, J.C. and Brown, G.J. 2025. *Graptemys pulchra*. The IUCN Red List of Threatened Species 2025: e.T170494A251714872.

Godwin, J.C. and Pearson, L.S. In press. *Pseudemys alabamensis*. The IUCN Red List of Threatened Species.

Godwin, J.C. and Selman, W. In press. *Graptemys nigrinoda*. The IUCN Red List of Threatened Species.

Godwin, J.C., Mays, J.D., and Hill, P. 2025. *Graptemys ernsti*. The IUCN Red List of Threatened Species 2025: e.T9500A251709703.

Hofmeyr, M.D. and Baard, E.H.W. 2015. *Psammobates geometricus*. The IUCN Red List of Threatened Species 2015: e.T18398A76918966.

Hofmeyr, M.D. and Baard, E.H.W. 2018. *Psammobates geometricus*. The IUCN Red List of Threatened Species 2018: e.T18398A111553007.

Hofmeyr, M.D. and Boycott, R.C. 2018. *Kinixys lobatsiana*. The IUCN Red List of Threatened Species 2018: e.T163454A115654759.

Hofmeyr, M.D., and Boycott, R.C. 2018. *Kinixys natalensis*. The IUCN Red List of Threatened Species 2018: e.T11004A115685642.

Hofmeyr, M.D. and Fritz, U. 2018. *Pelomedusa galeata*. The IUCN Red List of Threatened Species 2018: e.T113551736A113551752.

Hofmeyr, M.D. and Keswick, T. 2018. *Chersina angulata*. The IUCN Red List of Threatened Species 2018: e.T170519A115655918.

Hofmeyr, M.D. and Keswick, T. 2018. *Homopus areolatus*. The IUCN Red List of Threatened Species 2018: e.T170520A115656133.

Hofmeyr, M.D. and Keswick, T. 2018. *Homopus femoralis*. The IUCN Red List of Threatened Species 2018: e.T170522A115656613.

Hofmeyr, M.D., Loehr, V.J.T., Baard, E.H.W., and Juvik, J.O. 2018. *Chersobius boulengeri*. The IUCN Red List of Threatened Species 2018: e.T170521A115656360.

Hofmeyr, M.D., Loehr, V.J.T., and Baard, E.H.W. 2018. *Chersobius signatus*. The IUCN Red List of Threatened Species 2018: e.T10241A115650943.

Hofmeyr, M.D., Leuteritz, T., and Baard, E.H.W. 2018. *Psammobates tentorius*. The IUCN Red List of Threatened Species 2018: e.T170524A115656793.

Holcomb, K.L. and Butterfield, T.G. In press. *Rhinoclemmys rubida*. The IUCN Red List of Threatened Species.

Horne, B.D. and Bennett, E.L. 2025. *Chelodina mccordi* (Green Status assessment). The IUCN Red List of Threatened Species 2025: e.T123814489A123814489920251.

Horne, B.D., Chan, E.H., Platt, S.G., and Moll, E.O. 2016. *Batagur affinis*. The IUCN Red List of Threatened Species 2016: e.T170501A1315041.

Horne, B.D., Chan, E.H., Platt, S.G., and Moll, E.O. 2019. *Batagur affinis*. The IUCN Red List of Threatened Species 2019: e.T170501A152041284.

Horne, B.D., Das, I., Hamidy, A., Kusrini, M.D., Guntoro, J., and As-singkily, M. 2020. *Orlitia borneensis*. The IUCN Red List of Threatened Species 2020: e.T15509A724972.

Horne, B.D., Praschag, P., Choudhury, B.C., and Singh, S. 2020. *Melanochelys tricarinata*. The IUCN Red List of Threatened Species 2020: e.T13038A511526.

Horne, B.D., Kusrini, M.D., Hamidy, A., Platt, K., Guntoro, J., and Cota, M. 2021. *Siebenrockiella crassicollis*. The IUCN Red List of Threatened Species 2021: e.T39616A2930856.

Horne, B.D., McCormack, T., and Timmins, R.J. 2021. *Malayemys subtrijuga*. The IUCN Red List of Threatened Species 2021: e.T123770834A2929454.

Horne, B.D., Platt, K., and Praschag, P. 2021. *Nilssonia formosa*. The IUCN Red List of Threatened Species 2021: e.T14765A546244.

Horne, B.D., Praschag, P., and Platt, K. 2021. *Lissemys scutata*. The IUCN Red List of Threatened Species 2021: e.T12124A505402.

Horne, B.D., Platt, S.G., and Bennett, E.L. 2024. *Geochelone platynota* (Green Status assessment). The IUCN Red List of Threatened Species 2024: e.T9013A901320242.

Hughes, D.F. and Meshaka, W.E., Jr. 2025. *Kinosternon subrubrum*. The IUCN Red List of Threatened Species 2025: e.T236638832A236640172.

Iverson, J.B. 2025. *Kinosternon flavescens*. The IUCN Red List of Threatened Species 2025: e.T163421A161721376.

Iverson, J.B. In press. *Terrapene ornata*. The IUCN Red List of Threatened Species.

Jenkins, A.J. and Godwin, J.C. 2024. *Sternotherus depressus*. The IUCN Red List of Threatened Species 2024: e.T20824A242442967.

Johnston, G.R., Donini, J.T., and Dodd, C.K., Jr. 2025. *Kinosternon baurii*. The IUCN Red List of Threatened Species 2025: e.T163429A251955369.

Jones, M.T., Willey, L.L., Macip-Ríos, R., and Lara Resendiz, R.A. 2025. *Kinosternon stejnegeri*. The IUCN Red List of Threatened Species 2025: e.T63666A3127646.

Jones, M.T., Uc-Uc, M., Butterfield, T., Willey, L.L., and Díaz-Gamboa, L. In press. *Terrapene yucatana*. The IUCN Red List of Threatened Species.

Kiester, A.R., Palomo-Ramos, R., Ríos-Arana, J., and Goode, E.V. 2018. *Gopherus flavomarginatus*. The IUCN Red List of Threatened Species 2018: e.T9402A112660985.

Kubisch, E., Alcalde, L., and Waller, T. In press. *Chelonoidis chilensis*. The IUCN Red List of Threatened Species.

Kusrini, M.D., As-singkily, M., Light, C., and Stanford, C.B. 2021. *Indotestudo forstenii*. The IUCN Red List of Threatened Species 2021: e.T10825A499158.

Kusrini, M.D., Hamidy, A., Guntoro, J., Cota, M., and Schoppe, S. 2021. *Notochelys platynota*. The IUCN Red List of Threatened Species 2021: e.T14856A546301.

Lara Resendiz, R.A. In press. *Kinosternon alamosae*. The IUCN Red List of Threatened Species.

Lechowicz, C.J. 2025. *Kinosternon steindachneri*. The IUCN Red List of Threatened Species 2025: e.T236638080A236648309.

Lesher-Gordillo, J., Gallardo-Alvarez, M., Hernández-Marín, A., Reyes-Grajales, E., McKnight, D., Ligon, D., Marlin, J., Barrett, H., and Zenteno-Rúiz, C. In press. *Dermatemys mawii*. The IUCN Red List of Threatened Species.

Leuteritz, T. and Pedrono, M. (Madagascar Tortoise and Freshwater Turtle Red List Workshop). 2008. *Astrochelys yniphora*. The IUCN Red List of Threatened Species 2008: e.T9016A12950950.

Leuteritz, T. and Rioux Paquette, S. (Madagascar Tortoise and Freshwater Turtle Red List Workshop). 2008. *Astrochelys radiata*. The IUCN Red List of Threatened Species 2008: e.T9014A12950491.

Leuteritz, T. and Walker, R. (Madagascar Tortoise and Freshwater Turtle Red List Workshop). 2008. *Pyxis arachnoides*. The IUCN Red List of Threatened Species 2008: e.T19035A50987297.

Leuteritz, T. and Walker, R. 2014. *Pyxis arachnoides* (corrected version). The IUCN Red List of Threatened Species 2014: e.T19035A50987297.

Leuteritz, T. and Walker, R. 2020. *Pyxis arachnoides* (amended version of 2014 assessment). The IUCN Red List of Threatened Species 2020: e.T19035A177075588.

Leuteritz, T., Kuchling, G., Garcia, G., and Velosoa, J. (Madagascar Tortoise and Freshwater Turtle Red List Workshop). 2008. *Erymnochelys madagascariensis*. The IUCN Red List of Threatened Species 2008: e.T8070A12884059.

Leuteritz, T., Randriamahazo, H., and Lewis, R. (Madagascar Tortoise and Freshwater Turtle Red List Workshop). 2008. *Pyxis planicauda*. The IUCN Red List of Threatened Species 2008: e.T19036A8789990.

Li, P., McCormack, T., Rao, D.-Q., Shi, H., Stuart, B., and Wang, L. 2020. *Cuora galbinifrons*. The IUCN Red List of Threatened Species 2020: e.T97357437A123816666.

Li, P., Rao, D.-Q., and Wang, L. 2021. *Mauremys sinensis*. The IUCN Red List of Threatened Species 2021: e.T15026A547319.

Lindeman, P.V. 2025. *Graptemys gibbonsi*. The IUCN Red List of Threatened Species 2025: e.T184436A250638585.

Lindeman, P.V. 2025. *Graptemys pearlensis*. The IUCN Red List of Threatened Species 2025: e.T184437A251707426.

Loc-Barragán, J.A., Woolrich-Piña, G.A., López-Luna, M.A., and Iverson, J.B. In press. *Kinosternon cora*. The IUCN Red List of Threatened Species.

López-Luna, M.A. and Cázares-Hernández, E. In press. *Kinosternon herrerai*. The IUCN Red List of Threatened Species.

Luiselli, L. 2024. *Mauremys leprosa* (Europe assessment). The IUCN Red List of Threatened Species 2024: e.T158468A207995085.

Luiselli, L. 2024. *Testudo graeca* (Europe assessment). The IUCN Red List of

Threatened Species 2024: e.T21646A207669215.

Luiselli, L. 2024. *Testudo hermanni*. The IUCN Red List of Threatened Species 2024: e.T21648A2777071.

Luiselli, L. 2024. *Testudo marginata*. The IUCN Red List of Threatened Species 2024: e.T21653A2777760.

Luiselli, L. and Vamberger, M. 2024. *Emys orbicularis* (Europe assessment). The IUCN Red List of Threatened Species 2024: e.T7717A207667247.

Luiselli, L. and Vamberger, M. 2024. *Mauremys rivulata* (Europe assessment). The IUCN Red List of Threatened Species 2024: e.T158470A207668374.

Luiselli, L., Politano, E., and Lea, J. 2006. *Kinixys homeana*. The IUCN Red List of Threatened Species 2006: e.T11003A3238276.

Luiselli, L., Petrozzi, F., Akani, G.C., Segniagbeto, G.H., and Bour, R. 2019. *Pelusios niger*. The IUCN Red List of Threatened Species 2019: e.T163440A1008712.

Luiselli, L., Agyekumhene, A., Akani, G.C., Allman, P., Diagne, T., Eniang, E.A., Mifsud, D.A., Petrozzi, F., and Segniagbeto, G.H. 2021. *Kinixys homeana*. The IUCN Red List of Threatened Species 2021: e.T11003A18341580.

Luiselli, L., Capula, M., Di Vittorio, M., Marrone, F., and Vamberger, M. 2024. *Emys trinacris*. The IUCN Red List of Threatened Species 2024: e.T158469A778721.

Luiselli, L., Segniagbeto, G.H., Behangana, M., McGovern, P., Diagne, T., Eniang, E.A., and Petrozzi, F. In press. *Kinixys erosa*. The IUCN Red List of Threatened Species.

Luiselli, L., Pham, T.V., and Rhodin, A.G.J. In press. *Pelodiscus variegatus*. The IUCN Red List of Threatened Species.

Machkour-M'Rabet, S., Gallardo Alvarez, M.I., Lesher-Gordillo, J., and Bock, B.C. In press. *Trachemys grayi*. The IUCN Red List of Threatened Species.

Macip-Ríos, R. In press. *Kinosternon oaxacae*. The IUCN Red List of Threatened Species.

Macip-Ríos, R. and Iverson, J.B. In press. *Kinosternon hirtipes*. The IUCN Red List of Threatened Species.

Macip-Ríos, R. and Iverson, J.B. In press. *Kinosternon integrum*. The IUCN Red List of Threatened Species.

Mali, I. and Forstner, M.R.J. In press. *Gopherus berlandieri*. The IUCN Red List of Threatened Species.

Mali, I. and Forstner, M.R.J. In press. *Pseudemys gorzugi*. The IUCN Red List of Threatened Species.

Marques, T.S., Estrades, A., Fallabrino, A., Horta, G.d.F., Portelinha, T.C.G., Yves, A., Valadão, R.M., Cabrera, M.R., Florencia David, M., and Miorando, P.S. In press. *Acanthochelys spixii*. The IUCN Red List of Threatened Species.

Mays, J.D., Smith, L.L., Godwin, J.C., and Hill, P. 2025. *Graptemys barbouri*. The IUCN Red List of Threatened Species 2025: e.T9496A250305609.

McCormack, T. and Stuart, B. 2016. *Cuora bourreti*. The IUCN Red List of Threatened Species 2016: e.T163447A115303472.

McCormack, T. and Stuart, B. 2020. *Cuora bourreti*. The IUCN Red List of Threatened Species 2020: e.T163447A123815632.

McCormack, T., Shi, H., and Stuart, B. 2016. *Cuora galbinifrons*. The IUCN Red List of Threatened Species 2016: e.T97357437A3078734.

McCormack, T., Stuart, B., and Blanck, T. 2016. *Cuora picturata*. The IUCN Red List of Threatened Species 2016: e.T163463A115303820.

McCormack, T., Stuart, B., Blanck, T. 2020. *Cuora picturata*. The IUCN Red List of Threatened Species 2020: e.T163463A123815952.

McCormack, T., van Dijk, P.P., Roberton, S., and Dawson, J.S. 2020. *Mauremys annamensis*. The IUCN Red List of Threatened Species 2020: e.T12876A510309.

Miorando, P.S., Lacava, R.V., Ferrara, C.R., Andrade, P., and Valadao, R.M. In press. *Podocnemis sextuberculata*. The IUCN Red List of Threatened Species.

Moldowan, P.D. and van Dijk, P.P. 2025. *Chelydra serpentina*. The IUCN Red List of Threatened Species 2025: e.T163424A251347989.

Morales-Betancourt, M.A., Lasso, C.A., and Rojas-Runjaic, F.J.M. 2024. *Rhinoclemmys diademata*. The IUCN Red List of Threatened Species 2024: e.T170510A1315527.

Morales-Betancourt, M.A., Lasso, C.A., Rojas-Runjaic, F.J.M. In press. *Chelus orinocensis*. The IUCN Red List of Threatened Species

Mortimer, J.A. and Donnelly, M. (IUCN SSC Marine Turtle Specialist Group). 2008. *Eretmochelys imbricata*. The IUCN Red List of Threatened Species 2008: e.T8005A12881238.

Munscher, E.C., Morrison, M., Siders, Z., Weber, A.S., and Walde, A.D. 2025. *Apalone ferox*. The IUCN Red List of Threatened Species 2025: e.T165597A251135698.

Mwaya, R.T., Malonza, P.K., Ngwava, J.M., Moll, D., Schmidt, F.A.C., and Rhodin, A.G.J. 2019. *Malacochersus tornieri*. The IUCN Red List of Threatened Species 2019: e.T12696A508210.

Northwest Atlantic Leatherback Working Group. 2019. *Dermochelys coriacea* (Northwest Atlantic Ocean subpopulation). The IUCN Red List of Threatened Species 2019: e.T46967827A184748440.

Páez, V.P., Gallego-Garcia, N., and Restrepo, A. 2016. *Podocnemis lewyana*. The IUCN Red List of Threatened Species 2016: e.T17823A1528580.

Páez, V.P., Bock, B.C., Iverson, J.B., Cisneros-Heredia, D.F., Macip-Ríos, R., and Giraldo, A. In press. *Kinosternon leucostomum*. The IUCN Red List of Threatened Species.

Páez, V.P., Bock, B.C., Rhodin, A.G.J., Mittermeier, R.A., Cisneros-Heredia, D.F., and Rocha e Silva, R.d. In press. *Phrynops geoffroanus*. The IUCN Red List of Threatened Species.

Páez, V.P., Rivas, G.A., Rojas-Runjaic, F.J.M., Bock, B.C., Hernández, O., Lasso, C.A., Lasso-Alcalá, O., Morales-Betancourt, M.A., and Escalona, T. In press. *Podocnemis vogli*. The IUCN Red List of Threatened Species.

Páez, V.P., Carr, J.L., Bock, B.C., Garcés-Restrepo, M.F., Galvis-Rizo, C.A., Giraldo, A., and Echeverri-García, L.P. In press. *Rhinoclemmys melanosterna*. The IUCN Red List of Threatened Species.

Perälä, J. 2003a. *Testudo kleinmanni*. The IUCN Red List of Threatened Species 2003: e.T21652A9306908.

Perälä, J. 2003b. *Testudo werneri*. The IUCN Red List of Threatened Species 2003: previously online, now IUCN Red List SIS-archived document.

Petrozzi, F., Luiselli, L., Hema, E.M., Diagne, T., Segniagbeto, G.H., Eniang, E.A., Leuteritz, T.E.J., and Rhodin, A.G.J. 2021. *Centrochelys sulcata*. The IUCN Red List of Threatened Species 2021: e.T163423A1006958.

Pilcher, N.J., Chaloupka, M.Y., and Woods, E. 2012. *Chelonia mydas* (Hawaiian subpopulation). The IUCN Red List of Threatened Species 2012: e.T16285718A16285879.

Platt, K., Horne, B.D., and Praschag, P. 2019. *Batagur trivittata*. The IUCN Red List of Threatened Species 2019: e.T10952A152044061.

Platt, K., Rahman, S., Horne, B.D., and Praschag, P. 2020. *Heosemys depressa*. The IUCN Red List of Threatened Species 2020: e.T39596A2929864.

Platt, K., Praschag, P., and Horne, B.D. 2021. *Chitra vandijki*. The IUCN Red List of Threatened Species 2021: e.T170525A1316195.

Platt, K., Praschag, P., and Horne, B.D. 2021. *Cyclemys fusca*. The IUCN Red List of Threatened Species 2021: e.T170506A1315427.

Portelinha, T.C.G., Marques, T.S., Cunha, F.A.G., Brito, E.S., Valadão, R., Portilla, J.R.C., Páez, V.P., and Lamar, W.W. In press. *Chelus fimbriata*. The IUCN Red List of Threatened Species.

Praschag, P. and Ahmed, M.F. 2021. *Cyclemys gemeli*. The IUCN Red List of Threatened Species 2021: e.T170507A1315452.

Praschag, P. and Singh, S. 2019. *Batagur baska*. The IUCN Red List of Threatened Species 2019: e.T97358453A2788691.

Praschag, P., Ahmed, M.F., Das, I., and Singh, S. 2019. *Batagur kachuga*. The IUCN Red List of Threatened Species 2019: e.T10949A152043133.

Praschag, P., Ahmed, M.F., and Singh, S. 2019. *Geoclemys hamiltonii*. The IUCN Red List of Threatened Species 2019: e.T9029A152050337.

Praschag, P., Platt, K., and Horne, B.D. 2020. *Geochelone platynota*. The IUCN Red List of Threatened Species 2020: e.T9013A123815185.

Praschag, P., de Silva, A., and de Silva, P.K. 2021. *Lissemys ceylonensis*. The IUCN Red List of Threatened Species 2021: e.T123802224A123802334.

Praschag, P., Platt, K., and Horne, B.D. 2021. *Morenia ocellata*. The IUCN Red List of Threatened Species 2021: e.T13873A544192.

Praschag, P., Das, I., Choudhury, B.C., and Singh, S. 2021. *Nilssonia leithii*. The IUCN Red List of Threatened Species 2021: e.T2174A2778380.

Praschag, P., Ahmed, M.F., and Singh, S. 2021. *Nilssonia nigricans*. The IUCN Red List of Threatened Species 2021: e.T2173A2778172.

Praschag, P., Das, I., Ahmed, M.F., and Singh, S. 2021. *Pangshura sylhetensis*. The IUCN Red List of Threatened Species 2021: e.T10950A499618.

Rahman, S., Platt, K., Das, I., Choudhury, B.C., Ahmed, M.F., Cota, M., McCormack, T., Timmins, R.J., and Singh, S. 2019. *Indotestudo elongata*. The IUCN Red List of Threatened Species 2019: e.T10824A152051190.

Rahman, S., Ahmed, M.F., Choudhury, B.C., Praschag, P., and Singh, S.

2021. *Lissemys punctata*. The IUCN Red List of Threatened Species 2021: e.T123802477A3008930.

Reyes-Grajales, E. and Guichard-Romero, C. 2021. *Kinosternon abaxillare*. The IUCN Red Listof Threatened Species 2021: e.T192712123A192712129.

Reyes-Grajales, E., Walde, A.D., and Munscher, E.C. In press. *Chelydra rossignonii*. The IUCN Red List of Threatened Species.

Reyes-Grajales, E., Reynoso-Rosales, V.H., Vásquez-Cruz, M.L., Rivera Arroyo, R.C., Munscher, E., and Walde, A.D. In press. *Claudius angustatus*. The IUCN Red List of Threatened Species.

Reyes-Grajales, E., Munscher, E.C., Walde, A.D., and Iverson, J.B. In press. *Kinosternon acutum*. The IUCN Red List of Threatened Species.

Reyes-Grajales, E., Walde, A., Iverson, J.B., and Munscher, E. In press. *Staurotypus salvinii*. The IUCN Red List of Threatened Species.

Reyes-Grajales, E., Reynoso-Rosales, V.H., Vásquez Cruz, M.L., Rivera Arroyo, R.C., Munscher, E., and Walde, A.D. In press. *Staurotypus triporcatus*. The IUCN Red List of Threatened Species.

Reyes-Grajales, E., García-Caballero, F., and Butterfield, T. In press. *Trachemys yaquia*. The IUCN Red List of Threatened Species.

Rhodin, A.G.J. and Georges, A. 2020a. *Chelodina parkeri*. The IUCN Red List of Threatened Species 2020: e.T4608A21644618.

Rhodin, A.G.J. and Georges, A. 2020b. *Chelodina pritchardi*. The IUCN Red List of Threatened Species 2020: e.T4609A3004755.

Rhodin, A.G.J., Gibbs, J.P., Cayot, L.J., Kiester, A.R., and Tapia, W. 2017. *Chelonoidis phantasticus*. The IUCN Red List of Threatened Species 2017: e.T170517A1315907.

Rhodin, A.G.J., Vinke, T., Vinke, S., Métrailler, S., and Mittermeier, R.A. 2018. *Acanthochelys macrocephala*. The IUCN Red List of Threatened Species 2018: e.T97259978A3138205.

Rhodin, A.G.J., Richards, S., Georges, A., Amepou, Y., and Hamidy, A. 2018. *Pelochelys signifera*. The IUCN Red List of Threatened Species 2018: e.T170526A1316202.

Rhodin, A.G.J., Bressan, R.F., Buskirk, J.R., Cabrera, M.R., Carreira, S., Estrades, A., Mittermeier, R.A., Vinke, S., and Vinke, T. 2018. *Phrynops williamsi*. The IUCN Red List of Threatened Species 2018: e.T172024A1339018.

Rhodin, A.G.J., Villegas Raygoza, B.Y., Folt, B., and Mittermeier, R.A. In press. *Kinosternon angustipons*. The IUCN Red List of Threatened Species.

Ricardez, V., Gordon, M., DeChellis, D., Brown, G.J., and Franklin, C.J. 2025. *Sternotherus carinatus*. The IUCN Red List of Threatened Species 2025: e.T170492A251779169.

Rivas, G.A., Lasso-Alcalá, O., Barros, T.R., Franklin, C.J., Sibira, L.E., Viloria, A.L., Escalona, T., and Páez, V.P. In press. *Mesoclemmys zuliae*. The IUCN Red List of Threatened Species.

RLSPS [Red List Standards and Petitions Subcommittee]. 1996. *Natator depressus*. The IUCN Red List of Threatened Species 1996: e.T14363A4435952.

Roosenburg, W.M., Baker, P.J., Burke, R., Dorcas, M.E., and Wood, R.C. 2019. *Malaclemys terrapin*. The IUCN Red List of Threatened Species 2019: e.T12695A507698.

Rosen, P.C. and Stone, P.A. 2017. *Kinosternon sonoriense* ssp. *longifemorale*. The IUCN Red List of Threatened Species 2017: e.T96710001A91328680.

Santana, D.O. and Cunha, F.A.G. In press. *Mesoclemmys tuberculata*. The IUCN Red List of Threatened Species.

Schiavetti, A., Dias, I.R., Miranda, M.R., Montag, L.F.A., and Fraga, R.E. In press. *Acanthochelys radiolata*. The IUCN Red List of Threatened Species.

Schoppe, S., Diesmos, A.C., and Diesmos, M. 2021. *Siebenrockiella leytensis*. The IUCN Red List of Threatened Species 2021: e.T39599A2929929.

Scott, P.A., Iverson, J.B., and Brown, G.J. 2025. *Sternotherus intermedius*. The IUCN Red List of Threatened Species 2025: e.T218363413A218363420.

Seminoff, J.A. (Southwest Fisheries Science Center, U.S.). 2004. *Chelonia mydas*. The IUCN Red List of Threatened Species 2004: e.T4615A11037468.

Shepherd, C., Horne, B.D., Guntoro, J., and Cota, M. 2021. *Batagur borneoensis*. The IUCN Red List of Threatened Species 2021: e.T163458A1009824.

Stanford, C.B. and Hamidy, A. 2021. *Leucocephalon yuwonoi*. The IUCN Red List of Threatened Species 2021: e.T40761A2933851.

Stemle, L.R. and Howell, H.J. In press. *Gopherus polyphemus*. The IUCN Red List of Threatened Species.

Stone, P.A. 2025. *Kinosternon sonoriense*. The IUCN Red List of Threatened Species 2025: e.T11011A161718519.

Tavares, A.S., Morcatty, T.Q., Rojas-Runjaic, F.J.M., Rojano Bolaño, C., Fagundes, C.K., and Valadão, R.M. In press. *Chelonoidis denticulatus*. The IUCN Red List of Threatened Species.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Acanthochelys macrocephala*. The IUCN Red List of Threatened Species 1996: e.T74A13072810.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Acanthochelys pallidipectoris*. The IUCN Red List of Threatened Species 1996: e.T75A13074224.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Acanthochelys radiolata*. The IUCN Red List of Threatened Species 1996: e.T78A13078282.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Acanthochelys spixii*. The IUCN Red List of Threatened Species 1996: e.T76A13075785.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Actinemys marmorata*. The IUCN Red List of Threatened Species 1996: e.T4969A11104202.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Apalone spinifera* ssp. *ater*. The IUCN Red List of Threatened Species 1996: e.T1820A7781928.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Centrochelys sulcata*. The IUCN Red List of Threatened Species 1996: e.T163423A5605057.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Chelonoidis chilensis*. The IUCN Red List of Threatened Species 1996: e.T9007A12949680.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Chelonoidis denticulata*. The IUCN Red List of Threatened Species 1996: e.T9008A12949796.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Chelonoidis nigra*. The IUCN Red List of Threatened Species 1996: e.T9011A12950160.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Chelonoidis nigra* ssp. *abingdoni*. The IUCN Red List of Threatened Species 1996: e.T9017A12951337.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Chelonoidis nigra* ssp. *becki*. The IUCN Red List of Threatened Species 1996: e.T9018A12951383.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Chelonoidis nigra* ssp. *chathamensis*. The IUCN Red List of Threatened Species 1996: e.T9019A12951430.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Chelonoidis nigra* ssp. *darwini*. The IUCN Red List of Threatened Species 1996: e.T9020A12951477.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Chelonoidis nigra* ssp. *duncanensis*. The IUCN Red List of Threatened Species 1996: e.T9021A12951528.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Chelonoidis nigra* ssp. *guentheri*. The IUCN Red List of Threatened Species 1996: e.T9022A12951582.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Chelonoidis nigra* ssp. *hoodensis*. The IUCN Red List of Threatened Species 1996: e.T9024A12951677.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Chelonoidis nigra* ssp. *microphyes*. The IUCN Red List of Threatened Species 1996: e.T9025A12951724.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Chelonoidis nigra* ssp. *nigra*. The IUCN Red List of Threatened Species 1996: e.T9023A12951629.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Chelonoidis nigra* ssp. *porteri*. The IUCN Red List of Threatened Species 1996: e.T9026A12951777.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Chelonoidis nigra* ssp. *vandenburghi*. The IUCN Red List of Threatened Species 1996: e.T9027A12951826.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Chelonoidis nigra* ssp. *vicina*. The IUCN Red List of Threatened Species 1996: e.T9028A12951875.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Claudius angustatus*. The IUCN Red List of Threatened Species 1996: e.T4959A11102593.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Cyclanorbis elegans*. The IUCN Red List of Threatened Species 1996: e.T6004A12266357.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Cyclanorbis senegalensis*. The IUCN Red List of Threatened Species 1996: e.T6005A12275799.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Cycloderma frenatum*. The IUCN Red List of Threatened Species 1996: e.T6009A12241222.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Elseya bellii*. The IUCN Red List of Threatened Species 1996: e.T40758A10362671.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Elseya georgesi*. The IUCN Red List of Threatened Species 1996: e.T40760A10362891.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Elseya purvisi*. The IUCN Red List of Threatened Species 1996: e.T40759A10362750.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Elusor macrurus*. The IUCN Red List of Threatened Species 1996: e.T7664A12841291.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Emydura macquarii* ssp. *signata*. The IUCN Red List of Threatened Species 1996: e.T7710A12843893.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Emys orbicularis*. The IUCN Red List of Threatened Species 1996: e.T7717A12844431.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Geochelone gigantea*. The IUCN Red List of Threatened Species 1996: e.T9010A12949962.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Gopherus agassizii*. The IUCN Red List of Threatened Species 1996: e.T9400A12983037.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Gopherus berlandieri*. The IUCN Red List of Threatened Species 1996: e.T9401A12983179.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Gopherus polyphemus*. The IUCN Red List of Threatened Species 1996: e.T9403A12983629.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Hydromedusa maximiliani*. The IUCN Red List of Threatened Species 1996: e.T10309A3191766.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Kinixys erosa*. The IUCN Red List of Threatened Species 1996: e.T11002A3238083.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Kinixys natalensis*. The IUCN Red List of Threatened Species 1996: e.T11004A3238959.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Kinosternon acutum*. The IUCN Red List of Threatened Species 1996: e.T11010A3238234.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Kinosternon angustipons*. The IUCN Red List of Threatened Species 1996: e.T11005A3239054.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Kinosternon dunni*. The IUCN Red List of Threatened Species 1996: e.T11008A3237803.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Macrochelys temminckii*. The IUCN Red List of Threatened Species 1996: e.T12589A3362355.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Malaclemys terrapin*. The IUCN Red List of Threatened Species 1996: e.T12695A3373885.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Malacochersus tornieri*. The IUCN Red List of Threatened Species 1996: e.T12696A3373951.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Mesoclemmys dahli*. The IUCN Red List of Threatened Species 1996: e.T17080A6797374.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Mesoclemmys hogei*. The IUCN Red List of Threatened Species 1996: e.T17081A6797504.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Mesoclemmys vanderhaegei*. The IUCN Red List of Threatened Species 1996: e.T17084A6797906.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Mesoclemmys zuliae*. The IUCN Red List of Threatened Species 1996: e.T17083A6797773.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Peltocephalus dumerilianus*. The IUCN Red List of Threatened Species 1996: e.T16511A5972664.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Pelusios broadleyi*. The IUCN Red List of Threatened Species 1996: e.T16529A6000514.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Pelusios castanoides*. The IUCN Red List of Threatened Species 1996: e.T41601A10503545.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Pelusios rhodesianus*. The IUCN Red List of Threatened Species 1996: e.T16530A6002069.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Pelusios subniger*. The IUCN Red List of Threatened Species 1996: e.T41602A10503999.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Pelusios upembae*. The IUCN Red List of Threatened Species 1996: e.T16531A6002802.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Podocnemis erythrocephala*. The IUCN Red List of Threatened Species 1996: e.T17821A7498361.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Podocnemis expansa*. The IUCN Red List of Threatened Species 1996: e.T17822A7500662.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Podocnemis lewyana*. The IUCN Red List of Threatened Species 1996: e.T17823A7502824.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Podocnemis sextuberculata*. The IUCN Red List of Threatened Species 1996: e.T17824A7504825.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Podocnemis unifilis*. The IUCN Red List of Threatened Species 1996: e.T17825A7506933.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Pseudemydura umbrina*. The IUCN Red List of Threatened Species 1996: e.T18457A8294310.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Pseudemys alabamensis*. The IUCN Red List of Threatened Species 1996: e.T18458A8295960.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Rhinemys rufipes*. The IUCN Red List of Threatened Species 1996: e.T17082A6797634.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Rhinoclemmys annulata*. The IUCN Red List of Threatened Species 1996: e.T19501A8941417.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Rhinoclemmys funerea*. The IUCN Red List of Threatened Species 1996: e.T19503A8943143.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Rhinoclemmys nasuta*. The IUCN Red List of Threatened Species 1996: e.T19505A8944337.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Staurotypus salvinii*. The IUCN Red List of Threatened Species 1996: e.T20715A9218885.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Staurotypus triporcatus*. The IUCN Red List of Threatened Species 1996: e.T20716A9218927.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Terrapene nelsoni*. The IUCN Red List of Threatened Species 1996: e.T21643A9304626.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Testudo graeca*. The IUCN Red List of Threatened Species 1996: e.T21646A9305693.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Testudo horsfieldii*. The IUCN Red List of Threatened Species 1996: e.T21651A9306759.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Trachemys adiutrix*. The IUCN Red List of Threatened Species 1996: e.T40762A10363082.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Trachemys decorata*. The IUCN Red List of Threatened Species 1996: e.T22019A9346521.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Trachemys stejnegeri*. The IUCN Red List of Threatened Species 1996: e.T22026A9347196.

TFTSG [Tortoise and Freshwater Turtle Specialist Group]. 1996. *Trachemys terrapen*. The IUCN Red List of Threatened Species 1996: e.T22027A9347243.

Thomas, T.M., Enge, K.M., and Johnston, G.R. 2025. *Macrochelys suwanniensis*. The IUCN Red List of Threatened Species 2025: e.T232768492A232768500.

Timmins, R.J., Hoang, H., and McCormack, T. 2021. *Cyclemys atripons*. The IUCN Red List of Threatened Species 2021: e.T170504A1315080.

Timmins, R.J., Hoang, H., and McCormack, T. 2021. *Cyclemys oldhamii*. The IUCN Red List of Threatened Species 2021: e.T163415A1006544.

Timmins, R.J., Hoang, H., and McCormack, T. 2021. *Cyclemys pulchristriata*. The IUCN Red List of Threatened Species 2021: e.T170508A1315477.

Valenzuela-Ceballos, S., Stanford, C.B., and Castañeda Gaytán, G. In press. *Trachemys taylori*. The IUCN Red List of Threatened Species.

van Dijk, P.P. 2007. *Kinosternon durangoense*. The IUCN Red List of Threatened Species 2007: e.T63668A12704979.

van Dijk, P.P. 2009. *Emys trinacris*. The IUCN Red List of Threatened Species 2009: e.T158469A5199795.

van Dijk, P.P. 2011. *Apalone ferox*. The IUCN Red List of Threatened Species 2011: e.T165597A6065209.

van Dijk, P.P. 2011. *Apalone mutica*. The IUCN Red List of Threatened Species 2011: e.T165596A97398190.

van Dijk, P.P. 2011. *Apalone spinifera*. The IUCN Red List of Threatened Species 2011: e.T163451A5607536.

van Dijk, P.P. 2011. *Chrysemys picta*. The IUCN Red List of Threatened Species 2011: e.T163467A5608383.

van Dijk, P.P. 2011. *Clemmys guttata*. The IUCN Red List of Threatened Species 2011: e.T4968A11103766.

van Dijk, P.P. 2011. *Glyptemys muhlenbergii*. The IUCN Red List of Threatened Species 2013: e.T4967A11103317.

van Dijk, P.P. 2011. *Graptemys barbouri*. The IUCN Red List of Threatened Species 2013: e.T9496A12995762.

van Dijk, P.P. 2011. *Graptemys caglei*. The IUCN Red List of Threatened Species 2011: e.T9497A12996153.

van Dijk, P.P. 2011. *Graptemys ernsti*. The IUCN Red List of Threatened Species 2011: e.T9500A12997190.

van Dijk, P.P. 2011. *Graptemys flavimaculata*. The IUCN Red List of Threatened Species 2011: e.T9498A12996484.

van Dijk, P.P. 2011. *Graptemys geographica*. The IUCN Red List of Threatened Species 2011: e.T165598A6065540.

van Dijk, P.P. 2011. *Graptemys gibbonsi*. The IUCN Red List of Threatened Species 2011: e.T184436A8275938.

van Dijk, P.P. 2011. *Graptemys nigrinoda*. The IUCN Red List of Threatened Species 2011: e.T9502A12997533.

van Dijk, P.P. 2011. *Graptemys oculifera*. The IUCN Red List of Threatened Species 2011: e.T9499A12996892.

van Dijk, P.P. 2011. *Graptemys ouachitensis*. The IUCN Red List of Threatened Species 2011: e.T165599A6066039.

van Dijk, P.P. 2011. *Graptemys pearlensis*. The IUCN Red List of Threatened Species 2011: e.T184437A8276246.

van Dijk, P.P. 2011. *Graptemys pseudogeographica*. The IUCN Red List of Threatened Species 2011: e.T165600A6066439.

van Dijk, P.P. 2011. *Graptemys pulchra*. The IUCN Red List of Threatened Species 2011: e.T170494A6782009.

van Dijk, P.P. 2011. *Graptemys versa*. The IUCN Red List of Threatened Species 2011: e.T9503A12997853.

van Dijk, P.P. 2011. *Kinosternon baurii*. The IUCN Red List of Threatened Species 2011: e.T163429A5605837.

van Dijk, P.P. 2011. *Kinosternon sonoriense*. The IUCN Red List of Threatened Species 2011: e.T11011A97382186.

van Dijk, P.P. 2011. *Kinosternon flavescens*. The IUCN Red List of Threatened Species 2011: e.T163421A5604699.

van Dijk, P.P. 2011. *Kinosternon subrubrum*. The IUCN Red List of Threatened Species 2011: e.T163435A5606303.

van Dijk, P.P. 2011. *Mauremys reevesii*. The IUCN Red List of Threatened Species 2011: e.T170502A6783291.

van Dijk, P.P. 2011. *Pseudemys concinna*. The IUCN Red List of Threatened Species 2013: e.T163444A5606651.

van Dijk, P.P. 2011. *Pseudemys gorzugi*. The IUCN Red List of Threatened Species 2011: e.T18459A8297596.

van Dijk, P.P. 2011. *Pseudemys nelsoni*. The IUCN Red List of Threatened Species 2011: e.T170495A6782280.

van Dijk, P.P. 2011. *Pseudemys peninsularis*. The IUCN Red List of Threatened Species 2013: e.T170496A6782626.

van Dijk, P.P. 2011. *Pseudemys rubriventris*. The IUCN Red List of Threatened Species 2011: e.T18460A8299690.

van Dijk, P.P. 2011. *Pseudemys texana*. The IUCN Red List of Threatened Species 2011: e.T170497A6782942.

van Dijk, P.P. 2011. *Sternotherus carinatus*. The IUCN Red List of Threatened Species 2013: e.T170492A6781357.

van Dijk, P.P. 2011. *Sternotherus depressus*. The IUCN Red List of Threatened Species 2011: e.T20824A9231032.

van Dijk, P.P. 2011. *Sternotherus minor*. The IUCN Red List of Threatened Species 2011: e.T170493A6781671.

van Dijk, P.P. 2011. *Sternotherus odoratus*. The IUCN Red List of Threatened Species 2011: e.T163450A79816811.

van Dijk, P.P. 2011. *Terrapene carolina*. The IUCN Red List of Threatened Species 2011: e.T21641A9303747.

van Dijk, P.P. 2011. *Trachemys gaigeae*. The IUCN Red List of Threatened Species 2011: e.T22024A9346883.

van Dijk, P.P. 2012. *Chelydra serpentina*. The IUCN Red List of Threatened Species 2012: e.T163424A18547887.

van Dijk, P.P. 2016. *Cycloderma frenatum*. The IUCN Red List of Threatened Species 2016: e.T6009A3088072.

van Dijk, P.P. and Canseco-Marquez, L. 2007. *Kinosternon oaxacae*. The IUCN Red List of Threatened Species 2007: e.T11009A3237921.

van Dijk, P.P. and Castañeda Gaytán, G. 2024. *Gopherus flavomarginatus* (Green Status assessment). The IUCN Red List of Threatened Species 2024: e.T9402A940220242.

van Dijk, P.P. and Flores-Villela, O. 2007. *Gopherus flavomarginatus*. The IUCN Red List of Threatened Species 2007: e.T9402A12983328.

van Dijk, P.P. and Flores-Villela, O. 2007. *Trachemys taylori*. The IUCN Red List of Threatened Species 2007: e.T63662A12693739.

van Dijk, P.P. and Hammerson, G.A. 2011. *Terrapene ornata*. The IUCN Red List of Threatened Species 2011: e.T21644A9304752.

van Dijk, P.P. and Harding, J. 2011. *Glyptemys insculpta*. The IUCN Red List of Threatened Species 2011: e.T4965A11102820.

van Dijk, P.P. and Rhodin, A.G.J. 2011. *Emydoidea blandingii*. The IUCN Red List of Threatened Species 2011: e.T7709A12843518.

van Dijk, P.P. and Sindaco, R. 2004. *Emys orbicularis* (Europe regional). The IUCN Red List of Threatened Species 2004: e.T7717A12843950.

van Dijk, P.P., Mateo Miras, J.A., Cheylan, M., Joger, U., Sá-Sousa, P., and Pérez-Mellado, V. 2004. *Mauremys leprosa* (Europe regional). The IUCN Red List of Threatened Species 2004: e.T158468A5199555.

van Dijk, P.P., Lymberakis, P., Mousa Disi, A.M., Ajtic, R., Tok, V., Ugurtas, I., Sevinç, M., and Haxhiu, I. 2004. *Mauremys rivulata* (Europe regional). The IUCN Red List of Threatened Species 2004: e.T158470A5200041.

van Dijk, P.P., Corti, C., Mellado, V.P., and Cheylan, M. 2004. *Testudo graeca* (Europe regional). The IUCN Red List of Threatened Species 2004: e.T21646A9305080.

van Dijk, P.P., Corti, C., Mellado, V.P., and Cheylan, M. 2004. *Testudo hermanni*. The IUCN Red List of Threatened Species 2004: e.T21648A9306057.

van Dijk, P.P., Lymberakis, P., and Böhme, W. 2004. *Testudo marginata*. The IUCN Red List of Threatened Species 2004: e.T21653A9307262.

van Dijk, P.P., Tok, V., Ugurtas, I., Sevinç, M., Werner, Y., Hraoui-Bloquet, S., Sadek, R., and Baha El Din, S. 2004. *Trionyx triunguis* (Europe regional). The IUCN Red List of Threatened Species 2004: e.T62256A12584793.

van Dijk, P.P., Lee, J., Calderón Mandujano, R., Flores-Villela, O., López-Luna, M.A., and Vogt, R.C. 2007. *Chelydra rossignonii*. The IUCN Red List of Threatened Species 2007: e.T63660A97408221.

van Dijk, P.P., Ponce Campos, P., and Garcia Aguayo, A. 2007. *Kinosternon chimalhuaca*. The IUCN Red List of Threatened Species 2007: e.T63667A12695265.

van Dijk, P.P., Lee, J., and Calderón Mandujano, R. 2007. *Kinosternon creaseri*. The IUCN Red List of Threatened Species 2007: e.T11006A3239182.

van Dijk, P.P., Hammerson, G., Lavin, P., and Mendoza Quijano, F. 2007. *Kinosternon herrerai*. The IUCN Red List of Threatened Species 2007: e.T63669A12705142.

van Dijk, P.P., Hammerson, G., Vazquez Diaz, J., Quintero Diaz, G.E., Santos, G., and Flores-Villela, O. 2007. *Kinosternon hirtipes*. The IUCN Red List of Threatened Species 2007: e.T63670A12705290.

van Dijk, P.P., Hammerson, G., Vazquez Diaz, J., Quintero Diaz, G.E., Santos, G., and Flores-Villela, O. 2007. *Kinosternon integrum*. The IUCN Red List of Threatened Species 2007: e.T63671A12705506.

van Dijk, P.P., Lee, J., Calderón Mandujano, R., Flores-Villela, O., and López-Luna, M.A. 2007. *Rhinoclemmys areolata*. The IUCN Red List of Threatened Species 2007: e.T63664A12694597.

van Dijk, P.P., Canseco-Marquez, L., and Muñoz, A. 2007. *Rhinoclemmys*

rubida. The IUCN Red List of Threatened Species 2007: e.T19508A8941198.

van Dijk, P.P., Flores-Villela, O., and Howeth, J. 2007. *Terrapene coahuila*. The IUCN Red List of Threatened Species 2007: e.T21642A9304337.

van Dijk, P.P., Blanck, T., and Lau, M. 2010. *Cuora yunnanensis*. The IUCN Red List of Threatened Species 2010: e.T5957A11964406.

van Dijk, P.P., Harding, J., and Hammerson, G.A. 2011. *Trachemys scripta*. The IUCN Red List of Threatened Species 2011: e.T22028A9347395.

van Dijk, P.P., Rhodin, A.G.J., Cayot, L.J., and Caccone, A. 2017. *Chelonoidis niger*. The IUCN Red List of Threatened Species 2017: e.T9023A3149101.

van Dijk, P.P., Diagne, T., Luiselli, L., Baker, P.J., Turkozan, O., and Taskavak, E. 2017. *Trionyx triunguis*. The IUCN Red List of Threatened Species 2017: e.T62256A96894956.

Vargas-Ramírez, M., Cuadrado-Ríos, S., Gaviria-Hernández, J., and Ceballos, C.P. In press. *Trachemys medemi*. The IUCN Red List of Threatened Species.

Villegas Raygoza, B.Y., Rhodin, A.G.J., and Castañeda Gaytán, G. In press. *Trachemys hartwegi*. The IUCN Red List of Threatened Species.

Vinke, T. and Vinke, S. 2016. *Acanthochelys pallidipectoris*. The IUCN Red List of Threatened Species 2016: e.T75A3139283.

Vogt, R.C., Gonzalez-Porter, G.P., and van Dijk, P.P. 2006. *Dermatemys mawii*. The IUCN Red List of Threatened Species 2006: e.T6493A12783921.

Wallace, B.P., Tiwari, M., and Girondot, M. 2013. *Dermochelys coriacea*. The IUCN Red List of Threatened Species 2013: e.T6494A43526147.

WCMC [World Conservation Monitoring Centre]. 1996. *Cylindraspis indica*. The IUCN Red List of Threatened Species 1996: e.T6061A12383518.

WCMC [World Conservation Monitoring Centre]. 1996. *Cylindraspis inepta*. The IUCN Red List of Threatened Species 1996: e.T6062A12385198.

WCMC [World Conservation Monitoring Centre]. 1996. *Cylindraspis peltastes*. The IUCN Red List of Threatened Species 1996: e.T6063A12388776.

WCMC [World Conservation Monitoring Centre]. 1996. *Cylindraspis triserrata*. The IUCN Red List of Threatened Species 1996: e.T6064A12390055.

WCMC [World Conservation Monitoring Centre]. 1996. *Cylindraspis vosmaeri*. The IUCN Red List of Threatened Species 1996: e.T6065A12391587.

Wibbels, T. and Bevan, E. 2019. *Lepidochelys kempii*. The IUCN Red List of Threatened Species 2019: e.T11533A142050590.

Woolrich-Piña, G.A., Castañeda Gaytán, G., Rhodin, A.G.J., Iverson, J.B., and van Dijk, P.P. In press. *Kinosternon durangoense*. The IUCN Red List of Threatened Species.

Young-Valencia, K., Carr, J.L., Giraldo, A., Forero Medina, G., Botero Botero, A., and Ortega Guío, A. In press. *Chelydra acutirostris*. The IUCN Red List of Threatened Species.

APPENDIX 1

Eponyms Used for Turtle and Tortoise Genera, Subgenera, Species, and Subspecies

Alphabetical listing of all persons (single or grouped) who have been honored with eponyms for named Modern or Holocene turtle or tortoise genera, subgenera, species, or subspecies. The scientific names associated with them are also recorded, including *nomina nuda* or *nomina nova* (replacement or emended names). The listing currently has 346 entries covering 350 people. We would greatly appreciate receiving any updated or corrected information about any of these people, especially for those with imprecise lifetime dates or other missing or incorrect details.

Adanson, Michel (1727–1806), France; botanist who worked in Senegal and collected the type. *Emys adansonii* Schweigger 1812; *Chelys (Hydraspis) adamsonii* Gray 1830e.

Agassiz, Louis (1807–1873), Switzerland / USA; herpetologist at Harvard University who founded the Museum of Comparative Zoology. *Xerobates agassizii* Cooper 1861; *Chelonia agassizii* Bocourt 1868; *Podocnemis agassizii* Coutinho *in* Göldi 1886; *Platypeltis agassizii* Baur 1888c; *Claudius agassizii* Smith and Taylor 1950a.

Albury, Nancy and Michael (1955– and 1958–), USA / Bahamas; paleontologists and naturalists who helped curate and preserve the types. *Chelonoidis alburyorum* † Franz and Franz 2009.

Allen, Robert (c.1940–), USA; museum technician and amateur paleontologist who worked at the Florida Museum of Natural History, Gainesville. *Geochelone alleni* † Auffenberg 1966.

Anderson, John (1833–1900), Scotland, United Kingdom; physician, naturalist, and curator of the Indian Museum in Calcutta. *Lissemys punctata andersoni* Webb 1980a.

Annandale, Nelson (1876–1924), Scotland, United Kingdom; naturalist and herpetologist at the Indian Museum in Calcutta, who co-founded the Zoological Survey of India. *Cyclemys annandalii* Boulenger 1903a; *Hieremys annandalei* Smith 1916.

Arnold, E. Nicholas (1940–2023), United Kingdom; herpetologist at the British Museum of Natural History, London. *Dipsochelys arnoldi* Bour 1982a.

Aubry-Lecomte, Charles E. (1821–1879), France; civil servant and naturalist who worked in Gabon and collected the types. *Cryptopodus aubryi* Duméril 1856.

Babcock, Harold L. (1886–1953), USA; herpetologist and curator at the Museum of the Boston Society of Natural History. *Testudo pardalis babcocki* Loveridge 1935.

Baikie, William B. (1824–1864), United Kingdom; explorer of Africa who obtained the types. *Tetrathyra baikii* Gray 1865a; *Baikiea* Gray 1869a.

Bailey, Demetrius J.S. (1860–1908), United Kingdom / Malaysia; naval officer and naturalist in Sarawak (now part of Malaysia), who collected the type. *Hardella baileyi* Bartlett 1895b.

Baker, Sir William E. (1808–1881), United Kingdom / India; military officer and civil administrator in Sindh, India (now Pakistan), who presented the type. *Batagur bakeri* † Lydekker 1885.

Bangs, Outram (1863–1932), USA; mammalogist at the Museum of Comparative Zoology at Harvard University who discovered this disjunct population. *Pseudemys rubriventris bangsi* Babcock 1937.

Barbour, Roger W. (1919–1993), USA; vertebrate zoologist and herpetologist at the University of Kentucky. *Callopsis pulcherrima rogerbarbouri* Ernst 1978.

Barbour, Thomas (1884–1946), USA; herpetologist and curator at the Museum of Comparative Zoology at Harvard University. *Graptemys barbouri* Carr and Marchand 1942.

Bartlett, Edward (1836–1908), United Kingdom; ornithologist and herpetologist at the British Museum of Natural History, London, and at the Sarawak Museum in Malaysia, who collected the type species. *Bartlettia* Gray 1870f.

Bartram, William (1739–1823), USA; botanist, ornithologist, and explorer of the southern USA. *Testudo bartrami* Daudin 1801.

Baur, Georg (1859–1898), Czechoslovakia / Germany / USA; vertebrate morphologist and herpetologist at Yale University, Clark University, and University of Chicago. *Cinosternum baurii* Garman 1891; *Terrapene bauri* Taylor 1895.

Beale, Thomas (1775–1842), Scotland, United Kingdom; naturalist and merchant who lived and worked in China and obtained the type. *Cistuda bealei* Gray 1831d; *Emys bealii* Gray 1834a.

Beck, Rollo H. (1870–1950), USA; ornithologist and explorer who collected the type. *Testudo becki* Rothschild 1901.

Bélanger, Charles P. (1805–1881), France; botanist and mycologist who collected the type. *Emys belangeri* Lesson 1831b.

Bell, Thomas (1792–1880), United Kingdom; herpetologist, dental surgeon, and Professor of Zoology at King's College, London, who described many reptiles and turtles. *Testudo bellii* Gray 1828; *Emys bellii* Gray 1830e; *Testudo (Kinyxis) belliana* Gray 1830e; *Phrynops bellii* Gray 1844; *Rhinoclemys bellii* Gray 1863c; *Cyclemys bellii* Gray 1863e; *Bellia* Gray 1869a; *Trionyx bellii* Gray 1872a; *Bellemys* Williams 1950a.

Bennett, Edward T. (1797–1836), United Kingdom; physician, zoologist, and secretary of the Zoological Society of London, who obtained the types. *Emys bennettii* Gray 1844.

Bérard, Auguste (1796–1852), France; naval officer who presented the types. *Emys berardii* Duméril and Bibron *in* Duméril and Duméril 1851.

Berdmore, Thomas Matthew (1811–1859), United Kingdom; naturalist and soldier with the East India Company in Burma (now Myanmar), who obtained the types. *Emys berdmorei* Blyth 1859; *Morenia* Gray 1870c.

Berendt, Carl H. (1817–1878), Germany / USA; physician and naturalist who collected the types. *Cinosternum berendtianum* Cope 1865.

Berger, Carl (1871–1962), Germany; missionary in South West Africa (now Namibia), who collected the types. The entry for Berger in Beolens et al. (2011) is incorrect. *Homopus bergeri* Lindholm 1906.

Berlandier, Jean Louis (1803–1851), Belgium / Mexico; naturalist who worked in Mexico and collected the types. *Xerobates berlandieri* Agassiz 1857a.

Bertie, James (Earl of Abingdon) (1653–1699), England, United Kingdom; nobleman in whose honor Abingdon Island was also named. *Testudo abingdonii* Günther 1877.

Betta, Édoardo de (1822–1896), Italy; herpetologist and politician who provided comparative material from Venice for the description of the new species. *Testudo graeca bettai* Lataste 1881.

Bibron, Gabriel (1806–1848), France; herpetologist at the Muséum National d'Histoire Naturelle, Paris. *Trionyx (Gymnopus) bibroni* Owen 1853.

Blanding, William (1772–1857), USA; physician and naturalist who first collected the species. *Cistuda blandingii* Holbrook 1838b.

Bocourt, Marie Firmin (1819–1904), France; naturalist and artist who collected reptiles in Central America. *Rhinoclemmys bocourti* Gray 1873f.

Boddaert, Pieter (1730–1795), Netherlands; physician and naturalist who first described the species. *Testudo boddaerti* Schneider 1787.

Boettger (or Böttger), Oskar (1844–1910), Germany; paleontologist, herpetologist, and curator at the Senckenberg Museum, Frankfurt. *Testudo graeca boettgeri* Mojsisovics 1889; *Testudo boettgeri* Siebenrock 1904a; *Testudo oscarboettgeri* Lindholm 1929.

Bogdanov, Oleg P. (1925–2007), Soviet Union / Uzbekistan; herpetologist who collected the type. *Agrionemys bogdanovi* Chkhikvadze *in* Chkhikvadze, Brushko, and Kubykin 2008.

Boie, Heinrich (1794–1827), Germany / Netherlands; naturalist and herpetologist who worked at the Rijksmuseum van Natuurlijke Historie, Leiden. *Testudo boiei* Wagler 1830a.

Bomare, Jacques-Christophe Valmont de (1731–1807), France; botanist and naturalist who wrote an influential encyclopedia of natural history in the 1760s. *Testudo bomarii* Meyer 1790.

Bosc, Louis Augustin Guillaume (1759–1828), France; botanist and naturalist who served as French consul to the USA. *Terrapene boscii* Merrem 1820.

Bottego, Vittorio (1860–1897), Italy; military officer and explorer who collected the type. *Sternothaerus bottegi* Boulenger 1895a.

Boulenger, George Albert (1858–1937), Belgium / United Kingdom; herpetologist and curator at the British Museum of Natural History, London. *Chelys boulengerii* Baur 1890b; *Hydraspis boulengeri* Bohls 1895; *Homopus boulengeri* Duerden 1906.

Bour, Roger (1947–2020), France; herpetologist, paleontologist, and turtle taxonomist, former member of the Turtle Taxonomy Working Group, and expert on western Indian Ocean giant tortoises. *Astrochelys rogerbouri* † Kehlmaier, Graciá, Ali, Campbell, Chapman, Deepak, Ihlow, Jalil, Pierre-Huyet, Samonds, Vences, and Fritz 2023.

Bourret, René L. (1884–1957), France / Vietnam; naturalist and herpetologist who worked in French Indochina (now Vietnam). *Cuora galbinifrons bourreti* Obst and Reimann 1994.

Bouton, Louis S. (1800–1878), France / Mauritius; botanist and secretary of the Natural History Society of Mauritius. *Testudo boutonii* Günther 1875b.

Branderhorst, Bastiaan (1880–1971), Netherlands / Indonesia; physician and botanist who worked in Dutch New Guinea (now Papua, Indonesia) and Java, Netherlands Indies (now Indonesia), and collected the type. *Emydura branderhorsti* Ouwens 1914.

Broadley, Donald G. (1932–2016), United Kingdom / Zimbabwe; herpetologist and turtle researcher who lived and worked in Zimbabwe. *Pelusios broadleyi* Bour 1986.

Brongniart, Alexandre (1770–1847), France; geologist, paleontologist, and zoologist. *Trionyx brongniarti* Schweigger 1812.

Brooke, Charles (1829–1917), United Kingdom / Malaysia; Rajah of Sarawak (now part of Malaysia) and founder of the Sarawak Museum. *Kachuga brookei* Bartlett 1895a; *Brookeia* Bartlett 1896.

Brygoo, Édouard-Raoul (1920–2016), France / Madagascar; physician and herpetologist who worked in Madagascar. *Pyxoides brygooi* Vuillemin and Domergue 1972.

Buchanan-Hamilton, Francis (1762–1829), Scotland, United Kingdom; physician and botanist who served with the East India Company in India and Burma (now Myanmar). *Emys hamiltonii* Gray 1830e; *Trionyx buchanani* Theobald 1874; *Emys hamiltonoides* † Falconer and Cautley *in* Lydekker 1880.

Burnes, Sir Alexander (1805–1841), United Kingdom; explorer, military officer, and diplomat in Afghanistan who obtained the type. *Homopus burnesii* Blyth 1854.

Buxton, Patrick A. (1892–1955), United Kingdom / Iran; physician and entomologist who worked with the Royal Army Medical Corps in Persia (now Iran) and collected the type. *Testudo buxtoni* Boulenger 1921

Cagle, Fred R. (1915–1968), USA; herpetologist at Tulane University, Louisiana. *Graptemys caglei* Haynes and McKown 1974.

Cann, John R. (1938–), Australia; herpetologist who studied turtles extensively in Australia and New Guinea. *Emydura canni* Worrell 1970; *Chelymys johncanni* Wells and Wellington 1985; *Carettochelys insculpta canni* Wells 2002a; *Chelodina canni* McCord and Thomson 2002.

Cantor, Theodore E. (1809–1860), Denmark / India; physician and naturalist for the East India Company in Calcutta, who also accompanied British troops in China. *Pelochelys cantorii* Gray 1864b; *Cantorella* Gray 1870c; *Pelochelys cantoris* Boulenger 1889.

Capolongo, Domenico (1938–), Italy; geographer and zoologist at the University of Bari. *Emys orbicularis (galloitalica) capolongoi* Fritz 1995.

Carr, Archie F., Jr. (1909–1987), USA; herpetologist, conservationist, and turtle researcher at the University of Florida. *Chelonia mydas carrinegra* Caldwell 1962.

Carvalho, Antenor Leitão de (1910–1985), Brazil; herpetologist and ichthyologist at Museu Nacional, Rio de Janeiro. Common name Carvalho's Slider (*Trachemys adiutrix* Vanzolini 1995).

Chapin, James P. (1889–1964), USA; ornithologist and curator at the American Museum of Natural History, New York, who collected in the Congo. *Pelusios castaneus chapini* Laurent 1965.

Collie, Alexander (1793–1835), Scotland, United Kingdom; surgeon and botanist who worked and lived in Western Australia and collected the species. *Chelodina colliei* Gray 1856a.

Commerson, Philibert (1727–1773), France; botanist and explorer who studied the type. *Testudo commersoni* Vaillant 1898.

Cook, Robert (?–), Australia; amateur herpetologist. *Chelymys cooki* Wells and Wellington 1985.

Coutinho, João Martins da Silva (1830–1889), Brazil; military engineer and participant on the Thayer Amazon Expedition with Louis Agassiz. *Podocnemis coutinhii* Göldi 1886.

Creaser, Edwin P. (1907–1981), USA; biologist at the University of Michigan who collected the types. *Kinosternon creaseri* Hartweg 1934.

Cronwright, Samuel C. (1863–1936), South Africa; farmer, biographer, and naturalist who collected and donated several tortoises to the Albany Museum, Grahamstown. *Chersinella fiski cronwrighti* Hewitt 1934.

Cuming, Hugh (1791–1865), United Kingdom; businessman, naturalist, and collector who obtained the types. *Pelochelys cumingii* Gray 1864b.

Cuvier, Baron Georges (1769–1832), France; paleontologist, anatomist, and zoologist at the Muséum National d'Histoire Naturelle, Paris. *Trionyx (Tetraonyx) cuvieri* Gray 1830e.

Dahl, George (1905–1979), Sweden / Colombia; biologist and ichthyologist who collected the types. *Phrynops (Batrachemys) dahli* Zangerl and Medem 1958.

d'Albertis, Luigi M. (1841–1901), Italy; explorer and naturalist who worked in New Guinea and collected the types. *Emydura albertisii* Boulenger 1888b.

Damelip, Willy (1900s–2007), Vanuatu; archaeologist who worked at the Vanuatu Cultural Center and helped collect the species. ?*Meiolania damelipi* † White, Worthy, Hawkins, Bedford, and Spriggs 2010.

Damon, Robert (1814–1889), United Kingdom; naturalist and geologist who traded in natural historical and geological material and corresponded with Gray. *Damonia* Gray 1869a.

Darling, James J.F. (1859–1929), Ireland; gold prospector, medical officer, and naturalist who collected the type. *Homopus darlingi* Boulenger 1902b.

Darwin, Charles (1809–1882), United Kingdom; naturalist and proponent of the theory of evolution by natural selection who studied tortoises in the Galapagos Islands. *Testudo darwini* Van Denburgh 1907; *Darwintestudo* Antenbrink-Vetter and Vetter 1998.

Daudin, François Marie (1774–1804), France; zoologist, ornithologist, and herpetologist who described many new species of birds and reptiles. *Testudo daudinii* Duméril and Bibron 1835.

de Villiers, Cornelius G.S. (1894–1978), South Africa; physiologist and zoologist at Stellenbosch University. *Pelomedusa galeata devilliersi* Hewitt 1935.

De Vis, Charles Walter (1829–1915), United Kingdom / Australia; director of the Queensland Museum in Brisbane, who described several fossil turtle species. *Devisia* Ogilby 1905.

Diard, Pierre-Médard (1794–1863), France / Indonesia; naturalist, explorer, and collector who founded the Buitenzorg Botanical Gardens in Java. *Cistudo diardii* Duméril and Bibron 1835.

Dillwyn, Lewis L. (1814–1892), Wales, United Kingdom; industrialist, politician, and naturalist who studied the natural history of Labuan and Borneo. *Trionyx dillwynii* Gray 1873c.

Domergue, Charles A. (1914–2008), France / Madagascar; naturalist, ornithologist, and herpetologist who lived and worked in Madagascar most of his life. *Madakinixys domerguei* Vuillemin 1972b.

Donoso-Barros, Roberto (1922–1975), Chile; physician, biologist, and herpetologist at the Universidad de Concepción. *Geochelone donosobarrosi* Freiberg 1973.

d'Orbigny, Alcide (1802–1857), France; paleontologist and naturalist who collected the species. *Emys dorbigni* Duméril and Bibron 1835; *Clemmys (Rhinoclemmys) orbignyi* Fitzinger 1835; *Clemmys dorbignyi* Boulenger 1886b.

Dorr, Emile (1857–1907), France / Senegal; naval officer and naturalist stationed in Senegal who collected the types. *Cinixys dorri* Lataste 1888.

Dorrian, Chris (1942–2013), Australia; amateur herpetologist. *Elseya dorriani* Wells 2002b.

Dorse, Marc (1958–), Australia; educator and herpetologist. *Wollumbinia dorsii* Wells 2009.

Doubleday, Edward (1810–1849), United Kingdom; entomologist at the British Museum of Natural History, London, who collected the type. *Kinosternon (Kinosternon) doubledayii* Gray 1844.

Duerden, James E. (1865–1937), United Kingdom / South Africa; zoologist, herpetologist, and turtle researcher in South Africa. *Chersinella tentoria duerdeni* Hewitt 1933b.

Duméril, André Marie Constant (1774–1860), France; naturalist, herpetologist, and turtle researcher at the Muséum National d'Histoire Naturelle, Paris. *Emys dumeriliana* Schweigger 1812; *Chelys (Hydraspis) dumerilliana* Gray 1830e; *Dumerilia* Grandidier 1867.

Dunn, Emmett Reid (1894–1956), USA; herpetologist at Smith College and curator at the Academy of Natural Sciences, Philadelphia. *Kinosternon dunni* Schmidt 1947.

Durand, Sir Henry Marion (1812–1871), United Kingdom / India; military officer and Lieutenant Governor of Punjab, India, who helped collect the type. *Batagur durandi* † Lydekker 1885.

Dussumier, Jean-Jacques (1792–1883), France; shipowner in the merchant marine and naturalist collector. *Testudo dussumieri* Schlegel *in* Gray 1830e; *Testudo dussumieri* Gray 1831d; *Chelonia dussumierii* Duméril and Bibron 1835.

Duvaucel, Alfred (1792–1824), France; naturalist, explorer, and collector for the Muséum National d'Histoire Naturelle, Paris. *Emys duvaucelii* Duméril and Bibron 1835; *Gymnopus duvaucelii* Duméril and Bibron 1835.

Eden, Sir Ashley (1831–1887), United Kingdom / India / Myanmar; official and diplomat in British India and first civil governor of Burma (now Myanmar). *Melanochelys edeniana* Theobald 1876.

Effeldt, Rudolf (1821–1876), Germany; herpetologist and chiropterologist who collected the type specimen. *Cinosternon effeldtii* Peters 1873.

Eiselt, Josef (1912–2001), Austria; herpetologist who worked in Asia Minor and collected the type. *Emys orbicularis eiselti* Fritz, Baran, Budak, and Amthauer 1998b.

Elliot, Sir Walter (1830–1887), Scotland, United Kingdom / India; naturalist who served with the East India Company in India. *Batagur ellioti* Gray 1862b.

Elsey, Joseph R. (1834–1857), United Kingdom / Australia; surgeon and naturalist who explored northern Australia. *Chelymys elseyi* Gray 1864d; *Elseya* Gray 1867; *Chelymys elseya* Gray 1870c.

Emmott, Angus (1962–), Australia; grazier, conservationist, and naturalist. *Emydura macquarii emmotti* Cann, McCord, and Joseph-Ouni *in* McCord, Cann, and Joseph-Ouni 2003.

Emory, William H. (1811–1887), USA; army officer, civil engineer, and surveyor of the USA-Mexico border, who obtained the types. *Aspidonectes emoryi* Agassiz 1857a.

Enriques, Paolo (1878–1932), Italy; zoologist at the University of Padua who mentored the author. *Testudo enriquesi* Parenzan 1932.

Erhard, Andreas (1960–), Germany; historian, conservationist, and naturalist. *Mauremys leprosa erhardi* Schleich 1996a.

Ernst, Carl H. (1938–2018), USA; herpetologist and turtle researcher at George Mason University, Virginia. *Graptemys ernsti* Lovich and McCoy 1992.

Ernst, Evelyn M. (1948–), USA; educator and parasitologist and the senior author's wife. *Cuora evelynae* Ernst and Lovich 1990.

Falconer, Hugh (1808–1865), Scotland, United Kingdom / India; physician, paleontologist, and geologist who worked in India and described several fossil animals, including a giant tortoise. *Testudo (Scapia) falconeri* Gray 1869a; *Batagur falconeri* † Lydekker 1885.

Feliz, Gerson (1982–), Dominican Republic; field biology assistant in the Grupo Jaragua working on conservation of natural resources in Pedernales. *Chelonoidis gersoni* † Viñola-López and Almonte 2022.

Fisk, Reverend George H.R. (1829–1911), United Kingdom / South Africa; clergyman and naturalist who worked in South Africa in the 1870s and collected the types. *Testudo fiski* Boulenger 1886a.

Flower, Stanley S. (1871–1946), United Kingdom / Egypt; army officer, zoologist, and director of the Zoological Gardens at Giza, Egypt. *Testudo floweri* Bodenheimer 1935.

Ford, George H. (1809–1876), South Africa / United Kingdom; naturalist artist who illustrated for Sir Andrew Smith and the British Museum of Natural History, London. *Fordia* Gray 1869a.

Forsten, Eltio A. (1811–1843), Netherlands / Indonesia; naturalist and botanist who explored and collected in the Dutch East Indies (now Indonesia). *Testudo forstenii* Schlegel and Müller 1845.

Fraser, Louis (1810–1866), United Kingdom; naturalist and curator of the Museum of the Zoological Society of London, who collected the type. *Emys fraseri* Gray 1869c.

Frewer, Jason (1900s–), United Kingdom / USA; commercial turtle breeder and dealer in Florida. *Elseya (Hanwarachelys) freweri* Joseph-Ouni and McCord 2023b.

Gabb, William M. (1839–1878), USA / Costa Rica; paleontologist and geographer who collected the type. *Chelopus gabbii* Cope 1875a.

Gadow, Hans F. (1855–1928), Germany / United Kingdom; zoologist, ornithologist, and curator at Cambridge University. *Testudo gadowi* Van Denburgh 1914.

Gaige, Helen T. (1890–1976), USA; female herpetologist and curator of the Museum of Zoology at the University of Michigan, who collected the type. *Pseudemys scripta gaigeae* Hartweg 1939.

Gasconi, Alfred (1842–1929), France / Senegal; politician who obtained the original types. *Pelomedusa gasconi* Rochebrune 1884.

Gaudichaud-Beaupré, Charles (1789–1854), France; botanist who

collected the type. *Platemys gaudichaudii* Duméril and Bibron 1835.

Gemel, Richard (1948–), Austria; herpetologist who collected the type. *Cyclemys gemeli* Fritz, Guicking, Auer, Sommer, Wink, and Hundsdörfer 2008b.

Geoffroy Saint-Hilaire, Étienne (1772–1844), France; zoologist who obtained the type. *Emys geoffroana* Schweigger 1812; *Emys geoffroyana* Gray 1830e; *Platemys geoffreana* Duméril and Bibron 1835.

Georges, Arthur (1953–), Australia; geneticist and herpetologist at the University of Canberra and turtle researcher. *Elseya georgesi* Cann 1997a.

Gibbons, J. Whitfield (Whit) (1939–), USA; herpetologist and turtle researcher at Savannah River Ecology Laboratory, South Carolina. *Graptemys gibbonsi* Lovich and McCoy 1992.

Giebel, Christoph G.A. (1820–1881), Germany; zoologist and paleontologist at the University of Halle. *Cyclemys giebelii* Hubrecht 1881.

Goldman, Edward A. (1873–1946), USA; mammalogist and botanist who collected the type. *Terrapene goldmani* Stejneger 1933.

Goode, Eric V. (1957–), USA; hotelier, herpetologist, and conservationist who founded Turtle Conservancy and supported land purchase to protect this species as it was being described. *Gopherus evgoodei* Edwards, Karl, Vaughn, Rosen, Meléndez Torres, and Murphy 2016.

Goode, John (1927–2008), United Kingdom / Australia; journalist and amateur herpetologist. *Tropicochelymys goodei* Wells and Wellington 1985.

Gouffé, Alphonse A. (1845–1931), United Kingdom; ship and insurance broker who obtained the type. *Testudo gouffei* Rothschild 1906.

Grandidier, Alfred (1836–1921), France / Madagascar; naturalist and explorer who worked in Madagascar and first described this species. *Testudo grandidieri* † Vaillant 1885b.

Granger, Walter W. (1872–1941), USA; paleontologist, curator at the American Museum of Natural History, New York, and collector of the type. *Geoclemys grangeri* Schmidt 1925.

Grant, Major Chapman (1887–1983), USA; herpetologist and grandson of U.S. President Ulysses S. Grant, who collected the species. *Pseudemys granti* Barbour and Carr 1941.

Gray, John Edward (1800–1875), United Kingdom; herpetologist, turtle taxonomist, and curator at the British Museum of Natural History, London. *Testudo graii* Duméril and Bibron 1835; *Chersina grayi* Strauch 1865; *Clemmys grayi* Strauch 1865; *Emys grayi* Bocourt 1868; *Emys grayi* Günther 1869; *Trionyx grayii* Theobald 1875.

Gray, Samuel Oscar (1873–?), United Kingdom; grand-nephew of the author (Bour *in* Pritchard 2012:54). *Oscaria* Gray 1873g.

Grochowski, Inne (1900s), Soviet Union; female herpetologist who first collected the species. *Annamemys grochovskiae* Dao 1957.

Gronovius, Laurentius Theodorus (1730–1777), Netherlands; naturalist and ichthyologist who had a large private museum. *Emys gronovii* Merrem 1820.

Gunalen, Danny (1975–), Indonesia; reptile exporter and herpetologist who obtained the type. *Chelodina gunaleni* McCord and Joseph-Ouni 2007a; *Emydura gunaleni* Smales, McCord, Cann, and Joseph-Ouni 2019a.

Günther, Albert C.L.G. (1830–1914), Germany / United Kingdom; zoologist, herpetologist, ichthyologist, and curator at the British Museum of Natural History, London. *Dogania guentheri* Gray 1862c; *Testudo guntheri* Baur 1889; *Testudo guentheri* Gadow 1894; *Testudo guntheri* Gadow *in* Van Denburgh 1914; *Geochelone elephantopus guentheri* Baur *in* Pritchard 1971b.

Hamilton-Gordon, Arthur (1829–1912), United Kingdom / Trinidad; nobleman and Governor of Trinidad, who obtained the type. *Hydraspis gordoni* Gray 1868.

Hardwicke, Thomas (1756–1835), United Kingdom / India; military officer and naturalist who served with the East India Company in India. *Kachuga (Dongoka) hardwickii* Gray 1869a; *Hardella* Gray 1870c; *Testudo hardwickei* Cotton 1918.

Harlan, Richard (1796–1843), USA; physician, vertebrate paleontologist, and anatomist. *Trionyx harlani* Bell *in* Harlan 1835.

Hartweg, Norman E. (1904–1964), USA; herpetologist and curator at the Museum of Zoology at the University of Michigan. *Amyda spinifera hartwegi* Conant and Goin 1948; *Pseudemys scripta hartwegi* Legler 1990.

Haser, Brother Nicolas (1883–1900s), France / China; priest who worked in China and collected the species. *Amyda schlegelii haseri* Pavlov 1932.

Heinroth, Oskar (1871–1945), Germany; biologist, ethologist, and director of the Berlin Aquarium who presented living specimens of the species. *Sternothaerus heinrothi* Kanberg 1924.

Henry, T. Charlton (1825–1877), USA; surgeon with the U.S. Army who collected the type. *Kinosternum henrici* Le Conte 1860.

Hermann (or Herrmann), Johann (1738–1800), France; zoologist at the University of Strasbourg in Alsace, who first recognized this species as distinct. *Testudo hermanni* Gmelin 1789; *Emys herrmanni* Schweigger 1812.

Herrera López, Alfonso Luis (1868–1942), Mexico; zoologist and director of the National Museum of Mexico. *Kinosternon herrerai* Stejneger 1925.

Hilton, John W. (1904–1983), USA; landscape artist, geologist, and zoologist, who collected the types. *Pseudemys scripta hiltoni* Carr 1942.

Hoffmann, A. (1700s–1800s), Austria; naturalist who revised and published Funke's Naturhistorischer Atlas in Vienna in 1825. *Emys (Emys) hofmanni* Fitzinger 1835.

Hoge, Alphonse R. (1912–1982), Brazil; herpetologist at the Instituto Butantan, São Paulo. *Phrynops hogei* Mertens 1967a.

Holbrook, John Edwards (1794–1871), USA; physician, anatomist, and herpetologist at the Medical College of South Carolina. *Emys holbrookii* Gray 1844.

Home, Sir Everard (1756–1832), United Kingdom; physician, anatomist, and naturalist who presented the types to the British Museum of Natural History. *Kinixys homeana* Bell 1827.

Horsfield, Thomas (1773–1859), USA / Indonesia / India; physician and naturalist who worked in the Dutch East Indies (now Indonesia) and India and obtained the type. *Testudo horsfieldii* Gray 1844.

Hoy, Philo R. (1816–1892), USA; ornithologist and entomologist who collected the species. *Ptychemys hoyi* Agassiz 1857a.

Irwin, Steve (1962–2006), Australia; herpetologist, wildlife promoter, and television personality. *Elseya irwini* Cann 1997c.

Iverson, John B. (1949–), USA; herpetologist and turtle researcher at Earlham College, Indiana. *Mauremys iversoni* Pritchard and McCord 1991; *Trachemys venusta iversoni* McCord, Joseph-Ouni, Hagen, and Blanck 2010.

Jerdon, Thomas C. (1811–1872), United Kingdom / India; physician, zoologist, and ornithologist with the East India Company in India. *Jerdonella* Gray 1870c.

Johnston, C. Stuart (1900–1939), USA; paleontologist who collected the type and described the species in an unpublished dissertation. *Geochelone johnstoni* † Auffenberg 1962.

Jordan, E. Knowles (1800s–1900s), United Kingdom / Zambia; political official and naturalist in Northern Rhodesia (now Zambia) who collected the types. *Kinixys jordani* Hewitt 1931.

Jukes, Brian (?–), Australia; amateur herpetologist. *Elseya jukesi* Wells 2002b; *Elseya jukesi* Wells 2007b.

Kästle, Werner (1926–2019), Germany; amateur herpetologist. *Mauremys leprosa wernerkaestlei* Schleich 1996a.

Keegan, William F. (1959–), USA; archaeologist and curator at the Florida Museum of Natural History, Gainesville, who helped collect the types. *Geochelone alburyorum keegani* † Franz, Albury, and Steadman 2020.

Kemp, Richard M. (1825–1908), Bahamas / USA; fisherman who obtained the types. *Thalassochelys (Colpochelys) kempii* Garman 1880.

Klauber, Laurence M. (1883–1968), USA; electrical engineer, herpetologist, and authority on rattlesnakes, who inspired the author. *Terrapene klauberi* Bogert 1943.

Kleinmann, Édouard (1832–1901), France / Egypt; banker, naturalist, and director of Crédit Lyonnais in Egypt, who collected the types. *Testudo kleinmanni* Lortet 1883.

Kohn, Gustave (1837–1906), USA / France; amateur ornithologist and naturalist who collected the types. *Malacoclemmys kohnii* Baur 1890a.

Krauss, Christian Ferdinand (1812–1890), Germany; zoologist and director of the Staatliches Museum für Naturkunde, Stuttgart, who presented the types. *Onychochelys kraussi* Gray 1873i.

Krefft, Gerard (1830–1881), Germany / Australia; naturalist, herpetologist, and director at the Australian Museum, Sydney. *Chelymys krefftii* Gray 1871c.

Kuchling, Gerald (1952–), Austria / Australia; herpetologist, conservationist, and turtle researcher. *Chelodina kuchlingi* Cann 1997d.

Kuhl, Heinrich (1797–1821), Germany / Indonesia; zoologist and herpetologist who worked in the Netherlands and on Java in the Dutch East Indies (now Indonesia). *Emys kuhlii* Gray 1831d.

Kuznetsov, Valentin V. (1923–2006), Soviet Union / Kazakhstan; paleoherpetologist at the Institute of Zoology of Academy of Sciences of Kazakhstan. *Agrionemys kazachstanica kuznetzovi* Chkhikvadze, Ataev, Shammakov, and Zatoka *in* Chkhikvadze, Ataev, and Shammakov 2009.

Lacepède (or La Cepède), Baron B.G.E. de (1756–1825); France; naturalist, herpetologist, politician, and curator at the Muséum National d'Histoire Naturelle, Paris. *Testudo cepediana* Daudin 1801; *Caretta cepedii* Merrem 1820.

Lamare-Picquot, Christophe-Augustin (1785–1873), France; pharmacist, naturalist, and explorer who collected the type. *Emys piquotii* Lesson 1831a; *Emys picquotii* Lesson *in* Duméril and Bibron 1835.

Lambert, Michael R.K. (1941–2004), United Kingdom; entomologist and herpetologist at the Natural Resources Institute, University of Greenwich. *Testudo graeca lamberti* Pieh and Perälä 2004.

Lanza, Benedetto (1924–2016), Italy; herpetologist and chiropterologist at the University of Florence. *Emys orbicularis (galloitalica) lanzai* Fritz 1995.

Lasala, Colonel P. Perez de (1800s), Argentina; naval officer who presented the type. *Spatulemys lasalae* Gray 1872b.

Laurent, Raymond F. (1917–2005), Belgium; herpetologist who researched African turtles of the genus *Pelusios*. *Pelusios williamsi laurenti* Bour 1984a.

Lavarack, Sue and Jim (1926– and 1922–2014), Australia; amateur paleontologists who worked on and helped direct the Riversleigh Expedition and collected the type. *Emydura lavarackorum* † White and Archer 1994.

Le Loi (1385–1433), Vietnam; ancient king who, according to legend, returned a magic sword to the sacred turtle of Hoan Kiem Lake in Hanoi after his victory over Chinese occupiers. *Rafetus leloii* Ha 2000.

Leach, William Elford (1791–1836), United Kingdom; zoologist and marine biologist at the British Museum of Natural History, London. *Sternothaerus leachianus* Bell 1825a.

Leichhardt, Friedrich Wilhelm Ludwig (1813–1848), Germany / Australia; explorer and naturalist in Australia in whose honor the Leichhardt River was also named. *Tropicochelymys leichhardti* Wells and Wellington 1985.

Leith, Andrew H. (1827–1882), United Kingdom / India; physician, naturalist, and malacologist who worked in India on the Bombay Sanitary Commission and collected the type. *Testudo leithii* Günther 1869; *Pangshura leithii* Gray 1870c; *Trionyx leithii* Gray 1872a.

Lesson, René Primevère (1794–1849), France; surgeon, naturalist, ornithologist, and herpetologist. *Tetraonyx lessonii* Duméril and Bibron 1835.

LeSueur, Charles Alexander (1778–1846), France / USA; naturalist, artist, and explorer who collected turtles in the United States. *Emys lesueurii* Gray 1830e.

Lever, Sir Ashton (1729–1788), United Kingdom; naturalist collector and founder of the Leverian Museum in London that housed many specimens from the circumglobal voyages of James Cook. *Emys (Cistuda) amboinensis leveriana* Gray 1830e.

Lewy, Bernard (1817–1863), Denmark / France / Colombia; chemist and naturalist who worked in Bogotá, Colombia, and collected the type. *Podocnemis lewyana* Duméril 1852.

Licent, Père Émile (1876–1952), France / China; priest and naturalist who worked in China and collected the species. *Amyda schlegelii licenti* Pavlov 1932.

Lidth de Jeude, Theodoor Gerard van (1788–1863), Netherlands; physician, veterinarian, and zoologist at Utrecht University who presented the type to the British Museum. *Trionyx jeudi* Gray 1869a.

Liu, Mong-Song (?–), Taiwan; fossil collector who owns the types. *Trionyx liupani* † Tao 1986.

Livingstone, David (1813–1873), Scotland, United Kingdom; physician, missionary, and explorer of Africa, who collected and presented the type. *Aspidochelys livingstonii* Gray 1860a.

Llerena Sánchez, Don Fausto (1940–), Ecuador; park ranger at Galápagos National Park responsible for the care of the giant tortoises at the Charles Darwin Station. *Chelonoidis donfaustoi* Poulakakis, Edwards, and Caccone *in* Poulakakis, Edwards, Chiari, Garrick, Russello, Benavides, Watkins-Colwell, Glaberman, Tapia, Gibbs, Cayot, and Caccone 2015.

Loveridge, Arthur (1891–1980), United Kingdom / Kenya / USA; herpetologist and curator at first the Nairobi Museum and then the Museum of Comparative Zoology, Harvard University. *Testudo loveridgii* Boulenger 1920.

Lutz, Bertha (1894–1976), Brazil; female herpetologist, feminist, and politician, who collected the type. *Hydraspis lutzi* Ihering *in* Luederwaldt 1926.

Maack (or Maak), Richard K. (1825–1886), Estonia / Russia; educator and naturalist who explored the Amur River. *Trionyx maackii* Brandt 1857; *Trionyx maaki* Chkhikvadze 1987.

Mackay, A.C. (1800s–1900s), Australia; engineer who worked for the Australian Guano Company on Walpole Island and collected the types in 1922. *Meiolania mackayi* † Anderson 1925.

Macquarie, Lachlan (1762–1824), Scotland, United Kingdom / Australia; governor of New South Wales, Australia, in whose honor the Macquarie River was named. *Emys macquaria* Cuvier 1829; *Chelys (Hydraspis) macquarii* Gray 1830e; *Hydraspis macquarrii* Gray 1831d; *Platemys macquaria* Duméril and Bibron 1835; *Emydura macquariae* Boulenger 1889.

Malone, Japheth Vernon (J.V.) (1886–1973), Bahamas; politician from Abaco, Bahamas, and Commissioner for Inagua, who collected the types. *Pseudemys malonei* Barbour and Carr 1938.

Mann, William M. (1886–1960), USA; entomologist and director of the U.S. National Zoological Park, Washington, D.C. *Geoemyda manni* Dunn 1930.

Maran, Jérôme (1973–), France; amateur herpetologist and collector of the type. *Pelusios marani* Bour 2000.

Marcano Fondeur, Eugenio de Jesús (1923–2003), Dominican Republic; botanist, zoologist, and cave explorer, who first discovered giant tortoise fossils on Hispaniola. *Chelonoidis marcanoi* † Turvey, Almonte, Hansford, Scofield, Brocca, and Chapman 2017.

Martin, Joseph (1788–1826), France / French Guiana; botanist and director of the botanical garden in Cayenne, French Guiana, who collected the type. *Testudo martinella* Daudin 1802.

Matz, Gilbert (1934–), France; biologist and herpetologist at the University of Angers. *Pyxis arachnoides matzi* Bour 1979.

Maw, Solomon (1800s), United Kingdom; British naval officer who collected the type. *Dermatemys mawii* Gray 1847; *Dermatemys mavei* Cope 1865; *Dermatemys mawei* Neill and Allen 1959.

McCord, Christine, Karen, and Anne (1978–, 1979–, and 1950–), USA; the junior author's two daughters and wife. *Cuora chriskarannarum* Ernst and McCord 1987.

McCord, William P. (1950–), USA; veterinarian and herpetologist. *Cuora mccordi* Ernst 1988; *Chelodina mccordi* Rhodin 1994b.

Medem, Federico (Count Friedrich Johann von Medem) (1912–1984), Latvia / Germany / Colombia; nobleman, zoologist, herpetologist, and director of the Instituto Roberto Franco in Villavicencio. *Trachemys medemi* Vargas-Ramírez, del Valle, Ceballos, and Fritz 2017.

Meier, Elmar (1957–), Germany; turtle breeder and conservationist focused on the genus *Cuora*. *Cuora cyclornata meieri* Blanck, McCord, and Le 2006a.

Merklen, Monsieur (1900s) (given name not discovered), France / Vietnam; official and naturalist working for the Service Forestier de l'Indochine in Annam, French Indochina (now Vietnam), who collected the types. *Annamemys merkleni* Bourret 1940.

Mertens, Robert F.W. (1894–1975), Russia / Germany; herpetologist and director of the Senckenberg Museum, Frankfurt. *Testudo hermanni robertmertensi* Wermuth 1952; *Kinixys belliana mertensi* Laurent 1956.

Metter, Dean E. (1932–2001), USA; zoologist at the University of Missouri and mentor to the author. *Pseudemys concinna metteri* Ward 1984.

Milius, Baron Pierre Bernard (1773–1829), France / French Guiana; nobleman, naval officer, and Governor of French Guiana, who obtained the type specimen. *Platemys miliusii* Duméril and Bibron 1835.

Młynarski, Marian (1926–), Poland; paleontologist and herpetologist at the Institute of Systematics and Evolution of Animals, Polish Academy of Sciences. *Geochelone mlynarskii* † Auffenberg 1988.

Moll, Edward O. (1939–), USA; herpetologist at Eastern Illinois University and first Chair of the IUCN Freshwater Chelonian Specialist Group, precursor to the IUCN Tortoise and Freshwater Turtle Specialist Group. *Pseudemys scripta emolli* Legler 1990; *Batagur affinis edwardmolli* Praschag, Holloway, Georges, Päckert, Hundsdörfer, and Fritz 2009a.

Morafka, David J. (1946–2004), USA; zoologist and herpetologist at California State University. *Gopherus morafkai* Murphy, Berry, Edwards, Leviton, Lathrop, and Riedle 2011.

Morton, Samuel G. (1799–1851), USA; physician, paleontologist, and naturalist at the Academy of Natural Sciences, Philadelphia, who obtained the type. *Trionyx mortoni* Hallowell 1844.

Mouhot, Henri (1826–1861), France; naturalist and explorer in Thailand and Laos who collected the types. *Cyclemys mouhotii* Gray 1862a.

Mühlenberg, Gotthilf H.E. (1753–1815), Germany / USA; clergyman and botanist who discovered the species. *Testudo muhlenbergii* Schoepff 1801, *Geoclemmys mühlenburgii* Gray 1870c, *Glyptemys muehlenbergii* Frétey 2024.

Murray, Leo T. (1902–1958), USA; naturalist at Texas A&M University and director of the Strecker Museum. *Kinosternon murrayi* Glass and Hartweg 1951.

Nelson, Edward W. (1855–1934), USA; naturalist and ethnologist, Chief of the U.S. Biological Survey, who collected the type. *Terrapene nelsoni* Stejneger 1925.

Nelson, George (1876–1962), USA; naturalist and preparator at the Museum of Comparative Zoology, Harvard University, who collected the type. *Pseudemys nelsoni* Carr 1938a.

Néraud, Jules (1795–1855), France; botanist who collected bone fragments of this species on Mauritius. *Testudo neraudii* Gray 1831d.

Neumann, Oscar R. (1867–1946), Germany; ornithologist and mammalogist who did field work in East Africa and worked at the Museum für Naturkunde, Berlin. *Pelomedusa neumanni* Petzold, Vargas-Ramírez, Kehlmaier, Vamberger, Branch, Du Preez, Hofmeyr, Meyer, Schleicher, Široký, and Fritz 2014.

Newton, Francesco (1864–1909), Portugal; botanist who collected the type. *Trionyx cartilagineus newtoni* Ferreira 1897.

Nikolsky, Alexander M. (1858–1942), Russia / Soviet Union; naturalist and herpetologist at the Zoological Museum of the Academy of Sciences, St. Petersburg. *Testudo graeca nikolskii* Chkhikvadze and Tuniyev 1986.

Nilsson, Sven (1787–1883), Sweden; zoologist, naturalist, and director of the Naturhistoriska Riksmuseet, Stockholm. *Nilssonia* Gray 1872a.

Noguey, Gustave (1800s), France; naturalist of the Société Linnéenne de Bordeaux and a traveling companion to the author. *Homopus nogueyi* Lataste 1886.

Nutaphand (or Nutphand), Wirot (1932–2005), Thailand; airforce officer, naturalist, and herpetologist. *Geoemyda trijuga wiroti* Reimann *in* Nutaphand 1979; *Testudo nutapundi* Reimann and Nutaphand *in* Nutaphand 1979.

Nuttal, Thomas (1786–1859), United Kingdom / USA; naturalist and botanist. *Chrysemys nuttalii* Agassiz 1857a; *Chrysemys nuttalii* Agassiz 1857b.

Obst, Fritz-Jürgen (1939–2018), Germany; herpetologist and director of the Staatliches Museum für Tierkunde, Dresden. *Emys orbicularis fritzjuergenobsti* Fritz 1993; *Pyxidea mouhotii obsti* Fritz, Andreas, and Lehr 1998a.

Oelrich, Thomas M. (1924–2008), USA; anatomist, histologist, and paleoherpetologist at the University of Michigan Medical School. *Geochelone (Hesperotestudo) oelrichi* † Holman 1972.

Oldham, Thomas (1816–1878), Ireland / India; geologist and naturalist at the Geological Survey of India who collected turtles and mammals for the British Museum. The entry for Oldham in Beolens et al. (2011) is incorrect. *Cyclemys oldhamii* Gray 1863e; *Kachuga oldhami* Gray 1869a.

Oyama, Seiho (1912–1996), Japan; businessman and amateur archaeologist who first collected this species. *Manouria oyamai* † Takahashi, Otsuka, and Hirayama 2003.

Pallas, Peter Simon (1741–1811), Germany / Russia; zoologist and naturalist at the Imperial Academy of Sciences in St. Petersburg. *Testudo graeca pallasi* Chkhikvadze 1989; *Testudo graeca pallasi* Chkhikvadze and Bakradze 2002.

Pan, Chang-Wu (1932–2015), Taiwan; amateur paleontologist, fossil collector, and owner of the types. *Chinemys pani* † Tao 1985; *Trionyx liupani* † Tao 1986; *Ocadia sinensis changwui* † Tao 1988.

Pan, Liu (1900s), China; herpetologist at Shanghai Museum of Natural History, Shanghai. *Cuora pani* Song 1984.

Parker, Frederick S. (1941–), India / Australia / Papua New Guinea; government patrol officer (*kiap*) and herpetologist in Papua New Guinea, who first collected the species. *Chelodina parkeri* Rhodin and Mittermeier 1976.

Parker, Hampton W. (1897–1968), United Kingdom; herpetologist at the British Museum of Natural History, London, who first recognized the taxon as distinctive. *Melanochelys trijuga parkeri* Deraniyagala 1939.

Peck, Herbert W. (1860–1916), United Kingdom / Malaysia; administrative official in Sarawak (now part of Malaysia). *Trionyx pecki* Bartlett 1895a.

Peers, Bertram (1903–1939), South Africa; amateur archeologist and naturalist who collected the species. *Pseudomopus signatus peersi* Hewitt 1935.

Perrault, Claude (1613–1688), France; architect, physician, and anatomist, and the original describer of the species in the pre-Linnaean literature (Perrault 1676). *Testudo perraultii* Duméril and Bibron 1835.

Peters, James A. (1922–1972), USA; zoologist, herpetologist, and curator at the Smithsonian U.S. National Museum, D.C. *Geochelone petersi* Freiberg 1973.

Peters, Wilhelm K.H. (1815–1883), Germany; zoologist, herpetologist, and curator at the Museum für Naturkunde, Berlin. *Cyclanorbis petersii* Gray 1854a; *Batagur (Morenia) petersi* Anderson 1879.

Pfeiffer, Ida (1797–1858), Austria; explorer and the first European woman to cross Borneo who collected specimens of *Amyda cartilaginea* on Ambon and Seram (now part of Indonesia), and donated them to the British Museum. *Ida* Gray 1873h.

Phayre, Sir Arthur Purves (1812–1885), United Kingdom / Myanmar; army officer and eventual Commissioner in Burma (now Myanmar) who collected the type while a Captain serving in Arakan. *Testudo phayrei* Blyth 1854; *Trionyx phayrei* Theobald 1868b; *Trionyx phayrii* Boulenger 1889.

Philippen, Hans-Dieter (1957–2016), Germany; amateur herpetologist and turtle researcher. *Ocadia philippeni* McCord and Iverson 1992.

Pickering, Charles (1805–1878), USA; physician and naturalist who collected turtles for Holbrook. *Cistudo pickeringi* Holbrook *in* Duméril 1855.

Pilgrim, Henry G.E. (1875–1943), United Kingdom / India; geologist, paleontologist, and superintendent of the Geological Survey of India. *Geoemyda pilgrimi* † Prasad and Satsangi 1967.

Polyakov, Ivan S. (1845–1887), Russia; zoologist, anthropologist, and ethnographer at the Zoological Museum of the Russian Academy of Sciences, St. Petersburg, who collected the types. *Pelochelys poljakowii* Strauch 1890.

Porter, Captain David (1780–1843), USA; naval commander, the first to explore Santa Cruz, Galapagos, first named Porter's Island, and then later Indefatigable, who first noted differences among the tortoises from various Galapagos Islands (Porter 1815). *Testudo porteri* Rothschild 1903.

Praschag, Peter [1974–], Austria; turtle biologist who first recognized the species as distinct. *Cuora praschagi* Blanck, Gaillard, Protiva, Wheatley, Shi, Liu, Ray, and Anders 2023.

Pritchard, Peter C.H. (1943–2020), United Kingdom / USA; herpetologist and a Co-Chair of the IUCN Tortoise and Freshwater Turtle Specialist Group who founded Chelonian Research Institute, Florida. *Chelodina pritchardi* Rhodin 1994a; *Mauremys pritchardi* McCord 1998.

Procter, Joan B. (1897–1931), United Kingdom; female zoologist, herpetologist, and curator at the British Museum of Natural History, London. *Testudo procterae* Loveridge 1923.

Prouvençal de Saint-Hilaire, Augustin F.C. (1779–1853), France; botanist who collected in Brazil and Argentina. *Platemys hilarii* Duméril and Bibron 1835; *Hydraspis hilairii* Gray 1844.

Purgotti, Sebastiano (1799–1879), Italy; chemist, mathematician, and philosopher at the University of Perugia. *Testudo purgottii* † Ceselli 1846.

Purvis, Malcolm (?–), Australia; amateur herpetologist. *Elseya purvisi* Wells and Wellington 1985.

Racovitza (or Racoviță), Emil G. (1868–1947), Romania; biologist, zoologist, and president of the Romanian Academy. *Testudo ibera racovitzai* Călinescu 1931.

Rankin, Peter R. (1955–1979), Australia; amateur herpetologist. *Chelodina rankini* Wells and Wellington 1985; *Chelodina rankini* Wells 2007a.

Reeves, John R., Jr. (1804–1877), United Kingdom / China; naturalist who worked for the East India Company in China and collected turtles. The entry for Reeves in Beolens et al. (2011) is incorrect, referring instead to the father. *Emys reevesii* Gray 1831d.

Reimann, Michael J. (1952–), Germany; herpetologist who first recognized

the species as distinct. *Chelodina reimanni* Philippen and Grossmann 1990.

Retzius, Anders J. (1742–1821), Sweden; naturalist at Lund University. *Testudo retzii* Daudin 1801; *Monoclida retziana* Rafinesque 1832.

Rhodin, Anders G.J. (1949–), Sweden / USA; orthopedic surgeon and herpetologist, former Chair of the IUCN Tortoise and Freshwater Turtle Specialist Group and founder of Chelonian Research Foundation, who studied in Papua New Guinea and collected and first recognized the species as distinct. *Elseya (Hanwarachelys) rhodini* Thomson, Amepou, Anamiato, and Georges 2015.

Riggs, Elmer S. (1869–1963), USA; paleontologist and curator at the Field Museum of Natural History, Chicago. *Testudo riggsi* † Hibbard 1944.

Rossignon, Jules (c.1810–1883), France / Guatemala; scientist, agriculturist (coffee grower), and naturalist who lived and worked in Guatemala. *Emysaurus rossignonii* Bocourt 1868.

Rudolf, Crown Prince (1858–1889), Austria; royal nobleman for whom Lake Rudolf (now Lake Turkana) was also named. *Sternothaerus rudolphi* † Arambourg 1947.

Rüppell, Eduard (1794–1884), Germany; mammalogist, naturalist, and explorer. *Caouana ruppellii* Gray 1844.

Rustamov, Anver K. (1917–2005), Soviet Union / Turkmenistan; zoologist, ornithologist, and herpetologist. *Agrionemys horsfieldi rustamovi* Chkhikvadze, Amiranashvili, and Ataev 1990.

Sabin, Andrew (1945–), USA; conservationist and philanthropist. *Mesoclemmys sabiniparaensis* Cunha, Sampaio, Carneiro, Vogt, Mittermeier, Rhodin, and Andrade 2022.

Salvin, Osbert (1835–1898), United Kingdom; ornithologist, herpetologist, and explorer of Guatemala, who obtained the type. *Staurotypus (Stauremys) salvinii* Gray 1864c; *Dermatemys salvinii* Gray 1870c; *Emys (Clemmys) salvini* Günther 1885.

Sauzier, Théodore (1829–1904), France / Mauritius; naturalist and paleontologist who lived and worked on Réunion and Mauritius. *Testudo sauzieri* Gadow 1894.

Schlegel, Hermann (1804–1884), Germany / Austria / Netherlands; ornithologist and herpetologist who worked in Austria and the Netherlands and was director of the Rijksmuseum van Natuurlijke Historie, Leiden. *Trionyx schlegelii* Brandt 1857; *Sphargis coriacea schlegelii* Garman 1884.

Schlotheim, Ernst F. von (1764–1832), Germany; geologist and paleontologist. *Clemmys schlotheimii* † Fitzinger 1835; *Trionyx schlotheimii* † Fitzinger 1835.

Schmacker, Bernhard (1852–1896), Germany / China; malacologist who worked in China and collected the type. *Clemmys schmackeri* Boettger 1894.

Schneider, Johann G. (1750–1822), Germany; classicist, naturalist, and turtle taxonomist. *Emys schneideri* Schweigger 1812.

Schoepff (or Schöpf), Johann David (1752–1800), Germany; physician, botanist, herpetologist, and turtle taxonomist. *Testudo schoepfii* Fitzinger 1826; *Phrynops schöpfii* Fitzinger *in* Diesing 1839; *Testudo schoepfii* Rüppell *in* Gray 1873j; *Phrynops schoepffii* Fitzinger *in* Siebenrock 1904b.

Schonland (or Schönland), Selmar (1860–1940), Germany / South Africa; botanist and curator at the Albany Museum, Grahamstown. *Chersinella schonlandi* Hewitt 1934.

Schultze-Jena, Leonhard (1872–1955), Germany; explorer, zoologist, and anthropologist who explored in New Guinea and collected the types. *Emydura schultzei* Vogt 1911.

Schweigger, Augustus F. (1783–1821), Germany; naturalist, herpetologist, and turtle taxonomist. *Testudo schweiggeri* Fitzinger 1826; *Testudo schweigeri* Gray 1830e; *Testudo schweiggeri* Gray *in* Duméril and Bibron 1835; *Platemys schweiggerii* Duméril and Bibron 1835; *Geochelone (Geochelone) schweiggeri* Fitzinger 1835.

Schweinfurth, Georg A. (1836–1925), Germany; botanist, paleontologist, and African explorer who collected one of the types. *Pelomedusa schweinfurthi* Petzold, Vargas-Ramírez, Kehlmaier, Vamberger, Branch, Du Preez, Hofmeyr, Meyer, Schleicher, Široký, and Fritz 2014.

Seba, Albertus (1665–1736), Netherlands; pharmacist and naturalist with a large natural history collection. *Emys sebae* Gray 1831d; *Emys seba* Gray 1844; *Geoclemmys sebae* Gray 1869a; *Peltastes stellatus seba* Gray 1870c.

Seimund, Elbert C.H. (1878–1942), United Kingdom / South Africa; military medical technician and taxidermist who worked for the British Museum in South Africa and collected the type. *Testudo seimundi* Boulenger 1903b.

Selenka, Margarethe L. (1860–1922), Germany / Indonesia; female zoologist, paleontologist, and women's rights and peace activist, who worked at the Trinil site on Java, Netherlands Indies (now Indonesia). *Chitra selenkae* † Jaekel 1911.

Sellards, Elias H. (1875–1961), USA; paleontologist and geologist at the University of Florida, Gainesville. *Testudo sellardsi* † Hay 1916a.

Serié, Pedro (1865–1951), Argentina; zoologist, herpetologist, and curator at the Museo Argentino de Ciencias Naturales, Buenos Aires. *Kinosternon scorpioides seriei* Freiberg 1936.

Shaw, George (1751–1813), United Kingdom; naturalist at the British Museum of Natural History, London. *Kinosternon shavianum* Bell 1825a; *Cinosternon shawianum* Bocourt 1876a.

Shiu, Oscar (1959–), China; animal exporter in Hong Kong who obtained the types. *Platysternon megacephalum shiui* Ernst and McCord 1987.

Siebenrock, Friedrich (1853–1925), Austria; zoologist, herpetologist, and curator at the Naturhistorisches Museum, Vienna. *Chelodina siebenrocki* Werner 1901a; *Batagur siebenrocki* † Jaekel 1911; *Siebenrockiella* Lindholm 1929; *Mauremys caspica siebenrocki* Wischuf *in* Maran 1996; *Mauremys caspica siebenrocki* Wischuf and Fritz *in* Fritz and Wischuf 1997.

Sigriz, Adalbert von (1800s), Germany; physician and explorer who obtained turtles identified as *Clemmys caspica* at Ragusa (Sicily) on the Adriatic Sea, used by the author as comparison for his description of Spanish *C. sigriz*. *Clemmys sigriz* Michahelles 1829; *Emys sigritzii* Michahelles *in* Gray 1831d.

Simon, G. Eugène (1829–1896), France / China; agricultural engineer and French consul who collected specimens in China that were labeled as a new species by A.H.A. Duméril at the Muséum National d'Histoire Naturelle, Paris. *Gymnopus simonii* Duméril *in* David 1875.

Smith, Sir Andrew (1797–1872), Scotland, United Kingdom / South Africa; physician, administrator, and zoologist, who worked primarily in South Africa. *Batagur smithii* Gray 1863g; *Testudo smithi* Boulenger 1886a.

Smith-Stanley, Edward, 13th Earl of Derby (1775–1851), United Kingdom; nobleman and naturalist who obtained the type specimen. *Sternotherus derbianus* Gray 1844.

Sonnerat, Pierre (1748–1814), France; natural scientist and explorer who obtained the type. *Testudo sonnerati* Meyer 1790.

Speke, John H. (1827–1864), United Kingdom; explorer and first European to reach Lake Victoria, the source of the Nile River, and who collected the type. *Kinixys spekii* Gray 1863d.

Spengler, Lorenz (1720–1807), Switzerland / Denmark; educator and malacologist, and owner of the type in his private museum. *Testudo spengleri* Gmelin 1789.

Spix, Johann Baptist von (1781–1826), Germany; zoologist, herpetologist, turtle taxonomist, and explorer of the Brazilian Amazon. *Platemys spixii* Duméril and Bibron 1835.

Spooner, Charles S. (1885–1963), USA; zoologist and entomologist at Eastern Illinois State University. *Kinosternon flavescens spooneri* Smith 1951.

Spurrell, Herbert G.F. (1877–1918), United Kingdom; physician and zoologist who collected the type. *Cinosternum spurrelli* Boulenger 1913.

Steindachner, Franz (1834–1919), Austria; ichthyologist and director of the Naturhistorisches Museum, Vienna. *Sternothaerus steindachneri* Siebenrock 1902a; *Trionyx steindachneri* Siebenrock 1906a; *Cinosternum steindachneri* Siebenrock 1906b; *Chelodina steindachneri* Siebenrock 1914.

Stejneger, Leonhard H. (1851–1943), Norway / USA; herpetologist, ornithologist, curator at the United States National Museum, Washington, D.C. *Pseudemys stejnegeri* Schmidt 1928; *Kinosternon flavescens stejnegeri* Hartweg 1938.

Stirling, Stanley (1918–2012), Australia; amateur naturalist and private zoological park owner. *Elseya stirlingi* Wells and Wellington 1985; *Elseya sterlingi* Iverson, Thomson, and Georges 2001b; *Elseya stirlingi* Wells 2007b.

Stobaeus, Kilian (1690–1742), Sweden; physician, paleontologist, and naturalist. *Testudo stobaeana* Gmelin *in* Schoepff 1793a.

Stolley, George (1817–1911), Germany / USA; naturalist and geologist working with Agassiz who collected specimens of *Apalone spinifera*

emoryi. *Aspidonectes georgii* Gray 1870c.

Strauch, Alexander A. (1832–1893), Russia; herpetologist and director of the Zoological Museum of the Imperial Academy of Sciences at St. Petersburg. *Testudo strauchi* Lidth de Jeude 1893.

Sumeire, Benoit Camille (1850–1920), Mauritius; businessman and port station director who provided photos of the type. *Testudo sumeirei* Sauzier 1892.

Swinhoe, Consul Robert (1836–1877), India / United Kingdom / China; naturalist and diplomat who worked in Formosa (Taiwan) and China and collected birds, mammals, and reptiles. *Oscaria swinhoei* Gray 1873g; *Trionyx swinhonis* Boulenger 1889.

Taylor, Edward H. (1889–1978), USA / Philippines; zoologist and herpetologist at the University of Kansas who also worked in the Philippines. *Pseudemys scripta taylori* Legler 1960.

Temminck, Coenraad Jacob (1778–1858), Netherlands; zoologist and director of the Rijksmuseum van Natuurlijke Historie, Leiden. *Chelonura temminckii* Troost *in* Harlan 1835.

Terbish, Khayankhyarvaa (1952–), Mongolia; herpetologist at the National University of Mongolia who collected the type. *Agrionemys kazachstanica terbishi* Chkhikvadze 2009.

Theobald, William J. (1829–1908), United Kingdom / India; naturalist, herpetologist, and deputy superintendent of the Geological Survey of India. *Chaibassia theobaldi* Anderson 1879.

Thunberg, Carl Peter (1743–1828), Sweden; naturalist and curator of the Linnaean collection in Uppsala. *Caretta thunbergii* Merrem 1820.

Tornier, Gustav (1859–1938), Germany; zoologist, herpetologist, and curator at Humboldt University, Berlin. *Testudo tornieri* Siebenrock 1903b.

Trelease, William (1857–1945), USA; botanist, director of the Missouri Botanical Garden, and president of the Academy of Science of St. Louis. *Chrysemys treleasei* Hurter 1911.

Trimen, Roland (1840–1916), United Kingdom / South Africa; entomologist who worked in South Africa and became director of the South African Museum. *Testudo trimeni* Boulenger 1886a.

Tristram, Henry B. (1822–1906), United Kingdom; clergyman, Bible scholar, and ornithologist, who collected the type. *Emys tristrami* Gray 1869a.

Troost, Gerard (1776–1850), Netherlands / USA; geologist and naturalist and first president of the Academy of Natural Sciences, Philadelphia, who collected the types. *Emys troostii* Holbrook 1836.

Twente, John W. (1887–1966), USA; paleontologist at the Univeristy of Kansas who collected the type. *Emys twentei* † Taylor 1943.

Uhrig, Dennis W. (1952–), USA; amateur herpetologist and turtle breeder. *Trachemys venusta uhrigi* McCord, Joseph-Ouni, Hagen, and Blanck 2010.

Vaillant, Léon Louis (1834–1914), France; vertebrate zoologist at the Muséum National d'Histoire Naturelle, Paris. *Tetrathyra vaillantii* Rochebrune 1884.

Van Denburgh, John (1872–1924), USA; physician, herpetologist, and curator at the California Academy of Sciences, who studied Galapagos tortoises. *Testudo vandenburghi* DeSola 1930.

van Dijk, Peter Paul (1967–), Netherlands / Malaysia / USA; herpetologist, conservationist, and a Co-Chair of the IUCN Tortoise and Freshwater Turtle Specialist Group, who confirmed the occurrence of *Chitra* in Myanmar. *Chitra vandijki* McCord and Pritchard 2003a.

van Hasselt, Johan Conrad (1797–1823), Netherlands / Indonesia; physician and zoologist who worked on Java in the Dutch East Indies (now Indonesia). *Emys hasseltii* Boie *in* Fitzinger 1826.

Vanderhaege, Maurice (1938–), France; amateur herpetologist. *Phrynops tuberculatus vanderhaegei* Bour 1973.

Vanmeerhaeghe, Bernard (1950–1995), France; chemist and amateur herpetologist. *Mauremys leprosa vanmeerhaeghei* Bour and Maran 1999.

Verreaux, Jules P. (1807–1873), France / South Africa; botanist, ornithologist, and naturalist collector and trader in South Africa, who helped Smith found the South African Museum, Cape Town. *Testudo verroxii* Smith 1839b; *Peltastes verreauxii* Gray 1870b.

Victoria, Queen (1819–1901), United Kingdom; British monarch for whom the Victoria River in Australia was also named. *Hydraspis victoriae* Gray 1842.

Vijaya, Jagannathan (1959–1987), India; female herpetologist who studied the turtles of this genus. *Vijayachelys* Praschag, Schmidt, Fritzsch, Müller, Gemel, and Fritz 2006.

Viosca, Percy, Jr. (1892–1961), USA; biologist and herpetologist at Tulane University, Louisiana, who collected the types. *Pseudemys vioscana* Brimley 1928.

Vogel, Zdeněk (1913–1986), Czechoslovakia; writer and amateur herpetologist, who obtained the type. *Platysternon megacephalum vogeli* Wermuth 1969.

Vogl, Father Cornelius (1884–1959), Germany / Venezuela; priest and naturalist who worked in Venezuela and collected the type. *Podocnemis vogli* Müller 1935.

Vogt, Richard C. (1949–2021), USA / Mexico / Brazil; herpetologist and turtle ecology researcher who lived and worked in all three countries. *Kinosternon vogti* López-Luna, Cupul-Magaña, Escobedo-Galván, González-Hernández, Centenero-Alcala, Rangel-Mendoza, Ramírez-Ramírez, and Cazares-Hernández 2018.

Vosmaer, Arnout (1720–1799), Netherlands; naturalist and curator of a private natural history collection. *Testudo indica*. *Vosmaeri* Schoepff 1795; *Testudo indica vosmaeri* Suckow 1798.

Voznesensky, Ilya G. (1816–1871), Russia; naturalist and explorer who collected the type in California. *Clemmys wosnessenskyi* Strauch 1862.

Wagler, Johann G. (1800–1832), Germany; ornithologist and herpetologist at the Ludwig Maximilian University of München. *Platemys waglerii* Duméril and Bibron 1835; *Hydromedusa wagleri* Günther 1884.

Walbaum, Johann J. (1724–1799), Germany; physician, naturalist, and taxonomist. *Phrynops walbaumi* Fitzinger *in* Siebenrock 1904b.

Wallace, Alfred Russell (1823–1913), United Kingdom; naturalist who, with Charles Darwin, championed the idea of evolution by natural selection. *Testudo wallacei* Rothschild 1902.

Watson, Colonel John W. (1838–1889), United Kingdom / India; military officer in India who obtained the type. *Clemmys watsoni* † Lydekker 1886.

Weissinger, Heinz (1945–1993), Austria; amateur herpetologist. *Testudo marginata weissingeri* Trutnau 1994; *Testudo weissingeri* Bour 1996.

Wermuth, Heinz (1918–2002), Germany; herpetologist and curator at the Staatliches Museum für Naturkunde, Stuttgart. *Phrynops wermuthi* Mertens 1969b.

Werner, Franz (1867–1939), Austria; zoologist and herpetologist who worked as a professor at the University of Vienna. *Platemys werneri* Schnee 1900.

Werner, Yehudah L. (1931–), Germany / Israel; zoologist and herpetologist at the Hebrew University, Jerusalem. *Testudo werneri* Perälä 2001.

Wettstein-Westersheim, Otto von (1892–1967), Austria; herpetologist and curator at the Naturhistorisches Museum, Wien. *Pelomedusa subrufa wettsteini* Mertens 1937.

White, Reverend Gilbert (1720–1793), United Kingdom; naturalist and popular writer who obtained the type. *Testudo whitei* Bennett *in* White 1836.

Whitney, Francis L. (1878–1962), USA; paleontologist at Texas State University, Austin, who obtained the type. *Terrapene whitneyi* † Hay 1916b.

Wied-Neuwied, Prince Alexander Philipp Maximilian zu (1782–1867), Germany; nobleman, naturalist, and explorer of Brazil and USA. *Emys maximiliani* Mikan 1825; *Emys maximiliana* Gray 1830e; *Platemys neuwiedii* Duméril and Bibron 1835.

Williams, Ernest E. (1914–1998), USA; herpetologist and curator at the Museum of Comparative Zoology, Harvard University. *Pelusios williamsi* Laurent 1965; *Phrynops williamsi* Rhodin and Mittermeier 1983.

Wilson, John A. (1914–2008), USA; paleontologist and geologist at the University of Texas, Austin. *Testudo wilsoni* † Milstead 1956.

Worrell, Eric F.A. (1924–1987), Australia; naturalist and herpetologist. *Tropicochelymys worrelli* Wells and Wellington 1985.

Young, Reverend William Paulin (1886–1969), United Kingdom / South Africa; clergyman and naturalist who presented the types. *Kinixys youngi* Hewitt 1931.

Yuwono, Frank (1958–), Indonesia; animal exporter who obtained the types. *Geoemyda yuwonoi* McCord, Iverson, and Boeadi 1995.

Zarudny, Nikolai A. (1859–1919), Russia; zoologist, herpetologist, and explorer. *Testudo zarudnyi* Nikolsky 1896.

Zhou, Jiufa (1938–), China; economist, naturalist, and founder of the Nanjing Turtle Museum. *Cuora zhoui* Zhao *in* Zhao, Zhou, and Ye 1990.

Zug, George R. (1938–), USA; herpetologist and curator at the United States National Museum, Washington D.C. *Pseudemys concinna gorzugi* Ward 1984.

APPENDIX 2

Institutional (Museum) Acronyms Used for Turtle and Tortoise Type Specimens in this Checklist

AMF = Australian Museum Palaeontology Department, Sydney, Australia.
AMG = Albany Museum, Grahamstown, South Africa.
AMNH = American Museum of Natural History, New York, New York, USA.
AMPG = Athens Museum of Palaeontology and Geology, University of Athens, Athens, Greece.
AMS = Australian Museum, Sydney, New South Wales, Australia.
ANGSSR = see GNM.
ANSP = Academy of Natural Sciences, Philadelphia, Pennsylvania, USA.
ANU = Anhui Normal University Museum, Wuhu, Anhui, China.
AUM = Auburn University Museum of Natural History, Auburn, Alabama, USA.
AZM = Australian Zoological Museum (Private), Katoomba, New South Wales, Australia.
BEG = Bureau of Economic Geology, University of Texas, Austin, Texas, USA.
BMNH = see NHMUK.
BYU = Brigham Young University, Monte L. Bean Life Science Museum, Provo, Utah, USA.
CAS = California Academy of Sciences, San Francisco, California, USA.
CAS-SU = California Academy of Sciences, Stanford University Collection, San Francisco, California, USA.
CHM = Charleston Museum, Charleston, South Carolina, USA.
CIB = Chengdu Institute of Biology (formerly Sichuan Biological Research Institute, SBRI), Chengdu, Sichuan, China.
CJSHS = Chang Jung Senior High School, Tainan, Taiwan.
CM = Carnegie Museum of Natural History, Pittsburgh, Pennsylvania, USA.
CNAR = Colección Nacional de Anfibios y Reptiles, Instituto de Biología, Universidad Nacional Autónoma de México, Mexico City, Mexico.
CRES = Centre for Natural Resources Management and Environmental Studies, Hanoi National University, Hanoi, Vietnam.
CRI = see PCHP.
DBULPGC = Departamento de Biología, Universidad de Las Palmas de Gran Canaria, Spain.
EBD = Estación Biológica de Doñana, Sevilla, Spain.
EBRG = Estación Biológica de Rancho Grande, Museo, Maracay, Aragua, Venezuela.
ESK = Department of Earth Sciences, Kagoshima University, Kagoshima, Japan.
FGS = see USNM.
FMNH = Field Museum of Natural History, Chicago, Illinois, USA.
FU = Fudan University, Department of Biology, Shanghai, China.
GNM = Georgian National Museum (formerly Georgian State Museum [GSM] and Georgian Institute of Paleobiology [ANGSSR, IPANG, and IPGAS]), Tbilisi (Tiflis), Georgia.
GSI = Geological Survey of India, Kolkata, India.
GSM = see GNM.
GXUA = Guangxi Agricultural University, Nanning, Guangxi, China.
HNNU = Hunan Normal University, Animal Museum, College of Life Sciences, Changsha, Hunan, China.
HNU = Hainan Normal University, Turtle Museum, Haikou, Hainan, China.
HUJ = Hebrew University of Jerusalem, Zoological Museum, Jerusalem, Israel.
IAVH = Instituto de Investigacion de Recursos Biologicos Alexander von Humboldt, Villa de Leyva, Colombia.
IBH = Instituto de Biología, Universidad Nacional Autónoma de México, Mexico City, Mexico.
IEBR = Institute of Ecology and Biological Resources, Hanoi, Vietnam.
IGF = Institute of Geology, Firenze (now Museo di Storia Naturale, Sezione di Geologia e Paleontologia, Università degli Studi di Firenze), Florence, Italy.
INHS = Illinois Natural History Survey, University of Illinois, Champaign, Illinois, USA.
INPA = Instituto Nacional de Pesquisas da Amazônia, Manaus, Amazonas, Brazil.
IPANG = see GNM.
IPGAS = see GNM.
IRSNB = Institut Royal des Sciences Naturelles de Belgique, Bruxelles (Brussels), Belgium.
IVPP = Institute of Vertebrate Paleontology and Paleoanthropology, Chinese Academy of Sciences (Academia Sinica), Beijing, China.
JNU = Jinan University, Guangzhou, China.
KM = Kimberley Museum (now McGregor Museum), Kimberley, South Africa.
KU = University of Kansas Natural History Museum, Lawrence, Kansas, USA.
KUZ = Department of Zoology, Kyoto University, Kyoto, Japan.
KZM = Königsberger Zoologischen Museums, University of Königsberg, Königsberg, Germany (now Kaliningrad, Russia).
LACM = Natural History Museum of Los Angeles County, Los Angeles, California, USA.
LM = Leverian Museum (private, dispersed), London, England, United Kingdom.
LSUM = Louisiana State University Museum of Natural Science, Baton Rouge, Louisiana, USA.
LUZM = Lunds Universitet Zoologiska Museum (Lund University Biological Museum), Lund, Sweden.
MACN = Museo Argentino de Ciencias Naturales "Bernardino Rivadavia", Buenos Aires, Argentina.
MAF = see NRM and UPSZTY.
MB = see ZMB.
MBLSSC = Museum of the Biological Laboratory of the Science Society of China (now National Research Institute of Biology, Chinese Academy of Sciences), Nanjing, Jiangsu, China.
MCZ = Museum of Comparative Zoology, Harvard University, Cambridge, Massachusetts, USA.
MDG = see NRM and UPSZTY.
MGA = see NRM and UPSZTY.
MHD = Museo del Hombre Dominicano, Santo Domingo, Dominican Republic.
MHNG = Muséum d'Histoire Naturelle de la Ville de Genève, Geneva, Switzerland.
MHNL = Musée d'Histoire Naturelle de Lyon (now Musée des Confluences), Lyon, France.
MHNT = Muséum d'Histoire Naturelle de la Ville de Toulouse, Toulouse, France.
MLUH = Martin-Luther-Universität Halle-Wittenberg, Zentralmagazin Naturwissenschaftlicher Sammlungen (formerly Institut für Zoologie and Geiseltal Museum), Halle (Saale), Germany.
MMK = McGregor Museum, Kimberley, Cape Province, South Africa.
MMNHN = Metropolitan Museum of Natural History (now Nanjing Museum, Chinese Academy of Sciences (Academia Sinica), Nanjing, Jiangsu, China.
MNB = see ZMB.
MNG = Museums der Natur Gotha, Gotha, Germany.
MNHN = Muséum National d'Histoire Naturelle (Reptiles), Paris, France.
MNHN(P) = Muséum National d'Histoire Naturelle (Paleontology), Paris, France.
MNHNC = Museo Nacional de Historia Natural, Santiago, Chile.
MNHNSD = Museo Nacional de Historia Natural "Profesor Eugenio de Jesús Marcano," Santo Domingo, Dominican Republic.
MNHW = Muzeum Przyrodnicze Uniwersytetu Wrocławskiego (Museum of Natural History, University of Wroclaw), Wrocław (Breslau), Poland.
MNM = Museum für Naturkunde, Magdeburg, Germany.
MNUL = Museum für Natur und Umwelt, Lübecker Museen, Lübeck, Germany.
MPEG = Museu Paraense "Emilio Goeldi," Belém, Pará, Brazil.
MRAC = Musée Royal de l'Afrique Centrale (Koninklijk Museum voor Midden Afrika), Tervuren, Belgium.
MRHA = Museo Regional de Historia de Aguascalientes, Aguascalientes, Mexico.
MSNG = Museo Civico di Storia Naturale "Giacomo Doria" di Genova, Genoa, Italy.
MSNM = Museo Civico di Storia Naturale, Milano (Milan), Italy.
MSNVE = Museo di Storia Naturale di Venezia, Venice, Italy.
MSW = Menagerie zu Schönbrunn (now Tiergarten Schönbrunn), Wien (Vienna), Austria.
MTD = Museum für Tierkunde, Senckenberg Naturhistorische Sammlungen, Dresden, Germany.
MTKD = see MTD.
MUHNAC = Museu Nacional de História Natural e da Ciência, Universidade de Lisboa (formerly Museu Bocage and Secção Zoológica do Museu de Lisboa), Lisbon, Portugal.
MVZ = Museum of Vertebrate Zoology, University of California at Berkeley, California, USA.
MWNH = Museum Wiesbaden Naturhistorische Landessammlung, Wiesbaden, Germany (has also used acronym NWSW).
MZB = Museum Zoologicum Bogoriense (now Bidang Zoologi, Pusat Penelitian Biologi, Lembaga Ilmu Pengetahuan Indonesia), Bogor, Java, Indonesia.
MZFC-HE = Colección de Herpetología de la Facultad de Ciencias de la Universidad Nacional Autónoma de México, Mexico City, Mexico.

MZP = Museo di Zoologia di Padova, Italy.
MZT = Zoological Museum, Tartu (now University of Tartu Natural History Museum), Tartu, Estonia.
MZUF = Museo di Storia Naturale, Museo Zoologico "La Specola", Université di Firenze, Florence, Italy.
MZUS = Musée Zoologique de la Ville de Strasbourg, Université de Strasbourg, Strasbourg, France.
MZUSP = Museu de Zoologia da Universidade de São Paulo, São Paulo, Brazil.
NCSM = North Carolina Museum of Natural Sciences, Raleigh, North Carolina, USA.
NHM(P) = Natural History Museum (Paleontology) (formerly British Museum), London, England, United Kingdom.
NHMUK = Natural History Museum [formerly British Museum (Natural History)], London, England, United Kingdom.
NM = Natal Museum (now KwaZulu-Natal Museum), Pietermaritzburg, Natal, South Africa.
NMB = National Museum, Bloemfontein, South Africa.
NMNZ = Museum of New Zealand Te Papa Tongarewa, Wellington, New Zealand.
NMP6V = Národní Muzeum (National Museum), Prague, Czech Republic.
NMSZ = National Museums of Scotland (formerly Royal Scottish Museum, RSM), Edinburgh, Scotland, United Kingdom.
NMV = National Museum of Victoria, Melbourne, Victoria, Australia.
NMW = Naturhistorisches Museum Wien (Vienna), Austria.
NRM = Naturhistoriska Riksmuseet (Swedish Museum of Natural History) (formerly included Museum Adolphi Friderici [MAF], Museum Gustavi Adolphi [MGA], and Museum De Geer [MDG]), Department of Vertebrate Zoology, Stockholm, Sweden.
NTM = Northern Territory Museum (now Museums and Art Galleries of the Northern Territory), Darwin, Northern Territory, Australia.
NTM = National Taiwan Museums, Taipei, Taiwan.
NTM(P) = Nanjing Turtle Museum (Private), Nanjing, Jiangsu, China.
NTUM = National Taiwan University Museums, Institute of Zoology, Taipei, Taiwan.
NZMC = National Zoological Museum, Institute of Zoology, Chinese Academy of Sciences (formerly Fan Memorial Institute of Biology), Beijing, China.
OUM = Oxford University Museum, Oxford, England, United Kingdom.
PCHP (CRI) = Peter C.H. Pritchard Collection, Chelonian Research Institute, Oviedo, Florida, USA (now at Turtle Conservancy, Ojai, California, USA).
PEM = Port Elizabeth Museum, Bayworld, Port Elizabeth, South Africa.
PIK = Pedagogical Institute of Khovd, Khovd University, Mongolia.
PNGM = Papua New Guinea National Museum, Port Moresby, Papua New Guinea.
PPHM = Panhandle-Plains Historical Museum, Canyon, Texas, USA.
PUM = Punjab University, Museum of Centre of Advanced Study in Geology, Chandigarh, Punjab, India.
QM = Queensland Museum, Brisbane, Queensland, Australia.
RCSM = Royal College of Surgeons Museum, Hunterian Museum, London, England, United Kingdom.
RHK = Rua Hoan Kiem (= Hoan Kiem softshell turtle), stuffed specimen at Ngoc Son Temple [Hung Ky Temple], Hoan Kiem Lake, Hanoi, Vietnam.
RMNH = Rijksmuseum van Natuurlijke Historie (now Naturalis–Nationaal Natuurhistorisch Museum), Leiden, Netherlands.
RNM = Ratnapura National Museum, Ratnapura, Sri Lanka.
RSM = see NMSZ.
SBRI = see CIB.
SIZ = Shaanxi Institute of Zoology, Xian, Shaanxi, China.
SM(BCB) = Strecker Museum (now Mayborn Museum), Baylor University, Waco, Texas, USA.
SMF = Senckenberg Forschungsinstitut und Naturmuseum, Frankfurt, Germany.
SMK = Sarawak Natural History Museum (now Muzium Sejarah Semulajadi), Kuching, Sarawak, Malaysia.
SMNS = Staatliches Museum für Naturkunde, Stuttgart, Germany.
SNHM = Shanghai Natural History Museum, Jing'an, Shanghai, China.
SYS = Biological Museum of Sun Yat-Sen University (Zhongshan University), Guangzhou (Canton), China.
TCU = Texas Christian University Collection, Fort Worth, Texas, USA.
TCWC = Texas Cooperative Wildlife Collection, Texas A&M University, College Station, Texas, USA.
THNHM = Natural History Museum, Pathum Thani, Thailand.
TMM = Texas Memorial Museum (now Texas Vertebrate Paleontology Collections), Austin, Texas, USA.
TMP = Transvaal Museum of Natural History (now Ditsong National Museum of Natural History), Pretoria, Transvaal, South Africa.
TNHC = Texas Natural History Collections, Texas Memorial Museum, University of Texas at Austin, Texas, USA.
TNZ = Tianjin Natural History Museum (formerly Musée Hoang-ho Pai-ho), Tianjin, China.
TT = The Tortoise Trust (Private), London, England, United Kingdom.
TU = Tulane University Museum of Natural History, Belle Chasse, Louisiana, USA.
TxVP = Texas Vertebrate Paleontology Collections, Jackson Museum of Earth History, The University of Texas, Austin, Texas, USA.
UAHC = University of Alabama Herpetological Collection, Tuscaloosa, Alabama, USA.
UCM = University of Colorado, Museum of Natural History, Boulder, Colorado, USA.
UCMZ = University of California Museum of Zoology (now Museum of Vertebrate Zoology), Berkeley, California, USA.
UF = University of Florida, Florida Museum of Natural History, Gainesville, Florida, USA.
UIMNH = University of Illinois Museum of Natural History, Urbana, Illinois, USA.
UKZM = University of Karachi, Dept. of Zoology Museum (formerly Kurrachee Museum), Karachi, Pakistan.
UMMP = University of Michigan Museum of Paleontology, Ann Arbor, Michigan, USA.
UMMZ = University of Michigan Museum of Zoology, Ann Arbor, Michigan, USA.
UMZ = University Museum of Zoology, University of Cambridge, Cambridge, England, United Kingdom.
UNSW = University of New South Wales, Sydney, Australia.
UPSZTY = Uppsala Zoologiska Typsamling (Uppsala Zoological Type Collection), Museum of Evolution (formerly Uppsala University Zoological Museum, UUZM) (includes former Museum Adolphi Friderici [MAF], Museum Gustavi Adolphi [MGA], and Museum De Geer [MDG]), Uppsala University, Uppsala, Sweden.
USNM = United States National Museum (now National Museum of Natural History) (includes former Florida Geological Survey, FGS), Smithsonian Institution, Washington D.C., USA.
UU = University of Utah, Salt Lake City, Utah, USA.
UUZM = see UPSZTY.
UWZM = University of Wisconsin Zoological Museum, Madison, Wisconsin, USA.
VNUH = Vietnam National University, Hanoi, Vietnam.
VUM = Vinh University Museum, Vinh, Vietnam.
WAM = Western Australian Museum, Perth, Western Australia, Australia.
YPM = Peabody Museum of Natural History, Yale University, New Haven, Connecticut, USA.
YU = Yunnan University, Dept. of Biology, Kunming, Yunnan, China.
ZFMK = Zoologisches Forschungsmuseum Alexander Koenig, Bonn, Germany.
ZIN = Zoological Institute, Russian Academy of Sciences, St. Petersburg, Russia.
ZIUS = Zoologiska Institutionen, Stockholms Universitet, Stockholm, Sweden.
ZMB = Zoologischen Museum Berlin (now Museum für Naturkunde Berlin [MNB] and Paläontologisches Museum Berlin [MB]), Leibniz-Institut für Evolutions- und Biodiversitätsforschung an der Humboldt-Universität, Berlin, Germany.
ZMH = Zoological Museum Hamburg (Biozentrum Grindel und Zoologisches Museum), Universität Hamburg, Hamburg, Germany.
ZMMU = Zoological Museum of Moscow State University, Moscow, Russia.
ZMUG = Zoologisches Museum, Johann-Friedrich-Blumenbach-Institut für Zoologie und Anthropologie, Göttingen University, Göttingen, Germany (herpetology collection transferred to Zoologisches Forschungsmuseum Alexander Koenig [ZFMK] in 1977).
ZMUL = see LUZM.
ZMUP = see MZP.
ZSI = Zoological Survey of India, Kolkata (Calcutta), West Bengal, India.
ZSM = Zoologische Staatssammlung München (Munich), Germany.

Turtle Taxonomy Working Group – 2025
Author Biographies



Anders G.J. Rhodin, M.D., is a Swedish-American retired orthopedic surgeon as well as a turtle researcher and conservationist. Born in Sweden, he immigrated to the USA in 1958, received a B.A. from Dartmouth in 1971, an M.D. from the University of Michigan in 1977, did orthopedic surgical residency training at Yale, was in private orthopedic surgical practice in Massachusetts from 1982 to 2019, and now resides in Vermont. He has been doing research on turtles since 1972, initially as an Associate in Herpetology at the Museum of Comparative Zoology at Harvard. He is the Founder and Director of Chelonian Research Foundation, a philanthropic nonprofit organization established in 1992 for the production, publication, and support of worldwide turtle and tortoise research. He is Founding Editor and Publisher of *Chelonian Conservation and Biology*, a peer-reviewed leading scientific journal devoted exclusively to turtles and tortoises inaugurated in 1993, and *Chelonian Research Monographs*, a book-length series focused on turtle and tortoise research and conservation established in 1996. He was Co-Chair with John Behler and then Chair of the IUCN SSC Tortoise and Freshwater Turtle Specialist Group (TFTSG) from 2000 to 2012, and is currently its Executive Vice Chair and co-Red List Coordinator. He was awarded the Sir Peter Scott Award for Conservation Merit by the Species Survival Commission of the IUCN in 2012, the Behler Turtle Conservation Award by the TFTSG and Turtle Survival Alliance also in 2012, a Desert Tortoise Council Conservation Award in 2019, and a Lifetime Achievement Award by the International Sea Turtle Society in 2024. He served as Board Chair of the Turtle Conservancy from 2016 to 2024, and is the Founding Co-Chair of three turtle-focused philanthropic funds: the Turtle Conservation Fund (established 2002), the Congdon-Dickson Turtle Ecology Fund (established 2020), and the Turtle Taxonomy Fund (established 2021). He has been an Advisory Board member of the Mohamed bin Zayed Species Conservation Fund since 2012 and is also a founding Board Member of Turtle Survival Alliance and the African Aquatic Conservation Fund and its African Chelonian Institute.



John B. Iverson, Ph.D., is an American turtle and iguana ecologist, taxonomist, and conservationist. He was raised on a farm in Nebraska, holds a Ph.D. in Zoology from the University of Florida, and is a semi-retired Biology Research Professor at Earlham College in Indiana. Because of his interests in the natural history, ecology, and evolution of iguanas and turtles, he is currently on the steering committees of both the IUCN SSC Iguana Specialist Group, and the Tortoise and Freshwater Turtle Specialist Group (TFTSG). He has been involved with the Turtle Survival Alliance since its inception in 2001 (currently a Board member), is the Founding Co-Chair of the Turtle Taxonomy Fund, serves on the Advisory Review Board of the Turtle Conservation Fund, and is also on the Board of the International Iguana Foundation. He has been active in several herpetological societies, serving as Editor and President of the Herpetologists' League. He has maintained long-term field research programs with undergraduates on endangered iguanas in the Bahamas and on turtles in northern Indiana and western Nebraska. He has published nearly 400 papers, over 100 of which were co-authored by his undergraduate students, and over 700 photographs in publications other than his own. His awards include Phi Beta Kappa, the Sears-Roebuck Foundation Teaching Excellence and Campus Leadership Award, the George Mikisch Sutton Award in Conservation Research of the Southwestern Association of Naturalists, the Zoological Society of Japan's Zoological Science Award in Biodiversity and Evolution, Indiana Professor of the Year by the Carnegie Foundation for the Advancement of Teaching, and the Behler Turtle Conservation Award from the TFTSG and Turtle Survival Alliance in 2011. Since 1993, he has restored and managed a 76-acre woodlot/cornfield, planted 60,000 trees, constructed 55 amphibian wetlands, and permanently protected the property in a Conservation Easement.



Dr. Uwe Fritz is a German herpetologist, taxonomist, and biogeographer working on turtles and, to a lesser extent, Western Palearctic reptiles. He is a doctoral diplomate of Hohenheim University in Stuttgart. He worked for six years at Stuttgart Zoo (Wilhelma), where he was responsible for lower vertebrates and invertebrates. In 1996 he moved to the Museum für Tierkunde Dresden, now part of Senckenberg, a nationwide research institution specialized in natural history. As one of his first tasks in Dresden, he relocated the zoological collections of more than 6 million specimens from a war ruin into a new building. Subsequently, as Department Head and temporarily Director of the whole Dresden institute, he established new laboratories, among others molecular genetic facilities for sequencing historical and ancient DNA (hDNA and aDNA). He also initiated the relaunch of the scientific journals published by the museum (*Arthropod Systematics & Phylogeny*, *Vertebrate Zoology*) and established one of the largest collections of turtle skeletal specimens. He acts as Editor-in-Chief of *Vertebrate Zoology*, currently ranking among the top zoological journals in the Web of Science. For several years, he served as President and General Secretary of the German Society of Biological Systematics. In addition to his employment at the museum, he is also an Adjunct Professor at the University of Leipzig, where he has supervised several diploma, masters, and doctoral theses. His research interests include the diversity and geographic variation of widely distributed and controversial taxa. In this context, he has worked with turtles as well as water and grass snakes of the genus *Natrix*. Before the Nagoya Protocol came into force, he focused on widely distributed Asian and African turtle species. In recent years, using aDNA and hDNA, extinct giant tortoises have became another field of interest. Some recent projects have centered on the clarification of nomenclatural and systematic controversies using DNA from historical type specimens. He has for many years been a member of the IUCN SSC Tortoise and Freshwater Turtle Specialist Group and joined the Turtle Taxonomy Working Group for its 8th edition in 2017. In addition, he is a member of the Advisory Review Board for the Turtle Taxonomy Fund.



Natalia Gallego-García, Ph.D., is a Colombian biologist and conservation geneticist dedicated to studying and protecting freshwater turtles, primarily in Latin America. Born and raised in Colombia, she studied biology before moving to the Netherlands to pursue a Master's degree in Ecology and Evolution, specializing in computational biology. She earned a Ph.D. in Biological Sciences at the Universidad de los Andes in Bogotá, Colombia, in 2018, focusing on conservation genomics and the genetic management of endangered turtle species. Her work encompasses both field and laboratory settings, combining molecular science with applied conservation. She is passionate about using science to tackle real-world conservation challenges. Notable examples of her work include delineating conservation strategies to guide management plans, relocating confiscated individuals, assessing taxonomic uncertainty in key turtle species, and guiding breeding strategies in assurance colonies for the world's most endangered turtle species. She has worked with numerous organizations, including Conservation International and the Wildlife Conservation Society, and served as a postdoctoral researcher in Brad Shaffer's lab at UCLA, where she deepened her expertise in turtle population genetics and systematics. Since 2023, she has been the Director of Conservation Genetics for the Turtle Survival Alliance. She has been a member of the TFTSG since 2010 and serves on the Advisory Review Boards of the Turtle Conservation Fund and the Turtle Taxonomy Fund. She joined the Turtle Taxonomy Working Group in 2022 following the passing of former French member Roger Bour.



Arthur Georges, Ph.D., F.A.A., is an Australian ecologist and herpetologist. Born in Brisbane, Queensland, he studied mathematics at the University of Queensland until lured north by physicist Harry Messel as a volunteer on the crocodile research program in Arnhem Land. There he discovered the delights of field herpetology and so moved fields in his honours year to study thermoregulation in skinks. A Ph.D. on the turtles of Fraser Island cemented the transition, and there was no looking back. His research interests now lie in the evolution, ecology, and systematics of Australian reptiles and, in particular, freshwater turtles. A fundamental interest in these fascinating animals takes him into the field and the laboratory to learn more of their biology and to apply what he has learned in solving contemporary challenges for their conservation. He has published over 200 papers and is consistently in the top 2% of scientists in his discipline (Stanford Index). He has served as Dean of the Faculty of Applied Science at the University of Canberra and as President of the Australasian Wildlife Management Society and the Australian Society of Herpetologists. For many years he provided advice on threatened species and ecosystems to government as Chair of the ACT Scientific Committee. He is a founding Board Member of the Piku Biodiversity Network which coordinates conservation work in Papua New Guinea, and a member of the Advisory Review Board for the Turtle Taxonomy Fund. He was recently elected a Fellow of the Australian Academy of Science, and was honored with the Behler Turtle Conservation Award by the IUCN SSC Tortoise and Freshwater Turtle Specialist Group and Turtle Survival Alliance in 2025.



H. Bradley Shaffer (Brad), Ph.D., is an American geneticist and herpetologist and Professor at the University of California, Los Angeles (UCLA). He is Director of the UCLA La Kretz Center for California Conservation Science, and Distinguished Professor in both the Department of Ecology and Evolutionary Biology and the Institute of the Environment and Sustainability at UCLA. He earned a B.S. in Zoology from the University of California, Berkeley, in 1976, and a Ph.D. in Evolutionary Biology at the University of Chicago in 1982. He is a Fellow of the American Academy of Arts and Sciences, and was President of the American Association of Ichthyologists and Herpetologists and the American Genetics Association. He is on the Advisory Review Board for the Turtle Taxonomy Fund and was a co-founder of the Turtle Taxonomy Working Group with Anders Rhodin, James Parham, and Peter Paul van Dijk in 2005. He has published over 250 papers in the peer-reviewed scientific literature, many of which are on aspects of the systematics, conservation genomics, and field ecology of turtles and tortoises. He is also the Director and Principal Investigator of the California Conservation Genomics Project, a unique research effort to fully sequence the genomes of roughly 24,000 individual plants and animals from across California, and to use the results to predict, and protect, the most genetically resilient populations in the face of climate change. His team published the first complete turtle genome ever sequenced, and he hopes to continue using conservation genomic strategies to help stem the decline of turtles globally.



Peter Paul van Dijk, Ph.D., is a global turtle scientist and conservationist. Born and raised in the Netherlands, he was fascinated with turtles and other biodiversity, particularly aquatic animals, from a young age. He moved to Ireland to study zoology at University College Galway and was well on his way to becoming an Atlantic fisheries management biologist when some lucky circumstances created the opportunity to survey tortoises and freshwater turtles in Thailand. As a result, he spent most of the 1990s doing ecological research, field surveys, and conservation planning for the turtles of Thailand, as well as some initial surveys of the turtles of Myanmar. Seeing the impacts of the 'Asian Turtle Crisis' caused him to focus on turtle conservation through stricter regulation of exploitation and trade in wild turtles, including co-editing the Asian Turtle Trade monograph while working at TRAFFIC Southeast Asia in Malaysia. For the past 20 years, he has enjoyed residing in the mid-Atlantic USA turtle hotspot, working with Conservation International, Global Wildlife Conservation/Re:wild, and the Turtle Conservancy, as well as working with the IUCN SSC Tortoise and Freshwater Turtle Specialist Group (TFTSG) to assess the conservation status of tortoises and freshwater turtles for the IUCN Red List, and catalyzing better international trade regulation through CITES. He was Co-Chair of the TFTSG from 2012 to 2016, and is currently one of its Deputy Chairs. He was honored with the Behler Turtle Conservation Award by the TFTSG and Turtle Survival Alliance in 2017. He is a member of the Advisory Review Boards for the Turtle Conservation Fund, the Congdon-Dickson Turtle Ecology Fund, and the Turtle Taxonomy Fund. He has an enigmatic giant softshell from Myanmar named after him, and has had the privilege of seeing over 100 turtle species undisturbed in their native habitats.


CHELONIAN RESEARCH
C R F
FOUNDATION


TURTLE CONSERVANCY
TC


TURTLE TAXONOMY FUND
TTF


IUCN/SSC
TORTOISE AND
FRESHWATER TURTLE
SPECIALIST GROUP


TURTLE CONSERVATION FUND
TCF



IUCN



SSC
Species Survival Commission

(ISSN 1088-7105)

CHELONIAN RESEARCH MONOGRAPHS

Number 10 – 2025





ISBN: 978-0-9910368-6-8 (digital online); 978-0-9910368-7-5 (hard cover); 978-0-9910368-8-2 (soft cover)